Virus-Receptor Interactions of Glycosylated SARS-CoV-2 Spike and Human ACE2

Receptor

Peng Zhao[1,#], Jeremy L. Praissman[1,#], Oliver C. Grant[1,#], Yongfei Cai[2], Tianshu Xiao[2], Katelyn E.

Rosenbalm[1], Kazuhiro Aoki[1], Benjamin P. Kellman[3], Robert Bridger[1], Dan H. Barouch[4], Melinda

A. Brindley[5], Nathan E. Lewis[3,6], Michael Tiemeyer[1], Bing Chen[2], Robert J. Woods[1*], and Lance

Wells[1*]

[1] Complex Carbohydrate Research Center, Department of Biochemistry and Molecular

Biology, and Department of Chemistry, University of Georgia, Athens, Georgia, 30602, USA

[2] Division of Molecular Medicine, Children's Hospital and Department of Pediatrics, Harvard

Medical School, Boston, Massachusetts, 02115, USA

[3] Departments of Pediatrics and Bioengineering, University of California, San Diego, La Jolla,

California, 92093, USA

[4] Center for Virology and Vaccine Research, Beth Israel Deaconess Medical Center, Harvard

Medical School, Boston, Massachusetts, 02215, USA

[5] Department of Infectious Diseases, Department of Population Health, Center for Vaccines and

Immunology, College of Veterinary Medicine, University of Georgia, Athens, GA 30602, USA

[6] Novo Nordisk Foundation Center for Biosustainability at UC San Diego, La Jolla, California,

92093, USA

[#] Authors contributed equally

* To whom correspondence should be addressed: Robert J. Woods, rwoods@ccrc.uga.edu ,

and/or Lance Wells, lwells@ccrc.uga.edu

SUMMARY

The current COVID-19 pandemic is caused by the SARS-CoV-2 betacoronavirus, which utilizes its highly glycosylated trimeric Spike protein to bind to the cell surface receptor ACE2 glycoprotein and facilitate host cell entry. We utilized glycomics-informed glycoproteomics to characterize site-specific microheterogeneity of glycosylation for a recombinant trimer Spike mimetic immunogen and for a soluble version of human ACE2. We combined this information with bioinformatic analyses of natural variants and with existing 3D-structures of both glycoproteins to generate molecular dynamics simulations of each glycoprotein alone and interacting with one another. Our results highlight roles for glycans in sterically masking polypeptide epitopes and directly modulating Spike-ACE2 interactions. Furthermore, our results illustrate the impact of viral evolution and divergence on Spike glycosylation, as well as the influence of natural variants on ACE2 receptor glycosylation that, taken together, can facilitate immunogen design to achieve antibody neutralization and inform therapeutic strategies to inhibit viral infection.

Keywords: SARS-CoV-2; COVID-19; Spike protein; coronavirus; ACE2; glycoprotein; glycosylation; mass spectrometry; molecular dynamics; 3D-modeling

INTRODUCTION

The SARS-CoV-2 coronavirus, a positive-sense single-stranded RNA virus, is responsible for the severe acute respiratory syndrome referred to as COVID-19 that was first reported in China in December of 2019 (1). In approximately six months, this betacoronavirus has spread globally with more than 14 million people testing positive worldwide resulting in greater than 600,000 deaths as of July 20[th], 2020 (https://coronavirus.jhu.edu/map.html). The SARS-CoV-2 coronavirus is highly similar (nearly 80% identical at the genomic level) to SARS-CoV-1, which was responsible for the severe acute respiratory syndrome outbreak that began in 2002 (2,3). Furthermore, human SARS-CoV-2 at the whole genome level is >95% identical to a bat coronavirus (RaTG13), the natural reservoir host for multiple coronaviruses (1,4,5). Given the rapid appearance and spread of this virus, there is no current validated vaccine or SARS-CoV-2-specific targeting therapy clinically approved although statins, heparin, and steroids look promising for lowering fatality rates and antivirals likely reduce the duration of symptomatic disease presentation (6-12).

SARS-CoV-2, like SARS-CoV-1, utilizes the host angiotensin converting enzyme II (ACE2) for binding and entry into host cells (13,14). Like many viruses, SARS-CoV-2 utilizes a Spike glycoprotein trimer for recognition and binding to the host cell entry receptor and for membrane fusion (15). Given the importance of viral Spike proteins for targeting and entry into host cells along with their location on the viral surface, Spike proteins are often used as immunogens for vaccines to generate neutralizing antibodies and frequently targeted for inhibition by small molecules that might block host receptor binding and/or membrane fusion (15,16). In similar fashion, wildtype or catalytically-impaired ACE2 has also been investigated as a potential therapeutic biologic that might interfere with the infection cycle of ACE2 targeting coronaviruses (17,18). Thus, a detailed understanding of SARS-CoV-2 Spike binding to ACE2 is critical for elucidating mechanisms of viral binding and entry, as well as for undertaking the rational design of effective therapeutics.

The SARS-CoV-2 Spike glycoprotein consists of two subunits, a receptor binding subunit (S1)

and a membrane fusion subunit (S2) (1,2).   The Spike glycoprotein assembles into stable

homotrimers that together possess 66 canonical sequons for N-linked glycosylation (N-X-S/T,

where X is any amino acid except P) as well as a number of potential O-linked glycosylation sites

(19,20). Interestingly, coronaviruses virions bud into the lumen of the endoplasmic reticulum-

Golgi intermediate compartment, ERGIC, raising unanswered questions regarding the precise

mechanisms by which viral surface glycoproteins are processed as they traverse the secretory

pathway (21,22). While this and similar studies (19,23) analyze recombinant proteins, a previous

study on SARS-CoV-1 suggest that glycosylation of the Spike can be impacted by this

intracellular budding and remains to be investigated in SARS-CoV-2 (24). Nonetheless, it has

been proposed that this virus, and others, acquires a glycan coat sufficient and similar enough to

endogenous host protein glycosylation that it serves as a glycan shield, facilitating immune

evasion by masking non-self viral peptides with self-glycans (15,20-22). In parallel with their

potential masking functions, glycan-dependent epitopes can elicit specific, even neutralizing,

antibody   responses,   as   has   been   described   for   HIV-1   ((15,25-29),

https://www.biorxiv.org/content/10.1101/2020.06.30.178897v1).   Thus,   understanding   the

glycosylation of the viral Spike trimer is fundamental for the development of efficacious vaccines,

neutralizing antibodies, and therapeutic inhibitors of infection.

ACE2 is an integral membrane metalloproteinase that regulates the renin-angiotensin system

(30). Both SARS-CoV-1 and SARS-CoV-2 have co-opted ACE2 to function as the receptor by

which these viruses attach and fuse with host cells (13,14).   ACE2 is cleavable by ADAM

proteases at the cell surface (31), resulting in the shedding of a soluble ectodomain which can be

detected in apical secretions of various epithelial layers (gastric, airway, etc.) and in serum (32).

The N-terminal extracellular domain of ACE2 contains 6 canonical sequons for N-linked

glycosylation and several potential O-linked sites. Several nonsynonymous single-nucleotide

polymorphisms (SNPs) in the ACE2 gene have been identified in the human population and could

potentially alter ACE2 glycosylation and/or affinity of the receptor for the viral Spike protein (33).

Given that glycosylation can affect the half-life of circulating glycoproteins in addition to

modulating the affinity of their interactions with receptors and immune/inflammatory signaling

pathways (34,35), understanding the impact of glycosylation of ACE2 with respect to its binding

of SARS-CoV-2 Spike glycoprotein is of high importance. The proposed use of soluble

extracellular domains of ACE2 as decoy, competitive inhibitors for SARS-CoV-2 infection

emphasizes the critical need for understanding the glycosylation profile of ACE2 so that optimally

active biologics can be produced (17,18).

To accomplish the task of characterizing site-specific glycosylation of the trimer Spike of SARS-

CoV-2 and the host receptor ACE2, we began by expressing and purifying a stabilized, soluble

trimer Spike glycoprotein mimetic immunogen (that we define here and forward as S, (36)) and a

soluble version of the ACE2 glycoprotein from a human cell line. We utilized multiple mass

spectrometry-based approaches, including glycomic and glycoproteomic approaches, to

determine occupancy and site-specific heterogeneity of N-linked glycans. Occupancy (i.e. the

percent of any given residue being modified by a glycan) is an important consideration when

developing neutralizing antibodies against a glycan-dependent epitope. We also identified sites

of O-linked glycosylation and the heterogeneity of the O-linked glycans on S and ACE2. We

leveraged this rich dataset, along with existing 3D-structures of both glycoproteins, to generate

static and molecular dynamics models of S alone, and in complex with the glycosylated, soluble

ACE 2 receptor. By combining bioinformatic characterization of viral evolution and variants of the

Spike and ACE2 with molecular dynamics simulations of the glycosylated Spike-ACE2 interaction,

we identified important roles for glycans in multiple processes, including receptor-viral binding

and glycan-shielding of the Spike. Our rich characterization of the recombinant, glycosylated

Spike trimer mimetic immunogen of SARS-CoV-2 in complex with the soluble human ACE2

receptor provides a detailed platform for guiding rational vaccine, antibody, and inhibitor design.

RESULTS

Expression, Purification, and Characterization of SARS-CoV-2 Spike Glycoprotein Trimer and

Soluble Human ACE2

A trimer-stabilized, soluble variant of the SARS-CoV-2 Spike protein (S) that contains 22

canonical N-linked glycosylation sequons per protomer and a soluble version of human ACE2

that contains 6, lacking the most C-terminal 7th, canonical N-linked glycosylation sequons (**Fig.**

1A) were purified from the media of transfected HEK293 cells and the quaternary structure

confirmed by negative EM staining for the S trimer (**Fig. 1B**) and purity examined by SDS-PAGE

Coomassie G-250 stained gels for both (**Fig. 1C**). In addition, proteolytic digestions followed by

proteomic analyses confirmed that the proteins were highly purified (**Supplemental Table, Tab**

12). Finally, the N-terminus of both the mature S and the soluble mature ACE2 were empirically

determined via proteolytic digestions and LC-MS/MS analyses. These results confirmed that both

the secreted, mature forms of S protein and ACE2 begin with an N-terminal glutamine that has

undergone condensation to form pyroglutamine at residue 14 and 18, respectively (**Figs. 1D and**

S1). The N-terminal peptide observed for S also contains a glycan at Asn-0017 (**Fig. 1D**) and

mass spectrometry analysis of non-reducing proteolytic digestions confirmed that Cys-0015 of S

is in a disulfide linkage with Cys-0136 (**Fig. S2, Supplemental Table, Tab 2**). Given that SignalP

(37) predicts signal sequence cleavage between Cys-0015 and Val-0016 but we observed

cleavage between Ser-0013 and Gln-0014, we examined the possibility that an in-frame upstream

Methionine to the proposed start Methionine (**Fig. 1A**) might be used to initiate translation (**Fig.**

S3). If one examines the predicted signal sequence cleavage using the in-frame Met that is

encoded 9 amino acids upstream, SignalP now predicts cleavage between the Ser and Gln that

we observed in our studies (**Fig. S3**). To examine whether this impacted S expression, we

expressed constructs that contained or did not contain the upstream 27 nucleotides in a

pseudovirus (VSV) system expressing SARS-CoV2 S (**Fig. S4**) and in our HEK293 system (data

not shown). Both expression systems produced a similar amount of S regardless of which

expression construct was utilized (**Fig. S4**). Thus, while the translation initiation start site has still

not been fully defined, allowing for earlier translation in expression construct design did not have

a significant impact on the generation of S.

Glycomics Informed Glycoproteomics Reveals Site-Specific Microheterogeneity of SARS-CoV-2

S Glycosylation

We utilized multiple approaches to examine glycosylation of the SARS-CoV-2 S trimer. First, the

portfolio of glycans linked to SARS-CoV-2 S trimer immunogen was analyzed following their

release from the polypeptide backbone. N-glycans were released from protein by treatment with

PNGase F and O-glycans were subsequently released by beta-elimination. Following

permethylation to enhance detection sensitivity and structural characterization, released glycans

were analyzed by multi-stage mass spectrometry (MS^n) (38,39). Mass spectra were processed

by GRITS Toolbox and the resulting annotations were validated manually (40). Glycan

assignments were grouped by type and by additional structural features for relative quantification

of profile characteristics (**Fig. 2A**, **Supplemental Table, Tab 3**). This analysis quantified 49 N-

glycans and revealed that 55% of the total glycan abundance was of the complex type, 17% was

of the hybrid type, and 28% was high-mannose. Among the complex and hybrid N-glycans, we

observed a high degree of core fucosylation and significant abundance of bisected and LacDiNAc

structures. We also observed sulfated N-linked glycans using negative mode MS^n analyses

(**Supplemental Table, Tab 13**) though signal intensity was too low in positive ion mode (at least

10-fold lower than any of the non-sulfated glycans) for accurate quantification. In addition, we

detected 15 O-glycans released from the S trimer (**Fig. S5, Supplemental Table, Tab 4**).

To determine occupancy of N-linked glycans at each site, we employed a sequential

deglycoslyation approach using Endoglycosidase H and PNGase F in the presence of ^{18}O-H_2O

following tryptic digestion of S (28,41). Following LC-MS/MS analyses, the resulting data

confirmed that 19 of the canonical sequons had occupancies greater than 95% (**Supplemental**

Table, Tab 5). One canonical sequence, N0149, had insufficient spectral counts for quantification

by this method but subsequent analyses described below suggested high occupancy. The 2 most

C-terminal N-linked sites, N1173 and N1194, had reduced occupancy, 52% and 82% respectively.

Reduced occupancy at these sites may reflect hindered en-bloc transfer by the

oligosaccharyltransferase (OST) due to primary amino acid sequences at or near the N-linked

sequon.   Alternatively, this may reflect these two sites being post-translationally modified after

release of the protein by the ribosome by a less efficient STT3B-containing OST, either due to

activity or initial folding of the polypeptide, as opposed to co-translationally modified by the

STT3A-containing OST (42). None of the non-canoncial sequons (3 N-X-C sites and 4 N-G-L/I/V

sites, (43)) showed significant occupancy (>5%) except for N0501 that showed moderate (19%)

conversion to ^{18}O-Asp that could be due to deamidation that is facilitated by glycine at the +1

position (**Supplemental Table, Tab 5,** (44)). Further analysis of this site (see below) by direct

glycopeptide analyses allowed us to determine that N0501 undergoes deamidation but is not

glycosylated. Thus, all, and only the, 22 canonical sequences for N-linked glycosylation (N-X-S/T)

are utilized with only N1173 and N1194 demonstrating occupancies below 95%.

Next, we applied 3 different proteolytic digestion strategies to the SARS-CoV-2 S immunogen to

maximize glycopeptide coverage by subsequent LC-MS/MS analyses.   Extended gradient

nanoflow reverse-phase LC-MS/MS was carried out on a ThermoFisher Lumos™ Tribrid™

instrument using Step-HCD fragmentation on each of the samples (see STAR methods for details,

(25,26,28,41,45)). Following data analyses using pGlyco 2.2.2 (46), Byonic (47), and manual

validation of glycan compositions against our released glycomics findings (**Fig. 2A**,

Supplemental Table, Tab 3 and 13), we were able to determine the microheterogeneity at each

of the 22 canonical sites (**Fig. 2B-2E, Supplemental Table, Tab 6**). Notably, none of the non-

canonical consensus sequences, including N0501, displayed any quantifiable glycans. The N-

glycosites N0074 (**Fig. 2B**) and N0149 (**Fig. 2C**) are highly processed and display a typical

mammalian N-glycan profile. N0149 is, however, modified with several hybrid N-glycan structures

while N0074 is not. N0234 (**Fig. 2D**) and N0801 (**Fig. 2E**) have N-glycan profiles more similar to

those found on other viruses such as HIV (15) that are dominated by high-mannose structures.

N0234 (**Fig. 2D**) displays an abundance of Man7 - Man9 high-mannose structures suggesting

stalled processing by early acting ER and cis-Golgi mannosidases. In contrast, N0801 (**Fig. 2E**)

is processed more efficiently to Man5 high-mannose and hybrid structures suggesting that access

to the glycan at this site by MGAT1 and α-Mannosidase II is hindered. In general, for all 22 sites

(**Fig. 2B-2E, Supplemental Table, Tab 6**), we observed under processing of complex glycan

antennae (i.e. under-galactosylation and under-sialylation) and a high degree of core fucosylation

in agreement with released glycan analyses (**Fig. 2A**, **Supplemental Table, Tab 3**). We also

observed a small percent of sulfated N-linked glycans at several sites (**Supplemental Table, Tab**

6 and 8). Based on the assignments and the spectral counts for each topology, we were able to

determine the percent of total N-linked glycan types (high-mannose, hybrid, or complex) present

at each site (**Figure 3, Supplemental Table, Tab 7**). Notably, 3 of the sites (N0234, N0709, and

N0717) displayed more than 50% high-mannose glycans while 11 other sites (N0017, N0074,

N0149, N0165, N0282, N0331, N0657, N1134, N1158, N1173, and N1194) were more than 90%

complex when occupied. The other 8 sites were distributed between these 2 extremes. Notably,

only 1 site (N0717 at 45%), which also had greater than 50% high-mannose (55%), had greater

than 33% hybrid structures. To further evaluate the heterogeneity, we grouped all the topologies

into the 20 classes recently described by the Crispin laboratory with adding 2 categories (sulfated

and unoccupied) that we refer to here as the Oxford classification (**Supplemental Table, Tab 8,**

(19)). Among other features observed, this classification allowed us to observe that while most

sites with high mannose structures were dominated by the Man5GlcNAc2 structure, N0234 and

N0717 were dominated by the higher Man structures of Man8GlcNAc2 and Man7GlcNAc2,

respectively (**Fig. S7, Supplemental Table, Tab 8**). Limited processing at N0234 is in agreement

with a recent report suggesting that high mannose structures at this site help to stabilize the

receptor-binding domain of S (www.biorxiv.org/content/10.1101/2020.06.11.146522v1).

Furthermore, applying the Oxford classifications to our dataset clearly demonstrates that the 3

most C-terminal sites (N1158, N1173, and N1194), dominated by complex type glycans, were

more often further processed (i.e. multiple antennae) and elaborated (i.e. galactosylation and

sialylation) than other sites (**Supplemental Table, Tab 8**).

We also analyzed our generated mass spectrometry data for the presence of O-linked glycans

based on our glycomic findings (**Fig. S5, Supplemental Table, Tab 4**) and a recent manuscript

suggesting significant levels of O-glycosylation of S1 and S2 when expressed independently (23).

We were able to confirm sites of O-glycan modification with microheterogeneity observed for the

vast majority of these sites (**Supplemental Table, Tab 9**). However, occupancy at each site,

determined by spectral counts, was observed to be very low (below 4%) except for Thr0323 that

had a modestly higher but still low 11% occupancy (**Fig. S6, Supplemental Table, Tab 10**).

3D Structural Modeling of Glycosylated SARS-CoV-2 Trimer Immunogen Enables Predictions of

Epitope Accessibility and Other Key Features

A 3D structure of the S trimer was generated using a homology model of the S trimer described

previously (based on PDB code 6VSB, (48)). Onto this 3D structure, we installed explicitly defined

glycans at each glycosylated sequon based on one of three separate sets of criteria, thereby

generating three different glycoform models for comparison that we denote as "Abundance,"

"Oxford Class," and "Processed" models (**see Methods and Supplemental Table, Tab 1**). These

criteria were chosen in order to generate glycoform models that represent reasonable

expectations for glycosylation microheterogeneity and integrate cross-validating glycomic and

glycoproteomic characterization of S and ACE2.

The three glycoform models were subjected to multiple all atom MD simulations with explicit

water. Information from analyses of these structures is presented in **Figure 4A** along with the

sequence of the SARS-CoV-2 S protomer. We also determined variants in S that are emerging in

the virus that have been sequenced to date (**Supplemental Table, Tab 11**). The inter-residue

distances were measured between the most α-carbon-distal atoms of the N-glycan sites and

Spike glycoprotein population variant sites in 3D space (**Figure 4B**). Notable from this analysis,

there are several variants that don't ablate the N-linked sequon, but that are sufficiently close in

3-dimensional space to N-glycosites, such as D138H, H655Y, S939F, and L1203F, to warrant

further investigation.

The percentage of simulation time that each S protein residue is accessible to a probe that

approximates the size of an antibody variable domain was calculated for a model of the S trimer

using the Abundance glycoforms (**Supplemental Table, Tab 1,** (49)). The predicted antibody

accessibility is visualized across the sequence, as well as mapped onto the 3D surface, via color

shading **(Figure 4A, 4C, Supplemental Table, Tab 13, and Supplemental Movie A)**.

Additionally, the Oxford Class glycoforms model (**Supplemental Table, Tab 1)**, which is arguably

the most encompassing means for representing glycan microheterogeneity since it captures

abundant structural topologies (**Supplemental Table, Tab 8**), is shown with the sequence variant

information (Figure 4D, **Supplemental Table, Tab 11**). A substantial number of these variants

occur (directly by comparison to **Figure 4A** or visually by comparison to **Figure 4C**) in regions of

high calculated epitope accessibility (e.g. N74K, T76I, R78M, D138H, H146Y, S151I, D253G,

V483A, etc., **Supplemental Table, Tab 14**) suggesting potential selective pressure to avoid host

immune response. Also, it is interesting to note that 3 of the emerging variants would eliminate

N-linked sequons in S; N74K and T76I would eliminate N-glycosylation of N74 (found in the insert

variable region 1 of CoV-2 S compared to CoV-1 S), and S151I eliminates N-glycosylation of

N149 (found in the insert variable region 2) (**Fig. 4A, S7, Supplemental Table, Tab 11)**. Lastly,

the SARS-CoV-2 S Processed glycoform model is shown (**Supplemental Table, Tab 1**), along

with marking amino acid T0323 that has a modest (11% occupancy, **Fig. S6, Supplemental**

Table, Tab 10) amount of O-glycosylation to represent the most heavily glycosylated form of S

(**Figure 4E**).

Glycomics Informed Glycoproteomics Reveals Complex N-linked Glycosylation of ACE2

We also analyzed ACE2 glycosylation utilizing the same glycomic and glycoproteomic

approaches described for S protein. Glycomic analyses of released N-linked glycans (**Fig. 5A,**

Supplemental Table, Tab 3) revealed that the majority of glycans on ACE2 are complex with

limited high-mannose and hybrid glycans and we were unable to detect sulfated N-linked glycans.

Glycoproteomic analyses revealed that occupancy was high (>75%) at all 6 sites and significant

microheterogeneity dominated by complex N-glycans was observed for each site (**Fig. 5B-5G,**

Supplemental Table, Tabs 5-8). We also observed, consistent with the O-glycomics (**Fig. S5,**

Supplemental Table, Tab 4), that Ser 155 and several S/T residues at the C-terminus of ACE2

outside of the peptidase domain were O-glycosylated but stoichiometry was extremely low (less

than 2%, **Supplemental Table, Tab 9 and 10**).

3D Structural Modeling of Glycosylated, Soluble, ACE2 Highlighting Glycosylation and Variants.

We integrated our glycomics, glycoproteomics, and population variant analyses results with a 3D

model of Ace 2 (based on PDB code 6M0J (50), see methods for details) to generate two versions

of the soluble glycosylated ACE2 for visualization and molecular dynamics simulations. We

visualized the ACE2 glycoprotein with the Abundance glycoform model simulated at each site as

well as highlighting the naturally occurring variants observed in the human population (**Fig. 6A,**

Supplemental Movie B, Supplemental Table, Tab 11). Note, that the Abundance glycoform

model and the Oxford Class glycoform model for ACE2 are identical (**Supplemental Table, Tabs**

1 and 8). Notably, one site of N-linked glycosylation (N546) is predicted to not be present in 3

out of 10,000 humans based on naturally occurring variation in the human population

(**Supplemental Table, Tab 11**). We also modeled ACE2 using the Processed glycoform model

(**Fig. 6B**). In both models, the interaction domain with S is defined (**Fig. 6A-B, Supplemental**

Movie B) .

Molecular Dynamics Simulation of the Glycosylated Trimer Spike of SARS-CoV-2 in Complex

with Glycosylated, Soluble, Human Ace 2 Reveals Protein and Glycan Interactions

Molecular dynamics simulations were performed to examine the co-complex (generated from a

crystal structure of the ACE2-RBD co-complex, PDB code 6M0J, (50)) of glycosylated S with

glycosylated ACE2 with the 3 different glycoforms models (Abundance, Oxford Class, and

Processed, **Supplemental Table, Tab 1, Supplemental Simulations 1-3**). Information from

these analyses is laid out along the primary structure (sequence) of the SARS-CoV-2 S protomer

and ACE2 highlighting regions of glycan-protein interaction observed in the MD simulations

(**Supplemental Table, Tab 14**, **Supplemental Simulations 1-3**). Interestingly, two glycans on

ACE2 (at N090 and N322), that are highlighted in **Figure 7A** and shown in a more close-up view

in **Figure 7B**, are predicted to form interactions with the S protein (**Supplemental Table, Tab 15**).

The N322 glycan interaction with the S trimer is outside of the receptor binding domain, and the

interaction is observed across multiple simulations and throughout each simulation (**Fig. 7A-B**,

Supplemental Simulations 1-3). The ACE2 glycan at N090 is close enough to the S trimer

surface to repeatedly form interactions, however the glycan arms interact with multiple regions of

the surface over the course of the simulations, reflecting the relatively high degree of glycan

dynamics (**Fig. 7A-B, Supplemental Movie C**). Inter-molecule glycan-glycan interactions are

also observed repeatedly between the glycan at N546 of ACE2 and those in the S protein at

residues N0074 and N0165 (**Fig. 7D, Supplemental Table, Tab 16**). Finally, a full view of the

ACE2-S complex with Oxford class glycoforms on both proteins illustrates the extensive

glycosylation at the interface of the complex (**Fig. 7C, Supplemental Movie D**).

DISCUSSION

We have defined the glycomics-informed, site-specific microheterogeneity of 22 sites of N-linked

glycosylation per monomer on a SARS-CoV-2 trimer and the 6 sites of N-linked glycosylation on

a soluble version of its human ACE2 receptor using a combination of mass spectrometry

approaches coupled with evolutionary and variant sequence analyses to provide a detailed

understanding of the glycosylation states of these glycoproteins (**Figs. 1-6**). Our results suggest

essential roles for glycosylation in mediating receptor binding, antigenic shielding, and potentially

the evolution/divergence of these glycoproteins.

The highly glycosylated SARS-CoV-2 Spike protein, unlike several other viral proteins including

HIV-1 (15) but in agreement with another recent report (19), presents significantly more

processing of N-glycans towards complex glycosylation, suggesting that steric hindrance to

processing enzymes is not a major factor at most sites (**Figs. 2-3**). However, the N-glycans still

provide considerable shielding of the peptide backbone (**Fig. 4**). Our glycomics-guided

glycoproteomic data is in general in strong agreement with the trimer immunogen data recently

published by Crispin (19) though we also observed sulfated N-linked glycans, were able to

differentiate branching, bisected, and diLacNAc containing structures by glycomics, and observed

less occupancy on the 2 most C-terminal N-linked sites using a different approach. Our detection

of sulfated N-linked glycans at multiple sites on S is in agreement with a recent manuscript re-

analyzing the Crispin data   (https://www.biorxiv.org/content/10.1101/2020.05.31.125302v1).

Sulfated N-linked glycans could potentially play key roles in immune regulation and receptor

binding as in other viruses (51). This result is especially significant in that sulfated N-glycans

were not observed when we performed glycomics on ACE2. At each individual site, the glycans

we observed on our immunogen appear to be slightly more processed but the overlap between

our analysis and the Crispin's group results (19) at each site in terms of major features are nearly

superimposable. This agreement differs substantially when comparing our and Crispin's data (19)

to that of the Azadi group (23) that analyzed S1 and S2 that had been expressed individually.

When expressed as 2 separate polypeptides and not purified for trimers, several unoccupied

sites of N-linked glycosylation were observed and processing at several sites was significantly

different (23) than we and others (19) observed. Although O-glycosylation has recently been

reported for individually-expressed S1 and S2 domains of the Spike glycoprotein (23), in trimeric

form the level of O-glycosylation is extremely low, with the highest level of occupancy we observed

being 11% at T0323 (**Fig. 4E**). The low level of O-linked occupancy we observed is in agreement

with Crispin's analysis of a Spike Trimer immunogen (19) but differs significantly from Azadi's

analyses of individually expressed S1 and S2 (23). Thus, the context in which the Spike protein

is expressed and purified before analyses significantly alters the glycosylation of the protomer

that is reminiscent of previous studies looking at expression of the HIV-1 envelope Spike (15,52).

The soluble ACE2 protein examined here contains 6 highly utilized sites of N-linked glycosylation

dominated by complex type N-linked glycans (**Fig. 5**). O-glycans were also present on this

glycoprotein but at very low levels of occupancy at all sites (<2%).

Our glycomics-informed glycoproteomics allowed us to assign defined sets of glycans to specific

glycosylation sites on 3D-structures of S and ACE2 glycoproteins based on experimental

evidence (**Figs. 4, 6**). Similar to almost all glycoproteins, microheterogeneity is evident at most

glycosylation sites of S and ACE2; each glycosylation site can be modified with one of several

glycan structures, generating site-specific glycosylation portfolios. For modeling purposes,

however, explicit structures must be placed at each glycosylation site. In order to capture the

impact of microheterogenity on S and ACE2 molecular dynamics we chose to generate

glycoforms for modeling that represented reasonable portfolios of glycan types. Using 3

glycoform models for S (Abundance, Oxford Class, and Processed) and 2 models for ACE2

(Abundance, which was equivalent to Oxford Class, and Processed), we generated 3 molecular

dynamics simulations of the co-complexes of these 2 glycoproteins (**Fig. 7 and Supplemental**

Simulations 1-3). The observed interactions over time allowed us to evaluate glycan-protein

contacts between the 2 proteins as well as examine potential glycan-glycan interactions (**Fig. 7**).

We observed glycan-mediated interactions between the S trimer and glycans at N090, N322 and

N546 of ACE2. Thus, variations in glycan occupancy or processing at these sites, could alter the

affinity of the SARS-CoV-2 – ACE2 interaction and modulate infectivity. It is well established that

glycosylation states vary depending on tissue and cell type as well as in the case of humans, on

age (53), underlying disease (54,55) and ethnicity (56). Thus glycosylation portfolios may in part

be responsible for tissue tropism and individual susceptibility to infection. The importance of

glycosylation for S binding to ACE2 is even more emphatically demonstrated by the direct glycan-

glycan interactions observed (**Fig. 7**) between S glycans (at N0074 and N0165) and an ACE2

receptor glycan (at N546), adding an additional layer of complexity for interpreting the impact of

glycosylation on individual susceptibility.

Several emerging variants of the virus appear to be altering N-linked glycosylation occupancy by

disrupting N-linked sequons. Interestingly, the 2 N-linked sequons in SARS-CoV-2 S directly

impacted by variants, N0074 and N0149, are in divergent insert regions 1 and 2, respectively, of

SARS-CoV-2 S compared to SARS-CoV-1 S (**Fig. 4A**). The N0074, in particular, is one of the S

glycans that interact directly with ACE2 glycan (at N546, **Fig. 7**), suggesting that glycan-glycan

interactions may contribute to the unique infectivity differences between SARS-CoV-2 and SARS-

CoV-1. These sequon variants will also be important to examine in terms of glycan shielding that

could influence immunogenicity and efficacy of neutralizing antibodies, as well as interactions with

the host cell receptor ACE2.   Naturally-occurring amino acid-changing SNPs in the ACE2 gene

generate a number of variants including 1 variant, with a frequency of 3 in 10,000 humans, that

eliminates a site of N-linked glycosylation at N546 (**Fig. 6**). Understanding the impact of ACE2

variants on glycosylation and more importantly on S binding, especially for N546S which impacts

the glycan-glycan interaction between S and ACE2 (**Fig. 7**), should be prioritized in light of efforts

to develop ACE2 as a potential decoy therapeutic. Intelligent manipulation of ACE2 glycosylation

may lead to more potent biologics capable of acting as better competitive inhibitors of S binding.

The data presented here, and related similar recent findings (19,57,58), provide a framework to

facilitate the production of immunogens, vaccines, antibodies, and inhibitors as well as providing

additional information regarding mechanisms by which glycan microheterogeneity is achieved.

However, considerable efforts still remain in order to fully understand the role of glycans in SARS-

CoV-2 infection and pathogenicity. While HEK-expressed S and ACE2 provide a useful window

for understanding human glycosylation of these proteins, glycoproteomic characterization

following expression in cell lines of more direct relevance to disease and target tissue is sorely

needed. While site occupancy may change depending on presentation and cell type (59),

processing of N-linked glycans will almost certainly be altered in a cell-type dependent fashion.

Thus, analyses of the Spike trimer extracted from pseudoviruses, virion-like particles, and

ultimately from infectious SARS-CoV-2 virions harvested from airway cells or patients will provide

the most accurate view of how trimer immunogens reflect the true glycosylation pattern of the

virus. Detailed analyses of the impact of emerging variants in S and natural and designed-for-

biologics variants of ACE2 on glycosylation and binding properties are important next steps for

developing therapeutics. Finally, it will be important to monitor the slow evolution of the virus to

determine if existing sites of glycosylation are lost or new sites emerge with selective pressure

that might alter the efficacy of vaccines, neutralizing antibodies, and/or inhibitors.

ACKNOWLEDGEMENTS

The authors would like to thank Protein Metrics for providing licenses for their software used here

and the developers of pGlyco for productive discussions regarding their software. We would also

like to thank Galit Alter of the Ragon Institute for facilitating this collaborative effort. This effort

was facilitated by the ThermoFisher Scientific appointed Center of Excellence in Glycoproteomics

at the Complex Carbohydrate Research Center at the University of Georgia (co-directed by MT

and LW). This research is supported in part by R35GM119850 (NEL), NNF10CC1016517 (NEL),

R01AI139238 (MAB), R01AI147884-01A1S1 (BC), Massachusetts Consortium on Pathogen

Readiness (BC), U01CA207824 (R.J.W), P41GM103390 (R.J.W.), P41GM103490 (MT and LW),

U01GM125267 (MT), and R01GM130915 (LW). The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

Conceptualization: M.T., B.C., R.J.W. and L.W.; Methodology, Software, Validation, Formal

Analysis, Investigation, Resources, and Data Curation: P.Z., J.L.P., O.C.G., Y.C., T.X., K.E.R.,

K.A., B.P.K., R.B., D.H.B., M.A.B., N.E.L., M.T., B.C., R.J.W., and L.W.; Writing-Original Draft:

P.Z., J.L.P., and L.W.; Writing-Review & Editing: All authors; Visualization: P.Z., J.L.P., O.C.G,

Y.C., M.T., B.C., R.J.W., and L.W.; Supervision, Project Administration, and Funding Acquisition:

D.H.B., M.A.B., N.E.L., M.T., B.C., R.J.W., and L.W..

DECLARATION OF INTERESTS

The authors declare no competing interests.

450
451   **FIGURE LEGENDS**
452

Figure 1. Expression and Characterization of SARS-CoV-2 Spike Glycoprotein Trimer

Immunogen and Soluble Human ACE2. A) Sequences of SARS-CoV-2 S immunogen and

soluble human ACE2. The N-terminal pyroglutamines for both mature protein monomers are

bolded, underlined, and shown in green. The canonical N-linked glycosylation sequons are

bolded, underlined, and shown in red. Negative stain electron microscopy of the purified trimer

(B) and Coomassie G-250 stained reducing SDS-PAGE gels (C) confirmed purity of the SARS-

CoV-2 S protein trimer and of the soluble human ACE2. MWM = molecular weight markers. D)

A representative Step-HCD fragmentation spectrum from mass spectrometry analysis of a tryptic

digest of S annotated manually based on search results from pGlyco 2.2. This spectrum defines

the N-terminus of the mature protein monomer as (pyro-)glutamine 0014. A representative N-

glycan consistent with this annotation and our glycomics data (Fig. 2) is overlaid using the Symbol

Nomenclature For Glycans (SNFG) code. This complex glycan occurs at N0017. Note, that as

expected, the cysteine is carbamidomethylated and the mass accuracy of the assigned peptide

is 0.98 ppm. On the sequence of the N-terminal peptide and in the spectrum, the assigned b (blue)

and y (red) ions are shown. In the spectrum, purple highlights glycan oxonium ions and green

marks intact peptide fragment ions with various partial glycan sequences still attached. Note that

the green-labeled ions allow for limited topology to be extracted including defining that the fucose

is on the core and not the antennae of the glycopeptide.

Figure 2. Glycomics Informed Glycoproteomics Reveals Substantial Site-Specific

Microheterogeneity of N-linked Glycosylation on SARS-CoV-2 S. A) Glycans released from

SARS-CoV-2 S protein trimer immunogen were permethylated and analyzed by MSn. Structures

were assigned, grouped by type and structural features, and prevalence was determined based

on ion current. The pie chart shows basic division by broad N-glycan type. The bar graph provides

additional detail about the glycans detected. The most abundant structure with a unique

categorization by glycomics for each N-glycan type in the pie chart, or above each feature

category in the bar graph, is indicated. B – E) Glycopeptides were prepared from SARS-CoV-2

S protein trimer immunogen using multiple combinations of proteases, analyzed by LC-MSn, and

the resulting data was searched using several different software packages. Four representative

sites of N-linked glycosylation with specific features of interest were chosen and are presented

here. N0074 (B) and N0149 (C) are shown that occur in variable insert regions of S compared to

SARS-CoV and other related coronaviruses, and there are emerging variants of SARS-CoV-2

that disrupt these two sites of glycosylation in S. N0234 (D) contains the most high-mannose N-

linked glycans. N0801 (D) is an example of glycosylation in the S2 region of the immunogen and

displays a high degree of hybrid glycosylation compared to other sites. The abundance of each

composition is graphed in terms of assigned spectral counts. Representative glycans (as

determined by glycomics analysis) for several abundant compositions are shown in SNFG format.

The abbreviations used here and throughout the manuscript are N for HexNAc, H for Hexose, F

for Fucose, A for Neu5Ac, and S for Sulfation. Note that the graphs for the other 18 sites and

other graphs grouping the microheterogeneity observed by other properties are presented in

Supplemental Information.

Figure 3. SARS-CoV-2 S Immunogen N-glycan Sites are Predominantly Modified by

Complex N-glycans. N-glycan topologies were assigned to all 22 sites of the S protomer and

the spectral counts for each of the 3 types of N-glycans (high-mannose, hybrid, and complex) as

well as the unoccupied peptide spectral match counts at each site were summed and visualized

as pie charts. Note that only N1173 and N1194 show an appreciable amount of the unoccupied

amino acid.

Figure 4. 3D Structural Modeling of Glycosylated SARS-CoV-2 Spike Trimer Immunogen

Reveals Predictions for Antigen Accessibility and Other Key Features. Results from

glycomics and glycoproteomics experiments were combined with results from bioinformatics

analyses and used to model several versions of glycosylated SARS-CoV-2 S trimer immunogen.

A) Sequence of the SARS-CoV-2 S immunogen displaying computed antigen accessibility and

other information. Antigen accessibility is indicated by red shading across the amino acid

sequence. B) Emerging variants confirmed by independent sequencing experiments were

analyzed based on the 3D structure of SARS-CoV-2 S to generate a proximity chart to the

determined N-linked glycosylation sites. C) SARS-CoV-2 S trimer immunogen model from MD

simulation displaying abundance glycoforms and antigen accessibility shaded in red for most

accessible, white for partial, and black for inaccessible (see **Supplemental movie A**). D) SARS-

CoV-2 S trimer immunogen model from MD simulation displaying oxford class glycoforms and

sequence variants. * indicates not visible while the box represents 3 amino acid variants that are

clustered together in 3D space. E) SARS-CoV-2 S trimer immunogen model from MD simulation

displaying processed glycoforms plus shading of Thr-323 that has O-glycoslyation at low

stoichiometry in yellow.

Figure 5: Glycomics Informed Glycoproteomics of Soluble Human ACE2 Reveals High

Occupancy, Complex N-linked Glycosylation. A) Glycans released from soluble, purified

ACE2 were permethylated and analyzed by MSn. Structures were assigned, grouped by type and

structural features, and prevalence was determined based on ion current. The pie chart shows

basic division by broad N-glycan type. The bar graph provides additional detail about the glycans

detected. The most abundant structure with a unique categorization by glycomics for each N-

glycan type in the pie chart, or above each feature category in the bar graph, is indicated. B – G)

Glycopeptides were prepared from soluble human ACE2 using multiple combinations of

proteases, analyzed by LC-MSn, and the resulting data was searched using several different

software packages. All six sites of N-linked glycosylation are presented here. Displayed in the bar

graphs are the individual compositions observed graphed in terms of assigned spectral counts.

Representative glycans (as determined by glycomics analysis) for several abundant compositions

are shown in SNFG format. The abbreviations used here and throughout the manuscript are N

for HexNAc, H for Hexose, F for Fucose, and A for Neu5Ac. The pie chart (analogous to Figure

3 for SARS-CoV-2 S) for each site is displayed in the upper corner of each panel. B) N053. C)

N090. D) N103. E) N322. F) N432. G) N546, a site that does not exist in 3 in 10,000 people.

Figure 6: 3D Structural Modeling of Glycosylated Soluble Human ACE2. Results from
glycomics and glycoproteomics experiments were combined with results from bioinformatics
analyses and used to model several versions of glycosylated soluble human ACE2. A) Soluble
human ACE2 model from MD simulations displaying abundance glycoforms, interaction surface
with S, and sequence variants. N546 variant is boxed that would remove N-linked glycosylation
at that site (see **Supplemental movie B**). B) Soluble human ACE2 model from MD simulations
displaying processed glycoforms and interaction surface with S.

Figure 7: Interactions of Glycosylated Soluble Human ACE2 and Glycosylated SARS-CoV-
2 S Trimer Immunogen Revealed By 3D-Structural Modeling and Molecular Dynamics
Simulations. A) Molecular dynamics simulation of glycosylated soluble human ACE2 and
glycosylated SARS-CoV-2 S trimer immunogen interaction (see **Supplemental simulations 1-**
3). ACE2 (top) is colored red with glycans in pink while S is colored white with glycans in dark
grey. Highlighted are ACE2 glycans that interacts with S that are zoomed in on to the right. B)
Zoom in of ACE2-S interface highlighting ACE2 glycan interactions using 3D-SNFG icons (60)
with S protein (pink) as well as ACE2-S glycan-glycan interactions. C) Zoom in of dynamics
trajectory of glycans at the interface of soluble human ACE2 and S (see **Supplemental movies**
C and D).

STAR METHODS
KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, Peptides, and Recombinant Proteins		
SARS-CoV-2 S protein	This Study	N/A
Human ACE2 protein	This Study	N/A
2x Laemmli sample buffer	Bio-Rad	Cat#161-0737
Invitrogen NuPAGE 4 to 12%, Bis-Tris, Mini Protein Gel	Thermo Fisher Scientific	Cat#NP0321PK2
Coomassie Brilliant Blue G-250 Dye	Thermo Fisher Scientific	Cat#20279
Dithiothreitol	Sigma Aldrich	Cat#43815
Iodoacetamide	Sigma Aldrich	Cat#I1149

Trypsin	Promega	Cat#V5111
Lys-C	Promega	Cat#V1671
Arg-C	Promega	Cat#V1881
Glu-C	Promega	Cat#V1651
Asp-N	Promega	Cat#VA1160
Endoglycosidase H	Promega	Cat#V4871
PNGaseF	Promega	Cat#V4831
Chymotrypsin	Athens Research and Technology	Cat#16-19-030820
Alpha lytic protease	New England BioLabs	Cat#P8113
^{18}O water	Cambridge Isotope Laboratories	OLM-782-10-1
O-protease OpeRATOR	Genovis	Cat#G1-OP1-020
Deposited Data		
MS data for site-specific N-linked glycopeptides for SARS-Cov-2 S and human ACE2	This Study	PXD019937
MS data for site-specific O-linked glycopeptides for SARS-Cov-2 S and human ACE2	This Study	PXD019940
MS data for deglycosylated N-linked glycopeptides for SARS-Cov-2 S and human ACE2	This Study	PXD019938
MS data for disulfide bond analysis for SARS-Cov-2 S	This Study	PXD019939
MS data for N-linked glycomics deposited at GlycoPost	This Study	GPST000120
MS data for O-linked glycomics deposited at GlycoPost	This Study	GPST000121
Experimental Models: Cell Lines		
293-F Cells	Gibco	Cat#R79007
Experimental Models: Organisms/Strains		
VSV(G)-Pseudoviruses	This Study	
Software and Algorithms		
pGlyco	(43)	v2.2.2
Proteome Discoverer	Thermo Fisher Scientific	v1.4
Byonic	Protein Metrics Inc. (44)	v3.8.13
ProteoIQ	Premier Biosoft	v2.7
GRITS Toolbox	http://www.grits-toolbox.org (40)	v1.1
EMBOSS needle	(61)	v6.6.0
Biopython	https://biopython.org/	v1.76
Rpdb	https://rdrr.io/cran/Rpdb/	v2.3
SignalP	http://www.cbs.dtu.dk/services/SignalP/ (34)	v5.0
LibreOFFICE Writer	The Document Foundation	V6.4.4.2
GlyGen	https://www.glygen.org	V1.5

GNOme	https://github.com/glyg en-glycan- data/GNOme/blob/ma ster/README.md	V1.5.5
GlyTouCan	https://glytoucan.org	V3.1.0
Inkscape	Inkscape Developers	V1.0

LEAD CONTACT

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Peng Zhao (pengzhao@uga.edu) and/or Lance Wells

(lwells@ccrc.uga.edu).

METHOD DETAILS

Expression, Purification, and Characterization of SARS-CoV-2 S and Human ACE2

Proteins

To express a stabilized ectodomain of Spike protein, a synthetic gene encoding residues

1−1208 of SARS-CoV-2 Spike with the furin cleavage site (residues 682–685) replaced by a

"GGSG" sequence, proline substitutions at residues 986 and 987, and a foldon trimerization

motif followed by a C-terminal 6xHisTag was created and cloned into the mammalian

expression vector pCMV-IRES-puro (Codex BioSolutions, Inc, Gaithersburg, MD). The

expression construct was transiently transfected in HEK 293F cells using polyethylenimine

(Polysciences, Inc, Warrington, PA). Protein was purified from cell supernatants using Ni-NTA

resin (Qiagen, Germany), the eluted fractions containing S protein were pooled, concentrated,

and further purified by gel filtration chromatography on a Superose 6 column (GE Healthcare).

Negative stain electron microscopy (EM) analysis was performed as described (62). Briefly,

analysis was performed at room temperature with a magnification of 52,000x and a defocus

value of 1.5 μm following low-dose procedures, using a Philips Tecnai F20 electron microscope

(Thermo Fisher Scientific) equipped with a Gatan US4000 CCD camera and operated at voltage

of 200 kV. The DNA fragment encoding human ACE2 (1-615) with a 6xHis tag at C terminus

was synthesized by Genscript and cloned to the vector pCMV-IRES-puro. The expression

construct was transfected in HEK293F cells using polyethylenimine. The medium was discarded

and replaced with FreeStyle 293 medium after 6-8 hours. After incubation in 37 °C with 5.5%

CO_2 for 5 days, the supernatant was collected and loaded to Ni-NTA resin for purification. The

elution was concentrated and further purified by a Superdex 200 column.

In-Gel Analysis of SARS-CoV-2 S and Human ACE2 Proteins

A 3.5-µg aliquot of SARS-CoV-2 S protein as well as a 2-µg aliquot of human ACE2 were

combined with Laemmli sample buffer, analyzed on a 4-12% Invitrogen NuPage Bis-Tris gel

using the MES pH 6.5 running buffer, and stained with Coomassie Brilliant Blue G-250.

Analysis of N-linked and O-linked Glycans Released from SARS-Cov-2 S and Human

ACE2 Proteins

Aliquots of approximately 25-50 µg of S or ACE2 protein were processed for glycan analysis as

previously described (38,39). For N-linked glycan analysis, the proteins were digested with

trypsin. Following trypsinization, glycopeptides were enriched by C18 Sep-Pak and subjected to

PNGaseF digestion to release N-linked glycans. Following PNGaseF digestion, released

glycans were separated from residual glycosylated peptides bearing O-linked glycans by C18

Sep-Pak. O-glycosylated peptides were eluted from the Sep-Pak and subjected to reductive β-

elimination to release the O-glycans. Another 25-50 µg aliquot of each protein was denatured

with SDS and digested with PNGaseF to remove N-linked glycans. The de-N-glycosylated,

intact protein was precipitated with cold ethanol and then subjected to reductive β-elimination to

release O-glycans. The profiles of O-glycans released from peptides or from intact protein were

found to be comparable. N- and O-linked glycans released from glycoproteins were

permethylated with methyliodide according to the method of Anumula and Taylor prior to MS

analysis (63). Glycan structural analysis was performed using an LTQ-Orbitrap instrument

(Orbitrap Discovery, ThermoFisher). Detection and relative quantification of the prevalence of

individual glycans was accomplished using the total ion mapping (TIM) and neutral loss scan

(NL scan) functionality of the Xcalibur software package version 2.0 (Thermo Fisher Scientific)

as previously described (38,39). Mass accuracy and detector response was tuned with a

permethylated oligosaccharide standard in positive ion mode. For fragmentation by collision-

induced dissociation (CID in MS^2 and MSn), normalized collision energy of 45% was applied.

Most permethylated glycans were identified as singly or doubly charged, sodiated species in

positive mode. Sulfated N-glycans were detected as singly or doubly charged, deprotonated

species in negative ion mode. Peaks for all charge states were deconvoluted by the charge

state and summed for quantification. All spectra were manually interpreted and annotated. The

explicit identities of individual monosaccharide residues have been assigned based on known

human biosynthetic pathways. Graphical representations of monosaccharide residues are

consistent with the Symbol Nomenclature for Glycans (SNFG), which has been broadly adopted

by the glycomics community (64). The MS-based glycomics data generated in these analyses

and the associated annotations are presented in accordance with the MIRAGE standards and

the Athens Guidelines (65). Data annotation and assignment of glycan accession identifiers

were facilitated by GRITS Toolbox, GlyTouCan, GNOme, and GlyGen (40,66-68).

Analysis of Disulfide Bonds for SARS-Cov-2 S Protein by LC-MS

Two 10-µg aliquots of SARS-CoV-2 S protein were denatured by incubating with 20%

acetonitrile at room temperature and alkylated by 13.75 mM of iodoacetamide at room

temperature in dark. The two aliquots of proteins were then digested respectively using alpha

lytic protease, or a combination of trypsin, Lys-C and Glu-C. Following digestion, the proteins

were deglycosylated by PNGaseF treatment. The resulting peptides were separated on an

Acclaim PepMap RSLC C18 column (75 µm x 15 cm) and eluted into the nano-electrospray ion

source of an Orbitrap Fusion™ Lumos™ Tribrid™ mass spectrometer at a flow rate of 200

nL/min. The elution gradient consists of 1-40% acetonitrile in 0.1% formic acid over 370 minutes

followed by 10 minutes of 80% acetonitrile in 0.1% formic acid. The spray voltage was set to 2.2

kV and the temperature of the heated capillary was set to 280 °C. Full MS scans were acquired

from m/z 200 to 2000 at 60k resolution, and MS/MS scans following electron transfer

dissociation (ETD) were collected in the Orbitrap at 15k resolution. The raw spectra were

analyzed by Byonic (v3.8.13, Protein Metrics Inc.) with mass tolerance set as 20 ppm for both

precursors and fragments. The search output was filtered at 0.1% false discovery rate and 10

ppm mass error. The spectra assigned as cross-linked peptides were manually evaluated for

Cys0015 and Cys0136.

Analysis of Site-Specific N-linked Glycopeptides for SARS-Cov-2 S and Human ACE2

Proteins by LC-MS

Four 3.5-µg aliquots of SARS-CoV-2 S protein were reduced by incubating with 10 mM of

dithiothreitol at 56 °C and alkylated by 27.5 mM of iodoacetamide at room temperature in dark.

The four aliquots of proteins were then digested respectively using alpha lytic protease,

chymotrypsin, a combination of trypsin and Glu-C, or a combination of Glu-C and AspN. Three

10-µg aliquots of ACE2 protein were reduced by incubating with 5 mM of dithiothreitol at 56 °C

and alkylated by 13.75 mM of iodoacetamide at room temperature in dark. The three aliquots of

proteins were then digested respectively using alpha lytic protease, chymotrypsin, or a

combination of trypsin and Lys-C. The resulting peptides were separated on an Acclaim

PepMap RSLC C18 column (75 µm x 15 cm) and eluted into the nano-electrospray ion source

of an Orbitrap Fusion™ Lumos™ Tribrid™ mass spectrometer at a flow rate of 200 nL/min. The

elution gradient consists of 1-40% acetonitrile in 0.1% formic acid over 370 minutes followed by

10 minutes of 80% acetonitrile in 0.1% formic acid. The spray voltage was set to 2.2 kV and the

temperature of the heated capillary was set to 280 °C. Full MS scans were acquired from m/z

200 to 2000 at 60k resolution, and MS/MS scans following higher-energy collisional dissociation

(HCD) with stepped collision energy (15%, 25%, 35%) were collected in the Orbitrap at 15k

resolution. pGlyco v2.2.2 (43) was used for database searches with mass tolerance set as 20

ppm for both precursors and fragments. The database search output was filtered to reach a 1%

false discovery rate for glycans and 10% for peptides. Quantitation was performed by

calculating spectral counts for each glycan composition at each site. Any N-linked glycan

compositions identified by only one spectra were removed from quantitation. N-linked glycan

compositions were categorized into 22 classes (including Unoccupied):

HexNAc(2)Hex(9~5)Fuc(0~1) was classified as M9 to M5 respectively;

HexNAc(2)Hex(4~1)Fuc(0~1) was classified as M1-M4;

HexNAc(3~6)Hex(5~9)Fuc(0)NeuAc(0~1) was classified as Hybrid with

HexNAc(3~6)Hex(5~9)Fuc(1~2)NeuAc(0~1) classified as F-Hybrid; Complex-type glycans are

classified based on the number of antenna, fucosylation, and sulfation:

HexNAc(3)Hex(3~4)Fuc(0)NeuAc(0~1) is assigned as A1 with

HexNAc(3)Hex(3~4)Fuc(1~2)NeuAc(0~1) assigned as F-A1;

HexNAc(4)Hex(3~5)Fuc(0)NeuAc(0~2) is assigned as A2/A1B with

HexNAc(4)Hex(3~5)Fuc(1~5)NeuAc(0~2) assigned as F-A2/A1B;

HexNAc(5)Hex(3~6)Fuc(0)NeuAc(0~3) is assigned as A3/A2B with

HexNAc(5)Hex(3~6)Fuc(1~3)NeuAc(0~3) assigned as F-A3/A2B;

HexNAc(6)Hex(3~7)Fuc(0)NeuAc(0~4) is assigned as A4/A3B with

HexNAc(6)Hex(3~7)Fuc(1~3)NeuAc(0~4) assigned as F-A4/A3B;

HexNAc(7)Hex(3~8)Fuc(0)NeuAc(0~1) is assigned as A5/A4B with

HexNAc(7)Hex(3~8)Fuc(1~3)NeuAc(0~1) as F-A5/A4B; HexNAc(8)Hex(3~9)Fuc(0) is assigned

as A6/A5B with HexNAc(8)Hex(3~9)Fuc(1) assigned as F-A6/A5B; any glycans identified with a

sulfate are assigned as Sulfated.

Analysis of Deglycosylated SARS-Cov-2 S and Human ACE2 Proteins by LC-MS

Three 3.5-µg aliquots of SARS-CoV-2 S protein were reduced by incubating with 10 mM of

dithiothreitol at 56 °C and alkylated by 27.5 mM of iodoacetamide at room temperature in dark.

The three aliquots were then digested respectively using chymotrypsin, Asp-N, or a combination

of trypsin and Glu-C. Two 10-µg aliquots of ACE2 protein were reduced by incubating with 5

mM of dithiothreitol at 56 °C and alkylated by 13.75 mM of iodoacetamide at room temperature

in dark. The two aliquots were then digested respectively using chymotrypsin, or a combination

of trypsin and Lys-C. Following digestion, the proteins were deglycosylated by Endoglycosidase

H followed by PNGaseF treatment in the presence of ^{18}O water. The resulting peptides were

separated on an Acclaim PepMap RSLC C18 column (75 μm x 15 cm) and eluted into the nano-

electrospray ion source of an Orbitrap Fusion™ Lumos™ Tribrid™ mass spectrometer at a flow

rate of 200 nL/min. The elution gradient consists of 1-40% acetonitrile in 0.1% formic acid over

370 minutes followed by 10 minutes of 80% acetonitrile in 0.1% formic acid. The spray voltage

was set to 2.2 kV and the temperature of the heated capillary was set to 280 °C. Full MS scans

were acquired from m/z 200 to 2000 at 60k resolution, and MS/MS scans following collision-

induced dissociation (CID) at 38% collision energy were collected in the ion trap. The spectra

were analyzed using SEQUEST (Proteome Discoverer 1.4) with mass tolerance set as 20 ppm

for precursors and 0.5 Da for fragments. The search output was filtered using ProteoIQ (v2.7) to

reach a 1% false discovery rate at protein level and 10% at peptide level. Occupancy of each N-

linked glycosylation site was calculated using spectral counts assigned to the ^{18}O-Asp-

containing (PNGaseF-cleaved) and/or HexNAc-modified (EndoH-cleaved) peptides and their

unmodified counterparts.

Analysis of Site-Specific O-linked Glycopeptides for SARS-Cov-2 S and Human ACE2

Proteins by LC-MS

Three 10-μg aliquots of SARS-CoV-2 S protein and one 10-μg aliquot of ACE2 protein were

reduced by incubating with 5 mM of dithiothreitol at 56 °C and alkylated by 13.75 mM of

iodoacetamide at room temperature in dark. The four aliquots were then digested respectively

using trypsin, Lys-C, Arg-C, or a combination of trypsin and Lys-C. Following digestion, the

proteins were deglycosylated by PNGaseF treatment and then digested with O-protease

OpeRATOR®. The resulting peptides were separated on an Acclaim PepMap RSLC C18

column (75 μm x 15 cm) and eluted into the nano-electrospray ion source of an Orbitrap

Fusion™ Lumos™ Tribrid™ mass spectrometer at a flow rate of 200 nL/min. The elution

gradient consists of 1-40% acetonitrile in 0.1% formic acid over 370 minutes followed by 10

minutes of 80% acetonitrile in 0.1% formic acid. The spray voltage was set to 2.2 kV and the

temperature of the heated capillary was set to 280 °C. Full MS scans were acquired from m/z

200 to 2000 at 60k resolution, and MS/MS scans following higher-energy collisional dissociation

(HCD) with stepped collision energy (15%, 25%, 35%) or electron transfer dissociation (ETD)

were collected in the Orbitrap at 15k resolution. The raw spectra were analyzed by Byonic

(v3.8.13) with mass tolerance set as 20 ppm for both precursors and fragments. MS/MS filtering

was applied to only allow for spectra where the oxonium ions of HexNAc were observed. The

search output was filtered at 0.1% false discovery rate and 10 ppm mass error. The spectra

assigned as O-linked glycopeptides were manually evaluated. Quantitation was performed by

calculating spectral counts for each glycan composition at each site. Any O-linked glycan

compositions identified by only one spectra were removed from quantitation. Occupancy of each

O-linked glycosylation site was calculated using spectral counts assigned to any glycosylated

peptides and their unmodified counterparts from searches without MS/MS filtering.

Sequence Analysis of SARS-CoV-2 S and Human ACE2 Proteins

The genomes of SARS-CoV as well as bat and pangolin coronavirus sequences reported to be

closely related to SARS-CoV-2 were downloaded from NCBI. The S protein sequences from all

of those genomes were aligned using EMBOSS needle v6.6.0 (61) via the EMBL-EBI provided

web service (69). Manual analysis was performed in the regions containing canonical N-

glycosylation sequons (N-X-S/T). For further sequence analysis of SARS-CoV-2 S variants, the

genomes of SARS-CoV-2 were downloaded from NCBI and GISAID and further processed

using Biopython 1.76 to extract all sequences annotated as "surface glycoprotein" and to

remove any incomplete sequence as well as any sequence containing unassigned amino acids.

For sequence analysis of human ACE2 variants, the single nucleotide polymorphisms (SNPs) of

ACE2 were extracted from the NCBI dbSNP database and filtered for missense mutation entries

with a reported minor allele frequency. Manual analysis was performed on both SARS-CoV-2 S

and human ACE2 variants to further examine the regions containing canonical N-glycosylation

sequons (N-X-S/T). LibreOffice Writer was used to shade regions on the linear sequence of S

and ACE2.

3D Structural Modeling and Molecular Dynamics Simulation of Glycosylated SARS-CoV-2

S and Human ACE2 Proteins

SARS-CoV2 Spike (S) protein structure and ACE2 co-complex – A 3D structure of the prefusion

form of the S protein (RefSeq: YP_009724390.1, UniProt: P0DTC2 SPIKE_SARS2), based on a

Cryo-EM structure (PDB code 6VSB) (48), was obtained from the SWISS-MODEL server

(swissmodel.expasy.org). The model has 95% coverage (residues 27 to 1146) of the S protein.

The receptor binding domain (RBD) in the "open" conformation was replaced with the RBD from

an ACE2 co-complex (PDB code 6M0J) by grafting residues C336 to V524.

Glycoform generation – Glycans (detected by glycomics) were selected for installation on

glycosylated S and ACE2 sequons (detected by glycoproteomics) based on three sets of criteria

designed to reasonably capture different aspects of glycosylation microheterogeneity. We

denote the first of these glycoform models as "Abundance." The glycans selected for

installation to generate the Abundance model were chosen because they were identified as the

most abundant glycan structure (detected by glycomics) that matched the most abundant glycan

composition (detected by glycoproteomics) at each individual site. We denote the second

glycoform model as "Oxford Class." The glycans selected for installation to generate the Oxford

Class model were chosen because they were the most abundant glycan structure, (detected by

glycomics) that was contained within the most highly represented Oxford classification group

(detected by glycoproteomics) at each individual site (**Fig. S7, Supplemental Table, Tabs 1,8).**

Finally, we denote the third glycoform model as "Processed." The glycans selected for

installation to generate the Processed model were chosen because they were the most highly

trimmed, elaborated, or terminally decorated structure (detected by glycomics) that

corresponded to a composition (detected by glycoproteomics) which was present at $\geq 1/3^{rd}$ of

the abundance of the most highly represented composition at each site (**Supplemental Table,**

Tab 1). 3D structures of the three glycoforms (Abundance, Oxford Class, Processed) were

generated for the SARS-CoV2 S protein alone, and in complex with the glycosylated ACE2

protein. The glycoprotein builder available at GLYCAM-Web (www.glycam.org) was employed

together with an in-house program that adjusts the asparagine side chain torsion angles and

glycosidic linkages within known low-energy ranges (70) to relieve any atomic overlaps with the

core protein, as described previously (71,72).

Energy minimization and Molecular dynamics (MD) simulations – Each glycosylated structure

was placed in a periodic box of TIP3P water molecules with a 10 Å buffer between the solute

and the box edge. Energy minimization of all atoms was performed for 20,000 steps (10,000

steepest decent, followed by 10,000 conjugant gradient) under constant pressure (1 atm) and

temperature (300 K) conditions. All MD simulations were performed under nPT conditions with

the CUDA implementation of the PMEMD (73,74) simulation code, as present in the Amber14

software suite (University of California, San Diego). The GLYCAM06j force field (75) and

Amber14SB force field (76) were employed for the carbohydrate and protein moieties,

respectively. A Berendsen barostat with a time constant of 1 ps was employed for pressure

regulation, while a Langevin thermostat with a collision frequency of 2 ps^{-1} was employed for

temperature regulation. A nonbonded interaction cut-off of 8 Å was employed. Long-range

electrostatics were treated with the particle-mesh Ewald (PME) method (77). Covalent bonds

involving hydrogen were constrained with the SHAKE algorithm, allowing an integration time

step of 2 fs to be employed. The energy minimized coordinates were equilibrated at 300K over

400 ps with restraints on the solute heavy atoms. Each system was then equilibrated with

restraints on the Cα atoms of the protein for 1ns, prior to initiating 4 independent 250 ns

production MD simulations with random starting seeds for a total time of 1 μs per system, with

no restraints applied.

Antigenic surface analysis. A series of 3D structure snapshots of the simulation were taken at 1

ns intervals and analysed in terms of their ability to interact with a spherical probe based on the

average size of hypervariable loops present in an antibody complementarity determining region

(CDR), as described recently (https://www.biorxiv.org/content/10.1101/2020.04.07.030445v2).

The percentage of simulation time each residue was exposed to the AbASA probe was

calculated and plotted onto both the 3D structure and primary sequence.

793
Analysis of SARS-CoV-2 Spike VSV pseudoparticles (ppVSV-SARS-2-S)

293T cells were transfected with an expression plasmid encoding SARS-CoV-2 Spike

(pcDNAintron-SARS-2-SΔ19). To increase cell surface expression, the last 19 amino acids

containing the Golgi retention signal were removed. Two SΔ19 constructs were compared, one

started with Met1 and the other with Met2. Twenty-four hours following transfection, cells were

transduced with ppVSVΔG-VSV-G (particles that were pseudotyped with VSV-G in trans). One

hour following transduction cells were extensively washed and media was replaced.

Supernatant containing particles were collected 12-24 hour following transduction and cleared

through centrifugation. Cleared supernatant was frozen at -80°C for future use. Target cells

VeroE6 were seeded in 24-well plates (5×10^5 cells/mL) at a density of 80% coverage. The

following day, ppVSV-SARS-2-S/GFP particles were transduced into target cells for 60 minutes,

particles pseudotyped with VSV-G, Lassa virus GP, or no glycoprotein were included as

controls. 24 hours following transduction, transduced cells were released from the plate with

trypsin, fixed with 4% formaldehyde, and GFP-positive virus-transduced cells were quantified

using flow cytometry (Bectin Dickson BD-LSRII). To quantify the ability of various SARS-CoV-2

S mutants to mediate fusion, effector cells (HEK293T) were transiently transfected with the

indicated pcDNAintron-SARS-2-S expression vector or measles virus H and F (78). Effector

cells were infected with MVA-T7 four hours following transduction to produce the T7 polymerase

(79). Target cells naturally expressing the receptor ACE2 (Vero) or ACE2 negative cells

(HEK293T) were transfected with pTM1-luciferase, which encodes for firefly luciferase under the

control of a T7 promoter (80). 24 hours following transfection, the target cells were lifted and

added to the effector cells at a 1:1 ratio. 4 hours following co-cultivation, cells were washed,

lysed and luciferase levels were quantified using Promega's Steady-Glo substrate. To visualize

cell-to-cell fusion, Vero cells were co-transfected with pGFP and the pcDNAintron-SARS-2-S

constructs. 24 hours following transfection, syncytia was visualized by fluorescence microscopy.

DATA AVAILABILITY

The mass spectrometry proteomics data are available via ProteomeXchange with identifiers

listed in the KEY RESOURCES TABLE.

SUPPLEMENTAL INFORMATION

Tables (1, 16 tabs), Figures (7), Movies (4), and Simulations (3).

SUPPLEMENTAL LEGEND:

Supplemental Table Tab 1. Glycans modeled as Abundance, Oxford Class, and Processed.

Supplemental Table Tab 2. Cys0015-Cys0136 Disulfide Linked Peptide for SARS-CoV-2 S.

Supplemental Table Tab 3. Detection of N-linked glycans released from SARS-CoV-2 S and

human ACE2. Relative abundance (prevalence) of each species is calculated based on peak

intensity in full MS.

Supplemental Table Tab 4. Detection of O-linked glycans released from SARS-CoV-2 S and

human ACE2. Relative abundance (prevalence) of each species is calculated based on peak

intensity in full MS.

Supplemental Table Tab 5. N-linked glycan occupancy at each site of SARS-CoV-2 S and

human ACE2. Occupancy is calculated using spectral counts assigned to the 18O-Asp-

containing (PNGaseF-cleaved) and/or HexNAc-modified (EndoH-cleaved) peptides and their

unmodified counterparts. Sequon refers to the Asn-x-Ser/Thr/Cys, Asn-Gly-x sequences.

Supplemental Table Tab 6. N-linked glycan compositions identified at each site of SARS-CoV-

2 S and human ACE2. Asn(N)# indicates the numbers of asparagines in protein sequences. In

compositions: N=HexNAc, H=Hexose (Hex), F=Fucose (Fuc), A=Neu5Ac, S=Sulfate. In

fucosylation: NoFuc=No Fuc identified; 1Core=One Fuc identified at core position; 1Term=One

Fuc identified at terminal position; 1Core and 1Term=One Fuc identified as a mixture of core

and terminal positions; 1Core1Term=Two Fuc identified and one is at core and the other is at

terminal; 2Term=Two Fuc identified at terminal positions; 1Core1Term and 2Term=Two Fuc

identified as a mixture of core and terminal positions; 1Core2Term=Three Fuc identified and

one is at core and the others are at terminal; 3Term=Three Fuc identified at terminal positions;

1Core2Term and 3Term=Three Fuc identified as a mixture of core and terminal positions;

1Core3Term=Four Fuc identified and one is at core and the others are at terminal; 4Term=Four

Fuc identified at terminal positions; 1Core3Term and 4Term=Four Fuc identified as a mixture of

core and terminal positions; 1Core4Term=Five Fuc identified and one is at core and the others

are at terminal.

Supplemental Table Tab 7. N-linked glycan types identified at each site of SARS-CoV-2 S and

human ACE2. All N-linked glycans are categorized into 3 types: high-mannose, hybrid and

complex.

Supplemental Table Tab 8. N-linked glycan oxford classes identified at each site of SARS-

CoV-2 S and human ACE2. All N-linked glycan compositions are categorized into 22 classes:

M9 to M5 respectively is defined as HexNAc(2)Hex(9~5)Fuc(0~1); M1-M4 is defined as

HexNAc(2)Hex(4~1)Fuc(0~1); Hybrid is defined as HexNAc(3~6)Hex(5~9)Fuc(0)NeuAc(0~1)

and F-Hybrid is defined as HexNAc(3~6)Hex(5~9)Fuc(1~2)NeuAc(0~1). Complex-type glycans

are classified based on the number of antenna, fucosylation, and sulfation:

HexNAc(3)Hex(3~4)Fuc(0)NeuAc(0~1) is assigned as A1 with

HexNAc(3)Hex(3~4)Fuc(1~2)NeuAc(0~1) assigned as F-A1;

HexNAc(4)Hex(3~5)Fuc(0)NeuAc(0~2) is assigned as A2/A1B with

HexNAc(4)Hex(3~5)Fuc(1~5)NeuAc(0~2) assigned as F-A2/A1B;

HexNAc(5)Hex(3~6)Fuc(0)NeuAc(0~3) is assigned as A3/A2B with

HexNAc(5)Hex(3~6)Fuc(1~3)NeuAc(0~3) assigned as F-A3/A2B;

HexNAc(6)Hex(3~7)Fuc(0)NeuAc(0~4) is assigned as A4/A3B with

HexNAc(6)Hex(3~7)Fuc(1~3)NeuAc(0~4) assigned as F-A4/A3B;

HexNAc(7)Hex(3~8)Fuc(0)NeuAc(0~1) is assigned as A5/A4B with

HexNAc(7)Hex(3~8)Fuc(1~3)NeuAc(0~1) assigned as F-A5/A4B; HexNAc(8)Hex(3~9)Fuc(0) is

assigned as A6/A5B with HexNAc(8)Hex(3~9)Fuc(1) assigned as F-A6/A5B; any glycans

identified with a sulfate are assigned as Sulfated.

Supplemental Table Tab 9. O-linked glycan compositions identified at each site of SARS-CoV-

2 S and human ACE2. Ser/Thr# indicates the numbers of serines or threonines in protein

sequences. In compositions: N=HexNAc, H=Hexose (Hex), F=Fucose (Fuc), and A=Neu5Ac.

Supplemental Table Tab 10. O-linked glycan occupancy at each site of SARS-CoV-2 S and

human ACE2. Occupancy is calculated using spectral counts assigned to the glycosylated

peptides and their unmodified counterparts.

Supplemental Table Tab 11. SARS-CoV-2 S and human ACE2 variants.

Supplemental Table Tab 12. Proteomic Analyses of purified S and ACE2.

Supplemental Table Tab 13. Sulfated N-linked glycans released from SARS-CoV-2 S.

Following permethylation, almost all of the sulfated hybrid and complex N-glycans are recovered

in the organic phase despite their anionic charge. Organic phase permethylated glycans were

analyzed by mass spectrometry using negative ion mode. The indicated glycan structures are

consistent with the compositions detected at the m/z values shown.

Supplemental Table Tab 14. Surface Antigen Exposure of Abundance Glycosylated S. The

scale used is 0 (not accessible) to 1.0 (fully accessible).

Supplemental Table Tab 15. ACE2-Glycan-S-Peptide Interactions.

The scale used is 0 (no interaction) to 1.0 (interacted throughout entire simulation).

Supplemental Table Tab 16. S-Glycan-ACE2-Peptide Interactions.

The scale used is 0 (no interaction) to 1.0 (interacted throughout entire simulation).

Supplemental Figure S1. Defining N-terminus of ACE2 as pyro-glutamine at site Q0018.

Representative HCD MS2 spectrum shown.

Supplemental Figure S2. Disulfide bond formed between Cysteines 0015 and 0136 of SARS-

CoV-2 S. Representative EThcD MS2 spectrum shown.

Supplemental Figure S3. Signal P prediction of two different start methionines for SARS-CoV-

2 S.

Supplemental Figure S4. Functional characterization of various S constructs in Pseudovirus.

(A) Syncytia produced by SARS-CoV-2 S constructs in VeroE6 cells co-transfected with a GFP

plasmid to visualize cell-to-cell fusion. Quantification of fusion using a luciferase

complementation assay in 293T (B) or VeroE6 cells (C). (D) Transduction efficiency in Vero E6

cells of ppVSV-GFP particles coated in the indicated glycoprotein. Results suggest that start

methionine does not alter fusion or efficiency.

Supplemental Figure S5. Detection of O-linked glycans released from SARS-CoV-2 S and

human ACE2. The detected O-glycans were categorized based on their structures and types.

Relative abundance (prevalence) of each species is calculated based on peak intensity in full

MS.

Supplemental Figure S6. O-linked glycans detected at site T0323 of SARS-CoV-2 S.

Representative Step-HCD spectra shown for 6 glycoforms.

Supplemental Figure S7. Sequence alignments of SARS-CoV-1 and SARS-CoV-2 S variants

as well as alignment of multiple S proteins from related coronaviruses.

Supplemental Movie A: Linked to Figure 4C, Glycosylated S antigen accessibility

Supplemental Movie B: Linked to Figure 6A, Glycosylated ACE2 with variants

Supplemental Movie C: Linked to Figure 7C, Interface of ACE2-S Complex

Supplemental Movie D: Linked to Figure 7C, the glycosylated ACE2-S Complex

Simulation 1: Linked to Figure 7A, Abundance glycoforms on ACE2-S Complex

Simulation 2: Linked to Figure 7A, Oxford class glycoforms on ACE2-S Complex

Simulation 3: Linked to Figure 7A, Processed glycoforms on ACE2-S Complex

REFERENCES

1.   Zhou, P., Yang, X. L., Wang, X. G., Hu, B., Zhang, L., Zhang, W., Si, H. R., Zhu, Y., Li, B.,
Huang, C. L., Chen, H. D., Chen, J., Luo, Y., Guo, H., Jiang, R. D., Liu, M. Q., Chen, Y., Shen,
X. R., Wang, X., Zheng, X. S., Zhao, K., Chen, Q. J., Deng, F., Liu, L. L., Yan, B., Zhan, F. X.,
Wang, Y. Y., Xiao, G. F., and Shi, Z. L. (2020) A pneumonia outbreak associated with a
new coronavirus of probable bat origin. *Nature* **579**, 270-273
2.   Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., Wang, W., Song, H., Huang, B., Zhu, N., Bi,
Y., Ma, X., Zhan, F., Wang, L., Hu, T., Zhou, H., Hu, Z., Zhou, W., Zhao, L., Chen, J., Meng,
Y., Wang, J., Lin, Y., Yuan, J., Xie, Z., Ma, J., Liu, W. J., Wang, D., Xu, W., Holmes, E. C.,
Gao, G. F., Wu, G., Chen, W., Shi, W., and Tan, W. (2020) Genomic characterisation
and epidemiology of 2019 novel coronavirus: implications for virus origins and
receptor binding. *Lancet* **395**, 565-574
3.   Zhong, N. S., Zheng, B. J., Li, Y. M., Poon, Xie, Z. H., Chan, K. H., Li, P. H., Tan, S. Y.,
Chang, Q., Xie, J. P., Liu, X. Q., Xu, J., Li, D. X., Yuen, K. Y., Peiris, and Guan, Y. (2003)
Epidemiology and cause of severe acute respiratory syndrome (SARS) in
Guangdong, People's Republic of China, in February, 2003. *Lancet* **362**, 1353-1358
4.   Xia, X. (2020) Extreme genomic CpG deficiency in SARS-CoV-2 and evasion of host
antiviral defense. *Mol Biol Evol*
5.   Zhang, T., Wu, Q., and Zhang, Z. (2020) Probable Pangolin Origin of SARS-CoV-2
Associated with the COVID-19 Outbreak. *Curr Biol* **30**, 1346-1351 e1342
6.   Alijotas-Reig, J., Esteve-Valverde, E., Belizna, C., Selva-O'Callaghan, A., Pardos-Gea, J.,
Quintana, A., Mekinian, A., Anunciacion-Llunell, A., and Miro-Mur, F. (2020)
Immunomodulatory therapy for the management of severe COVID-19. Beyond the
anti-viral therapy: A comprehensive review. *Autoimmun Rev* **19**, 102569
7.   Beigel, J. H., Tomashek, K. M., Dodd, L. E., Mehta, A. K., Zingman, B. S., Kalil, A. C.,
Hohmann, E., Chu, H. Y., Luetkemeyer, A., Kline, S., Lopez de Castilla, D., Finberg, R.
W., Dierberg, K., Tapson, V., Hsieh, L., Patterson, T. F., Paredes, R., Sweeney, D. A.,
Short, W. R., Touloumi, G., Lye, D. C., Ohmagari, N., Oh, M. D., Ruiz-Palacios, G. M.,
Benfield, T., Fatkenheuer, G., Kortepeter, M. G., Atmar, R. L., Creech, C. B., Lundgren,
J., Babiker, A. G., Pett, S., Neaton, J. D., Burgess, T. H., Bonnett, T., Green, M.,
Makowski, M., Osinusi, A., Nayak, S., Lane, H. C., and Members, A.-S. G. (2020)
Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med*
8.   Beun, R., Kusadasi, N., Sikma, M., Westerink, J., and Huisman, A. (2020)
Thromboembolic events and apparent heparin resistance in patients infected with
SARS-CoV-2. *Int J Lab Hematol* **42 Suppl 1**, 19-20
9.   Dashti-Khavidaki, S., and Khalili, H. (2020) Considerations for Statin Therapy in
Patients with COVID-19. *Pharmacotherapy* **40**, 484-486
10.   Fedson, D. S., Opal, S. M., and Rordam, O. M. (2020) Hiding in Plain Sight: an
Approach to Treating Patients with Severe COVID-19 Infection. *mBio* **11**

11.   Shi, Q., Zhou, Q., Wang, X., Liao, J., Yu, Y., Wang, Z., Lu, S., Ma, Y., Xun, Y., Luo, X., Li, W.,
960   Fukuoka, T., Ahn, H. S., Lee, M. S., Luo, Z., Liu, E., Chen, Y., Li, Q., Yang, K., Guan, Q.,
961   Evidence, C.-., and Recommendations Working, G. (2020) Potential effectiveness and
962   safety of antiviral agents in children with coronavirus disease 2019: a rapid review
963   and meta-analysis. *Ann Transl Med* **8**, 624
964   12.   Tang, N., Bai, H., Chen, X., Gong, J., Li, D., and Sun, Z. (2020) Anticoagulant treatment
965   is associated with decreased mortality in severe coronavirus disease 2019 patients
966   with coagulopathy. *J Thromb Haemost* **18**, 1094-1099
967   13.   Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S.,
968   Schiergens, T. S., Herrler, G., Wu, N. H., Nitsche, A., Muller, M. A., Drosten, C., and
969   Pohlmann, S. (2020) SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
970   Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278
971   14.   Li, W., Moore, M. J., Vasilieva, N., Sui, J., Wong, S. K., Berne, M. A., Somasundaran, M.,
972   Sullivan, J. L., Luzuriaga, K., Greenough, T. C., Choe, H., and Farzan, M. (2003)
973   Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus.
974   *Nature* **426**, 450-454
975   15.   Watanabe, Y., Bowden, T. A., Wilson, I. A., and Crispin, M. (2019) Exploitation of
976   glycosylation in enveloped virus pathobiology. *Biochim Biophys Acta Gen Subj* **1863**,
977   1480-1497
978   16.   Li, F. (2016) Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu*
979   *Rev Virol* **3**, 237-261
980   17.   Lei, C., Qian, K., Li, T., Zhang, S., Fu, W., Ding, M., and Hu, S. (2020) Neutralization of
981   SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig. *Nat Commun* **11**,
982   2070
983   18.   Monteil, V., Kwon, H., Prado, P., Hagelkruys, A., Wimmer, R. A., Stahl, M., Leopoldi, A.,
984   Garreta, E., Hurtado Del Pozo, C., Prosper, F., Romero, J. P., Wirnsberger, G., Zhang,
985   H., Slutsky, A. S., Conder, R., Montserrat, N., Mirazimi, A., and Penninger, J. M. (2020)
986   Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-
987   Grade Soluble Human ACE2. *Cell* **181**, 905-913 e907
988   19.   Watanabe, Y., Allen, J. D., Wrapp, D., McLellan, J. S., and Crispin, M. (2020) Site-
989   specific glycan analysis of the SARS-CoV-2 spike. *Science*
990   20.   Watanabe, Y., Berndsen, Z. T., Raghwani, J., Seabright, G. E., Allen, J. D., Pybus, O. G.,
991   McLellan, J. S., Wilson, I. A., Bowden, T. A., Ward, A. B., and Crispin, M. (2020)
992   Vulnerabilities in coronavirus glycan shields despite extensive glycosylation. *Nat*
993   *Commun* **11**, 2688
994   21.   Stertz, S., Reichelt, M., Spiegel, M., Kuri, T., Martinez-Sobrido, L., Garcia-Sastre, A.,
995   Weber, F., and Kochs, G. (2007) The intracellular sites of early replication and
996   budding of SARS-coronavirus. *Virology* **361**, 304-315
997   22.   Ujike, M., and Taguchi, F. (2015) Incorporation of spike and membrane
998   glycoproteins into coronavirus virions. *Viruses* **7**, 1700-1725
999   23.   Shajahan, A., Supekar, N. T., Gleinich, A. S., and Azadi, P. (2020) Deducing the N- and
1000   O- glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2.
1001   *Glycobiology*
1002   24.   Ritchie, G., Harvey, D. J., Feldmann, F., Stroeher, U., Feldmann, H., Royle, L., Dwek, R.
1003   A., and Rudd, P. M. (2010) Identification of N-linked carbohydrates from severe
1004   acute respiratory syndrome (SARS) spike glycoprotein. *Virology* **399**, 257-269

1005   25.   Duan, H., Chen, X., Boyington, J. C., Cheng, C., Zhang, Y., Jafari, A. J., Stephens, T.,
1006   Tsybovsky, Y., Kalyuzhniy, O., Zhao, P., Menis, S., Nason, M. C., Normandin, E.,
1007   Mukhamedova, M., DeKosky, B. J., Wells, L., Schief, W. R., Tian, M., Alt, F. W., Kwong,
1008   P. D., and Mascola, J. R. (2018) Glycan Masking Focuses Immune Responses to the
1009   HIV-1 CD4-Binding Site and Enhances Elicitation of VRC01-Class Precursor
1010   Antibodies. *Immunity* **49**, 301-311 e305

1011   26.   Escolano, A., Gristick, H. B., Abernathy, M. E., Merkenschlager, J., Gautam, R., Oliveira,
1012   T. Y., Pai, J., West, A. P., Jr., Barnes, C. O., Cohen, A. A., Wang, H., Golijanin, J., Yost, D.,
1013   Keeffe, J. R., Wang, Z., Zhao, P., Yao, K. H., Bauer, J., Nogueira, L., Gao, H., Voll, A. V.,
1014   Montefiori, D. C., Seaman, M. S., Gazumyan, A., Silva, M., McGuire, A. T., Stamatatos, L.,
1015   Irvine, D. J., Wells, L., Martin, M. A., Bjorkman, P. J., and Nussenzweig, M. C. (2019)
1016   Immunization expands B cells specific to HIV-1 V3 glycan in mice and macaques.
1017   *Nature* **570**, 468-473

1018   27.   Seabright, G. E., Cottrell, C. A., van Gils, M. J., D'Addabbo, A., Harvey, D. J., Behrens, A.
1019   J., Allen, J. D., Watanabe, Y., Scaringi, N., Polveroni, T. M., Maker, A., Vasiljevic, S., de
1020   Val, N., Sanders, R. W., Ward, A. B., and Crispin, M. (2020) Networks of HIV-1
1021   Envelope Glycans Maintain Antibody Epitopes in the Face of Glycan Additions and
1022   Deletions. *Structure*

1023   28.   Yu, W. H., Zhao, P., Draghi, M., Arevalo, C., Karsten, C. B., Suscovich, T. J., Gunn, B.,
1024   Streeck, H., Brass, A. L., Tiemeyer, M., Seaman, M., Mascola, J. R., Wells, L.,
1025   Lauffenburger, D. A., and Alter, G. (2018) Exploiting glycan topography for
1026   computational design of Env glycoprotein antigenicity. *PLoS Comput Biol* **14**,
1027   e1006093

1028   29.   Pinto, D., Park, Y. J., Beltramello, M., Walls, A. C., Tortorici, M. A., Bianchi, S., Jaconi, S.,
1029   Culap, K., Zatta, F., De Marco, A., Peter, A., Guarino, B., Spreafico, R., Cameroni, E.,
1030   Case, J. B., Chen, R. E., Havenar-Daughton, C., Snell, G., Telenti, A., Virgin, H. W.,
1031   Lanzavecchia, A., Diamond, M. S., Fink, K., Veesler, D., and Corti, D. (2020) Cross-
1032   neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature*
1033   **583**, 290-295

1034   30.   Tikellis, C., Bernardi, S., and Burns, W. C. (2011) Angiotensin-converting enzyme 2 is
1035   a key modulator of the renin-angiotensin system in cardiovascular and renal
1036   disease. *Curr Opin Nephrol Hypertens* **20**, 62-68

1037   31.   Lambert, D. W., Yarski, M., Warner, F. J., Thornhill, P., Parkin, E. T., Smith, A. I.,
1038   Hooper, N. M., and Turner, A. J. (2005) Tumor necrosis factor-alpha convertase
1039   (ADAM17) mediates regulated ectodomain shedding of the severe-acute respiratory
1040   syndrome-coronavirus (SARS-CoV) receptor, angiotensin-converting enzyme-2
1041   (ACE2). *J Biol Chem* **280**, 30113-30119

1042   32.   Epelman, S., Shrestha, K., Troughton, R. W., Francis, G. S., Sen, S., Klein, A. L., and
1043   Tang, W. H. (2009) Soluble angiotensin-converting enzyme 2 in human heart failure:
1044   relation with myocardial function and clinical outcomes. *J Card Fail* **15**, 565-571

1045   33.   Li, W., Zhang, C., Sui, J., Kuhn, J. H., Moore, M. J., Luo, S., Wong, S. K., Huang, I. C., Xu, K.,
1046   Vasilieva, N., Murakami, A., He, Y., Marasco, W. A., Guan, Y., Choe, H., and Farzan, M.
1047   (2005) Receptor and viral determinants of SARS-coronavirus adaptation to human
1048   ACE2. *EMBO J* **24**, 1634-1643

1049   34.   Marth, J. D., and Grewal, P. K. (2008) Mammalian glycosylation in immunity. *Nat Rev*
1050   *Immunol* **8**, 874-887

35.	Varki, A. (2017) Biological roles of glycans. *Glycobiology* **27**, 3-49

36.	Yu, J., Tostanoski, L. H., Peter, L., Mercado, N. B., McMahan, K., Mahrokhian, S. H.,
Nkolola, J. P., Liu, J., Li, Z., Chandrashekar, A., Martinez, D. R., Loos, C., Atyeo, C.,
Fischinger, S., Burke, J. S., Slein, M. D., Chen, Y., Zuiani, A., FJ, N. L., Travers, M., Habibi,
S., Pessaint, L., Van Ry, A., Blade, K., Brown, R., Cook, A., Finneyfrock, B., Dodson, A.,
Teow, E., Velasco, J., Zahn, R., Wegmann, F., Bondzie, E. A., Dagotto, G., Gebre, M. S.,
He, X., Jacob-Dolan, C., Kirilova, M., Kordana, N., Lin, Z., Maxfield, L. F., Nampanya, F.,
Nityanandam, R., Ventura, J. D., Wan, H., Cai, Y., Chen, B., Schmidt, A. G., Wesemann,
D. R., Baric, R. S., Alter, G., Andersen, H., Lewis, M. G., and Barouch, D. H. (2020) DNA
vaccine protection against SARS-CoV-2 in rhesus macaques. *Science*

37.	Almagro Armenteros, J. J., Tsirigos, K. D., Sonderby, C. K., Petersen, T. N., Winther, O.,
Brunak, S., von Heijne, G., and Nielsen, H. (2019) SignalP 5.0 improves signal peptide
predictions using deep neural networks. *Nat Biotechnol* **37**, 420-423

38.	Aoki, K., Perlman, M., Lim, J. M., Cantu, R., Wells, L., and Tiemeyer, M. (2007)
Dynamic developmental elaboration of N-linked glycan complexity in the Drosophila
melanogaster embryo. *J Biol Chem* **282**, 9127-9142

39.	Aoki, K., Porterfield, M., Lee, S. S., Dong, B., Nguyen, K., McGlamry, K. H., and
Tiemeyer, M. (2008) The diversity of O-linked glycans expressed during Drosophila
melanogaster development reflects stage- and tissue-specific requirements for cell
signaling. *J Biol Chem* **283**, 30385-30400

40.	Weatherly, D. B., Arpinar, F. S., Porterfield, M., Tiemeyer, M., York, W. S., and
Ranzinger, R. (2019) GRITS Toolbox-a freely available software for processing,
annotating and archiving glycomics mass spectrometry data. *Glycobiology* **29**, 452-
460

41.	Wang, S., Voronin, Y., Zhao, P., Ishihara, M., Mehta, N., Porterfield, M., Chen, Y.,
Bartley, C., Hu, G., Han, D., Wells, L., Tiemeyer, M., and Lu, S. (2020) Glycan Profiles of
gp120 Protein Vaccines from Four Major HIV-1 Subtypes Produced from Different
Host Cell Lines under Non-GMP or GMP Conditions. *J Virol* **94**

42.	Ruiz-Canada, C., Kelleher, D. J., and Gilmore, R. (2009) Cotranslational and
posttranslational N-glycosylation of polypeptides by distinct mammalian OST
isoforms. *Cell* **136**, 272-283

43.	Zielinska, D. F., Gnad, F., Wisniewski, J. R., and Mann, M. (2010) Precision mapping of
an in vivo N-glycoproteome reveals rigid topological and sequence constraints. *Cell*
141, 897-907

44.	Palmisano, G., Melo-Braga, M. N., Engholm-Keller, K., Parker, B. L., and Larsen, M. R.
(2012) Chemical deamidation: a common pitfall in large-scale N-linked
glycoproteomic mass spectrometry-based analyses. *J Proteome Res* **11**, 1949-1957

45.	Zhou, T., Doria-Rose, N. A., Cheng, C., Stewart-Jones, G. B. E., Chuang, G. Y., Chambers,
M., Druz, A., Geng, H., McKee, K., Kwon, Y. D., O'Dell, S., Sastry, M., Schmidt, S. D., Xu,
K., Chen, L., Chen, R. E., Louder, M. K., Pancera, M., Wanninger, T. G., Zhang, B., Zheng,
A., Farney, S. K., Foulds, K. E., Georgiev, I. S., Joyce, M. G., Lemmin, T., Narpala, S.,
Rawi, R., Soto, C., Todd, J. P., Shen, C. H., Tsybovsky, Y., Yang, Y., Zhao, P., Haynes, B. F.,
Stamatatos, L., Tiemeyer, M., Wells, L., Scorpio, D. G., Shapiro, L., McDermott, A. B.,
Mascola, J. R., and Kwong, P. D. (2017) Quantification of the Impact of the HIV-1-
Glycan Shield on Antibody Elicitation. *Cell Rep* **19**, 719-732

46. Liu, M. Q., Zeng, W. F., Fang, P., Cao, W. Q., Liu, C., Yan, G. Q., Zhang, Y., Peng, C., Wu, J.
Q., Zhang, X. J., Tu, H. J., Chi, H., Sun, R. X., Cao, Y., Dong, M. Q., Jiang, B. Y., Huang, J. M.,
Shen, H. L., Wong, C. C. L., He, S. M., and Yang, P. Y. (2017) pGlyco 2.0 enables
precision N-glycoproteomics with comprehensive quality control and one-step mass
spectrometry for intact glycopeptide identification. *Nat Commun* **8**, 438
47. Bern, M., Kil, Y. J., and Becker, C. (2012) Byonic: advanced peptide and protein
identification software. *Curr Protoc Bioinformatics* **Chapter 13**, Unit13 20
48. Wrapp, D., Wang, N., Corbett, K. S., Goldsmith, J. A., Hsieh, C. L., Abiona, O., Graham, B.
S., and McLellan, J. S. (2020) Cryo-EM structure of the 2019-nCoV spike in the
prefusion conformation. *Science* **367**, 1260-1263
49. Ferreira, R. C., Grant, O. C., Moyo, T., Dorfman, J. R., Woods, R. J., Travers, S. A., and
Wood, N. T. (2018) Structural Rearrangements Maintain the Glycan Shield of an HIV-
1 Envelope Trimer After the Loss of a Glycan. *Sci Rep* **8**, 15031
50. Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., Zhang, Q., Shi, X., Wang, Q., Zhang, L., and
Wang, X. (2020) Structure of the SARS-CoV-2 spike receptor-binding domain bound
to the ACE2 receptor. *Nature* **581**, 215-220
51. Wang, C. C., Chen, J. R., Tseng, Y. C., Hsu, C. H., Hung, Y. F., Chen, S. W., Chen, C. M.,
Khoo, K. H., Cheng, T. J., Cheng, Y. S., Jan, J. T., Wu, C. Y., Ma, C., and Wong, C. H. (2009)
Glycans on influenza hemagglutinin affect receptor binding and immune response.
Proc Natl Acad Sci U S A **106**, 18137-18142
52. Behrens, A. J., Harvey, D. J., Milne, E., Cupo, A., Kumar, A., Zitzmann, N., Struwe, W. B.,
Moore, J. P., and Crispin, M. (2017) Molecular Architecture of the Cleavage-
Dependent Mannose Patch on a Soluble HIV-1 Envelope Glycoprotein Trimer. *J Virol*
91
53. Kristic, J., Vuckovic, F., Menni, C., Klaric, L., Keser, T., Beceheli, I., Pucic-Bakovic, M.,
Novokmet, M., Mangino, M., Thaqi, K., Rudan, P., Novokmet, N., Sarac, J., Missoni, S.,
Kolcic, I., Polasek, O., Rudan, I., Campbell, H., Hayward, C., Aulchenko, Y., Valdes, A.,
Wilson, J. F., Gornik, O., Primorac, D., Zoldos, V., Spector, T., and Lauc, G. (2014)
Glycans are a novel biomarker of chronological and biological ages. *J Gerontol A Biol*
Sci Med Sci **69**, 779-789
54. Pavic, T., Dilber, D., Kifer, D., Selak, N., Keser, T., Ljubicic, D., Vukic Dugac, A., Lauc, G.,
Rumora, L., and Gornik, O. (2018) N-glycosylation patterns of plasma proteins and
immunoglobulin G in chronic obstructive pulmonary disease. *J Transl Med* **16**, 323
55. Rudman, N., Gornik, O., and Lauc, G. (2019) Altered N-glycosylation profiles as
potential biomarkers and drug targets in diabetes. *FEBS Lett* **593**, 1598-1615
56. Gebrehiwot, A. G., Melka, D. S., Kassaye, Y. M., Rehan, I. F., Rangappa, S., Hinou, H.,
Kamiyama, T., and Nishimura, S. I. (2018) Healthy human serum N-glycan profiling
reveals the influence of ethnic variation on the identified cancer-relevant glycan
biomarkers. *PLoS One* **13**, e0209515
57. Wrobel, A. G., Benton, D. J., Xu, P., Roustan, C., Martin, S. R., Rosenthal, P. B., Skehel, J.
J., and Gamblin, S. J. (2020) SARS-CoV-2 and bat RaTG13 spike glycoprotein
structures inform on virus evolution and furin-cleavage effects. *Nat Struct Mol Biol*
58. Casalino, L., Gaieb, Z., Dommer, A. C., Harbison, A. M., Fogarty, C. A., Barros, E. P.,
Taylor, B. C., Fadda, E., and Amaro, R. E. (2020) Shielding and Beyond: The Roles of
Glycans in SARS-CoV-2 Spike Protein. *bioRxiv*

59.   Struwe, W. B., Chertova, E., Allen, J. D., Seabright, G. E., Watanabe, Y., Harvey, D. J.,
1142   Medina-Ramirez, M., Roser, J. D., Smith, R., Westcott, D., Keele, B. F., Bess, J. W., Jr.,
1143   Sanders, R. W., Lifson, J. D., Moore, J. P., and Crispin, M. (2018) Site-Specific
1144   Glycosylation of Virion-Derived HIV-1 Env Is Mimicked by a Soluble Trimeric
1145   Immunogen. *Cell Rep* **24**, 1958-1966 e1955

1146   60.   Thieker, D. F., Hadden, J. A., Schulten, K., and Woods, R. J. (2016) 3D implementation
1147   of the symbol nomenclature for graphical representation of glycans. *Glycobiology*
1148   **26**, 786-787

1149   61.   Rice, P., Longden, I., and Bleasby, A. (2000) EMBOSS: the European Molecular
1150   Biology Open Software Suite. *Trends Genet* **16**, 276-277

1151   62.   Shaik, M. M., Peng, H., Lu, J., Rits-Volloch, S., Xu, C., Liao, M., and Chen, B. (2019)
1152   Structural basis of coreceptor recognition by HIV-1 envelope spike. *Nature* **565**,
1153   318-323

1154   63.   Anumula, K. R., and Taylor, P. B. (1992) A comprehensive procedure for preparation
1155   of partially methylated alditol acetates from glycoprotein carbohydrates. *Anal*
1156   *Biochem* **203**, 101-108

1157   64.   Varki, A., Cummings, R. D., Aebi, M., Packer, N. H., Seeberger, P. H., Esko, J. D., Stanley,
1158   P., Hart, G., Darvill, A., Kinoshita, T., Prestegard, J. J., Schnaar, R. L., Freeze, H. H.,
1159   Marth, J. D., Bertozzi, C. R., Etzler, M. E., Frank, M., Vliegenthart, J. F., Lutteke, T.,
1160   Perez, S., Bolton, E., Rudd, P., Paulson, J., Kanehisa, M., Toukach, P., Aoki-Kinoshita, K.
1161   F., Dell, A., Narimatsu, H., York, W., Taniguchi, N., and Kornfeld, S. (2015) Symbol
1162   Nomenclature for Graphical Representations of Glycans. *Glycobiology* **25**, 1323-1324

1163   65.   Wells, L., Hart, G. W., and Athens Guidelines for the Publication of Glycomics, D.
1164   (2013) Glycomics: building upon proteomics to advance glycosciences. *Mol Cell*
1165   *Proteomics* **12**, 833-835

1166   66.   Kahsay, R., Vora, J., Navelkar, R., Mousavi, R., Fochtman, B. C., Holmes, X.,
1167   Pattabiraman, N., Ranzinger, R., Mahadik, R., Williamson, T., Kulkarni, S., Agarwal, G.,
1168   Martin, M., Vasudev, P., Garcia, L., Edwards, N., Zhang, W., Natale, D. A., Ross, K., Aoki-
1169   Kinoshita, K. F., Campbell, M. P., York, W. S., and Mazumder, R. (2020) GlyGen data
1170   model and processing workflow. *Bioinformatics* **36**, 3941-3943

1171   67.   Tiemeyer, M., Aoki, K., Paulson, J., Cummings, R. D., York, W. S., Karlsson, N. G.,
1172   Lisacek, F., Packer, N. H., Campbell, M. P., Aoki, N. P., Fujita, A., Matsubara, M.,
1173   Shinmachi, D., Tsuchiya, S., Yamada, I., Pierce, M., Ranzinger, R., Narimatsu, H., and
1174   Aoki-Kinoshita, K. F. (2017) GlyTouCan: an accessible glycan structure repository.
1175   *Glycobiology* **27**, 915-919

1176   68.   York, W. S., Mazumder, R., Ranzinger, R., Edwards, N., Kahsay, R., Aoki-Kinoshita, K.
1177   F., Campbell, M. P., Cummings, R. D., Feizi, T., Martin, M., Natale, D. A., Packer, N. H.,
1178   Woods, R. J., Agarwal, G., Arpinar, S., Bhat, S., Blake, J., Castro, L. J. G., Fochtman, B.,
1179   Gildersleeve, J., Goldman, R., Holmes, X., Jain, V., Kulkarni, S., Mahadik, R., Mehta, A.,
1180   Mousavi, R., Nakarakommula, S., Navelkar, R., Pattabiraman, N., Pierce, M. J., Ross, K.,
1181   Vasudev, P., Vora, J., Williamson, T., and Zhang, W. (2020) GlyGen: Computational
1182   and Informatics Resources for Glycoscience. *Glycobiology* **30**, 72-73

1183   69.   Madeira, F., Madhusoodanan, N., Lee, J., Tivey, A. R. N., and Lopez, R. (2019) Using
1184   EMBL-EBI Services via Web Interface and Programmatically via Web Services. *Curr*
1185   *Protoc Bioinformatics* **66**, e74

70. Nivedha, A. K., Makeneni, S., Foley, B. L., Tessier, M. B., and Woods, R. J. (2014)
Importance of ligand conformational energies in carbohydrate docking: Sorting the
wheat from the chaff. *J Comput Chem* **35**, 526-539
71. Grant, O. C., Xue, X., Ra, D., Khatamian, A., Foley, B. L., and Woods, R. J. (2016) Gly-
Spec: a webtool for predicting glycan specificity by integrating glycan array
screening data and 3D structure. *Glycobiology* **26**, 1027-1028
72. Peng, W., de Vries, R. P., Grant, O. C., Thompson, A. J., McBride, R., Tsogtbaatar, B.,
Lee, P. S., Razi, N., Wilson, I. A., Woods, R. J., and Paulson, J. C. (2017) Recent H3N2
Viruses Have Evolved Specificity for Extended, Branched Human-type Receptors,
Conferring Potential for Increased Avidity. *Cell Host Microbe* **21**, 23-34
73. Gotz, A. W., Williamson, M. J., Xu, D., Poole, D., Le Grand, S., and Walker, R. C. (2012)
Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 1.
Generalized Born. *J Chem Theory Comput* **8**, 1542-1555
74. Salomon-Ferrer, R., Gotz, A. W., Poole, D., Le Grand, S., and Walker, R. C. (2013)
Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 2.
Explicit Solvent Particle Mesh Ewald. *J Chem Theory Comput* **9**, 3878-3888
75. Kirschner, K. N., Yongye, A. B., Tschampel, S. M., Gonzalez-Outeirino, J., Daniels, C. R.,
Foley, B. L., and Woods, R. J. (2008) GLYCAM06: a generalizable biomolecular force
field. Carbohydrates. *J Comput Chem* **29**, 622-655
76. Maier, J. A., Martinez, C., Kasavajhala, K., Wickstrom, L., Hauser, K. E., and
Simmerling, C. (2015) ff14SB: Improving the Accuracy of Protein Side Chain and
Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696-3713
77. Darden, T. A., and Pedersen, L. G. (1993) Molecular modeling: an experimental tool.
Environ Health Perspect **101**, 410-412
78. Brindley, M. A., Chaudhury, S., and Plemper, R. K. (2015) Measles virus glycoprotein
complexes preassemble intracellularly and relax during transport to the cell surface
in preparation for fusion. *J Virol* **89**, 1230-1241
79. Paal, M., Heel, T., Schneider, R., and Auer, B. (2009) A novel Ecotin-Ubiquitin-Tag
(ECUT) for efficient, soluble peptide production in the periplasm of Escherichia coli.
Microb Cell Fact **8**, 7
80. Brindley, M. A., and Plemper, R. K. (2010) Blue native PAGE and biomolecular
complementation reveal a tetrameric or higher-order oligomer organization of the
physiological measles virus attachment protein H. *J Virol* **84**, 12174-12184

1219

A

Soluble Human ACE2

```
MSSSSWLLLS LVAVTAAQST IEEQAKTFLD KFNHEAEDLF YQSSLASWNY NTNITEENVQ   60
NMNNAGDKWS AFLKEQSTLA QMYPLQEIQN LTVKLQLQAL QQNGSSVLSE DKSKRLNTIL  120
NTMSTIYSTG KVCNPDNPQE CLLLEPGLNE IMANSLDYNE RLWAWESWRS EVGKQLRPLY  180
EEYVVLKNEM ARANHYEDYG DYWRGDYEVN GVDGYDYSRG QLIEDVEHTF EEIKPLYEHL  240
HAYVRAKLMN AYPSYISPIG CLPAHLLGDM WGRFWTNLYS LTVPFGQKPN IDVTDAMVDQ  300
AWDAQRIFKE AEKFFVSVGL PNMTQGFWEN SMLTDPGNVQ KAVCHPTAWD LGKGDFRILM  360
CTKVTMDDFL TAHHEMGHIQ YDMAYAAQPF LLRNGANEGF HEAVGEIMSL SAATPKHLKS  420
IGLLSPDFQE DNETEINFLL KQALTIVGTL PFTYMLEKWR WMVFKGEIPK DQWMKKWWEM  480
KREIVGVVEP VPHDETYCDP ASLFHVSNDY SFIRYYTRTL YQFQFQEALC QAAKHEGPLH  540
KCDISNSTEA GQKLFNMLRL GKSEPWTLAL ENVVGAKNMN VRPLLNYFEP LFTWLKDQNK  600
NSFVGWSTDW SPYADSGGSH HHHHH                                        625
```

SARS-CoV-2 S Trimer Immunogen

```
MFVFLVLLPL VSSQCVNLTT RTQLPPAYTN SFTRGVYYPD KVFRSSVLHS TQDLFLPFFS   60
NVTWFHAIHV SGTNGTKRFD NPVLPFNDGV YFASTEKSNI IRGWIFGTTL DSKTQSLLIV  120
NNATNVVIKV CEFQFCNDPF LGVYYHKNNK SWMESEFRVY SSANNCTFEY VSQPFLMDLE  180
GKQGNFKNLR EFVFKNIDGY FKIYSKHTPI NLVRDLPQGF SALEPLVDLP IGINITRFQT  240
LLALHRSYLT PGDSSSGWTA GAAAYYVGYL QPRTFLLKYN ENGTITDAVD CALDPLSETK  300
CTLKSFTVEK GIYQTSNFRV QPTESIVRFP NITNLCPFGE VFNATRFASV YAWNRKRISN  360
CVADYSVLYN SASFSTFKCY GVSPTKLNDL CFTNVYADSF VIRGDEVRQI APGQTGKIAD  420
YNYKLPDDFT GCVIAWNSNN LDSKVGGNYN YLYRLFRKSN LKPFERDIST EIYQAGSTPC  480
NGVEGFNCYF PLQSYGFQPT NGVGYQPYRV VVLSFELLHA PATVCGPKKS TNLVKNKCVN  540
FNFNGLTGTG VLTESNKKFL PFQQFGRDIA DTTDAVRDPQ TLEILDITPC SFGGVSVITP  600
GTNTSNQVAV LYQDVNCTEV PVAIHADQLT PTWRVYSTGS NVFQTRAGCL IGAEHVNNSY  660
ECDIPIGAGI CASYQTQTNS PGGSGSVASQ SIIAYTMSLG AENSVAYSNN SIAIPTNFTI  720
SVTTEILPVS MTKTSVDCTM YICGDSTECS NLLLQYGSFC TQLNRALTGI AVEQDKNTQE  780
VFAQVKQIYK TPPIKDFGGF NFSQILPDPS KPSKRSFIED LLFNKVTLAD AGFIKQYGDC  840
LGDIAARDLI CAQKFNGLTV LPPLLTDEMI AQYTSALLAG TITSGWTFGA GAALQIPFAM  900
QMAYRFNGIG VTQNVLYENQ KLIANQFNSA IGKIQDSLSS TASALGKLQD VVNQNAQALN  960
TLVKQLSSNF GAISSVLNDI LSRLDPPEAE VQIDRLITGR LQSLQTYVTQ QLIRAAEIRA 1020
SANLAATKMS ECVLGQSKRV DFCGKGYHLM SFPQSAPHGV VFLHVTYVPA QEKNFTTAPA 1080
ICHDGKAHFP REGVFVSNGT HWFVTQRNFY EPQIITTDNT FVSGNCDVVI GIVNNTVYDP 1140
LQPELDSFKE ELDKYFKNHT SPDVDLGDIS GINASVVNIQ KEIDRLNEVA KNLNESLIDL 1200
QELGKYEQGS GGYIPEAPRD GQAYVRKDGE WVLLSTFLGG SHHHHHH             1247
```

Q = Pyroglutamate, Mature N-terminus   N = Canonical N-Glycosylation Sequon Asparagine   *Italics* = "Tag"
~~Strikethrough~~ = Signal Peptide

B

100 nm

C

MWM   Spike   ACE2   MWM
kDa   kDa
250   250
150   150
100   100
75   75
                             50
50   37
37

D

Peptide (Pep): Q(Pyro-Glu)C(Carbamidomethyl)VNLTTR
MH^{2+} = m/z 1311.5477
ΔM = 0.98 ppm