Dense time-course gene expression profiling of the *Drosophila melanogaster* innate immune response

Florencia Schlamp[1], Sofie Y. N. Delbare[2], Angela M. Early[1], Martin T. Wells[2], Sumanta Basu[2], Andrew G. Clark[1,2]

[1] Molecular Biology and Genetics, Cornell University, Ithaca NY, United States

[2] Statistics and Data Science, Cornell University, Ithaca NY, United States

Corresponding authors: Florencia Schlamp (mfs97@cornell.edu), Andrew G. Clark (ac347@cornell.edu), Sumanta Basu (sb2457@cornell.edu).

ABSTRACT

Background

Immune responses need to be initiated rapidly, and maintained as needed, to prevent

establishment and growth of infections. At the same time, resources need to be balanced with

other physiological processes. On the level of transcription, studies have shown that this

balancing act is reflected in tight control of the initiation kinetics and shutdown dynamics of

specific immune genes.

Results

To investigate genome-wide expression dynamics and trade-offs after infection at a high

temporal resolution, we performed an RNA-seq time course on *D. melanogaster* with 20 time

points post Imd stimulation. A combination of methods, including spline fitting, cluster analysis,

and Granger causality inference, allowed detailed dissection of expression profiles, lead-lag

interactions, and functional annotation of genes through guilt-by-association. We identified Imd-

responsive genes and co-expressed, less well characterized genes, with an immediate-early

response and sustained up-regulation up to five days after stimulation. In contrast, stress

response and Toll-responsive genes, among which were Bomanins, demonstrated early and

transient responses. We further observed a strong trade-off with metabolic genes, which

strikingly recovered to pre-infection levels before the immune response was fully resolved.

Conclusions

This high-dimensional dataset enabled the comprehensive study of immune response dynamics

through the parallel application of multiple temporal data analysis methods. The well annotated

data set should also serve as a useful resource for further investigation of the *D. melanogaster*

innate immune response, and for the development of methods for analysis of a post-stress

transcriptional response time-series at whole-genome scale.

Keywords

Drosophila melanogaster, immune response, time course RNA-seq, Granger causality

BACKGROUND

Upon microbial infection, *Drosophila* launch rapid and efficient immune responses that are crucial to survival. However, immune responses are energetically costly (1) because they draw resources from other physiological processes (2, 3) such as metabolism, reproduction, and environmental stress responses. An excessive or overly prolonged immune response can lead to metabolic dysregulation, causing wasting in mammals and flies (4). Furthermore, it has been shown that allocating resources to the immune system reduces resources for mating (5, 6), and the opposite is also true, where mating reduces survivorship after infection and decreases resistance to infection (7-9). This represents a trade-off where limited resources need to be allocated to either the immune response or reproduction (10). Therefore, we expect that natural selection will act to tune the immune response to strike a balance between the advantage of a rapid and robust ability to fight infection, and the costly side-effects of an over-prolonged immune response. This tuning is likely to be mediated through a series of regulatory and feedback properties of the immune system of the fly.

While gene expression has been examined at several time points after infection in *Drosophila* (11-13), the dynamics of this immune response have not yet been studied with high temporal resolution. A high-resolution time-course analysis can help profile with more certainty the types of expression dynamics that different genes and pathways undergo after infection. Dense and extended time-course sampling of gene expression of the immune response can allow us to distinguish between transient and sustained expression patterns, where expression of genes with a transient response to perturbation will return back to normal after a certain period of time, while expression of genes with a sustained response will remain at a different level of expression compared to pre-perturbation levels. This kind of temporal profiling of the

immune response, coupled with computational modeling of gene interaction networks, can also

suggest candidates to examine for possible interactions and trade-offs between the immune

response and other physiological processes.

Statistical analysis of such high-dimensional longitudinal time-course omics data is not

straightforward. While the problems of detecting differentially expressed (DE) genes and

inferring gene interaction networks from gene expression data are common in genomics,

computational methods have focused primarily on cross-section rather than time-course data.

Most popular methods to analyze static RNA-seq data — such as edgeR (14) or DESeq2 (15)

— are not ideal for dealing with time-course RNA-seq data since they do not directly model the

correlation of genes between successive time points (16, 17). Smooth polynomial or spline

based models of temporal dependence in gene expression, such as those employed in Limma-

Voom (18) and maSigPro (19, 20), can fail to capture early impulses in stress response

situations, as we highlight in this paper. Also, joint network inference of temporal associations

among many genes requires tackling high-dimensionality, an aspect that has not received much

attention in the literature. Because there is not one consensus method for the analysis of time-

course RNA-seq data, it is important to ensure robustness of findings across different types of

computational modeling techniques.

In this study, we performed a dense time-course RNA-seq analysis of the *Drosophila*

transcriptional response to commercial *E. coli*-derived crude lipopolysaccharide (LPS), which

activates the Imd pathway (21), to better understand the dynamics of activation and resolution

of the innate immune response. Flies were sampled over 5 days generating a total of 20 time

points post-LPS injection with an additional time point pre-injection as a baseline control. We

analyzed the resulting longitudinal RNA-seq dataset using a broad range of statistical methods,

including a cross-sectional and a dynamic method for differential expression (DE), clustering,

and multivariate Granger causality (22), a method to investigate lead-lag relationships among

DE genes. We found that commercial LPS exposure has a major impact on the expression of

not only immune genes, but also genes involved in metabolism and replication stress.

Clustering analysis showed that both the onset and persistence of expression changes varied

across these DE genes. Clustering analysis further suggested a role in the immune response

and circadian rhythm for several previously uncharacterized genes. Finally, throughout our

analyses we observed a theme of interplay and trade-off between the immune response and

metabolism.

RESULTS

High-resolution profiling of gene expression after Imd stimulation

To generate a full transcriptional profile of gene expression dynamics in *Drosophila*

melanogaster after Imd stimulation, we injected adult male flies with commercial *E. coli*-derived

crude lipopolysaccharide (LPS). While pure *E.coli* LPS does not induce an immune response in

Drosophila, the peptidoglycan contamination in crude LPS preparations consistently activates

the Imd pathway (21). This was also confirmed using qPCR (see **Materials and Methods**). In

the remaining text, we refer to this treatment as "Imd stimulation". Using commercial LPS

instead of living bacteria gives the advantage of avoiding the confounding effects of a growing

and changing population of pathogens, and of the mechanisms the bacteria use to circumvent

immune responses (23).

Flies were sampled in duplicate for a total of 21 time points throughout the course of five

days, which includes an uninfected un-injected baseline sample as control at time zero, and 20

time points after injection. Since this is a perturbation-response experiment, denser sampling

occurred at early time points (16), with the first 13 time points taken within the first 24 hr (1, 2, 3,

4, 5, 6, 8, 10, 12, 14, 16, 20, and 24 hr). Sampling is also essential at later time points to know

how long it takes to return to baseline, and to differentiate between transient and sustained

responses (16). For this reason, sampling continued until day 5 after Imd stimulation, although

more sparsely (30, 36, 42, 48, 72, 96, 120 h) (**Figure 1A**). For this dataset we obtained 41 high-

quality libraries with an average of 23.5 million mapped reads per sample. After normalization of

libraries, only genes with more than 5 counts in at least 2 samples were kept, leaving 12,657

genes for further analysis.

Principal components analysis (PCA) on the 500 genes with highest row variance across

all time points revealed a horseshoe temporal trend, with the control samples clustering in the

middle, and the post-injection time points following a horseshoe-shaped track, consistent with a

pattern of many genes displaying a coordinated change over the five-day interval (**Figure 1B**).

This type of "horseshoe" or arch temporal trend in PCA has been seen in other time-series

experiments (18, 24-26), and is commonly seen in spatial population genetic variation (27) and

in ecological gradient data that varies in a non-linear manner (28). PC1, PC2, and PC3 captured

35, 15, and 14.5% of the variance in gene expression respectively, and the first six PCs account

for over 80% of the total variance in the data.

Proper normalization of the data was assessed by confirming the behavior of known

Drosophila housekeeping genes across time (Qiagen Housekeeping Genes RT^2 Profiler PCR

Array and (29)). As expected, housekeeping genes showed little change across time (**Figure**

S1A). The success of the Imd stimulation was confirmed by the immediate up-regulation of

known immune response genes within the first time points (**Figure S1B**).

A

B

Figure 1. **Transcriptional profiling of *Drosophila* immune response. (A**) Timeline of 21 time points,

including un-infected un-injected sample as control at time 0. Sampling was denser in the first 24 hr and

continued — although more sparsely — until day 5 (120 h). (**B**) Principal component analysis (PCA) of

the top 500 genes with highest row variance across all time points shows a coordinated change of gene

expression over five days. Both replicates are shown for all samples except for the time point at 3 h,

where one replicate was excluded from the analysis during RNA-seq data processing. The two samples

in blue clustering in the middle (marked with grey dashed circle) correspond to the control time point (0 h).

All other time points from 1 to 120 hr show a horseshoe temporal pattern around the controls.

Spline modeling and pairwise comparisons identify 951 genes that are differentially expressed over time following Imd stimulation

To identify genes whose expression levels were significantly altered across the time

course, we employed two methods. First, we used gene-wise linear models to fit cubic splines

with time, on both the first 8 hr and first 48 hr after commercial LPS exposure. Second, because

we noticed that certain expression patterns were not adequately described using cubic splines

(as discussed below), we also characterized the temporal patterns of expression by estimating

the differential expression of every gene at each time point, from 1 to 48 hr, compared to the un-

infected un-injected control samples at time zero (also referred to as baseline).

Cubic spline fits identified a total of 411 DE genes, based on a 5% False Discovery Rate

(FDR) using the Benjamini-Hochberg method (30) (**Table S1**). Of these 411 genes, 31 genes

were detected only using short spline fits on the first 8 hr post-injection. Long spline fits on the

first 48 hr post-injection identified 363 genes, and 17 genes were identified using both short and

long spline fits **(Figure 2A)**. Long spline fits excelled at identifying gradual changes and global

patterns, such as the ones shown by genes *Gale* and *Galk* (**Figure 2B**; note that these genes

are mentioned here only to illustrate the spline fit to expression dynamics, and not for their

biology). However, long spline fits failed to detect early impulse patterns, such as those

observed in the known immune response genes *AttA* and *DptB* (**Figure 2C**), which were better

captured by short spline fits on the first 8 hr post-injection. Still, even short spline fits failed to

identify additional known immune genes with early impulse patterns, such as *CecC and CecB*.

We also fit third degree polynomials using the R package maSigPro (19, 20). This approach

identified many DE genes that had been selected using spline modeling (**Figure S2**), but

similarly failed to adequately describe early impulse patterns. Based on these observations, we

also used pairwise comparisons to identify additional DE genes whose trajectories were not well

described using cubic splines. Pairwise comparisons identified 729 DE genes that were

significantly (FDR < 0.05) up- or down-regulated by a $\log_2$ fold change ($\log_2$FC) of at least 1 (i.e.:

2 times higher than baseline) in at least one time point throughout the first 48 hr after injections

(**Table S1**). Within this gene set, there were 214 genes that were up- or down-regulated by at

least 2 $\log_2$FC (4 times higher than baseline), in at least one time interval after injection (**Figure**

S3, **Table S1**). Of these 214 genes, 91 "core" DE genes underwent at least a 2 $\log_2$FC in

expression in at least two time intervals after injection, with a more stringent FDR < 0.01 (**Figure**

3A). Among the most strongly induced genes were known immune genes *DptB, AttC, Mtk, Dro,*

DptA and *edin* (bottom of **Figure 3A** and **Figure S4A**). These genes underwent an expression

change of approximately 5 $\log_2$FC (32 times higher than baseline) and remained elevated up

until 48 hr after Imd stimulation. Further investigation of the 91 core genes showed that the

number of up-regulated genes was much higher than the number of down-regulated genes

across all time points (**Figure 3B**). Eleven of the up-regulated genes at each time point were

known immune genes, as identified by a list of immune genes curated in Early *et al.* (31). Within

these 91 core DE genes, we also found circadian rhythm genes *period* (*per*), *timeless* (*tim*),

takeout (*to*), and *vrille* (*vri*), which when plotted against time exhibit the classic 24 hr periodic

expression of the circadian rhythm (**Figure S4B**).

Of a total of 951 DE genes, 189 genes were identified as differentially expressed using

both pairwise methods and spline modeling, but 762 out of 951 genes were identified using only

one of these methods, indicating the importance of using complementary methods for the

analysis of time course RNA-seq data (**Figure 2D)**.

Figure 2. Identification of time-dependent genes. (**A**) Genes that significantly change in expression across time according to spline analysis in the first 8 hr (yellow) vs 48 hr (blue). (**B**) Spline modeling of genes *Galk* and *Gale* when using first 48 hr (blue) and first 8 hr (yellow) compared to the pattern of normalized counts (green). Spline modeling over 8 hr misses the main change in pattern. (**C**) Spline modeling of two immune genes (*AttA* and *DptB*) when using first 48 hr (blue) and first 8 hr (yellow) compared to the pattern of normalized counts (green). Spline modeling over 48 hr smooths out the early impulse signal, drastically changing the inferred expression at time zero (indicated with arrows). (**D**) Comparing results from spline analysis (over 48 hr in blue and over 8 hr in yellow) vs. results from differential expression analysis (|$\log_2$FC| > 1 in green and |$\log_2$FC| > 2 in orange) at FDR < 0.05.

Figure 3. **Dynamics and functions of genes with changing expression patterns over time.** (**A**)
Heatmap of gene expression changes. Up-regulated genes are shown in orange, down-regulated genes
are shown in purple. These 91 genes were selected based on FDR correction of 0.01 and a |log$_2$FC| > 2
in at least two time points across 48 hr. The genes were ordered using Euclidean distance. (**B**) Number of
significantly up- and down-regulated genes, from the core 91 DE genes, at each timepoint (in red and

218   blue, correspondingly). Known immune genes are shaded over red. No down-regulated immune genes

219   were observed.

Gene Ontology and Gene Set Analysis demonstrate a divergence in expression between immune and metabolic processes after Imd stimulation

To understand the biological functions of genes whose expression is influenced by Imd

stimulation, we performed both a Gene Ontology (GO) and Gene Set Analysis. GO analysis is a

useful tool to illustrate the functions of genes with significant differential expression over time, in

this case 951 DE genes selected using spline fitting and/or pairwise contrasts. However,

focusing only on the top-scoring genes can lead to missing biologically relevant signals from

genes with modest expression changes. Furthermore, GO analysis does not take into account

expression changes over time. Both of these limitations are addressed by Gene Set Analysis,

which searches for enriched pathways (Gene Sets) across all 12,657 genes in the dataset,

guided by their $\log_2$ fold changes for all available time points.

GO analysis of the 951 DE genes using PANTHER identified a significant (FDR < 0.05)

overrepresentation of GO terms related to the immune and stress response, carbohydrate,

carboxylic acid and lipid metabolism, and proteolysis. Immune response related genes included

Attacins (*AttA*, *AttB*, *AttC*), Diptericins (*DptA*, *DptB*), Cecropins (*CecB*, *CecC*), Bomanins

(*BomS1, BomS2, BomS3, BomBc1*), genes encoding Daisho peptides (*Dso1, Dso2*), *IMPPP*

(also called *IM10*), *Drosocin* (*Dro*), *Drosomycin* and Drosomycin-like genes (*Drs*, *Drsl1*, *Drsl2*,

Drsl3), *Metchnikowin* (*Mtk*), Peptidoglycan Recognition Proteins (*PGRP-SB1*, *PGRP-SD*),

Diedel, *Relish* (*Rel*) and *elevated during infection* (*edin*), among others. DE genes known to

respond to stress included Turandots (*TotA*, *TotC, TotM*) and Heat Shock proteins (*Hsp70Aa*,

Hsp70Ab, *Hsp70Ba*, *Hsp70Bb*, *Hsp70Bbb*, *Hsp70Bc*).

Of the 951 DE genes, we identified 20 genes that encode known or putative transcription

factors, based on the FlyTF database (32) (**Table S2**). Seven of these twenty genes have a fast

impulse of up-regulation, reaching their maximum expression in the first two hours following Imd

stimulation (*Rel*, *Dif*, *CrebA*, *luna*, *Ets21C*, *Hr38* and *stripe;* **Figure 4A**). *Rel* and *Dif* encode

downstream components of the Imd and Toll pathways respectively, both involved in the

activation of the immune response (33-36). Of these two transcription factors, *Rel* undergoes

the strongest up-regulation, consistent with activation of the Imd pathway by Gram-negative

peptidoglycan in commercial LPS, and remains up-regulated at 1 $\log_2$FC even after 48 hr.

Ets21C encodes a stress-inducible transcription factor, and *Hr38* and *stripe* are the two most

robust activity-regulated genes (ARGs, defined as genes that are rapidly induced upon

stimulation of neurons, mostly within an hour) in *Drosophila* (37). Three genes encoding

transcription factors had oscillating expression patterns over time and are involved in the

regulation of the circadian clock (*vri*, *clk*, *Pdp1;* **Figure S5A**; (38, 39)). One gene of interest,

p53, involved in the response to genotoxic stress (**Figure S5B**; (40)), reached its maximum up-

regulation later, at 6 hr after injection.

Overall, the GO analysis indicates that the flies manifest a robust immune response, as

the gene expression changes are consistent with known expression profiles of immune

response deployment in *Drosophila* (11, 12). In addition, the GO analysis demonstrates that the

response to Imd stimulation also affects metabolic homeostasis.

Supporting these results, Gene Set analysis across all 12,657 genes and all time points

showed that the top up-regulated pathways were all related to immune response, defense

response to bacteria, and peptidoglycan functions (**Figure 4B**). Within these we found pathways

related to defense response against both Gram-negative and Gram-positive bacteria. While the

commercial LPS used for injections is derived from the outer membrane of Gram-negative

bacteria, which activates the Imd pathway, the injections themselves also result in septic injury,

which is known to activate both Gram-positive and Gram-negative immune pathways (Toll and

Imd pathways correspondingly) (41). Among down-regulated pathways we found many

metabolism-related functions. Three of these pathways (glycogen metabolic process,

triglyceride biosynthetic process, and gluconeogenesis) are highlighted in **Figure 4C-D** and **S5**.

The glycogen pathway down-regulation pattern was driven by genes *Fatty acid synthase 1*

(*FASN1*), and *UGP*, which encodes a UTP--glucose-1-phosphate uridylyltransferase (**Figure**

4C). Down-regulation of the triglyceride pathway was driven by *FASN1* and *minotaur* (*mino*), a

glycerol-3-phosphate 1-O-acyltransferase (**Figure 4D**). Finally, the gluconeogenesis pathway

down-regulation was driven by *fructose-1,6-bisphosphatase* (*fbp*), a rate limiting enzyme for

gluconeogenesis (42) (**Figure S6**). These metabolic genes reached their lowest expression

within the first 6 hr after Imd stimulation, and mostly recovered to baseline levels by hours 12-

24.

Figure 4. **Dynamics and functions of genes with changing expression patterns over time.** (**A**) Temporal

dynamics of DE transcription factors show that many transcription factors reach their maximal expression 1-2 hr after

Imd stimulation. **(B)** Heatmap showing most up- and down-regulated pathways (orange and purple respectively)

through the first 48 hr post-injections (absolute pathway score > 2.5 and *P*-value < 0.05 in at least one time point). (**C-**

D) **Gene Set Analysis identifies up- and down-regulated pathways.** Selected significantly down-regulated

metabolic pathways with corresponding gene memberships.

Clustering of temporal profiles highlights differences in the initiation and shutdown of immune and metabolic genes and demonstrates a regular rhythm of circadian clock genes

GO and Gene Set Analysis illuminated functions of genes that respond to Imd stimulation,

and indicated a trade-off between immune and metabolic processes. However, both GO and

Gene Set Analysis are based on prior knowledge of gene function. Clustering of genes based

only on their expression profiles is not influenced by prior annotations. Such an unbiased

approach can thus identify responses of poorly annotated genes. In addition, clustering can

illustrate how gene expression trajectories differ over time. We performed three analyses to

characterize temporal profiles. First, we performed hierarchical clustering based on Pearson

correlation on a set of 551 predominant time-dependent genes to identify major expression

patterns over time. These 551 genes included the 411 genes identified using spline modeling,

and 214 genes with at least a 2 $\log_2$ fold change in expression as identified using pairwise

comparisons (**Table S1**). Second, we performed clustering based on autocorrelation on these

551 genes. As opposed to Pearson correlation or Euclidean distance, an autocorrelation

function takes the ordering of time points into account, allowing us to identify more detailed

characteristics of gene expression profiles in time series. Third, because circadian rhythm

genes were not apparent in the clusters identified using the previous methods, but were

expected to be present in our dataset, we used the R package *JTK_Cycle* (43) to identify genes

with 24 hr cycling patterns among all genes in the dataset. We were interested in these patterns

since the circadian clock is known to regulate the expression of immune genes (44), and in turn,

infections are known to influence the flies' circadian rhythm (45).

First, expression profiles of the 551 predominant time-dependent genes fell into four main

hierarchical clusters (**Figure 5A**). Clusters 1 and 2 both had a strong increase in expression

after Imd stimulation (**Figure 5B**). Cluster 1 had a more immediate increase in expression,

reaching a maximum within the first 2 hr. Cluster 2, on the other hand, reached a maximum

expression later, at around 9 h. Cluster 1 showed significant enrichment of GO terms for

immune and stress response related processes, and contained Attacins and Cecropins, as well

as Heat Shock protein family genes. Cluster 2 was enriched for GO terms for abiotic stimulus

response, and contained among others *Bomanins*, *Daisho* genes, as well as genes from the

Turandot family (**Figure 5C)**. Clusters 3 and 4 were characterized by an initial decrease in

expression followed by an increase after 3 hr and 6 hr respectively (**Figure 5B**), with cluster 4

showing a stronger decrease in expression in the early hours after Imd stimulation. These

clusters had a significant enrichment of GO terms for biosynthetic, catabolic, and metabolic

processes (**Figure 5C**), and their down-regulation again indicates a trade-off between

metabolism and the initiation of an immune response.

Our second clustering analysis based on autocorrelation revealed additional differences

regarding the initiation and resolution of gene expression after Imd stimulation. First, we

identified a cluster of genes with an immediate and sustained up-regulation. This cluster was

characterized by a strong early induction with an up-regulation of ~2.5 to 6 log_2FC within the first

hour (6 to 64 times higher than baseline), reaching a maximum of 6 to 8.5 log_2FC (64 to 362

times higher than baseline), and maintaining persistent up-regulation of ~2.5 to 6 log_2FC

throughout 5 days (**Figure 6A**). This cluster contained canonical immune response genes

known to be activated downstream of Imd, such as *AttA*, *AttB*, *AttC*, *DptA*, *DptB*, *Dro*, *edin*, *Mtk*,

PGRP-SB1, *PGRP-SD.* The cluster further contained *IBIN* (*Induced by Infection*), whose exact

mode of action is unknown, but whose up-regulation stimulates starch catabolism as part of an

immune-induced metabolic switch, likely to make free glucose available to circulating immune

cells (46). Finally, this immediate-response cluster also contained *Mtk-like, CG43920* and

CG45045, which are less characterized transcripts known to be up-regulated after bacterial

infection (47).

335   Autocorrelation-based analysis also identified clusters of genes with transient responses to

336 infection (**Figure 6B-C**). One of these clusters was composed of Bomanins (*BomS1, BomS2,*

337 *BomS3, BomBc1*), *Daisho* genes (*Dso1, Dso2*), *IMPPP* and *CG33470* (**Figure 6B**). The

338 *Bomanins* are located in the 55C4 region of chromosome 2R, confer resistance against certain

339 bacteria and fungi and are regulated by Toll signaling (48). *Dso1 and Dso2* were also reported

340 to be regulated by Toll signaling and aid defense against specific filamentous fungi (49).

341 *CG33470* is an uncharacterized transcript located 3.3 kb downstream of *IMPPP* and might

342 belong to the same open reading frame, as both are sometimes referred to as *IM10* (50), and

343 show nearly identical gene counts in our dataset. This cluster was characterized by an early

344 induction (but not as immediate as the cluster containing known Imd-responsive genes in

345 **Figure 6A**) of ~2.5 to 3.5 $\log_2$FC (6 to 11 times higher than baseline) within the first two hours,

346 reaching a max of ~2.5 to 5 $\log_2$FC, and returning to a steady state after 3-5 days. Thus,

347 clustering analysis identified effector immune genes partitioned by Imd vs Toll signaling: Imd-

348 regulated genes showed an immediate early sustained up-regulation even after 5 days (**Figure**

349 **6A**), while Toll-regulated *Bomanins* and *Daisho* genes had an early up-regulation that eventually

350 returned to steady state levels (**Figure 6B**).

351   A final cluster illustrated a more complex expression pattern: many genes in this cluster

352 were down-regulated immediately, 1-2 hr after injection, after which they were up-regulated,

353 reaching their maximum expression after 8-12 hr, followed by a return to baseline after 2-3 days

354 (**Figure 6C**). This cluster was composed of genes from the stress-induced Turandot family

355 (*TotA, TotB, TotC, and TotX*) (51) as well as *Diedel*, *Grik*, *lectin-24A*, *NimB3*, *BomT2*, *CG11459*,

356 and *CG30287*. *Diedel* encodes an immunomodulatory cytokine known to down-regulate the Imd

357 pathway. *Grik* encodes a glutamate receptor, and *Lectin-24A* encodes a pattern recognition

358 receptor that mediates pathogen encapsulation by hemocytes (52). *Lectin-24A* has been shown

359 to be down-regulated in the first 2 hr following septic injury and then up-regulated 9 hr after (53),

360 consistent with the pattern we see in our data. *NimB3* is part of the Nimrod gene family, which is

involved in phagocytosis (54). *BomT2* is part of the Bomanin gene family (like *BomS1*, *BomS2*,

BomS3 and *BomBc1*, expressed in the previous cluster, **Figure 6B**). *CG11459* encodes a

predicted cathepsin-like peptidase induced by bacterial infection and injury (55). *CG30287*

encodes a predicted serine protease, a class of proteins that plays roles in immune response

proteolytic cascades (56).

Finally, using JTK_cycle, we identified 22 periodic genes with a 24 hr cycle, using a cutoff

of Benjamini-Hochberg corrected *Q*-value < 0.05 and amplitude > 0.5 (**Figure 7**). Among them

were four well characterized circadian genes, suggesting that their periodicity was not affected

by Imd stimulation: *period* (*per*), *takeout* (*to*), *vrille* (*vri*), and *PAR-domain protein 1* (*Pdp1*), as

well as eight genes which do not have assigned circadian functions but have evidence of cyclic

behavior in previous literature (**Table 1**), and 10 genes, of which 8 are uncharacterized, that

have not yet been reported to have cyclic expression outside this study (**Table 1**; *CG10560*,

Sgroppino, *CG15253*, *CG15254*, *CG18493*, *CG31321*, *CG33511*, *CG34134*, *CG42329, salt*).

Overall, the combination of clustering methods augmented by GO analysis allowed us to

identify strong temporal patterns that correspond to early and late induction of immune

processes, as well as both transient and sustained responses to infection, which point to a

trade-off between the immune response and metabolism. We found that genes that share

functions often have similar temporal expression patterns, suggesting co-regulation. This

observation further allowed us to assign putative functions to previously uncharacterized genes

that cluster together with well-studied genes.

Figure 5. Global dynamics of time-dependent genes show divergent patterns of expression. (**A**) Heatmap of the 551 most predominant time-dependent genes, identified by spline modeling over 48 and 8 hr (FDR < 0.05) and pairwise differential expression (with |log₂FC| > 1 and FDR < 0.05). Hierarchical clustering of the genes shows four main clusters characterized by time points in which the genes reach maximum and minimum expression across time. Z-score values of each gene are shown from dark purple (minimum expression across time) to dark orange (maximum expression across time). (**B**) Mean patterns of expression across time for genes within each of the four main clusters during the first 24 hr, displayed by their centered and scaled normalized counts. (**C**) Significant Gene Ontology terms (FDR < 0.05) for over-represented Biological Processes at each cluster.

Figure 6. Clusters of genes identified using autocorrelation. Network nodes represent genes; network edges

represent the distance between gene autocorrelations, based on ACF analysis using TSclust. (**A**) A cluster containing

multiple Imd-responsive genes shows sustained expression after Imd stimulation throughout 5 days (120 h). (**B**) Toll-

responsive genes show a transient response to Imd stimulation. (**C**) Other stress response genes return to steady

state by day 5 (120 h) post Imd stimulation.

397 **Figure 7. Top 22 genes identified by JTK_Cycle show 24 hr temporal cycling.**

405 **Table 1. Evidence of cyclic behavior for top genes identified by JTK_Cycle.**

Gene	Evidence of cyclic behavior	Source
Carbonic anhydrase 5 (*CAH5*)	upregulated DE gene in response to light stimulation	Adewoye *et al.* 2015 (59)
cypher (*cyr*)	downregulated DE gene in response to light stimulation	
CG17127	photoperiodic	Pegoraro and Tauber 2018 (60)
	rhythmically expressed in constant dark conditions	Ueda *et al.* 2002 (61)
Alcohol dehydrogenase *CG4842 and Cuticular protein 72Ec (Cpr72Ec)*	downregulated in the retina after *heme oxygenase* (*ho*) silencing in photoreceptor cells	Damulewicz *et al.* 2019 (62)
CG8170	cycling mRNA	Huang *et al.* 2013 (63)
Juvenile hormone esterase (*Jhe*)	cyclic hemolymph activity	Zhao and Zera 2004 (64)
Sodium-dependent multivitamin transporter (*Smvt*)	modulated by the circadian rhythm in mice	He *et al.* 2016 (65)
CG10560, Sgroppino (*Sgp*), *CG15253, CG15254, CG18493, CG31321, CG33511, CG34134, CG42329,* and sodium/solute symporter *salty dog* (*salt*)	none reported	

Gene interaction modeling of lead-lag patterns using Granger causality

Clustering methods based on a single gene's autocorrelation or cyclicity patterns can detect genes with similar expression profiles. However, these methods are not suitable for seeking causal relationships between genes that manifest in a lead-lag relationship, for instance when high expression of gene A results in a high expression of gene B shortly afterwards. Detecting such lead-lag patterns (**Figure 8A**) is a unique advantage of dense time-course experiments. Granger causality (GC), a statistical method popular in analysis of macroeconomic time series, provides an ideal framework for modeling such patterns and building directed networks among genes. The concept of GC is based on predictability. If the knowledge of the past of one time series improves the prediction of a second one, the first is said to be Granger causal (GC) for the second. *Bivariate* GC analysis between two genes A and B, as described above, does not account for possible confounding effects of other genes C, D, E which can also influence genes A and B (**Figure 8B**). *Multivariate* GC analysis alleviates this problem by explicitly accounting for the effects of the confounding genes by a joint modeling (57, 58), but does not account for high-dimensionality and consequently cannot jointly model hundreds of genes based on tens of data points. We used modern high-dimensional methods (viz LASSO (59) and de-biased LASSO (60, 61)) to address this problem and build lead-lag network models among 258 genes.

Figure 8. Diagram describing the process of constructing directed networks from Granger causality. (A)
Lagged correlated expression between two genes (Granger causality) leads to the construction of a directed edge
between two genes (nodes), which in turn is used to build directed lead-lag network models of putative interactions
among genes. Edges can be positive or negative, based on the sign of lead-lag correlation between the two genes.
(B) Bivariate associations are calculated between two genes at a time, while multivariate associations adjust for
potential indirect association from all other genes in the gene set.

We constructed directed GC edges and networks of putative interactions among a subset

of 258 genes (**Table S1**). These genes had a $|\log_2 FC| > 1$ (at least 2 times higher or lower than

baseline) across the time course and had available functional annotations. We performed

Granger causality analysis on sliding windows of 6 time points on the normalized counts of both

replicates using bivariate and multivariate methods (see **Materials and Methods**). We

investigated both positive and negative edges, reflecting positive and negative lagged

correlations between genes. The overall unfiltered GC network has a multitude of relationships

worth exploring, but limitations in the ability to distinguish different types of causality make

widespread conclusions from the network challenging. Here, we discuss several examples of

subnetworks which illustrate putative functional relationships among genes whose expression

changes in response to Imd stimulation.

Based on our interest in identifying trade-offs between biological processes in infected

animals, we first constructed a high-quality set of consistently significant GC edges of divergent

expression (negative edges). To this end we first filtered the subnetwork by (a) removing all

edges with a positive weight, (b) removing all nodes corresponding to cyclic genes identified

earlier through the JTK_Cycle method, (c) using only pairs of nodes with significant edges

(Benjamini Hochberg FDR < 0.05%) in at least 3 consecutive windows within the first 24 hr of

the time course. After filtering, the resulting high-quality GC network contained 51 nodes and 35

edges in 16 connected components (**Figure S7**). This network, by design, should include the

most interesting examples of divergent expression changes from our full dataset.

The largest connected component in this network (Component #1) is a multifunctional

chain of 6 genes, which connects the down-regulation of four metabolic genes with the up-

regulation of two genes that are involved in regulating proliferation and repair (**Figure 9A**). Two

of the metabolic genes, *Sorbitol dehydrogenase 1* (*Sodh-1*) and *UGP*, both lead the divergent

expression of *Claspin* (both 4 consecutive windows, 2 to 10 and 4 to 12, respectively) (**Figure

9C** & **S8A**). *Claspin* plays a role in DNA replication stress (62). It is known that there is an

interplay between host immune systems and replication stress (63). The immune system can

detect and respond to replication stress, which is an important feedback loop necessary to

remove defective cells (64). Furthermore, the activation of the immune response generates

reactive oxygen species (ROS) and reactive nitrogen species (RNS), and can promote chronic

inflammation, all of which can trigger DNA damage (65). *UGP* and *fbp* were identified earlier

during Gene Set Analysis to drive the down-regulation of metabolic pathways (**Figure 4B** and

4D), and in this cluster they are both negatively directed by *LpR2* (3 consecutive windows, 6 to

13) (**Figure 9D** and **S8B**). *LpR2* is a lipophorin receptor, known to regulate the innate immune

response by clearing serpin protease complexes from the hemolymph through endocytosis (66).

Lipophorin is a known humoral factor that contributes to clot formation (67, 68). Finally, *LpR2* is

also shown to negatively direct *juvenile hormone acid methyltransferase* (*jhamt*) (4 consecutive

windows, 1 to 9) (**Figure 9E**). JHAMT is an enzyme that activates juvenile hormone (JH)

precursors at the final step of the JH biosynthesis pathway in insects (69). JH is a known

hormonal immunosuppressor in *Drosophila* (70-72).

Interestingly, *Claspin* was identified to be part of the same pathway as *Orc1* in our

previous Gene Set Analysis, showing similar patterns and window of up-regulation (mitotic DNA

replication checkpoint pathway, **Figure S9**). In our network, *Orc1* is part of an isolated edge with

metabolic gene *ABGE* (Component #2, 4 consecutive windows, 4 to 12) (**Figure 9A** and **9E**).

These prioritized subnetwork components suggest an interplay between metabolic pathways

and other pathways such as proliferation and repair (**Figure 9B**), motivating follow-up studies to

determine which pathways might be regulating and trading off with each other in the hours

following Imd stimulation.

In addition to these purely negative edges, we detected highly significant positive and

negative edges among circadian rhythm genes. These included *cryptochrome* and *Smvt* (6

consecutive windows, 6 to 16) (**Figure S10A**), *vrille* and *takeout* (4 consecutive windows, 9 to

17) (**Figure S10B**), *period* and *takeout* (4 consecutive windows, 9 to 17) (**Figure S10C**), and

Smvt and *takeout* (4 consecutive windows, 9 to 17) (**Figure S10D**). *Smvt* is predicted to encode

a sodium-dependent multivitamin transporter, and *takeout* influences feeding behavior (73, 74).

Metabolic processes and feeding are known to be under circadian control (75). In addition, So *et*

al. (76) reported that *takeout* is regulated by the circadian clock, but with a phase shift relative to

period. This pattern is clearly visible in our dataset and was correctly identified using Granger

causality. This shows that Granger causality can be used to infer gene

dependencies/interactions using global gene expression behavior.

Finally, among genes connected only by positive edges, we identified an edge from

period, a regulator of the circadian clock (77, 78), to *Rhodopsin 5*, which encodes a G-protein-

coupled receptor involved in phototransduction (**Figure S11A**). *Rh5* mRNA levels are known to

demonstrate a cyclic pattern (79), indicating regulation by the circadian clock. We further

identified positive edges between genes that are likely co-regulated. These included edges

between up-regulated genes that respond to NF-κB signaling, such as edges from genes

encoding peptidoglycan recognition receptors *PGRP-SD* and *PGRP-SB1*, (regulated by Imd

signaling), to *DptB* and *AttC* (also regulated by Imd signaling) and to *BomS1*, *Dso1* and

BomBc1 (regulated by Toll signaling) (**Figure S11B-F**). We also observed edges between

down-regulated genes, such as from *Hao* (predicted to play a role in lactate oxidation) to *AGBE*

(a predicted hydrolase involved in glycogen synthesis) (**Figure S11G**). While the expression of

these genes likely responds to similar signals, the observed lags between these genes'

expression profiles suggest that there are differences in their transcriptional control, such as

regulation by cofactors, differences in promoter affinity for certain transcription factors, or other

variables that influence the rate of mRNA accumulation.

Figure 9. High-quality GC network components and their edges. (**A**) Components #1 and #2 from GC network (**Figure S4**). (**B**) Diagram summarizes interplay between main represented pathways on the selected components. (**C-F**) Selected edges from the components plotted against time. Significant windows colored in blue, non-significant colored in grey. Resulting overall consecutive windows are labeled in blue dashed rectangles. Individual windows represent 6 consecutive time points, but because time points are not at regular intervals, the windows have different time ranges, but identical numbers of samples.

DISCUSSION

We have produced a dense and high-quality time-course profiling of the *Drosophila*
524

transcriptome response after Imd stimulation through commercial LPS injection, using RNA-seq
525

sampling over 20 time points spanning 5 five days. This profiling provides a high-dimensional
526

dataset, which is available as a resource for the community. We analyzed this dataset using a
527

broad range of statistical methods, including Granger causality, to investigate lead-lag
528

relationships between genes. Because of the high dimensionality, it is not straightforward to
529

analyze as a time series, as illustrated by the partially distinct results of spline fitting and
530

pairwise comparisons. However, using a combination of analytical methods allowed us to
531

identify distinct patterns with high confidence, specifically responses to Imd stimulation with
532

divergent initiation and resolution dynamics, as well as cyclic patterns of gene expression, and
533

patterns of co-regulation and trade-offs. Below, we describe and discuss the main insights from
534

these analyses, as well as limitations and future steps.
535

Immune and stress response genes vary in their initiation and resolution
536
dynamics after Imd stimulation
537

Clusters of genes demonstrated distinct activation kinetics after stimulation of the
538

immune response. This phenomenon has been observed both in fly (11) and mammalian cells
539

(80), but as a result of the dense sampling, our dataset provides a highly detailed view of these
540

initiation dynamics. In addition, because we sampled up to five days post-injection, we could
541

also observe long-term responses to Imd stimulation.
542

First, a cluster of 13 genes showed the fastest up-regulation within the first 1-2 hr and
543

remained up-regulated during the entire five-day time course (**Figure 6A**). This cluster
544

contained 10 genes known to be regulated by Imd signaling (3 *Attacins*, 2 *Diptericins*, *Mtk*, *Dro*,
545

IBIN, *PGRP-SB1* and *PGRP-SD*), as well as *CG43920*, *CG45045 and Mtk-like.* Imd-regulation
546

of these 3 genes has not been experimentally validated, but their co-clustering pattern suggests

that they are regulated by the Imd pathway.

Second, a cluster containing Toll-regulated *Bomanins* and *Daisho* genes (81) reached its

highest point of expression at 5-8 hr and recovered to a baseline state after 2 to 5 days (**Figure**

6B). The later initiation of Toll-responsive genes relative to Imd-responsive genes is consistent

with Tanji *et al.* (82), who reported earlier peak expression of an *Attacin* and *Diptericin* (Imd-

responsive) versus *Drosomycin* (mostly Toll-responsive). The up-regulation of Toll-responsive

genes might be a response to the wounding that occurred during LPS injection, as Irving *et al.*

(83) reported that septic injury with Gram-negative *E. coli* induced responses to both Gram-

negative and Gram-positive bacteria, and Boutros *et al.* (11) reported that clean injury

experiments induced a set of genes that overlapped with those induced by a septic injury

experiment, but with a lower response magnitude, consistent with our observations (**Figure 6A-**

B; Imd-responsive genes reach a maximum of 6-8 $\log_2$FC while Toll-responsive genes reach a

maximum of 2.5-4.5 $\log_2$FC).

Third, stress response genes, among which were members of the Turandot family,

reached their highest point of expression at 10-12 hr, following a pattern of delayed response in

line with observations by Ekengren *et al.* (51). Similar to the Toll-regulated gene cluster, stress

response genes returned to a baseline state after 2 to 5 days (**Figure 6C**).

It is striking that the expression of Imd-regulated genes did not recover to pre-injection

levels even after five days. Since we stimulated the Imd pathway by injecting Gram-negative

peptidoglycan within commercial LPS, it is possible that the prolonged up-regulation of AMPs

and other Imd-regulated genes was due to remaining peptidoglycan in the flies. Alternatively, it

might be typical for these genes to be expressed at a higher level for a certain time even after

the peptidoglycan has been cleared. Troha *et al.* (47) also observed prolonged AMP up-

regulation after bacterial infection, even when levels of bacteria were below detection threshold.

On the other hand, Duneau *et al.* (84) reported that seven days after an initial infection with *E.*

coli, none of the surviving flies were completely free of bacteria - suggesting suppression rather

than clearance of the infection. These observations raise the question of whether one should

expect to see a return to the baseline gene expression levels. Rather than returning to a pre-

infection state, the prolonged up-regulation of Imd-responsive genes and the transcription factor

Rel (**Figure 4A**) might be required to prevent damage from dormant bacteria.

Initiation of the immune response coincides with a down-regulation of metabolic processes

Our dataset showed distinct global dynamics pointing to a divergence in expression

between immune and metabolic processes (both carbohydrate and lipid metabolism), with the

most up-regulated pathways related to immune and stress responses, and the most down-

regulated pathways related to metabolic functions (**Figure 5**).

Metabolic genes reached their maximum down-regulation at 5-8 hr (**Figure 4B-D, 5**).

FASN1, which showed the strongest down-regulation in both glycogen metabolic process and

triglyceride biosynthetic process (**Figure 4C-D**), is a lipogenic gene whose down-regulation

might indicate a need to have easily accessible nutrients instead of storing them. Indeed,

infections in mammals are known to induce adipose tissue lipolysis (85) and bacterial

peptidoglycan is a ligand that stimulates lipolysis as well (86). The gene with the strongest

down-regulation in the gluconeogenesis pathway was *fbp* (**Figure S6**), which codes for fructose-

1,6-bisphosphatase, the rate limiting enzyme for gluconeogenesis. This gene was significantly

down-regulated in a study that reported that *Listeria monocytogenes* infection in *Drosophila*

causes a decrease in energy stores, with reduced levels of triglycerides and glycogen (87).

Krejcova *et al.* (88) reported that an infection-induced switch to aerobic glycolysis within

macrophages coincides with a systemic depletion of glycogen stores and increased blood sugar

levels. The divergent dynamics detected in our dataset are thus in agreement with known

individual mechanisms characterized in the immune response.

598  We also observed that expression of metabolic genes and pathways recovered quickly to

599 a baseline state around 12-24 hr after Imd stimulation (**Figure 4B-D, 5**). The speedy recovery of

600 metabolic genes, despite sustained expression of Imd-regulated immune response genes,

601 suggests that the early stages of infection likely involve the greatest trade-offs, at least in

602 response to the commercial LPS injections. These dynamics might differ in flies infected with

603 live bacteria and might also differ depending on the strain of bacteria (47).

604  We further saw implications of functional interplays using the Granger Causal (GC)

605 network analysis. Main subnetwork components showed significant GC directional edges

606 between down-regulated metabolic genes (such as *Sodh-1*, *UGP*, *fbp*, and *AGBE*) and up-

607 regulated genes with cell proliferation and repair functions (*Claspin*, *LpR2*, and *Orc1*) (**Figure**

608 **9**). These results further suggest an underlying interplay between metabolic pathways and

609 proliferation and repair mechanisms such as regulation of DNA replication stress, endocytosis,

610 and clot formation. While functional genetics studies have demonstrated such trade-offs

611 previously (3, 89), our dataset reveals the extent and dynamics of these trade-offs on a

612 genome-wide scale.

613 Predicting function by association

614  Using clustering analysis, we identified several genes that lack a well-established function,

615 and that clustered tightly with well-studied genes. We can use this co-clustering to suggest

616 shared functions (90).

617  Temporal clustering analysis identified *Mtk-like, CG43920, IBIN* and *CG45045*, which

618 shared similar expression dynamics with Imd-regulated AMPs (**Figure 6A**). Suggesting AMP-

619 like functions to these less characterized genes can be supported by observations from

620 literature: All four genes were previously found to respond to infection (47, 55), and *Mtk-like* and

621 *CG43920* have been shown to encode small proteins predicted to be cationic (91), properties

622 shared by known AMPs (92). *IBIN* and *CG45045* were also predicted to physically interact with

antimicrobial peptide transcripts (91). On the other hand, Valanne *et al.* (46) found that *IBIN*

overexpression increased levels of hemocytes and hemolymph glucose, suggesting that *IBIN*

functions as a link between immunity and metabolism, instead of acting as an AMP itself.

The dense sampling nature of this time course allowed us to discern the clear cycling

patterns of differentially expressed genes such as *period*, *timeless*, *takeout*, *vrille*, and

cryptochrome, all of which have well-characterized circadian rhythm functions (38, 39, 76, 93,

94). Clustered with these genes, our analysis identified other genes that showcase cyclic

behavior but are not canonically circadian-associated genes. This includes eight genes which

do not have assigned circadian functions but do have some evidence of cyclic behavior in

previous literature. It also includes ten genes that had not been reported to exhibit any cyclic

expression before this study (**Table 1**). The identification of canonical circadian rhythm patterns

both validates our methods of data normalization and differential expression analysis, and

increases the certainty that we are accurately profiling novel temporal dynamics. It is important

to note, however, that proper validation of the cycling behavior of our novel cyclic genes should

be performed under normal *Drosophila* conditions, as we do not know whether Imd stimulation

affected their expression.

Overall, we were able to implicate these uncharacterized genes as potential members of

specific functional pathways due to the strong similarity of their expression dynamics. This is

impactful both in the functional implication of these genes, but also in demonstrating the

potential of this "guilt-by-association" method to assign putative function to other

uncharacterized genes through RNA expression time-course experiments.

Limitations and future steps

Limitations of our experimental design can be used as a guide to design future

experiments. This time-course design lacks time-matched controls to account for expression

changes associated with phenomena outside the Imd stimulation, such as aging. However, it is

still highly valuable to develop and improve methods for analyzing time-course transcriptional

data lacking time-matched controls, since these methods are needed to analyze processes

such as development, where such controls are inherently not possible.

This study also lacks a control for the wounding injury caused by the injection itself. As

discussed above, specific expression patterns such as those from the Toll-response genes

could be caused by the injection wounding rather than the Gram-negative peptidoglycan in

commercial LPS. Another important experimental design aspect for time series is choosing the

time frame within which sampling needs to occur, which can be difficult to establish in advance.

In our study Imd-regulated gene expression was sustained until five days after infection, and a

more prolonged sampling would have been necessary to determine the full duration of these

genes' up-regulation after Imd stimulation.

Further, our experiment sampled only males, while several previously published immune

response studies used females (e.g. (3, 21)). Overall, the observed gene expression changes

and trade-offs in our study agreed with functional and omics studies done using females. This

suggests that global gene expression dynamics after an Imd challenge are similar between

males and females, but future experiments would be needed to make direct comparisons. In

addition, multiple studies have reported a trade-off between immunity and reproduction, which

occurs in singly-mated females (9) and multiply mated males (95). Mating activity was outside

the scope of our design but can be included in future designs.

In our dataset, Granger causality analysis excelled at showcasing the relationships

between divergent gene pairs, but was overly sensitive to the extreme temporal correlation

between large groups of genes when analyzing positive edges. To avoid a prohibitively dense

network for analysis, we relied on heuristic network trimming criteria, which was effective, but is

likely not generalizable to other similar experiments. Developing co-integration methods that

take into account the specific bias found in high-dimensional RNA-seq datasets would provide a

more robust statistical analysis of the causal relationships observed in this type of data. The

biological interpretation of relationships identified using Granger causality also warrants further

investigation: Granger causality was successful at identifying what are likely the downstream

results of divergent regulation, and it was successful at identifying positive lead-lag relationships

between genes that likely respond to similar signals, but might differ in their exact (post-

)transcriptional control. However, these statistical causal relationships provide only hypotheses

that should be tested with direct experimental disruptions of a system to demonstrate biological

causality.

CONCLUSIONS

Our combination of analytical methods provides robust profiling of the innate immune

response in *Drosophila melanogaster* after Imd stimulation at the highest temporal resolution to

date, and serves as a proof of concept for high-density time-course RNA-seq analyses in other

systems. Further, it motivates innovation in computational and statistical methods for

longitudinal omics data, both to account for their inherent high-dimensionality and the complex

underlying architecture that contains both causal and spurious coordination. Specifically, the

development and application of multivariate Granger causality analysis highlights the potential

of time-course data to evaluate coordinated gene expression changes through lags and trade-

offs. While the immune response in *D. melanogaster* has been well studied, our research using

dense time-course gene expression data reveals genome-wide dynamic expression patterns at

higher temporal detail. Specifically, we reveal responses to Imd stimulation with divergent

initiation and resolution dynamics, cyclic patterns of gene expression, and patterns of co-

regulation and functional trade-offs, while also assigning putative gene functions to

uncharacterized genes through a temporal guilt-by-association method.

MATERIALS AND METHODS

Fly lines, injections, and sample collection

Male adult *Drosophila* of about 4 days old were sampled from an F1 cross from two

Drosophila melanogaster Genetic Reference Panel (DGRP) lines: line 379, which has shown to

have low bacterial resistance, and line 360, which has high bacterial resistance (96). Offspring

from these two DGRP lines were used to allow investigation of allele-specific expression after

Imd stimulation. Analysis indicated significant allele-specific differential expression across

multiple loci, including loci that contain immune genes, when aggregating data across all time

points (data not shown). This may reflect differential rates of transcription initiation, processivity,

RNA processing, or decay. However, allele-specific expression did not significantly change over

time after infection (data not shown). Because of the absence of time-dependent effects, this

analysis was not included in the manuscript.

Flies were kept on a 12:12 dark-light cycle, on standard yeast/glucose food. Flies were

injected in the abdomen with 9.2 μl of commercial lipopolysaccharide (LPS) (*Escherichia coli*

055:B5 Sigma) derived from the outer membrane of Gram-negative bacteria. Flies were injected

using a Nanoinjector (Nanoject II, catalog #3-000-204, Drummond), which allows high-

throughput fly injections with a constant injection volume. Injections were performed in the

abdomen, as it has been shown to be less detrimental to the fly compared to thorax injury (97).

Flies were sampled for a total of 21 time points throughout the course of five days, which

included an uninfected un-injected baseline sample as control at time zero, and 20 time points

after LPS injection (1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 20, 24, 30, 36, 42, 48, 72, 96, 120 hr). This

sampling was performed twice, using flies from the same stock, on two consecutive days.

Therefore, all time points have two replicates, giving a total of 42 samples. Since flies were

sampled from the same stock for both replicates, we consider the second replicate to control for

any effects of the injection technique. This sampling strategy was informed by experimental data

724   and theoretical analysis that show that under reasonable assumptions, sampling time points at

725   higher resolution is preferred over having more replicates (98), an important strategy to consider

726   when having a limited experimental budget.

727   During collection, a group of ~10 pooled flies corresponding to the sampled time point

728   were flash frozen in dry ice and stored at -80 C for later RNA extraction.

Experimental validation using qPCR

The Imd inducibility of commercial LPS was confirmed using qPCR. Adult male *Drosophila*

were injected with 9.2 µl or 40 µl of 1 mg/mL LPS and flash frozen at 8 and 24 hr for RNA

extraction. Uninfected un-injected flies were used as control. Each sampled time point consisted

of a group of ~10 pooled flies. Each sample had two replicates. Genes *AttA* and *DptB* were

measured to confirm Imd stimulation. Gene *Rp49* was used as a baseline for expression

normalization. Results showed a significant up-regulation of *AttA* and *DptB* at both volumes (9.2

µl and 40 µl) for both time points (8 and 24 hr). We decided to use 9.2 µl so as to cause the

least amount of disruption to flies during infections, while still eliciting an immune response.

RNA extraction, RNA sequencing, and quality control filtering

RNA extraction was performed using Trizol (Life Technologies) following the

manufacturer's instructions. cDNA libraries were prepared using the TruSeq RNA Sample

Preparation Kit (Illumina). RNA purity was assessed using a Nanodrop instrument. RNA

concentration was determined using a Qubit (Life Technologies) instrument. Sequencing was

performed on an Illumina Hi-Seq 2500, single-end, and a read length of 75 bp, at Cornell

Biotechnology Resource Center Genomics Facility.

Samples had an average of 24.8 M raw reads. Samples went through quality control using

FastQC (v0.11.5). Truseq adapter sequences were removed from any sample that showed any

level of adapter contamination using cutadapt (v1.14). Low quality bases in the beginning and

end of the reads were trimmed using fastx_trimmer (v0.0.13,

http://hannonlab.cshl.edu/fastx_toolkit/). Reads were mapped to the *Drosophila melanogaster*

genome (r6.17) using STAR (v2.5.2b). BAM files were generated using SAMtools (v1.3.2). Only

one sample (4B, at 3 hr) out of the original 42 failed to pass the quality thresholds, and all

subsequent analysis used the remaining 41 samples. An average of 92.97% reads per library

mapped uniquely to the *Drosophila melanogaster* genome. We ended up with an average of

23.4 million uniquely mapped reads per library.

Reads mapping to genes were counted using the R package *GenomicAlignments*. Genes

with zero counts across all samples were removed (923 genes out of 17,736). Samples were

normalized to library size. A "+1" count number was added to all genes before performing $\log_2$

transformation, to make sure values after transformation are finite, and stabilize the variance at

low expression end. After normalization and $\log_2$ transformation, only genes with more than 5

counts in at least 2 samples were kept (removing 4,156 genes). We ended up with 12,657

genes for downstream analysis.

A heatmap of the row Z-scores of normalized counts for all 12,657 genes indicated that

sample 6A (5 hr after commercial LPS injection) was an outlier for a subset of the 12,657 genes

(**Figure S12A**), even though sample 6A did not appear as an outlier using Principal Component

Analysis (see below and **Figure 1B**). We identified outlier genes in sample 6A by subtracting

replicate A row Z-scores of normalized counts from replicate B. While for most time points the

difference between samples A and B varied between -4 and 4, samples 6A and 6B

demonstrated larger differences for 1,439 genes (**Figure S12B**). These genes were enriched for

GO terms related to neuron signaling and development (based on PANTHER GO statistical

overrepresentation test with FDR 0.05). Only 40 of the 1,439 genes were annotated as immune

genes by Early *et al.* (31), and none of the 1,439 genes were among the DE genes detected as

differentially expressed using pairwise comparisons or spline fitting. Library sizes for samples

6A and 6B were similar (26,256,507 for replicate A and 22,980,006 for replicate B). We think the

difference between the replicates at 5 hr post-injection could be caused by unknown variation in

the flies' environment. It did not influence our data analysis and conclusions, but it is necessary

to be aware of when using this dataset for other analyses.

Principal component analysis

778   Principal components analysis (PCA) was performed using function *plotPCA* from the R

779   package *DESeq2* (15) after regularized-logarithm transformation of raw counts, using the design

780   ~time+time:time to create the DEseqDataSet. Genes with zero counts across all samples were

781   first removed. The default number of 500 top genes with highest row variance was used to

782   calculate the principal components.

Differential expression analysis

In order to identify genes that had differential expression over the time course, we adopted

the linear model-based methodology proposed in Law *et al.* (18) and available in the R package

limma. We first transformed the normalized RNA-seq read counts (before $\log_2$ transformation)

using the *voom* transformation, which estimates the heteroscedastic mean variance

relationships of log-counts and adds a precision weight to each observation to make them

amenable to the usual linear modeling pipelines that rely on normality. We used gene-wise

linear models to fit cubic splines (with 3 degrees of freedom) with time, TMM normalization

method (99), and standard empirical Bayes *F*-tests to select genes whose expression levels

were significantly altered across the time course in both replicates.

We also fit 3 degree polynomials across the first 48h using the R package maSigPro (19,

20). We used default parameters, including counts=F to model the data based on a normal

distribution, since we ran maSigPro on counts that were normalized using Limma-Voom. We

selected 169 genes as significantly time dependent if they had alfa (Benjamini-Hochberg

corrected) < 0.05 and a goodness of fit R^2 value of at least 0.6.

Next, we checked for differential expression of every gene between time point 0 (control)

and time point *t*, for *t* = 1, 2, …, 48 hr. For each test, a multiple testing correction at 5% False

Discovery Rate (FDR) using the Benjamini-Hochberg method (30) was adopted. Venn diagrams

to compare results were adapted from those generated using web tool Venny

(http://bioinfogp.cnb.csic.es/tools/venny/).

Functional annotation

Gene Ontology (GO) enrichment analysis was performed using PANTHER Statistical

Overrepresentation Test (http://pantherdb.org/, v14.1, released 2019/04/29) (100) using default

settings ("GO biological process complete" as annotation dataset, Fisher's Exact test, FDR <

0.05). Transcription factors were identified among differentially expressed genes based on a list

of 753 putative site-specific transcription factors available via the FlyTF database (v1) (32).

Gene set analysis was done using the R package *GSA*, which uses a Gene Set Analysis

algorithm (101) that improves the GSEA algorithm (102) by allowing testing for associations

between gene sets and time-dependent variables (101, 103). Gene set membership was

assigned from GO data downloaded from FlyBase.org in January 2019 (version FB2019_01).

Normalized counts for both replicates at each time point from 1 to 120 hr were compared

against both control replicates (0 h), using a two-class paired vector (-1, 1, -2, 2) which

corresponds to (control_replicateA, timepointX_replicateA, control_replicateB,

timepointX_replicateB). We used 100,000 permutations to estimate false discovery rates. Only

pathways with *P*-values below 0.05 and with 5 or more genes from our full dataset were kept. A

subset of most relevant pathways was compiled by selecting pathways that had at least one

gene from the subset of 551 most predominant time-dependent genes, and had a score of 2.5

or more in at least one time point from 1 to 48 hr. This gave us 41 unique pathways as shown in

Figure 9.

Cluster analyses

Hierarchical clustering of 91 core genes was performed using R package *hclust*, using

Euclidean correlation as a distance metric. Hierarchical clustering of 551 predominant time-

dependent genes was done using the default Pearson correlation with R package *pheatmap*.

Cluster membership assignment and mean patterns of expression across time for genes within

each cluster was performed as done in White *et al.* (26). Temporal clustering was performed

using the R package *TSclust* (104). Normalized counts of both replicates were clustered using

dissimilarity measures from Autocorrelation-based method (ACF), which computes the

dissimilarity between two time series as the distance between their estimated simple

autocorrelation coefficients (105). This method was used with a *P*-value cutoff of 0.05, and only

top 1% correlation edges were further explored.

Cyclic gene patterns were identified using the JTK_Cycle algorithm (43) available in R

package *JTK_Cycle*. Nine regularly distributed time points were subset from both replicates

every 6 hr (0, 6, 12, 18, 24, 30, 36, 42, 48 hr). The time point corresponding to 18 hr was

approximated by averaging normalized gene counts between time points 16 and 20 h. We

looked for rhythms between 18-30 hr (4 to 6 time points) with a cutoff of BH *Q*-value < 0.05 and

amplitude > 0.5.

Network inference

Granger causality-based methods (22) were used to construct putative interaction

networks among genes in the form of directed graphs with individual genes as nodes. A directed

edge from gene *A* to gene *B* is added if the time course of gene *A* Granger-causes the time

course of gene *B*. The notion of 'Granger causality' is popular in learning lead-lag relationships

among two or more time series. Formally, if the time series of gene *A*, given by x_t, has some

power in predicting the expression of gene *B* at time $t + 1$, called y_{t+1}, over and above y_t and

conditioned on an information set I_t, then gene *A* is said to exert a *Granger causal* effect on

gene *B*. Bivariate Granger causality uses a small information set $I_t = \{x_{1:t}, y_{1:t}\}$ and captures

Granger causal relationship from gene *A* to gene *B* by testing whether the regression coefficient

β in the following bivariate regression is different from zero:

$$y_{t+1} = \alpha\, y_t + \beta\, x_t + error_{t+1}$$

A master set of 258 genes was constructed from the 551 predominant time-dependent

genes by picking those that had available functional annotation and that had differential

expression of at least 2 $\log_2$ fold change. Using linear regression (function lm() in R), we

conducted bivariate (pairwise) Granger causality tests for every pair of genes among this set of

258 genes using data on sliding windows of $t = 6$ consecutive time points and the two replicates

(sample size = 12), and ranked them in order of increasing *P*-values (BH method used for

calculating FDR), keeping the top resulting edges (BHFDR < 5%).

A well-known critique of bivariate Granger causality is its use of a small information set

that does not contain any other factors except genes *A* and *B* (106). This failure to account for

other potential confounding variables can give rise to many spurious edges in our network

(106), where Granger causal effects from gene *A* to gene *B* is an artefact of gene *C*, which is

causal for one or both genes. To address this, we adopted multivariate (or network) Granger

causality (107), allowing us to avoid such spurious inferences through multiple linear regression.

In this framework, we start with *p* genes, and Granger causal relationship of Gene *A* on Gene *B*

is tested by regressing y_{t+1} on y_t, x_t and the time courses of the other *p* - 2 genes

$z_{1t}, z_{2t}, \ldots, z_{pt}$.

$$y_{t+1} = \alpha\, y_t + \beta\, x_t + \gamma_1 z_{1t} + \gamma_2 z_{2t} + \ldots + \gamma_{p-2}\, z_{p-2,tt} + error_{t+1}$$

For small sample size and large *p*, the above regression is not possible to run using

ordinary least squares (OLS), so we use LASSO (59) regression. To test if the regression

coefficient β in the above regression is different from zero, we used two different variants of de-

biased LASSO (60, 61), each of which corrects the bias of lasso and allows quantifying

uncertainty of regression coefficients one at a time. A non-zero coefficient in the above

multivariate regression suggests that gene *A* is Granger causal for gene *B*, even after

accounting for the effects of the other *p*-2 genes. Using this method on the master set of 258

genes, we reconstructed putative directed networks of multivariate Granger causality and

874 ranked the edges in increasing order of *P*-values, following the same parameters used in the

875 bivariate (pairwise) Granger causality method (sliding window of 6 consecutive time points in

876 both replicates, keeping the top resulting edges (BHFDR < 5%)).

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The RNA-seq data will be available at NCBI SRA, BioProject ID PRJNA641552. The pipeline scripts and intermediary data files will be made available through GitHub. A query interface will allow any user to query a gene of interest and download a table of raw counts, normalized counts, log2 fold change, and plots of gene trajectory over time.

Competing interests

The authors declare that they do not have any competing interests.

Funding

FS was supported by a Presidential Life Science Fellowship (PLSF) from Cornell University. SB, MTW, AGC and SYND were supported by NIH award (R01GM135926). In addition, SB was supported by NSF award DMS-1812128.

Authors' contributions

FS, AE, and AGC conceived the study. FS collected the samples and generated the time course data. FS, SB, MTW, and AGC conceived the computational and statistical analyses. FS, SYND, and SB performed the computational and statistical analyses. FS, SYND, SB, and AGC wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors want to thank Juan Felipe Beltrán, Amanda Manfredo, Keegan Kelsey, Yasir Ahmed-Braimah, and Elissa Cosgrove for all the help and advice provided. We also thank the three anonymous reviewers for their comments and suggestions.

SUPPORTING INFORMATION

Figure S1. Plots of normalized counts of housekeeping genes and immune response genes.

Figure S2. Venn Diagram showing overlap and differences of DE genes identified using limma-

voom spline fitting vs. maSigPro fitting of polynomials.

Figure S3. Heatmap of 214 genes

Figure S4. Temporal dynamics of gene expression of the most strongly up-regulated genes .

Figure S5. Expression profiles of DE genes encoding transcription factors.

Figure S6. Gluconeogenesis pathway

Figure S7. GC filtered network of negative edges

Figure S8. Negative GC edges

Figure S9. Pathway corresponding to 'mitotic DNA replication checkpoint'

Figure S10. GC edges of circadian rhythm genes plotted against time.

Figure S11. Positive GC edges

Figure S12. Outlier explanation

Table S1. List of differentially expressed genes identified using spline fitting and pairwise

comparisons.

Table S2. Genes that encode transcription factors and respond to commercial LPS injection.

SUPPLEMENTAL FIGURES

Figure S1. Plots of normalized counts of housekeeping genes (**A**) show little change across time as

expected under proper data normalization, while immune response genes (**B**) show up-regulation within

the first time points, as expected after a successful Imd stimulation.

958 **Figure S2.** Venn Diagram showing overlap and differences of DE genes identified using limma-voom

959 spline fitting vs. maSigPro fitting of polynomials.

960
Figure S3. Heatmap of 214 genes differentially expressed following injection with commercial LPS.

Scale is $\log_2$ fold change, with negative expression in purple and positive expression in dark orange,

compared to baseline.

Figure S4. (A) Temporal dynamics of the most strongly up-regulated genes (*DptB*, *AttC*, *Mtk*, *Dro*,

CG45045, *DptA*, *CG43920*, and *edin*) in the first 48 hr after injection. These genes are part of the 91

"core" DE genes with $\log_2$FC > 2 in at least two time intervals after injection, with an FDR < 0.01 (**Figure**

3). (**B**) Temporal dynamics of gene expression of circadian rhythm genes (*per*, *tim*, *to*, *vri*) show a classic

and well characterized 24 hr periodic expression.

Figure S5. Expression profiles of DE genes encoding transcription factors. (A) Genes involved in regulation of the circadian clock. **(B)** *p53,* which encodes a transcription factor involved in DNA repair and genotoxic stress.

Figure S6. Gluconeogenesis pathway

982 **Figure S7. GC filtered network.** The network was pruned to only the edges with at least three

983 consecutive windows of GC significance, those that have a negative GC relationship, and it excludes

984 genes identified with JTK_Cycle.

989 **Figure S8. Negative GC edges. (A) between** *Claspin* **and** *UGP.* **(B) between** *LpR2* **and** *fbp* Windows

990 in which a significant Granger Causal relationship is established are colored in blue and are contained in

991 a dashed blue box marking the plot area, non-significant windows are grey below the plot.

993   **Figure S9. Pathway corresponding to 'mitotic DNA replication checkpoint'**

Figure S10. GC edges of circadian rhythm genes plotted against time. (**A**) Positive lead-lag

correlation between *Smvt* and *cry*. (**B**) Negative lead-lag correlation between *to* and *vri*. (**C**) Negative

lead-lag correlation between *period* and *takeout*. (**D**) Positive lead-lag correlation between *Smvt* and

takeout. Significant windows are colored in blue, non-significant windows are colored in grey. Resulting

overall consecutive windows are labeled in blue dashed rectangles. Individual windows represent 6

consecutive time points. Note that time points are not regularly distributed with time, therefore windows

have different time ranges, but identical number of samples.

1005
Figure S11. Positive GC edges. Positive lead-lag correlation between (**A**) *period* and *Rh5*, (**B**) *BomS1*
and *PGRP-SD,* (**C**) *Dso1* and *PGRP-SD,* (**D**) *BomBc1* and *PGRP-SD,* (**E**) *DptB* and *PGRP-SD,* (**F**) *AttC*
and *PGRP-SB1,* and (**G**) *AGBE* and *Hao.* Significant windows are colored in blue, non-significant
windows are colored in grey. Resulting overall consecutive windows are labeled in blue dashed
rectangles. Individual windows represent 6 consecutive time points. Note that time points are not regularly
distributed with time, therefore windows have different time ranges, but identical number of samples.

1012

Figure S12. Sample 6A is an outlier for a subset of genes. (**A**) Heatmap showing row Z scores of normalized counts for all 12,657 genes in the filtered datasets. Each row represents a gene; each column represents a replicate, from pre-injection (0 hours) up to 6 hr post-injection. (**B**) Distribution of differences between normalized count Z scores for replicates A and B for each time point.

 Table S2. 20 genes that respond to commercial LPS injection encode transcription factors.

Symbol	Flybase ID	Flybase snapshot
E(spl)m3-HLH	FBgn0002609	Enhancer of split m3, helix-loop-helix (E(spl)m3-HLH) is a member of the enhancer of split gene (E(spl)) complex. E(spl)m3-HLH encodes a transcriptional repressor executing Notch-mediated cellular differentiation.
GATAe	FBgn0038391	GATAe (GATAe) encodes a endoderm-specific GATA factor. It regulates endoderm differentiation and intestinal stem cell maintenance.
CG10147	FBgn0035702	NA
exex	FBgn0041156	extra-extra (exex) encodes a homeodomain transcription factor that is expressed in and regulates the differentiation of motor neurons that project axons to ventral body wall muscles. Within the central nervous system, the product of exex negatively interacts with the products of Lim3 and eve to govern neuronal specification and differentiation. The roles of the product of exex include neuronal specification and differentiation.
CG2678	FBgn0014931	NA
vri	FBgn0016076	vrille (vri) encodes a bZIP transcription factor acting as an enhancer of dpp phenotypes both in embryo and in wing. It is involved in hair and cell growth and in tracheal development. Vri is a clock-controlled gene acting as a repressor of the products of Clk and cry.
Pdp1	FBgn0016694	PAR-domain protein 1 (Pdp1) encodes a member of the PAR domain bZip family of sequence-specific transcription factors. It regulates gene expression in muscles and in circadian clock neurons.
Clk	FBgn0023076	NA
unpg	FBgn0015561	NA
cbt	FBgn0043364	cabut (cbt) encodes a transcription factor that controls Dpp signaling and is involved in dorsal closure and wing disc morphogenesis.
sr	FBgn0003499	stripe (sr) encodes a transcription factor that induces the fate of tendon cells in the embryo as well as in the adult fly. It works upstream of tendon specific genes including Tsp, slow and Lrt.
CrebA	FBgn0004396	Cyclic-AMP response element binding protein A (CrebA) encodes a leucine-zipper transcription factor that upregulates genes encoding protein components of the canonical secretory pathway and tissue-specific genes in the salivary gland and epidermis.
Ets21C	FBgn0005660	Ets at 21C (Ets21C) encodes a stress-inducible transcription factor that binds specifically to purine-rich DNA motifs. It cooperates with transcription factors acting downstream of the MAP kinase pathways to fine tune gene expression in response to stress, such as infection or oncogene activation.
Dif	FBgn0011274	Dorsal-related immunity factor (Dif) encodes a transcription factor that contributes to zygotic function of the Toll pathway, notably the regulation of antimicrobial peptides, but it is not involved in dorsoventral patterning of the embryo.
Rel	FBgn0014018	Relish (Rel) encodes a transcription factor and the downstream component of the immune deficiency pathway, which regulates the antibacterial response and other less characterized cellular processes.
Hr38	FBgn0014859	Hormone receptor-like in 38 (Hr38) encodes a protein that can heterodimerize with the U adult cuticle and for the proper uptake and storage of glycogen in larvae.
luna	FBgn0040765	NA
p53	FBgn0039044	p53 (p53) encodes a transcriptional factor required for adaptive responses to genotoxic stress, including cell death, compensatory proliferation and DNA repair.
Hand	FBgn0032209	Hand (Hand) encodes a transcription factor that contributes to cardiogenesis, hemopoiesis, and muscle function.
Fer3	FBgn0037937	48 related 3 (Fer3) encodes a basic helix-loop-helix (bHLH) transcription factor expressed in the ventral nerve cord.

REFERENCES

1039

1.   Lazzaro BP, Galac Madeline R. Disease Pathology: Wasting Energy Fighting Infection.
Current Biology. 2006;16(22):R964-R5.
2.   Zerofsky M, Harel E, Silverman N, Tatar M. Aging of the innate immune response in
Drosophila melanogaster. Aging Cell. 2005;4(2):103-8.
3.   DiAngelo JR, Bland ML, Bambina S, Cherry S, Birnbaum MJ. The immune response
attenuates growth and nutrient storage in Drosophila by reducing insulin signaling. Proceedings
of the National Academy of Sciences of the United States of America. 2009;106(49):20853-8.
4.   Fitzpatrick M, Young SP. Metabolomics – A novel window into inflammatory disease.
Swiss medical weekly. 2013;143:w13743-w.
5.   McKean KA, Yourth CP, Lazzaro BP, Clark AG. The evolutionary costs of immunological
maintenance and deployment. BMC Evolutionary Biology. 2008;8(1):76.
6.   Howick VM, Lazzaro BP. Genotype and diet shape resistance and tolerance across
distinct phases of bacterial infection. BMC Evolutionary Biology. 2014;14(1):56.
7.   Fedorka KM, Linder JE, Winterhalter W, Promislow D. Post-mating disparity between
potential and realized immune response in Drosophila melanogaster. Proceedings of the Royal
Society B: Biological Sciences. 2007;274(1614):1211-7.
8.   Short SM, Lazzaro BP. Female and male genetic contributions to post-mating immune
defence in female Drosophila melanogaster. Proceedings of the Royal Society B: Biological
Sciences. 2010;277(1700):3649-57.
9.   Short SM, Wolfner MF, Lazzaro BP. Female *Drosophila melanogaster* suffer reduced
defense against infection due to seminal fluid components. Journal of insect physiology.
2012;58(9):1192-201.
10.   Schwenke RA, Lazzaro BP, Wolfner MF. Reproduction–Immunity Trade-Offs in Insects.
Annual review of entomology. 2016;61:239-56.
11.   Boutros M, Agaisse H, Perrimon N. Sequential Activation of Signaling Pathways during
Innate Immune Responses in Drosophila. Developmental Cell. 2002;3(5):711-22.
12.   De Gregorio E, Spellman PT, Rubin GM, Lemaitre B. Genome-wide analysis of the
Drosophila immune response by using oligonucleotide microarrays. Proceedings of the National
Academy of Sciences of the United States of America. 2001;98(22):12590-5.
13.   Sackton TB, Lazzaro BP, Clark AG. Genotype and Gene Expression Associations with
Immune Function in Drosophila. PLOS Genetics. 2010;6(1):e1000797.
14.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40.
15.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550.
16.   Bar-Joseph Z, Gitter A, Simon I. Studying and modelling dynamic biological processes
using time-series gene expression data. Nature Reviews Genetics. 2012;13:552.
17.   Spies D, Ciaudo C. Dynamics in Transcriptomics: Advancements in RNA-seq Time
Course and Downstream Analysis. Computational and Structural Biotechnology Journal.
2015;13:469-77.
18.   Law CW, Chen Y, Shi W, Smyth GKJGB. voom: precision weights unlock linear model
analysis tools for RNA-seq read counts. Genome Biology. 2014;15(2):R29.
19.   Conesa A, Nueda MJ, Ferrer A, Talón M. maSigPro: a method to identify significantly
differential expression profiles in time-course microarray experiments. Bioinformatics.
2006;22(9):1096-102.
20.   Nueda MJ, Tarazona S, Conesa A. Next maSigPro: updating maSigPro bioconductor
package for RNA-seq time series. Bioinformatics. 2014;30(18):2598-602.

21.   Kaneko T, Goldman WE, Mellroth P, Steiner H, Fukase K, Kusumoto S, et al.
Monomeric and Polymeric Gram-Negative Peptidoglycan but Not Purified LPS Stimulate the
Drosophila IMD Pathway. Immunity. 2004;20(5):637-49.
22.   Granger CWJ. Investigating causal relations by econometric models and cross-spectral
methods. Econometrica. 1969;37(3):424-38.
23.   Graham AL, Shuker DM, Pollitt LC, Auld SKJR, Wilson AJ, Little TJ. Fitness
consequences of immune responses: strengthening the empirical framework for
ecoimmunology. Functional Ecology. 2011;25(1):5-17.
24.   Deng Q, Ramsköld D, Reinius B, Sandberg R. Single-Cell RNA-Seq Reveals Dynamic,
Random Monoallelic Gene Expression in Mammalian Cells. Science. 2014;343(6167):193-6.
25.   Bendjilali N, MacLeon S, Kalra G, Willis SD, Hossian AKMN, Avery E, et al. Time-
Course Analysis of Gene Expression During the Saccharomyces cerevisiae Hypoxic Response.
G3: Genes|Genomes|Genetics. 2017;7(1):221-31.
26.   White RJ, Collins JE, Sealy IM, Wali N, Dooley CM, Digby Z, et al. A high-resolution
mRNA expression time course of embryonic development in zebrafish. eLife. 2017;6:e30860.
27.   Novembre J, Stephens M. Interpreting principal component analyses of spatial
population genetic variation. Nature genetics. 2008;40(5):646-9.
28.   Podani J, Miklós I. Resemblance Coefficients and The Horseshoe Effect in Principal
Coordinates Analysis. Ecology. 2002;83(12):3331-43.
29.   Lü J, Yang C, Zhang Y, Pan H. Selection of Reference Genes for the Normalization of
RT-qPCR Data in Gene Expression Studies in Insects: A Systematic Review. Frontiers in
Physiology. 2018;9:1560-.
30.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B
(Methodological). 1995;57(1):289-300.
31.   Early AM, Arguello JR, Cardoso-Moreira M, Gottipati S, Grenier JK, Clark AG. Survey of
Global Genetic Diversity Within the Drosophila Immune System. Genetics. 2017;205(1):353.
32.   Pfreundt U, James DP, Tweedie S, Wilson D, Teichmann SA, Adryan B. FlyTF:
improved annotation and enhanced functionality of the *Drosophila* transcription factor database.
Nucleic acids research. 2010;38(Database issue):D443-D7.
33.   Meng X, Khanuja BS, Ip YT. Toll receptor-mediated Drosophila immune response
requires Dif, an NF-κB factor. Genes & Development. 1999;13(7):792-7.
34.   Manfruelli P, Reichhart J-M, Steward R, Hoffmann JA, Lemaitre B. A mosaic analysis in
Drosophila fat body cells of the control of antimicrobial peptide genes by the Rel proteins Dorsal
and DIF. The EMBO Journal. 1999;18(12):3380-91.
35.   Myllymäki H, Valanne S, Rämet M. The *Drosophila* Imd Signaling Pathway. The Journal
of Immunology. 2014;192(8):3455.
36.   Mundorf J, Donohoe CD, McClure CD, Southall TD, Uhlirova M. Ets21c Governs Tissue
Renewal, Stress Tolerance, and Aging in the *Drosophila* Intestine. Cell Reports.
2019;27(10):3019-33.e5.
37.   Chen X, Rahman R, Guo F, Rosbash M. Genome-wide identification of neuronal activity-
regulated genes in Drosophila. eLife. 2016;5:e19942.
38.   Cyran SA, Buchsbaum AM, Reddy KL, Lin MC, Glossop NR, Hardin PE, et al. *vrille*,
Pdp1, and *dClock* form a second feedback loop in the *Drosophila* circadian clock. Cell.
2003;112(3):329-41.
39.   Collins B, Mazzoni EO, Stanewsky R, Blau J. *Drosophila* CRYPTOCHROME Is a
Circadian Transcriptional Repressor. Current Biology. 2006;16(5):441-9.
40.   Brodsky MH, Weinert BT, Tsang G, Rong YS, McGinnis NM, Golic KG, et al. *Drosophila*
melanogaster MNK/Chk2 and p53 Regulate Multiple DNA Repair and Apoptotic Pathways
following DNA Damage. Molecular and Cellular Biology. 2004;24(3):1219-31.

41.	Hoffmann JA, Reichhart J-M. Drosophila innate immunity: an evolutionary perspective.
Nature Immunology. 2002;3:121.
42.	Miyamoto T, Amrein H. Gluconeogenesis: An ancient biochemical pathway with a new
twist. Fly (Austin). 2017;11(3):218-23.
43.	Hughes ME, Hogenesch JB, Kornacker K. JTK_CYCLE: an efficient nonparametric
algorithm for detecting rhythmic components in genome-scale data sets. J Biol Rhythms.
2010;25(5):372-80.
44.	Cirelli C, LaVaute TM, Tononi G. Sleep and wakefulness modulate gene expression in
Drosophila. Journal of Neurochemistry. 2005;94(5):1411-9.
45.	Shirasu-Hiza MM, Dionne MS, Pham LN, Ayres JS, Schneider DS. Interactions between
circadian rhythm and immunity in *Drosophila melanogaster*. Current Biology. 2007;17(10):R353-
R5.
46.	Valanne S, Salminen TS, Järvelä-Stölting M, Vesala L, Rämet M. Correction: Immune-
inducible non-coding RNA molecule *lincRNA-IBIN* connects immunity and metabolism in
Drosophila melanogaster. PLOS Pathogens. 2019;15(10):e1008088.
47.	Troha K, Im JH, Revah J, Lazzaro BP, Buchon N. Comparative transcriptomics reveals
CrebA as a novel regulator of infection tolerance in D. melanogaster. PLoS Pathogens.
2018;14(2):e1006847.
48.	Clemmons AW, Lindsay SA, Wasserman SA. An Effector Peptide Family Required for
Drosophila Toll-Mediated Immunity. PLOS Pathogens. 2015;11(4):e1004876.
49.	Cohen LB, Lindsay SA, Xu Y, Lin SJH, Wasserman SA. The Daisho Peptides Mediate
Drosophila Defense Against a Subset of Filamentous Fungi. Frontiers in Immunology.
2020;11(9).
50.	Kenmoku H, Hori A, Kuraishi T, Kurata S. A novel mode of induction of the humoral
innate immune response in *Drosophila* larvae. Disease Models & Mechanisms.
2017;10(3):271.
51.	Ekengren S, Tryselius Y, Dushay MS, Liu G, Steiner H, Hultmark D. A humoral stress
response in Drosophila. Current Biology. 2001;11(9):714-8.
52.	Ao J, Ling E, Yu X-Q. Drosophila C-type lectins enhance cellular encapsulation. Mol
Immunol. 2007;44(10):2541-8.
53.	Keebaugh ES, Schlenke TA. Adaptive evolution of a novel *Drosophila* lectin induced by
parasitic wasp attack. Mol Biol Evol. 2012;29(2):565-77.
54.	Zsámboki J, Csordás G, Honti V, Pintér L, Bajusz I, Galgóczy L, et al. *Drosophila* Nimrod
proteins bind bacteria. Central European Journal of Biology. 2013;8(7):633-45.
55.	Katzenberger RJ, Ganetzky B, Wassarman DA. Age and Diet Affect Genetically
Separable Secondary Injuries that Cause Acute Mortality Following Traumatic Brain Injury in
Drosophila. G3: Genes|Genomes|Genetics. 2016;6(12):4151-66.
56.	Buchon N, Poidevin M, Kwon HM, Guillou A, Sottas V, Lee BL, et al. A single modular
serine protease integrates signals from pattern-recognition receptors upstream of the
Drosophila Toll pathway. Proc Natl Acad Sci U S A. 2009;106(30):12442-7.
57.	Fujita A, Severino P, Kojima K, Sato JR, Patriota AG, Miyano S. Functional clustering of
time series gene expression data by Granger causality. BMC Systems Biology. 2012;6(1):137.
58.	Finkle JD, Wu JJ, Bagheri N. Windowed Granger causal inference strategy improves
discovery of gene regulatory networks. Proceedings of the National Academy of Sciences.
2018;115(9):2252.
59.	Tibshirani R. Regression Shrinkage and Selection via the Lasso. Journal of the Royal
Statistical Society Series B (Methodological). 1996;58(1):267-88.
60.	Javanmard A, Montanari A. Confidence intervals and hypothesis testing for high-
dimensional regression. J Mach Learn Res. 2014;15(1):2869-909.
61.	Dezeure R, Buhlmann P, Meier L, Meinshausen N. High-Dimensional Inference:
Confidence Intervals, p-Values and R-Software hdi. Statist Sci. 2015;30(4):533-58.

62.   Lee E-M, Trinh TTB, Shim HJ, Park S-Y, Nguyen TTT, Kim M-J, et al. Drosophila
Claspin is required for the G2 arrest that is induced by DNA replication stress but not by DNA
double-strand breaks. DNA Repair. 2012;11(9):741-52.
63.   Ubhi T, Brown GW. Exploiting DNA Replication Stress for Cancer Treatment. Cancer
Research. 2019.
64.   Liu D, Shaukat Z, Saint RB, Gregory SL. Chromosomal instability triggers cell death via
local signalling through the innate immune receptor Toll. Oncotarget. 2015;6(36):38552-65.
65.   Nakad R, Schumacher B. DNA Damage Response and Immune Defense: Links and
Mechanisms. Front Genet. 2016;7:147-.
66.   Soukup SF, Culi J, Gubb D. Uptake of the necrotic serpin in Drosophila melanogaster
via the lipophorin receptor-1. PLoS genetics. 2009;5(6):e1000532-e.
67.   Karlsson C, Korayem AM, Scherfer C, Loseva O, Dushay MS, Theopold U. Proteomic
Analysis of the Drosophila Larval Hemolymph Clot. Journal of Biological Chemistry.
2004;279(50):52033-41.
68.   Krautz R, Arefin B, Theopold U. Damage signals in the insect immune response. Front
Plant Sci. 2014;5:342-.
69.   Shinoda T, Itoyama K. Juvenile hormone acid methyltransferase: A key regulatory
enzyme for insect metamorphosis. Proceedings of the National Academy of Sciences of the
United States of America. 2003;100(21):11986-91.
70.   Rolff J, Siva-Jothy MT. Copulation corrupts immunity: A mechanism for a cost of mating
in insects. Proceedings of the National Academy of Sciences of the United States of America.
2002;99(15):9916-8.
71.   Flatt T, Heyland A, Rus F, Porpiglia E, Sherlock C, Yamamoto R, et al. Hormonal
Regulation of the Humoral Innate Immune Response in Drosophila melanogaster. The Journal
of experimental biology. 2008;211(Pt 16):2712-24.
72.   Schwenke RA, Lazzaro BP. Juvenile Hormone Suppresses Resistance to Infection in
Mated Female Drosophila melanogaster. Current Biology. 2017;27(4):596-601.
73.   Sarov-Blat L, So WV, Liu L, Rosbash M. The *Drosophila takeout* Gene Is a Novel
Molecular Link between Circadian Rhythms and Feeding Behavior. Cell. 2000;101(6):647-56.
74.   Wong R, Piper MDW, Wertheim B, Partridge L. Quantification of Food Intake in
Drosophila. PLOS ONE. 2009;4(6):e6063.
75.   Giebultowicz JM. Circadian regulation of metabolism and healthspan in *Drosophila*. Free
Radical Biology and Medicine. 2018;119:62-8.
76.   So WV, Sarov-Blat L, Kotarski CK, McDonald MJ, Allada R, Rosbash M. *takeout*, a
Novel *Drosophila* Gene under Circadian Clock Transcriptional Regulation. Molecular and
Cellular Biology. 2000;20(18):6935.
77.   Smith RF, Konopka RJ. Effects of dosage alterations at the *per* locus on the period of
the circadian clock of *Drosophila*. Molecular and General Genetics MGG. 1982;185(1):30-6.
78.   Reddy P, Zehring WA, Wheeler DA, Pirrotta V, Hadfield C, Hall JC, et al. Molecular
analysis of the *period* locus in *Drosophila melanogaster* and identification of a transcript
involved in biological rhythms. Cell. 1984;38(3):701-10.
79.   Claridge-Chang A, Wijnen H, Naef F, Boothroyd C, Rajewsky N, Young MW. Circadian
Regulation of Gene Expression Systems in the *Drosophila* Head. Neuron. 2001;32(4):657-71.
80.   Bahrami S, Drabløs F. Gene regulation in the immediate-early response process.
Advances in Biological Regulation. 2016;62:37-49.
81.   Hanson MA, Lemaitre B. New insights on Drosophila antimicrobial peptide function in
host defense and beyond. Current Opinion in Immunology. 2020;62:22-30.
82.   Tanji T, Hu X, Weber ANR, Ip YT. Toll and IMD Pathways Synergistically Activate an
Innate Immune Response in Drosophila melanogaster. Molecular and
Cellular Biology. 2007;27(12):4578.

83.   Irving P, Troxler L, Heuer TS, Belvin M, Kopczynski C, Reichhart J-M, et al. A genome-
wide analysis of immune responses in Drosophila. Proceedings of the
National Academy of Sciences. 2001;98(26):15119.
84.   Duneau D, Ferdy J-B, Revah J, Kondolf H, Ortiz GA, Lazzaro BP, et al. Stochastic
variation in the initial phase of bacterial infection predicts the probability of survival in D.
melanogaster. eLife. 2017;6:e28298.
85.   Wolowczuk I, Verwaerde C, Viltart O, Delanoye A, Delacre M, Pot B, et al. Feeding our
immune system: impact on metabolism. Clin Dev Immunol. 2008;2008:639803.
86.   Chi W, Dao D, Lau TC, Henriksbo BD, Cavallari JF, Foley KP, et al. Bacterial
Peptidoglycan Stimulates Adipocyte Lipolysis via NOD1. PLOS ONE. 2014;9(5):e97675.
87.   Chambers MC, Song KH, Schneider DS. Listeria monocytogenes Infection Causes
Metabolic Shifts in Drosophila melanogaster. PLOS ONE. 2012;7(12):e50679.
88.   Krejčová G, Danielová A, Nedbalová P, Kazek M, Strych L, Chawla G, et al. Drosophila
macrophages switch to aerobic glycolysis to mount effective antibacterial defense. eLife.
2019;8:e50414.
89.   Clark Rebecca I, Tan Sharon WS, Péan Claire B, Roostalu U, Vivancos V, Bronda K, et
al. MEF2 Is an In Vivo Immune-Metabolic Switch. Cell. 2013;155(2):435-47.
90.   Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-
wide expression patterns. Proceedings of the National Academy of Sciences of the United
States of America. 1998;95(25):14863-8.
91.   Im JH. Functional and Population Genetics of *Drosophila* Innate Immunity: Cornell
University; 2018.
92.   Lemaitre B, Hoffmann J. The Host Defense of Drosophila melanogaster. Annu Rev
Immunol. 2007;25(1):697-743.
93.   Konopka RJ, Benzer S. Clock Mutants of *Drosophila melanogaster*. Proceedings of the
National Academy of Sciences. 1971;68(9):2112.
94.   Myers MP, Wager-Smith K, Rothenfluh-Hilfiker A, Young MW. Light-induced degradation
of TIMELESS and entrainment of the *Drosophila* circadian clock. Science.
1996;271(5256):1736-40.
95.   McKean KA, Nunney L. Increased sexual activity reduces male immune function in
Drosophila melanogaster. Proceedings of the National Academy of Sciences of the United
States of America. 2001;98(14):7904-9.
96.   Early AM, Shanmugarajah N, Buchon N, Clark AG. Drosophila Genotype Influences
Commensal Bacterial Levels. PLOS ONE. 2017;12(1):e0170332.
97.   Chambers MC, Jacobson E, Khalil S, Lazzaro BP. Thorax Injury Lowers Resistance to
Infection in Drosophila melanogaster. Infection and Immunity. 2014;82(10):4380-9.
98.   Sefer E, Kleyman M, Joseph Z-B. Tradeoffs between dense and replicate sampling
strategies for high throughput time series experiments. Cell systems. 2016;3(1):35-42.
99.   Robinson MD, Oshlack A. A scaling normalization method for differential expression
analysis of RNA-seq data. Genome Biology. 2010;11(3):R25.
100.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic
Acids Research. 2018;47(D1):D419-D26.
101.   Efron B, Tibshirani R. On testing the significance of sets of genes. Ann Appl Stat.
2007;1(1):107-29.
102.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene
set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression
profiles. Proceedings of the National Academy of Sciences. 2005;102(43):15545.
103.   Mullighan CG, Su X, Zhang J, Radtke I, Phillips LAA, Miller CB, et al. Deletion of IKZF1
and prognosis in acute lymphoblastic leukemia. The New England journal of medicine.
2009;360(5):470-80.

104.   Montero P, Vilar JA. TSclust: An R Package for Time Series Clustering. Journal of
Statistical Software. 2014;62(1):1-43.
105.   Galeano P, Peña D. Multivariate analysis in vector time series. Resenhas Do Instituto
De Matemática E Estatística Da Universidade De São Paulo. 2000;4(4):383-404.
106.   Mukhopadhyay ND, Chatterjee S. Causality and pathway search in microarray time
series experiment. Bioinformatics. 2007;23(4):442-9.
107.   Basu S, Shojaie A, Michailidis G. Network Granger Causality with Inherent Grouping
Structure. Journal of Machine Learning Research. 2015;16:417-53.
1297