**Title**: Megakaryopoiesis in Dengue virus infected K562 cell promotes viral replication which inhibits endomitosis and accumulation of ROS associated with differentiation

Jaskaran Kaur *[1], Yogita Rawat *[1], Vikas Sood [2], Deepak Rathore[1], Shrikant K. Kumar[1], Niraj K. Kumar[1] and Sankar Bhattacharyya[1]

[1] *Translational Health Science and Technology Institute, NCR Biotech Science Cluster, PO Box# 4, Faridabad-Gurgaon expressway, Faridabad, Haryana-121001, India*

[2] *Department of Biochemistry, School of Chemical and Life Sciences, Jamia Hamdard (Hamdard University) Hamdard Nagar, New Delhi - 110062, India*

*Equal contribution

*Email for correspondence*: sankar@thsti.res.in

**Keywords:** Dengue virus replication, Megakaryopoiesis, Reactive oxygen species, Endomitosis

**Abstract:** In the human host blood Monocytes and bone marrow Megakaryocytes are implicated as major sites supporting high replication. The human K562 cell line supports DENV replication and represent Megakaryocyte-Erythrocyte progenitors (MEP), replicating features of *in vivo* Megakaryopoiesis upon stimulation with Phorbol esters. In this article, we report results that indicate the mutual influence of Megakaryopoiesis and DENV replication on each other, through comparison of PMA-induced differentiation of either mock-infected or DENV-infected K562 cells. We present data showing PMA-induced differentiation to drastically increase DENV replication and a concomitant augmented secretion of infectious virus. Although the mechanism is not clear yet, we show that it is not through an increased uptake of virus by differentiated cells. On the other hand, DENV replication in cells undergoing PMA-induced differentiation, interferes with major differentiation markers of Megakaryopoiesis including activation of ERK1/2 MAP Kinase, endomitosis and surface expression of platelet-specific proteins without any drastic effect on cell death. Among signaling intermediaries of the JAK-STAT pathway, we observed infection associated degradation of SOC3 protein similar to earlier observations with STAT2. DENV infection leads to accumulation of Reactive-oxygen species (ROS) in different cells including K562. PMA-induced differentiation of uninfected K562 cells also leads to intracellular ROS accumulation. Interestingly, we observed ROS accumulation to be suppressed by concomitant DENV replication in K562 cells undergoing PMA-induced differentiation. This is the first report of a model system where DENV replication suppresses intracellular ROS accumulation. The implications of these results for Megakaryopoiesis and viral replication would be discussed.

## Introduction

Platelets play a unique role in tissue homeostasis and regulation of inflammation. The abundance of these small, anuclear cells produced from specialized cells called Megakaryocytes (MKs) in the bone marrow (BM) is controlled by a steady rate of biogenesis (about $1 \times 10^9$/day) and decay, with an average life of 7 days in humans (1). A perturbation of platelet homeostasis is associated with infection by multiple human pathogenic viruses (2). Infection by Dengue virus (DENV) a member of the Flaviviridae, can cause an acute febrile illness with the potential to turn fatal. Beside humans, DENV can infect mosquitoes and these transmit infectious virus along with salivary fluid during a blood meal. Only about a quarter of infected humans exhibit clinical symptoms that include high fever, retro-orbital pain, muscle pain, thrombocytopenia or drop in platelet level etc. that subside after 4-7 days. Subsequently a proportion of symptomatic individuals develop symptoms of severe Dengue characterized by leakage of fluid from the blood vessels, leading to reduction in blood volume in addition to acute thrombocytopenia (3). Multiple mechanisms working in parallel have been suggested to contribute to the thrombocytopenia associated with Dengue infection which include inhibition of platelet biogenesis by infection of Megakaryocytes, augmented platelet decay through either a direct interaction with the virus or upon binding of anti-platelet antibodies (4).

Depending on the cell type the entry of DENV can be mediated by different surface receptors and the complex is endocytosed into endosome following virus binding (5) Acidification of these endosomes by their membrane resident proton pumps induce a fusion between the viral and endosomal membranes leading to release of the viral genome into host cytosol. The genome of DENV is a ~11 kb long single-stranded plus-sense RNA coding for one large polyprotein which is then cleaved to generate the structural (Core or C, Envelop or E and precursor-Membrane or prM) and non-structural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5) of the virus. After few rounds of translation the viral genomic RNA undergoes replication in **r**eplication **c**omplexes (RC) that are associated with the host cell **E**ndoplasmic **r**eticulum (ER). The NS5 protein performs crucial function in replication of the genome by polymerizing both

negative and positive strand genomic RNA through asymmetric replication in favor of the later. Among

these only the plus-sense genomic RNA is capped by a Methyl-transferase (MTase) domain also harbored

in the NS5, followed by encapsidation of the genomic RNA by the Core or C protein. The newly assembled

virus particles contain a RNA-protein complex consisting of a single copy of the genomic RNA with

multiple copies of the C protein, encased in a membrane studded with E and prM protein. These transit

through the protein secretory pathway of the host cell where the prM protein is cleaved by a host protease

called Furin to produce M protein. The virion particles containing M protein along with E in the envelope

are released from the host cell without lysis (6).

DENV can infect multiple cell types in the human body including monocytes in the blood,

megakaryocytes in the BM and hepatocytes in the liver (5, 7). In addition to these DENV can activate

platelets, through a direct interaction with viral receptors on the cell surface (5, 8, 9). Platelets activated by

DENV secrete a number of cytokines and chemokines among which CXCL4 promotes viral replication in

infected Monocytes, upon binding to CXCR3 receptors on these cells (10). Myelosuppression or a reduction

in the mass of BM cells is one of the hallmarks of DENV infection and cells of Megakaryocytic cells have

been shown to be particularly permissive to viral replication (11-14). This suggested MK cells to

significantly contribute to viremia beside monocytes. In addition the survival of megakaryocytes have been

suggested to be negatively affected by infection, thereby leading to reduced platelet biogenesis which

contributes to DENV associated thrombocytopenia (15).

Hematopoietic stem cells (HSC) in the BM differentiate into multiple lineages producing all cell

types in the blood. One of such lineages are composed of bi-potential Megakaryocyte-Erythrocyte

progenitor (MEP) cells that can differentiate into either cell type (16, 17). Differentiation of MEP into a

MK involves profound cellular and molecular changes including cell expansion, endomitosis which

generates a multi-lobed and polyploid nucleus and expression of platelet specific surface markers on the

plasma membrane which will become part of the platelets membrane (18, 19). Platelets bud off from the

MK along with a bit of its cytoplasm, containing granules of different types and a host of specific mRNAs

(20). Differentiation of MEPs into MKs is promoted by a number of cytokines, the most characterized

among which is Thrombopoietin or TPO that binds to the TPO-receptor c-Mpl on the cell surface (21).

TPO binding to c-Mpl activates Janus Kinase 2 (JAK2) receptor tyrosine kinase which in turn transduces

the signal through specific **S**ignal **t**ransducer and **a**ctivator of **t**ranscription or STATs (STA3 and STAT5),

**Mi**togen-**a**ctivated **p**rotein **k**inases or MAPK (ERK1/2) and the Phosphatidyl Inosine-3 or PI-3 Kinase (22).

STAT3 and STAT5 have been shown to have opposing function in Megakaryopoiesis with the former

promoting it (23, 24). Activation of JAK2 pathway increases expression of pro-inflammatory genes and in

fact activating these genes through exogenous activation of Toll-like receptor 2 or TLR2 has been shown

to promote megakaryocyte development (25). In a similar manner, the activation of ERK1/2 MAPK and

inhibition of p38-MAPK promotes Megakaryopoiesis by playing a crucial role in endomitosis (26).

Transduction of the signal from JAK2 leads to activated transcription by a number of transcription factors

(TFs) increasing the expression of specific genes (27, 28). TFs like GATA-1 and RUNX1 play crucial role

in endomitosis through upregulation of CyclinD1 (29, 30). In addition to these different specific pathways

like ER-stress and Autophagy have been shown to play role in Megakaryopoiesis (31). A controlled form

of apoptosis is known to be crucial for production of platelets from megakaryocytes and ER-stress has been

suggested to play a role in this apoptosis (31-33). In the BM, HSCs occupy a relatively hypoxic niche

compared to more differentiated lineages, particularly the MKs which shed mature platelets into the blood

vasculature. This migration from hypoxic to normoxic environment increases the level of reactive oxygen

species (ROS) in these cells, which has been indicated as a crucial signaling molecules (34). In fact

increased oxygen level and augmented expression of ROS generating enzymes show positive correlation

with megakaryocyte development, corroborating ROS to serve a critical promoter of megakaryopoiesis (35-

37). ROS has been suggested to function at multiple levels which include increasing the level of tyrosine-

phosphorylation on some proteins or by inhibition certain tyrosine phosphatases or by maintenance of ERK

activation or modulation of cyclin levels or increasing expression of crucial TFs or promoting

differentiation associated apoptosis (35, 38-40).

K562 cells represent a bi-potential cell line which can be differentiated to either MK-like cells or

Erythroblast-like cells depending on differential pharmacological stimulation (41). Supplementation of

Phorbol esters like Phorbol-12 myristate-13 (PMA) in growth media of K562 cells drives differentiation

towards megakaryocytes, inducing cellular and molecular changes akin to MKs *in vivo,* through activation

of identical signaling axes (42-50). In addition to this K562 is permissible to DENV replication and is

therefore a good model system to study the effect of DENV infection on Megakaryopoiesis (51).   In this

report we show results suggesting that when DENV infected MEPs differentiate into megakaryocytes the

intracellular viral replication is promoted, by a mechanism that is not clear yet. This enhancement of viral

replication affects key signaling pathways that are known to be crucial for maturation of megakaryocytes,

without any significant effect on their survival. DENV infection is known to induce an accumulation of

ROS in the host cells. K562 cells induced to differentiate into megakaryocytes by treatment with PMA also

accumulate ROS. However surprisingly, infected cells undergoing differentiation accumulate significantly

less ROS when compared to uninfected controls, suggesting an active suppression of ROS by viral

replication. The implications of these observations with respect to Megakaryopoiesis and DENV replication

have been discussed.

**Results:**

**K562 cells recapitulate major events in Megakaryopoiesis upon PMA-treatment**

Differentiation of MK from an MEP follows elaborate cellular and molecular changes in distinct steps, a

few of which  been depicted in the schematic figure and involves dramatic increase in cell size and

endomitosis with consequential formation of a multi-lobed, polyploid nucleus (Fig. 1A.). K562 cells mimic

a bi-potential Megakaryocyte-Erythrocyte progenitor (MEP) and depending upon pharmacological

treatment can be differentiated into cells of either an Mk-lineage (Phorbol esters) or Erythroblast-lineage

(Sodium Butyrate or NaB) (41). Upon supplementation of Phorbol-12 Myristate-13 acetate (PMA) in

culture media, K562 cells stopped proliferation and enlarged in size (data not shown). Further, the cells

undergo endomitosis leading to generation of polyploid cells harboring nuclei that are multilobed (Fig.

1B.). When analyzed by flow cytometry, PMA-treated K562 cells also show a dramatic increase of

cytoplasmic granule content too as evidenced by an increase in side-scatter (data not shown). The platelet

plasma membrane is derived from that of the Mk mother cell and during differentiation platelet-specific

surface markers are expressed on the Mk surface. As a corroboration of Megakaryopoiesis induced by

PMA-treatment, the surface expression of three platelet specific markers were quantified in K562 cells and

the result showed a dramatic increase in the level of CD41/61, CD61 alone and a significant increase in

number of cells positive for CD42b expression (Fig. 1C.). As evidence of polyploidy following PMA-

treatment, we observed a gradual emergence of polyploid cells within 3 days, which did not show any

significant increase in the next 3 days (Fig. 1D.). As a control for these changes being specific to K562,

PMA-supplementation did not stall the growth or induce expression of platelet-specific surface markers in

cells of the hepatoma cell line Huh7 (data not shown). Since treatment with NaB induces differentiation

towards an Erythroblast-lineage, the differential expression of specific genes were compared between K562

cells treated with either PMA or NaB. The result showed a distinct pattern differentiating the two

differentiation lineages, especially with respect to transcripts of the gene **GATA2**, **EBP42**, **GYPA** and

**CD61** (Fig. 1E.).

**Differentiation of K562-MKs promote replication of Dengue virus**

Differentiation into megakaryocytes is associated with arrest in cell cycle and other molecular changes. In order to analyze how these changes might affect the ongoing intracellular replication process, DENV replication in infected K562 cells that have been treated with PMA soon after infection, was compared to simultaneously infected cells treated with vehicle (DMSO). For this purpose, the level of DENV envelope protein accumulated was compared between these conditions at 3 or 6 days post-infection by immunofluorescence followed by analysis in a flow cytometer. The result showed that although the protein accumulated at 3 days p.i was comparable between PMA-treated and DMSO-treated cells, a dramatic difference between them was observed at day 6 (Fig. 2A.). This difference was observed using different MOI of infection, although due to saturation of immunostaining the difference was drastic at higher MOI (data not shown). In order to check if higher accumulation of viral proteins correlate with viral replication, the level of intracellular viral RNA and infectious virus in culture supernatant was compared between infected K562 cells treated with either PMA or DMSO for either 3 or 6 days. The results showed a similar higher accumulation of intracellular viral genomic RNA and extracellular infectious virus, between these two conditions at 6 days post-infection (Fig. 2B and 2C). During virus maturation the prM protein on assembled virion particles undergoes a cleavage by the host protease Furin, a step necessary for conferring infectivity to the secreted virus (52). Therefore the comparatively higher infectious titer in culture supernatant of PMA-treated cells can result from either higher secretion of virus (more virion particles without any change in their infectivity) or more efficient Furin cleavage brought about by the cellular and biochemical changes accompanying differentiation (same number of virion particles as secreted by DMSO-treated cells but with higher infectivity). To dissect these possibilities, the viral genomic RNA from culture supernatant of cells kept with either PMA or DMSO was extracted and purified and their relative level compared by real-time PCR using DENV specific oligonucleotide primers. The comparison showed comparatively more viral genomic RNA in the supernatant of PMA-treated cells, suggesting that the differentiating cells secrete more virus that are produced by higher replication efficiency in these cells (data

not shown). However this indicated towards possibilities wherein the entry within differentiating cells

might be facilitated through overexpression of viral receptor. In order to address this, uninfected K562 cells

treated with either DMSO or PMA for 3 days were incubated with DENV inoculum and the viral entry

compared through comparison of the intracellular genomic RNA normalized to a host mRNA transcript.

The result showed a comparable entry into cells which are undifferentiated or differentiated for 3 days,

implying that the promotion of replication occurs post-entry (Fig. 2D).  As a negative control DENV-

infected Huh7 cells did not exhibit any differential replication depending on treatment with PMA and

DMSO (data not shown). Also, treatment of infected K562 cells with Na-butyrate which would initiate

erythropoiesis, did not exhibit a similar promotion of the viral replication (data not shown).

**Cellular and biochemical changes imposed by DENV infection in differentiating K562 cells**

Differentiation of MKs from HSCs is influenced by multiple cytokines the most characterized among which

is Thrombopoietin (TPO) that is secreted from liver cells and binds to the receptor c-Mpl (Ref). Interaction

of TPO with c-MPL activates Janus kinase 2 (JAK2) tyrosine kinase which initiates a cascade of

phosphorylation events culminating in modulating through multiple (Figure 3, panel A). Although an

important cytokine, TPO is not indispensable for Megakaryopoiesis since a knock-out of the gene (TPO$^{-/-}$)

is not lethal in mice albeit with the platelet level being at 10% of that in wild-type animals (Ref). Phorbol

esters like PMA are known to activate kinases like Protein kinase-C (PKC) that in its turn transduces the

signal through nodes many of which are shared with TPO signaling (Ref). In response to viral infection

mammalian cells secrete cytokines like Interferons that bind to cell surface receptors either in an autocrine

or paracrine manner and induce an 'anti-viral' state within the cell. JAK2 is also central to signaling via the

interferon receptor consequent to binding of type-I IFNs to the receptor (IFNR) (Ref). This implied that

Megakaryopoiesis even when it is PMA-induced would potentially create an antiviral state within the cell

and in that context it would be interesting to observe how DENV is still able to replicate to higher extent

compared to undifferentiated cells.  Since both of these pathways would also be potentially perturbed by a

virus infection, we investigated the status of different members by immunoblotting, in cells which were

stimulated to undergo PMA-dependent differentiation immediately after infection.

Phosphorylation of the MAPK ERK1/2 is known to play a crucial role in MK development and we observed

an increase in the ratio of p-ERK/ERK both subsequent time points when compared to day 0 (Fig. 3B,

compare lanes 2 and 5 with lane 1**)**. Interestingly, we observed that DENV infection to reverse the PMA-

induced ERK1/2 phosphorylation at 3 days post-infection (Fig. 3B, compare lanes 3 and 4 with lane 2).

However, when compared to day 3 the phosphorylation of ERK1/2 showed an increase in infected cells by

day 6 (Fig. 3B, compare lanes 6, 7 with lanes 3, 4 ). Nonetheless, the ratio of p-ERK/ERK at 6 days post-

infection was still lower when compared to uninfected cells treated with PMA for a similar duration (Fig.

3B, compare lanes 6 and 7 with lane 5**).** In a manner opposite to that of ERK1/2, a dephosphorylation of

p38 MAPK has been similarly reported to be crucial for MK development**.** We did not observe any

modulation in the phosphorylation status of p38 in DENV-infected cells treated with PMA for 3 or 6 days

(data not shown).

In addition to MAPKs we probed the status of different signaling protein in the JAK-STAT pathway in

uninfected or DENV-infected K562 cells treated with PMA. While some effect of virus infection was

observed with respect to a few of these signaling mediators, most of them were unaffected. We were

however unable to detect a few of these signaling proteins in our experiments (SOCS1, PIAS1, PIAS3 and

PIAS4). When compared to 0 hour the ratio of p-STAT1/STAT1 showed a significant decrease upon PMA

treatment at both time points (Fig. 3B, compare lanes 2 and 5 with lane 1). However, although DENV

replication in PMA-treated cells did not alter this ratio at 3 days p.i., by 6 days p.i. the ratio showed a

dramatic increase in infected cells when compared to uninfected controls (Fig. 3B, compare lanes 3, 4 with

2 and lanes 6, 7 with lane 5). Further, at 6 days p.i. the total STAT1 protein level was also reduced to

undetectable level in infected cells (Fig. 3B, compare lanes 6 and 7 with lane 5).

STAT3 has been suggested to promote Megakaryopoiesis and in support of this we observed an increase in

ratio of p-STAT3/STAT3 at 3 days after PMA treatment, which was maintained till 6 days (Fig. 3B,

compare lanes 2 and 5 with lane 1). However in DENV infected cells treated with PMA, the ratio of p-

STAT3/STAT3 at both 3 and 6 days post-treatment was significantly lower when compared to uninfected

controls (Fig. 3B, compare lane 3 with lane 2 and lanes 6,7 with lane 5). The lane 4 in this immunoblot is

missing due to the technical reason of sample insufficiency.

Although STAT2 protein level is not known to be modulated by PMA-treatment, it is known to be degraded

in host cells following DENV infection, in a NS5 dependent manner (Ref). We did not observe any

alteration in STAT2 levels with PMA-treatment at either time points (Fig. 3B, compare lanes 2 and 5 with

lane 1). As expected however, the protein was reduced to undetectable level in DENV infected cells. (Fig.

3B, compare lanes 3,4 with lane 2 and 6,7 with lane 5).

A downregulation of STAT5 protein has been shown to be essential for promoting development of MKs.

In concurrence with that, we observed a reduction in the level of this protein upon PMA-treatment starting

from day 0 to almost undetectable level by day 6 (Fig. 3B, compare lanes 2 and 5 with lane 1). No additional

modulatory effect was however observed by DENV infection of PMA-treated cells (Fig. 3B, compare lanes

3,4 with lane 2 and 6,7 with lane 5).

An activation of the JAK-STAT signaling upregulates inflammation, which eventually leads to expression

of a family of genes that suppresses this pathways and are called as Suppressor of cytokine signaling or

SOCS1, SOCS2 and SOCS3. In accordance with a crucial role for activated JAK-STAT pathway in MK

development, expression of the SOCS3 protein has been shown to have a negative effect on

Megakaryopoiesis. Upon investigation the SOCS3 level following PMA-treatment was observed to be

unchanged at either time points (Fig. 3B, compare lanes 2 and 5 with lane 1). Surprisingly, however SOCSs

was observed to be reduced to undetectable level at day 6 pi (Fig. 3B, compare lanes 6 and 7 with lane 5).

These results implied that DENV infection might be selectively affecting the innate antiviral and

inflammatory pathways in differentiating K562 cell through degradation of  STAT2 and SOCS3 level and

interference with STAT3 phosphorylation.

**DENV infection inhibits polyploidy and accumulation of ROS in differentiating cells without**

**affecting apoptosis**

The biochemical changes associated with Megakaryopoiesis affect the cellular modifications. Therefore we

checked PMA-induced surface marker expression and polyploidy in cells infected with DENV and

compared that with PMA-treated but mock infected cells. As above, cells that were either mock-infected or

infected with DENV were treated with PMA-supplemented growth media soon after infection and analyzed

at 3 or 6 days post-treatment. The results showed a significant reduction in the surface expression of

CD41/61, CD42b and CD61 in DENV infected cells (Fig. 4A). Phosphorylation of the ERK1/2 MAPK has

been shown to be an critical modulator of polyploidy during Megakaryopoiesis. Since we observed a

suppression of ERK1/2 phosphorylation at the early time point post-infection, a comparison of the

percentage of polyploid cells at both time points was quantified by PI staining. As expected the result

showed a significant reduction in the percentage of polyploid cells generated following DENV infection

(Fig. 4B). Generation of platelet from megakaryocytes has been demonstrated to be dependent on induction

of apoptosis in the mother cell and proceeds via a unique method of intrinsic apoptosis. In accordance with

PMA-induced Megakaryopoiesis in K562 cells we observed a significant increase in Annexin-V positive

cells upon PMA-treatment (data not shown). However, a comparison of mock-infected and DENV-infected

cells treated with PMA for either 3 or 6 days did not show any difference in the proportion of Annexin-V

positive cells (data not shown). However surprisingly, a comparison of Caspase3/7 activation showed

significant increase in cleavage activity at 6 days (Fig. 4C).

ROS is generated during Megakaryopoiesis and an accumulation is essential for differentiation (Ref).

DENV replication has been shown to induce generation of reactive oxygen species in infected cell (Ref).

An estimation of the ROS accumulated in K562 cells as a result of DENV infection showed slight

accumulation at 6 days post-infection (Fig. 5A, compare columns 2 to 1 and 6 to 5). In a similar manner,

when compared to levels at day 0 uninfected cells treated with PMA showed a significant accumulation of

ROS at both 3 and 6 days post-treatment (Fig. 5A, compare columns 3 to 1 and 7 to 5). We expected the

ROS level in DENV infected cells that have been treated with PMA to be higher than the uninfected PMA-

treated controls. However, surprisingly DENV infected cells treated with PMA did not exhibit additional

ROS accumulation at day 3 post-treatment (Fig. 5A, compare columns 3 to 4). Further, surprisingly at day

6 after PMA-treatment the ROS accumulation in DENV infected cells were remarkably lower compared to

uninfected PMA-treated cells (Fig. 5A, compare columns 7 to 8). This suggested DENV replication in

differentiating cells to impede accumulation of ROS. In corroboration we observed DENV infection to have

an effect similar to that of N-acetyl cysteine a biochemical known to quench ROS levels (Fig. 5B**).** In

conclusion, DENV infection do not seem to interfere with surface marker expression but significantly affect

accumulation of ROS and endomitosis which are associated with differentiation into megakaryocytes.

**PMA-induced transcriptome changes are reversed by DENV infection in differentiating cells**

Megakaryopoiesis is associated with profound alterations in the transcriptome of differentiating cells.

Similarly, innate antiviral response launched upon detection of a viral infection is also associated with

changes in the pattern of expression of many genes. Therefore, in order to study the interaction between

these two stimuli at the transcriptional level the transcriptome of mock-infected and DENV-infected cells

was compared after 6 days of PMA-induced differentiation. For this purpose, K562 cells either mock-

infected or DENV-infected were induced to differentiate using 100 nM PMA and the total RNA isolated

for analysis of transcriptome by next-generation sequencing. The transcriptome of both uninfected and

DENV-infected cells was compared to uninfected cells at 0 hour with respect to PMA-treatment. As shown

in Fig. 6A, the transcriptome in uninfected cells underwent a profound change in more than 5000 genes

showing different degree of deregulation. Interestingly, the comparison between uninfected and DENV-

infected cells after 6 days of differentiation showed reversal in the direction of PMA-induced deregulation

for a number of genes (Fig, 6B.). Since our earlier result showed that DENV infection suppressed ROS

accumulation in differentiating cells, ROS-associated genes the PMA-induced deregulation trend of which

was reversed by infection were analyzed further. A heatmap of these genes showed that although alteration

in transcript level was moderate for most of them, a few genes were dramatically affected (Fig. 6C). This

included GCH1, FOXO3, GLRX5, ETHE1, CTNS, EPAS1, GFOD1 and SNCA (Fig. 6C). Among these

transcripts corresponding to FOXO3, CTNS and EPAS1 were higher in infected cells compared to

uninfected ones, whereas the others showed lower levels in infected cells. FOXO3 and EPAS1 are

transcription factors among which the former is known to be pro-apoptotic (53). Among the downregulated

genes the protein corresponding to GCH1, ETHE1 and SNCA are enriched platelets (54-56). SNCA is also

known to function in an anti-apoptotic manner although the specific role of this gene in apoptosis associated

with megakaryocyte differentiation is not clear yet (57).

**Discussions:** DENV replicates in multiple cell types and causes a range of pathological symptoms. Thrombocytopenia, a prominent DENV associated pathological symptom is a consequence of both impaired biogenesis and altered stability of platelets. Megakaryocytes or platelets mother cells in the bone marrow are known to be permissive for DENV replication, which affects platelet biogenesis. In this article, we have explored the cellular and molecular effect of DENV infection on Megakaryocyte development using the phorbol ester PMA induced-differentiation model in human K562 cells. Our results show that differentiation of infected K562 cells promote the replication of intracellular virus by an unknown mechanism and viral replication interferes with cell signaling pathways crucial for Megakaryopoiesis, particularly involving the MAPK ERK1/2 and different factors in the JAK-STAT pathway. However high level of viral replication does not affect the survival of the cells. PMA-induced differentiation of K562 cells, as with *in vivo* Megakaryopoiesis, is associated with accumulation of intracellular ROS and our results show that DENV replication suppresses the accumulation of this signaling molecule by an unknown mechanism. Replication by multiple viruses is known to increase ROS in the host cell we report an unique observation about DENV replication suppressing ROS accumulation under specific conditions in the host cell.

The absolute dependence of virus life cycle on host biochemical pathways implies a profound influence of host cell metabolic status on viral replication. Depending on the virus being studies, PMA a known activator of protein kinase C, is known to have either positive or negative effect on virus replication (58-61). On the other hand, viral infection of human monocytes can induce differentiation of these cells into Dendritic cells although the implication of this is not clear yet (62). Cell cycle arrest in PMA-treated K562 cells happens through upregulation of p21 and p27 in a protein kinase C (PKC)-dependent but p53-independent manner (63). Inducing an arrest in the host cell cycle is quite common among human viruses with beneficial effect on their replication (64). The entry of DENV is known to be influenced by the cell-cycle stage of the host cell, albeit in a cell-type dependent manner (65, 66). Here we observed increased intracellular replication in K562 cells that have stopped dividing in response to PMA, although the halt in cell division does not

lower the metabolic activity as is obvious from increase in cell size and other cellular changes. In a similar

manner Parvovirus B19 has been observed to infect cells of erythroblast origin with replication of the viral

genome being promoted by differentiation of these cells into erythroblasts (67). Although it is not clear yet

if the PMA-induced cell cycle stall is responsible for increasing intracellular virus replication, single-cell

transcriptomics study by Zanini and coworkers suggested the cell cycle status of the host cell to have no

effect on DENV genome replication (68). In addition to DENV, MK cells are known to support replication

of other viruses like the Human Immunodeficiency Virus and Hantavirus, with the former inducing

apoptosis in addition to reducing the surface expression of c-Mpl (69-72). Interestingly, PMA-induced

differentiation of Hantavirus infected megakaryocytes augmented virus replication, as observed in this

study (71).

PMA induces rapid and profound changes in host gene expression and therefore it is possible that

specific promoters of viral replication are overexpressed upon PMA treatment as observed for Hepatitis C

virus which is benefitted by the liver cell enriched microRNA-122 (73). PMA-induced differentiation

correlates with increase in granularity of K562 cells, probably pertaining to platelets granules that are

synthesized in the megakaryocyte mother cell and transferred into platelets (74). Therefore the rate of

protein synthesis in these cells can be expected to be enhanced for synthesis of the proteins that will form

part of these granules. In face the protein content of K562 cells is significantly increased during PMA

induced differentiation (75).It is possible that translation of viral proteins benefits from this overall increase

in rate of protein synthesis. The DENV genomic RNA of DENV has a 5' end type-I cap identical to host

mRNAs that helps in recruiting ribosomes in addition to evading innate antiviral arsenal (76). A role for

enhanced host protein synthesis benefiting viral translation is corroborated from our observation that the

entry of DENV is not higher in differentiated cells.  Another possibility is that differentiation leads to

suppression of specific anti-viral genes,  thereby benefiting the replication.

Although TPO is a cytokine crucial for platelet biogenesis and mice knock-out for either the

cytokine or its receptor show drastically reduced levels of both mature MKs as well as circulating platelets,

residual level of both MKs and platelets in these mice suggest other factors to also contribute in MK

development and platelet formation (77, 78). Receptor interaction of TPO activates JAK2 kinase leading to

activation of the MAPK ERK1/2, an important regulator of cell cycle and proliferation that plays a crucial

role in Mk maturation (79, 80) Previous studies of Mk development using K562 cells showed PMA to

directly activate protein kinase C (PKC) which in turn induces ERK1/2 phosphorylation, an important

signaling event even for *in vivo* Megakaryopoiesis (26, 50, 81, 82). A concurrent inhibition of the MAPK

p38 on the other hand supports MK development, suggesting this kinase to oppose or retard ERK1/2

mediated changes (22). Dengue infection can activate the p38 MAPK which is responsible for enhancement

of pro-inflammatory cytokines and apoptosis (83, 84). We observed a reduction in the level of PMA-

induced ERK1/2 phosphorylation in DENV-infected cells, probably causing inhibition of endomitosis and

platelet-specific surface marker protein expression, although a concomitant increase in p38 phosphorylation

was not observed. Previous reports have conclusively shown DENV NS5 protein mediated degradation of

the host STAT2 infection as a major mechanism for suppression of the host innate-antiviral pathway (85).

We observed additional regulation of a few other members of the JAK-STAT pathway, namely STAT1,

STAT3 and STAT5, although it is yet clear if this is restricted to PMA-treated K562 cells or is a general

effect occurring in other cells too. STAT1 regulated by the transcription factor (TF) GATA-1  plays a

specific role in endomitosis by controlling expression of cell cycle genes like CCND1, CCND2 and

CCNE2TFs through the TF RUNX1 (86). PMA-induced Mk development in K562 has not been reported

earlier to mimic the *in vivo* differentiation with respect to regulation of STAT3 and STAT5. In a manner

corroborative of *in vivo* development, we observed a gradual reduction of STAT5 level in parallel with

progression of differentiation (24). STAT3 is phosphorylated initially on Tyr-705 which triggers

dimerization (87). In addition to this STAT3 can have other post-translational modifications e.g.

phosphorylation of S-727 or acetylation of K-685, although induction of these modifications by PMA in

K562 cells is not clear yet (88, 89). In our study we observed DENV infection to reverse the PMA-induced

STAT3 phosphorylation on Tyr-705, possibly through inhibition of the respective kinase or activation

specific phosphatases. In addition to phosphorylation STAT3 function can be also suppressed by Ubiquitin

mediated degradation or interaction with proteins that are activated downstream of the JAK-STAT pathway

in a negative feedback loop e.g. protein encoded by PIAS and SOCS genes (90-92). In fact SOCS3 is known

as a specific  inhibitor of STAT3 signaling pathway, for which we expected an upregulation in infected

cells (89). Unexpectedly however, we observed a reduction in SOCS3 protein level by DENV infection for

which either mechanism or implication with respect to viral life cycle is still unclear. One of the potential

role for STAT3 in Megakaryopoiesis is driving accumulation of intracellular ROS levels through

upregulation of NOX2 expression (93). Therefore, suppression of STAT3 phosphorylation might be

directly affecting accumulation of ROS as observed here. Further characterization of this observation would

probably shed more light on the implication of this regulation.

Reactive oxygen species or ROS, in the form of Hydroxyl radical ($\cdot$OH) or Singlet oxygen ($^1O_2$) or

Superoxide anion radical ($O_2^{\cdot}-$), are generated either in response to stimuli including metabolic

inflammation, exposure to pathogen or as secondary messengers in signal transduction pathway. The most

well characterized sources of intracellular ROS are two enzymes associated with the mitochondria namely

**N**icotinamide **A**denine **D**inucleotide-**Q**uinone (NADH-Q) reductase (Complex I), Q-cytochrome c

oxidoreductase (Complex III) and the plasma membrane associated homologs of NADPH-oxidase or NOX

(94-96). Intracellular ROS plays crucial role in regulation of protein tyrosine kinases and phosphatases

through post-translational modification (97, 98). It can also activate multiple cellular pathways including

MAPK, NFκB, Cell cycle genes (99). Intracellular ROS can either promote apoptosis by inducing damage

to the mitochondrial membrane or inhibit it through oxidation of catalytic site Cysteines on executioner

proteases like Caspase-3 (100). In our study suppression of ROS level coincided with activation of Caspase-

3 cleavage activity, although the mechanism would need further investigation to draw a direct correlation

between them.

In the BM, pluripotent hematopoietic stem cells (HSCs) reside in a relatively hypoxic area while

during differentiation megakaryocytes relocate to more oxygen-rich region near blood vessels from where

platelets can be shed into the vascular circulation (101). ROS serves a critical role in megakaryocyte

formation since a low level of these signaling molecules in differentiating HSCs disfavor generation of

(**M**egakaryocyte-**E**rythrocyte **p**rogenitors) MEP and in favor of (**G**ranulocyte-**M**onocyte **p**rogenitors) GMP

(102). Intracellular ROS levels are regulated through the interplay of generators (like NOXs) and mitigatory

pathways (like the KEAP1-NRF2 pathway)  (37, 103). PMA has been shown to induce accumulation of

ROS in K562 cells and ROS has been shown to be responsible for polyploidy and a reduction of ROS by

NAC directly impacts polyploidy (43, 47). Accumulation of ROS activates the KEAP1-NRF2 pathway and

activation of many genes that can suppress the cellular ROS levels. ROS is produced by cellular responses

like Unfolded-protein response (UPR) or innate-antiviral pathway (104, 105). We observed a relatively

higher level of intracellular ROS with respect to uninfected cells in K562 cells that are not undergoing

differentiation. Since PMA-induced differentiation of K562 cells also causes ROS to accumulate in cells,

we expected an additive effect of DENV replication on ROS accumulation in these cells. However,

surprisingly DENV replication quenched intracellular ROS levels through an unknown mechanism.

Although evidence from previous reports would strongly indicate this suppression of ROS to be major

contributor to inhibition in platelet biogenesis by infection of Mk mother cells by DENV, it is still not clear

if the viral replication in these cells is benefited from this or not. Future studies would be directed to address

the exact mechanism through which DENV infection suppresses ROS accumulation in these cells and the

physiological relevance of this suppression for the viral life cycle.

**Acknowledgement:** The authors would like to thank Dr. Manjula Kalia, Regional Centre for
Biotechnology, NCR Biotech Science Cluster for careful reading of the manuscript. The Translational
Health Science and Technology Institute is acknowledged for providing all support for equipment and other
infrastructure. JK is supported by fellowship from the University grants commission. YR is supported from
fellowship in the extramural research grant (BT/PR22985/MED/29/1168/2016) to SB from the Department
of Biotechnology, Govt of India. This work was supported by extramural research grant
(EMR/2016/005796) to SB from the Science and Engineering Research Board, Department of Science and
Technology, Govt. of India.

**Materials and methods**:

**Cell culture, Drugs and Virus**: K562 cells were procured from the American Type Culture Collection (ATCC) and cultured at $37^0$C, 5% $CO_2$ in Iscove's modified Dulbecco's media (IMDM) supplemented with Penicillin (100 U/ml), Streptomycin (0.1 mg/ml) and 10%-Fetal Bovine Serum (FBS). Vero and C6/36 cells were procured from the cell line repository of the National Centre for Cell Sciences (NCCS), India and cultured respectively in Minimum Essential Medium (MEM) and L15 cell culture medium supplemented with 10% FBS. C6/36 cells were maintained at conditions of $28^0$C and atmospheric $CO_2$.

Dengue virus serotype 2 (strain NGC) was amplified in C6/36 and the culture supernatant used as inoculum. The infectious titer of the virus was determined by Focus-forming unit (FFU) assay in Vero cells. For all infections, the inoculum was diluted in the respective culture media supplemented with 2% FBS, and the cells were incubated with the inoculum for 2 hours at the respective culture conditions of temperature and $CO_2$ concentration, with intermittent rocking.

Phorbol-12 Myristate-13 acetate (PMA, Sigma Aldrich) and N-Acetyl Cysteine were diluted in recommended vehicles and stored as single use aliquots at $-20^0$C .

**Focus-forming unit (FFU) assay**: Vero cell monolayers in 24-well plate were infected with 10-fold serial dilutions of virus inoculum. The inoculum was incubated with cells for 2 hours at $37^0$C and 5% $CO_2$ with intermittent rocking. The inoculum was then discarded and complete MEM added to each followed by incubation for 48 hours at $37^0$C and 5% $CO_2$. Subsequently the cells were washed with phosphate-buffered saline (PBS), fixed with 2% para-formaldehyde (PFA) and permeabilized with PBS supplemented with 0.1 % Triton-X-100 and 1% bovine serum albumin (BSA). The permeabilized cells were washed with PBS and sequentially incubated anti-DENV primary antibody (dilution 1:400 of Mab8705, Millipore) and anti-Mouse Alexa-488 conjugated secondary antibody (dilution 1:500; ThermoFisher Scientific) both diluted in PBS supplemented with 1% BSA. After incubation with primary antibody, the cells were washed twice

with PBS before addition of secondary antibody dilutions. The fluorescent foci were visualized and

manually counted using a fluorescence microscope and the titer calculated as FFU/ml.

**Flow Cytometry Analysis**: All flow cytometry analysis were performed in a BD FACS Canto II flow

cytometer (BD Biosciences) under standard conditions and all raw data analyzed using Flow-Jo software.

For immune staining of intracellular DENV antigen, K562 cells were washed with ice-cold PBS before

fixation with PBS supplemented with 2% PFA and permeabilization with the same buffer containing 0.1%

Triton-X-100. The permeabilized cells were sequentially stained with dilution of 4G2 (purified IgG2a mAb

produced in the lab from Hybridoma-HB112) antibody and anti-mouse Alexa-488 conjugated secondary

antibody. For immune staining of surface markers, cells were washed with PBS and incubated with dilutions

of fluor-conjugated primary antibodies, specific to either human CD61-Phycoerythrin (PE) (BioLegend) or

CD41/61-(Allophycocyanin) APC (BioLegend) or human CD42b-(BD Horizon brilliant violet 421) Bv421

(BioLegend). Antibodies were diluted in staining buffer (PBS with 1%BSA and 0.02% Sodium Azide) and

cells were incubated for 1h at 4°C . Subsequently the cells were washed with PBS and analyzed by flow

cytometry.

Propidium iodide staining for polyploidy analysis: Cells were harvested by centrifugation, washed

twice with ice-cold phosphate-buffered saline (PBS) and fixed with 70% ethanol at 4°C for 30 min. After

a PBS wash cells were treated with 100μl PBS supplemented with RNase-A (200μg/ml) and incubated at

37°C for 30 min. Cells were then washed using PBS and the genomic DNA stained with Propidium iodide

(50μg/ml).

Annexin-V and PI staining: Apoptotic detection assays were carried out by surface labeling with the Ca2+

dependent phosphatidylserine-binding protein Annexin-V. Cells were harvested by centrifugation, washed

twice with cold PBS and then re-suspended in 1x binding buffer(Apoptosis detection Kit I # 556578, BD

Pharmingen) at 1x $10^6$ cells/ml. A 100μl aliquot was stained by addition of 5μl FITC Annexin V and 5μl

PI.  The reagents were mixed by gentle vortex and incubated for 15 min at RT in the dark. 400µl of 1x

binding buffer was added to each tube and the cells analyzed by flow cytometry within 1 hr.

**RNA extraction and Quantitative RT-PCR**: Total RNA from $1x10^6$ K562 cells were extracted using

Trizol (Takara) and purified with Qiagen RNAeasy mini kit (Qiagen), as per manufacturer's instructions.

Viral RNA from cell-free culture supernatant was isolated using QIAmp viral RNA mini kit (Qiagen) as

per manufacturer's instructions. 1.0 ug of total RNA was reverse-transcribed with ImProm-II reverse-

transcriptase (Promega, USA) and random hexamers (Sigma) as per manufacturer's protocol, and the cDNA

diluted with nuclease-free water before use for real-time PCR. Real-time PCR was performed with 2x

SYBR mix (Takara) in a QuantStudio-6 Flex Real-Time PCR System (Applied Biosystems) using the

default run program.

**Confocal microscopy**: K562 cells were washed with PBS, fixed and permeabilized as described earlier for

immunostaining purposes. The cells were stained with 2.5µl of Alexa Fluor-568 Phalloidin (Invitrogen,

Thermo fisher scientific) at RT for 20 min to stain for F-Actin followed by PBS wash. Cells were then

mounted onto glass slides using ProLong Gold Antifade Mountant supplemented with DAPI (Invitrogen,

Thermo fisher scientific). The fluorescence was observed and imaged in a FLUOVIEW FV3000 confocal

microscope (Olympus).

**Intracellular Caspase-3/7 activity assay**: The intracellular caspase-3/7 activity was measured using

Caspase-Glo 3/7 Assay (Promega). For this 100 µl of cell suspension and 50 µl Caspase-Glo® 3/7 reagent

were mixed in wells of an opaque white 96-well plate.  The plates were incubated  at RT for 3 hours and

the luminescence measured using a Orion II microplate luminometer (Berthold).  The luminescence from

culture media without cells was used as negative control.

**Measurement of intracellular Reactive Oxygen species (ROS)**: The intracellular ROS accumulated was

measure using CM-H2DCFDA probe (Life Technologies) as per manufacturer's instructions. Briefly, K562

cells were washed with PBS and then incubated in PBS supplemented with CM-H2DCFDA as per

manufacturer's instructions. The cells were then incubated at 37 °C for 30 min and washed twice with PBS.

The fluorescence intensity was quantified in a BD FACS Canto II flow cytometer (BD Biosciences) under

standard conditions and the raw data analyzed using Flow-Jo software.

**Electrophoresis and Immunoblotting**: Cells were harvested by centrifugation and washed with PBS.

Whole-cell lysates were prepared using cell lysis solution [250 mmol/L NaCl, 20 mmol/L Tris-HCl (pH

7.4), , 1% Triton X-100, 1 mmol/L EDTA,20µl PMSF, 10µl PI], followed by centrifugation (13 000×g, 10

min). The proteins in 40µg of whole-cell lysate was denatured, resolved in a SDS-10% gel and transferred

to nitrocellulose blotting membrane. The membrane was blocked with either 5% skimmed milk or 5%

bovine-serum albumin (BSA) and incubated with dilution of respective antibodies as per manufacturer's

instructions (Cell Signaling technologies). The membrane was washed, incubated with dilution of HRP-

conjugated secondary antibody and the bands visualized by ECL chemiluminescence.

**References:**

1.    Patel SR, Hartwig JH, Italiano JE, Jr. 2005. The biogenesis of platelets from megakaryocyte
proplatelets. J Clin Invest 115:3348-54.
2.    Hottz ED, Bozza FA, Bozza PT. 2018. Platelets in Immune Response to Virus and Immunopathology
of Viral Infections. Front Med (Lausanne) 5:121.
3.    Martina BE, Koraka P, Osterhaus AD. 2009. Dengue virus pathogenesis: an integrated view. Clin
Microbiol Rev 22:564-81.
4.    de Azeredo EL, Monteiro RQ, de-Oliveira Pinto LM. 2015. Thrombocytopenia in Dengue:
Interrelationship between Virus and the Imbalance between Coagulation and Fibrinolysis and
Inflammatory Mediators. Mediators Inflamm 2015:313842.
5.    Fang S, Wu Y, Wu N, Zhang J, An J. 2013. Recent advances in DENV receptors.
ScientificWorldJournal 2013:684690.
6.    Rodenhuis-Zybert IA, Wilschut J, Smit JM. 2010. Dengue virus life cycle: viral and host factors
modulating infectivity. Cell Mol Life Sci 67:2773-86.
7.    Acosta EG, Kumar A, Bartenschlager R. 2014. Revisiting dengue virus-host cell interaction: new
insights into molecular and cellular virology. Adv Virus Res 88:1-109.
8.    Simon AY, Sutherland MR, Pryzdial EL. 2015. Dengue virus binding and replication by platelets.
Blood 126:378-85.
9.    Ojha A, Nandi D, Batra H, Singhal R, Annarapu GK, Bhattacharyya S, Seth T, Dar L, Medigeshi GR,
Vrati S, Vikram NK, Guchhait P. 2017. Platelet activation determines the severity of
thrombocytopenia in dengue infection. Sci Rep 7:41697.
10.   Ojha A, Bhasym A, Mukherjee S, Annarapu GK, Bhakuni T, Akbar I, Seth T, Vikram NK, Vrati S, Basu
A, Bhattacharyya S, Guchhait P. 2019. Platelet factor 4 promotes rapid replication and
propagation of Dengue and Japanese encephalitis viruses. EBioMedicine 39:332-347.
11.   Noisakran S, Onlamoon N, Hsiao HM, Clark KB, Villinger F, Ansari AA, Perng GC. 2012. Infection of
bone marrow cells by dengue virus in vivo. Exp Hematol 40:250-259 e4.
12.   Vogt MB, Lahon A, Arya RP, Spencer Clinton JL, Rico-Hesse R. 2019. Dengue viruses infect human
megakaryocytes, with probable clinical consequences. PLoS Negl Trop Dis 13:e0007837.
13.   Clark KB, Noisakran S, Onlamoon N, Hsiao HM, Roback J, Villinger F, Ansari AA, Perng GC. 2012.
Multiploid CD61+ cells are the pre-dominant cell lineage infected during acute dengue virus
infection in bone marrow. PLoS One 7:e52902.
14.   Sridharan A, Chen Q, Tang KF, Ooi EE, Hibberd ML, Chen J. 2013. Inhibition of megakaryocyte
development in the bone marrow underlies dengue virus-induced thrombocytopenia in
humanized mice. J Virol 87:11648-58.
15.   Basu A, Jain P, Gangodkar SV, Shetty S, Ghosh K. 2008. Dengue 2 virus inhibits in vitro
megakaryocytic colony formation and induces apoptosis in thrombopoietin-inducible
megakaryocytic differentiation from cord blood CD34+ cells. FEMS Immunol Med Microbiol 53:46-
51.
16.   Akashi K, Traver D, Miyamoto T, Weissman IL. 2000. A clonogenic common myeloid progenitor
that gives rise to all myeloid lineages. Nature 404:193-7.
17.   Kaushansky K. 2008. Historical review: megakaryopoiesis and thrombopoiesis. Blood 111:981-6.
18.   Deutsch VR, Tomer A. 2006. Megakaryocyte development and platelet production. Br J Haematol
134:453-66.
19.   Schulze H, Shivdasani RA. 2004. Molecular mechanisms of megakaryocyte differentiation. Semin
Thromb Hemost 30:389-98.

20.   Cecchetti L, Tolley ND, Michetti N, Bury L, Weyrich AS, Gresele P. 2011. Megakaryocytes
differentially sort mRNAs for matrix metalloproteinases and their inhibitors into platelets: a
mechanism for regulating synthetic events. Blood 118:1903-11.
21.   Matsumura I, Kanakura Y. 2002. Molecular control of megakaryopoiesis and thrombopoiesis. Int
J Hematol 75:473-83.
22.   Severin S, Ghevaert C, Mazharian A. 2010. The mitogen-activated protein kinase signaling
pathways: role in megakaryocyte differentiation. J Thromb Haemost 8:17-26.
23.   Kirito K, Osawa M, Morita H, Shimizu R, Yamamoto M, Oda A, Fujita H, Tanaka M, Nakajima K,
Miura Y, Ozawa K, Komatsu N. 2002. A functional role of Stat3 in in vivo megakaryopoiesis. Blood
99:3220-7.
24.   Olthof SG, Fatrai S, Drayer AL, Tyl MR, Vellenga E, Schuringa JJ. 2008. Downregulation of signal
transducer and activator of transcription 5 (STAT5) in CD34+ cells promotes megakaryocytic
development, whereas activation of STAT5 drives erythropoiesis. Stem Cells 26:1732-42.
25.   Beaulieu LM, Lin E, Morin KM, Tanriverdi K, Freedman JE. 2011. Regulatory effects of TLR2 on
megakaryocytic cell function. Blood 117:5963-74.
26.   Racke FK, Lewandowska K, Goueli S, Goldfarb AN. 1997. Sustained activation of the extracellular
signal-regulated kinase/mitogen-activated protein kinase pathway is required for megakaryocytic
differentiation of K562 cells. J Biol Chem 272:23366-70.
27.   Goldfarb AN. 2007. Transcriptional control of megakaryocyte development. Oncogene 26:6795-
802.
28.   Tijssen MR, Ghevaert C. 2013. Transcription factors in late megakaryopoiesis and related platelet
disorders. J Thromb Haemost 11:593-604.
29.   Muntean AG, Pang L, Poncz M, Dowdy SF, Blobel GA, Crispino JD. 2007. Cyclin D-Cdk4 is regulated
by GATA-1 and required for megakaryocyte growth and polyploidization. Blood 109:5199-207.
30.   Lordier L, Bluteau D, Jalil A, Legrand C, Pan J, Rameau P, Jouni D, Bluteau O, Mercher T, Leon C,
Gachet C, Debili N, Vainchenker W, Raslova H, Chang Y. 2012. RUNX1-induced silencing of non-
muscle myosin heavy chain IIB contributes to megakaryocyte polyploidization. Nat Commun
3:717.
31.   Lopez JJ, Palazzo A, Chaabane C, Albarran L, Polidano E, Lebozec K, Dally S, Nurden P, Enouf J,
Debili N, Bobe R. 2013. Crucial role for endoplasmic reticulum stress during megakaryocyte
maturation. Arterioscler Thromb Vasc Biol 33:2750-8.
32.   You T, Wang Q, Zhu L. 2016. Role of autophagy in megakaryocyte differentiation and platelet
formation. Int J Physiol Pathophysiol Pharmacol 8:28-34.
33.   De Botton S, Sabri S, Daugas E, Zermati Y, Guidotti JE, Hermine O, Kroemer G, Vainchenker W,
Debili N. 2002. Platelet formation is the consequence of caspase activation within
megakaryocytes. Blood 100:1310-7.
34.   Chen S, Su Y, Wang J. 2013. ROS-mediated platelet generation: a microenvironment-dependent
manner for megakaryocyte proliferation, differentiation, and maturation. Cell Death Dis 4:e722.
35.   Sardina JL, Lopez-Ruano G, Sanchez-Abarca LI, Perez-Simon JA, Gaztelumendi A, Trigueros C,
Llanillo M, Sanchez-Yague J, Hernandez-Hernandez A. 2010. p22phox-dependent NADPH oxidase
activity is required for megakaryocytic differentiation. Cell Death Differ 17:1842-54.
36.   Mostafa SS, Miller WM, Papoutsakis ET. 2000. Oxygen tension influences the differentiation,
maturation and apoptosis of human megakaryocytes. Br J Haematol 111:879-89.
37.   McCrann DJ, Eliades A, Makitalo M, Matsuno K, Ravid K. 2009. Differential expression of NADPH
oxidases in megakaryocytes and their role in polyploidy. Blood 114:1243-9.
38.   Sattler M, Winkler T, Verma S, Byrne CH, Shrikhande G, Salgia R, Griffin JD. 1999. Hematopoietic
growth factors signal through the formation of reactive oxygen species. Blood 93:2928-35.

39.   Eliades A, Papadantonakis N, Ravid K. 2010. New roles for cyclin E in megakaryocytic
polyploidization. J Biol Chem 285:18909-17.
40.   Siddiqui NF, Shabrani NC, Kale VP, Limaye LS. 2011. Enhanced generation of megakaryocytes from
umbilical cord blood-derived CD34(+) cells expanded in the presence of two nutraceuticals,
docosahexaenoic acid and arachidonic acid, as supplements to the cytokine-containing medium.
Cytotherapy 13:114-28.
41.   Witt O, Sand K, Pekrun A. 2000. Butyrate-induced erythroid differentiation of human K562
leukemia cells involves inhibition of ERK and activation of p38 MAP kinase pathways. Blood
95:2391-6.
42.   Kim KW, Kim SH, Lee EY, Kim ND, Kang HS, Kim HD, Chung BS, Kang CD. 2001. Extracellular signal-
regulated kinase/90-KDA ribosomal S6 kinase/nuclear factor-kappa B pathway mediates phorbol
12-myristate 13-acetate-induced megakaryocytic differentiation of K562 cells. J Biol Chem
276:13186-91.
43.   Ojima Y, Duncan MT, Nurhayati RW, Taya M, Miller WM. 2013. Synergistic effect of hydrogen
peroxide on polyploidization during the megakaryocytic differentiation of K562 leukemia cells by
PMA. Exp Cell Res 319:2205-15.
44.   Chaman N, Iqbal MA, Siddiqui FA, Gopinath P, Bamezai RN. 2015. ERK2-Pyruvate Kinase Axis
Permits Phorbol 12-Myristate 13-Acetate-induced Megakaryocyte Differentiation in K562 Cells. J
Biol Chem 290:23803-15.
45.   Lee CH, Yun HJ, Kang HS, Kim HD. 1999. ERK/MAPK pathway is required for changes of cyclin D1
and B1 during phorbol 12-myristate 13-acetate-induced differentiation of K562 cells. IUBMB Life
48:585-91.
46.   Nurhayati RW, Ojima Y, Nomura N, Taya M. 2014. Promoted megakaryocytic differentiation of
K562 cells through oxidative stress caused by near ultraviolet irradiation. Cell Mol Biol Lett 19:590-
600.
47.   Huang R, Zhao L, Chen H, Yin RH, Li CY, Zhan YQ, Zhang JH, Ge CH, Yu M, Yang XM. 2014.
Megakaryocytic differentiation of K562 cells induced by PMA reduced the activity of respiratory
chain complex IV. PLoS One 9:e96246.
48.   Colosetti P, Puissant A, Robert G, Luciano F, Jacquel A, Gounon P, Cassuto JP, Auberger P. 2009.
Autophagy is an important event for megakaryocytic differentiation of the chronic myelogenous
leukemia K562 cell line. Autophagy 5:1092-8.
49.   Limb JK, Yoon S, Lee KE, Kim BH, Lee S, Bae YS, Jhon GJ, Kim J. 2009. Regulation of megakaryocytic
differentiation of K562 cells by FosB, a member of the Fos family of AP-1 transcription factors. Cell
Mol Life Sci 66:1962-73.
50.   Rojnuckarin P, Drachman JG, Kaushansky K. 1999. Thrombopoietin-induced activation of the
mitogen-activated protein kinase (MAPK) pathway in normal megakaryocytes: role in
endomitosis. Blood 94:1273-82.
51.   Lin YL, Liao CL, Chen LK, Yeh CT, Liu CI, Ma SH, Huang YY, Huang YL, Kao CL, King CC. 1998. Study
of Dengue virus infection in SCID mice engrafted with human K562 cells. J Virol 72:9729-37.
52.   Elshuber S, Allison SL, Heinz FX, Mandl CW. 2003. Cleavage of protein prM is necessary for
infection of BHK-21 cells by tick-borne encephalitis virus. J Gen Virol 84:183-191.
53.   Obexer P, Hagenbuchner J, Unterkircher T, Sachsenmaier N, Seifarth C, Bock G, Porto V, Geiger K,
Ausserlechner M. 2009. Repression of BIRC5/survivin by FOXO3/FKHRL1 sensitizes human
neuroblastoma cells to DNA damage-induced apoptosis. Mol Biol Cell 20:2041-8.
54.   Caparros-Perez E, Teruel-Montoya R, Lopez-Andreo MJ, Llanos MC, Rivera J, Palma-Barqueros V,
Blanco JE, Vicente V, Martinez C, Ferrer-Marin F. 2017. Comprehensive comparison of neonate
and adult human platelet transcriptomes. PLoS One 12:e0183042.

55.    Majek P, Reicheltova Z, Stikarova J, Suttnar J, Sobotkova A, Dyr JE. 2010. Proteome changes in
661             platelets activated by arachidonic acid, collagen, and thrombin. Proteome Sci 8:56.

662    56.    Park SM, Jung HY, Kim HO, Rhim H, Paik SR, Chung KC, Park JH, Kim J. 2002. Evidence that alpha-
663             synuclein functions as a negative regulator of Ca(++)-dependent alpha-granule release from
664             human platelets. Blood 100:2506-14.

665    57.    da Costa CA, Ancolio K, Checler F. 2000. Wild-type but not Parkinson's disease-related ala-53 -->
666             Thr mutant alpha -synuclein protects neuronal cells from apoptotic stimuli. J Biol Chem
667             275:24065-9.

668    58.    Polyak SJ, Rawls WE, Harnish DG. 1991. Characterization of Pichinde virus infection of cells of the
669             monocytic lineage. J Virol 65:3575-82.

670    59.    Seamone ME, Wang W, Acott P, Beck PL, Tibbles LA, Muruve DA. 2010. MAP kinase activation
671             increases BK polyomavirus replication and facilitates viral propagation in vitro. J Virol Methods
672             170:21-9.

673    60.    Wang W, Wang Y, Debing Y, Zhou X, Yin Y, Xu L, Herrera Carrillo E, Brandsma JH, Poot RA, Berkhout
674             B, Neyts J, Peppelenbosch MP, Pan Q. 2017. Biological or pharmacological activation of protein
675             kinase C alpha constrains hepatitis E virus replication. Antiviral Res 140:1-12.

676    61.    Golding H, Manischewitz J, Vujcic L, Blumenthal R, Dimitrov DS. 1994. The phorbol ester phorbol
677             myristate acetate inhibits human immunodeficiency virus type 1 envelope-mediated fusion by
678             modulating an accessory component(s) in CD4-expressing cells. J Virol 68:1962-9.

679    62.    Hou W, Gibbs JS, Lu X, Brooke CB, Roy D, Modlin RL, Bennink JR, Yewdell JW. 2012. Viral infection
680             triggers rapid differentiation of human blood monocytes into dendritic cells. Blood 119:3128-31.

681    63.    Zeng YX, el-Deiry WS. 1996. Regulation of p21WAF1/CIP1 expression by p53-independent
682             pathways. Oncogene 12:1557-64.

683    64.    Bagga S, Bouchard MJ. 2014. Cell cycle regulation during viral infection. Methods Mol Biol
684             1170:165-227.

685    65.    Phoolcharoen W, Smith DR. 2004. Internalization of the dengue virus is cell cycle modulated in
686             HepG2, but not Vero cells. J Med Virol 74:434-41.

687    66.    Helt AM, Harris E. 2005. S-phase-dependent enhancement of dengue virus 2 replication in
688             mosquito cells, but not in human cells. J Virol 79:13218-30.

689    67.    Bua G, Manaresi E, Bonvicini F, Gallinella G. 2016. Parvovirus B19 Replication and Expression in
690             Differentiating Erythroid Progenitor Cells. PLoS One 11:e0148547.

691    68.    Zanini F, Pu SY, Bekerman E, Einav S, Quake SR. 2018. Single-cell transcriptional dynamics of
692             flavivirus infection. Elife 7.

693    69.    Zauli G, Catani L, Gibellini D, Re MC, Milani D, Borgatti P, Bassini A, La Placa M, Capitani S. 1995.
694             The CD4 receptor plays essential but distinct roles in HIV-1 infection and induction of apoptosis in
695             primary bone marrow GPIIb/IIIa+ megakaryocytes and the HEL cell line. Br J Haematol 91:290-8.

696    70.    Zauli G, Catani L, Gibellini D, Re MC, Vianelli N, Colangeli V, Celeghini C, Capitani S, La Placa M.
697             1996. Impaired survival of bone marrow GPIIb/IIa+ megakaryocytic cells as an additional
698             pathogenetic mechanism of HIV-1-related thrombocytopenia. Br J Haematol 92:711-7.

699    71.    Lutteke N, Raftery MJ, Lalwani P, Lee MH, Giese T, Voigt S, Bannert N, Schulze H, Kruger DH,
700             Schonrich G. 2010. Switch to high-level virus replication and HLA class I upregulation in
701             differentiating megakaryocytic cells after infection with pathogenic hantavirus. Virology 405:70-
702             80.

703    72.    Zhang M, Evans S, Yuan J, Ratner L, Koka PS. 2010. HIV-1 determinants of thrombocytopenia at
704             the stage of CD34+ progenitor cell differentiation in vivo lie in the viral envelope gp120 V3 loop
705             region. Virology 401:131-6.

706    73.    Henke JI, Goergen D, Zheng J, Song Y, Schuttler CG, Fehr C, Junemann C, Niepmann M. 2008.
707             microRNA-122 stimulates translation of hepatitis C virus RNA. EMBO J 27:3300-10.

708    74.    Ambrosio AL, Di Pietro SM. 2019. Mechanism of platelet alpha-granule biogenesis: study of cargo
709           transport and the VPS33B-VPS16B complex in a model system. Blood Adv 3:2617-2626.
710    75.    Baatout S, Chatelain B, Staquet P, Symann M, Chatelain C. 1999. Protein content and number of
711           nucleolar organizer regions are enhanced during phorbol ester-induced differentiation of cultured
712           human megakaryocytic cells. Anticancer Res 19:3229-35.
713    76.    Daffis S, Szretter KJ, Schriewer J, Li J, Youn S, Errett J, Lin TY, Schneller S, Zust R, Dong H, Thiel V,
714           Sen GC, Fensterl V, Klimstra WB, Pierson TC, Buller RM, Gale M, Jr., Shi PY, Diamond MS. 2010. 2'-
715           O methylation of the viral mRNA cap evades host restriction by IFIT family members. Nature
716           468:452-6.
717    77.    Gurney AL, de Sauvage FJ. 1996. Dissection of c-Mpl and thrombopoietin function: studies of
718           knockout mice and receptor signal transduction. Stem Cells 14 Suppl 1:116-23.
719    78.    Kaushansky K. 2005. The molecular mechanisms that control thrombopoiesis. J Clin Invest
720           115:3339-47.
721    79.    Roux PP, Blenis J. 2004. ERK and p38 MAPK-activated protein kinases: a family of protein kinases
722           with diverse biological functions. Microbiol Mol Biol Rev 68:320-44.
723    80.    Fichelson S, Freyssinier JM, Picard F, Fontenay-Roupie M, Guesnu M, Cherai M, Gisselbrecht S,
724           Porteu F. 1999. Megakaryocyte growth and development factor-induced proliferation and
725           differentiation are regulated by the mitogen-activated protein kinase pathway in primitive cord
726           blood hematopoietic progenitors. Blood 94:1601-13.
727    81.    Herrera R, Hubbell S, Decker S, Petruzzelli L. 1998. A role for the MEK/MAPK pathway in PMA-
728           induced cell cycle arrest: modulation of megakaryocytic differentiation of K562 cells. Exp Cell Res
729           238:407-14.
730    82.    Whalen AM, Galasinski SC, Shapiro PS, Nahreini TS, Ahn NG. 1997. Megakaryocytic differentiation
731           induced by constitutive activation of mitogen-activated protein kinase kinase. Mol Cell Biol
732           17:1947-58.
733    83.    Nagila A, Netsawang J, Suttitheptumrong A, Morchang A, Khunchai S, Srisawat C, Puttikhunt C,
734           Noisakran S, Yenchitsomanus PT, Limjindaporn T. 2013. Inhibition of p38MAPK and CD137
735           signaling reduce dengue virus-induced TNF-alpha secretion and apoptosis. Virol J 10:105.
736    84.    Sreekanth GP, Chuncharunee A, Sirimontaporn A, Panaampon J, Noisakran S, Yenchitsomanus PT,
737           Limjindaporn T. 2016. SB203580 Modulates p38 MAPK Signaling and Dengue Virus-Induced Liver
738           Injury by Reducing MAPKAPK2, HSP27, and ATF2 Phosphorylation. PLoS One 11:e0149486.
739    85.    Ashour J, Laurent-Rolle M, Shi PY, Garcia-Sastre A. 2009. NS5 of dengue virus mediates STAT2
740           binding and degradation. J Virol 83:5408-18.
741    86.    Huang Z, Richmond TD, Muntean AG, Barber DL, Weiss MJ, Crispino JD. 2007. STAT1 promotes
742           megakaryopoiesis downstream of GATA-1 in mice. J Clin Invest 117:3890-9.
743    87.    Yoshimura A, Naka T, Kubo M. 2007. SOCS proteins, cytokine signalling and immune regulation.
744           Nat Rev Immunol 7:454-65.
745    88.    Yuan ZL, Guan YJ, Chatterjee D, Chin YE. 2005. Stat3 dimerization regulated by reversible
746           acetylation of a single lysine residue. Science 307:269-73.
747    89.    Rebe C, Vegran F, Berger H, Ghiringhelli F. 2013. STAT3 activation: A key factor in tumor
748           immunoescape. JAKSTAT 2:e23010.
749    90.    Wei J, Yuan Y, Jin C, Chen H, Leng L, He F, Wang J. 2012. The ubiquitin ligase TRAF6 negatively
750           regulates the JAK-STAT signaling pathway by binding to STAT3 and mediating its ubiquitination.
751           PLoS One 7:e49567.
752    91.    Shuai K. 2000. Modulation of STAT signaling by STAT-interacting proteins. Oncogene 19:2638-44.
753    92.    Krebs DL, Hilton DJ. 2001. SOCS proteins: negative regulators of cytokine signaling. Stem Cells
754           19:378-87.

93. Corzo CA, Cotter MJ, Cheng P, Cheng F, Kusmartsev S, Sotomayor E, Padhya T, McCaffrey TV,
McCaffrey JC, Gabrilovich DI. 2009. Mechanism regulating reactive oxygen species in tumor-
induced myeloid-derived suppressor cells. J Immunol 182:5693-701.
94. Lambeth JD. 2004. NOX enzymes and the biology of reactive oxygen. Nat Rev Immunol 4:181-9.
95. Zorov DB, Juhaszova M, Sollott SJ. 2014. Mitochondrial reactive oxygen species (ROS) and ROS-
induced ROS release. Physiol Rev 94:909-50.
96. Turrens JF. 2003. Mitochondrial formation of reactive oxygen species. J Physiol 552:335-44.
97. Salmeen A, Andersen JN, Myers MP, Meng TC, Hinks JA, Tonks NK, Barford D. 2003. Redox
regulation of protein tyrosine phosphatase 1B involves a sulphenyl-amide intermediate. Nature
423:769-73.
98. Lee SR, Kwon KS, Kim SR, Rhee SG. 1998. Reversible inactivation of protein-tyrosine phosphatase
1B in A431 cells stimulated with epidermal growth factor. J Biol Chem 273:15366-72.
99. Zhang J, Wang X, Vikash V, Ye Q, Wu D, Liu Y, Dong W. 2016. ROS and ROS-Mediated Cellular
Signaling. Oxid Med Cell Longev 2016:4350965.
100. Choi K, Ryu SW, Song S, Choi H, Kang SW, Choi C. 2010. Caspase-dependent generation of reactive
oxygen species in human astrocytoma cells contributes to resistance to TRAIL-mediated
apoptosis. Cell Death Differ 17:833-45.
101. Huang X, Trinh T, Aljoufi A, Broxmeyer HE. 2018. Hypoxia Signaling Pathway in Stem Cell
Regulation: Good and Evil. Curr Stem Cell Rep 4:149-157.
102. Shinohara A, Imai Y, Nakagawa M, Takahashi T, Ichikawa M, Kurokawa M. 2014. Intracellular
reactive oxygen species mark and influence the megakaryocyte-erythrocyte progenitor fate of
common myeloid progenitors. Stem Cells 32:548-57.
103. Motohashi H, Kimura M, Fujita R, Inoue A, Pan X, Takayama M, Katsuoka F, Aburatani H, Bresnick
EH, Yamamoto M. 2010. NF-E2 domination over Nrf2 promotes ROS accumulation and
megakaryocytic maturation. Blood 115:677-86.
104. Schwarz KB. 1996. Oxidative stress during viral infection: a review. Free Radic Biol Med 21:641-9.
105. Zeeshan HM, Lee GH, Kim HR, Chae HJ. 2016. Endoplasmic Reticulum Stress and Associated ROS.
Int J Mol Sci 17:327.

**Figure legends:**

**Figure 1: Panel A: Schematic diagram of Megakaryopoiesis**: The schematic diagram shows some of the lineages that are produced during differentiation of Hematopoietic stem cells (HSC) to Megakaryocytes (Mk). The formation of multilobed nucleus has also been depicted. **Panel B: Visualization of endomitosis and multi-lobed nuclear formation following PMA-treatment.** K562 cells treated with either DMSO or 100 nM PMA for 6 days were stained with Alexa Fluor-568 Phalloidin, mounted onto glass slides using mountant supplemented with DAPI and visualized in a confocal microscope. The red color indicates the plasma membrane and blue color shows nuclei. The inset shows a cell with multilobed nucleus. **Panel C**: **Expression of platelet specific surface markers:** K562 cells treated with PMA for 0, 3 or 6 days were immune-stained for expression of the indicated surface marker protein using fluor-conjugated primary antibodies. The mean fluorescence intensity (MFI) corresponding to each day was quantified. For each surface marker the MFI at day 0 was arbitrarily set to 1 and those at day 3 and day 6, expressed as the relative fold change with respect to that. The error bars represent standard deviation. The significance was calculated by Student's t-test (*, **, *** respectively indicate P-values <0.5, <0.1 and <0.01). **Panel D**: K562 cells treated with PMA for 0 or 3 or 6 days were fixed, permeabilized and stained for intracellular DNA using Propidium iodide (PI). The PI stain was quantified by flow cytometry and mean number of cells having ploidy of >4N was plotted. The error bars represent standard deviation. **Panel E:** The total RNA from K562 cells treated with either DMSO or 100nM PMA or Sodium Butyrate (NaB) for 3 or 6 days was extracted and purified. The total RNA was reverse transcribed using random hexamers and the cDNA obtained used for real-time PCR estimation of mRNA transcripts from indicated genes. The $C_T$ value corresponding to each was normalized to that of GAPDH. The normalized value at day 0 was taken arbitrarily as 1 and those at day 3 and 6 expressed as fold-change with respect to that.

**Figure 2: DENV replication is increased upon PMA treatment of infected K562 cells: Panel A:** K562

cells infected with DENV at an MOI of 0.1 were maintained in growth media supplemented with either

DMSO or 100nM PMA and fixed with PFA at 0 hour or 3 days or 6 days post-infection. The cells were

permeabilized and immune-stained for intracellular DENV antigen and the fluorescence quantified by flow

cytometry. The fluorescence in cells fixed at 0 hour was used to gate for DENV antigen positive cells. The

number of antigen positive cells were quantified and plotted. The error bars represent standard deviation

and the significance was calculated by Student's t-test (*** indicates P <0.001). **Panel B**: Total RNA from

K562 cells treated similarly as in panel A was extracted, purified and reverse-transcribed using random

hexamers. The cDNA was used for real-time PCR comparison of DENV RNA level at 3 or 6 days to that

at 0 hour, normalized to that of GAPDH mRNA. The value at 0 hour was taken arbitrarily as 1 and that at

other time points expressed as fold-change over that. The error bars represent standard deviation and the

significance was calculated by Student's t-test (** indicates P <0.01). **Panel C**: The culture supernatant of

cell infected with MOI of 0.1 and subsequently treated as in panel A was collected and used for

quantification of focus-forming unit/ml (FFU/ml) on Vero cell monolayers. The FFU/ml for each sample

was transformed to their logarithmic value to the base of 10 and plotted. The error bars represent standard

deviation and the significance was calculated by Student's t-test (** indicates P <0.01). **Panel D**:

Uninfected K562 cells treated with either DMSO or PMA for 3 days were washed with PBS and incubated

in either mock- or DENV inoculum (10 MOI) on ice for 2 hours. The cells were then transferred to $37^0$C

and further incubated for 1 hour. Subsequently cells were washed, treated with proteinase K and the total

RNA extracted and purified. 1.0 ug of total RNA was reverse-transcribed with random hexamers and the

cDNA used in real-time PCR for comparison of DENV genomic RNA normalized to GAPDH transcripts.

The level in mock-infected cells was taken as 1 and that in DENV infected cells calculated as fold-change

over that. The relative enrichment of DENV RNA compared to mock-infected cells was quantified,

transformed to their logarithmic value to the base of 2 and plotted. The error bars show standard deviation

and the statistical significance was calculated by Student's t-test (ns= not significant).

**Figure 3: DENV replication in differentiating K562 cells interfere with PMA-induced signaling:**

**Panel A:** Schematic showing comparison of signaling activated upon TPO binding to c-Mpl and by PMA.

**Panel B: Immunoblot for signaling protein activated by PMA in K562 cells**. K562 cells either mock-infected or DENV-infected were incubated in PMA-supplemented growth media and the total protein extracted after 3 or 6 days. The protein in each lysate was denatured, resolved in SDS-PAGE and immunoblotted for proteins indicated at right. Immunoblot for b-actin was performed for each batch of lysate used. The relative molecular mass of the marker protein is indicated on the left. **Panel C: Quantification of immunoblot bands in panel B**. the intensity of the immunoblot band for indicated proteins were quantified, normalized to that of b-actin and the ratio calculated before plotting. The error bars represent standard deviation of the ratio derived from at least two independent experiments, performed in multiple replicates.

**Figure 4: Effect of DENV replication on cell surface marker, polyploidy and apoptosis in differentiating K562 cells. Panel A: Suppression of platelet-specific surface marker protein**. K562 infected with DENV at two different MOI, were incubated in PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were immune-stained for the indicated surface marker proteins and the fluorescence quantified by flow cytometry. The mean fluorescence intensity was plotted. The error bars show standard deviation derived from at least two independent experiments in multiple replicates. **Panel B: DENV replication inhibits endomitosis**. DENV infected K562 were incubated in PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were fixed and stained with propidium iodide and the fluorescence intensity measured by flow cytometry. The number of cells with ploidy of >4N were quantified and plotted. The error bars show standard deviation of the mean from two independent experiments performed in multiple replicates. **Panel C: Caspase cleavage assay in differentiated DENV infected cells**. Mock-infected or DENV-infected K562 were incubated in either DMSO or PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were incubated with an substrate that can be cleaved

by intracellular Caspase3/7 to generate a luminescent product. The luminescence was measured in a

luminometer and plotted. The total luminescence at day 3 in DMSO-treated uninfected cells was arbitrarily

taken as 1 and the rest plotted as fold-change over that. The error bars show standard deviation of the mean

from two independent experiments performed in multiple replicates.

**Figure 5: DENV infection suppresses ROS generated during PMA-induced Megakaryopoiesis in**

**K562 cells: Panel A**. Mock-infected or DENV-infected K562 cells were incubated in either DMSO or

PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were stained with $H_2DCFDA$ and

the fluorescence quantified by flow cytometry. The mean fluorescence intensity in mock-infected and

DMSO-treated cells at 3 days was arbitrarily taken as 1 and the rest calculated as fold-change over that.

The error bars show standard deviation of the mean from two independent experiments performed in

multiple replicates. **Panel B**. K562 cells either mock-infected or DENV-infected were immediately

incubated in growth media supplemented with PMA in addition to either 0 or 1 or 3 mM N-Acetyl Cysteine

(NAC) as indicated. At 3 or 6 days post-infection the cells were stained with $H_2DCFDA$ and the

fluorescence quantified by flow-cytometry. The mean fluorescence intensity in mock-infected cells treated

with 100nM PMA and 0 mM NAC was arbitrarily taken as 1 and that in the rest expressed as fold change

with respect to that. The error bars show standard deviation of the mean from two independent experiments

performed in multiple replicates.

**Figure 5: DENV replication interferes with PMA-induced transcriptome changes. Panel A**: The

transcriptome in K562 cells treated with 100nM PMA for 6 days was compared to that after 0 hour by Next-

generation sequencing. The genes differentially expressed after 6 days of treatment, either upregulated by

$\geq 1.5$ fold or downregulated by $\leq 0.5$ fold with a P-value of $\leq 0.05$, when compared to 0 hour were used to

draw a Volcano plot prepared using in-house written Python script. The X-axis represents the logarithmic

value to the base of 2 of the fold-change in gene expression. The Y-axis represents the negative logarithmic

to the base of 10 of the P-value. **Panel B**: The transcriptome in K562 cells, either uninfected or DENV-

infected and treated with 100nM for 6 days was compared to uninfected cells at 0 hour by NGS to find

differentially expressed genes. The level of differential expression of each gene was then compared between

uninfected and DENV-infected cells to draw the list of genes the PMA-induced deregulation of which is

influenced by infection. A volcano plot was prepared from this list using in-house written Python script.

The X-axis represents the logarithmic value to the base of 2 of the fold-change in gene expression. The Y-

axis represents the negative logarithmic to the base of 10 of the P-value. **Panel C**: **Heatmap of genes**

**associated with ROS-pathway**. The genes that are associated with the ROS-pathway in panel B were

selected and enriched using the Metascape software. The PMA-induced fold-change in the transcript level

of these genes after 6 days when compared to uninfected cells treated with PMA for 0 hour (column 1) in

either uninfected cells (column 2) or DENV-infected cells (column 3), was represented by a heatmap

prepared using the Morpheus algorithm (https://software.broadinstitute.org/morpheus). The colors

represent the trend of deregulation corresponding to each gene shown on the right.

**Figure 1**

**Figure 2**

A

B

C

D

**Figure 3**

A

B

C

**Figure 4**

**Figure 5.**

A

B

**Figure 6.**

A

B

C