Title: Megakaryopoiesis in Dengue virus infected K562 cell promotes viral replication which inhibits
endomitosis and accumulation of ROS associated with differentiation

Jaskaran Kaur *[1], Yogita Rawat *[1], Vikas Sood [2], Deepak K. Rathore[1], Shrikant K. Kumar[1], Niraj K. Kumar[1]
and Sankar Bhattacharyya[1]

[1] *Translational Health Science and Technology Institute, NCR Biotech Science Cluster, PO Box# 4,*
Faridabad-Gurgaon expressway, Faridabad, Haryana-121001, India

[2] *Department of Biochemistry, School of Chemical and Life Sciences, Jamia Hamdard (Hamdard*
University) Hamdard Nagar, New Delhi - 110062, India

*Equal contribution

Email for correspondence: sankar@thsti.res.in

Keywords: Dengue virus replication, Megakaryopoiesis, Reactive oxygen species, Endomitosis

Abstract: In the human host blood Monocytes and bone marrow Megakaryocytes are implicated as major sites supporting high replication. The human K562 cell line supports DENV replication and represent Megakaryocyte-Erythrocyte progenitors (MEP), replicating features of *in vivo* Megakaryopoiesis upon stimulation with Phorbol esters. In this article, we report results that indicate the mutual influence of Megakaryopoiesis and DENV replication on each other, through comparison of PMA-induced differentiation of either mock-infected or DENV-infected K562 cells. We present data showing PMA-induced differentiation to drastically increase DENV replication and a concomitant augmented secretion of infectious virus. Although the mechanism is not clear yet, we show that it is not through an increased uptake of virus by differentiated cells. On the other hand, DENV replication in cells undergoing PMA-induced differentiation, interferes with major differentiation markers of Megakaryopoiesis including activation of ERK1/2 MAP Kinase, endomitosis and surface expression of platelet-specific proteins without any drastic effect on cell death. Among signaling intermediaries of the JAK-STAT pathway, we observed infection associated degradation of SOC3 protein similar to earlier observations with STAT2. DENV infection leads to accumulation of Reactive-oxygen species (ROS) in different cells including K562. PMA-induced differentiation of uninfected K562 cells also leads to intracellular ROS accumulation. Interestingly, we observed ROS accumulation to be suppressed by concomitant DENV replication in K562 cells undergoing PMA-induced differentiation. This is the first report of a model system where DENV replication suppresses intracellular ROS accumulation. The implications of these results for Megakaryopoiesis and viral replication would be discussed.

Introduction

Platelets play a unique role in tissue homeostasis and regulation of inflammation. The abundance of these small, anuclear cells produced from specialized cells called Megakaryocytes (MKs) in the bone marrow (BM) is controlled by a steady rate of biogenesis (about 1×10^9/day) and decay, with an average life of 7 days in humans (1). A perturbation of platelet homeostasis is associated with infection by multiple human pathogenic viruses (2). Infection by Dengue virus (DENV) a member of the Flaviviridae, can cause an acute febrile illness with the potential to turn fatal. Beside humans, DENV can infect mosquitoes and these transmit infectious virus along with salivary fluid during a blood meal. Only about a quarter of infected humans exhibit clinical symptoms that include high fever, retro-orbital pain, muscle pain, thrombocytopenia or drop in platelet level etc. that subside after 4-7 days. Subsequently a proportion of symptomatic individuals develop symptoms of severe Dengue characterized by leakage of fluid from the blood vessels, leading to reduction in blood volume in addition to acute thrombocytopenia (3). Multiple mechanisms working in parallel have been suggested to contribute to the thrombocytopenia associated with Dengue infection which include inhibition of platelet biogenesis by infection of Megakaryocytes, augmented platelet decay through either a direct interaction with the virus or upon binding of anti-platelet antibodies (4).

Depending on the cell type the entry of DENV can be mediated by different surface receptors and the complex is endocytosed into endosome following virus binding (5) Acidification of these endosomes by their membrane resident proton pumps induce a fusion between the viral and endosomal membranes leading to release of the viral genome into host cytosol. The genome of DENV is a ~11 kb long single-stranded plus-sense RNA coding for one large polyprotein which is then cleaved to generate the structural (Core or C, Envelop or E and precursor-Membrane or prM) and non-structural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5) of the virus. After few rounds of translation the viral genomic RNA undergoes replication in **r**eplication **c**omplexes (RC) that are associated with the host cell **E**ndoplasmic **r**eticulum (ER). The NS5 protein performs crucial function in replication of the genome by polymerizing both

negative and positive strand genomic RNA through asymmetric replication in favor of the later. Among

these only the plus-sense genomic RNA is capped by a Methyl-transferase (MTase) domain also harbored

in the NS5, followed by encapsidation of the genomic RNA by the Core or C protein. The newly assembled

virus particles contain a RNA-protein complex consisting of a single copy of the genomic RNA with

multiple copies of the C protein, encased in a membrane studded with E and prM protein. These transit

through the protein secretory pathway of the host cell where the prM protein is cleaved by a host protease

called Furin to produce M protein. The virion particles containing M protein along with E in the envelope

are released from the host cell without lysis (6).

DENV can infect multiple cell types in the human body including monocytes in the blood,

megakaryocytes in the BM and hepatocytes in the liver (5, 7). In addition to these DENV can activate

platelets, through a direct interaction with viral receptors on the cell surface (5, 8, 9). Platelets activated by

DENV secrete a number of cytokines and chemokines among which CXCL4 promotes viral replication in

infected Monocytes, upon binding to CXCR3 receptors on these cells (10). Myelosuppression or a reduction

in the mass of BM cells is one of the hallmarks of DENV infection and cells of Megakaryocytic cells have

been shown to be particularly permissive to viral replication (11-14). This suggested MK cells to

significantly contribute to viremia beside monocytes. In addition the survival of megakaryocytes have been

suggested to be negatively affected by infection, thereby leading to reduced platelet biogenesis which

contributes to DENV associated thrombocytopenia (15).

Hematopoietic stem cells (HSC) in the BM differentiate into multiple lineages producing all cell

types in the blood. One of such lineages are composed of bi-potential Megakaryocyte-Erythrocyte

progenitor (MEP) cells that can differentiate into either cell type (16, 17). Differentiation of MEP into a

MK involves profound cellular and molecular changes including cell expansion, endomitosis which

generates a multi-lobed and polyploid nucleus and expression of platelet specific surface markers on the

plasma membrane which will become part of the platelets membrane (18, 19). Platelets bud off from the

MK along with a bit of its cytoplasm, containing granules of different types and a host of specific mRNAs

(20). Differentiation of MEPs into MKs is promoted by a number of cytokines, the most characterized

among which is Thrombopoietin or TPO that binds to the TPO-receptor c-Mpl on the cell surface (21).

TPO binding to c-Mpl activates Janus Kinase 2 (JAK2) receptor tyrosine kinase which in turn transduces

the signal through specific **S**ignal **t**ransducer and **a**ctivator of **t**ranscription or STATs (STA3 and STAT5),

Mitogen-**a**ctivated **p**rotein **k**inases or MAPK (ERK1/2) and the Phosphatidyl Inosine-3 or PI-3 Kinase (22).

STAT3 and STAT5 have been shown to have opposing function in Megakaryopoiesis with the former

promoting it (23, 24). Activation of JAK2 pathway increases expression of pro-inflammatory genes and in

fact activating these genes through exogenous activation of Toll-like receptor 2 or TLR2 has been shown

to promote megakaryocyte development (25). In a similar manner, the activation of ERK1/2 MAPK and

inhibition of p38-MAPK promotes Megakaryopoiesis by playing a crucial role in endomitosis (26).

Transduction of the signal from JAK2 leads to activated transcription by a number of transcription factors

(TFs) increasing the expression of specific genes (27, 28). TFs like GATA-1 and RUNX1 play crucial role

in endomitosis through upregulation of CyclinD1 (29, 30). In addition to these different specific pathways

like ER-stress and Autophagy have been shown to play role in Megakaryopoiesis (31). A controlled form

of apoptosis is known to be crucial for production of platelets from megakaryocytes and ER-stress has been

suggested to play a role in this apoptosis (31-33). In the BM, HSCs occupy a relatively hypoxic niche

compared to more differentiated lineages, particularly the MKs which shed mature platelets into the blood

vasculature. This migration from hypoxic to normoxic environment increases the level of reactive oxygen

species (ROS) in these cells, which has been indicated as a crucial signaling molecules (34). In fact

increased oxygen level and augmented expression of ROS generating enzymes show positive correlation

with megakaryocyte development, corroborating ROS to serve a critical promoter of megakaryopoiesis (35-

37). ROS has been suggested to function at multiple levels which include increasing the level of tyrosine-

phosphorylation on some proteins or by inhibition certain tyrosine phosphatases or by maintenance of ERK

activation or modulation of cyclin levels or increasing expression of crucial TFs or promoting

differentiation associated apoptosis (35, 38-40).

K562 cells represent a bi-potential cell line which can be differentiated to either MK-like cells or

Erythroblast-like cells depending on differential pharmacological stimulation (41). Supplementation of

Phorbol esters like Phorbol-12 myristate-13 (PMA) in growth media of K562 cells drives differentiation

towards megakaryocytes, inducing cellular and molecular changes akin to MKs *in vivo,* through activation

of identical signaling axes (42-50). In addition to this K562 is permissible to DENV replication and is

therefore a good model system to study the effect of DENV infection on Megakaryopoiesis (51).   In this

report we show results suggesting that when DENV infected MEPs differentiate into megakaryocytes the

intracellular viral replication is promoted, by a mechanism that is not clear yet. This enhancement of viral

replication affects key signaling pathways that are known to be crucial for maturation of megakaryocytes,

without any significant effect on their survival. DENV infection is known to induce an accumulation of

ROS in the host cells. K562 cells induced to differentiate into megakaryocytes by treatment with PMA also

accumulate ROS. However surprisingly, infected cells undergoing differentiation accumulate significantly

less ROS when compared to uninfected controls, suggesting an active suppression of ROS by viral

replication. The implications of these observations with respect to Megakaryopoiesis and DENV replication

have been discussed.

Results:

K562 cells recapitulate major events in Megakaryopoiesis upon PMA-treatment

Differentiation of Mk from an MEP follows elaborate cellular changes involving a dramatic increase in cell

size and endomitosis with consequential formation of a multi-lobed, polyploid nucleus as depicted in the

schematic figure (Fig. 1A.). K562 cells mimic a bi-potential Megakaryocyte-Erythrocyte progenitor (MEP)

and depending upon pharmacological treatment can be differentiated into cells of either Mk-lineage

(Phorbol esters) or Erythroblast-lineage (Sodium Butyrate or NaB) (41). Upon supplementation of Phorbol-

12 Myristate-13 acetate (PMA) in culture media, K562 cells stopped proliferation and enlarged in size (data

not shown). Further, the cells underwent endomitosis leading to generation of polyploid cells harboring

multilobed nuclei (Fig. 1B.). When analyzed by flow cytometry, the side scatter of PMA-treated K562 cells

showed a significant increase indicating accumulation of cytoplasmic granules (data not shown). The

platelet plasma membrane is derived from that of the Mk mother cell and during differentiation platelet-

specific surface markers are expressed on the Mk surface. As a corroboration of megakaryopoiesis induced

by PMA-treatment, the surface expression of three platelet specific markers were quantified in K562 cells

and the result showed a dramatic increase in the level of CD41/61, CD61 alone and a significant increase

in number of cells positive for CD42b expression (Fig. 1C.). Analysis of the cell ploidy following PMA-

treatment showed an almost complete disappearance of cells in the S-phase and emergence of polyploid

cells with ploidy of 8N and 16N (Fig. 1D.). As a control for these changes being specific to K562, PMA-

supplementation did not stall the growth or induce expression of platelet-specific surface markers in cells

of the hepatoma cell line Huh7 (data not shown). Since treatment with NaB induces differentiation towards

an Erythroblast-lineage, the differential expression of specific genes were compared between K562 cells

treated with either PMA or NaB. The result showed a distinct pattern of gene expression induced by these

pharmacological treatments differentiating the two lineages, especially with respect to transcripts of the

gene GATA2, EBP42, GYPA and CD61 (Fig. 1E.).

Differentiation of K562-MKs promote replication of Dengue virus

Differentiation into megakaryocytes is associated with arrest in cell cycle and other molecular changes. In order to analyze how these changes might affect the ongoing intracellular replication process, DENV replication in infected K562 cells that have been treated with PMA soon after infection, was compared to simultaneously infected cells treated with vehicle (DMSO). For this purpose, the level of DENV envelope protein accumulated was compared between these conditions at 3 or 6 days post-infection by immunofluorescence followed by analysis in a flow cytometer. The result showed that although the viral protein accumulated at 3 days p.i was comparable between PMA-treated and DMSO-treated cells, a dramatic difference between them was observed at day 6 (Fig. 2A.). This difference was observed using different MOI of infection, although due to saturation of immunostaining the difference was less dramatic at higher MOI (data not shown). In order to check if higher accumulation of viral protein correlates positively with viral replication, the level of intracellular viral RNA and infectious virus in culture supernatant was compared between infected K562 cells treated with either PMA or DMSO for 3 or 6 days. The results showed a similar higher accumulation of intracellular viral genomic RNA and extracellular infectious virus, between these two conditions at 6 days post-infection (Fig. 2B and 2C). During virus maturation the prM protein on assembled virion particles undergoes a cleavage by the host protease Furin, a step necessary for conferring infectivity to the secreted virus (52). Therefore the comparatively higher infectious titer in culture supernatant of PMA-treated cells can result from either higher secretion of virus (more virion particles without any change in their infectivity) or more efficient Furin cleavage brought about by the cellular and biochemical changes accompanying differentiation (same number of virion particles as secreted by DMSO-treated cells but with higher infectivity). To dissect these possibilities, the viral genomic RNA from culture supernatant of cells kept with either PMA or DMSO was extracted and purified and their relative level compared by real-time PCR using DENV specific oligonucleotide primers. The comparison showed comparatively more viral genomic RNA in the supernatant of PMA-treated cells,

suggesting that the differentiating cells secrete more virus that are produced by higher replication efficiency

in these cells (data not shown). However this indicated towards possibilities wherein the entry within

differentiating cells might be facilitated e.g. through overexpression of viral receptor. In order to address

this, uninfected K562 cells treated with either DMSO or PMA for 3 days were incubated with DENV

inoculum and the viral entry compared by comparison of the internalized genomic RNA immediately after

entry. The result showed a comparable entry into cells irrespective of no differentiation or differentiation

for 3 days, implying that the promotion of replication occurs post-entry (Fig. 2D). As a negative control

DENV-infected Huh7 cells did not exhibit any differential replication depending on treatment with PMA

and DMSO (data not shown). Also, treatment of infected K562 cells with Na-butyrate which would initiate

erythropoiesis, did not exhibit a similar promotion of the viral replication (data not shown).

Cellular and biochemical changes imposed by DENV infection in differentiating K562 cells

Differentiation of Mk from HSCs is influenced by multiple cytokines, the most characterized among which

is Thrombopoietin (TPO) which is secreted from liver cells and binds to the surface receptor c-Mpl.

Interaction of TPO with c-MPL activates Janus kinase 2 (JAK2) tyrosine kinase initiating a cascade of

phosphorylation events (Fig. 3A). Although an important cytokine, TPO is not indispensable for

megakaryopoiesis since a knock-out of the gene encoding TPO (TPO$^{-/-}$) is not lethal in mice, albeit with the

platelet level being at 10% of that in wild-type animals. Phorbol esters like PMA are known to activate

kinases like Protein kinase-C (PKC) that in its turn transduces the signal through nodes many of which are

shared with TPO signaling. In response to viral infection mammalian cells secrete cytokines like Interferons

that bind to cell surface receptors either in an autocrine or paracrine manner and induce an 'anti-viral' state

within the cell. JAK2 is also central to signaling via the interferon receptor consequent to binding of type-

I IFNs to the receptor (IFNR). This implied that megakaryopoiesis even when it is PMA-induced would

potentially create an antiviral state within the cell and in that context it would be interesting to observe how

DENV is still able to replicate to higher extent compared to undifferentiated cells. Since both of these

pathways would also be potentially perturbed by a virus infection, we investigated the status of different

members by immunoblotting, in cells which were stimulated to undergo PMA-dependent differentiation

immediately after infection.

Phosphorylation of the MAPK ERK1/2 plays a crucial role in MK development and expectedly we observed

an increase in the ratio of p-ERK/ERK both subsequent time points when compared to day 0 (Fig. 3B,

compare lanes 2 and 5 with lane 1). Interestingly, we observed that DENV infection to reverse the PMA-

induced ERK1/2 phosphorylation at 3 days post-infection (Fig. 3B, compare lanes 3 and 4 with lane 2).

However, by day 6 the phosphorylation of ERK1/2 recovered significantly in infected cells when compared

to uninfected controls (Fig. 3B, compare lanes 6, 7 with lanes 3, 4). Nonetheless, the ratio of p-ERK/ERK

at 6 days post-infection was still lower in virus infected cells when compared to uninfected cells treated

with PMA for a similar duration (Fig. 3B, compare lanes 6 and 7 with lane 5). In a manner opposite to that

of ERK1/2, a dephosphorylation of p38 MAPK has been similarly reported to be crucial for MK

development. We did not observe any modulation in the phosphorylation status of p38 in DENV-infected

cells treated with PMA for 3 or 6 days (data not shown).

In addition to MAPKs we compared the status of different signaling protein in the JAK-STAT pathway in

uninfected or DENV-infected K562 cells treated with PMA. While some effect of virus infection was

observed with respect to a few of these signaling mediators, most of them were unaffected. A few more

JAK-STAT pathway intermediaries were probed by immunoblotting but remained undetectable under

presently used conditions (SOCS1, PIAS1, PIAS3 and PIAS4). When compared to 0 hour the ratio of p-

STAT1/STAT1 showed a significant decrease upon PMA treatment at both time points (Fig. 3B, compare

lanes 2 and 5 with lane 1). However, although DENV replication in PMA-treated cells did not alter this

ratio at 3 days p.i., by 6 days p.i. the ratio showed a dramatic increase in infected cells when compared to

uninfected controls (Fig. 3B, compare lanes 3, 4 with 2 and lanes 6, 7 with lane 5). Further, at 6 days p.i.

the total STAT1 protein level was also reduced to undetectable level in infected cells (Fig. 3B, compare

lanes 6 and 7 with lane 5).

STAT3 has been suggested to promote Megakaryopoiesis and in support of this we observed an increase in

ratio of p-STAT3/STAT3 at 3 days after PMA treatment, which was maintained till 6 days (Fig. 3B,

compare lanes 2 and 5 with lane 1). However in DENV infected cells treated with PMA, the ratio of p-

STAT3/STAT3 at both 3 and 6 days post-treatment was significantly lower when compared to uninfected

controls (Fig. 3B, compare lane 3 with lane 2 and lanes 6,7 with lane 5). The lane 4 in this immunoblot is

missing due to the technical reason of sample insufficiency.

Although STAT2 protein level is not known to be modulated by PMA-treatment, it is known to be degraded

in host cells following DENV infection, in a NS5 dependent manner (Ref). We did not observe any

alteration in STAT2 levels with PMA-treatment of uninfected cells at either time point (Fig. 3B, compare

lanes 2 and 5 with lane 1). As expected however, the protein was reduced to undetectable level in DENV

infected cells. (Fig. 3B, compare lanes 3,4 with lane 2 and 6,7 with lane 5).

A downregulation of STAT5 protein has been shown to be essential for promoting development of MKs.

In concurrence with that, we observed a reduction in the level of this protein upon PMA-treatment starting

from day 0 to almost undetectable level by day 6 (Fig. 3B, compare lanes 2 and 5 with lane 1). No additional

modulatory effect was however observed by DENV infection of PMA-treated cells (Fig. 3B, compare lanes

3,4 with lane 2 and 6,7 with lane 5).

An activation of the JAK-STAT signaling upregulates inflammation, which eventually leads to expression

of a family of genes that suppresses this pathways and are called as Suppressor of cytokine signaling or

SOCS1, SOCS2 and SOCS3. In accordance with a crucial role for activated JAK-STAT pathway in Mk

development, expression of the SOCS3 protein has been shown to have a negative effect on

megakaryopoiesis. However, the SOCS3 level following PMA-treatment was observed to be unchanged at

either time points (Fig. 3B, compare lanes 2 and 5 with lane 1). Surprisingly, in DENV-infected cells SOCSs

was observed to be reduced to undetectable level by day 6 (Fig. 3B, compare lanes 6 and 7 with lane 5).

These results implied that DENV infection might be selectively affecting the innate antiviral and

inflammatory pathways in differentiating K562 cell through degradation of STAT2 and SOCS3 level and

interference with STAT3 phosphorylation.

252 DENV infection inhibits polyploidy and accumulation of ROS in differentiating cells without

253 affecting apoptosis

The biochemical changes associated with Megakaryopoiesis affect the cellular modifications. Therefore we

quantified the PMA-induced surface marker expression and polyploidy was compared between uninfected

or DENV-infected cells treated with PMA. As above, cells that were either mock-infected or infected with

DENV were treated with PMA-supplemented growth media soon after infection and analyzed at 3 or 6 days

post-treatment. The results showed a significant reduction in the surface expression of CD41/61, CD42b

and CD61 in DENV infected cells (Fig. 4A). Phosphorylation of the ERK1/2 MAPK has been shown to be

an critical modulator of polyploidy during Megakaryopoiesis. Since we observed a suppression of ERK1/2

phosphorylation at the early time point post-infection, a comparison of the percentage of polyploid cells at

both time points was quantified by PI staining. As expected the result showed a significant reduction in the

percentage of polyploid cells generated following DENV infection (Fig. 4B). Generation of platelet from

megakaryocytes has been demonstrated to be dependent on induction of apoptosis in the mother cell and

proceeds via a unique method of intrinsic apoptosis. In accordance with PMA-induced Megakaryopoiesis

in K562 cells we observed a significant increase in Annexin-V positive cells upon PMA-treatment (data

not shown). However, a comparison of mock-infected and DENV-infected cells treated with PMA for either

3 or 6 days did not show any difference in the proportion of Annexin-V positive cells (data not shown).

However surprisingly, a comparison of Caspase3/7 activation showed significant increase in cleavage

activity in virus infected cells when compared to uninfected controls at 6 days p.i. (Fig. 4C).

Intracellular ROS is known to be generated during megakaryopoiesis and an accumulation is

essential for differentiation (47). DENV replication has been shown to induce generation of reactive oxygen

species in infected cell (53). An estimation of the ROS accumulated in K562 cells as a result of DENV

infection showed slight accumulation at 6 days post-infection (Fig. 5A, compare columns 2 to 1 and 6 to

5). In a similar manner, when compared to levels at day 0, uninfected cells treated with PMA showed a

significant accumulation of ROS at both 3 and 6 days post-treatment (Fig. 5A, compare columns 3 to 1 and

7 to 5). We expected the ROS level in DENV infected cells that have been treated with PMA to be higher

than the uninfected PMA-treated controls. However, surprisingly DENV infected cells treated with PMA

did not exhibit additional ROS accumulation at day 3 post-treatment (Fig. 5A, compare columns 3 to 4).

Further, surprisingly at day 6 after PMA-treatment the ROS accumulation in DENV infected cells were

remarkably lower compared to uninfected PMA-treated cells (Fig. 5A, compare columns 7 to 8). This

suggested DENV replication in differentiating cells to impede accumulation of ROS. In corroboration we

observed DENV infection to have an effect similar to that of N-acetyl cysteine a biochemical known to

quench ROS levels (Fig. 5B). In conclusion, DENV infection interferes with surface marker expression in

addition to significantly affecting accumulation of ROS and endomitosis which are associated with

differentiation into megakaryocytes.

PMA-induced transcriptome changes are reversed by DENV infection in differentiating cells

Megakaryopoiesis is associated with profound alterations in the transcriptome of differentiating cells.

Similarly, innate antiviral response launched upon detection of a viral infection is also associated with

changes in the pattern of expression of many genes. Therefore, in order to study the interaction between

these two stimuli at the transcriptional level the transcriptome of mock-infected and DENV-infected cells

was compared after 6 days of PMA-induced differentiation. For this purpose, K562 cells either mock-

infected or DENV-infected were induced to differentiate using 50 nM PMA and the total RNA isolated for

analysis of transcriptome by next-generation sequencing. The transcriptome of both uninfected and DENV-

infected cells was compared to uninfected cells at 0 hour with respect to PMA-treatment. As shown in Fig.

6A, the transcriptome in uninfected cells underwent a profound change in more than 5000 genes showing

different degree of deregulation. Interestingly, the comparison between uninfected and DENV-infected

cells after 6 days of differentiation showed reversal in the direction of PMA-induced deregulation for a

number of genes (Fig. 6B.). Since our earlier result showed that DENV infection suppressed ROS

accumulation in differentiating cells, ROS-associated genes the PMA-induced deregulation trend of which

was reversed by infection were analyzed further. A heatmap of these genes showed that although alteration

in transcript level was moderate for most of them, a few genes were dramatically affected (Fig. 6C). This

included GCH1, FOXO3, GLRX5, ETHE1, CTNS, EPAS1, GFOD1 and SNCA (Fig. 6C). Among these

transcripts corresponding to FOXO3, CTNS and EPAS1 were higher in infected cells compared to

uninfected ones, whereas the others showed a relatively lower level in infected cells. FOXO3 and EPAS1

are transcription factors among which the former is known to be pro-apoptotic (54). Among the

downregulated genes the protein corresponding to GCH1, ETHE1 and SNCA are enriched platelets (55-

57). SNCA is also known to function in an anti-apoptotic manner although the specific role of this gene in

apoptosis associated with megakaryocyte differentiation is not clear yet (58).

Discussions: DENV replicates in multiple cell types and causes a range of pathological symptoms.
Thrombocytopenia, a prominent DENV associated pathological symptom is a consequence of both
impaired biogenesis and altered stability of platelets. Megakaryocytes or platelets mother cells in the bone
marrow are known to be permissive for DENV replication, which affects platelet biogenesis. In this article,
we have explored the cellular and molecular effect of DENV infection on Megakaryocyte development
using the phorbol ester PMA induced-differentiation model in human K562 cells. Our results show that
differentiation of infected K562 cells promote the replication of intracellular virus by an unknown
mechanism and viral replication interferes with cell signaling pathways crucial for Megakaryopoiesis,
particularly involving the MAPK ERK1/2 and different factors in the JAK-STAT pathway. However high
level of viral replication does not affect the survival of the cells. PMA-induced differentiation of K562
cells, as with *in vivo* Megakaryopoiesis, is associated with accumulation of intracellular ROS and our results
show that DENV replication suppresses the accumulation of this signaling molecule by an unknown
mechanism. Replication by multiple viruses is known to increase ROS in the host cell we report an unique
observation about DENV replication suppressing ROS accumulation under specific conditions in the host
cell.

The absolute dependence of virus life cycle on host biochemical pathways implies a profound influence of
host cell metabolic status on viral replication. Depending on the virus being studies, PMA a known activator
of protein kinase C, is known to have either positive or negative effect on virus replication (59-62). On the
other hand, viral infection of human monocytes can induce differentiation of these cells into Dendritic cells
although the implication of this is not clear yet (63). Cell cycle arrest in PMA-treated K562 cells happens
through upregulation of p21 and p27 in a protein kinase C (PKC)-dependent but p53-independent manner
(64). Inducing an arrest in the host cell cycle is quite common among human viruses with beneficial effect
on their replication (65). The entry of DENV is known to be influenced by the cell-cycle stage of the host
cell, albeit in a cell-type dependent manner (66, 67). Here we observed increased intracellular replication
in K562 cells that have stopped dividing in response to PMA, although the halt in cell division does not

lower the metabolic activity as is obvious from increase in cell size and other cellular changes. In a similar

manner Parvovirus B19 has been observed to infect cells of erythroblast origin with replication of the viral

genome being promoted by differentiation of these cells into erythroblasts (68). Although it is not clear yet

if the PMA-induced cell cycle stall is responsible for increasing intracellular virus replication, single-cell

transcriptomics study by Zanini and coworkers suggested the cell cycle status of the host cell to have no

effect on DENV genome replication (69). In addition to DENV, Mk cells are known to support replication

of other viruses like the Human Immunodeficiency Virus and Hantavirus, with the former inducing

apoptosis in addition to reducing the surface expression of c-Mpl (70-73). Interestingly, PMA-induced

differentiation of Hantavirus infected megakaryocytes augmented virus replication, as observed in this

study (72).

PMA induces rapid and profound changes in host gene expression and therefore it is possible that

specific promoters of viral replication are overexpressed upon PMA treatment as observed for Hepatitis C

virus which is benefitted by the liver cell enriched microRNA-122 (74). PMA-induced differentiation

correlates with increase in granularity of K562 cells, probably pertaining to platelets granules that are

synthesized in the megakaryocyte mother cell and transferred into platelets (75). Therefore the rate of

protein synthesis in these cells can be expected to be enhanced for synthesis of the proteins that will form

part of these granules. In face the protein content of K562 cells is significantly increased during PMA

induced differentiation (76).It is possible that translation of viral proteins benefits from this overall increase

in rate of protein synthesis. The DENV genomic RNA of DENV has a 5' end type-I cap identical to host

mRNAs that helps in recruiting ribosomes in addition to evading innate antiviral arsenal (77). A role for

enhanced host protein synthesis benefiting viral translation is corroborated from our observation that the

entry of DENV is not higher in differentiated cells. Another possibility is that differentiation leads to

suppression of specific anti-viral genes, thereby benefiting the replication.

Although TPO is a cytokine crucial for platelet biogenesis and mice knock-out for either the

cytokine or its receptor show drastically reduced levels of both mature MKs as well as circulating platelets,

residual level of both MKs and platelets in these mice suggest other factors to also contribute in MK

development and platelet formation (78, 79). Receptor interaction of TPO activates JAK2 kinase leading to

activation of the MAPK ERK1/2, an important regulator of cell cycle and proliferation that plays a crucial

role in Mk maturation (80, 81) Previous studies of Mk development using K562 cells showed PMA to

directly activate protein kinase C (PKC) which in turn induces ERK1/2 phosphorylation, an important

signaling event even for *in vivo* Megakaryopoiesis (26, 50, 82, 83). A concurrent inhibition of the MAPK

p38 on the other hand supports MK development, suggesting this kinase to oppose or retard ERK1/2

mediated changes (22). Dengue infection can activate the p38 MAPK which is responsible for enhancement

of pro-inflammatory cytokines and apoptosis (84, 85). We observed a reduction in the level of PMA-

induced ERK1/2 phosphorylation in DENV-infected cells, probably causing inhibition of endomitosis and

platelet-specific surface marker protein expression, although a concomitant increase in p38 phosphorylation

was not observed. Previous reports have conclusively shown DENV NS5 protein mediated degradation of

the host STAT2 infection as a major mechanism for suppression of the host innate-antiviral pathway (86).

We observed additional regulation of a few other members of the JAK-STAT pathway, namely STAT1,

STAT3 and STAT5, although it is yet clear if this is restricted to PMA-treated K562 cells or is a general

effect occurring in other cells too. STAT1 regulated by the transcription factor (TF) GATA-1 plays a

specific role in endomitosis by controlling expression of cell cycle genes like CCND1, CCND2 and

CCNE2TFs through the TF RUNX1 (87). PMA-induced Mk development in K562 has not been reported

earlier to mimic the *in vivo* differentiation with respect to regulation of STAT3 and STAT5. In a manner

corroborative of *in vivo* development, we observed a gradual reduction of STAT5 level in parallel with

progression of differentiation (24). STAT3 is phosphorylated initially on Tyr-705 which triggers

dimerization (88). In addition to this STAT3 can have other post-translational modifications e.g.

phosphorylation of S-727 or acetylation of K-685, although induction of these modifications by PMA in

K562 cells is not clear yet (89, 90). In our study we observed DENV infection to reverse the PMA-induced

STAT3 phosphorylation on Tyr-705, possibly through inhibition of the respective kinase or activation

specific phosphatases. In addition to phosphorylation STAT3 function can be also suppressed by Ubiquitin

mediated degradation or interaction with proteins that are activated downstream of the JAK-STAT pathway

in a negative feedback loop e.g. protein encoded by PIAS and SOCS genes (91-93). In fact SOCS3 is known

as a specific inhibitor of STAT3 signaling pathway, for which we expected an upregulation in infected

cells (90). Unexpectedly however, we observed a reduction in SOCS3 protein level by DENV infection for

which either mechanism or implication with respect to viral life cycle is still unclear. One of the potential

role for STAT3 in Megakaryopoiesis is driving accumulation of intracellular ROS levels through

upregulation of NOX2 expression (94). Therefore, suppression of STAT3 phosphorylation might be

directly affecting accumulation of ROS as observed here. Further characterization of this observation would

probably shed more light on the implication of this regulation.

Reactive oxygen species or ROS, in the form of Hydroxyl radical ($^{\cdot}OH$) or Singlet oxygen ($^{1}O_2$) or

Superoxide anion radical ($O_2^{\cdot}\,-$), are generated either in response to stimuli including metabolic

inflammation, exposure to pathogen or as secondary messengers in signal transduction pathway. The most

well characterized sources of intracellular ROS are two enzymes associated with the mitochondria namely

Nicotinamide **A**denine **D**inucleotide-**Q**uinone (NADH-Q) reductase (Complex I), Q-cytochrome c

oxidoreductase (Complex III) and the plasma membrane associated homologs of NADPH-oxidase or NOX

(95-97). Intracellular ROS plays crucial role in regulation of protein tyrosine kinases and phosphatases

through post-translational modification (98, 99). It can also activate multiple cellular pathways including

MAPK, NFκB, Cell cycle genes (100). Intracellular ROS can either promote apoptosis by inducing damage

to the mitochondrial membrane or inhibit it through oxidation of catalytic site Cysteines on executioner

proteases like Caspase-3 (101). In our study suppression of ROS level coincided with activation of Caspase-

3 cleavage activity, although the mechanism would need further investigation to draw a direct correlation

between them.

In the BM, pluripotent hematopoietic stem cells (HSCs) reside in a relatively hypoxic area while

during differentiation megakaryocytes relocate to more oxygen-rich region near blood vessels from where

platelets can be shed into the vascular circulation (102). ROS serves a critical role in megakaryocyte

formation since a low level of these signaling molecules in differentiating HSCs disfavor generation of

(**M**egakaryocyte-**E**rythrocyte **p**rogenitors) MEP and in favor of (**G**ranulocyte-**M**onocyte **p**rogenitors) GMP

(103). Intracellular ROS levels are regulated through the interplay of generators (like NOXs) and mitigatory

pathways (like the KEAP1-NRF2 pathway) (37, 104). PMA has been shown to induce accumulation of

ROS in K562 cells and ROS has been shown to be responsible for polyploidy and a reduction of ROS by

NAC directly impacts polyploidy (43, 47). Accumulation of ROS activates the KEAP1-NRF2 pathway and

activation of many genes that can suppress the cellular ROS levels. ROS is produced by cellular responses

like Unfolded-protein response (UPR) or innate-antiviral pathway (105, 106). We observed a relatively

higher level of intracellular ROS with respect to uninfected cells in K562 cells that are not undergoing

differentiation. Since PMA-induced differentiation of K562 cells also causes ROS to accumulate in cells,

we expected an additive effect of DENV replication on ROS accumulation in these cells. However,

surprisingly DENV replication quenched intracellular ROS levels through an unknown mechanism.

Although evidence from previous reports would strongly indicate this suppression of ROS to be major

contributor to inhibition in platelet biogenesis by infection of Mk mother cells by DENV, it is still not clear

if the viral replication in these cells is benefited from this or not. Future studies would be directed to address

the exact mechanism through which DENV infection suppresses ROS accumulation in these cells and the

physiological relevance of this suppression for the viral life cycle.

Acknowledgement: The authors would like to thank Dr. Manjula Kalia, Regional Centre for
Biotechnology, NCR Biotech Science Cluster for careful reading of the manuscript. The Translational
Health Science and Technology Institute is acknowledged for providing all support for equipment and other
infrastructure. The Core Technology Research Initiative (CoTeRI) in the National Institute for Biomedical
Genomics, India is acknowledged for providing Next-generation sequencing facility. JK is supported by
fellowship from the University grants commission. YR is supported from fellowship in the extramural
research grant (BT/PR22985/MED/29/1168/2016) to SB from the Department of Biotechnology, Govt of
India. This work was supported by extramural research grant (EMR/2016/005796) to SB from the Science
and Engineering Research Board, Department of Science and Technology, Govt. of India.

Materials and methods:

Cell culture, Drugs and Virus: K562 cells were procured from the American Type Culture Collection

(ATCC) and cultured at 37^0C, 5% CO_2 in Iscove's modified Dulbecco's media (IMDM) supplemented with

Penicillin (100 U/ml), Streptomycin (0.1 mg/ml) and 10%-Fetal Bovine Serum (FBS). Vero and C6/36 cells

were procured from the cell line repository of the National Centre for Cell Sciences (NCCS), India and

cultured respectively in Minimum Essential Medium (MEM) and L15 cell culture medium supplemented

with 10% FBS. C6/36 cells were maintained at conditions of 28^0C and atmospheric CO_2.

Dengue virus serotype 2 (strain NGC) was amplified in C6/36 and the culture supernatant used as inoculum.

The infectious titer of the virus was determined by Focus-forming unit (FFU) assay in Vero cells. For all

infections, the inoculum was diluted in the respective culture media supplemented with 2% FBS, and the

cells were incubated with the inoculum for 2 hours at the respective culture conditions of temperature and

CO_2 concentration, with intermittent rocking.

Phorbol-12 Myristate-13 acetate (PMA, Sigma Aldrich) and N-Acetyl Cysteine were diluted in

recommended vehicles and stored as single use aliquots at -20^0C . Fresh dilutions of N-Acetyl-L-Cysteine

(Sigma) were used for every experiment.

Focus-forming unit (FFU) assay: Vero cell monolayers in 24-well plate were infected with 10-fold serial

dilutions of virus inoculum. The inoculum was incubated with cells for 2 hours at 37^0C and 5% CO_2 with

intermittent rocking. The inoculum was then discarded and complete MEM added to each followed by

incubation for 48 hours at 37^0C and 5% CO_2. Subsequently the cells were washed with phosphate-buffered

saline (PBS), fixed with 2% para-formaldehyde (PFA) and permeabilized with PBS supplemented with 0.1

% Triton-X-100 and 1% bovine serum albumin (BSA). The permeabilized cells were washed with PBS and

sequentially incubated anti-DENV primary antibody (dilution 1:400 of Mab8705, Millipore) and anti-

Mouse Alexa-488 conjugated secondary antibody (dilution 1:500; ThermoFisher Scientific) both diluted in

PBS supplemented with 1% BSA. After incubation with primary antibody, the cells were washed twice

with PBS before addition of secondary antibody dilutions. The fluorescent foci were visualized and

manually counted using a fluorescence microscope and the titer calculated as FFU/ml.

Flow Cytometry Analysis: All flow cytometry analysis were performed in a BD FACS Canto II flow

cytometer (BD Biosciences) under standard conditions and all raw data analyzed using Flow-Jo software.

For immune staining of intracellular DENV antigen, K562 cells were washed with ice-cold PBS before

fixation with PBS supplemented with 2% PFA and permeabilization with the same buffer containing 0.1%

Triton-X-100. The permeabilized cells were sequentially stained with dilution of 4G2 (purified IgG2a mAb

produced in the lab from Hybridoma-HB112) antibody and anti-mouse Alexa-488 conjugated secondary

antibody. For immune staining of surface markers, cells were washed with PBS and incubated with dilutions

of fluor-conjugated primary antibodies, specific to either human CD61-Phycoerythrin (PE) (BioLegend) or

CD41/61-(Allophycocyanin) APC (BioLegend) or human CD42b-(BD Horizon brilliant violet 421) Bv421

(BioLegend). Antibodies were diluted in staining buffer (PBS with 1%BSA and 0.02% Sodium Azide) and

cells were incubated for 1h at 4°C . Subsequently the cells were washed with PBS and analyzed by flow

cytometry.

Propidium iodide staining for polyploidy analysis: Cells were harvested by centrifugation, washed

twice with ice-cold phosphate-buffered saline (PBS) and fixed with 70% ethanol at 4°C for 30 min. After

a PBS wash cells were treated with 100µl PBS supplemented with RNase-A (200µg/ml) and incubated at

37°C for 30 min. Cells were then washed using PBS and the genomic DNA stained with Propidium iodide

(50µg/ml).

Annexin-V and PI staining: Apoptotic detection assays were carried out by surface labeling with the Ca2+

dependent phosphatidylserine-binding protein Annexin-V. Cells were harvested by centrifugation, washed

twice with cold PBS and then re-suspended in 1x binding buffer(Apoptosis detection Kit I # 556578, BD

Pharmingen) at 1x 10^6 cells/ml. A 100µl aliquot was stained by addition of 5µl FITC Annexin V and 5µl

PI. The reagents were mixed by gentle vortex and incubated for 15 min at RT in the dark. 400µl of 1x

binding buffer was added to each tube and the cells analyzed by flow cytometry within 1 hr.

RNA extraction and Quantitative RT-PCR: Total RNA from 1×10^6 K562 cells were extracted using

Trizol (Takara) and purified with Qiagen RNAeasy mini kit (Qiagen), as per manufacturer's instructions.

Viral RNA from cell-free culture supernatant was isolated using QIAmp viral RNA mini kit (Qiagen) as

per manufacturer's instructions. 1.0 ug of total RNA was reverse-transcribed with ImProm-II reverse-

transcriptase (Promega, USA) and random hexamers (Sigma) as per manufacturer's protocol, and the cDNA

diluted with nuclease-free water before use for real-time PCR. Real-time PCR was performed with 2x

SYBR mix (Takara) in a QuantStudio-6 Flex Real-Time PCR System (Applied Biosystems) using the

default run program.

Confocal microscopy: K562 cells were washed with PBS, fixed and permeabilized as described earlier for

immunostaining purposes. The cells were stained with 2.5µl of Alexa Fluor-568 Phalloidin (Invitrogen,

Thermo fisher scientific) at RT for 20 min to stain for F-Actin followed by PBS wash. Cells were then

mounted onto glass slides using ProLong Gold Antifade Mountant supplemented with DAPI (Invitrogen,

Thermo fisher scientific). The fluorescence was observed and imaged in a FLUOVIEW FV3000 confocal

microscope (Olympus).

Intracellular Caspase-3/7 activity assay: The intracellular caspase-3/7 activity was measured using

Caspase-Glo 3/7 Assay (Promega). For this, 1×10^4 cells in 50 µl were mixed with an equal volume of

Caspase-Glo3/7 reagent in wells of 96-well luminometer plate. The plate was incubated at room

temperature for 3 hours and the luminescence measured using a Orion II microplate luminometer

(Berthold). The luminescence from culture media without cells was used as negative control.

Measurement of intracellular Reactive Oxygen species (ROS): The intracellular ROS accumulated was

measure using CM-H2DCFDA probe (Life Technologies) as per manufacturer's instructions. Briefly, K562

cells were washed with PBS and then incubated in PBS supplemented with CM-H2DCFDA as per

manufacturer's instructions. The cells were then incubated at 37 °C , 5% CO_2 for 30 minutes and washed

twice with PBS. The fluorescence intensity was quantified in a BD FACS Canto II flow cytometer (BD

Biosciences) under standard conditions and the raw data analyzed using Flow-Jo software.

Electrophoresis and Immunoblotting: Cells were harvested by centrifugation at 4^0C and washed with ice

-cold PBS. Cells were suspended in lysis solution [250 mM NaCl, 20 mM Tris-HCl (pH 7.4), , 1% Triton

X-100, 1 mM EDTA, supplemented with 0.1mM PMSF and protease inhibitor cocktail] and incubated on

ice for 15 minutes. The lysate was clarified by centrifugation (13 000×g for 10 min at 4^0C) and the

supernatant collected. The total protein in the lysate was estimated using Bradford's method. 40µg of

whole-cell lysate was denatured, resolved in a SDS-10% gel and transferred to nitrocellulose blotting

membrane. The membrane was blocked with either 5% skimmed milk or 5% bovine-serum albumin (BSA)

and incubated with dilution of respective antibodies as per manufacturer's instructions (Cell Signaling

technologies). The membrane was washed, incubated with dilution of HRP-conjugated secondary antibody

and the bands visualized by ECL chemiluminescence.

Next-generation sequencing: The total RNA from K562 cells either uninfected or infected with DENV at

an MOI of 10 was extracted and purified. Total RNA from similar number of uninfected cells, harvested

for RNA extraction immediately after addition of PMA was used as 0 hour control. The total RNA was

subjected to ribosomal RNA depletion before a sequencing library was prepared for 2x100 bp end-paired

sequencing in Illumina HiSeq2500 platform, generating 50 million reads per sample.

References:

1. Patel SR, Hartwig JH, Italiano JE, Jr. 2005. The biogenesis of platelets from megakaryocyte proplatelets. J Clin Invest 115:3348-54.
2. Hottz ED, Bozza FA, Bozza PT. 2018. Platelets in Immune Response to Virus and Immunopathology of Viral Infections. Front Med (Lausanne) 5:121.
3. Martina BE, Koraka P, Osterhaus AD. 2009. Dengue virus pathogenesis: an integrated view. Clin Microbiol Rev 22:564-81.
4. de Azeredo EL, Monteiro RQ, de-Oliveira Pinto LM. 2015. Thrombocytopenia in Dengue: Interrelationship between Virus and the Imbalance between Coagulation and Fibrinolysis and Inflammatory Mediators. Mediators Inflamm 2015:313842.
5. Fang S, Wu Y, Wu N, Zhang J, An J. 2013. Recent advances in DENV receptors. ScientificWorldJournal 2013:684690.
6. Rodenhuis-Zybert IA, Wilschut J, Smit JM. 2010. Dengue virus life cycle: viral and host factors modulating infectivity. Cell Mol Life Sci 67:2773-86.
7. Acosta EG, Kumar A, Bartenschlager R. 2014. Revisiting dengue virus-host cell interaction: new insights into molecular and cellular virology. Adv Virus Res 88:1-109.
8. Simon AY, Sutherland MR, Pryzdial EL. 2015. Dengue virus binding and replication by platelets. Blood 126:378-85.
9. Ojha A, Nandi D, Batra H, Singhal R, Annarapu GK, Bhattacharyya S, Seth T, Dar L, Medigeshi GR, Vrati S, Vikram NK, Guchhait P. 2017. Platelet activation determines the severity of thrombocytopenia in dengue infection. Sci Rep 7:41697.
10. Ojha A, Bhasym A, Mukherjee S, Annarapu GK, Bhakuni T, Akbar I, Seth T, Vikram NK, Vrati S, Basu A, Bhattacharyya S, Guchhait P. 2019. Platelet factor 4 promotes rapid replication and propagation of Dengue and Japanese encephalitis viruses. EBioMedicine 39:332-347.
11. Noisakran S, Onlamoon N, Hsiao HM, Clark KB, Villinger F, Ansari AA, Perng GC. 2012. Infection of bone marrow cells by dengue virus in vivo. Exp Hematol 40:250-259 e4.
12. Vogt MB, Lahon A, Arya RP, Spencer Clinton JL, Rico-Hesse R. 2019. Dengue viruses infect human megakaryocytes, with probable clinical consequences. PLoS Negl Trop Dis 13:e0007837.
13. Clark KB, Noisakran S, Onlamoon N, Hsiao HM, Roback J, Villinger F, Ansari AA, Perng GC. 2012. Multiploid CD61+ cells are the pre-dominant cell lineage infected during acute dengue virus infection in bone marrow. PLoS One 7:e52902.
14. Sridharan A, Chen Q, Tang KF, Ooi EE, Hibberd ML, Chen J. 2013. Inhibition of megakaryocyte development in the bone marrow underlies dengue virus-induced thrombocytopenia in humanized mice. J Virol 87:11648-58.
15. Basu A, Jain P, Gangodkar SV, Shetty S, Ghosh K. 2008. Dengue 2 virus inhibits in vitro megakaryocytic colony formation and induces apoptosis in thrombopoietin-inducible megakaryocytic differentiation from cord blood CD34+ cells. FEMS Immunol Med Microbiol 53:46-51.
16. Akashi K, Traver D, Miyamoto T, Weissman IL. 2000. A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. Nature 404:193-7.
17. Kaushansky K. 2008. Historical review: megakaryopoiesis and thrombopoiesis. Blood 111:981-6.
18. Deutsch VR, Tomer A. 2006. Megakaryocyte development and platelet production. Br J Haematol 134:453-66.
19. Schulze H, Shivdasani RA. 2004. Molecular mechanisms of megakaryocyte differentiation. Semin Thromb Hemost 30:389-98.

577   20.   Cecchetti L, Tolley ND, Michetti N, Bury L, Weyrich AS, Gresele P. 2011. Megakaryocytes
578   differentially sort mRNAs for matrix metalloproteinases and their inhibitors into platelets: a
579   mechanism for regulating synthetic events. Blood 118:1903-11.
580   21.   Matsumura I, Kanakura Y. 2002. Molecular control of megakaryopoiesis and thrombopoiesis. Int
581   J Hematol 75:473-83.
582   22.   Severin S, Ghevaert C, Mazharian A. 2010. The mitogen-activated protein kinase signaling
583   pathways: role in megakaryocyte differentiation. J Thromb Haemost 8:17-26.
584   23.   Kirito K, Osawa M, Morita H, Shimizu R, Yamamoto M, Oda A, Fujita H, Tanaka M, Nakajima K,
585   Miura Y, Ozawa K, Komatsu N. 2002. A functional role of Stat3 in in vivo megakaryopoiesis. Blood
586   99:3220-7.
587   24.   Olthof SG, Fatrai S, Drayer AL, Tyl MR, Vellenga E, Schuringa JJ. 2008. Downregulation of signal
588   transducer and activator of transcription 5 (STAT5) in CD34+ cells promotes megakaryocytic
589   development, whereas activation of STAT5 drives erythropoiesis. Stem Cells 26:1732-42.
590   25.   Beaulieu LM, Lin E, Morin KM, Tanriverdi K, Freedman JE. 2011. Regulatory effects of TLR2 on
591   megakaryocytic cell function. Blood 117:5963-74.
592   26.   Racke FK, Lewandowska K, Goueli S, Goldfarb AN. 1997. Sustained activation of the extracellular
593   signal-regulated kinase/mitogen-activated protein kinase pathway is required for megakaryocytic
594   differentiation of K562 cells. J Biol Chem 272:23366-70.
595   27.   Goldfarb AN. 2007. Transcriptional control of megakaryocyte development. Oncogene 26:6795-
596   802.
597   28.   Tijssen MR, Ghevaert C. 2013. Transcription factors in late megakaryopoiesis and related platelet
598   disorders. J Thromb Haemost 11:593-604.
599   29.   Muntean AG, Pang L, Poncz M, Dowdy SF, Blobel GA, Crispino JD. 2007. Cyclin D-Cdk4 is regulated
600   by GATA-1 and required for megakaryocyte growth and polyploidization. Blood 109:5199-207.
601   30.   Lordier L, Bluteau D, Jalil A, Legrand C, Pan J, Rameau P, Jouni D, Bluteau O, Mercher T, Leon C,
602   Gachet C, Debili N, Vainchenker W, Raslova H, Chang Y. 2012. RUNX1-induced silencing of non-
603   muscle myosin heavy chain IIB contributes to megakaryocyte polyploidization. Nat Commun
604   3:717.
605   31.   Lopez JJ, Palazzo A, Chaabane C, Albarran L, Polidano E, Lebozec K, Dally S, Nurden P, Enouf J,
606   Debili N, Bobe R. 2013. Crucial role for endoplasmic reticulum stress during megakaryocyte
607   maturation. Arterioscler Thromb Vasc Biol 33:2750-8.
608   32.   You T, Wang Q, Zhu L. 2016. Role of autophagy in megakaryocyte differentiation and platelet
609   formation. Int J Physiol Pathophysiol Pharmacol 8:28-34.
610   33.   De Botton S, Sabri S, Daugas E, Zermati Y, Guidotti JE, Hermine O, Kroemer G, Vainchenker W,
611   Debili N. 2002. Platelet formation is the consequence of caspase activation within
612   megakaryocytes. Blood 100:1310-7.
613   34.   Chen S, Su Y, Wang J. 2013. ROS-mediated platelet generation: a microenvironment-dependent
614   manner for megakaryocyte proliferation, differentiation, and maturation. Cell Death Dis 4:e722.
615   35.   Sardina JL, Lopez-Ruano G, Sanchez-Abarca LI, Perez-Simon JA, Gaztelumendi A, Trigueros C,
616   Llanillo M, Sanchez-Yague J, Hernandez-Hernandez A. 2010. p22phox-dependent NADPH oxidase
617   activity is required for megakaryocytic differentiation. Cell Death Differ 17:1842-54.
618   36.   Mostafa SS, Miller WM, Papoutsakis ET. 2000. Oxygen tension influences the differentiation,
619   maturation and apoptosis of human megakaryocytes. Br J Haematol 111:879-89.
620   37.   McCrann DJ, Eliades A, Makitalo M, Matsuno K, Ravid K. 2009. Differential expression of NADPH
621   oxidases in megakaryocytes and their role in polyploidy. Blood 114:1243-9.
622   38.   Sattler M, Winkler T, Verma S, Byrne CH, Shrikhande G, Salgia R, Griffin JD. 1999. Hematopoietic
623   growth factors signal through the formation of reactive oxygen species. Blood 93:2928-35.

39. Eliades A, Papadantonakis N, Ravid K. 2010. New roles for cyclin E in megakaryocytic
polyploidization. J Biol Chem 285:18909-17.
40. Siddiqui NF, Shabrani NC, Kale VP, Limaye LS. 2011. Enhanced generation of megakaryocytes from
umbilical cord blood-derived CD34(+) cells expanded in the presence of two nutraceuticals,
docosahexaenoic acid and arachidonic acid, as supplements to the cytokine-containing medium.
Cytotherapy 13:114-28.
41. Witt O, Sand K, Pekrun A. 2000. Butyrate-induced erythroid differentiation of human K562
leukemia cells involves inhibition of ERK and activation of p38 MAP kinase pathways. Blood
95:2391-6.
42. Kim KW, Kim SH, Lee EY, Kim ND, Kang HS, Kim HD, Chung BS, Kang CD. 2001. Extracellular signal-
regulated kinase/90-KDA ribosomal S6 kinase/nuclear factor-kappa B pathway mediates phorbol
12-myristate 13-acetate-induced megakaryocytic differentiation of K562 cells. J Biol Chem
276:13186-91.
43. Ojima Y, Duncan MT, Nurhayati RW, Taya M, Miller WM. 2013. Synergistic effect of hydrogen
peroxide on polyploidization during the megakaryocytic differentiation of K562 leukemia cells by
PMA. Exp Cell Res 319:2205-15.
44. Chaman N, Iqbal MA, Siddiqui FA, Gopinath P, Bamezai RN. 2015. ERK2-Pyruvate Kinase Axis
Permits Phorbol 12-Myristate 13-Acetate-induced Megakaryocyte Differentiation in K562 Cells. J
Biol Chem 290:23803-15.
45. Lee CH, Yun HJ, Kang HS, Kim HD. 1999. ERK/MAPK pathway is required for changes of cyclin D1
and B1 during phorbol 12-myristate 13-acetate-induced differentiation of K562 cells. IUBMB Life
48:585-91.
46. Nurhayati RW, Ojima Y, Nomura N, Taya M. 2014. Promoted megakaryocytic differentiation of
K562 cells through oxidative stress caused by near ultraviolet irradiation. Cell Mol Biol Lett 19:590-
600.
47. Huang R, Zhao L, Chen H, Yin RH, Li CY, Zhan YQ, Zhang JH, Ge CH, Yu M, Yang XM. 2014.
Megakaryocytic differentiation of K562 cells induced by PMA reduced the activity of respiratory
chain complex IV. PLoS One 9:e96246.
48. Colosetti P, Puissant A, Robert G, Luciano F, Jacquel A, Gounon P, Cassuto JP, Auberger P. 2009.
Autophagy is an important event for megakaryocytic differentiation of the chronic myelogenous
leukemia K562 cell line. Autophagy 5:1092-8.
49. Limb JK, Yoon S, Lee KE, Kim BH, Lee S, Bae YS, Jhon GJ, Kim J. 2009. Regulation of megakaryocytic
differentiation of K562 cells by FosB, a member of the Fos family of AP-1 transcription factors. Cell
Mol Life Sci 66:1962-73.
50. Rojnuckarin P, Drachman JG, Kaushansky K. 1999. Thrombopoietin-induced activation of the
mitogen-activated protein kinase (MAPK) pathway in normal megakaryocytes: role in
endomitosis. Blood 94:1273-82.
51. Lin YL, Liao CL, Chen LK, Yeh CT, Liu CI, Ma SH, Huang YY, Huang YL, Kao CL, King CC. 1998. Study
of Dengue virus infection in SCID mice engrafted with human K562 cells. J Virol 72:9729-37.
52. Elshuber S, Allison SL, Heinz FX, Mandl CW. 2003. Cleavage of protein prM is necessary for
infection of BHK-21 cells by tick-borne encephalitis virus. J Gen Virol 84:183-191.
53. Olagnier D, Peri S, Steel C, van Montfoort N, Chiang C, Beljanski V, Slifker M, He Z, Nichols CN, Lin
R, Balachandran S, Hiscott J. 2014. Cellular oxidative stress response controls the antiviral and
apoptotic programs in dengue virus-infected dendritic cells. PLoS Pathog 10:e1004566.
54. Obexer P, Hagenbuchner J, Unterkircher T, Sachsenmaier N, Seifarth C, Bock G, Porto V, Geiger K,
Ausserlechner M. 2009. Repression of BIRC5/survivin by FOXO3/FKHRL1 sensitizes human
neuroblastoma cells to DNA damage-induced apoptosis. Mol Biol Cell 20:2041-8.

55.   Caparros-Perez E, Teruel-Montoya R, Lopez-Andreo MJ, Llanos MC, Rivera J, Palma-Barqueros V,
672   Blanco JE, Vicente V, Martinez C, Ferrer-Marin F. 2017. Comprehensive comparison of neonate
673   and adult human platelet transcriptomes. PLoS One 12:e0183042.

674   56.   Majek P, Reicheltova Z, Stikarova J, Suttnar J, Sobotkova A, Dyr JE. 2010. Proteome changes in
675   platelets activated by arachidonic acid, collagen, and thrombin. Proteome Sci 8:56.

676   57.   Park SM, Jung HY, Kim HO, Rhim H, Paik SR, Chung KC, Park JH, Kim J. 2002. Evidence that alpha-
677   synuclein functions as a negative regulator of Ca(++)-dependent alpha-granule release from
678   human platelets. Blood 100:2506-14.

679   58.   da Costa CA, Ancolio K, Checler F. 2000. Wild-type but not Parkinson's disease-related ala-53 -->
680   Thr mutant alpha -synuclein protects neuronal cells from apoptotic stimuli. J Biol Chem
681   275:24065-9.

682   59.   Polyak SJ, Rawls WE, Harnish DG. 1991. Characterization of Pichinde virus infection of cells of the
683   monocytic lineage. J Virol 65:3575-82.

684   60.   Seamone ME, Wang W, Acott P, Beck PL, Tibbles LA, Muruve DA. 2010. MAP kinase activation
685   increases BK polyomavirus replication and facilitates viral propagation in vitro. J Virol Methods
686   170:21-9.

687   61.   Wang W, Wang Y, Debing Y, Zhou X, Yin Y, Xu L, Herrera Carrillo E, Brandsma JH, Poot RA, Berkhout
688   B, Neyts J, Peppelenbosch MP, Pan Q. 2017. Biological or pharmacological activation of protein
689   kinase C alpha constrains hepatitis E virus replication. Antiviral Res 140:1-12.

690   62.   Golding H, Manischewitz J, Vujcic L, Blumenthal R, Dimitrov DS. 1994. The phorbol ester phorbol
691   myristate acetate inhibits human immunodeficiency virus type 1 envelope-mediated fusion by
692   modulating an accessory component(s) in CD4-expressing cells. J Virol 68:1962-9.

693   63.   Hou W, Gibbs JS, Lu X, Brooke CB, Roy D, Modlin RL, Bennink JR, Yewdell JW. 2012. Viral infection
694   triggers rapid differentiation of human blood monocytes into dendritic cells. Blood 119:3128-31.

695   64.   Zeng YX, el-Deiry WS. 1996. Regulation of p21WAF1/CIP1 expression by p53-independent
696   pathways. Oncogene 12:1557-64.

697   65.   Bagga S, Bouchard MJ. 2014. Cell cycle regulation during viral infection. Methods Mol Biol
698   1170:165-227.

699   66.   Phoolcharoen W, Smith DR. 2004. Internalization of the dengue virus is cell cycle modulated in
700   HepG2, but not Vero cells. J Med Virol 74:434-41.

701   67.   Helt AM, Harris E. 2005. S-phase-dependent enhancement of dengue virus 2 replication in
702   mosquito cells, but not in human cells. J Virol 79:13218-30.

703   68.   Bua G, Manaresi E, Bonvicini F, Gallinella G. 2016. Parvovirus B19 Replication and Expression in
704   Differentiating Erythroid Progenitor Cells. PLoS One 11:e0148547.

705   69.   Zanini F, Pu SY, Bekerman E, Einav S, Quake SR. 2018. Single-cell transcriptional dynamics of
706   flavivirus infection. Elife 7.

707   70.   Zauli G, Catani L, Gibellini D, Re MC, Milani D, Borgatti P, Bassini A, La Placa M, Capitani S. 1995.
708   The CD4 receptor plays essential but distinct roles in HIV-1 infection and induction of apoptosis in
709   primary bone marrow GPIIb/IIIa+ megakaryocytes and the HEL cell line. Br J Haematol 91:290-8.

710   71.   Zauli G, Catani L, Gibellini D, Re MC, Vianelli N, Colangeli V, Celeghini C, Capitani S, La Placa M.
711   1996. Impaired survival of bone marrow GPIIb/IIa+ megakaryocytic cells as an additional
712   pathogenetic mechanism of HIV-1-related thrombocytopenia. Br J Haematol 92:711-7.

713   72.   Lutteke N, Raftery MJ, Lalwani P, Lee MH, Giese T, Voigt S, Bannert N, Schulze H, Kruger DH,
714   Schonrich G. 2010. Switch to high-level virus replication and HLA class I upregulation in
715   differentiating megakaryocytic cells after infection with pathogenic hantavirus. Virology 405:70-
716   80.

717   73.   Zhang M, Evans S, Yuan J, Ratner L, Koka PS. 2010. HIV-1 determinants of thrombocytopenia at
718   the stage of CD34+ progenitor cell differentiation in vivo lie in the viral envelope gp120 V3 loop
719   region. Virology 401:131-6.
720   74.   Henke JI, Goergen D, Zheng J, Song Y, Schuttler CG, Fehr C, Junemann C, Niepmann M. 2008.
721   microRNA-122 stimulates translation of hepatitis C virus RNA. EMBO J 27:3300-10.
722   75.   Ambrosio AL, Di Pietro SM. 2019. Mechanism of platelet alpha-granule biogenesis: study of cargo
723   transport and the VPS33B-VPS16B complex in a model system. Blood Adv 3:2617-2626.
724   76.   Baatout S, Chatelain B, Staquet P, Symann M, Chatelain C. 1999. Protein content and number of
725   nucleolar organizer regions are enhanced during phorbol ester-induced differentiation of cultured
726   human megakaryocytic cells. Anticancer Res 19:3229-35.
727   77.   Daffis S, Szretter KJ, Schriewer J, Li J, Youn S, Errett J, Lin TY, Schneller S, Zust R, Dong H, Thiel V,
728   Sen GC, Fensterl V, Klimstra WB, Pierson TC, Buller RM, Gale M, Jr., Shi PY, Diamond MS. 2010. 2'-
729   O methylation of the viral mRNA cap evades host restriction by IFIT family members. Nature
730   468:452-6.
731   78.   Gurney AL, de Sauvage FJ. 1996. Dissection of c-Mpl and thrombopoietin function: studies of
732   knockout mice and receptor signal transduction. Stem Cells 14 Suppl 1:116-23.
733   79.   Kaushansky K. 2005. The molecular mechanisms that control thrombopoiesis. J Clin Invest
734   115:3339-47.
735   80.   Roux PP, Blenis J. 2004. ERK and p38 MAPK-activated protein kinases: a family of protein kinases
736   with diverse biological functions. Microbiol Mol Biol Rev 68:320-44.
737   81.   Fichelson S, Freyssinier JM, Picard F, Fontenay-Roupie M, Guesnu M, Cherai M, Gisselbrecht S,
738   Porteu F. 1999. Megakaryocyte growth and development factor-induced proliferation and
739   differentiation are regulated by the mitogen-activated protein kinase pathway in primitive cord
740   blood hematopoietic progenitors. Blood 94:1601-13.
741   82.   Herrera R, Hubbell S, Decker S, Petruzzelli L. 1998. A role for the MEK/MAPK pathway in PMA-
742   induced cell cycle arrest: modulation of megakaryocytic differentiation of K562 cells. Exp Cell Res
743   238:407-14.
744   83.   Whalen AM, Galasinski SC, Shapiro PS, Nahreini TS, Ahn NG. 1997. Megakaryocytic differentiation
745   induced by constitutive activation of mitogen-activated protein kinase kinase. Mol Cell Biol
746   17:1947-58.
747   84.   Nagila A, Netsawang J, Suttitheptumrong A, Morchang A, Khunchai S, Srisawat C, Puttikhunt C,
748   Noisakran S, Yenchitsomanus PT, Limjindaporn T. 2013. Inhibition of p38MAPK and CD137
749   signaling reduce dengue virus-induced TNF-alpha secretion and apoptosis. Virol J 10:105.
750   85.   Sreekanth GP, Chuncharunee A, Sirimontaporn A, Panaampon J, Noisakran S, Yenchitsomanus PT,
751   Limjindaporn T. 2016. SB203580 Modulates p38 MAPK Signaling and Dengue Virus-Induced Liver
752   Injury by Reducing MAPKAPK2, HSP27, and ATF2 Phosphorylation. PLoS One 11:e0149486.
753   86.   Ashour J, Laurent-Rolle M, Shi PY, Garcia-Sastre A. 2009. NS5 of dengue virus mediates STAT2
754   binding and degradation. J Virol 83:5408-18.
755   87.   Huang Z, Richmond TD, Muntean AG, Barber DL, Weiss MJ, Crispino JD. 2007. STAT1 promotes
756   megakaryopoiesis downstream of GATA-1 in mice. J Clin Invest 117:3890-9.
757   88.   Yoshimura A, Naka T, Kubo M. 2007. SOCS proteins, cytokine signalling and immune regulation.
758   Nat Rev Immunol 7:454-65.
759   89.   Yuan ZL, Guan YJ, Chatterjee D, Chin YE. 2005. Stat3 dimerization regulated by reversible
760   acetylation of a single lysine residue. Science 307:269-73.
761   90.   Rebe C, Vegran F, Berger H, Ghiringhelli F. 2013. STAT3 activation: A key factor in tumor
762   immunoescape. JAKSTAT 2:e23010.

91. Wei J, Yuan Y, Jin C, Chen H, Leng L, He F, Wang J. 2012. The ubiquitin ligase TRAF6 negatively
regulates the JAK-STAT signaling pathway by binding to STAT3 and mediating its ubiquitination.
PLoS One 7:e49567.
92. Shuai K. 2000. Modulation of STAT signaling by STAT-interacting proteins. Oncogene 19:2638-44.
93. Krebs DL, Hilton DJ. 2001. SOCS proteins: negative regulators of cytokine signaling. Stem Cells
19:378-87.
94. Corzo CA, Cotter MJ, Cheng P, Cheng F, Kusmartsev S, Sotomayor E, Padhya T, McCaffrey TV,
McCaffrey JC, Gabrilovich DI. 2009. Mechanism regulating reactive oxygen species in tumor-
induced myeloid-derived suppressor cells. J Immunol 182:5693-701.
95. Lambeth JD. 2004. NOX enzymes and the biology of reactive oxygen. Nat Rev Immunol 4:181-9.
96. Zorov DB, Juhaszova M, Sollott SJ. 2014. Mitochondrial reactive oxygen species (ROS) and ROS-
induced ROS release. Physiol Rev 94:909-50.
97. Turrens JF. 2003. Mitochondrial formation of reactive oxygen species. J Physiol 552:335-44.
98. Salmeen A, Andersen JN, Myers MP, Meng TC, Hinks JA, Tonks NK, Barford D. 2003. Redox
regulation of protein tyrosine phosphatase 1B involves a sulphenyl-amide intermediate. Nature
423:769-73.
99. Lee SR, Kwon KS, Kim SR, Rhee SG. 1998. Reversible inactivation of protein-tyrosine phosphatase
1B in A431 cells stimulated with epidermal growth factor. J Biol Chem 273:15366-72.
100. Zhang J, Wang X, Vikash V, Ye Q, Wu D, Liu Y, Dong W. 2016. ROS and ROS-Mediated Cellular
Signaling. Oxid Med Cell Longev 2016:4350965.
101. Choi K, Ryu SW, Song S, Choi H, Kang SW, Choi C. 2010. Caspase-dependent generation of reactive
oxygen species in human astrocytoma cells contributes to resistance to TRAIL-mediated
apoptosis. Cell Death Differ 17:833-45.
102. Huang X, Trinh T, Aljoufi A, Broxmeyer HE. 2018. Hypoxia Signaling Pathway in Stem Cell
Regulation: Good and Evil. Curr Stem Cell Rep 4:149-157.
103. Shinohara A, Imai Y, Nakagawa M, Takahashi T, Ichikawa M, Kurokawa M. 2014. Intracellular
reactive oxygen species mark and influence the megakaryocyte-erythrocyte progenitor fate of
common myeloid progenitors. Stem Cells 32:548-57.
104. Motohashi H, Kimura M, Fujita R, Inoue A, Pan X, Takayama M, Katsuoka F, Aburatani H, Bresnick
EH, Yamamoto M. 2010. NF-E2 domination over Nrf2 promotes ROS accumulation and
megakaryocytic maturation. Blood 115:677-86.
105. Schwarz KB. 1996. Oxidative stress during viral infection: a review. Free Radic Biol Med 21:641-9.
106. Zeeshan HM, Lee GH, Kim HR, Chae HJ. 2016. Endoplasmic Reticulum Stress and Associated ROS.
Int J Mol Sci 17:327.

 Figure legends:

Figure 1: Panel A: Schematic diagram of Megakaryopoiesis: The schematic diagram shows some of the

lineages that are produced during differentiation of Hematopoietic stem cells (HSC) to Megakaryocytes

(Mk). The formation of multilobed nucleus has also been depicted. **Panel B: Visualization of endomitosis**

and multi-lobed nuclear formation following PMA-treatment. K562 cells treated with either DMSO or

50 nM PMA for 6 days were stained with Alexa Fluor-568 Phalloidin, mounted onto glass slides using

mountant supplemented with DAPI and visualized in a confocal microscope. The red color indicates the

plasma membrane and blue color shows nuclei. The inset shows a cell with multilobed nucleus. **Panel C**:

Expression of platelet specific surface markers: K562 cells treated with PMA for 0, 3 or 6 days were

immune-stained for expression of the indicated surface marker protein using fluor-conjugated primary

antibodies. The mean fluorescence intensity (MFI) corresponding to each day was quantified. For each

surface marker the MFI at day 0 was arbitrarily set to 1 and those at day 3 and day 6, expressed as the

relative fold change with respect to that. The error bars represent standard deviation. The significance was

calculated by Student's t-test (*, **, *** respectively indicate P-values <0.5, <0.1 and <0.01). **Panel D**:

K562 cells treated with PMA for 0 or 3 or 6 days were fixed, permeabilized and stained for intracellular

DNA using Propidium iodide (PI). The PI stain was quantified by flow cytometry and mean number of

cells having ploidy of >4N was plotted. The error bars represent standard deviation. **Panel E:** The total

RNA from K562 cells treated with either DMSO or 50 nM PMA or 1mM Sodium Butyrate (NaB) for 3 or

6 days was extracted and purified. The total RNA was reverse transcribed using random hexamers and the

cDNA obtained used for real-time PCR estimation of mRNA transcripts from indicated genes. The C_T value

corresponding to each was normalized to that of GAPDH. The normalized value at day 0 was taken

arbitrarily as 1 and those at day 3 and 6 expressed as fold-change with respect to that.

Figure 2: DENV replication is increased upon PMA treatment of infected K562 cells: Panel A: K562

cells infected with DENV at an MOI of 0.1 were maintained in growth media supplemented with either

DMSO or 50nM PMA and fixed with PFA at 0 hour or 3 days or 6 days post-infection. The cells were

permeabilized and immune-stained for intracellular DENV antigen and the fluorescence quantified by flow

cytometry. The fluorescence in cells fixed at 0 hour was used to gate for DENV antigen positive cells. The

number of antigen positive cells were quantified and plotted. The error bars represent standard deviation

and the significance was calculated by Student's t-test (*** indicates P <0.001). **Panel B**: Total RNA from

K562 cells treated similarly as in panel A was extracted, purified and reverse-transcribed using random

hexamers. The cDNA was used for real-time PCR comparison of DENV RNA level at 3 or 6 days to that

at 0 hour, normalized to that of GAPDH mRNA. The value at 0 hour was taken arbitrarily as 1 and that at

other time points expressed as fold-change over that. The error bars represent standard deviation and the

significance was calculated by Student's t-test (** indicates P <0.01). **Panel C**: The culture supernatant of

cell infected with MOI of 0.1 and subsequently treated as in panel A was collected and used for

quantification of focus-forming unit/ml (FFU/ml) on Vero cell monolayers. The FFU/ml for each sample

was transformed to their logarithmic value to the base of 10 and plotted. The error bars represent standard

deviation and the significance was calculated by Student's t-test (** indicates P <0.01). **Panel D**:

Uninfected K562 cells treated with either DMSO or PMA for 3 days were washed with PBS and incubated

in either mock- or DENV inoculum (10 MOI) on ice for 2 hours. The cells were then transferred to 37^0C

and further incubated for 2 hours. Subsequently cells were washed, treated with 0.25% trypsin for 5 minutes

and the total RNA extracted and purified. 1.0 µg of total RNA was reverse-transcribed with random

hexamers and the cDNA used in real-time PCR for comparison of DENV genomic RNA normalized to

GAPDH transcripts. The level in mock-infected cells was taken as 1 and that in DENV infected cells

calculated as fold-change over that. The relative enrichment of DENV RNA compared to mock-infected

cells was quantified, transformed to their logarithmic value to the base of 2 and plotted. The error bars show

standard deviation and the statistical significance was calculated by Student's t-test (ns= not significant).

Figure 3: DENV replication in differentiating K562 cells interfere with PMA-induced signaling:
Panel A: Schematic showing comparison of signaling activated upon TPO binding to c-Mpl and by PMA.
Panel B: Immunoblot for signaling protein activated by PMA in K562 cells. K562 cells either mock-infected or DENV-infected were incubated in PMA-supplemented growth media and the total protein extracted after 3 or 6 days. The protein in each lysate was denatured, resolved in SDS-PAGE and immunoblotted for proteins indicated at right. Immunoblot for b-actin was performed for each batch of lysate used. The relative molecular mass of the marker protein is indicated on the left. **Panel C: Quantification of immunoblot bands in panel B**. the intensity of the immunoblot band for indicated proteins were quantified, normalized to that of b-actin and the ratio calculated before plotting. The error bars represent standard deviation of the ratio derived from at least two independent experiments, performed in multiple replicates.

Figure 4: Effect of DENV replication on cell surface marker, polyploidy and apoptosis in differentiating K562 cells. Panel A: Suppression of platelet-specific surface marker protein. K562 infected with DENV at two different MOI, were incubated in PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were immune-stained for the indicated surface marker proteins and the fluorescence quantified by flow cytometry. The mean fluorescence intensity was plotted. The error bars show standard deviation derived from at least two independent experiments in multiple replicates. **Panel B: DENV replication inhibits endomitosis**. DENV infected K562 were incubated in PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were fixed and stained with propidium iodide and the fluorescence intensity measured by flow cytometry. The number of cells with ploidy of >4N were quantified and plotted. The error bars show standard deviation of the mean from two independent experiments performed in multiple replicates. **Panel C: Caspase cleavage assay in differentiated DENV infected cells**. Mock-infected or DENV-infected K562 were incubated in either DMSO or PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were incubated with an substrate that can be cleaved

by intracellular Caspase3/7 to generate a luminescent product. The luminescence was measured in a

luminometer and plotted. The total luminescence at day 3 in DMSO-treated uninfected cells was arbitrarily

taken as 1 and the rest plotted as fold-change over that. The error bars show standard deviation of the mean

from two independent experiments performed in multiple replicates.

Figure 5: DENV infection suppresses ROS generated during PMA-induced Megakaryopoiesis in

K562 cells: Panel A. Mock-infected or DENV-infected K562 cells were incubated in either DMSO or

PMA-supplemented growth media for 3 or 6 days. Subsequently the cells were stained with H_2DCFDA and

the fluorescence quantified by flow cytometry. The mean fluorescence intensity in mock-infected and

DMSO-treated cells at 3 days was arbitrarily taken as 1 and the rest calculated as fold-change over that.

The error bars show standard deviation of the mean from two independent experiments performed in

multiple replicates. **Panel B**. K562 cells either mock-infected or DENV-infected were immediately

incubated in growth media supplemented with 50 nM PMA in addition to either 0 or 1 or 3 mM N-Acetyl

Cysteine (NAC) as indicated. At 3 or 6 days post-infection the cells were stained with H_2DCFDA and the

fluorescence quantified by flow-cytometry. The mean fluorescence intensity in mock-infected cells treated

with PMA and 0 mM NAC was arbitrarily taken as 1 and that in the rest expressed as fold change with

respect to that. The error bars show standard deviation of the mean from two independent experiments

performed in multiple replicates.

Figure 5: DENV replication interferes with PMA-induced transcriptome changes. Panel A: The

transcriptome in K562 cells treated with 50 nM PMA for 6 days was compared to that after 0 hour by Next-

generation sequencing. The genes differentially expressed after 6 days of treatment, either upregulated by

≥ 1.5 fold or downregulated by ≤ 0.5 fold with a P-value of ≤ 0.05, when compared to 0 hour were used to

draw a Volcano plot prepared using in-house written Python script. The X-axis represents the logarithmic

value to the base of 2 of the fold-change in gene expression. The Y-axis represents the negative logarithmic

to the base of 10 of the P-value. **Panel B**: The transcriptome in K562 cells, either uninfected or DENV-

infected and treated with 50 nM PMA or 6 days was compared to uninfected cells at 0 hour by NGS to find

differentially expressed genes. The level of differential expression of each gene was then compared between

uninfected and DENV-infected cells to draw the list of genes the PMA-induced deregulation of which is

influenced by infection. A volcano plot was prepared from this list using in-house written Python script.

The X-axis represents the logarithmic value to the base of 2 of the fold-change in gene expression. The Y-

axis represents the negative logarithmic to the base of 10 of the P-value. **Panel C**: **Heatmap of genes**

associated with ROS-pathway. The genes that are associated with the ROS-pathway in panel B were

selected and enriched using the Metascape software. The PMA-induced fold-change in the transcript level

of these genes after 6 days when compared to uninfected cells treated with PMA for 0 hour (column 1) in

either uninfected cells (column 2) or DENV-infected cells (column 3), was represented by a heatmap

prepared using the Morpheus algorithm (https://software.broadinstitute.org/morpheus). The colors

represent the trend of deregulation corresponding to each gene shown on the right.

Figure 1

A

B

C

D

E

Figure 2

Figure 3

A

B

C

Mock infected   DENV infected

Figure 4

Figure 5

A

B

Figure 6