Deep ancestral introgression shapes evolutionary history of dragonflies and

damselflies

Anton Suvorov[1$], Celine Scornavacca[2,*], M. Stanley Fujimoto[3], Paul Bodily[4], Mark

Clement[3], Keith A. Crandall[5], Michael F. Whiting[6,7], Daniel R. Schrider[1,*] and Seth M.

Bybee[6,7,*]

[1]Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC

[2]Institut des Sciences de l'Evolution Université de Montpellier, CNRS, IRD, EPHE CC 064,

Place Eugène Bataillon, 34095 Montpellier Cedex 05, France.

[3]Department of Biology, Brigham Young University, Provo, UT

[4]Department of Computer Science, Idaho State University, Pocatello, ID

[5]Computational Biology Institute, Department of Biostatistics and Bioinformatics, Milken

Institute School of Public Health, George Washington University, Washington, DC

[6]Department of Biology, Brigham Young University, Provo, UT

[7]M.L. Bean Museum, Brigham Young University, Provo, UT

$Correspondence: antony.suvorov@med.unc.edu

*Equal contributions

SUMMARY

Introgression is arguably one of the most important biological processes in the evolution of

groups of related species, affecting at least 10% of the extant species in the animal kingdom.

Introgression reduces genetic divergence between species, and in some cases can be highly

beneficial, facilitating rapid adaptation to ever-changing environmental pressures. Introgression

also significantly impacts inference of phylogenetic species relationships where a strictly binary

tree model cannot adequately explain reticulate net-like species relationships. Here we use

phylogenomic approaches to understand patterns of introgression along the evolutionary history

of a unique, non-model insect system: dragonflies and damselflies (Odonata). We demonstrate

that introgression is a pervasive evolutionary force across various taxonomic levels within

Odonata. In particular, we show that the morphologically "intermediate" species of

Anisozygoptera (one of the three primary suborders within Odonata besides Zygoptera and

Anisoptera), which retain phenotypic characteristics of the other two suborders, experienced high

levels of introgression likely coming from zygopteran genomes. Additionally, we found evidence

for multiple cases of deep inter-superfamilial ancestral introgression.

INTRODUCTION

In recent years there has been a large body of evidence amassed demonstrating that multiple

parts of the Tree of Life did not evolve according to a strictly bifurcating phylogeny [1, 2].

Instead, many organisms experience reticulate network-like evolution that is caused by an

exchange of inter-specific genetic information via various biological processes. In particular,

lateral gene transfer, incomplete lineage sorting (ILS) and introgression can result in gene trees

that are discordant with the species tree [3, 4]. Lateral transfer and introgression both involve

gene flow following speciation, thereby producing "reticulate" phylogenies. Incomplete lineage

sorting (ILS), on the other hand, occurs when three or more lineages coalesce in their ancestral

population. This often arises in an order that conflicts with that of the true species tree, and thus

involves no post-speciation gene flow and therefore does not contribute to reticulate evolution.

Phylogenetic species-gene tree incongruence observed in empirical data can provide insight into

underlying biological factors that shape the evolutionary trajectories of a set of taxa. The major

source of reticulate evolution for eukaryotes is introgression where it affects approximately 25%

of flowering plant and 10% of animal species [1, 5]. Introgressed alleles can be fitness-neutral,

deleterious [6] or adaptive [7, 8]. For example, adaptive introgression has been shown to provide

an evolutionary rescue from polluted habitats in gulf killifish (*Fundulus grandis*) [7], yielded

mimicry adaptations among *Heliconius* butterflies [9] and archaic introgression has facilitated

adaptive evolution of altitude tolerance [10], immunity and metabolism in modern humans [11].

Additionally, hybridization and introgression are important and often overlooked mechanisms of

invasive species establishment and spread [12].

Odonata, the insect order that contains dragonflies and damselflies, lacks a strongly

supported backbone tree to clearly resolve higher-level phylogenetic relationships [13, 14].

Current evidence places odonates together with Ephemeroptera (mayflies) as the living

representatives of the most ancient insect lineages to have evolved wings and active flight [15].

Odonates possess unique anatomical and morphological features such as a specialized body

form, specialized wing venation, a distinctive form of muscle attachment to the wing base [16]

allowing for direct flight and accessory (secondary) male genitalia that support certain unique

behaviors (e.g., sperm competition). They are among the most adept flyers of all animals and are

exclusively carnivorous insects relying primarily on vision [17, 18] to capture prey. During their

immature stage they are fully aquatic and spend much of their adult life in flight.

Biogeographically, odonates exhibit species ranges varying from worldwide dispersal [19] to

island-endemic. Odonates also play crucial ecological roles in local freshwater communities,

being a top invertebrate predator as both adults and immatures [20]. Due to this combination of

characteristics, odonates are quickly becoming model organisms to study specific questions in

ecology, physiology and evolution [21, 22]. However, the extent of introgression at the genomic

scale within Odonata remains largely unknown.

Two early attempts to tackle introgression/hybridization patterns within Odonata were

undertaken in [23, 24]. The studies showed that two closely related species of damselflies,

Ischnura graellsii and *I. elegans,* can hybridize under laboratory conditions and that genital

morphology of male hybrids shares features with putative hybrids from *I. graellsii–I. elegans*

natural allopatric populations [23]. The existence of abundant hybridization and introgression in

natural populations of *I. graellsii* and *I. elegans* has received further support from an analysis of

microsatellite data [25]. Putative hybridization events have also been identified in a pair of

calopterygoid damselfly species, *Mnais costalis* and *M. pruinosa* based on the analyses of two

molecular loci (mtDNA and nucDNA) [26], and between *Calopteryx virgo* and *C. splendens*

using 16S ribosomal DNA and 40 random amplified polymorphic DNA (RAPD) markers [27]. A

more recent study identified an inter-specific hybridization between two cordulegasterid

dragonfly species, *Cordulegaster boltonii* and *C. trinacriae* using two molecular markers

(mtDNA and nucDNA) and geometric morphometrics [28]. Furthermore, in various biological

systems, the empirical evidence shows that hybridization can potentially lead to intermediate

phenotypes [29] observed at molecular level (e.g. semidominant expression in inter-specific

hybrids [30]) as well as organismal morphology (e.g. [31-33]). The Anisozygoptera suborder,

which contains only three extant species, retains traits shared with both dragonflies and

damselflies (hence its taxonomic name), ranging from morphology and anatomical structures

[34] to behavior and flight biomechanics [35]. These characteristics could suggest a hybrid origin

of this suborder. The potential introgression scenario for Anisozygoptera is yet to be formally

tested using genome-wide data.

Here we present a comprehensive analysis of transcriptomic data from 83 odonate

species. First, we reconstruct a robust phylogenetic backbone using up to 4341 genetic loci for

the order and discuss its evolutionary history spanning from the Carboniferous period (~360

mya) to present day. Furthermore, in light of the "intermediate" phenotypic nature of

Anisozygoptera, we investigate phylogenetic signatures of introgression within Odonata. Most

notably, we identify a strong signal of deep introgression in the Anisozygoptera suborder,

species of which possess traits of both main suborders, Anisoptera and Zygoptera. Although the

strongest signatures of introgression are found in Anisozygoptera, we find evidence that

introgression was pervasive in Odonata throughout its entire evolutionary history.

RESULTS

Phylogenetic Inference

We compiled transcriptomic data for 83 odonate species including 49 new transcriptomes

sequenced for this study (Data S1). To assess effects of various steps of our phylogenetic

pipeline on species tree inference, we examined different methods of sequence homology

detection, multiple sequence alignment strategies, postprocessing filtering procedures and tree

estimation methods. Specifically, three types of homologous loci (gene clusters) were used to

develop our supermatrices, namely 1603 conserved single-copy orthologs (CO), 1643 all single-

copy orthologs (AO) and 4341 paralogy-parsed orthologs (PO) with ≥ 42 (~50%) species present

(for more details, see Material and Methods). To date, our data represents the most

comprehensive resource available for Odonata in terms of gene sampling. Each gene cluster was

aligned, trimmed and concatenated resulting in five main supermatrices, CO (DNA/AA), AO

(AA) and PO (DNA/AA), which included 2,167,861 DNA (682,327 amino acid (AA) sites),

882,417 AA sites, 6,202,646 DNA (1,605,370 AA) sites, respectively. Thus, the largest

alignment that we used to infer the odonate phylogeny consists of 4341 loci concatenated into a

supermatrix with >6 million nucleotide sites. All supermatrices are summarized in Data S2; the

inferred odonate relationships are shown in Figure S1A whereas topologies of all inferred

phylogenies are plotted in Figure S1B and topologies of 1603 CO gene trees are shown in Figure

S1C. Additionally, we performed nodal dating of the inferred phylogeny using 20 fossil

calibration points (Data S3).

The inferred ML phylogenetic tree of Odonata using DNA supermatrix of 1603 BUSCO

loci (Figure 1) was used as a primary phylogenetic hypothesis throughout this study as it agrees

with the majority of relationships inferred by other methods (Figures S1A, S1B). Divergence of

Zygoptera and Epiprocta (Anisozygoptera+Anisoptera) from the Most Recent Common Ancestor

(MRCA) occurred in the Middle Triassic ~242 mya (Figure 1), which is in line with recent

estimates [15, 36]. Comprehensive phylogenetic coestimation of subordinal relationships within

Odonata showed that the suborders were well supported (Figure S1A), as they were consistently

recovered as monophyletic clades in all analyses. In several previous studies, paraphyletic

relationships of Zygoptera had been proposed based on wing vein characters derived from fossil

odonatoids and extant Odonata [37], analysis of 12S [38], analysis of 18S, 28S, Histone 3 (H3)

and morphological data [39] and analysis of 16S and 28S data [40]. In most of these studies,

Lestidae was inferred to be sister to Anisoptera. Functional morphology comparisons of flight

systems, secondary male genitalia and ovipositors also supported a non-monophyletic Zygoptera

with uncertain phylogenetic placement of multiple groups [41]. Nevertheless, the relationships

inferred from these previous datasets seem to be highly unlikely due to apparent morphological

differentiation (e.g., eye spacing, body robustness, wing shape) between the suborders and

support for monophyletic Anisoptera and Zygoptera from more recent morphological [34],

molecular [15, 18, 42, 43] and combined studies using both data types [44]. Our analyses recover

Zygoptera as monophyletic consistently (Figure S1A).

Divergence time estimates suggest a TMRCA of Anisoptera and Anisozygoptera (we

occasionally refer these two suborders as "Epiprocta") in the Late Triassic (~205 mya; Figure 1).

Epiprocta as well as Anisoptera were consistent with more recent studies and recovered as

monophyletic with very high support. We also note here that our divergence time estimates of

Anisoptera tend to be younger than those found in ref. [45]. The fossil-calibration approach

based on penalized likelihood has been shown to overestimate true nodal age [46] preventing

direct comparison between our dates and those estimated in ref. [45].

The phylogenetic position of Gomphidae and Petaluridae, both with respect to each other

and the remaining anisopteran families, has long been difficult to resolve. Several phylogenetic

hypothesis have been proposed in the literature based on molecular and morphological data

regarding the placement of Gomphidae as sister to the remaining Anisoptera [47] or to

Libelluloidea [48]. Petaluridae has exhibited stochastic relationships with different members of

Anisoptera, including sister to Gomphidae [48], sister to Libelluloidea [43], sister to

Chlorogomphidae+Cordulegasteridae [44] and sister to all other Anisoptera [37, 49]. The most

recent analyses of the major anisopteran lineages using several molecular markers [14] suggest

Gomphidae and Petaluridae as a monophyletic group, but without strong support. Here the

majority of our supermatrix analyses (Figure S1A) strongly support a sister relationship between

the two families, and in our phylogeny (Figure 1) they split from the MRCA ~165 mya in the

Middle Jurassic (Figure 1); however, almost all the coalescent-based species trees reject such a

relationship with high confidence (Figure S1A). In the presence of incomplete lineage sorting,

concatenation methods can be statistically inconsistent [50] leading to an erroneous species tree

topology with unreasonably high support [51]. Thus, inconsistency in the recovery of a sister

group relationship between Gomphidae and Petaluridae can be explained by elevated levels of

incomplete lineage sorting between the families and/or possible introgression events [52] (see

below).

New zygopteran lineages originated in the Early Jurassic ~191 mya with the early split of

Lestoidea and the remaining Zygoptera (Figure 1). Subsequent occurrence of two large

zygopteran groups, Calopterygoidea and Coenagrionoidea, was estimated within the Cretaceous

(~145-66 mya) and culminated with the rapid radiation of the majority of extant lineages in the

Paleogene (~66-23 mya). Our calibrated divergences generally agree with estimates in [15].

However, any further comparisons are precluded by the lack of comprehensive divergence time

estimation for Odonata in the literature. The backbone of the crown group Calopterygoidea that

branched off from Coenagrionoidea ~129 mya in the Early Cretaceous was well supported as

monophyletic in most of our inferred phylogenies (Figures 1, S1A). Previous analyses struggled

to provide convincing support for the monophyly of the superfamily [13, 43, 44], whereas only

11 out of 48 phylogenetic reconstructions rejected Calopterygoidea (Figure S1A).

183   We used quartet sampling [53] to provide additional information about nodal support and

investigate biological explanations for alternative evolutionary histories that received some

support. We found that for most odonate key radiations, the majority of quartets (i.e. Frequency

> 0.5) support the proposed phylogenetic hypothesis with Quartet Concordance (QC) scores > 0

(Figure S2) across all estimated putative species trees (Figure S1B). The few exceptions consist

of the Gomphidae+Petaluridae split and the A+B split, where we have Frequency < 0.5 and QC

< 0, which suggests that alternative relationships are possible. Quartet Differential (QD),

inspired by f and D statistics for introgression, provides an indicator of how much the

proportions of discordant quartets are skewed (i.e. whether one of the two discordant

relationships is more common than the other), suggestive of introgression and/or introgression or

substitution rate heterogeneity [53]. Interestingly, we identified skewness (i.e. QD < 0.5) for

almost every major radiation (Figure S2), which suggests that alternative relationships can be a

result of underlying biological processes (e.g. introgression) rather than ILS alone [53]; however,

this score may not be highly informative if the majority of quartets agree with the focal topology

(i.e Frequency > 0.5 and QC > 0). For both the Gomphidae+Petaluridae and the A+B splits, we

have Frequency > 0.5, QC < 0 and QD < 0.5, implying that alternative phylogenetic relationships

are plausibly not only due to ILS but also possible ancestral introgression.

Major Trends in Evolutionary History of Odonata

Investigation of diversification rates in Odonata highlighted two major trends correlated with two

mass extinction events in the Permian-Triassic (P-Tr) ~251 mya and Cretaceous-Paleogene (K-

Pg) ~66 mya. First, it appears likely that the ancient diversity of Odonata was largely eliminated

after the P-Tr mass extinction event (see the temporal distribution of fossil samples in Figure 1)

as was also the case for multiple insect lineages [54]. According to the fossil record at least two

major odonatoid lineages went extinct (Protodonata and Protanisoptera [55]) and likely many

genera from other lineages as well (e.g., Kargalotypidae from Triadophlebimorpha [56]). The

establishment of major odonate lineages was observed during the Cretaceous starting ~135 mya

(Figure 1, red line). This coincided with the radiation of angiosperm plants that, in turn, triggered

the formation of herbivorous insect lineages [36]. Odonates are exclusively carnivorous insects

and their diversification was likely driven by the aforementioned sequence of events.

Interestingly, molecular adaptations in the odonate visual systems are coupled with their

diversification during the Cretaceous as well [18].

Figure 1. Evolutionary History of Odonata

Fossil calibrated ML phylogenetic tree of Odonata using a DNA supermatrix consisting of 1603 BUSCO genes with a total of 2,167,861 aligned sites. The blue densities at each node represent posterior distributions of ages estimated in MCMCTree using 20 fossil calibration points. The red dashed vertical line indicates the beginning of establishment for major odonate lineages originating in and spanning the Cretaceous. The histogram (blue bars) represents temporal distribution of fossil Odonatoptera samples. The black dashed vertical lines mark major extinction events, namely Permian-Triassic (P-Tr, ~ 251 mya), Triassic-Jurassic (Tr-J, ~ 201.3 mya) and Cretaceous-Paleogene (K-Pg, ~66 mya)

Overview of introgression hypotheses tested

Figure 2. Detection of Introgression/Hybridization Trajectories by Different Methods in Anisozygoptera.
Three site pattern-based (D statistic, HyDe and D_{FOIL}), gene tree count/branch length-based (χ^2 test/BLT and QuIBL) and network ML inference (PhyloNet) methods were used to test for introgression/hybridization. Arrows denote introgression. The figure panel represents larval and adult stages for three Odonata suborders. Species from top to bottom: *Lestes australis, Epiophlebia superstes* and *Anax junius*. Image credit: *Epiophlebia superstes* adult by Christian Dutto Engelmann; *Lestes australis* and *Anax junius* adults by John Abbott.

The scope of introgression within Odonata remains largely unknown, where previous studies

looked for its patterns only within certain species relying on inference from a limited number of

genetic loci. Thus, we searched for signatures of introgression using genome-scale datasets

between lineages at several different taxonomic levels: between different suborders, between

superfamilies and within superfamilies. We used six different methods to test for introgression

within Odonata, as exemplified for the Anisozygoptera suborder in Figure 2. Specifically, we

searched for signatures of ancestral inter-superfamilial introgression within the Zygoptera and

Anisoptera suborders. Also, we tested the hypothesis of inter-subordinal gene flow between

Anisozygoptera and Zygoptera. Finally, we tested introgression within superfamilies of

Zygoptera and Anisoptera that included several species (Lestoidea, Calopterygoidea,

Coenagrionoidea, Aeshnoidea (Aeshnidae), Aeshnoidea (Gomphidae+Petaluridae),

Cordulegastroidea and Libelluloidea; Figure 1). The introgression results for the entire Odonata

order either comprise of all tests performed within the entire phylogeny (HyDe) or of a union of

tests performed within Anisoptera, Zygoptera and between Anisozygoptera-Zygoptera (D_{FOIL},

QuiBL and χ^2 count test/Branch Length Test (BLT)).

Site-pattern-based methods strongly suggest multiple instances of introgression within

Odonata

Initially, we tested the above hypotheses of introgression in quartet topologies (Figure S3) within

Odonata using two site-pattern-based methods: the ABBA-BABA test [57] and HyDe [58]. The

ABBA-BABA test and HyDe rely on computation of D and γ statistics, respectively, where their

significant deviation from 0 may indicate the presence of introgression between the tested pair of

taxa. Additionally, estimated γ and $(1-\gamma)$ of HyDe's hybrid speciation model [58] corresponds to

the parental fractions in a putatively hybrid genome. Note that HyDe's hybrid speciation model

is appropriate for detecting introgression with sufficient statistical power and can produce

reasonable estimates of γ [59, 60]. The analysis of the ABBA-BABA test results revealed

Figure 3. Distributions of HyDe γ and Quartets Fractions Supporting Introgression Across Odonate Taxonomic Levels.
(A) Distribution of significant (Bonferroni corrected $P < 10^{-6}$) γ values for each quartet estimated by HyDe. In general, γ values that are not significantly different from 0 denote no relation of a putative hybrid species to either of the parental species P_1 (1-γ) or P_2 (γ) in a quartet. Asterisks indicate significantly greater (red) or lower (blue) γ averages of various tested cases compared to γ average of the entire order.
(B) Proportions of quartets that support introgression based on simultaneous significance of D statistics and γ. Asterisks indicate significantly greater (red) and smaller (blue) fraction of quartets that support introgression compared to the entire order.

possible gene flow events throughout the entire evolutionary history of Odonata (Figure S4 A,

Data S4, SI Results). We also highlight the positive relationship between the values of D and γ

statistics (Spearman's rank correlation test, $\rho = 0.308$, $P = 0$), demonstrating their broad

concordance in identifying signatures of introgression (Figure S4 B).

There are a variety of other test statistics that have been developed to detect introgression

(e.g., [61-64]). Because, like D and γ, many of these statistics are computed from different

invariants, we attempted to visualize the relationships between all 15 site patterns computed by

HyDe using t-distributed stochastic neighbor embedding (tSNE; [65]) for dimensionality

reduction, along with the corresponding values of D and γ (Figures S4 C and D, respectively).

We found that the clustering of quartets with significant introgression according to a two-

dimensional representation of their site patterns may suggest the presence of additional site-

pattern signatures of introgression (Figure S4 C-D).

In order to assess the extent of preservation of ancestrally introgressed genetic material

within contemporary taxa, we compared inferred average values of significant γ statistic from

Odonata with the averages derived from different intra- and inter-superfamilial taxonomic levels

(Figure 3A). We found significantly higher values of γ for several intra- and inter-superfamilial

comparisons including those that involve Anisozygoptera (Wilcoxon rank-sum test [WRST], all

$P < 0.05$, Figure 3A, Data S4, SI Results). Additionally, Anisozygoptera exhibits the largest

average γ (0.27) across all the inter-superfamilial comparisons (Figure 3A, Data S4). We also

found an excess (Fisher exact test [FET], all $P < 0.05$, Data S1) of significant triplets that support

introgression (Figure 3B) based on both the ABBA-BABA and HyDe hybrid speciation model

(Figure S3) tests for Anisozygoptera, Aeshnoidea (Aeshnidae), Lestoidea and Calopterygoidea

(between and within). Additionally, HyDe analysis yielded highly similar results for D and γ

using only the 1st and 2nd codon positions (Figure S5) suggesting that the potential saturation

effect in 3rd codon position had a minor effect on introgression inference.

Further, we tested introgression within Odonata using an alternative invariant-based

method, D_{FOIL} (Figure S6 A, SI Results), which is capable of detecting ancestral as well as inter-

group introgression and inferring its polarization (see Materials and Methods). Specifically, we

observed a highly skewed distribution of D_{FOIL} statistics for Anisozygoptera (Figure S6 B, SI

Results) that may suggest specific directionality of introgression: all 61 (1157 out of 1158

without FDR correction) quintets with positive evidence for ancestral introgression suggest that

Anisozygoptera is the recipient lineage whereas Zygoptera is the donor (Figure 2, red and blue

arrows). Additionally, 1030 out of 1091 (4473 out of 4738 without FDR correction) evaluated

quintets with significant inter-group introgression are indicative of one-way introgression from

Zygoptera lineages to Anisozygoptera as well, whereas the directionality of remaining 59 (254

without FDR correction) quintets support Anisozygoptera as a donor and Zygoptera as a

recipient and only two (11 without FDR correction) show introgression between Zygoptera and

Anisoptera.

Signatures of introgression revealed by phylogenetic gene tree-based discordance methods

Besides specific phylogenetic invariants, introgression will also generate certain patterns of gene

tree-species tree discordance and reduce the genetic divergence between introgressing taxa,

which is reflected in gene tree branch lengths. Thus, the footprints of introgression can be

detected using phylogenetic discordance methods. First, for a triplet of species, under ILS alone

one would expect equal proportions of gene tree topologies supporting the two topologies

disagreeing with the species tree, and any imbalance may suggest introgression (Figure S7).

Thus, deviation from equal frequencies of gene tree counts among discordant gene trees can be

assessed using a χ^2 test, similar to the method proposed in [66, 67]. One would expect the

average distance between putatively introgressing taxa in discordant trees to be significantly

smaller than the distances derived from the concordant as well as alternative discordant triplets

(see Materials and Methods and Figure S7). We used both the χ^2 test of discordant gene tree

counts and a test based on the distribution of branch lengths for concordant and discordant gene

tree triplets (BLT) to identify introgression within Odonata testing different scenarios (Figure

4A). With this combination of methods, we identified a significant fraction of triplets that

support ancestral introgression for scenario 1 that involves inter-subordinal gene flow between

Anisozygoptera and Zygoptera as well as scenarios 2 through 4, which correspond to inter-

superfamilial instances of introgression within Epiprocta (Figure 4B). Within superfamilies we

found signatures of introgression for Calopterygoidea and Libelluloidea (Figure 4B). For

scenario 1 (introgression between Zygoptera and Anisozygoptera), examination of the genetic

divergence distributions (Figure 4C) for concordant and discordant triplets showed that

discordant triplets that may have resulted from gene flow between Anisozygoptera and

Zygoptera (the topology labeled "discord2"), have markedly smaller average divergence between

these two taxa, as expected in the presence of introgression. Similarly, based on the distribution

of mean divergence between putatively introgressing taxa (Figure S8) as well as fraction of

significant triplets (Figure 4B) gene flow is supported for scenarios 2, 3 and 4.

An alternative gene tree branch length-based approach, QuIBL [68], also detected

multiple instances of introgression within the entire order (Figure S9). We note that particularly

for introgression involving Anisozygoptera we observed a larger fraction of triplets suggestive of

introgression than for any other scenario tested.

Figure 4. Results of the χ^2 Count-Branch Length Test (BLT) for Odonate Taxonomic Levels and PhyloNet result for Anisozygoptera.
(A) Scenarios of deep (numbered red arrows) and intra-superfamilial (white triangles) introgression in Odonata tested using χ^2 test and BLT. Red arrows mark the location of ancestral introgression events that were tested between lineages (e.g. for scenario 8, we tested whether contemporary species of Platystictoidea shared introgressed genetic material with either Coenagrionoidea or Calopterygoidea).
(B) Classification of triplets based on the χ^2 test and BLT results. Introgression+ILS cases are those significant according to both the χ^2 test and BLT (FDR corrected $P < 0.05$); all the remaining cases were those where any discordance was inferred to be due to ILS alone.
(C) Normalized genetic divergence between sister taxa (shown in triplets above the panel) averaged across all BUSCO gene trees supporting each topology involving three lineages representing Anisoptera, Anisozygoptera, and Zygoptera.
(D) Phylogenetic network estimated from a set of ML gene trees using maximum likelihood. *Epiophlebia superstes*, the sole representative of Anisozygoptera in our study, was specified to be involved in a reticulation with two other (unspecified) lineages. Blue lines indicate the reticulation event and are labeled with PhyloNet's estimate of γ. The number above the network indicates the log-likelihood score.

Phylogenetic network analyses support Anisozygoptera-Zygoptera introgression

As an alternative approach to identify the lineage experiencing gene flow with Anisozygoptera, we performed phylogenetic network inference in PhyloNet [69, 70]. For this analysis we specified that Anisozygoptera was involved in a reticulation event—the only such event occurring on the tree—and inferred for the other two nodes that were most likely to be involved in the reticulation as well as the value of γ, the fraction of Anisozygoptera's genetic material derived from this reticulation. The full maximum likelihood (Figure 4D) and maximum pseudolikelihood (Figure S10) approaches recovered topologically similar networks with comparable values of γ. Pseudolikelihood analysis suggests a reticulation event between Aeshnidae (a family within Anisoptera) and Zygoptera with 38% of genetic material coming from Zygoptera (Figure S10), whereas full likelihood infers reticulation between Anisoptera and Zygoptera suborders 33% (Figure 4D) of genetic material from Zygoptera. Overall, we note that the full likelihood approach returned networks with higher log-likelihood scores. This observation is most likely due to the fact that we performed full likelihood analysis in an exhaustive manner (Data S6) testing every possible reticulation event within a focal clade topology congruent with the species tree (Figure 1), whereas pseudolikelihood analysis used the hill-climbing algorithm to search the full network space but is not guaranteed to retrieve the most optimal solution. Moreover, differences in objective function within likelihood and pseudolikelihood frameworks could also lead to distinct network topology and γ estimate.

Methodological consensus corroborates strong evidence of introgression

We examined the agreement across site- and gene tree-based methods for detecting introgression (i.e. Hyde/D, and D_{FOIL} and the χ^2 test/BLT) and summarize the results in (Figure S11). Specifically, we compared sets of unique introgressing species pairs that were identified with significant pattern of introgression. Overall, all methods individually were able to identify abundant introgression within the entire Odonata order with strong agreement identified by the significance of overlap using an exact test of multi-set interactions [71]. We found strong overlap across methods in their support of introgression within Epiprocta (scenarios 2 and 3), gene flow involving Anisozygoptera (scenario 1), and intra-superfamilial introgression within

Libelluloidea. Within Zygoptera our methods showed strong overlap in identifying signatures of

introgression for Calopterygoidea only. Notably, our χ^2 test /BLT produced very few predictions

that were not in agreement with Hyde, D and/or D_{FOIL} across different comparisons.

DISCUSSION

Resolving main radiations on Odonata phylogeny

Several competing hypotheses of evolutionary relationships within Odonata have been proposed

by multiple authors regarding various taxonomic levels (Figure S12A) [13, 14, 18, 37, 38, 40, 43,

44, 47, 48, 72, 73]. Our phylogenetic analyses recovered Epiprocta (Anisoptera+Anisozygoptera)

and Zygoptera as monophyletic with high support (Figure S1A) agreeing with other recent

estimates [18, 43, 44, 72]. Our estimated superfamilial relationships within Zygoptera support

hypothesis of Dijkstra, et al. [13] recovering monophyly of Lestoidea, Platystictoidea,

Coenagrionoidea and Calopterygoidea (Figure S12A) with high support (Figure S1A). Inferred

higher-level phylogenetic classification of anisopteran families was highly congruent with [14]

and well-supported (Figure S1A) with the exception of Gomphidae and Petaluridae radiations.

The phylogenetic position of Gomphidae and Petaluridae, both with respect to each other and the

remaining anisopteran families, has long been difficult to resolve (Figure S12C). The most recent

analyses of major anisopteran lineages, using several molecular markers [14], suggest

Gomphidae and Petaluridae as a monophyletic group, but without strong branch support. Here

the majority of the supermatrix analyses strongly support a sister relationship between the two

families (Figure S1A); however, almost all the ASTRAL species tree analyses reject such a

relationship with high confidence (Figure S1A). In the presence of incomplete lineage sorting

concatenation methods can be statistically inconsistent [50] leading to an erroneous species tree

topology with unreasonably high support [51]. Thus, inconsistency in the recovery of a sister

group relationship between Gomphidae and Petaluridae could be a result of elevated levels of

incomplete lineage sorting between the families.

Widespread introgression within Odonata

Introgression among Odonata has been previously thought to be uncommon [74, 75] as a result

of probable reproductive isolation mechanisms such as ethological barriers, phenotypic

divergence (e.g., variable morphology of genitalia [76, 77]) and habitat and temporal isolation.

Most of the introgression and hybridization research in Odonata has been done within the

zygopteran Coenagrionidae family and especially between populations of genus *Ishnura*

focusing on mechanisms of reproductive isolation (e.g. [25, 78]). These studies established

"hybridization thresholds" for these closely related species and showed positive correlation

between isolation and genetic divergence. Furthermore, rapid karyotype evolution can also

contribute to post-mating isolation [79]; however, recent cytogenetic studies across Odonata

phylogeny indicate very stable chromosome number with most prevalent karyotype of $2n = 25$

[80]. Additionally, in recent years there has been a growing body of evidence for introgression

throughout evolutionary histories of various groups. This includes more recent hybridization

events observed in *Heliconius* butterflies [68], cats [81], cichlid fishes [82] as well as deeper

ancestral introgression in primates [83], *Drosophila* [84] and vascular plants [53].

On the shallow evolutionary timescales, the rate of gene flow and maintenance of

introgressed variation can be approximated by the fraction of introgressed material (i.e. by γ in

our case) in a focal genome [85]. However, with the increased divergence time the fixation or

loss of the ancestrally introduced genetic material will primarily depend on the scope of selection

[6] via its adaptive value [7, 8]. Here, our taxon sampling allowed to test ultra-deep ancestral

introgression scenarios where the MRCA of tested taxa can be traced as far back as to the

Triassic period (~251 mya to ~201 mya). We argue that for the cases where we infer that a large

amount of introgressed genetic material was preserved (e.g. >25% in Anisozygoptera) the

ancestral effective migration rates [85] were high (similar conclusions can be made for

Aeshnoidea (Aeshnidae), Calopterygoidea and within Libelluloidea and Lestoidea; Figure 3A);

whereas for the remaining taxa with lower or non-significant deviations of γ (Figure 3A) the

rates of introgression may have been substantially lower or its signatures were purged from the

genome by selection.

We found compelling evidence for patterns of ancestral introgression (Figure S11)

among distantly related odonate taxa. Our conservative analysis of agreement between different

introgression tests shows presence of gene flow within the entire order (Figure S11). This

observation may be explained by reduced sexual selection pressures in the early stages of

odonate evolution that may have inhibited rapid genital divergence [86], which is probably a

primary source of reproductive isolation in Odonata [87].

Ancestral introgression as an evolutionary driver in Anisozygoptera

Perhaps the most compelling evidence that we uncovered was for deep ancestral introgression

between the Zygoptera and Anisozygoptera suborders, which based on the fossil record became

genetically isolated after the Lower Jurassic [88]. Species of Anisozygoptera exhibit anatomical

characteristics of both Anisoptera and Zygoptera suborders. Some general features of

Anisozygoptera that relate them to Zygoptera include dorsal folding of wings during perching in

adults, characteristic anatomy of proventriculus (a part of digestive system that is responsible for

grinding of food particles) and absence of specific muscle groups in the larval rectal gills;

whereas abdominal tergite shape, rear wing geometry and larval structures are similar to

Anisoptera [89]. More recent studies also revealed that Anisozygoptera ovipositor morphology

shares similarity with Zygoptera [90]; muscle composition of the head resemble characteristics

of both Anisoptera and Zygoptera [91]; thoracic musculature of Anisozygoptera larva exhibit

similarity between Anisoptera and Zygoptera [34]. Thus, Anisozygoptera represent a

morphological and behavioral "intermediate" [92], which is supported by our findings where we

consistently recovered introgression between Zygoptera and Anisozygoptera (Figures 2-4).

Moreover, according to the D_{FOIL} method the Anisozygoptera was inferred to be a recipient taxon

from a zygopteran donor, though we cannot rule out a lesser degree of gene flow in the opposite

direction as well. Strikingly, the average HyDe probability parameter $\gamma \sim 0.27$ (Figure 3A)

inferred from multiple quartets is very similar to PhyloNet's inheritance probability $\gamma \sim 0.33$.

This may suggest that a sizeable fraction of the Anisozygoptera genome descends from

zygopteran lineages. Taken together, these observations strongly suggest xenoplasious origin

[93] of Anisozygopteran traits (i.e. traits introduced to a recipient taxon via introgression) that

are shared with Zygoptera. However, we do not reject the possibility that some trait hemiplasy

may have resulted from ILS [94]. Based on the gathered evidence for introgression, we suggest

that ancestral lineages that gave rise to modern day Anisozygoptera and Zygoptera taxa

experienced introgression in their past evolutionary history.

According to the hybrid swarm hypothesis [95], introgression can trigger a rapid

cladogenesis resulting in the establishment of ecologically different species with niche-specific

adaptive phenotypes. Some of the notable examples can include adaptive radiations caused by

ancestral introgression in cichlid fishes [96], and intricate inter-specific introgression patterns in

big cats that could be linked to rapid diversification of modern-day species in that lineage [97].

Interestingly, the large fraction of introgressed material estimated within the Anisozygopteran

lineage does not appear to have facilitated any adaptive radiation. This proposition is supported

by an extremely low species diversification within the suborder (just three extant species) and

also by the lack of fossil record for Anisozygoptera.

Conclusions

Our findings provide the first insight into global patterns of introgression that were

pervasive throughout the evolutionary history of Odonata. Further work based on full genome

sequencing data, including non-coding regions, and denser taxon sampling will be necessary to

more accurately estimate the fraction of the introgressed genome in various species. We also

stress the importance of further method development to allow for more accurate and scalable

inference of phylogenetic networks and enable researchers to pin-point instances of inter-

taxonomic introgression on the phylogeny.

ACKNOWLEDGMENTS

We thank Gavin Martin, Nathan Lord and Camilla Sharkey for the generation of sequence data.

We also thank the DNA Sequencing Center and Fulton Supercomputer Lab (BYU) for

assistance. We also thank Aaron Comeault for his invaluable comments on the manuscript. This

work was supported by the NSF grant DEB-1265714 to SMB and NIH grants R00HG008696

and R35GM138286 to DRS. CS was funded by grant ANR-19-CE45-0012 from French Agence

Nationale de la Recherche.

AUTHOR CONTRIBUTIONS

AS, CS, DRS and SMB conceived the research. AS, CS, MSF and PB analyzed the data. MC,

KAC, MFW and DRS supervised the project. AS, CS, DRS and SMB wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Mallet, J., Besansky, N., and Hahn, M.W. (2016). How reticulated are species? Bioessays *38*, 140-149.
2. Hallstrom, B.M., and Janke, A. (2010). Mammalian evolution may not be strictly bifurcating. Molecular biology and evolution *27*, 2804-2816.
3. Degnan, J.H., and Rosenberg, N.A. (2009). Gene tree discordance, phylogenetic inference and the multispecies coalescent. Trends in ecology & evolution *24*, 332-340.
4. Posada, D., and Crandall, K.A. (2001). Intraspecific gene genealogies: trees grafting into networks. Trends in ecology & evolution *16*, 37-45.
5. Mallet, J. (2005). Hybridization as an invasion of the genome. Trends in ecology & evolution *20*, 229-237.
6. Petr, M., Paabo, S., Kelso, J., and Vernot, B. (2019). Limits of long-term selection against Neandertal introgression. Proc Natl Acad Sci U S A *116*, 1639-1644.
7. Oziolor, E.M., Reid, N.M., Yair, S., Lee, K.M., Guberman VerPloeg, S., Bruns, P.C., Shaw, J.R., Whitehead, A., and Matson, C.W. (2019). Adaptive introgression enables evolutionary rescue from extreme environmental pollution. Science *364*, 455-457.
8. Norris, L.C., Main, B.J., Lee, Y., Collier, T.C., Fofana, A., Cornel, A.J., and Lanzaro, G.C. (2015). Adaptive introgression in an African malaria mosquito coincident with the increased usage of insecticide-treated bed nets. Proc Natl Acad Sci U S A *112*, 815-820.
9. Heliconius Genome, C. (2012). Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. Nature *487*, 94-98.
10. Huerta-Sanchez, E., Jin, X., Asan, Bianba, Z., Peter, B.M., Vinckenbosch, N., Liang, Y., Yi, X., He, M., Somel, M., et al. (2014). Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. Nature *512*, 194-197.
11. Gouy, A., and Excoffier, L. (2020). Polygenic Patterns of Adaptive Introgression in Modern Humans Are Mainly Shaped by Response to Pathogens. Molecular biology and evolution *37*, 1420-1433.
12. Perry, W.L., Lodge, D.M., and Feder, J.L. (2002). Importance of hybridization between indigenous and nonindigenous freshwater species: an overlooked threat to North American biodiversity. Syst Biol *51*, 255-275.
13. Dijkstra, K.D., Kalkman, V.J., Dow, R.A., Stokvis, F.R., and Van Tol, J. (2014). Redefining the damselfly families: a comprehensive molecular phylogeny of Zygoptera (Odonata). Syst Entomol *39*, 68-96.
14. Carle, F.L., Kjer, K.M., and May, M.L. (2015). A molecular phylogeny and classification of Anisoptera (Odonata). Arthropod Syst Phylo *73*, 281-301.
15. Thomas, J.A., Trueman, J.W., Rambaut, A., and Welch, J.J. (2013). Relaxed phylogenetics and the palaeoptera problem: resolving deep ancestral splits in the insect phylogeny. Syst Biol *62*, 285-297.

16. Busse, S., Genet, C., and Hornschemeyer, T. (2013). Homologization of the flight
musculature of zygoptera (insecta: odonata) and neoptera (insecta). PloS one *8*, e55787.
17. Chauhan, P., Hansson, B., Kraaijeveld, K., de Knijff, P., Svensson, E.I., and
Wellenreuther, M. (2014). De novo transcriptome of Ischnura elegans provides insights
into sensory biology, colour and vision genes. BMC Genomics *15*, 808.
18. Suvorov, A., Jensen, N.O., Sharkey, C.R., Fujimoto, M.S., Bodily, P., Wightman, H.M.,
Ogden, T.H., Clement, M.J., and Bybee, S.M. (2017). Opsins have evolved under the
permanent heterozygote model: insights from phylotranscriptomics of Odonata.
Molecular ecology *26*, 1306-1322.
19. Troast, D., Suhling, F., Jinguji, H., Sahlen, G., and Ware, J. (2016). A Global Population
Genetic Study of Pantala flavescens. PloS one *11*, e0148949.
20. Dijkstra, K.D., Monaghan, M.T., and Pauls, S.U. (2014). Freshwater biodiversity and
aquatic insect diversification. Annual review of entomology *59*, 143-163.
21. Cordoba-Aguilar, A. (2008). Dragonflies and damselflies : model organisms for
ecological and evolutionary research, (Oxford ; New York: Oxford University Press).
22. Bybee, S., Córdoba-Aguilar, A., Duryea, M.C., Futahashi, R., Hansson, B., Lorenzo-
Carballa, M.O., Schilder, R., Stoks, R., Suvorov, A., Svensson, E.I., et al. (2016).
Odonata (dragonflies and damselflies) as a bridge between ecology and evolutionary
genomics. Frontiers in zoology *13*, 46.
23. Monetti, L., SÁNchez-GuillÉN, R.A., and Rivera, A.C. (2002). Hybridization between
Ischnura graellsii (Vander Linder) and I. elegans (Rambur) (Odonata: Coenagrionidae):
are they different species? Biological Journal of the Linnean Society *76*, 225-235.
24. SÁNchez-GuillÉN, R.A., Van Gossum, H., and Cordero Rivera, A. (2005). Hybridization
and the inheritance of female colour polymorphism in two ischnurid damselflies
(Odonata: Coenagrionidae). Biological Journal of the Linnean Society *85*, 471-481.
25. Sanchez-Guillen, R.A., Wellenreuther, M., Cordero-Rivera, A., and Hansson, B. (2011).
Introgression and rapid species turnover in sympatric damselflies. BMC evolutionary
biology *11*, 210.
26. Hayashi, F., Dobata, S., and Futahashi, R. (2005). Disturbed population genetics:
suspected introgressive hybridization between two Mnais damselfly species (Odonata).
Zoological science *22*, 869-881.
27. Tynkkynen, K., Grapputo, A., Kotiaho, J.S., Rantala, M.J., Väänänen, S., and Suhonen, J.
(2008). Hybridization in Calopteryx damselflies: the role of males. Animal behaviour *75*,
1431-1439.
28. Solano, E., Hardersen, S., Audisio, P., Amorosi, V., Senczuk, G., and Antonini, G.
(2018). Asymmetric hybridization in Cordulegaster (Odonata: Cordulegastridae):
Secondary postglacial contact and the possible role of mechanical constraints. Ecology
and evolution *8*, 9657-9671.
29. Runemark, A., Vallejo-Marin, M., and Meier, J.I. (2019). Eukaryote hybrid genomes.
PLoS genetics *15*, e1008404.
30. Landry, C.R., Wittkopp, P.J., Taubes, C.H., Ranz, J.M., Clark, A.G., and Hartl, D.L.
(2005). Compensatory cis-trans evolution and the dysregulation of gene expression in
interspecific hybrids of Drosophila. Genetics *171*, 1813-1822.
31. Lemmon, E.M., and Lemmon, A.R. (2010). Reinforcement in chorus frogs: lifetime
fitness estimates including intrinsic natural selection and sexual selection against hybrids.
Evolution; international journal of organic evolution *64*, 1748-1761.

32.   Rothfels, C.J., Johnson, A.K., Hovenkamp, P.H., Swofford, D.L., Roskam, H.C., Fraser-
564   Jenkins, C.R., Windham, M.D., and Pryer, K.M. (2015). Natural hybridization between
565   genera that diverged from each other approximately 60 million years ago. Am Nat *185*,
566   433-442.
567   33.   Kaldy, J., Mozsar, A., Fazekas, G., Farkas, M., Fazekas, D.L., Fazekas, G.L., Goda, K.,
568   Gyongy, Z., Kovacs, B., Semmens, K., et al. (2020). Hybridization of Russian Sturgeon
569   (Acipenser gueldenstaedtii, Brandt and Ratzeberg, 1833) and American Paddlefish
570   (Polyodon spathula, Walbaum 1792) and Evaluation of Their Progeny. Genes (Basel) *11*.
571   34.   Busse, S., Helmker, B., and Hornschemeyer, T. (2015). The thorax morphology of
572   Epiophlebia (Insecta: Odonata) nymphs--including remarks on ontogenesis and
573   evolution. Scientific reports *5*, 12835.
574   35.   Ruppell, G., and Hilfert, D. (1993). The flight of the relict dragonfly Epiophlebia
575   superstes in comparison with that of the modern Odonata. Odonatologica *22*, 295-309.
576   36.   Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen, P.B.,
577   Ware, J., Flouri, T., Beutel, R.G., et al. (2014). Phylogenomics resolves the timing and
578   pattern of insect evolution. Science *346*, 763-767.
579   37.   Trueman, J.W.H. (1996). A preliminary cladistic analysis of odonate wing venation.
580   Odonatologica *25*, 59-72.
581   38.   Saux, C., Simon, C., and Spicer, G.S. (2003). Phylogeny of the dragonfly and damselfly
582   order odonata as inferred by mitochondrial 12S ribosomal RNA sequences. Annals of the
583   Entomological Society of America *96*, 693-699.
584   39.   Ogden, T.H., and Whiting, M.F. (2003). The problem with "the Paleoptera Problem:"
585   sense and sensitivity. Cladistics *19*, 432-442.
586   40.   Hasegawa, E., and Kasuya, E. (2006). Phylogenetic analysis of the insect order Odonata
587   using 28S and 16S rDNA sequences: a comparison between data sets with different
588   evolutionary rates. Entomological Science *9*, 55-66.
589   41.   Pfau, H.K. (1991). Contributions of functional morphology to the phylogenetic
590   systematics of Odonata. Advances in Odonatology *5*, 109-141.
591   42.   Kim, M.J., Jung, K.S., Park, N.S., Wan, X., Kim, K.-G., Jun, J., Yoon, T.J., Bae, Y.J.,
592   Lee, S.M., and Kim, I. (2014). Molecular phylogeny of the higher taxa of Odonata
593   (Insecta) inferred from COI, 16S rRNA, 28S rRNA, and EF1-α sequences. Entomological
594   Research *44*, 65-79.
595   43.   Carle, F.L., Kjer, K.M., and May, M.L. (2008). Evolution of Odonata, with special
596   reference to Coenagrionoidea (Zygoptera). Arthropod Systematics and Phylogeny *66*, 37-
597   44.
598   44.   Bybee, S.M., Ogden, T.H., Branham, M.A., and Whiting, M.F. (2008). Molecules,
599   morphology and fossils: a comprehensive approach to odonate phylogeny and the
600   evolution of the odonate wing. Cladistics *24*, 477-514.
601   45.   Letsch, H., Gottsberger, B., and Ware, J.L. (2016). Not going with the flow: a
602   comprehensive time-calibrated phylogeny of dragonflies (Anisoptera: Odonata: Insecta)
603   provides evidence for the role of lentic habitats on diversification. Molecular ecology *25*,
604   1340-1353.
605   46.   Britton, T., Anderson, C.L., Jacquet, D., Lundqvist, S., and Bremer, K. (2007).
606   Estimating divergence times in large phylogenetic trees. Syst Biol *56*, 741-752.

47.   Blanke, A., Greve, C., Mokso, R., Beckmann, F., and Misof, B. (2013). An updated
phylogeny of Anisoptera including formal convergence analysis of morphological
characters. Syst Entomol *38*, 474-490.
48.   Misof, B., Rickert, A.M., Buckley, T.R., Fleck, G., and Sauer, K.P. (2001). Phylogenetic
signal and its decay in mitochondrial SSU and LSU rRNA gene fragments of Anisoptera.
Molecular biology and evolution *18*, 27-37.
49.   Rehn, A.C. (2003). Phylogenetic analysis of higher-level relationships of Odonata. Syst
Entomol *28*, 181-240.
50.   Roch, S., and Steel, M. (2014). Likelihood-based tree reconstruction on a concatenation
of aligned sequence data sets can be statistically inconsistent. Theoretical population
biology *100C*, 56-62.
51.   Kubatko, L.S., and Degnan, J.H. (2007). Inconsistency of phylogenetic estimates from
concatenated data under coalescence. Syst Biol *56*, 17-24.
52.   Maddison, W.P. (1997). Gene Trees in Species Trees. Syst Biol *46*, 523-536.
53.   Pease, J.B., Brown, J.W., Walker, J.F., Hinchliff, C.E., and Smith, S.A. (2018). Quartet
Sampling distinguishes lack of support from conflicting support in the green plant tree of
life. Am J Bot *105*, 385-403.
54.   Labandeira, C.C., and Sepkoski, J.J., Jr. (1993). Insect diversity in the fossil record.
Science *261*, 310-315.
55.   Grimaldi, D.A., and Engel, M.S. (2005). Evolution of the insects, (Cambridge, UK ; New
York, NY: Cambridge University Press).
56.   Nel, A., Bechly, G., Martinez-Delclos, X., and Fleck, G. (2001). A new family of
Anisoptera from the Upper Jurassic of Karatau in Kazakhstan (Insecta: Odonata:
Juragomphidae n. fam.). Stuttgarter Beitraege zur Naturkunde Serie B (Geologie und
Palaeontologie) *314*, 1-9.
57.   Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T.,
Webster, T., and Reich, D. (2012). Ancient admixture in human history. Genetics *192*,
1065-1093.
58.   Meng, C., and Kubatko, L.S. (2009). Detecting hybrid speciation in the presence of
incomplete lineage sorting using gene tree incongruence: a model. Theor Popul Biol *75*,
35-45.
59.   Blischak, P.D., Chifman, J., Wolfe, A.D., and Kubatko, L.S. (2018). HyDe: A Python
Package for Genome-Scale Hybridization Detection. Syst Biol *67*, 821-829.
60.   Kong, S., and Kubatko, L.S. (2020). Comparative Performance of Popular Methods for
Hybrid Detection using Genomic Data. bioRxiv, 2020.2007.2027.224022.
61.   Durand, E.Y., Patterson, N., Reich, D., and Slatkin, M. (2011). Testing for ancient
admixture between closely related populations. Molecular biology and evolution *28*,
2239-2252.
62.   Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M., Patterson,
N., Li, H., Zhai, W., Fritz, M.H., et al. (2010). A draft sequence of the Neandertal
genome. Science *328*, 710-722.
63.   Martin, S.H., Davey, J.W., and Jiggins, C.D. (2015). Evaluating the use of ABBA-BABA
statistics to locate introgressed loci. Molecular biology and evolution *32*, 244-257.
64.   Kubatko, L.S., and Chifman, J. (2019). An invariants-based method for efficient
identification of hybrid species from large-scale genomic data. BMC evolutionary
biology *19*, 112.

65.   van der Maaten, L., and Hinton, G. (2008). Visualizing Data using t-SNE. Journal of
Machine Learning Research *9*, 2579-2605.

66.   Huson, D.H., Klöpper, T., Lockhart, P.J., and Steel, M.A. (2005). Reconstruction of
Reticulate Networks from Gene Trees. In Research in Computational Molecular Biology,
S. Miyano, J. Mesirov, S. Kasif, S. Istrail, P.A. Pevzner and M. Waterman, eds. (Berlin,
Heidelberg: Springer Berlin Heidelberg), pp. 233-249.

67.   Degnan, J.H., and Rosenberg, N.A. (2009). Gene tree discordance, phylogenetic
inference and the multispecies coalescent. Trends in ecology & evolution *24*, 332-340.

68.   Edelman, N.B., Frandsen, P.B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R.B., Garcia-
Accinelli, G., Van Belleghem, S.M., Patterson, N., Neafsey, D.E., et al. (2019). Genomic
architecture and introgression shape a butterfly radiation. Science *366*, 594-599.

69.   Than, C., Ruths, D., and Nakhleh, L. (2008). PhyloNet: a software package for analyzing
and reconstructing reticulate evolutionary relationships. BMC bioinformatics *9*, 322.

70.   Wen, D., Yu, Y., Zhu, J., and Nakhleh, L. (2018). Inferring Phylogenetic Networks Using
PhyloNet. Syst Biol *67*, 735-740.

71.   Wang, M., Zhao, Y., and Zhang, B. (2015). Efficient Test and Visualization of Multi-Set
Intersections. Scientific reports *5*, 16923.

72.   Dumont, H.J., Vierstraete, A., and Vanfleteren, J.R. (2010). A molecular phylogeny of
the Odonata (Insecta). Syst Entomol *35*, 6-18.

73.   Ware, J., May, M., and Kjer, K. (2007). Phylogeny of the higher Libelluloidea
(Anisoptera: Odonata): an exploration of the most speciose superfamily of dragonflies.
Molecular phylogenetics and evolution *45*, 289-310.

74.   Tennessen, K. (1982). Review of reproductive isolating barriers in Odonata. Advances in
Odonatology *1*, 251-265.

75.   Lowe, C., Harvey, I., Thompson, D., and Watts, P. (2008). Strong genetic divergence
indicates that congeneric damselflies Coenagrion puella and C. pulchellum (Odonata:
Zygoptera: Coenagrionidae) do not hybridise. Hydrobiologia *605*, 55-63.

76.   Hosken, D.J., and Stockley, P. (2004). Sexual selection and genital evolution. Trends in
ecology & evolution *19*, 87-93.

77.   Barnard, A.A., Fincke, O.M., McPeek, M.A., and Masly, J.P. (2017). Mechanical and
tactile incompatibilities cause reproductive isolation between two young damselfly
species. Evolution; international journal of organic evolution *71*, 2410-2427.

78.   Sanchez-Guillen, R.A., Cordoba-Aguilar, A., Cordero-Rivera, A., and Wellenreuther, M.
(2014). Genetic divergence predicts reproductive isolation in damselflies. Journal of
evolutionary biology *27*, 76-87.

79.   Lai, Z., Nakazato, T., Salmaso, M., Burke, J.M., Tang, S., Knapp, S.J., and Rieseberg,
L.H. (2005). Extensive chromosomal repatterning and the evolution of sterility barriers in
hybrid sunflower species. Genetics *171*, 291-303.

80.   Kuznetsova, V.G., and Golub, N.V. (2020). A checklist of chromosome numbers and a
review of karyotype variation in Odonata of the world. Comp Cytogenet *14*, 501-540.

81.   Li, G., Davis, B.W., Eizirik, E., and Murphy, W.J. (2016). Phylogenomic evidence for
ancient hybridization in the genomes of living cats (Felidae). Genome Res *26*, 1-11.

82.   Svardal, H., Quah, F.X., Malinsky, M., Ngatunga, B.P., Miska, E.A., Salzburger, W.,
Genner, M.J., Turner, G.F., and Durbin, R. (2020). Ancestral Hybridization Facilitated
Species Diversification in the Lake Malawi Cichlid Fish Adaptive Radiation. Molecular
biology and evolution *37*, 1100-1113.

83.   Vanderpool, D., Minh, B.Q., Lanfear, R., Hughes, D., Murali, S., Harris, R.A.,
700   Raveendran, M., Muzny, D.M., Hibbins, M.S., Williamson, R.J., et al. (2020). Primate
701   phylogenomics uncovers multiple rapid radiations and ancient interspecific introgression.
702   PLoS Biol *18*, e3000954.
703   84.   Suvorov, A., Kim, B.Y., Wang, J., Armstrong, E.E., Peede, D., D'Agostino, E.R.R.,
704   Price, D.K., Wadell, P., Lang, M., Courtier-Orgogozo, V., et al. (2021). Widespread
705   introgression across a phylogeny of 155 Drosophila genomes.
706   bioRxiv, 2020.2012.2014.422758.
707   85.   Martin, S.H., and Jiggins, C.D. (2017). Interpreting the genomic landscape of
708   introgression. Current opinion in genetics & development *47*, 69-74.
709   86.   Eberhard, W.G. (2004). Rapid divergent evolution of sexual morphology: comparative
710   tests of antagonistic coevolution and traditional female choice. Evolution; international
711   journal of organic evolution *58*, 1947-1970.
712   87.   Cordero Rivera, A., Andres, J.A., Cordoba-Aguilar, A., and Utzeri, C. (2004). Postmating
713   sexual selection: allopatric evolution of sperm competition mechanisms and genital
714   morphology in calopterygid damselflies (Insecta: Odonata). Evolution; international
715   journal of organic evolution *58*, 349-359.
716   88.   Carle, F. (2012). A new Epiophlebia (Odonata: Epiophlebioidea) from China with a
717   review of epiophlebian taxonomy, life history, and biogeography. Arthropod Systematics
718   & Phylogeny *70*, 75–83.
719   89.   Asahina, S., Asahina, S., and Nihon Gakujutsu, S. (1954). A Morphological Study of a
720   Relic Dragonfly Epiophlebia Superstes Selys: (Odonata, Anisozygoptera), (Japan Society
721   for the Promotion of Science).
722   90.   Matushkina, N.A. (2008). The ovipositor of the relic dragonfly Epiophlebia superstes: a
723   morphological re-examination (Odonata: Epiophlebiidae). International Journal of
724   Odonatology *11*, 71-80.
725   91.   Blanke, A., Beckmann, F., and Misof, B. (2013). The head anatomy of Epiophlebia
726   superstes (Odonata: Epiophlebiidae). Organisms Diversity & Evolution *13*, 55-66.
727   92.   Liu, J., Yu, L., Arnold, M.L., Wu, C.H., Wu, S.F., Lu, X., and Zhang, Y.P. (2011).
728   Reticulate evolution: frequent introgressive hybridization among Chinese hares (genus
729   lepus) revealed by analyses of multiple mitochondrial and nuclear DNA loci. BMC
730   evolutionary biology *11*, 223.
731   93.   Wang, Y., Cao, Z., Ogilvie, H.A., and Nakhleh, L. (2020). Phylogenomic assessment of
732   the role of hybridization and introgression in trait evolution. bioRxiv,
733   2020.2009.2016.300343.
734   94.   Guerrero, R.F., and Hahn, M.W. (2018). Quantifying the risk of hemiplasy in
735   phylogenetic inference. Proc Natl Acad Sci U S A *115*, 12787-12792.
736   95.   Seehausen, O. (2004). Hybridization and adaptive radiation. Trends in ecology &
737   evolution *19*, 198-207.
738   96.   Meier, J.I., Marques, D.A., Mwaiko, S., Wagner, C.E., Excoffier, L., and Seehausen, O.
739   (2017). Ancient hybridization fuels rapid cichlid fish adaptive radiations. Nature
740   communications *8*, 14363.
741   97.   Figueiro, H.V., Li, G., Trindade, F.J., Assis, J., Pais, F., Fernandes, G., Santos, S.H.D.,
742   Hughes, G.M., Komissarov, A., Antunes, A., et al. (2017). Genome-wide signatures of
743   complex introgression and adaptive evolution in the big cats. Sci Adv *3*, e1700299.

744   98.   Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I.,
745   Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length
746   transcriptome assembly from RNA-Seq data without a reference genome. Nature
747   biotechnology *29*, 644-652.
748   99.   Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J.,
749   Couger, M.B., Eccles, D., Li, B., Lieber, M., et al. (2013). De novo transcript sequence
750   reconstruction from RNA-seq using the Trinity platform for reference generation and
751   analysis. Nature protocols *8*, 1494-1512.
752   100.   Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment
753   using DIAMOND. Nat Methods *12*, 59-60.
754   101.   Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: accelerated for clustering
755   the next-generation sequencing data. Bioinformatics *28*, 3150-3152.
756   102.   Simao, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., and Zdobnov, E.M.
757   (2015). BUSCO: assessing genome assembly and annotation completeness with single-
758   copy orthologs. Bioinformatics *31*, 3210-3212.
759   103.   Li, L., Stoeckert, C.J., Jr., and Roos, D.S. (2003). OrthoMCL: identification of ortholog
760   groups for eukaryotic genomes. Genome Res *13*, 2178-2189.
761   104.   Yang, Y., and Smith, S.A. (2014). Orthology inference in nonmodel organisms using
762   transcriptomes and low-coverage genomes: improving accuracy and matrix occupancy
763   for phylogenomics. Molecular biology and evolution *31*, 3081-3092.
764   105.   Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLoS computational biology *7*,
765   e1002195.
766   106.   Fujimoto, M.S., Suvorov, A., Jensen, N.O., Clement, M.J., and Bybee, S.M. (2016).
767   Detecting false positive sequence homology: a machine learning approach. BMC
768   bioinformatics *17*, 101.
769   107.   Fujimoto, M.S., Suvorov, A., Jensen, N.O., Clement, M.J., Snell, Q., and Bybee, S.M.
770   (2016). The OGCleaner: filtering false-positive homology clusters. Bioinformatics.
771   108.   Hellmuth, M., Wieseke, N., Lechner, M., Lenhof, H.P., Middendorf, M., and Stadler, P.F.
772   (2015). Phylogenomics with paralogs. Proc Natl Acad Sci U S A *112*, 2058-2063.
773   109.   Mirarab, S., Nguyen, N., Guo, S., Wang, L.S., Kim, J., and Warnow, T. (2015). PASTA:
774   Ultra-Large Multiple Sequence Alignment for Nucleotide and Amino-Acid Sequences.
775   Journal of computational biology : a journal of computational molecular cell biology *22*,
776   377-386.
777   110.   Nguyen, L.T., Schmidt, H.A., von Haeseler, A., and Minh, B.Q. (2015). IQ-TREE: a fast
778   and effective stochastic algorithm for estimating maximum-likelihood phylogenies.
779   Molecular biology and evolution *32*, 268-274.
780   111.   Loytynoja, A., and Goldman, N. (2008). Phylogeny-aware gap placement prevents errors
781   in sequence alignment and evolutionary analysis. Science *320*, 1632-1635.
782   112.   Loytynoja, A. (2014). Phylogeny-aware alignment with PRANK. Methods in molecular
783   biology *1079*, 155-170.
784   113.   Misof, B., and Misof, K. (2009). A Monte Carlo approach successfully identifies
785   randomness in multiple sequence alignments: a more objective means of data exclusion.
786   Syst Biol *58*, 21-34.
787   114.   Mirarab, S., Reaz, R., Bayzid, M.S., Zimmermann, T., Swenson, M.S., and Warnow, T.
788   (2014). ASTRAL: genome-scale coalescent-based species tree estimation. Bioinformatics
789   *30*, i541-548.

115.   Wickett, N.J., Mirarab, S., Nguyen, N., Warnow, T., Carpenter, E., Matasci, N.,
Ayyampalayam, S., Barker, M.S., Burleigh, J.G., Gitzendanner, M.A., et al. (2014).
Phylotranscriptomic analysis of the origin and early diversification of land plants. Proc
Natl Acad Sci U S A *111*, E4859-4868.
116.   Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T., and Calcott, B. (2016).
PartitionFinder 2: New Methods for Selecting Partitioned Models of Evolution for
Molecular and Morphological Phylogenetic Analyses. Molecular biology and evolution.
117.   Lanfear, R., Calcott, B., Kainer, D., Mayer, C., and Stamatakis, A. (2014). Selecting
optimal partitioning schemes for phylogenomic datasets. BMC evolutionary biology *14*,
82.
118.   Minh, B.Q., Nguyen, M.A., and von Haeseler, A. (2013). Ultrafast approximation for
phylogenetic bootstrap. Molecular biology and evolution *30*, 1188-1195.
119.   Gadagkar, S.R., Rosenberg, M.S., and Kumar, S. (2005). Inferring species phylogenies
from multiple genes: concatenated sequence tree versus consensus gene tree. J Exp Zool
B Mol Dev Evol *304*, 64-74.
120.   Aberer, A.J., Kobert, K., and Stamatakis, A. (2014). ExaBayes: massively parallel
bayesian tree inference for the whole-genome era. Molecular biology and evolution *31*,
2553-2556.
121.   Lakner, C., van der Mark, P., Huelsenbeck, J.P., Larget, B., and Ronquist, F. (2008).
Efficiency of Markov chain Monte Carlo tree proposals in Bayesian phylogenetics. Syst
Biol *57*, 86-103.
122.   Brooks, S.P., and Gelman, A. (1998). General Methods for Monitoring Convergence of
Iterative Simulations. Journal of Computational and Graphical Statistics *7*, 434-455.
123.   Lanfear, R., Hua, X., and Warren, D.L. (2016). Estimating the Effective Sample Size of
Tree Topologies from Bayesian Phylogenetic Analyses. Genome biology and evolution *8*,
2319-2332.
124.   Sayyari, E., and Mirarab, S. (2016). Fast Coalescent-Based Computation of Local Branch
Support from Quartet Frequencies. Molecular biology and evolution *33*, 1654-1668.
125.   Yi, H., and Jin, L. (2013). Co-phylog: an assembly-free phylogenomic approach for
closely related organisms. Nucleic acids research *41*, e75.
126.   Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Molecular
biology and evolution *24*, 1586-1591.
127.   dos Reis, M., and Yang, Z. (2011). Approximate likelihood calculation on a phylogeny
for Bayesian estimation of divergence times. Molecular biology and evolution *28*, 2161-
2172.
128.   Puttick, M.N. (2019). MCMCtreeR: functions to prepare MCMCtree analyses and
visualize posterior ages on trees. Bioinformatics *35*, 5321-5322.
129.   Pease, J.B., and Hahn, M.W. (2015). Detection and Polarization of Introgression in a
Five-Taxon Phylogeny. Syst Biol *64*, 651-662.
130.   Hahn, M.W., and Hibbins, M.S. (2019). A Three-Sample Test for Introgression.
Molecular biology and evolution *36*, 2878-2882.
831

MATERIALS AND METHODS

Taxon sampling and RNA-seq

In this study, we used distinct 85 species (83 ingroup and 2 outgroup taxa). 35 RNA-seq libraries were obtained from NCBI (see Data S1). The remaining 58 libraries were sequenced in Bybee Lab (some species have several RNA-seq libraries; Data S1). Total RNA was extracted for each taxon from eye tissue using NucleoSpin columns (Clontech) and reverse-transcribed into cDNA libraries using the Illumina TruSeq RNA v2 sample preparation kit that both generates and amplifies full-length cDNAs. Prepped mRNA libraries with insert size of ~200bp were multiplexed and sequenced on an Illumina HiSeq 2000 producing 101-bp paired-end reads by the Microarray and Genomic Analysis Core Facility at the Huntsman Cancer Institute at the University of Utah, Salt Lake City, UT, USA. Quality scores, tissue type and other information about RNA-seq libraries are summarized in Data S1.

Transcriptome assembly and CDS prediction

RNA-seq libraries were trimmed and de novo assembled using Trinity [98, 99] with default parameters. Then only the longest isoform was selected from each gene for downstream analyses using Trinity utility script. In order to identify potentially coding regions within the transcriptomes, we used TransDecoder with default parameters specifying to predict only the single best ORF. Each predicted proteome was screened for contamination using DIMOND BLASTP [100] with an E-value cutoff of 10^{-10} against custom protein database. Non-arthropod hits were discarded from proteomes (amino acid, AA sequences) and corresponding CDSs. To mitigate redundancy in proteomes and CDSs, we used CD-HIT [101] with the identity threshold of 0.99. Such a conservative threshold was used to prevent exclusion of true paralogous sequences; thus, reducing possible false positive detection of 1:1 orthologs during homology searches.

Homology assessment

In the present study three types of homologous loci (gene clusters) namely conserved single-copy orthologs (CO), all single-copy orthologs (AO) and paralogy-parsed orthologs (PO) identified by BUSCO v1.22 [102], OrthoMCL [103] and Yang's [104] pipelines respectively were used in phylogenetic inference.

BUSCO arthropod Hidden Markov Model Profiles of 2675 single-copy orthologs were

used to find significant COs matches within CDS datasets by HMMER's *hmmersearch* v3 [105]

with group-specific expected bit-score cutoffs. BUSCO classifies loci into complete [duplicated]

and fragmented. Thus, only complete single-copy loci were extracted from CDS datasets and

corresponding AA sequences for further phylogenetic analyses. Since loci were identified as true

orthologs if they score above expected bit-score and complete if their lengths lie within ~95% of

BUSCO group mean length, many partial erroneously assembled sequences were filtered out.

OrthoMCL v2.0.9 [103] was used to compute AOs in all species using predicted AA

sequences by TransDecoder. AA sequences were used in an all-vs-all BLASTP with an E-value

cutoff of 10^{-10} to find putative orthologs and paralogs. The Markov Cluster algorithm (MCL)

inflation point parameter was set to 2. Only 1:1 orthologs were used in further analyses. In order

to exclude false-positive homology clusters identified by OrthoMCL, we applied machine

learning filtering procedure [106] implemented in OGCleaner software v1.0 [107] using a

metaclassifier with logistic regression.

Finally, to identify additional clusters, we used Yang's tree-based orthology inference

pipeline [104] that was specifically designed for non-model organisms using transcriptomic data.

Yang's algorithm is capable of parsing paralogous gene families into "orthology" clusters that

can be used in phylogenetic analyses. It has been shown that paralogous sequences encompass

useful phylogenetic information [108]. First, the Transdecoder-predicted AA sequences were

trimmed using CD-HIT with the identity threshold of 0.995. Then, all-vs-all BLASTP with an E-

value cutoff of 10^{-5} search was implemented. The raw BLASTP output was filtered by hit

fraction of 0.4. Then, MCL clustering was performed with inflation point parameter of 2. Each

cluster was aligned using iterative algorithm of PASTA [109] and then was used to infer a

maximum-likelihood (ML) gene tree using IQ-TREE v1.5.2 [110] with an automatic model

selection. Tree tips that were longer than relative and absolute cutoffs of 0.4 and 1 respectively

were removed. Mono- and paraphyletic tips that belonged to the same species were masked as

well. To increase quality of homology clusters realignment, tree inference and tip masking steps

were iterated with more stringent relative and absolute masking cutoffs of 0.2 and 0.5

respectively. Finally, POs (AA sequences and corresponding CDSs) were extracted by rooted

ingroups (RI) procedure using *Ephemera danica* as an outgroup (for details see [104]).

Cluster alignment, trimming and supermatrix assembly

For most of the analyses only clusters with $\geq$ 42 (~50%) species present were retained. In total,

we obtained five cluster types, namely DNA (CDS) and AA COs, AA AOs and DNA and AA

POs. Each cluster was aligned using PASTA [109] for the DNA and AA alignments and PRANK

v150803 [111, 112] for the codon alignments and alignments with removed either 1st and 2nd or

3rd codon positions. In order to reduce the amount of randomly aligned regions, we implemented

ALISCORE v2.0 [113] trimming procedure (for PASTA alignments) followed by masking any

site with $\geq$ 42 gap characters (for both PASTA and PRANK alignments). Also, sequence

fragments with >50% gap characters were removed from clusters that were subjected to

ASTRAL v4.10.12 [114] estimation since fragmentary data may have a negative effect on

accuracy of gene and hence species tree inference [115]. For each of the cluster type, we

assembled supermatrices from trimmed gene alignments. Additionally, completely untrimmed

supermatrices were generated from DNA and AA COs with $\geq$ 5 species present.

Phylogenetic tree reconstruction

Four contrasting tree building methods (ML:IQTREE, Bayesian:ExaBayes, Supertree:ASTRAL,

Alignment-Free (AF): Co-phylog) were used to infer odonate phylogenetic relationships using

different input data types (untrimmed and trimmed supermatrices, codon supermatrices, codon

supermatrices with 1st and 2nd or 3rd positions removed, gene trees and assembled

transcriptomes). In total we performed 48 phylogenetic analyses and compared topologies to

identify stable and conflicting relationships (Data S2).

We inferred phylogenetic ML trees from each supermatrix using IQTREE implementing

two partitioning schemes: single partition and those identified by PartitionFinder v2.0 (three

GTR models for DNA and a large array of protein models for AA) [116] with relaxed

hierarchical clustering option [117]. In the first case, IQTREE was run allowing model selection

and assessing nodal support with 1000 ultrafast bootstrap (UFBoot) [118] replicates. In the

second case, IQTREE was run with a given PartitionFinder partition model applying gene and

site resampling to minimize false-positives [119] for 1000 UFBoot replicates.

For Bayesian analyses implemented in ExaBayes [120], we used highly trimmed

(retaining sites only with occupancy of $\leq$ 5 gap characters) and original DNA and AA CO

supermatrices assuming a single partition. We initiated 4 independent runs with 4 Markov Chain

Monte Carlo (MCMC) coupled chains sampling every 500^{th} iteration. Due to high computational

demands of the procedure, only the GTR and JTT substitution model priors were applied to DNA

and AA CO supermatrices respectively with the default topology, rate heterogeneity and branch

lengths priors. However, all supported protein substitution models as a prior was specified for the

trimmed AA CO supermatrix. For convergence criteria an average standard deviation of split

frequencies (ASDSF) [121], a potential scale reduction factor (PSRF) [122] and an effective

sample size (ESS) [123] were utilized. Values of $0\% < ASDSF < 1\%$ and $1\% < ASDSF < 5\%$

indicate excellent and acceptable convergence respectively; $ESS > 100$ and $PSRF \sim 1$ represent

good convergence (see ExaBayes manual, [120]).

ASTRAL analyses were conducted using two input types: (i) gene trees obtained by IQ-

TREE allowing model selection for fully trimmed DNA and AA clusters and (ii) gene trees

obtained from the alignment-tree coestimation process in PASTA. Nodal support was assessed

by local posterior probabilities [124]. ASTRAL is a statistically consistent supertree method

under the multispecies coalescent model with better accuracy than other similar approaches

[114].

In addition to standard phylogenetic inferential approaches, we applied an alignment-free

(AF) species tree estimation algorithm using Co-phylog [125]. Raw Transdecoder CDS outputs

were used in this analysis using k-mer size of 9 as the half context length required for Co-phylog.

Bootstrap replicate trees were obtained by running Co-Phylog with the same parameter settings

on each subsampled with replacement CDS Transdecoder libraries and were used to assess nodal

support.

Assessment of phylogenetic support via quartet sampling

As an additional phylogenetic support, we implemented quartet sampling (QS) approach [53].

Briefly, it provides three scores for internal nodes: (i) the quartet concordance (QC) score gives

an estimate of how sampled quartet topologies agree with the putative species tree; (ii) quartet

differential (QD) estimates frequency skewness of the discordant quartet topologies, which can

be indicative of introgression if a skewed frequency is observed and (iii) quartet informativeness

(QI) quantifies how informative sampled quartets are by comparing likelihood scores of

alternative quartet topologies. Finally, QS provides a quartet fidelity (QF) score for terminal

nodes that measures a taxon "rogueness". We performed QS analysis with all 48 putative species

phylogenies using SuperMatrix_50BUSCO_dna_pasta_ali_trim supermatrix, specifying an IQ-

TREE engine for quartet likelihood calculations with 100 replicates (i.e. number of quartet draws

per focal branch).

Fossil dating

A Bayesian algorithm of MCMCTREE [126] with approximate likelihood computation was

implemented to estimate divergence times within Odonata using 20 crown group fossil

constraints with corresponding prior distributions (Data S3). First, we estimated branch length by

ML and then the gradient and Hessian matrix around these ML estimates in MCMCTREE using

SuperMatrix_50BUSCO_dna_pasta_ali_trim supermatrix. Second, we used the gradient and

Hessian matrix which constructs an approximate likelihood function by Taylor expansion [127]

to perform fossil calibration in MCMC framework under the uncorrelated clock model. For this

step we specified GTR+Γ substitution model with 4 gamma categories; birth, death and sampling

parameters of 1, 0.5 and 0.01, respectively. To ensure convergence, the analysis was run

independently 10 times for 10^6 generations, logging every 50th generation and then removing

10% as burn-in. Visualization of the calibrated tree was performed in R using MCMCtreeR

package [128].

Analyses of introgression

In order to address the scope of possible reticulate evolution across odonate phylogeny, we used

various methods of introgression detection such as HyDe/D [59], D_{FOIL} [129], χ^2 goodness-of-fit

test, QuIBL [68] and PhyloNet [69, 70].

The HyDe framework allows detection of hybridization events which relies on

quantification of phylogenetic site patterns (invariants). HyDe estimates whether a putative

hybrid population (or taxon) H is sister to either population P1 with probability γ or to P2 with

probability 1-γ in a 4-taxon (quartet) tree ((P1,H,P2),O), where O denotes an outgroup. Then, it

conducts a formal statistical test of H$_0$: $\gamma = 0$ vs. H$_1$: $\gamma > 0$ using Z-test, where $\gamma = 0$ (=1) is

indicative of non-significant introgression. We applied HyDe to the concatenated supermatrix

SuperMatrix_50BUSCO_dna_pasta_ali_trim of 1603 BUSCO genes under default parameters

specifying *Ephemera danica* as an outgroup. Under this setup HyDe evaluates all possible taxa

quartets. Since HyDe only allows indication of a single outgroup taxon (i.e. *Ephemera danica*),

we excluded all quartets that contained the *Isonychia kiangsinensis* outgroup from the HyDe

output. Additionally, we calculated Patterson's *D* statistic [57] for every quartet from the

frequency (*p*) of ABBA-BABA site patterns estimated by HyDe as $D = \frac{p_{ABBA} - p_{BABA}}{p_{ABBA} + p_{BABA}}$. To test

significance of *D* statistics we used a χ^2 test to assess whether the proportions p_{ABBA} and p_{BABA}

were significantly different. To minimize effect of false positive cases (type I error) in the

output, we first applied a Bonferroni correction to the *P* values derived from *Z*- and χ^2 tests and

then filtered the results based on a significance level of 0.05 and 10^{-6} for *D* and γ, respectively.

Also, we noticed that negative values of *D* were always associated with nonsignificant γ values,

thus all of the *D* values remained positive after filtering was applied. Additionally, we excluded

all quartets that did not match species topology. Further, we ran HyDe on

SuperMatrix_50BUSCO_dna_prank_trim with the excluded 3rd codon position to investigate a

potential impact of the saturation effect on introgression inference.

D_{FOIL} is an alternative site pattern-based approach that detects introgression using

symmetric 5-taxon (quintet) trees, i.e. (((P1,P2),(P3,P4)),O). D_{FOIL} represents a collection of

similar to *D* statistics applied to quintet trees and, if considered simultaneously, they provide a

powerful approach to identify introgression including ancestral as well as donor and recipient

taxa (i.e. introgression directionality). Moreover, D_{FOIL} exhibits exceptionally low false positive

rates [129]. Since the number of possible quintet topologies for a phylogeny of 85 taxa is >

32×10^6, for analysis we extracted them only for every odonate suborder individually with the

custom R scripts. Note, that for Anisozygoptera we only considered quintets that can be formed

between Anisozygoptera, Anisoptera, and Zygoptera taxonomic groups. As the number of

Anisozygoptera quintets is highly disproportional (34619 out of all 72971 tested Odonata

quintets), for downstream analyses we randomly selected 4000 Anisozygoptera quintets which

approximately matches the number of quintets for an individual species. Analogously to HyDe,

we applied D_{FOIL} to the concatenated supermatrix   SuperMatrix_50BUSCO_dna_pasta_ali_trim

of 1603 BUSCO genes under default parameters specifying *Ephemera danica* as an outgroup.

Also, since D_{FOIL} requires that every quintet has a symmetric topology we considered only those

quintets within our phylogeny that met this criterion (Figure 1). Additionally, D_{FOIL} requires that

the divergence time of P3 and P4 precedes divergence of P1 and P2, i.e. $T_2 > T_1$, thus we filtered

out quintets that violated this assumption using divergence times from our fossil calibrated

phylogeny. In order to correct the P values resulted from D_{FOIL} analysis for multiple testing, we

applied the Benjamini-Hochberg procedure at a false discovery rate cutoff (FDR) of 0.05.

1021   As an alternative test for introgression, we performed a simple yet conservative χ^2

goodness-of-fit test on the gene tree count values for each triplet. First, we tested the three count

values to see whether their proportions significantly deviate from 1/3, and if the null hypothesis

is *not* rejected it will be indicative of extreme levels of ILS. Second, we tested the two count

values that originated from discordant gene trees to see whether their proportions significantly

deviate from 1/2, and if the null hypothesis is rejected, it will suggest a presence of introgression.

In order to correct the P values resulted from the χ^2 test for multiple testing, we applied the

Benjamini-Hochberg procedure at a false discovery rate cutoff (FDR) of 0.05. Second, we used a

branch length test (BLT) to identify cases of introgression [84]. This test examines branch

lengths to estimate the age of the most recent coalescence event (measured in substitutions per

site). Introgression should result in more recent coalescences than expected under the concordant

topology with complete lineage sorting, while ILS yields older coalescence events. Importantly,

ILS alone is not expected to result in different coalescence times between the two discordant

topologies, and this forms the null hypothesis for the BLT. For a given triplet, for each gene tree

we calculated the distance d (a proxy for the divergence time between sister taxa) by averaging

the external branch lengths leading to the two sister taxa under that gene tree topology. We

calculated d for each gene tree and denote values of d from the first discordant topology d_{T1} and

those from the second discordant topology d_{T2}. We then compared the distributions of d_{T1} and

d_{T2} using a Wilcoxon Rank Sum Test. Under ILS alone the expectation is that $d_{\text{T1}} = d_{\text{T2}}$, while in

the presence of introgression $d_{\text{T1}} < d_{\text{T2}}$ (suggesting introgression consistent with discordant

topology T_1) or $d_{\text{T1}} > d_{\text{T2}}$ (suggesting introgression with consistent with topology discordant T_2).

The BLT is conceptually similar to the D3 test [130], which transforms the values of d_{T1} and d_{T2}

in a manner similar to the D statistic for detecting introgression. As with the DCT, we performed

the BLT on all triplets within a clade and used a Benjamini-Hochberg correction with a false

discovery rate cutoff (FDR) of 0.05. We note that both the χ^2 test and BLT will be conservative

in cases where there is bidirectional introgression, with the extreme case of equal rates of

introgression in both directions resulting in a complete loss of power.

1048   QuIBL is based on the analysis of branch length distributions across gene trees to infer

putative introgression patterns. Briefly, under coalescent theory internal branches of rooted gene

trees for a set of 3 taxa (triplet) can be viewed as a mixture of two distributions with the

underlying parameters. Each mixture component generates branch lengths corresponding to

either ILS or introgression/speciation. Thus, estimated mixing proportions (π_1 for ILS and π_2 for

introgression/speciation; $\pi_1 + \pi_2 = 1$) of those distribution components show what fraction of the

gene trees were generated through ILS or non-ILS processes. For a given triplet, QuIBL

computes frequency of gene trees that support three alternative topologies. Then for every

alternative topology QuIBL estimates mixing proportions along with other relevant parameters

via Expectation-Maximization and computes Bayesian Information Criterion (BIC) scores for

ILS-only and introgression models. For concordant topologies elevated values of π_2 are expected

whereas for discordant ones π_2 can vary depending on the severity of ILS/intensity of

introgression. In extreme cases when the gene trees were generated exclusively under ILS, π_2

will approach to zero and the expected gene tree frequency for each alternative topology of a

triplet will be approximately 1/3. To identify significant cases of introgression here we used a

stringent cutoff of ΔBIC < -30 [68]. We ran QuIBL on every triplet individually under default

parameters with number of steps (`numsteps` parameter) is equal to 50 and specifying one of

the Ephemeroptera species for triplet rooting. For computational efficiency we extracted triplets

only from the odonate superfamilies in a similar manner as we did for D_{FOIL} (see above). For this

analysis we used 1603 ML gene trees estimated from CO orthology clusters.

1068   In order to estimate phylogenetic networks from the 1603 ML gene trees estimated from

CO orthology clusters, we used pseudolikelihood (InferNetwork_MPL) and likelihood

(CalGTProb) approaches implemented in PhyloNet. For both analyses we only selected gene

trees that had at least one of the outgroup species (*Isonychia kiangsinensis* and *Ephemera*

danica) and at least three ingroup taxa. We performed network searches for our focal clades,

namely Anisozygoptera, Aeshnidae, Gomphidae+Petaluridae and Libellulidae. For

pseudolikelihood analysis, we ran PhyloNet on every clade individually allowing a single

reticulation event, with the starting tree that corresponds to the species phylogeny (`-s` option),

100 iterations (`-x` option), 0.9 bootstrap threshold for gene trees (`-b` option) and optimization of

branch lengths and inheritance probabilities on the inferred networks (`-po` option). To ensure

convergence, the network searches were repeated 3 times. For Anisozygoptera, we explicitly

indicated *Epiophlebia superstes* as a putative hybrid (`-h` option). For the full likelihood

estimation, we fixed the topology of a focal clade (equivalent to the species tree topology) and

calculated likelihood scores for possible networks with a single reticulation (generated with a

custom script) using CalGTProb. Additionally, to assess significance of networks, we used

difference of BIC scores (ΔBIC) derived from network without reticulation (i.e. tree) and a

network with a reticulation (Data S6).

1085

Dimensionality reduction and visualization

To uncover and visualize complex relationships between site pattern frequencies and Patterson's

D statistic and Hyde γ parameter we implemented a dimensionality reduction technique t-

distributed stochastic neighbor embedding (tSNE) [65] under default parameters in R.

Specifically we estimated tSNE maps from counts of 15 quartet site patterns calculated by HyDe

("AAAA" ,"AAAB" ,"AABA" , "AABB" , "AABC" , "ABAA" , "ABAB", "ABAC" , "ABBA"

, "BAAA" ,"ABBC" , "CABC" , "BACA" , "BCAA" , "ABCD").

1093

1094 **QUANTIFICATION AND STATISTICAL ANALYSES**

1095 All statistical analyses were performed in R (https://www.R-project.org/). Statistical significance

1096 was tested using the Wilcoxon Rank Sum test, Fisher Exact test and/or χ^2 test.

1097

1098 **DATA AND SOFTWARE AVAILABILITY**

1099

All raw RNA-seq read libraries generated for this study have been uploaded to the Short Read

Archive (SRA) (SRR12086708-SRR12086765). All assembled transcriptomes, alignments,

inferred gene and species trees, fossil calibrated phylogenies (including paleontological record

assessed from PaleoDB used to plot the fossil distribution in Figure 1), PhyloNet results, QS

results and introgression results are available on figshare

(https://doi.org/10.6084/m9.figshare.12518660). Custom scripts that were used to create various

input files as well as the analysis pipeline are available on GitHub

https://github.com/antonysuv/odo_introgression.