Linkage mapping and QTL analysis of flowering time using ddRAD sequencing with

genotype error correction in *Brassica napus*

Armin Scheben[1a], Anita Severn-Ellis[1], Dhwani Patel[1], Aneeta Pradhan[1], Stephen Rae[2],

Jacqueline Batley[1], David Edwards[1*]

[1]School of Biological Sciences and Institute of Agriculture, The University of Western

Australia, Perth, WA, Australia

[2]BASF Agricultural Solutions Belgium NV, BASF Innovation Center Gent, Technologiepark-

Zwijnaarde 101, 9052 Gent

[a] Current affiliation: Simons Center for Quantitative Biology, Cold Spring Harbor Laboratory,

Cold Spring Harbor, NY, 11724, United States of America

[*] Correspondence to: David Edwards (dave.edwards@uwa.edu.au)

Abstract

Brassica napus is an important oilseed crop cultivated worldwide. During domestication and

breeding of *B. napus*, flowering time has been a target of selection because of its substantial

impact on yield. Here we use double digest restriction-site associated DNA sequencing

(ddRAD) to investigate the genetic basis of flowering in *B. napus*. An F_2 mapping population

was derived from a cross between an early-flowering spring type and a late-flowering winter

type. Flowering time in the mapping population differed by up to 25 days between individuals.

High genotype error rates persisted after initial quality controls, as suggested by a genotype

discordance of ~12% between biological sequencing replicates. After genotype error

correction, a linkage map spanning 3,605.70 cM and compromising 14,630 single nucleotide

polymorphisms (SNPs) was constructed. A quantitative trail locus (QTL) on chromosome C2

was detected in the vicinity of flowering time genes including *FT* and *FLC*. These findings

demonstrate the effectiveness of the ddRAD approach to sample the *B. napus* genome. Our

results also suggest that ddRAD genotype error rates can be higher than expected in F_2

populations. Quality filtering and genotype correction and imputation can substantially reduce

these error rates and allow effective linkage mapping and QTL analysis.

Introduction

Genotyping-by-sequencing (GBS) is a powerful tool for high-throughput discovery of genetic polymorphisms in crops (*1-5*). GBS comprises a range of library preparation and sequencing approaches that differ in their costs, methodical biases and the type and amount of data produced (*1, 6*). Restriction site-associated DNA sequencing (RAD) is a GBS method that can be used to cost-effectively calibrate the number and coverage of genotyped loci and single nucleotide polymorphisms (SNPs) by varying the enzymes used and the sequencing depth. A recent comparative analysis of single enzyme RAD and two enzyme double digest RAD (ddRAD) used a range of enzyme combinations in different plants and suggested that the enzyme combination of HinfI and HpyCH4IV was promising for maximising genome coverage breadth across a range of species (*7*).

GBS has been used for marker discovery, linkage mapping and QTL analysis in a range of crops (*8-10*), including the important oilseed crop *Brassica napus* (*11-13*). Over 20 high density linkage maps have been generated for *B. napus* using RNA sequencing (*14*), the Brassica 60K genotyping array (*15, 16*) and ddRAD sequencing (*12*). Combined with phenotypic data, these linkage maps provide a powerful basis for identification of genes underlying agronomic traits, which can then be introduced into crop germplasm (*17, 18*). Crop yield in *B. napus* is strongly dependent on flowering time, making this trait a key breeding target.

Flowering time genetic pathways have been elucidated in Arabidopsis and most flowering time genes are known to be conserved between Arabidopsis and *B. napus* (*19-21*). Many QTL and associated SNPs for flowering time have been detected in *B. napus* (*22-29*). Despite this progress in understanding the genetic underpinnings of *B. napus* flowering time, a substantial proportion of flowering time variation remains to be explained.

There are three *B. napus* oilseed rape (OSR) growth types with considerable variation in flowering time: spring, semi-winter and winter. Spring OSR and semi-winter OSR have a low requirement for vernalization to flower and are early-flowering, whilst winter OSR has a strong vernalization requirement and is late-flowering. In *B. napus* breeding, the flowering traits of spring OSR decrease its generation time compared to winter OSR, allowing more rapid breeding cycles. Reducing vernalization requirements in winter OSR by introducing spring OSR alleles would be one approach to allow breeders to accelerate winter OSR breeding. In addition, *B. napus* hybrids are generally higher yielding than open pollinated varieties due to heterosis (*30, 31*). If flowering time can be efficiently managed, heterosis could be exploited

from hybridization of spring OSR and winter OSR. Identifying flowering time loci that

distinguish spring OSR and winter OSR therefore has important breeding applications. Here,

to identify these loci, we crossed a spring OSR and a winter OSR to generate an F_2 mapping

population. We genotyped the progeny and parental lines using ddRAD sequencing. Finally,

we constructed a high-density linkage map and carried out QTL analysis of flowering time and

the related trait budding time. We present candidate regions for flowering time and budding

time and discuss the use of error-prone ddRAD genotyping in heterozygous breeding

populations.

68 Results

69 Pre-processing and aligning sequencing reads

70 A single sample ("146") was excluded from further analysis as it had fewer than one million

71 reads after trimming. In the remaining 206 samples, a mean of 13.14 million raw paired

72 sequences were generated per sample. A mean of 56.2% of reads were uniquely aligned with

73 high quality. The mean coverage depth at covered bases was 9.41 × and the mean coverage

74 breadth of the genome was 18.03%.

75 SNP filtering and genotype correction

A total of 4,841,931 biallelic SNPs were identified in the mapping population and parental

individuals. For further analysis, the seven parental individuals were excluded from the SNP

set. Filtering by individual missingness, genotype depth, minor allele frequency (MAF), and

genotype missingness reduced the number of SNPs to 124,804. Of the 199 progeny, 192 were

retained after filtering individuals with high genotype missingness. Of the 124,804 SNPs,

50,856 did not have a heterozygous genotype in any parental individual. The SNPs with

heterozygous genotypes in the parental individuals may be caused by mismapping or remaining

heterozygosity in the parental genomes and were therefore excluded. Next, removing 16,647

SNPs that were monomorphic between parents and 5,957 that showed segregation distortion

(p < 0.01), generated a set of 28,252 SNPs. Segregation distorted SNPs were distributed

relatively evenly across chromosomes, with noticeable hotspots at the ends of chromosomes

A1 and C5 (Figure S1). Genotype-Corrector quality control removed 13,509 further SNPs after

filtering homozygous SNPs located within heterozygous regions. A total of 4.94% of genotypes

were corrected using Genotype-Corrector and 94.76% of missing genotypes were imputed
(Figure 1). The most frequent genotype corrections were B to AB (29.56%) and A to AB
(23.48%).

In the parental replicate individuals, analysis of pairwise genotype concordance identified a
mean genotype discordance of 12.28% (Figure 1). Discordance between homozygous
genotypes (A vs. B) was rare, with conflicts between homozygous and heterozygous genotypes
(A vs. AB, B vs. AB) making up 97.51% of genotype discordance.

Figure 1. A) Mean pairwise genotype concordance between replicate individuals for the parental lines.
The bars for the category 'missing' include all genotype pairs where at least one pair had a missing
genotype call. Genotypes shown are BnSOSR as 'A', BnWOSR as 'B' and heterozygous as 'AB'. B)
Comparison of corrected and uncorrected genotypes. An example of genotype correction and
imputation using 200 SNPs on chromosome A3 is given for six representative individuals denoted as I-
VI (samples 1, 100, 102, 103, 104, and 105). Genotypes are encoded in three colours (A: red; B: green;
AB: blue) and missing markers are shown in white. For each individual, the lower row shows the
genotypes before the correction and imputation step and the upper row shows the genotypes after this
step.

Linkage mapping

107 A linkage map spanning 3,605.70 cM and compromising 14,630 markers was constructed using
108 ASMap with the corrected and imputed markers (Figure 2 and Table S2). The A genome map
109 was 1,771.53 cM with 8,587 markers and the C genome was 1,661.98 cM with 6,043 markers.
110 The highest mean marker density was found on chromosome A10, with 47.90 markers per Mb.

Mean marker density per Mb was higher on the A genome (28.90) than the C genome (12.18). A supplementary map was constructed using uncorrected markers, which showed a high inflation of genetic distances with a total map length over 30,000 cM (Figure S2). Compared to six published genetic maps that were generated with different approaches, the corrected genetic map still showed some indications of inflation (Table S3).

Figure 2. Genetic map based on ddRAD markers. Black lines indicate marker positions and colours along the chromosomes indicate markers density, with extremely dense regions appearing as black due to tightly clustered markers. Figure rendered using LinkageMapView 2.1.2 (*32*).

The correlation between genetic and physical map order provides information about the
consistency between the genetic map and the reference genome. The mean Spearman's rank
correlation for marker order per chromosome was 1.0 (Table 1). Several minor inconsistencies
in marker order were observed (Figure S3). All chromosomes showed mean correlations over
0.98. Mean individual crossover frequency per chromosome was 2.79.

Table 1. Summary of the genetic map. Spearman's rank correlation (rho) was calculated for the genetic
marker positions and the physical marker positions on the reference genomes.

Chr	Length (Mb)	Length (cM)	Number of markers	rho
A1	31.16	242.40	740	0.98
A2	31.34	185.10	630	1.00
A3	39.49	300.82	1,581	1.00
A4	23.31	58.35	748	1.00
A5	28.6	228.92	892	1.00
A6	31.9	34.53	729	1.00
A7	28.9	194.30	954	1.00
A8	21.74	109.17	365	1.00
A9	46.72	223.68	990	1.00
A1	19.96	194.26	958	0.99
C1	47.95	172.18	739	1.00
C2	58.66	195.33	832	0.99
C3	71.85	276.37	900	1.00
C4	61.04	235.26	861	0.99
C5	52.72	219.41	483	1.00
C6	44.61	141.83	504	1.00
C7	52.5	168.96	612	1.00
C8	46.29	195.51	569	1.00
C9	60.21	229.31	543	1.00
Total	798.95	3,605.70	14,630	-

QTL mapping

BnSOSR flowered 20 days earlier (range: 10 to 28) and went to bud 17 days earlier (range: 12 to 20) on average than winter type BnWOSR (Table S4). In the F_2 progeny, flowering times were distributed within the parental range (Figure 3). A single significant ($p<0.05$) overlapping QTL region for the traits flowering time and budding time was detected on chromosome C2 (Figure 3). The physical region of the QTL spanned 20.57 Mb for flowering time and 0.77 Mb for time to bud (Table S5). The flowering time QTL contained 10 flowering time homologs including *FT* (Table S6), and the budding time QTL contained no known flowering time homologs. Carrying the BnWOSR allele at the QTL peak SNP led to an increase in the days to bud and flower (Figure S4). The percentage of phenotypic variance explained for the identified QTL was 9.08% for flowering time and 8.08% for budding time. Suggestive LOD peaks are also noticeable on A2, A3 and C9.

Figure 3. A) A photograph comparing development in parental individuals 126 six days after sowing. B) Histogram of mapping population phenotypes for budding time. C) Histogram of mapping population phenotypes for flowering time. Parental individuals S1-S4 of BnSOSR and W1-W3 of BnWOSR are indicated. Phenotype data is provided in Table S7. D) Genome-wide LOD plots for single QTL mapping of flowering time and budding time. Significant (p<0.05) thresholds based on permutation testing shown with broken lines.

Discussion

Optimising genotyping-by-sequencing strategies

Genotyping with ddRAD was effective at generating a set of 14,630 high-quality SNPs for linkage and QTL mapping. The findings of this study can help calibrate the number of sequencing reads and genomic loci required for a range of research goals in *B. napus* and related species. For GBS, researchers often aim to optimize the genome coverage by controlling the ratio of reads sequenced to the number of loci generated. Here, the expected maximum genome coverage breadth based on in silico digestion with the enzymes HpyCH4IV and HinfI was 24.6%. However, the observed mean genome coverage breadth was lower at 18.03%. An even greater inconsistency was reported when using HpyCH4IV and HinfI in Arabidopsis and *Glycine max*, where the expected genome coverage breadth was 29.4% and 23.1% but the observed experimental values were 4.45% and 3.33% respectively (*7*). The inconsistency has been explained as a product of fragment size selection bias, redundant in silico loci, and insufficient sequencing reads (*7*). In the *B. napus* population used here, the most important factor determining coverage was the amount of sequencing reads available for each sample. Indeed, 24 samples with high sequencing effort showed genome coverage breadth greater than the expected 24.6% up to a maximum value of 36.68%. This is particularly surprising, as coverage was calculated based only on reads aligned with high quality, which is expected to substantially reduce coverage breadth. These findings suggest that, at least in *B. napus*, igCoverage can underestimate the maximum achievable genome coverage breadth.

The high genome coverage breadth achieved using HpyCH4IV and HinfI indicates that these enzymes are well suited for high-density sampling of genome-wide diversity in *B. napus*. However, when sequencing effort is uneven between samples, high genome coverage breadth can increase genotype missingness through allele dropout. If a locus is not sequenced in enough individual samples (here the cut off was 50%) at sufficient depth, it is removed during SNP calling or SNP filtering and becomes a missing genotype call. High levels of missingness are a common characteristic of reduced representation sequencing (*3*) and can limit the usefulness of genotype data in studies where genotype imputation is not possible (*33*). Nevertheless, as shown in this study, by combining imputation with a high-density sampling of the genome, the limitations of genotype missingness in a mapping population can be overcome.

Linkage mapping

The correlation of physical and genetic maps was high, indicating that the map is accurate and collinear with the reference genome. Similarly, collinear maps with only minor inconsistencies were also found by earlier linkage mapping studies in *B. napus* (*12, 34*). The linkage map constructed here was on average 1.9× larger in cM compared to six published *B. napus* linkage maps generated using different approaches with relatively similar marker densities (*12, 14, 16, 34-36*). Although our mapping population is derived from two highly divergent parental lines and may enable us to sample more crossovers than other studies, some residual genotype errors or segregation distortion is expected to lead to some map inflation. In contrast to our study, which relied on an F_2 population, five of the compared studies used recombinant inbred lines or doubled haploid populations, which will likely suffer less genotyping errors due to their inherent lack of heterozygous alleles. Using a genotyping array, which is less error-prone than ddRAD genotyping, in a *B. napus* F_2 population also led to a smaller genetic map of roughly half the size (*16*). This suggests that ddRAD genotyping errors, and not the population type, are the main reason for the genetic map inflation. Nevertheless, the high-density and accuracy of the linkage map presented here suggest that it is useful for localizing QTL.

Flowering time QTL on chromosome C2

Flowering time QTL in *B. napus* are mostly found in regions syntenic with Arabidopsis chromosome 5 (Parkin et al., 2005) on *B. napus* chromosomes A2, A3, A9, A10, C2 and C3. Here, we identified a single significant locus for flowering time and budding time on C2. These two phenotypes were significantly correlated, explaining the shared QTL that explained ~9% of variance. This modest amount of variance explained is typical for traits that are controlled by many loci spread over multiple chromosomes, with each making minor contributions to the phenotypic effect. Additionally, the size of the QTL region (LOD confidence interval) differed for flowering time and budding time. The size of the QTL region is important, as it reflects the level of mapping resolution that was achieved. In F_2 mapping populations, the confidence interval of a QTL can be large (>1Mb), and these populations often represent the starting point for fine-mapping of candidate genes. In this study we used the recommended LOD threshold of 1.5 units for 95% coverage of the confidence interval (*37*). However, the width of the confidence interval depends on how steep the QTL peak is, which can depend on a range of factors including marker density (*38, 39*).

The QTL overlaps with a locus identified in an earlier study, which found that the 60K array SNP Bn-scaff_18507_1-p889927 was associated with a QTL on C2 at position 33,936,984 on v81 (position on Darmor v41: 26,548,393) explaining 6.36% of flowering time variation (*26*). However, the QTL LOD peaks identified here are distant from this location. Among the known flowering time genes on C2, *FT* (*40*), *FLC* (*41*) and *FY* (*42*) may have particularly substantial effects. *FT* has been implicated in *B. napus* flowering time divergence (*28, 43*) and *FLC* has been found to explain ~23% of flowering time variation in *B. napus* (*44*). *FY* is a suppressor of the transcription factor *FLC* (*42, 45*), and lies in close physical proximity to the LOD peaks identified in this study. However, the *FT* homolog on C2 is expressed at low levels in all ecotypes tested (*46*), which suggests it may not be the candidate gene. Further, none of the flowering time gene homologs occur within the budding time QTL region. Based on the QTL results of this study, we therefore cannot conclusively determine whether the C2 homologs and/or regulatory elements of *FY*, *FLC*, *FT* or another gene are underlying the detected QTL. The F_2 experimental design is only the start of the discovery process because lack of recombination between closely linked regions can hinder high-resolution mapping. Our results show one significant QTL and additional suggestive regions of interest. It is likely that several of these loci would need to be transferred into the desired genetic background to exploit heterosis between spring and winter varieties.

Genotype errors and correction

We detected high pairwise genotype discordance within the duplicate parental samples. Because genotype errors in either of the compared duplicate samples can lead to discordance, the genotype error rate can be roughly estimated as half of the discordance (~6%). In line with GBS results in a rice F_2 population (*47*), most errors can be attributed to undercalling of heterozygous alleles (97.51%). The parental lines are homozygous, though residual heterozygosity and mismapping can lead to heterozygous allele calls. The true error rate in the progeny is therefore likely higher than in the parents, because F_2 populations contain an expected 50% heterozygous alleles.

Despite the range of optimal depths reported in the literature, it is clear that calling heterozygous SNPs accurately requires depths substantially higher than those required for calling homozygous SNPs (*48, 49*). The moderate sequencing depths used in this study (9.41×) may thus lead to inflation of sequencing noise and insufficient allele sampling, which can result

in undercalling of heterozygous alleles (*50*). The percentage of errors attributed to undercalling
of heterozygotes may even be an underestimate, as errors between apparently homozygous
alleles may be caused by conflicting erroneous genotype calls of a heterozygous allele.

The genotype error rates found here are higher than error rates reported in the literature, even
for heterozygous populations. For example, Malmberg et al. (2018a) analysed a heterozygous
B. napus panel with different skim sequencing coverages and filtered genotype calls using a
minimum read depth of 5. The authors found error rates of 2.1% error at 2 × sequencing
coverage and 4.2% error at 1 × sequencing coverage (*50*). Similarly, an error rate of 3% was
found using GBS in a bovine population with a minimum read depth of 5 (*51*). In a ddRAD
genotyping study using a mapping population of cichlid fishes, an investigation of genotype
errors found error rates of 4.41% at 8 × coverage (*52*). This suggests that GBS can lead to
higher genotype error rates than expected in samples such as F_2 mapping populations.

A substantial effect of genotype errors on linkage mapping was found. Cumulative inflation of
linkage map length is often caused by genotype errors that introduce spurious double
recombination events into the map (*53*). In addition, high levels of missing data and markers
with segregation distortion may also affect the mapping distance as these alter the calculated
recombination rate (*54*). It has been estimated that every 1% error rate in a marker adds
approximately two cM to the linkage map (*55*). Linkage map inflation has been previously
reported for GBS data in wheat (*53, 54*) and rice (*47*). In one of the studies on wheat, errors
inflated the linkage by up to 11 times (*53*). In a linkage and QTL mapping study of *B. rapa*
based on SNPs derived from GBS, high error rates (19.6%) were found and the resulting A
genome linkage map was inflated, spanning 4,802.52 cM (*56*).

Linkage and QTL mapping in major crops are commonly carried out using highly accurate
commercial genotyping arrays such as the Illumina Infinium Brassica 60K array (*57*).
Genotyping arrays may introduce sampling bias because they only genotype previously known
SNPs. A particular advantage of GBS over genotyping arrays is that regions missing from the
reference genome can be genotyped and used for linkage mapping. However, genotyping arrays
have the important advantages of more accurate calling of heterozygous genotypes and low
missing data. Although genotype arrays also produce errors that frequently involve
heterozygous sites, the error rates are likely lower at 1-2% (*58, 59*). Our results suggest that in
heterozygous populations, genotyping arrays will generate markers with substantially higher
accuracy than GBS. Here, to increase genotype accuracy for GBS data, genotype correction

was applied. We find that genotype correction substantially decreased genetic map inflation,
underlining the value of a correction step in heterozygous populations analysed using GBS at
low to moderate sequencing coverage.

Conclusion

We report a QTL on chromosome C2 for flowering time and budding time in a *B. napus* winter
type *x* spring type cross. This QTL and the additional suggestive loci can be fine-mapped and
backcrossed into the parental varieties to facilitate flowering time control in hybrid spring type
x winter type varieties. An optimised combination of enzymes was also identified using in
silico analysis, and the resulting number of empirical ddRAD loci and SNPs demonstrate the
effectiveness of the enzyme combination HinfI and HpyCH4IV. In addition, we show that
ddRAD generates high levels of genotype errors that can impact linkage map construction. By
filtering SNPs and applying genotype imputation and correction an accurate map could be
constructed, allowing effective QTL analysis of flowering time. Further investigation of the
loci controlling flowering time and maturation will allow *B. napus* breeders to better exploit
variation in winter and spring types.

Materials and methods

Plant material and phenotyping

The mapping population resulted from a cross between an early-flowering spring line (BnSOSR) and a late-flowering winter line (BnWOSR) carried out by BASF. The F_2 population consisting of 200 individuals as well as 4 BnSOSR and 3 BnWOSR parental replicates was sown directly into 10x10x15 cm pots in a phytotron at the University of Western Australia, Perth. Temperature was maintained between 18-22 °C. To ensure flowering occurred, vernalization was initiated 63 days after sowing and plants were moved to a controlled environment chamber with a constant temperature of 4 °C and an 8-hour light period. Plants were watered twice a week by hand. After approximately 6 weeks, and a total of 108 days after sowing, the plants were returned to the phytotron. This 6 week vernalization period is sufficiently long to ensure that variation in flowering time in the F_2 population is not driven by variation in vernalization response (*60, 61*). Each pot was provided with a dripper and connected to the irrigation system. Plants were watered twice a day for 1 minute. A total of 125 ml of fertilizer with micro minerals was provided by hand every two weeks. The time of first floral buds appearing and the date of first flower opening were recorded. All plants were grown until seed set. To investigate whether phenotype data was normally distributed, Shapiro-Wilke's tests were carried and histograms plotted using ggpubr 0.2.1 (*62*).

Restriction enzyme selection and digestion

DNA fragmentation was carried out by simultaneous digestion using two restriction enzymes. Suitable restriction enzyme pairs, that created sticky or overhanging ends, were selected based on their reaction buffer and incubation temperature compatibility to allow simultaneous digestion. The software IgCoverage 1.0 (*7*) was used to carry out in silico digestion of the *B. napus* genome (Darmor-bzh v8.1; *63*) using these enzyme pairs. The number of fragments within the 100-600 bp size range with different ends (LengthDeFrag100-600), as well as the expected percentage of genome coverage breadth generated by the selected 26 restriction enzyme pairs, were then compared. Out of the 26 suitable enzyme pairs selected, 18 pairs showed coverage breadth > 20% (Table S1). The enzyme pair HinfI and HpyCH4IV (New England Biolabs, Ipswich, USA) was selected based on the number of fragments, genome coverage, availability and cost per sample. This pair was predicted to generate 840,663

fragments with different ends within the 100-600 bp range, which covered 24.6% of the

genome. The suitability of the selected restriction enzyme pair was confirmed by digesting 400

ng of genomic DNA using 5 Units of each restriction enzyme and NEB CutSmart™ buffer

(10×) (New England Biolabs (NEB), Ipswich, USA). The reaction was incubated for four hours

at 37°C and the results were visualised using the LabChip GX Touch 24 (PerkinElmer,

Waltham, USA).

Adapter design

Adapters for the ddRAD protocol were designed based on the adapters and indexed primers

used by Peterson et al. (2012). Barcoded adapters were modified to create a complementary

overhang for the HpyCH4IV restriction enzyme, while the common adapter was altered to

create a complementary overhang for the frequent-cutter HinfI. The adapters were assembled

by annealing 10 µM forward and reverse strand oligos as described in Peterson et al. (2012).

The adapter concentrations to be used in the ligation step for the barcoded and common

adapters were determined using the molarity calculator described by Peterson et al. (2012). The

average distance between the restriction sites required for the calculation was calculated using

the estimated in silico digestion results obtained using the IgCoverage package.

ddRAD library preparation

Genomic DNA was extracted from leaf material using the DNeasy Plant Mini Kit (QIAGEN,

Hilden, Germany) according to the manufacturer's protocol. DNA concentrations were

quantified using the broad range Qubit 3.0 Fluorometric assay (Invitrogen, Carlsbad, USA),

while DNA quality was assessed with the LabChip GX Touch 24 (PerkinElmer, Waltham,

USA). Modified versions of the Peterson et al. (2012) and Clark et al. (2014) protocols were

used to construct the ddRAD libraries. The extracted gDNA was normalised at 50 ng/µL and

200 ng of DNA of each sample was digested in a 20 µL reaction volume containing restriction

enzyme/s and recommended buffer. Digestion for the preparation of the ddRAD libraries was

carried out using HpyCH4IV (5 U) and Hinfl (5 U) in NEB CutSmart™ buffer. The reaction

was incubated at 37°C for 4 hours.

The digested DNA was ligated respectively to the unique barcoded and common adapters using

T4 DNA ligase (Thermo Invitrogen, Carlsbad, USA). An 18 µL master mix containing ligation

buffer, 200 U of T4 ligase and the common adapter was prepared and added directly to the 20

µL digest reaction, after which the individual barcoded adapters were added. The reaction was

incubated at 22°C for two hours, followed by 65°C for 20 minutes, then cooled to 4°C at a

ramp rate of 2°C per 90 seconds. To accommodate variation in DNA concentration and quality

the samples were not pooled after ligation but individually purified and double size selected to

enable enrichment of fragments between 250 bp and 800 bp. The total volume of the samples

was adjusted to 100 µL by adding 60 µL of nuclease free water. Double size selection was

carried out by adding 50 µL of a 1:4 (0.5X) mixture of AMPure XP Beads (Beckman Coulter,

Brea, USA) to PEG buffer (20% PEG w/v, 2.5M NaCl) to remove fragments >800 bp. The

supernatant was transferred to 20 µL of a 1:1 (0.7X) Ampure XP bead to PEG buffer mixture

to collect fragments >250 bp. The beads were washed using 80% ethanol and the fragments

eluted in 30 µL nuclease free water.

A 10 µL aliquot of the size selected DNA was used for PCR amplification. A 40 µL master mix

of Phusion Hot-Start High-Fidelity Master Mix Polymerase (Thermo Fisher Scientific,

Walthan, USA) and the Forward (0.5 µM) and Reverse primers (0.5 µM) was prepared.

Samples were amplified at 98°C for 2 minutes, followed by 15 cycles of 98°C for 15 seconds,

62°C for 30 seconds, 72°C for 30 seconds, and a final extension for 5 minutes at 72°C.

Amplified libraries were cleaned using 1.5X Ampure XP Beads to sample volume to remove

primer dimers. The resulting library DNA concentrations were determined using the High

Sensitivity (HS) Qubit 3.0 Fluoro metric assay. Library quality and fragment size distribution

were visualised using the LabChip GX Touch 24. Equimolar amounts of the prepared libraries

were pooled and loaded on a 1.5% agarose gel to enrich and select fragments between 300-700

bp. The DNA was recovered using the QIAquick Gel Extraction Kit (QIAGEN, Hilden,

Germany). The final library concentration, quality and size distribution were assessed again

and adjusted to 20 nM DNA using 10 nM Tris Buffer (pH 8.5, 0.1% Tween 20, 10 nM). The

final libraries were sent to the KCCG Core facility at the Garvan Institute for Medical Research

for paired end sequencing on the HiSeq X Ten platform.

Adapter trimming and quality trimming

The Illumina bcl2fastq 2.20.0.422 pipeline (*64*) was used to convert base call files to FASTQ

format. Paired-end sequencing reads were demultiplexed using sabre 1.0 (*65*) with a single

mismatch allowed. Raw FASTQ files were trimmed of adapter sequences and low quality bases

with Trimmomatic 0.36 (*66*). For adapter trimming, a maximum mismatch score of 2 was used

for the adapter sequence, together with a palindrome clip score threshold of 30 and a simple

clip score threshold of 10. Low quality bases with a Phred+33 score below 3 were trimmed

from the start and end of the read. Sliding window trimming was carried out using a 4-base

wide window, trimming the bases when the average quality per base was below 15. The

Illumina TruSeq3-PE adapter list provided with Trimmomatic was used for adapter trimming.

All reads with fewer than 36 bases after trimming were discarded. Reads that were unpaired

after trimming were also discarded. Following read pre-processing, the untrimmed and

trimmed reads were analysed using the diagnostic tool fastQC (*67*). The results of fastQC for

each sample were then aggregated and summarised using multiQC (*68*). The multiQC report

was used to verify that adapters had been removed and read quality was high.

Aligning sequencing reads

Trimmed reads were mapped using BWA 0.7.17 with the BWA-MEM algorithm (*69*) to the *B.*

napus Darmor-bzh v8.1 assembly (*63*) using default parameters. After alignment, SAM files

were converted to BAM format using SAMtools 1.8 (*70*). Unmapped reads, supplementary

alignments and reads with a mapping quality below 20 were discarded. This filter removes

multi-mapping reads, which commonly occur in *B. napus* due to the homeologous regions of

its polyploid genome. The mapping results were analysed using SAMtools stats and mosdepth

0.2.3 (*71*). The number of ddRAD loci were calculated from mosdepth per-base output using

BEDTools 2.26.0 (*72*) to merge neighboring loci within 100bp.

Calling single nucleotide variants

Variants were called using GATK 3.6 (*73*). First, BAM alignments were indexed using

SAMtools, then HaplotypeCaller was used to call SNPs for each individual sample.

Genotyping was carried out using GATK GenotypeGVCF using default setting with auto index

creation and locking when reading rods disabled. Results per chromosome were merged using

GATK CatVariants. Variants were filtered using VCFtools 0.1.15 (*74*). Indels and multiallelic

SNPs were excluded (*--remove-indels --max-alleles 2 --min-alleles 2*). Before filtering SNPs

individuals with >0.9 missing genotypes were removed. To reduce the rate of heterozygous

alleles incorrectly called as homozygous alleles due to insufficient read depth, genotypes with

a depth <5 (*--minDP 5*) were set to missing. SNPs were discarded if they displayed a minor

allele frequency <0.05 (*--maf 0.05*) or when genotypes were not present in >80% of all

individuals (*--max-missing 0.8*). Genotype discordance was calculated with snpEff 4.3t (*75*)

using the duplicate samples for the parents (spring type BnSOSR with n=4, winter type

BnWOSR with n=3) with pairwise comparisons of genotypes for SNPs passing the above

filters. Heterozygosity per individual was calculated using VCFtools.

The parentage assignment and filtering of distorted SNPs was carried out using the custom

script vcf2gt.py (*76*), which uses cyvcf2 0.8.0 (*77*) to parse VCF files. A chi-square test

implemented in scipy 1.2.0 (*78*) was carried out to identify and discard SNPs with significant

segregation distortion (p<0.01) based on the expected F_2 segregation ratio of 1:2:1. Further

filtering removed SNPs that were heterozygous in at least one of the parents or that were not

polymorphic between the parents. The script also converted the SNPs in VCF format to a

genotype matrix in AB format (A: homozygous allele from Parent 1, B: homozygous allele

from Parent 2; AB: heterozygous allele; -: missing allele).

Genotypes were imputed and corrected using Genotype-Corrector 1.0 (*79*). This software uses

the order of SNPs on the genome reference and a sliding-window approach to impute and

correct genotypes based on neighboring genotypes in F_2 populations. Before correction, up to

eight consecutive homozygous SNPs within 150 bp genomic intervals in heterozygous regions

were binned into a single SNP with Genotype-Corrector qc_hetero. This helps prevent

miscorrection of heterozygous genotypes to homozygous genotypes when using the sliding

window approach. With the 20% missing SNPs used here, the expected accuracy of Genotype-

Corrector is >95%, based on empirical testing in crop mapping populations (*79*).

Linkage mapping

Linkage mapping was carried out using the MSTMap algorithm (*80*) implemented in the R

package ASMap (*81*). The sum of recombination events objective function was used to find the

optimal sequence of loci, and the p-value threshold for clustering markers into linkage groups

was set to $1e^{-23}$ based on evaluation of a range of values from $1e^{-14}$ to $1e^{-29}$. The kosambi

distance function was used to estimate genetic distances between SNPs, and rare recombination

events were treated as errors (*detectBadData=True*). Linkage groups were assigned

chromosome names based on marker positions on the reference genome. Small linkage groups

that did not represent an entire chromosome, were merged with chromosomal linkage groups

and genetic distances recalculated if an unambiguous assignment was possible using physical

marker positions. Linkage groups with <7 markers were discarded. The estimated pairwise

recombination fractions between markers were calculated using the rqtl function plotRF (*37*).
Recombination fractions were used to identify outlier markers that are not in LD with
neighboring markers and to manually correct marker order using physical marker positions. A
total of 95 outlier markers were removed from further analysis. To ascertain the quality of the
genetic map, the correlation between marker order on the genetic map and the reference
genome was calculated using a Spearman's rank correlation test in R. Crossover frequency was
estimated using the rqtl function locateXO and a custom python script crossover.py (*76*).

QTL mapping

QTL mapping was conducted with rqtl scanone using a single QTL model and the non-
parametric model for flowering and budding time because these traits did not follow a normal
distribution. Genome-wide significance thresholds for logarithm of the odds scores (LOD)
were estimated using a permutation test with 1,000 iterations (*82*). The 1.5-LOD drop interval
of each QTL position was estimated using rqtl lodint. The percentage of variance explained for
each QTL was calculated using rqtl fitqtl with Haley-Knott regression. The custom interactive
R script used for QTL mapping in Rstudio 1.1.456 (*83*) was based on the rqtl manual (*37*).

Flowering time genes

A total of 306 Arabidopsis flowering time (FT) genes from the FLOweRing Interactive
Database (*84*) was downloaded from The Arabidopsis Information Resource (*85*). These genes
include homologs of the known *B. napus* flowering time genes. BLAST+ 2.2.29 (*86, 87*)
analysis of the FT genes against the reference genome was carried out to find gene homologs
using a cut-off of $1e^{-6}$ (following *88*). Overlapping hits were merged using BEDtools. The
gene names in the v81 annotation (*63*) were identified using BEDtools to obtain gene
annotations overlapping the BLAST alignments.

Acknowledgements

A.S. was supported by an IPRS awarded by the Australian government. This work was supported by the Australia Research Council (Projects LP160100030, LP140100537 and LP130100925) and BASF. Bayer transferred this work with the CropScience business to BASF in 2018. This work was supported by resources provided by the Pawsey Supercomputing Centre with funding from the Australian Government and the Government of Western Australia.

Author contributions

DE, JB and SR conceived and supervised the project. ASE grew the plants and carried out phenotyping. ASE, AP and DP prepared the sequencing libraries. ASE wrote the phenotyping and library preparation sections of the Materials and Methods. AS carried out the analyses and drafted the manuscript. AS, DE, JB and SR wrote the final manuscript and all authors provided comments and edits.

Data availability

All sequences have been deposited in SRA (PRJNA640838) with individual accessions listed in Table S8.

Code availability

Scripts used to for this study are available at https://github.com/ascheben/bn_gbs/.

References

1. K. R. Andrews, J. M. Good, M. R. Miller, G. Luikart, P. A. Hohenlohe (2016) Harnessing the power of RADseq for ecological and evolutionary genomics. Nature Reviews Genetics 17, 81-92. https://doi.org/10.1038/nrg.2015.28

2. K. Voss-Fels, R. J. Snowdon (2016) Understanding and utilizing crop genome diversity via high-resolution genotyping. Plant Biotechnol. J. 14, 1086-1094. https://doi.org/10.1111/pbi.12456

3. J. A. Poland, T. W. Rife (2012) Genotyping-by-sequencing for plant breeding and genetics. The Plant Genome 5, 92-102. https://doi.org/10.3835/plantgenome2012.05.0005

4. S. Deschamps, V. Llaca, G. D. May (2012) Genotyping-by-sequencing in plants. Biology 1, 460-483. https://doi.org/10.3390/biology1030460

5. A. Scheben, J. Batley, D. Edwards (2017) Genotyping-by-sequencing approaches to characterize crop genomes: choosing the right tool for the right application. Plant Biotech. J. 15, 149-161. https://doi.org/10.1111/pbi.12645

6. X. Xu, G. Bai (2015) Whole-genome resequencing: changing the paradigms of SNP detection, molecular mapping and gene discovery. Mol. Breed. 35, 33-33. https://doi.org/10.1007/s11032-015-0240-6

7. Y. B. Fu, G. W. Peterson, Y. Dong (2016) Increasing genome sampling and improving SNP genotyping for Genotyping-by-sequencing with new combinations of restriction enzymes. G3: Genes|Genomes|Genetics 6, 845. https://doi.org/10.1534/g3.115.025775

8. J. G. Uitdewilligen, A. M. Wolters, B. D'Hoop B, T. J. Borm, R. G. Visser, H. J. van Eck (2013) A next-generation sequencing method for genotyping-by-sequencing of highly heterozygous autotetraploid potato. PLoS One 8, e62355. https://doi.org/10.1371/journal.pone.0062355

9. H. Sonah, L. O'Donoughue, E. Cober, I. Rajcan, F. Belzile (2015) Identification of loci governing eight agronomic traits using a GBS-GWAS approach and validation by QTL mapping in soya bean. Plant Biotechnol. J. 13, 211-221. https://doi.org/10.1111/pbi.12249

10. M. M. Malmberg, L. W. Pembleton, R. C. Baillie, M. C. Drayton, S. Sudheesh, S. Kaur, H. Shinozuka, P. Verma, G. C. Spangenberg, H. D. Daetwyler *et al.* (2018) Genotyping-by-sequencing through transcriptomics: implementation in a range of crop species with varying reproductive habits and ploidy levels. Plant Biotechnol. J. 16, 877-889. https://doi.org/10.1111/pbi.12835

11. S. J. Yang, B. Zhang, G. Liu, B. H. Hong, J. S. Xu, X. Chen, B. Wang, Z. K. Wu, F. Hou, X. P. Yue *et al.* (2018) A comprehensive and precise set of intervarietal substitution lines to identify candidate genes and quantitative trait loci in oilseed rape (*Brassica napus* L.). Theor. Appl. Genet. 131, 2117-2129. https://doi.org/10.1007/s00122-018-3140-4

525   12.   X. Chen, X. Li, B. Zhang, J. Xu, Z. Wu, B. Wang, H. Li, M. Younas, L. Huang, Y. Luo
526   *et al.* (2013) Detection and genotyping of restriction fragment associated
527   polymorphisms in polyploid crops with a pseudo-reference sequence: a case study in
528   allotetraploid *Brassica napus*. BMC Genomics 14, 346-346.
529   https://doi.org/10.1186/1471-2164-14-346

530   13.   Q. H. Zhou, C. Zhou, W. Zheng, A. S. Mason, S. Y. Fan, C. J. Wu, D. H. Fu, Y. J.
531   Huang (2017) Genome-wide SNP markers based on SLAF-seq uncover breeding traces
532   in rapeseed (*Brassica napus* L.). Frontiers in Plant Science 8.
533   https://doi.org/10.3389/fpls.2017.00648

534   14.   I. Bancroft, C. Morgan, F. Fraser, J. Higgins, R. Wells, L. Clissold, D. Baker, Y. Long,
535   J. L. Meng, X. W. Wang *et al.* (2011) Dissecting the genome of the polyploid crop
536   oilseed rape by transcriptome sequencing. Nat. Biotechnol. 29, 762-766.
537   https://doi.org/10.1038/nbt.1926

538   15.   B. Li, J. Gao, J. Chen, Z. Wang, W. Shen, B. Yi, J. Wen, C. Ma, J. Shen, T. Fu *et al.*
539   (2019) Identification and fine mapping of a major locus controlling branching in
540   *Brassica napus*. Theor. Appl. Genet. https://doi.org/10.1007/s00122-019-03506-x

541   16.   F. Sun, J. Liu, W. Hua, X. Sun, X. Wang, H. Wang (2016) Identification of stable QTLs
542   for seed oil content by combined linkage and association mapping in *Brassica napus*.
543   Plant Sci. 252, 388-399. https://doi.org/10.1016/j.plantsci.2016.09.001

544   17.   A. Abe, S. Kosugi, K. Yoshida, S. Natsume, H. Takagi, H. Kanzaki, H. Matsumura, K.
545   Yoshida, C. Mitsuoka, M. Tamiru *et al.* (2012) Genome sequencing reveals
546   agronomically important loci in rice using MutMap. Nat. Biotechnol. 30, 174-178.
547   https://doi.org/10.1038/nbt.2095

548   18.   D. Edwards, J. Batley, R. J. Snowdon (2013) Accessing complex crop genomes with
549   next-generation sequencing. Theor. Appl. Genet. 126, 1-11.
550   https://doi.org/10.1007/s00122-012-1964-x

551   19.   M. Tadege, C. C. Sheldon, C. A. Helliwell, P. Stoutjesdijk, E. S. Dennis, W. J. Peacock
552   (2001) Control of flowering time by *FLC* orthologues in *Brassica napus*. Plant J. 28,
553   545-553. https://doi.org/10.1046/j.1365-313X.2001.01182.x

554   20.   I. A. Parkin, S. M. Gulden, A. G. Sharpe, L. Lukens, M. Trick, T. C. Osborn, D. J.
555   Lydiate (2005) Segmental structure of the *Brassica napus* genome based on
556   comparative analysis with *Arabidopsis thaliana*. Genetics 171, 765-781.
557   https://doi.org/10.1534/genetics.105.042093

558   21.   L. S. Robert, F. Robson, A. Sharpe, D. Lydiate, G. Coupland (1998) Conserved
559   structure and function of the Arabidopsis flowering time gene *CONSTANS* in *Brassica*
560   *napus*. Plant Mol. Biol. 37, 763-772. https://doi.org/10.1023/A:1006064514311

561   22.   H. J. Jian, A. X. Zhang, J. Q. Ma, T. Y. Wang, B. Yang, L. S. Shuang, M. Liu, J. N. Li,
562   X. F. Xu, A. H. Paterson *et al.* (2019) Joint QTL mapping and transcriptome sequencing
563   analysis reveal candidate flowering time genes in *Brassica napus* L. BMC Genomics
564   20, 21. https://doi.org/10.1186/s12864-018-5356-8

565   23.   B. J. Li, W. G. Zhao, D. R. Li, H. B. Chao, X. P. Zhao, N. Ta, Y. H. Li, Z. B. Guan, L.
566   X. Guo, L. N. Zhang *et al.* (2018) Genetic dissection of the mechanism of flowering
567   time based on an environmentally stable and specific QTL in *Brassica napus*. Plant Sci.
568   277, 296-310. https://doi.org/10.1016/j.plantsci.2018.10.005

24. H. Raman, R. Raman, P. Eckermann, N. Coombes, S. Manoli, X. X. Zou, D. Edwards,
J. L. Meng, R. Prangnell, J. Stiller *et al.* (2013) Genetic and physical mapping of
flowering time loci in canola (*Brassica napus* L.). Theor. Appl. Genet. 126, 119-132.
https://doi.org/10.1007/s00122-012-1966-8

25. Y. S. Shen, Y. Xiang, E. S. Xu, X. H. Ge, Z. Y. Li (2018) Major co-localized QTL for
plant height, branch initiation height, stem diameter, and flowering time in an alien
introgression derived *Brassica napus* DH population. Frontiers in Plant Science 9, 390.
https://doi.org/10.3389/fpls.2018.00390

26. L. P. Xu, K. N. Hu, Z. Q. Zhang, C. Y. Guan, S. Chen, W. Hua, J. N. Li, J. Wen, B. Yi,
J. X. Shen *et al.* (2016) Genome-wide association study reveals the genetic architecture
of flowering time in rapeseed (*Brassica napus* L.). DNA Res. 23, 43-52.
https://doi.org/10.1093/dnares/dsv035

27. M. N. Nelson, R. Rajasekaran, A. Smith, S. Chen, C. P. Beeck, K. H. M. Siddique, W.
A. Cowling (2014) Quantitative trait loci for thermal time to flowering and photoperiod
responsiveness discovered in summer annual-type *Brassica napus* L. PLoS One 9,
e102611. https://doi.org/10.1371/journal.pone.0102611

28. H. Raman, R. Raman, Y. Qiu, A. S. Yadav, S. Sureshkumar, L. Borg, M. Rohan, D.
Wheeler, O. Owen, I. Menz *et al.* (2019) GWAS hints at pleiotropic roles for
FLOWERING LOCUS T in flowering time and yield-related traits in canola. BMC
Genomics 20, 636. https://doi.org/10.1186/s12864-019-5964-y

29. T. C. Osborn, C. Kole, I. A. Parkin, A. G. Sharpe, M. Kuiper, D. J. Lydiate, M. Trick
(1997) Comparison of flowering time genes in *Brassica rapa*, *B. napus* and *Arabidopsis
thaliana*. Genetics 146, 1123-1129.

30. K. P. Starmer, J. Brown, J. B. Davis (1998) Heterosis in spring canola hybrids grown
in   northern   Idaho.   Crop   Sci.   38,   376-380.
https://doi.org/10.2135/cropsci1998.0011183X003800020018x

31. H. Cutforth, B. McConkey, S. Brandt, Y. Gan, G. Lafond, S. Angadi, D. Judiesch
(2009) Fertilizer N response and canola yield in the semiarid Canadian prairies. Can J
Plant Sci 89, 501-503. https://doi.org/10.4141/Cjps08128

32. L. A. Ouellette, R. W. Reid, S. G. Blanchard, Jr., C. R. Brouwer (2017)
LinkageMapView - Rendering High Resolution Linkage and QTL Maps.
Bioinformatics 34, 306-307. https://doi.org/10.1093/bioinformatics/btx576

33. H. Alipour, G. H. Bai, G. R. Zhang, M. R. Bihamta, V. Mohammadi, S. A. Peyghambari
(2019) Imputation accuracy of wheat genotyping-by-sequencing (GBS) data using
barley   and   wheat   genome   references.   PLoS   One   14,   e0208614.
https://doi.org/10.1371/journal.pone.0208614

34. X. D. Wang, K. J. Yu, H. G. Li, Q. Peng, F. Chen, W. Zhang, S. Chen, H. L. Maolong,
J. F. Zhang (2015) High-density SNP Map construction and QTL identification for the
apetalous character in *Brassica napus* L. Frontiers in Plant Science 6, 1164.
https://doi.org/10.3389/fpls.2015.01164

35. W. Ecke, A. Kampouridis, K. Ziese-Kubon, A. C. Hirsch (2015) Identification and
genetic characterization by high-throughput SNP analysis of intervarietal substitution
lines of rapeseed (*Brassica napus* L.) with enhanced embryogenic potential. Theor.
Appl. Genet. 128, 587-603. https://doi.org/10.1007/s00122-015-2455-7
614

36. W. E. Clarke, E. E. Higgins, J. Plieske, R. Wieseke, C. Sidebottom, Y. Khedikar, J.
Batley, D. Edwards, J. L. Meng, R. Y. Li *et al.* (2016) A high-density SNP genotyping
array for *Brassica napus* and its ancestral diploid species based on optimised selection
of single-locus markers in the allotetraploid genome. Theor. Appl. Genet. 129, 1887-
1899. https://doi.org/10.1007/s00122-016-2746-7

37. K. W. Broman, H. Wu, S. Sen, G. A. Churchill (2003) R/qtl: QTL mapping in
experimental crosses. Bioinformatics 19, 889-890.
https://doi.org/10.1093/bioinformatics/btg112

38. J. Dupuis, D. Siegmund (1999) Statistical methods for mapping quantitative trait loci
from a dense set of markers. Genetics 151, 373-386.

39. B. Mangin, B. Goffinet, A. Rebai (1994) Constructing confidence intervals for QTL
location. Genetics 138, 1301-1308.

40. F. Turck, F. Fornara, G. Coupland (2008) Regulation and identity of florigen:
FLOWERING LOCUS T moves center stage. Annu. Rev. Plant Biol. 59, 573-594.
https://doi.org/10.1146/annurev.arplant.59.032607.092755

41. S. D. Michaels, R. M. Amasino (1999) *FLOWERING LOCUS C* encodes a novel
MADS domain protein that acts as a repressor of flowering. Plant Cell 11, 949-956.
https://doi.org/10.1105/tpc.11.5.949

42. G. G. Simpson, P. P. Dijkwel, V. Quesada, I. Henderson, C. Dean (2003) *FY* is an RNA
3 ' end-processing factor that interacts with *FCA* to control the *Arabidopsis* floral
transition. Cell 13, 777-787. https://doi.org/10.1016/S0092-8674(03)00425-2

43. D. Z. Wu, Z. Liang, T. Yan, Y. Xu, L. J. Xuan, J. Tang, G. Zhou, U. Lohwasser, S. J.
Hua, H. Y. Wang *et al.* (2019) Whole-genome resequencing of a worldwide collection
of rapeseed accessions reveals the genetic basis of ecotype divergence. Molecular Plant
12, 30-43. https://doi.org/10.1016/j.molp.2018.11.007

44. R. Raman, S. Diffey, J. Carling, R. B. Cowley, A. Kilian, D. J. Luckett, H. Raman
(2016) Quantitative genetic analysis of grain yield in an Australian *Brassica napus*
doubled-haploid population. Crop Pasture Sci 67, 298-307.
https://doi.org/10.1071/Cp15283

45. W. Feng, S. D. Michaels (2011) Dual roles for *FY* in the regulation of *FLC*. Plant
Signaling & Behavior 6, 703-705.

46. J. Wang, C. J. Hopkins, J. N. Hou, X. X. Zou, C. N. Wang, Y. Long, S. Kurup, G. J.
King, J. L. Meng (2012) Promoter variation and transcript divergence in Brassicaceae
lineages of *FLOWERING LOCUS T*. PLoS One 7, e47127.
https://doi.org/10.1371/journal.pone.0047127

47. T. Furuta, M. Ashikari, K. K. Jena, K. Doi, S. Reuscher (2017) Adapting genotyping-
by-sequencing for rice F2 populations. G3-Genes Genom Genet 7, 881-893.
https://doi.org/10.1534/g3.116.038190

48. D. R. Bentley, S. Balasubramanian, H. P. Swerdlow, G. P. Smith, J. Milton, C. G.
Brown, K. P. Hall, D. J. Evers, C. L. Barnes, H. R. Bignell *et al.* (2008) Accurate whole
human genome sequencing using reversible terminator chemistry. Nature 456, 53-59.
https://doi.org/10.1038/nature07517

49. T. Maruki, M. Lynch (2017) Genotype calling from population-genomic sequencing
data. G3-Genes Genom Genet 7, 1393-1404. https://doi.org/10.1534/g3.117.039008

50.   M. M. Malmberg, D. M. Barbulescu, M. C. Drayton, M. Shinozuka, P. Thakur, Y. O.
660   Ogaji, G. C. Spangenberg, H. D. Daetwyler, N. O. I. Cogan (2018) Evaluation and
661   recommendations for routine genotyping using skim whole genome re-sequencing in
662   canola. Frontiers in Plant Science 9, 1809. https://doi.org/10.3389/fpls.2018.01809

663   51.   J. S. Brouard, B. Boyle, E. M. Ibeagha-Awemu, N. Bissonnette (2017) Low-depth
664   genotyping-by-sequencing (GBS) in a bovine population: strategies to maximize the
665   selection of high quality genotypes and the accuracy of imputation. BMC Genet. 18,
666   32. https://doi.org/10.1186/s12863-017-0501-y

667   52.   F. Henning, H. J. Lee, P. Franchini, A. Meyer (2014) Genetic mapping of horizontal
668   stripes in Lake Victoria cichlid fishes: benefits and pitfalls of using RAD markers for
669   dense linkage mapping. Mol. Ecol. 23, 5224-5240. https://doi.org/10.1111/mec.12860

670   53.   P. Bajgain, M. N. Rouse, J. A. Anderson (2016) Comparing genotyping-by-sequencing
671   and single nucleotide polymorphism chip genotyping for quantitative trait loci mapping
672   in wheat. Crop Sci. 56, 232-248. https://doi.org/10.2135/cropsci2015.06.0389

673   54.   W. Hussain, P. S. Baenziger, V. Belamkar, M. J. Guttieri, J. P. Venegas, A. Easterly,
674   A. Sallam, J. Poland (2017) Genotyping-by-sequencing derived high-density linkage
675   map and its application to QTL mapping of flag leaf traits in bread wheat. Sci. Rep. 7,
676   16394. https://doi.org/10.1038/s41598-017-16006-z

677   55.   C. Saintenac, D. Y. Jiang, S. C. Wang, E. Akhunov (2013) Sequence-based mapping
678   of the polyploid wheat genome. G3-Genes Genom Genet 3, 1105-1114.
679   https://doi.org/10.1534/g3.113.005819

680   56.   F. Q. Yu, X. G. Zhang, G. Peng, K. C. Falk, S. E. Strelkov, B. D. Gossen (2017)
681   Genotyping-by-sequencing reveals three QTL for clubroot resistance to six pathotypes
682   of *Plasmodiophora brassicae* in *Brassica rapa*. Sci. Rep. 7, 4516.
683   https://doi.org/10.1038/s41598-017-04903-2

684   57.   A. S. Mason, E. E. Higgins, R. J. Snowdon, J. Batley, A. Stein, C. Werner, I. A. Parkin
685   (2017) A user guide to the *Brassica* 60K Illumina Infinium SNP genotyping array.
686   Theor. Appl. Genet. 130, 621-633. https://doi.org/10.1007/s00122-016-2849-1

687   58.   N. A. Tinker, S. M. Chao, G. R. Lazo, R. E. Oliver, Y. F. Huang, J. A. Poland, E. N.
688   Jellen, P. J. Maughan, A. Kilian, E. W. Jackson (2014) A SNP Genotyping Array for
689   Hexaploid Oat. Plant Genome-Us 7. https://doi.org/10.3835/plantgenome2014.03.0010

690   59.   H. X. Hong, L. Xu, J. Liu, W. D. Jones, Z. Q. Su, B. T. Ning, R. Perkins, W. G. Ge, K.
691   Miclaus, L. Zhang *et al.* (2012) Technical reproducibility of genotyping SNP arrays
692   used in genome-wide association studies. PLoS One 7, e44483.
693   https://doi.org/10.1371/journal.pone.0044483

694   60.   J. C. Richter, C. Mollers (2018) Genetic variation for vernalization requirement of
695   winter oilseed rape. Acta Hortic 1202, 87-91.
696   https://doi.org/10.17660/ActaHortic.2018.1202.13

697   61.   M. Rapacz, A. Markowski (1999) Winter hardiness, frost resistance and vernalization
698   requirement of European winter oilseed rape (Brassica napus var. oleifera) cultivars
699   within the last 20 years. J Agron Crop Sci 183, 243-253. https://doi.org/10.1046/j.1439-
700   037x.1999.00346.x

701   62.   A. Kassambara (2018) 'ggplot2' Based Publication Ready Plots. Accessed: 19 March
702   2018. https://github.com/kassambara/ggpubr

63. P. E. Bayer, B. Hurgobin, A. A. Golicz, C. K. Chan, Y. Yuan, H. Lee, M. Renton, J.
Meng, R. Li, Y. Long *et al.* (2017) Assembly and comparison of two closely related
Brassica napus genomes. Plant Biotechnol. J. 15, 1602-1610.
https://doi.org/10.1111/pbi.12742

64. Illumina Inc. (2017) Accessed: 7 May 2019.
http://sapac.support.illumina.com/downloads/bcl2fastq-conversion-software-v2-
20.html

65. Najoshi (2013) sabre. Accessed: 7 May 2019. https://github.com/najoshi/sabre

66. A. M. Bolger, M. Lohse, B. Usadel (2014) Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics 30, 2114-2120.
https://doi.org/10.1093/bioinformatics/btu170

67. S. R. Andrews (2010) FastQC: a quality control tool for high throughput sequence data.
Accessed: 4 September 2018.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc

68. P. Ewels, M. Magnusson, S. Lundin, M. Kaller (2016) MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinformatics 32, 3047-3048.
https://doi.org/10.1093/bioinformatics/btw354

69. H. Li (2013) Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. arXiv, 1303.3997. https://doi.org/

70. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.
Abecasis, R. Durbin (2009) The Sequence Alignment/Map format and SAMtools.
Bioinformatics 25, 2078-2079. https://doi.org/10.1093/bioinformatics/btp352

71. B. S. Pedersen, A. R. Quinlan (2018) Mosdepth: quick coverage calculation for
genomes and exomes. Bioinformatics 34, 867-868.
https://doi.org/10.1093/bioinformatics/btx699

72. A. R. Quinlan, I. M. Hall (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics 26, 841-842.
https://doi.org/10.1093/bioinformatics/btq033

73. M. A. DePristo, E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire, C. Hartl, A. A.
Philippakis, G. del Angel, M. A. Rivas, M. Hanna *et al.* (2011) A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat.
Genet. 43, 491-498. https://doi.org/10.1038/ng.806

74. P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.
Handsaker, G. Lunter, G. T. Marth, S. T. Sherry *et al.* (2011) The variant call format
and VCFtools. Bioinformatics 27, 2156-2158.
https://doi.org/10.1093/bioinformatics/btr330

75. P. Cingolani, A. Platts, L. L. Wang, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Y.
Lu, D. M. Ruden (2012) A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*
strain w(1118); iso-2; iso-3. Fly 6, 80-92. https://doi.org/10.4161/fly.19695

76. A. Scheben (2020) Accessed: 25 June 2020. https://github.com/ascheben/bn_gbs

77. B. S. Pedersen, A. R. Quinlan (2017) cyvcf2: fast, flexible variant analysis with Python.
Bioinformatics 33, 1867-1869. https://doi.org/10.1093/bioinformatics/btx057

78.   E. Jones, T. Oliphant, P. Peterson (2001) SciPy: Open Source Scientific Tools for
Python. Accessed: 19 January 2019. http://www.scipy.org/

79.   C. Miao, J. Fang, D. Li, P. Liang, X. Zhang, J. Yang, J. C. Schnable, H. Tang (2018)
Genotype-Corrector: improved genotype calls for genetic mapping in F2 and RIL
populations. Sci. Rep. 8, 10088. https://doi.org/10.1038/s41598-018-28294-0

80.   Y. H. Wu, P. R. Bhat, T. J. Close, S. Lonardi (2008) Efficient and accurate construction
of genetic linkage maps from the minimum spanning tree of a graph. PLoS Genet. 4,
e1000212. https://doi.org/10.1371/journal.pgen.1000212

81.   J. Taylor, D. Butler (2017) R package ASMap: efficient genetic linkage map
construction and diagnosis. J Stat Softw 79, 1-29. https://doi.org/10.18637/jss.v079.i06

82.   G. A. Churchill, R. W. Doerge (1994) Empirical threshold values for quantitative trait
mapping. Genetics 138, 963-971.

83.   RStudio Team (2015) RStudio: Integrated Development for R. Accessed: 11 November
2017. http://www.rstudio.com/

84.   F. Bouche, G. Lobet, P. Tocquin, C. Perilleux (2016) FLOR-ID: an interactive database
of flowering-time gene networks in *Arabidopsis thaliana*. Nucleic Acids Res. 44,
D1167-1171. https://doi.org/10.1093/nar/gkv1054

85.   P. Lamesch, T. Z. Berardini, D. H. Li, D. Swarbreck, C. Wilks, R. Sasidharan, R.
Muller, K. Dreher, D. L. Alexander, M. Garcia-Hernandez *et al.* (2012) The
Arabidopsis Information Resource (TAIR): improved gene annotation and new tools.
Nucleic Acids Res. 40, D1202-D1210. https://doi.org/10.1093/nar/gkr1090

86.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman (1990) Basic local
alignment search tool. J. Mol. Biol. 215, 403-410. https://doi.org/10.1016/s0022-
2836(05)80360-2

87.   C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T. L.
Madden (2009) BLAST plus : architecture and applications. BMC Bioinformatics 10,
421. https://doi.org/10.1186/1471-2105-10-421

88.   J. H. Yang, K. Osman, M. Iqbal, D. J. Stekel, Z. W. Luo, S. J. Armstrong, F. C. H.
Franklin (2013) Inferring the *Brassica rapa* interactome using protein-protein
interaction data from *Arabidopsis thaliana*. Frontiers in Plant Science 3, 297.
https://doi.org/10.3389/fpls.2012.00297

Supplemental tables

Supplemental tables

Table S1. In silico double enzyme digest analysis. Frag: Total number of fragments; FragDe:

total number of different end fragments; LengthDeFrag: total length in bases of different end

fragments; DeFrag100-600: total number of different end fragments between 100 bases and

600 bases; LengthDeFrag100-600: total length in bases of different end fragments between 100

bases and 600 bases; % Coverage: percentage of the reference genome covered by fragments

between 100 bases and 600 bases.

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S1.xlsx

Table S2. Linkage map constructed using ASMap with corrected and imputed ddRAD markers

derived from an BnSOSR x BnWOSR F2 population.

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S2.xlsx

Table S3. Mean markers and linkage map sizes across maps generated in this study.

Abbreviations: doubled haploid (DH), recombinant inbred lines (RIL), amplified fragment

length polymorphisms (AFLP), Brassica 60K genotyping array (Brassica 60k).

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S3.xlsx

Table S4. Phenotypic variation in flowering time and time to bud in the F2 mapping population

and the parental lines. Phenotypes of spring type BnSOSR and winter type BnWOSR parental

lines are shown as means of all replicates of each type (n=4, n=3, respectively).

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S4.xlsx

Table S5. QTL identified using genome-wide single QTL scans for budding time (B) and

flowering time (FT). For each QTL, the phenotypic variance explained (PVE) and additive

effect (AE) are shown.

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S5.xlsx

Table S6. Candidate flowering time genes within the QTL interval detected on chromosome

C2. https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S6.xlsx

Table S7. Phenotype data for flowering time and budding time in the mapping population and

parents. https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S7.xlsx

Table S8. Sample list for the 192 progeny and 7 parental individuals used in this study.

https://github.com/ascheben/bn_gbs/blob/master/supplemental_tables/Table_S8.xlsx

Supplemental figures

808
809 Figure S1. Segregation distorted loci across all chromosomes. The significance threshold
810 (p>0.01) is shown as a red dotted line.
811

Figure S2. Comparison of linkage group sizes in an uncorrected genetic map and a corrected

genetic map. Corrected linkage groups are aligned centrally to uncorrected groups.

Figure S3. Physical (x-axis) and genetic (y-axis) marker positions on all chromosomes in Mb

and cM respectively. Spearmans's rank correlation test result shown in the top left corner of

each plot.

Figure S4. Effect plot for the budding time (A) and flowering time (B) QTL on C2 at positions

4,673,904 and 4,655,461 respectively. The 'AA' genotype is BnSOSR and the 'BB' genotype

is BnWOSR.