Full Title: An *in vitro* human-based fracture gap model – Mimicking the crosstalk

between bone and immune cells

Authors: Moritz Pfeiffenberger[1,2], Alexandra Damerau[1,2], Igor Ponomarev[3], Christian H.

Bucher[4,5], Yuling Chen[1,2], Dirk Barnewitz[3], Christa Thöne-Reineke[6], Paula Hoff[1,7], Frank

Buttgereit[1,2,4], Timo Gaber[1,2,4], Annemarie Lang[1,2,4]

Affiliations

[1]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin,

Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department of Rheumatology

and Clinical Immunology, Berlin, Germany

[2]German Rheumatism Research Centre (DRFZ) Berlin, a Leibniz Institute, Berlin, Germany

[3]Research Center of Medical Technology and Biotechnology, Bad Langensalza, Germany

[4]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-

Universität zu Berlin, and Berlin Institute of Health, Berlin Institute of Health Center for

Regenerative Therapies, Berlin, Germany

[5]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-

Universität zu Berlin, and Berlin Institute of Health, Julius Wolff Institute, Berlin, Germany

[6]Institute of Animal Welfare, Animal Behavior and Laboratory Animal Science, Department of

Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

[7]Endokrinologikum Berlin, MVZ am Gendarmenmarkt, Berlin, Germany

Correspondence

Dr. med. vet. Annemarie Lang, PhD

Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-

Universität zu Berlin, and Berlin Institute of Health

Department of Rheumatology and Clinical Immunology

Charitéplatz 1, 10117 Berlin

annemarie.lang@charite.de

Abstract

The interaction between the bone and immune cells plays a crucial role in bone pathologies such as disturbed fracture healing. After a trauma, the initially formed fracture hematoma in the fracture gap contains all important components (immune/stem cells, mediators) to directly induce bone regeneration and is therefore of great importance but most susceptible to negative influences. Thus, reliable *in vitro* models are needed to study the underlying mechanisms and to predict the efficiency of novel therapeutic approaches. Since common bioengineering approaches exclude the immune component, we introduce an *in vitro* 3D fracture gap model which combines scaffold-free bone-like constructs with a fracture hematoma model consisting of human peripheral blood (immune cells) and bone marrow-derived mesenchymal stromal cells. Our *in vitro* 3D fracture gap model provides all osteogenic cues to induce the initial bone healing processes, which were further promoted by applying the osteoinductive deferoxamine (DFO). Thus, we were able to distinctly mimic processes of the initial fracture phase and demonstrated the importance of including the crosstalk between bone and immune cells.

Key words: bone; fracture; fracture hematoma; immune cells; *in vitro* model

Introduction

Bone is an essential part of the musculoskeletal system shaping the body and enabling locomotion and stability. Bone pathologies such as osteoporosis or disturbed fracture healing lead to pain, immobility, inflexibility, considerable loss of life quality and even mental illnesses[1]. Traumatic events can result in bone fracturing accompanied by vessel ruptures and the opening of the bone marrow channel. The pivotal event in the initial phase of fracture healing is the formation of the fracture hematoma, which mainly consists of immune and progenitor cells[2,3]. Negative influences on the initial phase by medications or comorbidities such as diabetes, rheumatoid arthritis or immunosuppression can lead to disturbed fracture healing occurring in approximately 10% of patients with fractures[4,5]. Recent treatment strategies have achieved high technology standards with regard to fixation systems such as plates or implants, regenerative approaches using autologous bone graft (gold-standard) or the additional application of stem cells and/or growth factors[6]. Therefore, preclinical studies are highly needed to tackle the unmet clinical need, especially with respect to an aging population and the increase of comorbidities.

The surgical removal of the fracture hematoma results in a prolonged healing process, while transplantation in an ectopic location elevates bone formation[7]. Formation of the fracture hematoma and the constricted interplay of pro- and anti-inflammatory processes are considered as the starting point of bone regeneration[8-10]. Since the bone marrow cavity is opened during the fracture, the bone marrow acts as a resource for chondro- and osteo-progenitor cells such as mesenchymal stromal cells (MSCs). Therefore, it can be hypothesized that osteogenic induction within the fracture gap is directly induced and controlled by signals from bone components in the vicinity of the fracture gap[11,12]. Hence, the crosstalk between immune cells from peripheral blood (after vessel rupture) and the bone marrow, and osteo-progenitor cells is essential and needs to be considered in preclinical studies[13]. The today's gold-standard of preclinical drug, compound screening and risk assessment is the use of animal models - mainly rodents (mice and rats) - which is in accordance with most national legal requirements. Nevertheless, trans-species differences may be responsible for the limited transferability of findings to the human patient[14,15]. Mimicking the *in-patient* situation in preclinical studies is highly encouraged and evading cross-species differences by novel *in vitro* approaches is of great interest. During the past decades, conventional *in vitro* cell culture systems have been revised and improved to provide more physiological and human-relevant features. This development was mainly driven by the triumph of regenerative medicine relying on improvements in tissue engineering to produce e.g. huge batches of primary cells, 3D nature-resembling artificial tissues or biocompatible biomaterials. Furthermore, the rapid technical evolution allocating sophisticated biomaterials, bioreactors and microfluidic platforms

allows the development of innovative human-relevant *in vitro* systems as alternative or
predictive support to animal testing[16].

Current *in vitro* systems focus on mimicking bone development, endochondral
ossification or the bone homoeostasis itself by using spheroids, scaffold-based or scaffold-free
model systems. Common cell sources are either primary bone-related cells such as
osteoblasts, osteocytes, osteoclasts or MSCs as progenitor cells[16,17]. To mimic fracture
healing, models mainly focus on later stages of the regeneration processes particularly
endochondral ossification or remodeling. We have previously described the development of a
fracture hematoma model consisting of human whole blood and a certain amount of human
(h)MSCs[18]. However, to study the initiated processes in an interconnected manner and more
adequate experimental setting, the combination of the bone components with the fracture
hematoma (immune component) remains elusive.

Within our study, we have developed and characterized scaffold-free bone-like
constructs (SFBCs) based on mesenchymal condensation which exceeded the dimensions of
spheroids. These SFBCs were co-cultivated with *in vitro* fracture hematoma (FH) models to (i)
confirm the capability of the SFBCs to act as an osteogenic inducer, (ii) to mimic the initial
phase of fracture healing with adequate culture conditions and (iii) to use the system as a
platform to test potential therapeutics (DFO - deferoxamine).

Results

SFBCs are characterized by permeating mineralization

MSCs are well-known for their pronounced osteogenic capacity, especially when cultivated *in*
vitro[17,19]. However, in a first step, we wanted to know if it is possible to employ mesenchymal
condensation as a macroscale approach with consistent 3D self-organization and permeating
mineralization. Thus, SFBCs were generated by hMSC condensation and treatment with
osteogenic medium until analysis at week 12 (**Fig 1a**). Macroscopic observation indicated
comparable generation of SFBCs from different hMSC donors with a diameter of approx. 1 cm
and a thickness of 0.5 cm (**Fig. 1b**). To verify the mineralization, *in vitro* computed micro-
tomography (µCT) was performed, showing a consistently high mineralization in the outer area
which was slightly reduced towards the center (**Fig. 1c**). 3D reconstruction yielded the
presence of mineralized tissue as indicated by parameters such as bone volume (BV; mean =
5.1 ± 3.7 mm^3) and bone surface (BS; mean = 276.1 ± 195.4 mm^2) (**Fig. 1d**). To quantify the
connectedness of the mineralized areas, we additionally examined the trabecular pattern factor
(TBPf). This parameter was originally invented to evaluate trabecular bone. Although no
osteoclasts were present in the SFBCs, thus the formation of clear trabeculae was not
expected, we use this parameter to distinguish between concave (= connected) and convex (=
isolated) structures. Low or even negative values represent hereby high connected tissue

which was found in at least 4 (TBPf < 2) out of 9 SFBCs (**Fig. 1d**). To evaluate the structural
morphology of the SFBCs in further detail, we used scanning electron microscopy (SEM) and
found similar morphology when compared to human native bone (**Fig. 1e, f**). In detail, the top
view shows strong matrix formation and a closed superficial layer with certain, isolated crystal-
like depositions while the cut face revealed a layer-like structure.

To get a more detailed overview on the matrix composition, assorted histological and
immunohistochemical stainings were applied. Cells were homogenously distributed within the
SFBCs as indicated by hematoxylin and eosin (H&E) staining while Alizarin Red S and von
Kossa staining confirmed the deposition of calcium and phosphate, respectively throughout
the tissue (**Fig. 2a**). Interestingly, SFBCs which showed a higher amount of negative stained
area, showed a more pronounced layer-like structure while the layers itself were strongly
mineralized at the borders (**Fig. 2a**; lower row). Negative controls were cultivated without
osteogenic medium (**Fig. 2b**). In addition, we observed the expression of *collagen type I* (Col
I) and *alkaline phosphatase* (ALP), which are typical markers for osteogenic processes, while
no Col II, a typical marker of chondrogenesis, was found (**Fig. 2c**).

Figure 1: Characterization of the SFBCs with respect to mineralization and structure. (**a**) Study design; (**b**)
Exemplary images of SFBCs. Scale bar indicates 1 cm. (**c**) 3D reconstruction of μCT. Exemplary images of n= 9.
(**d**) Quantitative results from μCT analysis. Data are shown as mean ± SD. n= 9. (**e**) Structural/morphological
examination of the SFBCs in comparison to (**f**) native bone using scanning electron microscopy. Exemplary images
of n= 3 SFBCs and n= 2 native human bone pieces. Scale bars are indicated in the images. Arrows mark isolated
crystal-like depositions.

Figure 2: Histological and immunohistochemical examination of the SFBCs. (**a**) Exemplary images of H&E, Alizarin red staining and von Kossa stains. Upper row is exemplary for fully calcified SFBCs and lower row for less fully calcified SFBCs. n= 9. (**b**) Negative control for Alizarin red and von Kossa staining. (**d**) Immunohistochemical stainings for Col I, Col II and ALP. All scale bars indicate 200 µm. All images are exemplary for n=9.

SFBCs show profound expression of bone-specific markers

In order to confirm the previous findings on a molecular level, immunofluorescence staining for osteopontin (OPN) and osteocalcin (OC), two non-collagenous bone matrix proteins were analyzed and yielded the evenly distributed, distinct protein expression (**Fig. 3a, Fig. 3b**). While OPN is mainly produced by immature osteoblasts, OC is a marker of late stage osteoblasts indicating the presence of different cell states within the SFBCs. mRNA expression analysis showed high levels of osteogenic marker genes such as *secreted phosphoprotein 1* (*SPP1*) and *distal-less homeobox 5* (*DLX5;* **Fig. 3**). *SPP1* was significantly higher expressed (10-fold), while *DLX5* was higher expressed by trend (8-fold) when compared to monolayer MSCs. In contrast the early osteogenic transcription factor *runt-related transcription factor 2*

(*RUNX2*) and the chondrogenic marker *SRY-box transcription factor 9* (*SOX9*) were expressed
on a more basal level. Interestingly, *receptor activator of NF-κB ligand* (*RANKL)* was highly
expressed (**Fig. 3c**). Moreover, *vascular endothelial growth factor* (*VEGFA*) was significantly
higher expressed (10-fold), although other hypoxia-inducible factor (HIF1) target genes such
as *phosphoglycerate kinase 1* (*PGK1*), *lactate dehydrogenase A (LDHA)*, *endothelial PAS*
domain-containing protein 1 (*EPAS*) and *HIF1* were comparably or lower (*LDHA*) expressed
as in monolayer MSCs (**Fig. 3d**).

167
Figure 3: Immunofluorescence and mRNA expression analysis of SFBCs. (**a**-**b**) Immunofluorescence staining
of osteocalcin (OC, **a**) and osteopontin (OPN, **b**). White asterisks highlight cells with high OPN production. To reveal
the nuclei of present cells, all slides were counter-stained with DAPI. All scale bars indicate 200 µm. Images are
exemplary for n=3. (**c**) qPCR results of mature osteogenic markers (*SPP1*, *DLX5*) (n=8-10), markers indicative for
osteoprogenitors (*RUNX2*, *SOX9*), and (**d**) metabolic marker *VEGFA* is highly expressed while other metabolic
markers (*PGK1, LDHA, HIF1, EPAS*) remain at a basal level (n=10). Data are shown as mean ± SEM. For statistical
analysis the results were compared to the expression level of monolayer hMSC and the Wilcoxon signed rank test
was used (Table S1); *$p<0.05$, **$p<0.01$.

Co-cultivation indicates biological functionality of SFBCs

To confirm the functional activity of the SFBCs and to mimic the initial phase of fracture healing
in an adequate experimental setting, SFBCs were directly co-cultivated with FH models under
hypoxic conditions (5% CO_2, ~ 1% O_2) for 48h, since the initial phase of fracture healing in
humans takes place within the first 72h[2] (**Fig. 4a**).
Although we have not fixed both models to each other by any technical measures, we
observed a close contact of the FH with the corresponding SFBC allowing direct cell-cell-
contact and crosstalk between the models. After co-cultivation for 48h, H&E staining revealed
the typical cell morphology with no obvious alterations within the SFBCs, while the cells in the
FH seemed to be evenly distributed. The calcification throughout the SFBC was reconfirmed

via von Kossa staining with no obvious calcification in the border areas of the FH (**Fig. 4b**).
We have previously shown that most interesting observations in our *in vitro* FH were visible
between 12-48h with respect to immune cell survival and activity[18]. Thus, in a first experiment,
mRNA expression in the FH and the SFBC was analyzed after 12 and 48h and normalized to
the mean expression at 0h (**Fig. S1 and S2**, respectively). In addition, we compared the
cultivation with normal medium (NM) and osteogenic medium (OM, control) to analyze the
effect of the additional osteogenic impact on the co-culture system. In short, analysis of the
mRNA-expression in the FH indicated a time-consistent expression of almost all genes with
only slight differences between NM and OM (**Fig. S1**). The main differences were observed
for *matrix metalloproteinase* (*MMP2*), significantly higher expressed at 48h when cultivated
with OM. The inflammatory markers *interleukin 6* (*IL6*) and *IL8* were slightly lower expressed
after 12h, although gene expression was upregulated in NM and OM after 48h (**Fig. S1**). Within
the SFBCs, *SPP1* was elevated after 12h of incubation with NM and was marginally lower
expressed in NM and higher expressed in OM at 48h compared to 0h. *MMP2*, *VEGFA* and *IL8*
were highly expressed at both time points with no differences between the cultivation medium.
(**Fig. S2**).

Based on this data, further expression analysis and experiments were conducted after
48h of co-cultivation. To verify the osteoinductive potential and biological functionality of the
SFBC, we compared the results from the co-cultivated FH model with a FH only control group
(treated with OM under hypoxic conditions). The normalization of the data to the starting point
(0h) and the FH control revealed a substantial higher expression of *MMP2, VEGF, PGK1* and
IL6 in both groups (NM and OM) when compared to the FH control (**Fig. 4c, d**). While *RUNX2*
and *IL8* were higher expressed in the NM group compared to the OM and FH control group
SPP1 was lower expressed in both groups and *LDHA* only in the NM group. Based on these
findings, we concluded that the SFBCs show a comparable osteoinductive capacity as the OM
medium and biological functionality (**Fig. 4c, d**).

Regarding the protein release, we confirmed the secretion of the pro-inflammatory IL-
6 and the pro-inflammatory/-angiogenic IL-8. The pro-inflammatory granulocyte/macrophage
stimulating factor (GM-CSF) and the macrophage inflammatory protein MIP were released
(**Fig. 5**).

Figure 4: Co-cultivation of SFBCs and *in vitro* FH. (**a**) Experimental setup for the co-cultivation study. (**b**)
Exemplary images of H&E (1, 2) and von Kossa staining (3, 4). Images in the right row (2, 4) are magnifications.
Scale bars show 10 and 5 µm. (**c, d**) qPCR results of the FH model of the co-cultivation system after 48h co-
cultivation under hypoxic conditions. Data is normalized to *EF1A*, 0h and the FH only control group cultured without
SFBC in OM under hypoxic conditions (median). Data are presented as mean ± SEM (n= 3). Statistical significance
was determined using the Wilcoxon signed rank test (matched-pairs and unpaired to hypothetical value = 1; Table
S2 - S5).

Figure 5: Analysis of the supernatant after co-cultivation. Supernatants were collected after 24h and 48h and analyzed via Multiplex assay. Statistical significance was determined using the Kruskal Wallis test with Dunn's multiple comparisons test (Table S4). Data is shown as mean ± SD.

DFO treatment intensifies pro-inflammatory processes

For evaluation of the model suitability as a platform to test potential therapeutics, we supplemented 250 µM DFO for 48h and analyzed the changes in mRNA expression and protein release. DFO is an iron chelator which inhibits prolylhydroxylases to chemically stabilizes HIF and is a well-known osteo-inductive substance[20]. After co-cultivation for 48h in NM, we found that DFO triggered the expression of osteogenic, angiogenic and hypoxia-related genes in the *in vitro* FH with barely any effect on the SFBCs (**Fig. 6a, b**). In detail, the expression of the osteogenic marker *SPP1* in the FH was higher expressed compared to the untreated control, while DFO had barely any effect on the expression of *RUNX2*. The inflammatory markers *MMP2* and *IL6* were additionally elevated while the pro-angiogenic factor *VEGFA* was highly expressed under the influence of DFO compared to the untreated control group, while *LDHA* was also elevated compared to the untreated control (**Fig. 6a**). These results confirm the biological activation capacity of DFO and the possibility to monitor these effects within a short time period. Referring to the mRNA-expression within the SFBCs, we observed a pattern very similar to untreated conditions, which were expected due to the short treatment period (**Fig. 6b**). However, when analyzing the samples treated with OM, we did not see any on the gene expression compared to the untreated control leading to the assumption that the SFBC as natural trigger is more favorable than providing another more artificial (OM) medium (**Fig. 6c, d**). Finally, the protein release of IL-6 and IL-8 was highly induced after 24h NM and DFO treatment (**Fig. 6e**).

Discussion

Our data show that the presented 3D *in vitro* fracture gap model is able to distinctly recapitulate key features of the initial phase of fracture healing. Therefore, we first developed and characterized SFBCs based on mesenchymal condensation, which were subsequently co-cultivated with FH models.

Improved tissue engineering approaches employ mesenchymal condensation as natural form of 3D self-assembly or self-organization consisting exclusively of the cells and their own produced extracellular matrix (ECM)[21]. MSC from the bone marrow, but also the adipose tissue as well as primary cells (osteoblasts) or induced pluripotent stem cells have been used depending on the availability and phenotype stability. It has been described that bone marrow-derived MSCs tend to rather mineralize than undergo chondrogenesis after

mesenchymal condensation[17,19]. Several different techniques are exerted such as the use of
low attachment plates to induce spontaneous MSC aggregation, membrane-based
aggregation (e.g. chitosan) or forced aggregation (via centrifugation)[22].

The applied technique patented by Ponomarev *et al.*[23] exploits the capacity of MSC to
undergo mesenchymal condensation on a macroscale to produce macro-tissues in a highly
reproducible standardized manner without causing necrosis formation in the center[24]. The
advantage of such macroscale approaches is the physiologically relevant size, geometry, low
cell number and density compared to the matrix and mechanical properties when compared to
e.g. spheroids[24]. However, the generation and examination of these constructs are time-
consuming and require high numbers of cells limiting the throughput of the system but enabling
the performance of several analyses from one model. Here, we reported the bone-like structure
and ongoing calcification/mineralization of the SFBCs that were verified by *in vitro* µCt (**Fig. 1**)
and histology/immunohistochemistry, showing pronounced expression of ALP and Col I in the
absence of Col II (**Fig. 2**). These findings are comparable to other studies using MSC
aggregates analyzed after 1 to 3 weeks[25,26]. The morphology and structure of the SFBCs
resemble immature woven bone with regard to the body structure. To identify bone-
representative cell types, we applied immunofluorescence staining. Since osteoblasts and
osteocytes are the characteristic cells in native bone, we stained for OPN as well as OC. OPN
- a glycol-phosphoprotein[27] - is mainly expressed in immature osteoblasts and OC can be
found in late stage osteoblasts currently transforming to osteocytes. Thus, we have
heterogenic differentiation and maturation states within the SFBCs, indicating a heterogeneous
cell population in functional balance. With respect to the mRNA expression, we observed
augmented expression of osteogenic relevant markers such as *SPP1*, *DLX5* and *VEGFA* (**Fig.**
3). *SPP1*, the coding gene for OPN, is hereby the most distinct upregulated gene, coherent
with the expression of OPN within the immunofluorescence staining. Muraglie *et al.* observed
comparable trends including a two to eightfold increase in OC and OPN expression in MSC
aggregates cultivated in low attachment plates[28]. Since *RUNX2* is an early upstream
transcription factor, high expression on mRNA-level is expected 3 to 7 days after osteogenic
induction[26]. After three weeks, the process of ossification and cellular differentiation towards
the osteogenic lineage is already in an advanced state[29,30]. *VEGFA* is an essential coordinator,
not only of angiogenetic processes and important for fracture healing, but also in the process
of endochondral ossification[31] and is known to enhance osteogenic differentiation *in vitro*[32].
Interestingly, other HIF1 target genes such as *PGK1*, *EPAS* and *HIF1* were comparably
expressed as in monolayer MSCs, also *LDHA* was downregulated (**Fig. 3e**). Since the SFBCs
were not cultivated under hypoxic conditions, the increased expression of *VEGFA* might result
from an alternative pathway, e.g. induced by transforming growth factor beta 1 (TGF-β1)[33].
DLX5, an important transcription factor in osteogenesis and bone development[34], is also highly

expressed within the SFBCs, also indicating an intense ossification process. RANKL is also
expressed in mature osteoblasts, differentiating into osteocytes and also regulates
osteoblastogenesis indicating the presence of late-stage osteoblasts[35].

After characterizing their bone-like quantities, we co-cultivated the SFBCs with
in vitro FHs in order to evaluate the capability of the SFBCs to act as an osteogenic inducer
and to recapitulate key features of the initial phase of fracture healing closely to the *in vivo*
situation. Previously, we developed an *in vitro* FH model, incubated in osteogenic induction
medium, which closely reflects the *in vivo* situation[18,36]. One of the main findings was the
importance of hypoxia. Therefore, we included hypoxic conditions in our co-cultivation setup.
The co-cultivation of the *in vitro* FH and SFBCs in NM for up to 48h under hypoxic conditions
revealed significant initiation of ongoing cellular processes (mRNA-level) for adaptation to
hypoxia and osteogenic induction within the FH (**Fig. S1, S2**). These findings are in accordance
with results from an *ex vivo* study and an *in vitro* FH model conducted in our group (**Table**
1)[18,37].

Table 1: Gene expression data from an *ex vivo* study using primary human fracture hematomas obtained between
48 and 72 h after trauma (n=40), the i*n vitro* human FH model (n=12, 48 h incubation under hypoxia and the FHs
of the *in vitro* fracture gap model (n=3, 48 h incubation under hypoxia in normal medium).

Gene symbol	*ex vivo* FHs (< 72 h)	*in vitro* human FHs (48 h hypoxia)	*in vitro* fracture gap model (48 h hypoxia)
RUNX2	↑	↑**	↑
SPP1	↑*	↑****	↑
VEGFA	↑*	↑***	↑
IL8	↑***	↑**	↑
IL6	↑***	↑	↑
LDHA	↑**	↑***	↑
MMP2	n.a.	↑*	↑

332
Based on the comparative analysis with an FH control, we concluded that the SFBCs show a
comparable osteoinductive capacity as the OM medium and biological functionality indicated
by e.g. *VEGF* and *MMP2* (**Fig. 4**). Interestingly, *SPP1* was noticeably lower expressed in the
FH when co-cultivated with SFBCs compared to the FH control, which could be explained by
either the higher amount of OPN provided by the SFBC (**Fig. 3**) or the considerable high
expression/concentration of IL-6 (**Fig. 4 and 5**), as previously reported[38]. The level of pro-
inflammatory cytokines (e.g. IL-6 and IL-8) was abundant, which has been also observed in *ex*
vivo samples from patients (**Fig. 5**)[2,5,37]. The microenvironment of the fracture hematoma is
described by hypoxia, high lactate and low pH due to the disruption of vessels, cell death of
e.g. erythrocytes and the lack of nutrients. This cytotoxic environment needs to be counter-
regulated to allow the invasion of regenerative cells. Therefore, we included whole human

blood instead of isolated peripheral blood mononuclear cells (PBMCs). At the initial stage, the

microenvironment in the fracture hematoma is acidotic and switched to neutral and slightly

alkaline during the regeneration process[39] which can be triggered by hypoxia. Interestingly,

upon co-cultivation *LDHA* was highly upregulated in the FH model (**Fig. S1**).

Furthermore, the effect of certain immune cells during the initial phase of fracture

healing has been studied in detail during the last decade. Lymphocytes play a crucial role as

shown in RAG1(-/-) mice supposing a detrimental effect of adaptive immune cells[40]. A negative

impact was reported for the presence of CD8$^+$ cytotoxic T-cells in humans and mice[41], while

CD4$^+$ cells have been shown to enhance osteogenic differentiation *in vitro* and upregulated

osteogenic markers e.g. *RUNX2* or *OC*[42]. This clearly indicates the need to combine bone and

immune cells in *in vitro* approaches to recapitulate the crosstalk, environment and key features

of the *in vivo* situation.

With respect to the DFO treatment, we found an upregulation of *HIF*-target genes

(*VEGFA* and *LDHA*) indicating the effectivity of DFO to stabilize *HIF* (**Fig. 6**). In addition, pro-

inflammatory processes were more pronounced, which is in accordance with current findings

in a mouse-osteotomy-model revealing the activation of e.g. *C-X-C motif chemokine ligand* 3

(*Cxcl3*) or *metallothionein 3* (*Mt3*) expression at day 3 after application of DFO in the fracture

gap[20]. We did not expect a strong upregulation of osteogenic markers or changes in the

SFBCs, which normally require longer treatment periods[29,43].

Nevertheless, there are many ways to foster the current approach. In order to improve

the current protocol to generate SFBC, different approaches such as the addition of specific

growth factors, e.g. BMPs, fibroblast growth factors (FGFs) or VEGFs, enhancing the

differentiation and bone formation *in vitro* can be considered to fasten up the generation period

in the future. Furthermore, bioreactors can be used to approximate the environment to the

actual *in vivo* situation by dynamic culturing and restrained environment, while overcoming the

lack of nutrient transfer and combining cells with scaffolds[44]. Bioreactors also provide the

possibility to withdraw toxic and cell apoptotic signals perhaps allowing a longer cultivation

period of the fracture gap model.

Taken together, within our 3D *in vitro* fracture gap model, we have been able to

distinctly mimic key features of the initial phase of fracture healing which can be used i) to

study potential underlying mechanism of fracture healing disorders, especially with respect to

immunologically restricted patients[4,5], requiring the crosstalk between immune cells and bone

and ii) as a prediction tool for potential new therapeutic strategies actively implementing the

3R principle.

Material & Methods

Bone marrow derived MSC isolation, cultivation and characterization

Human mesenchymal stromal cells (hMSC) were isolated from bone marrow of patients undergoing total hip replacement (provided by the Center for Musculoskeletal Surgery, Charité-Universitätsmedizin Berlin; donor list in **Table 2**). All protocols were approved by the Charité-Universitätsmedizin Ethics Committee and performed according to the Helsinki Declaration (ethical approval EA1/012/13). MSC isolation was performed as described in detail before[18,45]. Briefly, bone marrow was transferred cell culture flask, covered with normal expansion medium (NM) containing DMEM+GlutaMAX (Gibco), 10 (v/v) % FCS (Biowest), 1 (v/v) % Penicillin/Streptomycin (Gibco), 20 (v/v) % StemMACS MSC Expansion Media XF (Miltenyi Biotech) and incubated at 37°C in 5% CO_2 atmosphere (app. 18% O2). Medium was changed after 3-4 days when cells became adherent. Isolated cells were expanded in at 37 °C, 5% CO_2. Medium exchange was performed weekly and passaging with Trypsin-EDTA (Gibco) was conducted at a cellular confluency of 80-90%. For characterization, MSCs were evaluated at passage 3 for their differentiation potential (osteogenic, adipogenic) and the presence and absence of specific cell surface markers (MSC Phenotyping Kit, Miltenyi Biotech) as described in detail before[18,45]. Human EDTA- blood was collected from healthy donors with written consent[18].

Table 2: hMSC and blood donor information and conducted experiments.

Donor	Age	Sex	Type of experiments	Used methods
MSC 1	71	m		µCt, gene expression analysis, histology, immunofluorescence
MSC 2	77	m		
MSC 3	69	m		
MSC 4	69	w	Characterization of SFBCs	
MSC 5	81	w		
MSC 6	49	w		µCt, gene expression analysis, histology
MSC 7	59	m		
MSC 8	80	w		
MSC 9	51	m		
MSC 10	75	m	Characterization of SFBCs, Co-cultivation experiments	µCt, gene expression analysis, histology, gene expression analysis, DFO treatment studies
MSC 11	68	w		Gene expression analysis, DFO treatment studies
MSC 12	56	m		
Blood 1	37	M	Co-cultivation experiments	
Blood 2	26	M		
Blood 3	38	M		

Fabrication of the 3D bone-like scaffold-free constructs (SFBCs)

SFBCs were produced based on a patented protocol (patent no.: EP1550716B1) [46] which was modified by applying osteogenic medium to induce osteogenic differentiation after 1 week and no application of biomechanical loading to avoid matrix destruction due to mineral formation. The osteogenic medium for generation and maturation contained DMEM/F-12, 10% (v/v) FCS, 1% (v/v) streptomycin/penicillin, 10 mM β-glycerophosphate and 10 nM dexamethasone (Sigma Aldrich). Approx. $10\text{-}20 \times 10^5$ hMSC/cm^2 were cultivated in expansion medium until reaching confluency and forming a monolayer cell sheet. These cell sheets were detached and centrifuged at 350 g for 15 min at RT. Afterwards, the resulting cell aggregates were cultivated for up to one week with medium exchange every day. Cell aggregates were then transferred to a multi-well plate and cultured for up 12 weeks until performing characterization or co-cultivation with the FH model.

Generation of FH and co-cultivation – The fracture gap model

The FH models were generated as described previously [18]. In brief, per FH 2.5×10^5 hMSCs per well were centrifuged for 3 min at 300 g in a 96-well plate (U-bottom). Afterwards, the cell pellet was resuspended in 100 µL of EDTA-blood and subsequently mixed with a 10 mM CaCl$_2$ solution (solved in PBS). After an incubation time of 30 min at 37 °C, 5% CO$_2$ the FH were placed on a SFBC with direct contact and transferred into either NM or osteogenic medium (OM) containing NM supplemented with 10 mM β-glycerophosphate and 0.1 mM L-ascorbic acid-2-phosphate.

For the treatment study, 250 µM DFO (Sigma Aldrich) was supplemented to the medium. The generated fracture gap models were incubated under hypoxic (37 °C, 5% CO$_2$ and ~ 1% O$_2$ - flushed with N$_2$) conditions in a humidified atmosphere for up to 48h.

In vitro micro computed tomography (µCT)

SFBCs were scanned at a nominal resolution of 8 µm, with a SkyScan 1172 high-resolution microCT (Bruker). X-ray tube voltage was set at 80 kV, 124 µA with maximized power of 10W and a 0.5 mm aluminum filter was employed to reduce beam hardening effects. The scan orbit was 360 degrees with a rotation step of 0.2 degree. For reconstruction the Bruker NRecon software accelerated by GPU was used and Gaussian smoothing, ring artifact reduction, misalignment compensation, and beam hardening correction were applied. XY alignment was corrected with a reference scan to determine the thermal shift during scan time. The CTAn software (Bruker) was used to analyze the total VOI of SFBCs. The threshold for bony tissue was set globally (determined by the Otsu method) and kept constant for all SFBCs. For analysis we used the bone volume (BV), the bone surface (BS) and the trabecular pattern factor (TBPf)[47] as measured and calculated by the software.

Scanning electron microscopy (SEM)

Samples were fixated with 2.5 (v/v) % glutaraldehyde (fixation for 10 min), dehydrated with increasing alcohol concentration 30 (v/v) % - 100 (v/v) % (5 steps) and incubation with 100% hexamethyldisilazane (all Sigma Aldrich). Subsequently, the samples were transferred to a sample holder and gold coating was performed with a fine gold coater JFC-1200 (JEOL). For electron microscopy, the JCM-6000 Plus NeoScope (JEOL) was used for imaging, high vacuum was adjusted.

Histological stainings

The embedding and slice preparation of the SFBCs was conducted according to the Kawamoto *et al.* [48] method to prepare slices of undecalcified bones. In detail, the SFBCs were fixated for 6h in a 4 (v/v) % paraformaldehyde solution (PFA; Carl Roth) followed by an ascending sucrose solution treatment (10 (w/v) %, 20 (w/v) % and 30 (w/v) %) for 24h, respectively and afterwards cryo-embedded with SCEM medium (Sectionlab). Slices were produced with a cryotom using cryofilms (Sectionlab) and afterwards air dried for 20 min and fixated with 4 (v/v) % PFA prior to every histological or immunhistological staining on a microscope slide.

H&E staining was performed as described previously[24]. In short, slices were fixed with 4 (v/v) % PFA (10 min), washed with distilled water and stained with Harris's hematoxylin solution (Merck Millipore). Staining was followed by several washing steps, a differentiation step (0.25 (v/v) % concentrated HCl) and a second staining step in 0.2 (w/v) % eosin (Chroma Waldeck). Staining was finished by differentiation in 96 (v/v) % and 100 (v/v) % ethanol, fixation with xylol and covering with Vitro-Clud (R. Langenbrinck GmbH).

Alizarin Red S staining was conducted by applying slices after fixation and washing to the 2 (w/v) % Alizarin Red S staining solution (Sigma Aldrich; pH = 4.1-4.3) for 10 min. Afterwards slices were washed in distilled water and differentiation was performed in 0.1 (v/v) % HCL solved in ethanol and fixed by washing two times with 100% ethanol before xylol fixation and covering.

Von Kossa staining was conducted according to the following protocol: air drying, fixation and washing as described above, 3% (w/v) silver nitrate solution (10 min), washing step with distilled water, sodium carbonate formaldehyde solution (2 min), washing step with tap water, 5% (w/v) sodium thiosulphate solution (5 min), washing step with tap water and distilled water, ascending ethanol series (70 (v/v) % - 100 (v/v) %), fixation in xylol and covering.

For Col I, II and ALP staining, slices were rehydrated with phosphate buffered saline (PBS) treated with 3 (v/v) % H_2O_2 (30 min), washed with PBS, blocked with 5% normal horse or goat serum (Vector Laboratories) in 2 (w/v) % bovine serum albumin (BSA), and incubated overnight with primary antibodies at 4 °C (Col I antibody: ab6308, 1:500, Abcam; Col II antibody, 1:10, Quartett Immunodiagnostika; ALP antibody, ab95462, Abcam). Afterwards,

slices were washed and treated with 2 (v/v) % secondary antibody (biotinylated horse anti-
mouse IgG antibody – Col I and Col II; biotinylated goat anti-rabbit IgG antibody - ALP, Vector
Laboratories) diluted in 2 (v/v) % normal horse/goat serum/2 (v/v) % BSA/PBS (30 min),
washed with PBS, incubated with avidin-biotin complex (Vectastain Elite ABC HRP Kit, Vector
Laboratories) (50 min), washed with PBS, incubated with DAB under microscopic control with
time measurement (DAB peroxidase (HRP) Substrate Kit, Vector Laboratories) and stopped
with PBS. For counterstaining slices were washed with distilled water and stained with Mayer's
hematoxylin (Sigma Aldrich), washed with tap water and covered with Aquatex (Merck
Millipore). Pictures were taken with the Axioskop 40 optical microscope (Zeiss) using the
corresponding AxioVision microscopy software. Von Kossa staining was quantified using
ImageJ and the threshold tool to mark positive (black) and negative (brown) stained areas.

Immunofluorescence

For the immunofluorescence staining, the slides were rehydrated with PBS and blocked with
PBS with 5 (v/v) % FCS for 30 min at RT. Primary osteopontin antibody (mouse anti-human,
Abcam) or osteocalcin (rabbit anti-human, Abcam) was diluted 1:50 in PBS/5 (v/v) % FCS/0.1
(v/v) % Tween 20 and incubated for 2h. After washing with PBS/0.1% Tween 20, the secondary
antibody (donkey anti-goat A568; Life Technologies/Thermo Scientific) was diluted 1:500 in
PBS/5 (v/v) % FCS/0.1 (v/v) % Tween 20 and applied for 1h. Pictures were taken with a
Keyence fluorescence microscope BZ 9000 (Keyence) using the DAPI, TexasRed and Cy5
channels.

RNA isolation and quantitative PCR (qPCR)

SFBCs were transferred to RLT-buffer (Qiagen, Germany) with 1% 2-Mercaptoethanol (Serva)
and disrupted using the Qiagen Tissue Ruptor (Qiagen). Total RNA was extracted using the
RNeasy Fibrous Tissue Mini Kit (Qiagen) according to the manufacturers´ instructions and the
RNA concentration was determined using the Nanodrop ND-1000 (Peqlab). RNA was stored
at -80 °C until further processing. The same RNA isolation method was preceded for the
SFBCs after co-cultivation.
After co-cultivation, the cells of the FH were filtered through a cell strainer (Corning). After
centrifugation for 10 min at 300 *g*, the cell pellet was resuspended in 350 µL RLT buffer with
3.5 µL 2-Mercaptoethanol and total RNA was extracted using the Rneasy Fibrous Tissue Mini
Kit according to the manufacturers´ instructions.
The cDNA for both the FH and the SFBCs was synthesized by reverse transcription using
TaqMan Reverse Transcription Reagents (Applied Biosystems). qPCR was performed using
the DyNAmo Flash SYBR Green qPCR Kit (Thermo Fisher) and the Stratagene Mx3000P
(Agilent Technologies). Initial denaturation was for 7 min at 98 °C. Afterwards 50 cycles with 5
sec at 98 °C, 7 sec at 56 °C and 9 sec at 72 °C were performed. The melting curve was

analyzed through stepwise increasing the temperature from 50 °C to 98 °C every 30 sec. All
primers were purchased from TIB Molbiol (**Table S10**). For the gene expression of the SFBCs,
data were normalized to the expression of *eukaryotic translation elongation factor 1 alpha 1*
(EF1A) and to the corresponding MSC culture in 2D, using the delta-delta-Ct-method. For the
gene expression in the fracture gap model, data were normalized to the expression of *EF1A*
using the deltaCt method and 0h using the deltadeltaCt method.

Cytokine and chemokine quantification in supernatants

Supernatants were immediately stored at -80 °C after 48h co-cultivation. The concentration
[pg/mL] of cytokines and chemokines was determined using multiplex suspension assay (Bio-
Rad Laboratories) following the manufacturers´ instructions. Following cytokines and
chemokines (lower detection limit) were measured: IL-1β (7.55 pg/mL), IL-2 (18.99 pg/mL), IL-
4 (4.13 pg/mL), IL-5 (20.29 pg/mL), IL-6 (25.94 pg/mL), IL-7 (16.05 pg/mL), IL-8 (37.9 pg/mL),
IL-10 (37.9 pg/mL), IL-13 (7.21 pg/mL), IL-17 (24.44 pg/mL), interferon-gamma (IFNγ, 56.32
pg/mL), tumor necrosis factor-alpha (TNFα, 59.53 pg/mL), monocyte chemotactic protein-1
(MCP-1, 27.02 pg/mL), macrophage inflammatory protein MIP-1β (6.27 pg/mL), granulocyte
colony-stimulating factor (G-CSF, 50.98 pg/mL) and granulocyte-macrophage colony-
stimulating factor (GM-CSF, 11.82 pg/mL)

Statistical analysis

Statistical tests were performed using GraphPad Prism Software version 8. Statistical
differences towards a hypothetical value were determined by Wilcoxon signed rank test
(unpaired). With respect to the co-cultivation studies, differences between two groups were
determined with Wilcoxon matched-pairs signed rank test or between more groups with the
Kruskal Wallis test with Dunn's multiple comparisons test. Probability values of $p<0.05$ were
considered to be statistically significant (***$p<0.001$, **$p<0.01$, *$p<0.05$). Details on the
statistics per Figure are displayed in the Supplementary Information.

Acknowledgments

The authors would like to thank Manuela Jakstadt for excellent technical assistance. AL, FB,
AD, MP and TG are members of Berlin-Brandenburg research platform BB3R and Charité 3[R].
This study was funded by the German Federal Ministry for Education and Research (BMBF)
(project no. 031A334). AL is currently being supported by the Joachim Herz Foundation (Add-
on Fellowship 2019). The work of TG was funded by the Deutsche Forschungsgemeinschaft
(353142848). Funding bodies did not have any role in designing the study, in collecting,
analyzing and interpreting the data, in writing this manuscript, and in deciding to submit it for
publication.

Contributions

Study design: AL, TG, MP; Data collection and analysis: MP, AL, IP, AD, CB, YC; Data discussion and interpretation: FB, CTR, AL, TG, MP; Drafting manuscript: MP, AL, TG; Revising manuscript: FB, IP, CTR, PH.

Conflict of interest

The authors declare no conflict of interests.

References

Claes, L., Recknagel, S. & Ignatius, A. Fracture healing under healthy and inflammatory conditions. *Nature reviews. Rheumatology* **8**, 133-143, doi:10.1038/nrrheum.2012.1 (2012).

Hoff, P. *et al.* Immunological characterization of the early human fracture hematoma. *Immunologic research* **64**, 1195-1206, doi:10.1007/s12026-016-8868-9 (2016).

Oe, K. *et al.* An in vitro study demonstrating that haematomas found at the site of human fractures contain progenitor cells with multilineage capacity. *The Journal of bone and joint surgery. British volume* **89**, 133-138, doi:10.1302/0301-620x.89b1.18286 (2007).

Hoff, P. *et al.* Immunologically restricted patients exhibit a pronounced inflammation and inadequate response to hypoxia in fracture hematomas. *Immunologic research* **51**, 116-122, doi:10.1007/s12026-011-8235-9 (2011).

Hoff, P. *et al.* A Pronounced Inflammatory Activity Characterizes the Early Fracture Healing Phase in Immunologically Restricted Patients. *International journal of molecular sciences* **18**, DOI: 10.3390/ijms18030583, doi:10.3390/ijms18030583 (2017).

Winkler, T., Sass, F. A., Duda, G. N. & Schmidt-Bleek, K. A review of biomaterials in bone defect healing, remaining shortcomings and future opportunities for bone tissue engineering: The unsolved challenge. *Bone & Joint Research* **7**, 232-243, doi:10.1302/2046-3758.73.BJR-2017-0270.R1 (2018).

Grundnes, O. & Reikeras, O. The importance of the hematoma for fracture healing in rats. *Acta orthopaedica Scandinavica* **64**, 340-342, doi:10.3109/17453679308993640 (1993).

Kolar, P. *et al.* The early fracture hematoma and its potential role in fracture healing. *Tissue engineering. Part B, Reviews* **16**, 427-434, doi:10.1089/ten.TEB.2009.0687 (2010).

Schmidt-Bleek, K., Petersen, A., Dienelt, A., Schwarz, C. & Duda, G. N. Initiation and early control of tissue regeneration - bone healing as a model system for tissue regeneration. *Expert opinion on biological therapy* **14**, 247-259, doi:10.1517/14712598.2014.857653 (2014).

Schmidt-Bleek, K. *et al.* Inflammatory phase of bone healing initiates the regenerative healing cascade. *Cell and tissue research* **347**, 567-573, doi:10.1007/s00441-011-1205-7 (2012).

Bernhardt, A., Lode, A., Peters, F. & Gelinsky, M. Optimization of culture conditions for osteogenically-induced mesenchymal stem cells in beta-tricalcium phosphate ceramics with large interconnected channels. *J Tissue Eng Regen Med* **5**, 444-453, doi:10.1002/term.331 (2011).

Burska, A. N. *et al.* Dynamics of Early Signalling Events during Fracture Healing and Potential Serum Biomarkers of Fracture Non-Union in Humans. *J Clin Med* **9**, doi:10.3390/jcm9020492 (2020).

13 Schlundt, C. *et al.* Immune modulation as a therapeutic strategy in bone regeneration.
Journal of experimental orthopaedics **2**, 1, doi:10.1186/s40634-014-0017-6 (2015).
14 Haffner-Luntzer, M. *et al.* Review of Animal Models of Comorbidities in Fracture-
Healing Research. *J Orthop Res* **37**, 2491-2498, doi:10.1002/jor.24454 (2019).
15 Peric, M. *et al.* The rational use of animal models in the evaluation of novel bone
regenerative therapies. *Bone* **70**, 73-86, doi:10.1016/j.bone.2014.07.010 (2015).
16 Scheinpflug, J. *et al.* Journey into Bone Models: A Review. *Genes* **9**,
doi:10.3390/genes9050247 (2018).
17 Dickhut, A. *et al.* Calcification or dedifferentiation: requirement to lock mesenchymal
stem cells in a desired differentiation stage. *Journal of cellular physiology* **219**, 219-
226, doi:10.1002/jcp.21673 (2009).
18 Pfeiffenberger, M. *et al.* The in vitro human fracture hematoma model - a tool for
preclinical drug testing   *ALTEX - Alternatives to animal experimentation.*,
doi:10.14573/altex.1910211 (2020).
19 Farrell, M. J. *et al.* Functional properties of bone marrow-derived MSC-based
engineered cartilage are unstable with very long-term in vitro culture. *Journal of
biomechanics* **47**, 2173-2182, doi:10.1016/j.jbiomech.2013.10.030 (2014).
20 Lang, A., Helfermeier, S., Stefanowski, J., Kuppe, A., Sunkara, V.,, Pfeiffenberger, M.,
Wolter, A., Damerau, A., Hemmati-Sadeghi, J., Ringe, J., Hauser, A.E., Löhning, M.,
Perka, C., Duda, G.N., Hoff, P., Schmidt-Bleek, K., Gaber, T., Buttgereit, F. HIF-
stabilization as possible treatment towards fracture healing disorders. *in preparation for
submission* (2020).
21 DuRaine, G. D., Brown, W. E., Hu, J. C. & Athanasiou, K. A. Emergence of scaffold-
free approaches for tissue engineering musculoskeletal cartilages. *Ann Biomed Eng*
43, 543-554, doi:10.1007/s10439-014-1161-y (2015).
22 Sart, S., Tsai, A.-C., Li, Y. & Ma, T. Three-dimensional aggregates of mesenchymal
stem cells: cellular mechanisms, biological properties, and applications. *Tissue
engineering. Part B, Reviews* **20**, 365-380, doi:10.1089/ten.TEB.2013.0537 (2014).
23 Ponomarev, I. V., Kochneva, L. M. & Barnewitz, D. Effect of 3D chondrocyte culturing
conditions on the formation of extracellular matrix in cartilage tissue-engineering
constructs. *Bulletin of experimental biology and medicine* **156**, 548-555,
doi:10.1007/s10517-014-2394-3 (2014).
24 Weber, M.-C. *et al.* In vitro and in silico modeling of cellular and matrix-related changes
during the early phase of osteoarthritis. *bioRxiv*, 725317, doi:10.1101/725317 (2019).
25 Frith, J. E., Thomson, B. & Genever, P. G. Dynamic three-dimensional culture methods
enhance mesenchymal stem cell properties and increase therapeutic potential. *Tissue
Eng Part C Methods* **16**, 735-749, doi:10.1089/ten.TEC.2009.0432 (2010).
26 Kabiri, M. *et al.* 3D mesenchymal stem/stromal cell osteogenesis and autocrine
signalling.   *Biochem   Biophys   Res   Commun*   **419**,   142-147,
doi:10.1016/j.bbrc.2012.01.017 (2012).
27 Standal, T., Borset, M. & Sundan, A. Role of osteopontin in adhesion, migration, cell
survival and bone remodeling. *Exp. Oncol.* **26**, 179-184 (2004).
28 Muraglia, A. *et al.* Formation of a chondro-osseous rudiment in micromass cultures of
human bone-marrow stromal cells. *J Cell Sci* **116**, 2949-2955, doi:10.1242/jcs.00527
(2003).
29 Hoshiba, T., Kawazoe, N., Tateishi, T. & Chen, G. Development of stepwise
osteogenesis-mimicking matrices for the regulation of mesenchymal stem cell
functions. *J Biol Chem* **284**, 31164-31173, doi:10.1074/jbc.M109.054676 (2009).
30 Xu, J., Li, Z., Hou, Y. & Fang, W. Potential mechanisms underlying the Runx2 induced
osteogenesis of bone marrow mesenchymal stem cells. *Am J Transl Res* **7**, 2527-2535
(2015).
31 Gerber, H. P. *et al.* VEGF couples hypertrophic cartilage remodeling, ossification and
angiogenesis during endochondral bone formation. *Nat. Med.* **5**, 623-628 (1999).
32 Zhang, Y., Madhu, V., Dighe, A. S., Irvine, J. N., Jr. & Cui, Q. Osteogenic response of
human adipose-derived stem cells to BMP-6, VEGF, and combined VEGF plus BMP-

6 in vitro. *Growth factors (Chur, Switzerland)* **30**, 333-343,
doi:10.3109/08977194.2012.720574 (2012).

33 Chae, K. S. *et al.* Opposite functions of HIF-α isoforms in VEGF induction by TGF-β1
under non-hypoxic conditions. *Oncogene* **30**, 1213-1228, doi:10.1038/onc.2010.498
(2011).

34 Chung, I. H., Han, J., Iwata, J. & Chai, Y. Msx1 and Dlx5 function synergistically to
regulate frontal bone development. *Genesis (New York, N.Y. : 2000)* **48**, 645-655,
doi:10.1002/dvg.20671 (2010).

35 Nakashima, T. *et al.* Evidence for osteocyte regulation of bone homeostasis through
RANKL expression. *Nat Med* **17**, 1231-1234, doi:10.1038/nm.2452 (2011).

36 Pfeiffenberger, M. *et al.* Hypoxia and mesenchymal stromal cells as key drivers of initial
fracture healing in an equine in vitro fracture hematoma model. *PLOS ONE* **14**,
e0214276, doi:10.1371/journal.pone.0214276 (2019).

37 Hoff, P. *et al.* Human immune cells' behavior and survival under bioenergetically
restricted conditions in an in vitro fracture hematoma model. *Cellular & Molecular*
Immunology **10**, 151-158, doi:10.1038/cmi.2012.56 (2013).

38 Scutera, S. *et al.* Adaptive Regulation of Osteopontin Production by Dendritic Cells
Through the Bidirectional Interaction With Mesenchymal Stromal Cells. *Front Immunol*
9, 1207, doi:10.3389/fimmu.2018.01207 (2018).

39 Schell, H. *et al.* The haematoma and its role in bone healing. *Journal of experimental*
orthopaedics **4**, 5, doi:10.1186/s40634-017-0079-3 (2017).

40 El Khassawna, T. *et al.* T Lymphocytes Influence the Mineralization Process of Bone.
Front Immunol **8**, 562-562, doi:10.3389/fimmu.2017.00562 (2017).

41 Reinke, S. *et al.* Terminally differentiated CD8(+) T cells negatively affect bone
regeneration in humans. *Science translational medicine* **5**, 177ra136,
doi:10.1126/scitranslmed.3004754 (2013).

42 Grassi, F. *et al.* T cell subsets differently regulate osteogenic differentiation of human
mesenchymal stromal cells in vitro. *J Tissue Eng Regen Med* **10**, 305-314,
doi:10.1002/term.1727 (2016).

43 Qu, Z. H., Zhang, X. L., Tang, T. T. & Dai, K. R. Promotion of osteogenesis through
beta-catenin signaling by desferrioxamine. *Biochem Biophys Res Commun* **370**, 332-
337, doi:10.1016/j.bbrc.2008.03.092 (2008).

44 Kashte, S., Jaiswal, A. K. & Kadam, S. Artificial Bone via Bone Tissue Engineering:
Current Scenario and Challenges. *Tissue Engineering and Regenerative Medicine* **14**,
1-14, doi:10.1007/s13770-016-0001-6 (2017).

45 Gaber, T. *et al.* Impact of Janus Kinase Inhibition with Tofacitinib on Fundamental
Processes of Bone Healing. *International journal of molecular sciences* **21**,
doi:10.3390/ijms21030865 (2020).

46 Ponomarev, I. & Wilke, I. Manufacturing process of three dimensional tissue structures
and structures obtainable thereby. (2004).

47 Hahn, M., Vogel, M., Pompesius-Kempa, M. & Delling, G. Trabecular bone pattern
factor—a new parameter for simple quantification of bone microarchitecture. *Bone* **13**,
327-330, doi:https://doi.org/10.1016/8756-3282(92)90078-B (1992).

48 Kawamoto, T. & Kawamoto, K. Preparation of thin frozen sections from nonfixed and
undecalcified hard tissues using Kawamot's film method (2012). *Methods in molecular*
biology (Clifton, N.J.) **1130**, 149-164, doi:10.1007/978-1-62703-989-5_11 (2014).

Supplementary Information

693 **Table S1:** Results from Wilcoxon signed rank test for Figure 3c-e

Specification	Wilcoxon signed rank test Hypothetical median = 1 *p*-value	95% CI
RUNX2	*0.7344*	-0.50 – 68.05
SPP1	***0.0039***	0.82 – 27.34
DLX5	*0.1563*	-0.54 – 19.25
SOX9	*0.7422*	-0.99 - 10.51
RANKL	***0.0391****	-0.99 – 351.10
VEGF	***0.002***	0.17 – 25.72
PGK1	*0.2754*	-0.45 – 2.22
LDHA	***0.0195****	-0.66 – 0.20
HIF	*0.5566*	-0.76 – 2.18
EPAS	*0.7695*	-0.82 – 0.03

Figure S1: Co-cultivation of SFBCs and *in vitro* fracture hematomas. qPCR results of the fracture hematoma
model or the after 12 and 48h co-cultivation under hypoxic conditions. Data are normalized to *EF1A* and 0h and
presented as mean ± SEM (n= 3). Statistical significance was determined using the Kruskal Wallis test with Dunn's
multiple comparisons test and Wilcoxon signed rank test (Table S2 - S5). **p*<0.05.

Figure S2: Co-cultivation of SFBCs and *in vitro* fracture hematomasqPCR results of the SFBCs after 12 and
48h co-cultivation under hypoxic conditions. Data are normalized to *EF1A* and 0h and presented as mean ± SEM
(n= 3). Statistical significance was determined using the Kruskal Wallis test with Dunn's multiple comparisons test
and Wilcoxon signed rank test (Table S2 - S5). *p<0.05.

Table S2: Results from Wilcoxon signed rank test for Figure 4c,d

Specification including 0h	Wilcoxon signed rank test hypothetical median = 1 *p*-value
48h NM - RUNX2	*0.25*
48h OM - RUNX2	*0.25*
48h NM - SPP1	*0.25*
48h OM - SPP1	*0.25*
48h NM - MMP2	*0.25*
48h OM - MMP2	*0.25*
48h NM - PGK1	*0.25*
48h OM - PGK1	*0.50*
48h NM - LDHA	*0.50*
48h OM - LDHA	*0.25*
48h NM - VEGF	*>0.99*
48h OM - VEGF	*0.25*
48h NM - IL8	*0.50*
48h OM - IL8	*0.25*
48h NM - IL6	*0.50*
48h OM - IL6	*0.25*

711   **Table S3:** Results from Wilcoxon signed rank test for Figure 4c,d

Specification including 0h	Wilcoxon matched-pairs signed rank test *p*-value
NM vs. OM - RUNX2	*0.50*
NM vs. OM - SPP1	*0.50*
NM vs. OM - MMP2	*0.25*
NM vs. OM - PGK1	*0.50*
NM vs. OM - LDHA	*0.25*
NM vs. OM - VEGF	*0.25*
NM vs. OM - IL8	*0.50*
NM vs. OM - IL6	*0.75*

713   **Table S4:** Results from Kruskal Wallis test with Dunn's multiple comparisons test for Figure 5

Specification	Kruskal-Wallis test		Adjusted *p*-value if applicable
	H	***p*-value**	
IL-6	1.77	*0.67*	-
IL-8	1.97	*0.63*	-
GM-CSF	4.74	*0.21*	-
MIP	0.64	*0.92*	-

715   **Table S5:** Results from Wilcoxon signed rank test for Figure 6a

Specification including 0h	Wilcoxon signed rank test hypothetical median = 1 *p*-value
RUNX2	*>0.99*
SPP1	*0.25*
MMP2	*0.50*
PGK1	*0.75*
LDHA	*0.50*
VEGF	*0.50*
IL8	*>0.99*
IL6	*0.50*

718 **Table S6:** Results from Wilcoxon signed rank test for Figure 6b

Specification including 0h	Wilcoxon signed rank test hypothetical median = 1 *p*-value
SPP1	0.75
MMP2	0.50
VEGF	0.50
IL8	0.75

719 **Table S7:** Results from Wilcoxon signed rank test for Figure 6c

Specification including 0h	Wilcoxon signed rank test hypothetical median = 1 *p*-value
RUNX2	0.50
SPP1	0.50
MMP2	0.25
PGK1	0.75
LDHA	>0.99
VEGF	0.75
IL8	0.25
IL6	0.25

720 **Table S8:** Results from Wilcoxon signed rank test for Figure 6d

Specification including 0h	Wilcoxon signed rank test hypothetical median = 1 *p*-value
SPP1	0.25
MMP2	>0.99
VEGF	0.75
IL8	>0.99

721 **Table S9:** Results from Kruskal Wallis test with Dunn's multiple comparisons test for Figure 6c

Specification	Kruskal-Wallis test		Adjusted *p*-value if applicable
	H	*p*-value	
IL-6	1.15	0.81	-
IL-8	2.54	0.52	-
GM-CSF	2.69	0.49	-
MIP	1.68	0.68	-

724 **Table S10:** Sequences of primers used for qPCR.

	Gene	Sequence of forward primer	Sequence of reverse primer
SPP1	*Secreted phosphoprotein 1*	GCCGAGGTGATAGTGTGGTT	TGAGGTGATGTCCTCGTCGTCTG
VEGFA	*Vascular endothelial growth factor A*	AGCCTTGCCTTGCTGCTCTA	GTGCTGGCCTTGGTGAGG
DLX5	*Distal-Less Homeobox 5*	GCTGGGATTGACACAAACAC	AGGCACCATTGAAAGTGTCC
RUNX2	*Runt-related transcription factor 2*	TTACTTACACCCCGCCAGTC	TATGGAGTGCTGCTGGTCTG
EF1A	*Elongation factor 1-alpha*	GTTGATATGGTTCCTGGCAAGC	TTGCCAGCTCCAGCAGCCT
MMP2	*Matrix metalloproteinase-2*	GATACCCCTTTGACGGTAAGGA	CCTTCTCCCAAGGTCCATAGC
IL8	*Interleukin 8*	GGACCCCAAGGAAAACTGG	CAACCCTACAACAGACCCACAC
IL6	*Interleukin 6*	TACCCCCAGGAGAAGATTCC	TTTTCTGCCAGTGCCTCTTT
RANKL	*Receptor Activator of NF-κB Ligand*	CTCAGCCTTTTGCTCATCTCACT	CCAAGAGGACAGACTCACTTTATGG
SPI1	*Transcription factor PU.1*	GTGCCCTATGACACGGATCTA	AGTCCCAGTAATGGTCGCTAT
SOX9	*SRY (sex determining region Y)-box 9*	CGCCTTGAAGATGGCGTTG	GCTCTGGAGACTTCTGAACGA
PGK1	*Phosphoglycerate kinase 1*	ATGGATGAGGTGGTGAAAGC	CAGTGCTCACATGGCTGACT
LDHA	*Lactate dehydrogenase A*	ACCCAGTTTCCACCATGATT	CCCAAAATGCAAGGAACACT
HIF1A	*Hypoxia-inducible factor 1-alpha*	CCATTAGAAAGCAGTTCCGC	TGGGTAGGAGATGGAGATGC
EPAS	*Endothelial PAS domain-containing protein 1*	TCGGCTTTTTGCCATCTGTG	TGTCCAAATGTGCCGTGTGA