Normal mammary epithelia accumulate DNA methylation changes with age that are

consistent with breast cancer methylomes

Parijat Senapati[1], Masaru Miyano[2], Rosalyn W. Sayaman[2,4], Mudaser Basam[1,2], Candi Trac[1],

Amy Leung[1], Mark A. LaBarge[2,3], Dustin E. Schones*[1,3]

[1]Department of Diabetes Complications and Metabolism, Beckman Research Institute, City of Hope,

Duarte, CA, 91010, USA

[2]Department of Population Sciences, Beckman Research Institute, City of Hope, Duarte, CA, 91010, USA

[3]Irell & Manella Graduate School of Biological Sciences, City of Hope, Duarte, CA, USA

[4]Department of Laboratory Medicine, Helen Diller Family Comprehensive Cancer Center, University of

California, San Francisco, San Francisco, CA, USA

*Corresponding Author:

Dustin E. Schones, Ph.D.

Department of Diabetes Complications and Metabolism

City of Hope

Duarte, CA, 91010

Email: dschones@coh.org

Phone: 626-218-1319

Keywords: aging, DNA methylation, human mammary epithelial cells, luminal epithelial cells,

breast cancer, transposable elements

Running Title: DNA methylome characteristics of aging mammary luminal cells

Significance: Mammary luminal epithelial cells lose lineage-specific expression with age and

accumulate stochastic methylation changes that lead to loss of silencing at transposable

elements. These age-dependent events potentially promote breast cancer susceptibility.

Abstract

Luminal epithelial cells (LEps) are a key cell lineage implicated in age-related luminal breast

cancers. Alterations to the epigenome are a hallmark of aging cells. However, the extent of age-

associated DNA methylation alterations in LEps, and the corresponding functional

consequences of these alterations, have remained unclear. We report here that aging leads to

distinct methylation changes in LEps. Luminal lineage-specific genes gain promoter methylation,

whereas myoepithelial-specific genes lose promoter methylation. Regulatory elements display

methylation changes at lineage-specific TF binding sites consistent with the loss of lineage

fidelity. CpG islands (CGIs) and transposable elements (TEs) have stochastic methylation gain

and loss, respectively. PRC2 target genes that are hypermethylated in luminal breast cancer

exhibit stochastic methylation increase with age. TEs with stochastic methylation loss are

activated in breast cancer and potentially function as regulatory elements contributing to the loss

of lineage fidelity with age. Each of these classes of methylation changes impact the regulation

of genes associated with luminal breast cancer. Altogether, our results indicate that aging leads

to DNA methylation changes that could determine breast cancer susceptibility.

Introduction

Over 75% of breast cancer diagnoses are in women over the age of 50, making age the most significant risk factor for breast cancer (1). This is especially true for luminal subtypes of all age-related breast cancers (2). The mammary epithelium is composed of two principal lineages, myoepithelial cells (MEps) and luminal epithelial cells (LEps), with the latter being the chief suspect cell of origin for luminal breast cancers (3,4). Most breast cancers are sporadic with no known inherited mutation or family history. A better understanding of how age-dependent molecular changes contribute to breast cancer risk is critical for designing better therapies and prevention strategies.

Aging in the breast tissue is associated with significant physiological changes that lead to changes in molecular properties and the proportion of the different cell populations with age (5-8). Human mammary epithelial cells (HMECs) from older post-menopausal women, when transformed *in vitro*, exhibit gene expression patterns characteristic of luminal breast cancer subtypes. In contrast, transformed HMECs from younger pre-menopausal women show more properties of basal breast cancers (9). Striking age-dependent changes in abundance and phenotype occur in LEps, which gain expression of characteristic MEp genes (6,10,11) and signaling molecules consistent with those of immortalized luminal cells (10), suggesting that age-associated epigenetic changes in luminal cells are directly related to elevated cancer risk.

DNA methylation is a relatively stable component of the epigenome that gets established in early development and is maintained through successive cycles of cell division to preserve cell identity (12). DNA methylation patterns, however, are known to change with age (13-16) and DNA methylation levels of specific CpG sites show a significant correlation with chronological age (14). This 'epigenetic clock' can be used to predict the biological age of diverse human tissues (14). However, this pan-tissue epigenetic clock is "poorly calibrated" in breast tissue, even for women without cancer, possibly due to physiological factors such as changes in

hormonal status with age (14). Previous studies examining DNA methylation changes in breast

tissue with age were either performed in bulk mammary tissue with heterogeneous cell

populations or interrogated only a subset of CpGs in the genome (17,18). A complete

understanding of age-associated DNA methylation changes in human mammary LEps and how

it relates to age-related breast cancer has been lacking.

We performed whole-genome bisulfite sequencing (WGBS) and Assay for Transposase-

Accessible Chromatin (ATAC-seq) in non-senescent primary human mammary luminal epithelial

cells from pre-menopausal (young) and post-menopausal (old) women. We find that LEps from

older women display DNA methylation changes at regulatory elements associated with altered

expression and activity of lineage-specific transcription factors such as ELF5 and TP63. We

used chromatin state predictions from mammary epithelial cell lineages to show that MEp

lineage-specific regulatory elements lose methylation, whereas luminal-specific regions gain

methylation. Additionally, PRC2 (polycomb repressive complex 2) regulated genes gain DNA

methylation at their promoters with age, consistent with these genes having methylated

promoters in luminal breast cancer (19). We also observed increased methylation disorder with

age, resulting in stochastic methylation gain at CpG islands (CGIs) and loss at evolutionarily

recent transposable elements (TEs). Using breast cancer WGBS data, we show that TEs that

lose methylation with age were also hypomethylated in breast cancers. In addition, we used

ATAC-seq and RNA-seq data from the TCGA-BRCA cohort and RNA-seq done on DCIS (ductal

carcinoma in situ) samples to show that TEs with age-dependent methylation loss are activated

in breast cancer and can function as regulatory elements for nearby genes. Overall, our results

suggest that aging mammary luminal epithelial cells undergo DNA methylation alterations at

regulatory elements, CGIs, and evolutionarily recent TEs, similar to methylation changes

observed in luminal breast cancer.

Results

Aging leads to altered DNA methylation patterns in mammary luminal epithelial cells.

We performed whole-genome bisulfite sequencing (WGBS-seq) on human mammary LEps from younger (<30y) and older (>55y) women (**Figure S1A, Table S1**). The age groups were chosen to represent samples from women who were clearly pre-menopausal or post-menopausal to avoid confounding issues of peri-menopause. We previously demonstrated that LEps from these primary cultures maintain gene expression and DNA methylation profiles that mirror those in tissue (8,11). Moreover, these are non-senescent cells and therefore are a valuable resource that can be used to interrogate aging-dependent changes in the breast tissue – distinct from those arising from cellular senescence.

WGBS libraries were sequenced to ~30X genome coverage, generating ~ 1B 150-bp paired-end reads (~86 Gb uniquely mapped sequence per sample) (**Table S2**) to obtain a representation of about 95% of individual CpG sites in our samples (see Methods and **Tables S2, S3**). We compared our DNA methylation profiles with previously published WGBS studies on distinct human mammary epithelial cell types from reduction mammoplasty (RM) tissue of a young (19y old) donor (20) and found that the cultured LEps resemble the ER-negative (ER⁻) LEp population (**Figure S1B**). Young and old LEPs clustered separately indicating differences in methylomes (**Figure S1B**) that were not apparent when considering total methylation levels (**Figure S1C, D**). Pairwise comparisons of methylomes also showed higher number of differentially methylated regions (DMRs) between young and old pairs (**Figure S1E**). Comparing the methylomes of old versus young LEps, we identified 18,860 differentially methylated regions (DMRs) (see Methods), with 7,207 showing methylation loss and 11,653 showing methylation gain in old LEps. Most DMRs were 100-500bp in length (**Figure S1F**). The methylation changes at DMRs were confirmed with heatmaps showing methylation levels across samples (**Figure**

1A), as well as kernel smoothed methylation plots of representative DMRs (**Figure 1B**). The

majority of DNA methylation changes were observed at non-promoter regions (**Figure S1G-H**).

We assigned DMRs to genes based on overlap with either promoters or gene body (exonic

or intronic) (**Table S4A-B**). We used the Molecular Signatures Database (MSigDB) to identify

the gene sets or pathways over-represented among these genes (**Table S5A-D**). Genes with

methylation loss at promoters belong to gene sets expressed in mammary stem cells, cell-cell

anchoring junctions epithelial-to-mesenchymal transition, and genes downregulated in luminal

breast cancer (**Figure 1C, Table S5A**). Genes with promoter methylation gain were enriched for

luminal specific genes upregulated in luminal breast cancer and Polycomb (PRC2) target genes

(**Figure 1D, Table S5C**, BENPORATH gene sets). We also noted that genes expressed in

mammary stem cell enriched basal compartment showed methylation loss (**Figure S2A**)

whereas luminal lineage restricted genes showed methylation gain (**Figure S2B**). As an

example, *FOXP1*, a mammary stem cell specific gene essential for differentiation into various

mammary epithelial lineages (21), shows methylation loss at several sites its promoter and gene

body (**Figure 1E**). *ALDH1A3*, on the other hand, is an ER$^-$ luminal lineage restricted gene (22)

which gains methylation at several sites including the promoter in old LEps (**Figure 1F**). DNA

methylation changes with age suggest that luminal cells increase stem cell related properties

while repressing genes that are specifically related to maintenance of the luminal lineage.

Aging-associated DNA methylation changes occur at lineage-specific transcription factor

binding sites

To investigate whether DNA methylation changes were associated with transcription

factor (TF) binding signatures, we performed motif enrichment analysis (see Supplemental

Methods for details) using position weight matrices (PWMs) from the JASPAR database (23).

DMRs that lose methylation are enriched for binding sites of TFs such as EGR1, TP63, TP73,

TEAD1, TEAD4, and RUNX1 (**Figure 2A left panel, Table S6A**). Several of these TF binding

sites, including TEAD, TP63, TP73, NFI, and EGR1, are enriched at active regulatory elements

in basal/myoepithelial cells (20). To investigate whether these TFs change expression with age,

we checked their expression using RNA-sequencing data from LEps and MEps of young and

old individuals (11). *TP63* and *NFIA (NFI family TF; binds to NFI half-sites)* showed increased

expression in old compared to young LEps (**Figure 2B and S3A**). We also checked whether

these TFs showed any DNA methylation changes. *EGR1* showed significant methylation loss at

its promoter (**Figure S3B**) whereas *NFIA* and *TP63* exhibit methylation loss at specific sites

within their gene bodies (**Table S4A**) suggesting altered methylation at regulatory elements.

Although not all TFs showed an increase in expression with age, we noted that *TP63, NFIA,*

EGR1, and TP73 have relatively higher expression in MEps as compared to LEps (**Figure 2A**

right panel and 2E). The expression patterns of TFs and the hypomethylation of their targets

are concordant with the known roles of these TFs. For example, young LEps do not express

TP63, which is a MEp lineage-specific transcription factor (8). Furthermore, TEAD1 and TEAD4

are DNA binding partners for YAP (Yes-associated protein), a transcriptional co-activator

regulated by the Hippo signaling pathway. YAP, typically a MEp specific protein, is sequestered

in the cytoplasm through activity of kinases downstream of the Hippo pathway (24). Old LEps

exhibit increased *TP63* expression and nuclear YAP localization (7,8), which leads to increased

TEAD binding and YAP activity. YAP is also known to bind p73 (TP73 protein) and RUNX2 (24).

These results suggest that increased expression/activity of myoepithelial-specific TFs in old

LEps lead to increased binding and loss of DNA methylation in these regions (**Figure 2F**).

DMRs with methylation gain are enriched for binding sites of the ETS family of

transcription factors EHF, ELF3, ELF5, ELF4, ETV1, ETS1, ETV4, GABPA, and NFI half and full

sites (**Figure 2C left panel, Table S6B**). ELF5 is a luminal lineage-specific transcription factor

that decreases expression with age (**Figure 2D**) (8). Both *ELF3* and *ELF5* genes also show

significant methylation gain at their promoter and gene locus (**Figure S3C-D**). In addition, *ELF5*, *EHF* and *ELF3* exhibit relatively higher expression in LEps as compared to MEps (**Figure 2C right panel and 2E**). These results further suggest that decreased expression of luminal lineage-specific TFs may lead to a decrease in binding and protection of binding sites from DNA methyltransferases (DNMTs). This, in turn, may result in the methylation and repression of these luminal lineage-specific regions (**Figure 2F**). We also examined the known DNA methylation binding sensitivities of TFs (25) with enriched binding sites. TFs with binding sites showing methylation loss either do not have a CpG at their respective site or their binding is unaffected by DNA methylation (**Table S7**). Presumably they recruit other factors such as TETs to bring about DNA methylation changes. However, TFs with binding sites showing methylation gain are sensitive to DNA methylation (**Table S7**). Generally, sites of methylation loss were enriched for myoepithelial-specific TF binding motifs (**Figure 2E**), whereas sites of methylation gain were enriched for luminal lineage-specific TF binding motifs (**Figure 2E**). For example, SOX4 sites are enriched in DMRs with methylation gain, and it is a luminal-specific TF (**Figure 2E**). SOX15, on the other hand, is a myoepithelial-specific TF and its binding sites are enriched at DMRs with methylation loss (**Figure 2E**). Expression of *SOX4* decreases, whereas *SOX15* increases with age in old LEps (**Figure S3E-F**), indicating loss of expression of luminal lineage-specific TFs and increased expression of myoepithelial-specific TFs. Collectively, we observe that aging-associated DNA methylation changes in LEps occur at lineage-specific TF binding sites that seem to contribute to loss of cell identity. Enticingly this is a characteristic of cancer cells whereby even the so-called luminal subtype cancer cells exhibit expression of genes and proteins that are characteristic of basal cells.

Lineage-specific regulatory regions are hotspots for methylation changes with age.

Given that aging-associated DNA methylation changes occurred at regions containing

lineage-specific TF binding sites, we examined whether these binding sites are functional by

measuring chromatin accessibility using ATAC-seq in young and old LEps (**Table S1**). Open

chromatin region at a TF binding site is a reliable indicator of a regulatory element with bound

TFs as not all TF binding sites in a given condition in a cell are bound. DMRs with methylation

loss showed corresponding increased accessibility in older LEps (**Figure 3A-B**). On the other

hand, DMRs with methylation gain showed a decrease in chromatin accessibility (**Figure 3C-D**).

These results indicate that functional regulatory elements undergo DNA methylation changes

with age.

To further investigate the functional relevance of the age-dependent DNA methylation

changes in the breast tissue, we used published histone modification ChIP-seq datasets in

mammary epithelial lineages to predict chromatin states at DMRs (20). To discern whether age-

dependent DNA methylation changes were occurring at lineage-specific regulatory elements,

we predicted chromatin states in ER^+ luminal, ER^- luminal, and basal (MEp) cells from RM tissue

of a young (19y old) donor (20). We classified DMRs into seven different chromatin states that

are characteristic of i) Active TSS, ii) TSS flanking/promoter, iii) transcribed, iv) Enhancer, v)

Heterochromatin, vi) Repressed and vii) None/unclassified regions based on the combinatorial

histone modification levels in each cell type (see Supplemental Methods; **Figure S4A-C**). DMRs

with methylation loss in old LEps overlap primarily with MEp-specific promoter and enhancer

regions (~80% of methylation loss DMRs) (**Figure 3E**). Because these sites also contain binding

sites for TFs enriched at MEp-specific active enhancers (20), these results indicate that MEp-

specific regulatory elements lose methylation in old LEps. Active enhancer sites are

hypomethylated (26,27) and methylation loss at MEp-specific regulatory elements might indicate

an epigenetic shift towards MEp gene expression patterns in LEps with age.

Conversely, DMRs with methylation gain overlap putative promoter and enhancer regions

in ER- luminal cells, accounting for about 42% of methylation gain DMRs (**Figure 3F**). As these

DMRs also contain binding sites for TFs enriched at luminal lineage-specific active enhancers

(20), these results indicate that luminal-specific regulatory regions gain DNA methylation in aging

LEps. Together, these results suggest that aging LEps acquire permissive chromatin states at

MEp-specific regions and repress ER- luminal-specific chromatin regions through changes in

DNA methylation.

Polycomb repressive complex 2 target genes gain DNA methylation with age.

The Polycomb repressive complex 2 (PRC2) is a histone methyltransferase complex that

trimethylates lysine 27 of histone H3 (H3K27me3) and mediates transcriptional silencing of

developmentally important genes. As per current understanding PRC2 is targeted to

unmethylated CpG-dense regions, primarily CpG islands (CGIs) that occur at gene promoters

(28).   Earlier we found that genes with methylation gain at their promoters were enriched in

PRC2 target genes identified in embryonic stem cells (ESCs) (**Figure 1D**). Moreover, about 15%

of the DMRs with methylation gain overlapped with Polycomb repressed regions marked by

H3K27me3 (**Figure 3F and S4A-C**). PRC2 target gene promoters have been described to gain

DNA methylation with age in various tissues as well as in mice (13,29-32). They are also known

to be hypermethylated in diverse types of cancer, some among them are known tumor

suppressor genes (33). Since the MSigDB PRC2 target gene sets are ESC-specific, we used

H3K27me3 ChIP-seq data from ER- and ER+ luminal cells, and basal cells (20) to determine if

genes with promoter methylation gain are PRC2 targets in mammary epithelial lineages. We

found that 39-46% of promoters that gain methylation with age are marked by H3K27me3 across

the three cell lineages (**Figure S5A**). In contrast, only 14-27% of promoters with methylation loss

are enriched for H3K27me3 (**Figure S5B**). Promoters with DNA methylation gain also have

conserved H3K27me3 occupancy in all three epithelial lineages (**Figure S5C**) as compared to

promoters with DNA methylation loss that show H3K27me3 only in the luminal lineage (**Figure**

S5D). These results indicate that promoters with age-dependent DNA methylation gain are

indeed *bona fide* PRC2 target genes in mammary epithelial cells. These results are consistent

with PRC2 protecting unmethylated CpG-rich regions and that erosion of the PRC2 targeting

with aging (32) may be fundamentally related to increased methylation at its target genes.

Stochastic methylation loss at transposable elements (TE) and methylation gain at CpG

islands (CGIs) with age.

Aging is associated with increased transcriptional variability, cell-to-cell heterogeneity,

and DNA methylation variability (34-37). Because we use a sorted luminal lineage cell

population, we can discern DNA methylation variability in LEps eliminating confounding

problems such as changes in cell proportions with age. To examine whether there is an increase

in methylation heterogeneity in older individuals, we calculated methylation entropy, a measure

of methylation heterogeneity between adjacent CpGs, for 150 bp non-overlapping windows

across the genome (38). There are two modes by which a given region could increase

methylation entropy. An unmethylated region can gain methylation stochastically or a methylated

region can lose methylation stochastically (**Figure 4A-B**). Considering these regions, we

observed an overall increase in methylation entropy in the old LEps. We identified regions with

greater methylation entropy in older individuals and found a net decrease in methylation levels

at these regions (**Figure S6A-D**), consistent with the fact that most of the genome is methylated

in normal cells. Of these, unmethylated regions with increased methylation entropy were mainly

at CGIs (69.2%; 1017 of 1469; **Figure 4C**). Conversely, methylated regions with increased

entropy were found mainly at repeat regions (70.3%; 76575 of 108877; **Figure 4D**), which are

known to be enriched with TEs. These results indicate that increased DNA methylation

heterogeneity with age leads to stochastic methylation gain at CGIs and methylation loss at repeat regions.

To examine the potential of stochastic methylation changes to have consequences for breast cancer, we used a published WGBS dataset of 30 primary tumors containing ER+ (luminal breast cancer) and ER- (basal breast cancer) samples (39). At CGIs with increased entropy, luminal breast tumors showed an even greater increase in methylation and entropy than observed in old LEps (**Figure 4E**). In contrast, basal tumors did not show an increase in CGI methylation or entropy (**Figure 4F**), indicating that this behavior is specific to the luminal subtypes of breast cancer. We identified the subset of entropy-gained CGIs that were hypermethylated in at least 20% (n >= 6) of the breast tumors. These CGIs were enriched at PRC2 target genes, which are known to be hypermethylated in luminal breast cancer (19) and other types of cancer (**Figure S6E**). Of note, PRC2 target genes are repressed in normal tissues through H3K27me3, and although they show DNA methylation gain at their CGI promoters, expression changes are not expected. Apart from PRC2 target genes, we also found gene sets that are upregulated in basal breast cancer and downregulated in luminal B breast cancer to be enriched at these CGIs (**Figure S6E**). For example, *FOXC1* gene is expressed in LEps and normal breast tissue but downregulated specifically in luminal and Her2-enriched breast cancers (**Figure 4G**). Taken together, these results indicate that increased DNA methylation heterogeneity with age leads to stochastic methylation gain at CGIs which is further exacerbated in luminal breast cancer.

Increased DNA methylation variability between old individuals at TEs.

We next tested whether aging would lead to increased interindividual variability in addition to increased methylation entropy. We indeed do observe increased variability of CpG methylation in older individuals (**Figure S7A**). Since some amount of DNA methylation variability

is expected in human populations and differences at individual CpG sites may not have

functional consequences, we first identified systemic regions of variable methylation across all

10 individuals using the Systemic Interindividual Variation (SIV) approach (40) (see

Supplemental Methods). We identified 23,921 regions that were variably methylated only in old

samples and 25,585 regions that were specific to the young samples (**Figure S7B**). Regions

variably methylated in old samples mainly showed methylation loss compared methylation levels

in younger individuals (17,780 of 23,921; 74.3%). The variable regions were ~150p-450bp in

length with some spanning 2-2.5kb (**Figure S7C and S7D**). We next examined whether variable

regions contained TEs and found that 68.7% of the regions variable in young samples contained

TEs as compared to 67.59% of control non-variable regions. 71.4% regions variable in old

samples overlapped with TEs as compared to 67.65% of control non-variable regions (**Figure**

S7E). Among the TEs that are variably methylated in old samples, about 76.9% lose methylation

and 23.13% gain methylation (**Figure S7F**). We also confirmed that variable methylation at TEs

was not due to mapping issues (**Figure S7G**).

One of the main functions of DNA methylation is to silence TEs, especially those that are

evolutionarily recent and more likely to be capable of activity (41,42). To investigate whether the

TEs that are variably methylated in old samples are evolutionarily recent, we calculated the

Jukes-Cantor evolutionary age of each TE copy. Among the variably methylated TEs, the

evolutionarily recent elements (inserted <100 million years ago; (43)) tended towards DNA

methylation loss, whereas the older TE elements (>100 million years) tended to gain methylation

in old cells (**Figure 5A-D)**. These results indicate that evolutionarily recent TEs lose methylation

with age in a stochastic manner. While evolutionarily recent Alu elements such as AluY and AluS

are highly abundant in the genome, a few AluS subfamilies such as AluSz and AluSx were

enriched above background for stochastic loss of methylation (**Figure 5E, S7H**). Other

evolutionarily recent TE subfamilies were enriched at regions with variable methylation loss as

well, including both L1PA (primate-specific LINE-1) and HERVH elements (**Figure 5E-F)**.

Many TEs in the human genome have retained latent regulatory potential and can serve

as regulatory elements when not suppressed (44-46). We therefore examined the TF motifs

enriched at specific TE subfamilies with variable DNA methylation loss in old. We found that Alu

subfamilies contain motifs for Homeobox TFs such as SIX4, BAPX1 and nuclear receptor TFs

such as RARA which are upregulated in breast cancer (**Figure 5G**). Moreover, the younger

L1PA subfamilies L1PA(2-4) have motifs for many TFs; the most relevant among them being

binding sites for GATA, FOXA1 TFs and EREs which are bound by Estrogen receptor (ESR1)

(**Figure 5G**). GATA3, FOXA1 and ESR1 are highly upregulated in luminal breast cancers (47).

These results indicate that stochastic loss of methylation at these TEs can lead to binding of TFs

expressed during breast cancer development.

To examine whether TEs similarly contribute TF binding sites to DMR regions associated

with loss of lineage fidelity, we identified the TEs that overlap DMRs. Although many DMRs

overlap TEs, given the high number of copies in the genome only a few TE subfamilies,

especially those of the LTR class, were enriched over background. Several LTR subfamilies,

including HERVH, MER52A, MER52D and LTR27E were enriched at DMRs with methylation

loss in old LEps (**Figure S8A**) while LTR10C, HERVE, and MER11D (**Figure S8B**) were

enriched in DMRs with methylation gain. Among the TEs that lose methylation, HERVH elements

showed the highest enrichment for TEAD and RUNX TF binding sites (**Figure S8C**), motifs that

were also found in non-repetitive DMRs with methylation loss (**Figure 2A**) and known binding

partners of the MEp specific YAP. TP63 binding sites, which were also found at DMRs with

methylation loss (**Figure 2A**) were enriched at AluSx, AluSp, L2, and L3 elements. ETS family

TF binding sites that gain methylation were found within MIRb, MIR subfamilies of SINE

elements and L2, L2a-c LINE subfamilies (**Figure S8D**), similar to non-repetitive DMRs with

methylation gain (**Figure 2C**). SOX family TF binding sites that also occur in DMRs with methylation gain (**Figure 2E**) were present in AluY, LTR10 and MER11D elements with methylation gain (**Figure S8D**). Most of these DMR-associated TEs are evolutionarily older, indicating that they have been co-opted by lineage-specific TFs and contribute to aging-dependent DNA methylation changes in LEps.

Transposable elements that lose methylation with age are activated in breast cancer.

Given our observations that evolutionarily recent transposons lose methylation stochastically with age and contain motifs for TFs overexpressed in breast cancer, we examined whether breast cancers show loss of epigenetic suppression at these elements. Loss of TE suppression can lead to aberrant transcription of TE-derived transcripts and alternative isoforms due to splicing with neighboring genes in cancer cells (48). To investigate the potential of this happening in breast cancer, we examined published RNA-seq data from high-grade ductal carcinoma in situ (DCIS), a form of pre-cancer (49). We used the SQuIRE pipeline (50) to measure gene and TE expression changes in DCIS as compared to normal tissues; 554 age-hypomethylated TEs showed higher expression in DCIS. Of these, 468 were variably hypomethylated with age and 86 were within DMRs with methylation loss. As with the loci with variable methylation loss in older samples, HERVH elements and L1PA2 elements were most common among the TEs that showed aberrant expression. A representative example of a HERVH element that loses methylation in aging LEps and breast tumors is shown (**Figure 6A**). Aberrant transcription from this HERVH element was observed only in DCIS samples (**Figure 6A**). We also found examples of exonization of TEs, where a MIRb element was included within the exon of a non-coding RNA DRAIC expressed in DCIS (**Figure S9A**). While aberrant expression of hypomethylated TEs in DCIS was clear, we did not observe breast cancer subtype specificity in the expression of these TEs.

We next examined whether TEs with methylation loss function as regulatory elements in

breast cancer. To this end we utilized ATAC-seq data from The Cancer Genome Atlas (TCGA)

breast cancer cohort (TCGA-BRCA) and ChIP-seq datasets from breast cancer cell lines (51).

We identified 1197 TCGA-BRCA ATAC-seq peaks that overlap TEs with methylation loss in

aging LEps (**Figure 6B**). Although some TEs were accessible in basal subtype tumors (418 of

1197), most of them were accessible across luminal tumors (779 of 1197 peaks) (**Figure 6B**).

Some luminal tumors that were diagnosed as lobular carcinomas had particularly high chromatin

accessibility across all TEs (**Figure 6B**). These results indicate that TEs hypomethylated with

age may function as regulatory elements in breast cancer.

To determine whether loss of methylation at TEs can lead to changes in expression of

genes that are thought to be important in breast cancer, we mapped the 554 age-

hypomethylated TEs to genes within $\pm$500kb and calculated the correlation of TE expression

with each gene mapped within this window in DCIS samples (49). We identified 406 genes that

were significantly correlated with the expression of a nearby TE (**Table S8**). A representative

example of an L1PA4 element with significant expression correlation (correlation: 0.8, *p*-val:

$7.29e^{-9}$) with the nearby *ZNF92* gene is shown (**Figure S10A**). We find that *ZNF92* is also highly

expressed in luminal subtype breast tumors (Luminal A and Luminal B) as compared to basal

tumors (**Figure 6C**). These results suggest that TEs located at deleterious locations in the

genome are reawakened through stochastic loss of methylation causing expression of genes

that are associated with breast cancers. Overall suggesting that reawakening of TEs is directly

related to increased cancer susceptibility with age.

Discussion

Previous studies analyzing aging-associated DNA methylation changes in human

mammary luminal epithelial cells have either used array-based technologies (8,11,17,18) or

were performed using composite tissues that represented a mixture of cells (17,18). Array

probes are mainly enriched at genic regions and intergenic elements including repeats and distal

regulatory elements are under-represented (52). It has furthermore been shown that different

cell lineages have unique responses to aging (8,53), making pure cell populations critical for

interpretation. Our use of WGBS on sorted normal human mammary epithelial cells addresses

both shortcomings.

We find that aging leads to DNA methylation changes at lineage-specific gene regulatory

elements in mammary luminal epithelial cells, in-line with previous studies that show loss of

lineage fidelity in luminal cells with age (8). Specifically, luminal cells show DNA methylation loss

at myoepithelial-specific regulatory elements containing MEp-specific TF binding sites.

Moreover, luminal-specific regulatory regions containing LEp-specific TF binding sites are

repressed by DNA methylation with age. This loss of lineage-fidelity has been shown to occur

with age and in mammary tissue that is susceptible to breast cancer (11,54).

Previous DNA methylome studies in human cells suggested a global loss of methylation

with age (16,38,55). Those studies were performed using CD4$^+$ T cells from a neonate and a

centenarian (16), sorted CD4$^+$ T cells (38) and skin keratinocytes (55) from young (<25 y) and

old (>70 y) individuals. The global methylation loss with age was accompanied by a global

increase in entropy or methylation discordance (38). We did not observe a global methylation

loss or a global increase in entropy in mammary luminal epithelial cells with age. This could be

due to higher fidelity of DNA methylation maintenance in luminal cells or due to a lesser degree

of separation between the age of young (<30 years) and old (> 55 years) individuals in our study.

DNA methylome studies in aging mouse livers also showed no global loss of methylation, but

specific changes at regulatory elements were detected (56). Whereas global methylation loss

was not detected, we did detect methylation changes at regulatory elements and variable

methylation changes at evolutionarily recent transposons.

Global loss of methylation and focal increase in methylation, termed 'epigenetic drift', has

been reported to occur with age and in cancer (57). The focal increase in methylation has been

reported to occur mainly at CpG islands (CGIs). We find CGIs showing both gain and loss in

methylation reflecting changes in lineage-specific gene expression. However, we do observe

increased methylation disorder/entropy at CGIs that lead to methylation gains with age, which is

further enhanced during the transition to luminal breast cancer. These CGIs were predominantly

PRC2 targets that are known to gain methylation in other cancer types (58). PRC2 targets were

also found among genes that showed methylation gain at promoters. Loss/repositioning of PRC2

has been proposed to cause DNA methylation gain at its targets. Presumably, CGIs with

increased methylation entropy are in the process of becoming hypermethylated in luminal breast

cancer. Hypermethylation of PRC2 targets seems to be specific to the luminal subtype of breast

cancer (19), although the reasons for this are not currently clear.

DNA methylation is an important mechanism for silencing of transposable elements.

Reactivation of transposable elements has been shown in cancer cells, coincident with a global

loss of DNA methylation (59,60). Previous research comparing DNA methylation levels of CD4[+]

T cells between a neonate and a centenarian reported that 12% of DMRs that lost methylation

with age were located on Alu elements (16), whereas changes in epigenetic silencing at TEs

with aging has not been systematically investigated. Increased expression of retrotransposons

and active retrotransposition have been reported in aging mouse tissues (61,62) and increased

LINE-1 expression has been reported in senescent human fibroblasts *in vitro* (63). However,

these studies used bulk quantitative strategies such as RT-qPCR, which are not ideal due to

high sequence similarity between multiple active and inactive copies. Our approach allowed us

to investigate the impact of aging on DNA methylation at specific TEs using non-senescent

mammary epithelial cells. Our results demonstrate that evolutionarily recent (e.g primate and

hominid specific) transposons are more likely to be silenced by DNA methylation in young and

show increased variability and methylation loss with age. These TEs could be an important

component of understanding differential susceptibility to cancer as not all aged individuals

develop cancer. Our data also underscore the need for use of human cells or organoids for future

studies as these cannot be recapitulated in mice.

We further show that the methylation loss observed at TEs with age becomes

exacerbated in frank luminal breast cancers. The derepressed TEs can lead to aberrant

transcription or function as regulatory elements in breast cancer cells. We demonstrate that

hypomethylated TEs could regulate genes associated with luminal subtype breast cancers. TEs

also contribute lineage-specific TF binding sites to regulatory elements associated with loss of

lineage fidelity with age. Retrotransposon activation has been widely observed in nearly every

human cancer, including breast cancer (64,65), and even in premalignant lesions (60). This

suggests that early events in tumorigenesis may be driven by retrotransposon activity. We show

here, for the first time, that loss of transposon silencing is evident in aging luminal epithelial cells

that could predispose towards breast cancer. Future studies are needed to characterize the role

of these TEs in the early events of mammary cancer initiation. Altogether, our results showing

DNA methylation alterations at regulatory elements, CGIs and TEs with aging provide further

evidence in support of the observation that epigenetic changes with age could determine

susceptibility towards breast cancer.

Materials and Methods

Isolation of mammary luminal epithelial cells

LEps were isolated from 4^{th} passage (p4) cultures of pre-stasis finite-lifespan HMECs from

normal reduction mammoplasty (RM) tissues (6). Primary non-immortalized HMECs were

generated and maintained, as described previously (66). HMEC strains were established in the

laboratory of Dr. Martha Stampfer (Lawrence Berkeley National Laboratory) who has all required

IRB approvals to distribute these samples. Cells were grown in M87A medium with cholera toxin

and oxytocin at 0.5 ng/ml and 0.1nM, respectively. 4^{th} passage HMECs were stained with anti-

human CD133-PE (Biolegend, clone 7) and anti-human CD271 (Biolegend, clone ME20.4) by

following standard flow cytometry protocol. LEps and MEps were enriched using Aria III (Becton

Dickinson) flow sorter as $CD133^+/CD271^-$ and $CD133^-/CD271^+$ fractions, respectively. The

different HMEC strains used are described in Table S1.

WGBS

We examined the DNA methylome of primary LEps from younger (< 30y) pre-menopausal and

older (> 55y) post-menopausal women. Genomic DNA from FACS enriched LEps were extracted

using Quick-DNA Microprep kit (Zymo Research). Whole-genome bisulfite sequencing (WGBS-

seq) was performed at Admera Health Biopharma Services. Genomic DNA was bisulfite

converted using the Zymo EZ DNA methylation kit (Zymo Research) per manufacturer's protocol.

Library preparation was then performed using Accel-NGS® Methyl-Seq DNA Library kit (Swift

Biosciences, Michigan, USA) per manufacturer's recommendations. Equimolar pooling of

libraries was performed based on QC values. Samples were sequenced on an Illumina®

NovaSeq S4 (Illumina, California, USA) with a read length configuration of 150bp paired-end.

We obtained about 38X coverage per genome to an average of 414 million paired-end reads

(Table S2).

Sequencing reads were hard trimmed 15 bp from both ends using Trim Galore

(https://github.com/FelixKrueger/TrimGalore) (version 0.4.5) to remove unwanted methylation

bias arising from the library preparation. Adapter contamination was also removed using Trim

Galore using --illumina option. Trimmed reads were aligned to the human reference genome

(hg19) using Bismark (version 0.19.1) (67) with the --bowtie1 option (68). Bismark

(deduplicate_bismark) to remove PCR duplicates. Methpipe (69) (version 3.4.3) was used to

determine the methylation status of individual CpG sites. The hg19 reference genome contains

about 56.4 million CpG loci (hg19) or 28.2 million CpG dyads. Methylation levels were calculated

for individual CpG dyads and are henceforth referred to as CpG sites.

DMR identification

DMRfinder (70) was used to identify DMRs. Briefly, CpG sites with at least 4X coverage in at

least 6 out of 10 samples were determined. Genomic regions with at least 4 CpG sites within a

distance of 100 bp were identified. The maximum length of regions was set to 500 bp. Of these

regions, ones that had less than 20 reads covering them were discarded. The final set of regions

was then used to calculate differentially methylated regions with greater than 20% change (*P*-

value < 0.05) between pooled young and old samples.

ATAC-seq and analysis

50,000 live FACS sorted young (*n* = 4) and old (*n* = 4) LEps were used for Tn5 transposition in

duplicates using the Omni-ATAC method (71). Libraries were sequenced to a depth of 40 million

paired reads each on an Illumina HiSeq 2500 system. Raw sequencing reads were trimmed

using Trim Galore (https://github.com/FelixKrueger/TrimGalore) (version 0.4.5) to remove

adapters. Trimmed reads were aligned to the human reference genome (hg19) using bowtie (68)

with the --best -k 1 -X 2000 --mm --chunkmbs 1024 parameters. Unmapped reads, mates as

well as low quality reads and secondary alignments were removed using samtools. PCR

duplicates were identified using Picard and removed using samtools. The reads were shifted by

4 bases on the sense strand and 5 bases on the anti-sense strand to account for Tn5 bias.

Reads from technical replicates were pooled to generate pseudoreplicates. Peaks were called

using MACS2 version 2.1.1.20160309 using -p 0.01 --nomodel --shift -37 --extsize 73 --SPMR -

-keep-dup all --call-summits as parameters. Peak calls from technical replicates and pseudo-

replicates were compared using IDR to get the most conservative set of peaks for each sample.

Enrichment of DMRs and variably methylated regions at TE subfamilies

The extent of overlap of DMRs and variable regions at repat subfamilies (RepeatMasker

annotation) was calculated using bedtools intersect. Randomly shuffled control regions of the

same length were generated 1000 times, and the expected number of overlaps calculated using

bedtools intersect. Enrichment values were calculated by dividing the number of DMR overlaps

(observed) with the average number of random overlaps (expected). *P* values were calculated

by dividing the number of extreme values (at either tail of distribution) from random overlaps by

1000. Significantly enriched repeat subfamilies had $p < 0.05$, enrichment > 1.5-fold, and > 5

copies overlapping DMRs or variable regions.

Availability of data and materials

All raw and processed sequencing data files have been deposited at NCBI's Gene Expression

Omnibus (GEO) under accession GSE153696 (WGBS and ATAC-seq) and GSE182338 (RNA-

seq).

Funding

This work was supported by the National Institutes of Health, grants R01DK112041,

R01CA220693 (D.E.S.); K01DK104993 (A.L.); Department of Defense/Army Breast Cancer Era

of Hope Scholar Award BC141351, and Expansion Award BC181737, Conrad N. Hilton

Foundation, and City of Hope Center for Cancer and Aging (M.A.L.); the National Cancer

Institute (NCI) Cancer Metabolism Training Program Postdoctoral Fellowship T32CA221709

(R.W.S.) and City of Hope Program in Molecular and Cellular Biology pilot award (D.E.S and

M.A.L.). The research reported in this publication included work performed in the Integrative

Genomics and Analytical Cytometry Cores supported by the National Cancer Institute of the

National Institutes of Health under award number P30CA033572. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. The funding bodies had no role in the design of the study, data collection,

analysis, and interpretation, decision to publish of data or in writing the manuscript.

Authors' contributions

PS, MAL and DES designed the study. PS, MM and CT performed experiments. PS, RWS, MB

and AL analyzed and interpreted the data. PS, MAL and DES prepared the manuscript. All

authors read and approved the final manuscript.

Acknowledgements

We thank other members of the Schones laboratory for helpful discussions and comments, and

the continued involvement of patient advocates Susan Samson and Sandy Preto. The results

shown and referenced here are based in part upon data generated by the TCGA Research

Network: https://www.cancer.gov/tcga, and the ENCODE Consortium:

https://www.encodeproject.org/.

Conflict of Interest Statement

The authors declare that they have no potential conflicts of interest.

References

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. CA Cancer J Clin **2021**;71:7-
33
2. Jenkins EO, Deal AM, Anders CK, Prat A, Perou CM, Carey LA*, et al.* Age-specific changes in
intrinsic breast cancer subtypes: a focus on older women. Oncologist **2014**;19:1076-83
3. Prat A, Perou CM. Mammary development meets cancer genomics. Nat Med **2009**;15:842-4
4. Fu NY, Nolan E, Lindeman GJ, Visvader JE. Stem Cells and the Differentiation Hierarchy in
Mammary Gland Development. Physiol Rev **2020**;100:489-523
5. Benz CC. Impact of aging on the biology of breast cancer. Crit Rev Oncol Hematol **2008**;66:65-
74

6.   Garbe JC, Pepin F, Pelissier FA, Sputova K, Fridriksdottir AJ, Guo DE, *et al.* Accumulation of
multipotent progenitors with a basal differentiation bias during aging of human mammary
epithelia. Cancer Res **2012**;72:3687-701
7.   Pelissier FA, Garbe JC, Ananthanarayanan B, Miyano M, Lin C, Jokela T, *et al.* Age-related
dysfunction in mechanotransduction impairs differentiation of human mammary epithelial
progenitors. Cell Rep **2014**;7:1926-39
8.   Miyano M, Sayaman RW, Stoiber MH, Lin CH, Stampfer MR, Brown JB, *et al.* Age-related
gene expression in luminal epithelial cells is driven by a microenvironment made from
myoepithelial cells. Aging (Albany NY) **2017**;9:2026-51
9.   Lee JK, Garbe JC, Vrba L, Miyano M, Futscher BW, Stampfer MR, *et al.* Age and the means of
bypassing stasis influence the intrinsic subtype of immortalized human mammary epithelial cells.
Front Cell Dev Biol **2015**;3:13
10. Pelissier Vatter FA, Schapiro D, Chang H, Borowsky AD, Lee JK, Parvin B, *et al.* High-
Dimensional Phenotyping Identifies Age-Emergent Cells in Human Mammary Epithelia. Cell
Rep **2018**;23:1205-19
11. Sayaman RW, Miyano M, Senapati P, Shalabi S, Zirbes A, Todhunter ME, *et al.* Epigenetic
changes with age primes mammary luminal epithelia for cancer initiation. bioRxiv
2021:2021.02.12.430777
12. Bird A. DNA methylation patterns and epigenetic memory. Genes Dev **2002**;16:6-21
13. Maegawa S, Hinkal G, Kim HS, Shen L, Zhang L, Zhang J, *et al.* Widespread and tissue specific
age-related DNA methylation changes in mice. Genome Res **2010**;20:332-40
14. Horvath S. DNA methylation age of human tissues and cell types. Genome Biol **2013**;14:R115
15. Rakyan VK, Down TA, Maslau S, Andrew T, Yang TP, Beyan H, *et al.* Human aging-associated
DNA hypermethylation occurs preferentially at bivalent chromatin domains. Genome Res
2010;20:434-9
16. Heyn H, Li N, Ferreira HJ, Moran S, Pisano DG, Gomez A, *et al.* Distinct DNA methylomes of
newborns and centenarians. Proc Natl Acad Sci U S A **2012**;109:10522-7
17. Hofstatter EW, Horvath S, Dalela D, Gupta P, Chagpar AB, Wali VB, *et al.* Increased epigenetic
age in normal breast tissue from luminal breast cancer patients. Clin Epigenetics **2018**;10:112
18. Castle JR, Lin N, Liu J, Storniolo AMV, Shendre A, Hou L, *et al.* Estimating breast tissue-
specific DNA methylation age using next-generation sequencing data. Clin Epigenetics
2020;12:45
19. Holm K, Hegardt C, Staaf J, Vallon-Christersson J, Jonsson G, Olsson H, *et al.* Molecular
subtypes of breast cancer are associated with characteristic DNA methylation patterns. Breast
Cancer Res **2010**;12:R36
20. Pellacani D, Bilenky M, Kannan N, Heravi-Moussavi A, Knapp D, Gakkhar S, *et al.* Analysis of
Normal Human Mammary Epigenomes Reveals Cell-Specific Active Enhancer States and
Associated Transcription Factor Networks. Cell Rep **2016**;17:2060-74
21. Fu NY, Pal B, Chen Y, Jackling FC, Milevskiy M, Vaillant F, *et al.* Foxp1 Is Indispensable for
Ductal Morphogenesis and Controls the Exit of Mammary Stem Cells from Quiescence. Dev
Cell **2018**;47:629-44 e8
22. Nguyen QH, Pervolarakis N, Blake K, Ma D, Davis RT, James N, *et al.* Profiling human breast
epithelial cells using single cell RNA sequencing identifies cell diversity. Nat Commun
2018;9:2028
23. Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, *et al.*
JASPAR 2020: update of the open-access database of transcription factor binding profiles.
Nucleic Acids Res **2020**;48:D87-D92
24. Badouel C, Garg A, McNeill H. Herding Hippos: regulating growth in flies and man. Curr Opin
Cell Biol **2009**;21:837-43

25. Yin Y, Morgunova E, Jolma A, Kaasinen E, Sahu B, Khund-Sayeed S, *et al.* Impact of cytosine
methylation on DNA binding specificities of human transcription factors. Science **2017**;356
26. Stadler MB, Murr R, Burger L, Ivanek R, Lienert F, Scholer A, *et al.* DNA-binding factors shape
the mouse methylome at distal regulatory regions. Nature **2011**;480:490-5
27. Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, *et al.* Human DNA
methylomes at base resolution show widespread epigenomic differences. Nature **2009**;462:315-
22
28. Blackledge NP, Klose RJ. The molecular principles of gene regulation by Polycomb repressive
complexes. Nat Rev Mol Cell Biol **2021**;22:815-33
29. Beerman I, Bock C, Garrison BS, Smith ZD, Gu H, Meissner A, *et al.* Proliferation-dependent
alterations of the DNA methylation landscape underlie hematopoietic stem cell aging. Cell Stem
Cell **2013**;12:413-25
30. Sun D, Luo M, Jeong M, Rodriguez B, Xia Z, Hannah R, *et al.* Epigenomic profiling of young
and aged HSCs reveals concerted changes during aging that reinforce self-renewal. Cell Stem
Cell **2014**;14:673-88
31. Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus SJ, Weisenberger DJ, Shen H, *et al.*
Age-dependent DNA methylation of genes that are suppressed in stem cells is a hallmark of
cancer. Genome Res **2010**;20:440-6
32. Jung M, Pfeifer GP. Aging and DNA methylation. BMC Biol **2015**;13:7
33. Widschwendter M, Fiegl H, Egle D, Mueller-Holzner E, Spizzo G, Marth C, *et al.* Epigenetic
stem cell signature in cancer. Nat Genet **2007**;39:157-8
34. Hernando-Herraez I, Evano B, Stubbs T, Commere PH, Jan Bonder M, Clark S, *et al.* Ageing
affects DNA methylation drift and transcriptional cell-to-cell variability in mouse muscle stem
cells. Nat Commun **2019**;10:4361
35. Bahar R, Hartmann CH, Rodriguez KA, Denny AD, Busuttil RA, Dolle ME, *et al.* Increased
cell-to-cell variation in gene expression in ageing mouse heart. Nature **2006**;441:1011-4
36. Cheung P, Vallania F, Warsinske HC, Donato M, Schaffert S, Chang SE, *et al.* Single-Cell
Chromatin Modification Profiling Reveals Increased Epigenetic Variations with Aging. Cell
2018;173:1385-97 e14
37. Martinez-Jimenez CP, Eling N, Chen HC, Vallejos CA, Kolodziejczyk AA, Connor F, *et al.*
Aging increases cell-to-cell transcriptional variability upon immune stimulation. Science
2017;355:1433-6
38. Jenkinson G, Pujadas E, Goutsias J, Feinberg AP. Potential energy landscapes identify the
information-theoretic nature of the epigenome. Nat Genet **2017**;49:719-29
39. Brinkman AB, Nik-Zainal S, Simmer F, Rodriguez-Gonzalez FG, Smid M, Alexandrov LB, *et*
al. Partially methylated domains are hypervariable in breast cancer and fuel widespread CpG
island hypermethylation. Nat Commun **2019**;10:1749
40. Gunasekara CJ, Scott CA, Laritsky E, Baker MS, MacKay H, Duryea JD, *et al.* A genomic atlas
of systemic interindividual epigenetic variation in humans. Genome Biol **2019**;20:105
41. Edwards JR, Yarychkivska O, Boulard M, Bestor TH. DNA methylation and DNA
methyltransferases. Epigenetics Chromatin **2017**;10:23
42. Greenberg MVC, Bourc'his D. The diverse roles of DNA methylation in mammalian
development and disease. Nat Rev Mol Cell Biol **2019**;20:590-607
43. Burns KH, Boeke JD. Human transposon tectonics. Cell **2012**;149:740-52
44. Michalak EM, Nacerddine K, Pietersen A, Beuger V, Pawlitzky I, Cornelissen-Steijger P, *et al.*
Polycomb group gene Ezh2 regulates mammary gland morphogenesis and maintains the luminal
progenitor pool. Stem Cells **2013**;31:1910-20

45.   Xie M, Hong C, Zhang B, Lowdon RF, Xing X, Li D, *et al.* DNA hypomethylation within
672   specific transposable element families associates with tissue-specific enhancer landscape. Nat
673   Genet **2013**;45:836-41

674   46.   Pehrsson EC, Choudhary MNK, Sundaram V, Wang T. The epigenomic landscape of
675   transposable elements across normal human development and anatomy. Nat Commun
676   **2019**;10:5640

677   47.   Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, *et al.* Molecular portraits of
678   human breast tumours. Nature **2000**;406:747-52

679   48.   Jang HS, Shah NM, Du AY, Dailey ZZ, Pehrsson EC, Godoy PM, *et al.* Transposable elements
680   drive widespread expression of oncogenes in human cancers. Nat Genet **2019**;51:611-7

681   49.   Abba MC, Gong T, Lu Y, Lee J, Zhong Y, Lacunza E, *et al.* A Molecular Portrait of High-Grade
682   Ductal Carcinoma In Situ. Cancer Res **2015**;75:3980-90

683   50.   Yang WR, Ardeljan D, Pacyna CN, Payer LM, Burns KH. SQuIRE reveals locus-specific
684   regulation of interspersed repeat expression. Nucleic Acids Res **2019**;47:e27

685   51.   Franco HL, Nagari A, Malladi VS, Li W, Xi Y, Richardson D, *et al.* Enhancer transcription
686   reveals subtype-specific gene expression programs controlling breast cancer pathogenesis.
687   Genome Res **2018**;28:159-70

688   52.   Bell CG, Lowe R, Adams PD, Baccarelli AA, Beck S, Bell JT, *et al.* DNA methylation aging
689   clocks: challenges and recommendations. Genome Biol **2019**;20:249

690   53.   Kimmel JC, Penland L, Rubinstein ND, Hendrickson DG, Kelley DR, Rosenthal AZ. Murine
691   single-cell RNA-seq reveals cell-identity- and tissue-specific trajectories of aging. Genome Res
692   **2019**;29:2088-103

693   54.   Shalabi SF, Miyano M, Sayaman RW, Lopez JC, Jokela TA, Todhunter ME, *et al.* Evidence for
694   accelerated aging in mammary epithelia of women carrying germline BRCA1 or BRCA2
695   mutations. Nature Aging **2021**;1:838-49

696   55.   Vandiver AR, Irizarry RA, Hansen KD, Garza LA, Runarsson A, Li X, *et al.* Age and sun
697   exposure-related widespread genomic blocks of hypomethylation in nonmalignant skin. Genome
698   Biol **2015**;16:80

699   56.   Cole JJ, Robertson NA, Rather MI, Thomson JP, McBryan T, Sproul D, *et al.* Diverse
700   interventions that extend mouse lifespan suppress shared age-associated epigenetic changes at
701   critical gene regulatory regions. Genome Biol **2017**;18:58

702   57.   Issa JP. Aging and epigenetic drift: a vicious cycle. J Clin Invest **2014**;124:24-9

703   58.   Easwaran H, Johnstone SE, Van Neste L, Ohm J, Mosbruger T, Wang Q, *et al.* A DNA
704   hypermethylation module for the stem/progenitor cell signature of cancer. Genome Res
705   **2012**;22:837-49

706   59.   Burns KH. Transposable elements in cancer. Nat Rev Cancer **2017**;17:415-24

707   60.   Ishak CA, De Carvalho DD. Reactivation of Endogenous Retroelements in Cancer Development
708   and Therapy. Annual Review of Cancer Biology **2020**;4

709   61.   De Cecco M, Criscione SW, Peterson AL, Neretti N, Sedivy JM, Kreiling JA. Transposable
710   elements become active and mobile in the genomes of aging mammalian somatic tissues. Aging
711   (Albany NY) **2013**;5:867-83

712   62.   Benayoun BA, Pollina EA, Singh PP, Mahmoudi S, Harel I, Casey KM, *et al.* Remodeling of
713   epigenome and transcriptome landscapes with aging in mice reveals widespread induction of
714   inflammatory responses. Genome Res **2019**;29:697-709

715   63.   De Cecco M, Ito T, Petrashen AP, Elias AE, Skvir NJ, Criscione SW, *et al.* L1 drives IFN in
716   senescent cells and promotes age-associated inflammation. Nature **2019**;566:73-8

717   64.   Rodic N, Sharma R, Sharma R, Zampella J, Dai L, Taylor MS, *et al.* Long interspersed element-
718   1 protein expression is a hallmark of many human cancers. Am J Pathol **2014**;184:1280-6

65.   Rooney MS, Shukla SA, Wu CJ, Getz G, Hacohen N. Molecular and genetic properties of
tumors associated with local immune cytolytic activity. Cell **2015**;160:48-61
66.   Labarge MA, Garbe JC, Stampfer MR. Processing of human reduction mammoplasty and
mastectomy tissues for cell culture. J Vis Exp **2013**
67.   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics **2011**;27:1571-2
68.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biol **2009**;10:R25
69.   Song Q, Decato B, Hong EE, Zhou M, Fang F, Qu J, *et al.* A reference methylome database and
analysis pipeline to facilitate integrative and comparative epigenomics. PLoS One
2013;8:e81148
70.   Gaspar JM, Hart RP. DMRfinder: efficiently identifying differentially methylated regions from
MethylC-seq data. BMC Bioinformatics **2017**;18:528
71.   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, *et al.*
An improved ATAC-seq protocol reduces background and enables interrogation of frozen
tissues. Nat Methods **2017**;14:959-62

735
736

Figures

Figure 1: Mammary luminal epithelial cells from older individuals show DNA methylation changes at distinct genomic loci. (A) Methylation levels of identified DMRs across all samples. Samples are in rows, and DMRs are in columns. The color intensity represents the Z-score scaled from the mean. Values ranging from higher to lower methylation are shown as purple to green. (B) Kernel smoothed line plots of representative DMRs±500bp with methylation gain (left panel) and methylation loss (right panel). Blue and red lines respectively represent young and old samples. DMRs are highlighted in pink and black ticks mark each CpG position. (C) MSigDB gene sets enriched in genes with age dependent DNA methylation loss at promoters (TSS+/-2kb). (D) MSigDB gene sets enriched in genes with age dependent DNA methylation gain at promoters (TSS+/-2kb). Relevant gene sets are highlighted in green. C-D. UP: upregulated, DN: downregulated. (E) Genome browser tracks showing DMRs with methylation loss in old LEps at the *FOXP1* gene locus. (F) Genome browser tracks showing methylation gain at the CpG island promoter of *ALDH1A3* gene. E-F. DMR positions are shown as black boxes.

Figure 2: Differentially methylated regions show enrichment of lineage-specific transcription factor binding sites. (A) Top ten TF binding sites enriched in the DMRs that lose methylation with age are shown. Dots in the left panel show -log10(p-value) for the enrichment of the transcription factor binding site. Dots in the right panel show expression levels (RPKM) of the respective transcription factor in young myoepithelial (MEp) and luminal (LEp) cells. (B) Boxplot showing expression levels of myoepithelial-specific TF TP63 in young and old LEps (n=8, *p*-value < 0.01; two-tailed unpaired t test with Welch's correction). (C) Top ten TF binding sites enriched in the DMRs that gain methylation with age are shown. Dots in the left panel show -log10(p-value) for the enrichment of the TF binding site. Dots in the right panel show expression levels (RPKM) of the respective transcription factor in young myoepithelial (MEp) and luminal (LEp) cells. (D) Boxplot showing expression levels of luminal lineage-specific TF ELF5 in young and old LEps (n=8, *p*-value < 0.05; two-tailed Mann Whitney test). (E) Volcano plot showing expression levels of all TFs expressed in LEPs and MEPs (RPKM > 0.1). TFs with motifs enriched in DMRs with methylation loss and gain are highlighted in blue and red respectively. *P*-value < 0.05. (F) Schematic showing the role of myoepithelial-specific transcription factors in causing loss of methylation. Conversely, loss of expression of luminal-specific transcription factors leads to methylation of unbound sites and methylation gain.

795
Figure 3: Differentially methylated regions are enriched at lineage-specific regulatory elements.
(A) Genome browser tracks showing a representative example of a DMR with methylation loss
accompanied with increase in chromatin accessibility in old LEPs. DMRs are highlighted in green. Bottom
tracks show location of MEp-specific TF binding sites. (B) Volcano plot showing log2 fold change (log2fc)
of ATAC-seq peaks overlapping DMRs with methylation loss in old versus young LEPs. Positive log2 fold
change indicates increase in accessibility. (C) Genome browser tracks showing a representative example
of a DMR with methylation gain accompanied with decrease in chromatin accessibility in old LEPs. DMRs
are highlighted in green. Bottom tracks show location of LEp-specific TF binding sites. (D) Volcano plot
showing log2 fold change (log2fc) of ATAC-seq peaks overlapping DMRs with methylation gain in old
versus young LEPs. Negative log2 fold change indicates decrease in accessibility. (E) Percentage
number of DMRs with methylation loss and (F) methylation gain that overlap with chromHMM features of
ER⁻, ER⁺ luminal cells, and basal cells (rows) is shown. The color key indicates the various chromatin
states identified from ChIP-seq datasets.
809

Figure 4: Aging leads to stochastic methylation loss at transposable elements and methylation gain at CpG islands (CGIs). (A) Schematic depicting changes in entropy associated with stochastic methylation gain at CGIs. (B) Schematic depicting changes in entropy associated with stochastic methylation loss at fully methylated regions such as repeat elements. (C) Methylation and entropy levels at unmethylated CGIs that show increase in entropy. (D) Alu methylation and entropy levels at fully methylated Alu elements that show increase in entropy. C-D. Significant differences in distributions were tested using Kolmogorov–Smirnov test. (E) Aggregate profiles of methylation and entropy levels at unmethylated CGIs that show increase in entropy with age. Luminal breast cancer samples (red) show an increase in DNA methylation as well as entropy at these CGIs. (F) Aggregate profiles of methylation and entropy levels at unmethylated CGIs that show increase in entropy with age. Basal breast cancer samples (pink) do not show an increase in DNA methylation or entropy at these CGIs. (G) Expression of *FOXC1* in normal tissues and five PAM50 subtypes of breast tumors from the TCGA BRCA collection. Luminal A (LumA) versus Basal p-value < 2.2x10^{-16}; using two-tailed Wilcoxon signed-rank test. Luminal B (LumB) versus Basal (p-value < 2.2x10^{-16}; using two-tailed Wilcoxon signed-rank test).

Figure 5: Variably methylated regions with methylation loss are enriched at primate-specific transposable elements (A-D) Evolutionary age of (A) all TE elements (B) SINE, (C) LINE, and (D) LTR/ERV elements that are variably methylated with methylation loss or gain. Evolutionary ages of only the significantly enriched TE subfamilies are plotted. (E) TE subfamilies enriched in regions with variable methylation loss. Bars show observed/expected ratio for each TE subfamily. Expected number of overlaps in shuffled controls was used to calculate the ratio. Colors in the observed/expected graph highlight primate-specific or younger elements and grey bars indicate older elements. (F) Genome browser tracks showing a representative example of a HERVH-int element showing variable methylation loss in old samples. Bottom tracks are from ATAC-seq data showing gain of accessibility at the variably methylated region in the corresponding sample. (G) TF binding motifs enriched within each TE subfamily with variable methylation loss. Color scale is –log10(p-value) of TF motif enrichment.

Figure 6: Transposable elements with age-dependent methylation loss are activated in breast cancer. (A) A representative HERVH element showing variable loss of methylation in old samples and

in breast cancer (BC) samples. PAM50 subtypes and estrogen receptor (ER) expression status of the
tumors are shown for the BC WGBS tracks. Bottom tracks are RNA-seq tracks from DCIS samples
showing cryptic transcription of the HERVH element. All RNA-seq tracks are scaled to the same value
and PAM50 subtype is indicated on the right. (B) Heatmap showing signals across 74 TCGA-BRCA
samples for ATAC-seq peaks that overlap TEs hypomethylated with age. ATAC-seq peaks (rows) and
TCGA BRCA samples (columns) are split into 3 clusters each using k-means clustering. (C) Expression
of *ZNF92* in normal tissues and five PAM50 subtypes of breast tumors from the TCGA BRCA collection
(p-value Luminal A (LumA) versus Basal and Luminal B (LumB) versus Basal (p-value < 2.2×10^{-16}; using
two-tailed Wilcoxon signed-rank test).