Spatial Variation of Microtubule Depolymerization in Large Asters Suggests Regulation by MAP Depletion

Keisuke Ishihara[1,2,3,4], Franziska Decker[1,2,3,4], Paulo Caldas[5], James F. Pelletier[6,7,8], Martin Loose[5], Jan Brugués[1,2,3,4], Timothy J. Mitchison[6,7]

Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

Max Planck Institute for the Physics of Complex Systems, Dresden, Germany

Center for Systems Biology Dresden, Dresden, Germany

Cluster of Excellence Physics of Life, TU Dresden, Dresden, Germany

Institute of Science and Technology Austria, Klosterneuburg, Austria

Department of Systems Biology, Harvard Medical School, Boston, United States

Cell Division Group, Marine Biological Laboratory, Woods Hole, United States

Department of Physics, Massachusetts Institute of Technology, Cambridge, United States

Abstract

Microtubule plus end depolymerization rate is a potentially important target of physiological regulation, but it has been challenging to measure, so its role in spatial organization is poorly understood. Here we apply a method for tracking plus ends based on time difference imaging to measure depolymerization rates in large interphase asters growing in *Xenopus* egg extract. We observed strong spatial regulation of depolymerization rates, which were almost two-fold higher in the aster interior compared to the periphery, and much less regulation of polymerization or catastrophe rates. We interpret these data in terms of a limiting component model, where aster growth results in lower levels of soluble tubulin and MAPs in the interior cytosol compared to that at the periphery. The steady-state polymer fraction of tubulin was ~30%, so tubulin is not strongly depleted in the aster interior. We propose that the limiting component for microtubule assembly is a MAP that inhibits depolymerization, and that egg asters are tuned to low microtubule density.

Introduction

Microtubules undergo rapid polymerization dynamics in many cell types, as first revealed by polarization microscopy (Inoué and Sato 1967). Polymerization dynamics occur mostly on plus ends, and are well-approximated by a two-state dynamic model of dynamic instability (Dogterom and Leibler 1993). Plus end behavior is described by four parameters in this model: polymerization, depolymerization, catastrophe and rescue rates. For the robust organization of intracellular structures, these parameters are regulated in both space and time, which also allows for a rapid response to internal and external cues. Most recent studies on the biology and mechanisms of dynamics regulation focused on polymerization and catastrophe rates, in part because multiple factors are known to regulate these rates, and in part because they are the easiest to measure in cells. Tracking of the the plus-tip binding proteins EB1/3 allows measurement of polymerization rate with high reliability, and inference of catastrophe rate from comet lifetime with somewhat lower reliability. In contrast, depolymerization rates are harder to measure. Most studies reporting depolymerization rates focused on pure tubulin systems or on manual measurements of small numbers of microtubules at the periphery of living cells (Desai and Mitchison 1997; Brouhard and Rice 2018). As a result, we know rather little about regulation of depolymerization rates in complex microtubule structures such as the mitotic spindle or microtubule asters. Here, we apply imaging-based methods to systematically measure microtubule depolymerization rates in *Xenopus* egg extract, and report spatial regulation of depolymerization for the first time.

One important emergent property from the four dynamic instability factors, combined with nucleation, is the fraction of polymerized tubulin at steady state, which we will call the "polymer fraction". In a closed system like a cell, an increase in microtubule mass leads to depletion of soluble subunits which limits further assembly giving rise to a negative feedback of microtubule growth. When continuously supplied with GTP, the system comes to a steady state where the polymer fraction is approximately constant over time, and sufficient tubulin remains in solution to power robust polymerization of GTP-capped plus ends. The steady-state polymer fraction in interphase tissue culture cells was found to be 60-80% (Zhai and Borisy 1994; Kim, Peshkin, and Mitchison 2012). What determines this value, and its signficance for regulation, are largely unknown. Negative feedback from microtubule mass on further assembly in a closed system was conceptualized in the "component limitation" model for mitotic spindle size (Mitchison et al. 2015). A feedback of this kind is inevitable, but its precise mechanism is non-obvious. For pure tubulin, polymerization rate is a linear function of soluble dimer concentration (Desai and Mitchison 1997; Brouhard and Rice 2018). Thus, increased microtubule mass in a closed system decreases polymerization rate by subunit consumption. Catastrophe rate should increase in parallel, assuming it negatively correlates with polymerization rate as in most models (Walker et al. 1988; Gardner et al. 2008). Depolymerization is commonly considered to be a zero-order process whose rate is independent of soluble subunit concentration, though this assumption has been questioned (Gardner et al. 2011). In

physiological contexts, tubulin need not be the only limiting component in a closed system.
Microtubule-associated proteins (MAPs) that promote microtubule polymerization are also
depleted when microtubule mass increases, which could limit further polymerization. It is
unclear how MAP-limitation would regulate individual dynamic instability parameters.

Xenopus egg extract provides undiluted, metabolically-active cytoplasm that reconstitutes
physiological polymerization dynamics and is useful for investigating regulatory
mechanisms (Field and Mitchison 2018). It allows microscopy-based scoring of large
numbers of dynamic microtubules in a relatively homogeneous environment which is ideal
for quantitative analysis. Here, we focus on interphase asters, which model the egg-
spanning asters that position centrosomes, nuclei and cleavage furrows during early
divisions (Wühr et al. 2010). These asters grow to hundreds of microns in radius at a rate
of ~20 microns/min. EB1 tracking showed that the microtubules within them are relatively
short, ~16 microns on average. Their plus end dynamics are in the bounded regime of
dynamic instability, so they polymerize transiently, but are biased towards eventual
depolymerization with a half-life of ~1 min (Ishihara, Korolev, and Mitchison 2016). The
aster as a whole grows because each microtubule nucleates more than one daughter
microtubule during its lifetime by a poorly understood branching process. The density of
plus ends is approximately homogeneous within interphase asters, and there are two
interesting boundaries: the stationary MTOC and the growing aster periphery. The MTOC
nucleates continuously, and is thought to provide the information that directs microtubule
polarity. The periphery controls the rate of aster expansion and whether expansion is
bounded or unbounded. Plus ends at the periphery polymerize into a microtubule-free
environment, while those in the aster interior polymerize into an environment that is
dense in microtubules, and as we will show below, partially depleted of soluble subunits.
Many questions remain about these asters, including the polymer fraction, minus end
behavior, and whether dynamics differ between the interior and periphery. These
dynamical properties likely have important implications for aster mechanics and function.
Here, we unexpectedly observed that plus end depolymerization rate is subject to strong
spatial regulation. We interpret this observation in terms of a new model for regulation of
tubulin polymer fraction by MAP depletion.

Results

Polymerization rate increases slightly at the aster periphery

We first investigated the spatial variation of plus end polymerization and catastrophe rates
using standard EB1 comet tracking methods (Fig. 1A, Movie 1). We found that the
polymerization rate remained relatively constant at 30.9 µm/min (N=9348 tracks, SD 5.52,
SEM 0.06) as a function of distance from the center of asters (Fig. 1B). We detected
a ~10% increase in polymerization rate at the aster periphery 33.7 ± 10.9 µm/min (mean,
std) compared to the interior 30.0 ± 9.5 µm/min (mean, std). The plus end density is lower
at the periphery compared to the aster center (Fig. 1C), and resulted in a slight anti-
correlation between plus end density and polymerization rate (Fig. 1D). This anti-

correlation was much smaller than reported in an independent study (Geisterfer et al.
2019).

Next, we asked if the plus end catastrophe rate showed spatiotemporal variation using EB1
comet lifetimes as a proxy. The average EB1 comet persisted for 19 seconds which
corresponds to a catastrophe rate of 3.1 per minute. This value did not show much
variation as a function of location or plus end density (Fig. 1E). These measurements
performed with wide-field imaging were consistent with our previous measurements with
spinning disc confocal microscopy (Ishihara, Korolev, and Mitchison 2016). Therefore, we
expect our measurements to be sensitive to spatial variation in catastrophe rate. In
summary, our analysis of EB1 comet imaging demonstrated a modest 10% increase in
polymerization rate at the aster periphery, where local plus end density is relatively low,
and no measurable differences in catastrophe rate.

Fig. 1 *EB1 tracking-based measurements of microtubule plus end polymerization rate*
and catastrophe rate during aster growth. (A) Time lapse movies of EB1-GFP comets

are analyzed by particle tracking. Region in the white box is magnified on the right.
Purple spots show the location of individual EB1 comets in this frame, while the trailing
lines show the trajectory of the corresponding plus end for the preceding 10 seconds.
The velocity of EB1 comets report microtubule polymerization rate. (B) Polymerization
rate as a function radial distance from the center of the aster. N=9348 EB1 tracks are
represented as blue dots. Filled black circles and the shaded region indicate the mean
and the standard deviation of polymerization rate for each spatial bin. The estimated
mean difference in the interior (distance>300 μm) vs. periphery (50 μm<distance<280
μm) for t =1.5 min was 3.7 μm/min (95%CI=[2.4, 4.9]). (C) EB1 comet density as a
function radial distance. (D) Polymerization rate vs. EB1 comet density. Filled black
circles and error bars indicate the mean and the standard deviation of polymerization
rate for each EB1 density bin. The different colors correspond to the same time points
as in the previous panel. The estimated mean difference in the dense (0.066 1/μm^2)
vs. sparse (0.022 1/μm^2) was 3.8 μm/min (95%CI=[2.6, 5.0]). (E) Catastrophe rate is
calculated from the duration of EB1 tracks and plotted over radial distance. Filled black
circles and the shaded region indicate the mean and the corresponding 95% confidence
interval for the catastrophe rate for each spatial bin. Original data used for this analysis
is Movie 1. Scale bars 100 μm and 10 μm.

Spatial regulation of depolymerization rates

To measure microtubule depolymerization rates, we applied tubulin intensity difference analysis. This method relies on collecting high quality tubulin images at frequent intervals using TIRF microscopy, subtracting the intensity of sequential frames to generate time difference images, then applying a tracking algorithm. It allows tracking of polymerizing and depolymerizing ends even when most microtubules are in small bundles as is the case in egg extract asters. Our previous analysis of growing and shrinking ends used manual kymograph analysis of difference images and was restricted to the interior of the interphase asters (Ishihara et al. 2014). Here, we applied a newly developed automated workflow (Caldas et al. 2019; Caldas et al. 2020), and systematically compared spatial differences in polymerization dynamics (Fig. 2A). In brief, this method applies a low pass spatial filter to the difference images, followed by particle based tracking to quantify many microtubule ends. Using this workflow, the polymerization rate was found to have a mean of 32.0 ± 4.5 μm/min (mean, std) in the aster interior, and 32.0 ± 4.0 μm/min (mean, std) in the periphery (Fig. 2B, upper panel). These values of polymerization rates are consistent with EB1 comet tracking (Fig. 1). Polymerization rates exhibited a sharp, unimodal histogram, which overlapped for the interior and periphery with no statistical difference. In contrast, the depolymerization rate showed a striking difference in the aster interior vs. periphery. The depolymerization rate was 36.3 ± 7.9 μm/min (mean, std) in the aster interior, compared to 29.2 ± 8.9 μm/min (mean, std) at the aster periphery. The distribution of the depolymerization rates was spread out and had a positive skew in the aster interior, and a negative skew in the aster periphery, with some hint of a bimodal distribution. In summary, our analysis based on an improved intensity difference analysis showed that polymerization rates are similar in the aster periphery relative to the interior, while depolymerization rates show strong spatial variation and are faster inside asters.

Tubulin difference imaging revealed other useful information on polymerization dynamics.
Plus end polymerization and depolymerization rates were both variable, but over fairly
narrow ranges, so the data supported a two-state model of dynamic instability over
alternatives such as a biased random walk (Needleman et al. 2010). Our tracking was not
accurate enough to score rate fluctuations within bouts of depolymerization, and we did
not try to distinguish stochastic switching between states from more complex dynamics.
We observed very few outward-moving depolymerization events, suggesting that minus
ends are not depolymerizing, ie there is no significant treadmilling. We also saw very few
inward moving polymerization events, suggesting minus ends are static, and perhaps
capped.

Fig. 2 *Intensity difference-based measurement of polymerization and depolymerization*
rates in the interior and the periphery of microtubule asters. (A) TIRF microscopy
movies of fluorescent tubulin were subjected to intensity difference analysis, which
revealed a spatial map of polymerization and depolymerization for the interior (less than
100 μm away from the MTOC) and peripheral (more than 200 μm from the MTOC with
few microtubules) region of microtubule asters. These images correspond to Movies 2
and 3. (B) The distributions of polymerization and depolymerization rates measured
from N=6 interior movies and N=6 peripheral movies. The polymerization rate (median
32.0 vs. 32.0 μm/min) showed no statistical difference in the interior vs. periphery (p-
value = 0.819, t-test). The depolymerization rate (median 37.6 vs. 26.7 μm/min) was
higher in the interior (p-value = 4.44e-16, two sample Kolmogorov-Smirnov test) with
estimated difference of mean and median depolymerization rates of 7.1 μm/min (95% CI
= [5.9, 8.3]) and 11.0 μm/min (95% CI = [8.4, 12.4]). Scale bar, 20 μm.

Laser ablation induced depolymerization rates

In the conventional model of dynamic instability, growing plus ends lose their structural stability prior to the switch from polymerization to depolymerization. To ask if such physiological catastrophe events are required for the difference in depolymerization rates, we used femtosecond laser ablation to artificially induce depolymerization. Ablation along a line normal to the microtubule axis triggers a spatially defined, synchronous wave of depolymerization (Decker et al. 2018; Decker and Brugués 2015). Using circular patterns , we ablated microtubules at a fixed distance from the center of asters (Fig. 3A, Movie 4). Intensity difference analysis (Fig. 3B) revealed a resulting wave of depolymerization that moved inward at a constant rate (Fig. 3C). This single inward wave is consistent with the fact that (a) the majority of microtubules are oriented with plus ends outward, (b) cut plus ends immediately depolymerize, and (c) newly formed minus ends are stable. By applying this laser ablation assay to multiple asters, we measured an average depolymerization wave velocity of 33.5 ± 9.4 (31.5) um/min (mean, std, median), which was comparable to the modal value of the depolymerization rates in the aster interior (Fig. 2). We found no correlation between the rate of depolymerization wave and the distance at which the ablation was induced (15-56 μm). We were not able to reliably cut and measure at the aster periphery due to the low microtubule density. Overall, these observations suggest that depolymerization dynamics are similar for plus ends following a natural catastrophe vs. ablation in the aster interior.

Laser ablation experiments offer insights into the organization of microtubule structures (Brugués et al. 2012; Decker et al. 2018). We found that the strength of the depolymerization wave (i.e. the area under the curves of the summed differential intensities) decreased with progression (Fig. 3E). This is explained by the fact that depolymerization is halted when individual plus ends reach their corresponding minus ends. Thus, this observation provides evidence that minus ends exist throughout the aster, and confirms our previous proposal that asters are built of short microtubules (Ishihara, Korolev, and Mitchison 2016).

Fig. 3 *Measurement of depolymerization rates following laser ablation. (A) An interphase aster was subjected to a circular laser cut with radius 56 µm (left), which induced a wave of microtubule depolymerization Movie 4. The resulting movie was used to construct a differential intensity movie (right and Movie 4). Dotted wedge region indicates region excluded from image analysis. (B) Radial sum of differential intensities at different time points (from dark to light blue) of the same laser cut experiment. The area under each curve equals the mass of microtubules depolymerized per time interval of 5 s. Vertical dotted line indicates the location of the laser cut. (C) For each cut, the peak position of the differential intensity travels at constant speed. (D) Depolymerization rates measured from N = 16 laser cuts positioned at 15-56 µm from the center of the aster. (E) The area under differential intensity curves decreases as the depolymerization wave travels inward. This example corresponds to the cut depicted in panel A-C. Vertical dotted line indicates the location of the laser cut.*

Aster growth results in spatial gradients of soluble tubulin and EB1

Spatial regulation of depolymerization rate could be due to complex biochemical schemes, such as activation of a dynamics-regulating kinase in the aster interior. We cannot rule out this kind of hypothesis, but we prefer an interpretation based on component limitation, in part because this is a mechanistically simpler hypothesis, and in part because component limitation is inevitable in a closed system. Component limitation is a relevant consideration when comparing microtubule dynamics between the inside and periphery of large asters because aster cytosol is not well mixed by diffusion on relevant time and length scales. Specifically, the aster grows ballistically into fresh cytosol outside the aster much faster

than its interior can diffusively equilibrate with that fresh cytosol. A simple Péclet number
calculation serves to make this point. Consider the dimensionless Péclet number: Pe =
$\frac{\text{advective transport}}{\text{diffusive transport}} = \frac{Lu}{D}$. While a typical protein molecule is transported in a diffusive process
with cytoplasmic diffusion coefficient $D = 360 \frac{\mu m^2}{min}$ (Salmon et al. 1984), aster growth is a
ballistic or advective process, i.e. radius increases linearly with time, with a velocity $u =$
$20 \frac{\mu m}{min}$ (Ishihara et al. 2014). Given a relevant length scale L, we evaluate the magnitude of
the Péclet number. The extremes are: Pe>>1, peripheral microtubules grow so fast that
they will always grow into fresh cytoplasm with plenty of available material. Pe<<1, aster
growth is limited by the diffusive transport of material to the aster periphery. At a depth
of $L = 200$ μm from the aster periphery, which is typical for our internal measurements, Pe
= 11, so diffusion makes little contribution to supplying subunits. Thus, we expect the aster
interior to effectively constitute a closed system, shut off from fresh cytosol supplied by
diffusion from the outwardly growing aster boundary.

To seek evidence for depletion of soluble subunits by growing asters we imaged
fluorescently labeled tubulin and analyzed its intensity profile as a proxy of local tubulin
concentration, summing both polymer and soluble forms (Fig. 4A). The center of asters
showed the highest signal, reflecting the highest microtubule polymer density. In addition,
we noticed an annular region at the aster periphery with ~4% lower value of fluorescence
signal relative to the background cytosol levels (Fig. 4B). We interpret this as evidence for
depletion of soib el tubulin by aster growth. Considering the potential contribution of
background signal, we interpret the 4% decrease of tubulin intensity as a lower bound for
the degree of depletion. The EB1-GFP intensity profile showed a similar spatial pattern,
suggesting that microtubule associated proteins may also be depleted to some degree.
Finally, we compared the relative position of tubulin/EB1 intensity profiles to that of the
EB1 comet density profile (Fig. 4B). This suggested that even the most peripheral plus ends
exist in a cytosolic environment that is 4% (or more) depleted of tubulin and EB1
compared to unconsumed cytosol. The half-width of the depleted zone extended ~50
microns beyond the growing aster periphery, approximately consistent with our Péclet
number estimate. This analysis indicated that soluble protein levels may vary within
growing asters due to subunit consumption.

Fig. 4 *Fluorescent intensity profiles of tubulin and EB1 indicate spatial variation of these species within growing interphase asters. (A) Wide field images of a growing aster visualized with Alexa647-labelled tubulin and EB1-GFP. For both images, the contrast is adjusted to emphasize the zone of low fluorescence intensity. (B) Corresponding quantification of the fluorescence intensity profiles averaged over the quadrant and normalized to the intensity outside the aster (top) and EB1 comet density profile (bottom). We define the aster as the region that has EB1 density higher than the background level + standard deviation. Scale bar, 100 μm.*

Discussion

Limiting component hypothesis for the spatial regulation of depolymerization rate

The initial motivation of this study was to test our previous assumption that microtubule dynamics was spatially homogenous during aster growth (Ishihara, Korolev, and Mitchison 2016). EB1 tracking experiments showed that the catastrophe rate was largely constant over hundreds of microns (Fig. 1), and the polymerization rate was ~10% higher in the periphery than the interior. Intensity difference imaging showed that the depolymerization rate varied from modal values of ~20 μm/min at the aster periphery to ~40 μm/min in the interior (Fig. 2). Average values were closer, 29 vs. 36 μm/min, but the effect size was evident. This may be the first report of systematic spatial variation in this parameter of dynamic instability.

Multiple hypotheses could be considered to account for the observed spatial regulation of depolymerization rates. We will focus on component depletion models because this is the

simplest hypothesis; the growth of structures in confined cytoplasm necessarily consumes
the available building blocks and limits further growth (see Introduction). Consistent with
such a model, we found evidence for depletion of soluble tubulin and EB1 inside asters by
imaging (Fig. 4) and a Péclet number calculation showed that the cytosol inside and
outside growing asters do not mix by diffusion once the distance between them is larger
than ~50 microns.

The first candidate for a limiting component in a microtubule system is tubulin itself. As
tubulin concentration decreases, the literature predicts that plus ends should grow more
slowly and catastrophe more often (see Introduction). However, we observed little spatial
variation in these parameters, though EB1-measured growth rates trended in the expected
direction (Fig. 1). One reason for a lack of effect of depletion on polymerization rate may be
that the steady state polymer fraction in our system is relatively low, ~30% (Methods,
Tubulin polymer fraction estimate), which is the upper bound for difference in soluble
tubulin concentration inside and outside asters. Given this small difference and lack of
literature on how soluble tubulin levels regulate depolymerization, we explore alternative
candidates for the limiting component that is responsible for the spatial regulation of
depolymerization rate.

Several MAPs have been shown to slow down depolymerization rates (Drechsel et al. 1992;
Andersen and Karsenti 1997). We propose that one or mores MAPs that are present at low
concentrations relative to microtubule polymer binds to microtubules at lower density in
the interior compared to the periphery, resulting in faster depolymerization in the interior
(Fig. 5A). The key parameters that determines whether a given MAP can cause spatial
regulation in this way are concentration in cytosol and affinity to the microtubule lattice,
which together determine whether the MAP is significantly depleted by microtubule
binding. We present a toy model to illustrate how these parameters may help us predict
which MAPs regulate depolymerization (see Method, A model of single MAP species
binding to microtubule lattice). The model predicts the fraction of freely diffusing MAP
molecules $\frac{M_{\text{free}}}{M_{\text{total}}}$ and the site occupancy of the microtubule lattice $\theta = \frac{M_{\text{bound}}}{P}$, both of
which decrease with microtubule polymer concentration P as expected (see Fig. 5B). For a
hypothetical MAP whose abundance is 0.2 µM and associates with microtubules with a
relatively tight dissociation constant of 100 nM, the fractional occupancy of the
microtubule lattice decreases from 0.67 to 0.12 when the polymer concentration increases
from 0 to 1.6 µM. We argue that such 5.6-fold difference in microtubule lattice occupancy
may be sufficient to cause differences in depolymerization rate that account for spatial
regulation. Soluble tubulin, in contrast, only decreases by 0.7-fold. In Table 1, we
summarize how different MAPs present in the frog egg are predicted to change their lattice
occupancies. Based on the predicted fold change lattice occupancies, we favor EML4, MAP7
and its homologs MAP7D1 and MAP7D2, and MAP1B as candidate regulators of
depolymerization that are subject to component limitation effects in large interphase
asters, leading to faster depolymerization in the interior.

Fig. 5 *Proposed model for the regulation of depolymerization rate by a MAP species as the limiting component. (A) We consider the equilibrium of a single MAP species that associates with the microtubule lattice with dissociation constant $K_d = \frac{k_2}{k_1}$. We hypothesize that the degree at which a microtubule is bound by this MAP species slows down depolymerization rate. This effect is greater in the aster periphery, where the microtubule density is lower. (B) Fraction of free MAP and occupied microtubule lattice as a function of microtubule concentration for a hypothetical MAP present at 0.2 μM.*

Gene Symbol	Protein Description	abundance [nM]	Kd [nM]	reference	theta at low P	theta at P = 1.6 μM	theta fold change
MAP4	Microtubule-associated protein, XMAP230	964.5	300	Tokuraku et al. JBC 2003, Andersen et al. JCB 1994	0.76	0.45	1.69
EML2	Echinoderm microtubule-associated protein-like 2	578.18	?				
EML4	EML4 protein	504.2	180	Eichenmuller et al. Cytoskeleton 2001	0.74	0.27	2.74
EML1	Echinoderm microtubule-associated protein-like 1	135.67	?				
MAP7	Ensconsin	91.3	460	Monroy et al. Nat. Comm. 2018	0.17	0.04	4.25
MAP1B	Microtubule-associated protein 1B, XMAP310	84.68	200	Andersen et al. JCB 1997	0.3	0.05	6.0
MAP1S	MAP1S light chain	66.04	?				

| MAP7D1 | MAP7 domain-containing protein 1 | 46.97 | ? | | | | |
| MAP7D2 | MAP7 domain containing 2 protein varaint 2 (Fragment) | 8.55 | ? | | | | |

Table 1 Summary of microtubule associated proteins (MAPs) that may slow down plus
end depolymerization rate. Protein abundance in the frog egg is from proteome data
in (Wühr et al. 2014). For those with reported dissociation constants (Eichenmüller et al.
2001; Andersen and Karsenti 1997; Andersen et al. 1994; Monroy et al. 2018; Tokuraku
et al. 2003), we provide the estimated value of θ, the fractional occupancy of the
microtubule lattice at low (i.e. excess MAP) and high polymer concentrations (P = 1.6
µM).

Our proposal that a spatially varying MAP:microtubule ratio regulates depolymerization
comes with several limitations. First, while a single, low-abundance, tight binding MAP was
considered in our model, multiple MAPs may reglated depolymerization rate in eggs. The
total abundance of all MAP species easily exceeds 2 µM. Thus, in a more realistic scenario,
multiple MAP species may compete for microtubule binding and their order of association
is governed by their abundances and affinities. Kinases such as Cdk1, MARK, and Nek may
regulate the affinity of these MAPs (Adib et al. 2018; Ookata et al. 1997; Chang et al. 2001;
Drewes et al. 1992; Illenberger et al. 1996; Drewes et al. 1997). Our current model is a
useful starting point for future studies that consider complex multi-species dynamics of
cytoskeletal self-organization including the mitotic spindle (Hazel et al. 2013; Good et al.
2013; Rieckhoff, Ishihara, and Brugués 2019; Rieckhoff et al. 2020).

Second, while most MAPs that slow down depolymerization have been reported to
accelerate polymerization in minimal conditions (Drechsel et al. 1992; Andersen et al.
1994; Andersen and Karsenti 1997), we observed little regulation of polymerization rate in
our aster reactions. We speculate that, in a cellular context where the synergistic effects of
XMAP215 and EB1 play a prominent role in modulating plus end dynamics (Kinoshita et al.
2001; Zanic et al. 2013), our hypothetical MAP may not further accelerate polymerization.
It will be interesting to carefully revisit depolymerization rates in other cellular and
reconstitution systems.

Implications of low tubulin polymer fraction for aster mechanics

Finally, what are the implications of our observations for the biology of interphase asters in
large eggs? Asters function to separate centrosomes and nuclei at the end of mitosis, and to
position cleavage furrows, both of which require that they grow to egg-spanning
size. Previously, we modeled aster growth from the interplay of polymerization dynamics
and autocatalytic nucleation at the periphery (Ishihara, Korolev, and Mitchison 2016). To
prevent the uncontrolled exponential growth expected from an autocatalytic process, we
had to assume negative feedback from growth to further assembly, which we implemented

as a logistic function that made no mechanistic assumptions. Our current findings suggest
that this feedback occurs mainly by increased depolymerization rate in the aster interior,
and that it leads to a steady state with a relatively low polymer fraction, estimated as ~30%
tubulin polymerized. This low polymer density may be important for how dynein exert
forces for aster positioning (Wühr et al. 2010). Egg asters are gels comprising a network of
relatively short microtubules entangled with F-actin and organelles (Pelletier et al. 2020).
Microtubules are the stiffest component of this composite gel, so the bulk stiffness of the
aster is likely to depend strongly on the tubulin polymer fraction. We speculate that a low
polymer fraction allows the aster gel to deform in response to forces from dynein and the
ingressing furrow while maintaining sufficient connectivity to transmit forces across large
length scales.

Materials and Methods

Aster Reconstitution in Xenopus Egg Extract

396 Interphase microtubule asters were reconstituted in Xenopus egg extract with anti-AurkA
397 coated beads as microtubule organizing centers (Field, Pelletier, and Mitchison 2017; Field
398 et al. 2014). All reactions were supplemented with 0.04 mg/mL p150-CC1 fragment of
399 dynactin to inhibit dynein motor activity. Fluorescently labelled tubulin or EB1-GFP was
400 used to visualize microtubules.

Measurement of polymerization and catastrophe rates from EB1 comets

Wide-field images were acquired on a Nikon Eclipse Ni-E upright microscope equipped
with a CFI Plan Apochromat Lambda 20× 0.75 N.A. objective lens (Nikon), Nikon motorized
XY stage, and Hamamatsu ORCA-Flash4.0 LT scientific CMOS camera, driven by NIS-
Elements. EB1-GFP was supplemented to the reactions at a final concentration of 200-400
nM. EB1 comets were imaged at 2 second intervals, and analyzed with TrackMate (Tinevez
et al. 2017). For identifying spots, the LoG detector was used with with 2.0 micron spot
diameter, threshold 10.0, no median filter, sub-pixel localization enabled. For linking
tracks, the LAP tracker was applied with max search radius of 3 micron without gap
closing. The results were further analyzed by a custom script written in Python to calculate
the polymerization velocity (average of the frame-to-frame velocity for each track) and the
catastrophe rate (fitting an exponential function to the comet duration distribution). The
DABEST-Python package was used for statistics and effect size estimation (Ho et al. 2019).

Measurement of polymerization and depolymerization rates of microtubule ends

417 Physiological microtubule polymerization and depolymerization rates were measured by
418 TIRF microscopy and tubulin intensity difference analysis. Briefly, asters labeled with
419 Alexa647-labelled bovine tubulin were assembled under K-casein coated coverslips and

imaged with TIRF microscopy (Ishihara et al. 2014). The Nikon Ti-E motorized inverted microscope was equipped with a Nikon motorized TIRF illuminator, Perfect Focus, a Prior Proscan II motorized stage, Agilent MLC400B laser launch (488 nm, 561 nm, 647 nm), and an Andor DU-897 EM-CCD camera. Alexa 647-labeled bovine tubulin was imaged with a 100× CFI Apo 1.49 N.A. TIRF objective lens (Nikon) with or without 1.5x Optovar. Stream acquisition of images at 500-ms intervals was performed using the RAM capture mode in NIS-Elements software (Nikon Instruments). We defined the aster interior as FOVs with the artificial MTOC in the corner (=center of the FOV less than 100 microns from the center of the aster), while peripheral regions were chosen as the leading edge of the growing aster with very few microtubules in the field of view (=center of the FOV at least 200 microns from the center of the aster).

To obtain spatial information regarding polymerization and depolymerization rates, we applied a recently developed workflow used to quantify treadmilling dynamics in bacterial cytoskeletal filaments (Caldas et al. 2019; Caldas et al. 2020). This method allows to track hundreds of growing and shrinking microtubule ends in an automated fashion, overcoming the limitations of a standard manual kymograph analysis. We first constructed differential time-lapse movies by subtracting the intensity differences between each two consecutive frames. This image-processing step removes static objects from the movie while regions containing intensity fluctuations give rise to either a positive or negative signal (speckles), which corresponds to growth (polymerization) or shrinkage (depolymerization) rates, respectively. Next, the particle-tracking software for ImageJ, TrackMate (Tinevez et al. 2017), was used to automatically detect and follow the resultant moving speckles to rebuild trajectories for analysis. Finally, a custom python script is used to quantify densities and rates of all the detect spots along their trajectories. For TrackMate, we used the LoG (Laplacian Gaussian) detector with an estimated diameter of 1 μm to detect the moving spots. To discard potential false positives, we considered only particles with a signal-to-noise ratio lower than 0.8 and a track displacement distance larger than 1 μm. To build the final trajectories, we used the "Simple LAP tracker" with a "Max Linking Distance" of 1 μm, a "Maximal gap-closing distance" of 1 μm and "Max frame Gap" of 5 frames. Later, we only considered for analysis trajectories longer than 5 sec. To construct the distributions in Fig. 2B, we pooled the frame-to-frame velocities for each track without averaging. The DABEST-Python package was used for statistics and effect size estimation (Ho et al. 2019).

Measurement of laser cutting-induced depolymerization rates

During the cutting experiments, interphase asters were labeled with Atto565 porcine tubulin and imaged using a Nikon microscope (Ti Eclipse) with Yokogawa CSU-X1 confocal spinning disk, an EMCCD camera (Andor iXon DU-888), a 60x 1.2 NA water immersion objective, and the software AndorIQ for image acquisition. Laser cutting and image analysis were done as described in (Decker et al. 2018) with the only difference that the laser pulse repetition rate was reduced by a pulse picker (APE pulseSelect) from 80 MHz to 20 kHz for some of the cuts, which reduced the probability of destroying the asters. In brief, the sample stage was moved according to the desired cut shape in multiple z planes to reach a cut depth of ~2 μm around the focal plane. The resulting inward traveling microtubule

depolymerization wave was recorded at 2-5 frames/s. The measurement of the polymerization speed involved calculating differential intensities from the raw images with a time difference of 2-3 s (see Fig. 3, Movie 4). These differential intensities were integrated with respect to the radius leading to a Gaussian-shaped integrated intensity profile plotted over the radius for each time point. Using the peak position of fitted Gaussians to these profiles, we determined the distance that the depolymerization wave travelled over time. The slope of this travelled distance over time plot equals the depolymerization speed of the cut microtubules, which was constant over the entire time that the wave was visible (in agreement with depolymerization in metaphase spindles and monopoles (Decker et al. 2018)).

Fluorescence intensity-based estimation of local depletion of tubulin and EB1

Wide-field images were acquired on a Nikon Eclipse Ni-E upright microscope equipped with a CFI Plan Apochromat Lambda 20× 0.75 N.A. objective lens (Nikon), Nikon motorized XY stage, and Andor Zyla 4.2 Plus scientific CMOS camera, driven by NIS-Elements. Alexa647-labelled bovine tubulin and EB1-GFP were supplemented to the reactions. Dark current subtraction and flat field correction were applied to both channels. A custom script written in Matlab was used to quantify the fluorescent intensity as a function of radial distance from the MTOC (Pelletier et al. 2020). Spot detection of EB1 comets was performed with TrackMate as described above.

Tubulin polymer fraction estimate

To estimate the fraction of tubulin that is in polymer state, we performed a simple calculation based on our measurements. Specifically, we calculated the total concentration of microtubule polymer in the interior of our aster reactions:

$$\left[\mathrm{MT_{polymer}}\right] = \frac{\rho \cdot L}{p \cdot N_A}$$

where

ρ: number of microtubules per volume [#/μm^3]

L: average length of a microtubule – 16 μm (Ishihara 2016)

p: pitch of tubulin dimers – 8 nm/13 heterodimers

N_A: Avogadro's number – 6*10^23 heterodimers/mole

By analyzing a z-stack of EB1-GFP images, we measured the thickness of a reaction prepared by squashing 4 μl of extract under a 18x18 mm cover slip as ~10 micron. Further, we found that a single focal plane of a 20x 0.75 NA lens captures as much as 30-40% of all EB1 comets that are axially distributed in such reaction. Thus, we estimate the microtubule density from $\rho = \frac{\sigma}{d\phi} \cdot \frac{f_{\mathrm{cat}}+f_{\mathrm{res}}}{f_{\mathrm{res}}}$.

σ: number of microtubule per area in a single focal plane – 0.08 per μm^2

d: Thickness of the cover slip reaction – 10 μm

ϕ: fraction of EB1 comet detected in a single focal plane - 0.35

f_{cat}: catastrophe rate - 3.3 [1/min] (Ishihara, Korolev, and Mitchison 2016)

f_{res}: rescue rate - 2.0 [1/min] (Ishihara, Korolev, and Mitchison 2016)

The ratio of catastrophe and rescue rates allow us to account for the shrinking plus ends,
which are not detected by EB1 comets. We estimate that asters contain [polymer] = 2.65
μM of polymerized tubulin. Comparing this value to abundance of tubulin heterodimers in
the frog egg, 8.6 μM (Wühr et al. 2015), we estimate that ∼31 % of tubulin is in the polymer
form.

A model of single MAP species binding to microtubule lattice

509   We consider a single MAP species M (total concentration M_{total}) that binds and unbinds
510   from the microtubule lattice (total concentration P). Let S denote the concentration of free
511   lattice sites and k_1 and k_2 denote the binding and unbinding rates, respectively.

512
$$M_{free} + S \rightleftharpoons M_{bound}$$

513   Conservation of MAP species: $M_{total} = M_{free} + M_{bound}$

514   Conservation of microtubule lattice sites: $P = S + M_{bound}$

515   The equation for mass action kinetics is:

516
$$\frac{dM_{bound}}{dt} = k_1 M_{free} S - k_2 M_{bound} \,.$$

517   Solving for the steady state solution $\frac{dM_{bound}}{dt} = 0$, we find:

518
$$M_{bound} = \frac{1}{2}\left(P + M_{total} + K_D - \sqrt{(P + M_{total} + K_D)^2 - 4PM_{total}}\right)$$

519   This allows us to calculate, (fraction free MAP) $= \frac{M_{free}}{M_{total}} = 1 - \frac{M_{bound}}{M_{total}}$

520   and (fractional lattice occupancy) $= \theta = \frac{M_{bound}}{P}$.

Author Contributions

KI and TJM designed the research and wrote the manuscript with help from all
authors. TJM acquired the data and KI performed the analysis for EB1 comets. KI, PC, and
ML performed the tubulin intensity difference analysis. KI, FD, and JFB performed the laser
ablation experiments and analyzed the data. KI and JFP performed the analysis to infer

local protein depletion from wide field microscopy images. KI developed the theoretical
model.

Acknowledgements

The authors thank the members of Mitchison, Brugues, and Jay Gatlin groups (Uni. of
Wyoming) for discussions. We thank Heino Andreas (MPI-CBG) for frog maintenance. We
thank Nikon Inc. for microscopy support at MBL. KI was supported by fellowships from the
Honjo International Scholarship Foundation and Center of Systems Biology Dresden. FD
was supported by the DIGGS-BB fellowship provided by the DFG. PC is supported by a
Boehringer Ingelheim Fonds (BIF) PhD fellowship. JFP was supported by a fellowship from
the Fannie and John Hertz Foundation. ML's research is supported by a European Research
Council (ERC) grant ERC-2015-StG-679239. JB's research is supported by the Human
Frontiers Science Program (CDA00074/2014). TJM's research is supported by NIH grant
GM39565 and by MBL summer fellowships.

Competing interests

540   All the authors declare no competing interests.

References

Inoué, S, and H Sato. 1967. "Cell Motility by Labile Association of Molecules. The Nature of
Mitotic Spindle Fibers and Their Role in Chromosome Movement.". *J Gen Physiol* 50:
Suppl:259–92.

Dogterom, M, and S Leibler. 1993. "Physical Aspects of the Growth and Regulation of
Microtubule Structures.". *Phys Rev Lett* 70: 1347–50.

Desai, A, and TJ Mitchison. 1997. "Microtubule Polymerization Dynamics.". *Annu Rev Cell
Dev Biol* 13: 83–117.

Brouhard, GJ, and LM Rice. 2018. "Microtubule Dynamics: an Interplay of Biochemistry and
Mechanics.". *Nat Rev Mol Cell Biol* 19: 451–63.

Zhai, Y, and GG Borisy. 1994. "Quantitative Determination of the Proportion of Microtubule
Polymer Present during the Mitosis-Interphase Transition.". *J Cell Sci* 107 (Pt 4): 881–90.

Kim, S, L Peshkin, and TJ Mitchison. 2012. "Vascular Disrupting Agent Drug Classes Differ in
Effects on the Cytoskeleton.". *PLoS One* 7: e40177.

Mitchison, TJ, K Ishihara, P Nguyen, and M Wühr. 2015. "Size Scaling of Microtubule
Assemblies in Early Xenopus Embryos.". *Cold Spring Harb Perspect Biol* 7: a019182.

Walker, RA, ET O'Brien, NK Pryer, MF Soboeiro, WA Voter, HP Erickson, and ED Salmon. 1988. "Dynamic Instability of Individual Microtubules Analyzed by Video Light Microscopy: Rate Constants and Transition Frequencies.". *J Cell Biol* 107: 1437–48.

Gardner, MK, AJ Hunt, HV Goodson, and DJ Odde. 2008. "Microtubule Assembly Dynamics: New Insights at the Nanoscale.". *Curr Opin Cell Biol* 20: 64–70.

Gardner, MK, BD Charlebois, IM Jánosi, J Howard, AJ Hunt, and DJ Odde. 2011. "Rapid Microtubule Self-Assembly Kinetics.". *Cell* 146: 582–92.

Field, CM, and TJ Mitchison. 2018. "Assembly of Spindles and Asters in <i>Xenopus</i> Egg Extracts.". *Cold Spring Harb Protoc* 2018.

Wühr, M, ES Tan, SK Parker, HW 3rd Detrich, and TJ Mitchison. 2010. "A Model for Cleavage Plane Determination in Early Amphibian and Fish Embryos.". *Curr Biol* 20: 2040–45.

Ishihara, K, KS Korolev, and TJ Mitchison. 2016. "Physical Basis of Large Microtubule Aster Growth.". *Elife* 5.

Geisterfer, Zachary M., Daniel Zhu, Timothy Mitchison, John Oakey, and Jesse C. Gatlin. 2019. "Microtubule Growth Rates Are Sensitive to Global and Local Changes in Microtubule plus-End Density", November. https://doi.org/10.1101/849190.

Ishihara, K, PA Nguyen, AC Groen, CM Field, and TJ Mitchison. 2014. "Microtubule Nucleation Remote from Centrosomes May Explain How Asters Span Large Cells.". *Proc Natl Acad Sci U S A* 111: 17715–22.

Caldas, P, M López-Pelegrín, DJG Pearce, NB Budanur, J Brugués, and M Loose. 2019. "Cooperative Ordering of Treadmilling Filaments in Cytoskeletal Networks of FtsZ and Its Crosslinker ZapA.". *Nat Commun* 10: 5744.

Caldas, Paulo, Philipp Radler, Christoph Sommer, and Martin Loose. 2020. "Computational Analysis of Filament Polymerization Dynamics in Cytoskeletal Networks". In *Methods in Cell Biology*. Elsevier. https://doi.org/10.1016/bs.mcb.2020.01.006.

Needleman, DJ, A Groen, R Ohi, T Maresca, L Mirny, and T Mitchison. 2010. "Fast Microtubule Dynamics in Meiotic Spindles Measured by Single Molecule Imaging: Evidence That the Spindle Environment Does Not Stabilize Microtubules.". *Mol Biol Cell* 21: 323–33.

Decker, F, D Oriola, B Dalton, and J Brugués. 2018. "Autocatalytic Microtubule Nucleation Determines the Size and Mass of <i>Xenopus Laevis</i> Egg Extract Spindles.". *Elife* 7.

Decker, F, and J Brugués. 2015. "Dissecting Microtubule Structures by Laser Ablation.". *Methods Cell Biol* 125: 61–75.

Brugués, J, V Nuzzo, E Mazur, and DJ Needleman. 2012. "Nucleation and Transport Organize Microtubules in Metaphase Spindles.". *Cell* 149: 554–64.

Salmon, ED, WM Saxton, RJ Leslie, ML Karow, and JR McIntosh. 1984. "Diffusion Coefficient of Fluorescein-Labeled Tubulin in the Cytoplasm of Embryonic Cells of a Sea Urchin: Video

Image Analysis of Fluorescence Redistribution after Photobleaching.". *J Cell Biol* 99: 2157–
64.

Drechsel, DN, AA Hyman, MH Cobb, and MW Kirschner. 1992. "Modulation of the Dynamic
Instability of Tubulin Assembly by the Microtubule-Associated Protein Tau.". *Mol Biol Cell*
3: 1141–54.

Andersen, SS, and E Karsenti. 1997. "XMAP310: a Xenopus Rescue-Promoting Factor
Localized to the Mitotic Spindle.". *J Cell Biol* 139: 975–83.

Wühr, M, RM Jr Freeman, M Presler, ME Horb, L Peshkin, S Gygi, and MW Kirschner. 2014.
"Deep Proteomics of the Xenopus Laevis Egg Using an MRNA-Derived Reference Database.".
Curr Biol 24: 1467–75.

Eichenmüller, B, DP Ahrens, Q Li, and KA Suprenant. 2001. "Saturable Binding of the
Echinoderm Microtubule-Associated Protein (EMAP) on Microtubules, but Not Filamentous
Actin or Vimentin Filaments.". *Cell Motil Cytoskeleton* 50: 161–72.

Andersen, SS, B Buendia, JE Domínguez, A Sawyer, and E Karsenti. 1994. "Effect on
Microtubule Dynamics of XMAP230, a Microtubule-Associated Protein Present in Xenopus
Laevis Eggs and Dividing Cells.". *J Cell Biol* 127: 1289–99.

Monroy, Brigette Y., Danielle L. Sawyer, Bryce E. Ackermann, Melissa M. Borden, Tracy C.
Tan, and Kassandra M. Ori-McKenney. 2018. "Competition between Microtubule-Associated
Proteins Directs Motor Transport". *Nature Communications* 9 (1).

Tokuraku, K, K Matsushima, T Matui, H Nakagawa, M Katsuki, R Majima, and S Kotani. 2003.
"The Number of Repeat Sequences in Microtubule-Associated Protein 4 Affects the
Microtubule Surface Properties.". *J Biol Chem* 278: 29609–18.

Adib, Rozita, Jessica M. Montgomery, Joseph Atherton, Laura O'Regan, Mark W. Richards,
Kees R. Straatman, Daniel Roth, et al. 2018. "Mitotic Phosphorylation by Nek6 and Nek7
Reduces Microtubule Affinity of EML4 to Alter Spindle Dynamics and Promote
Chromosome Congression", November. https://doi.org/10.1101/466979.

Ookata, K, S Hisanaga, M Sugita, A Okuyama, H Murofushi, H Kitazawa, S Chari, JC Bulinski,
and T Kishimoto. 1997. "MAP4 Is the in Vivo Substrate for CDC2 Kinase in HeLa Cells:
Identification of an M-Phase Specific and a Cell Cycle-Independent Phosphorylation Site in
MAP4.". *Biochemistry* 36: 15873–83.

Chang, W, D Gruber, S Chari, H Kitazawa, Y Hamazumi, S Hisanaga, and JC Bulinski. 2001.
"Phosphorylation of MAP4 Affects Microtubule Properties and Cell Cycle Progression.". *J
Cell Sci* 114: 2879–87.

Drewes, G, B Lichtenberg-Kraag, F Döring, EM Mandelkow, J Biernat, J Goris, M Dorée, and E
Mandelkow. 1992. "Mitogen Activated Protein (MAP) Kinase Transforms Tau Protein into
an Alzheimer-like State.". *EMBO J* 11: 2131–38.

Illenberger, S, G Drewes, B Trinczek, J Biernat, HE Meyer, JB Olmsted, EM Mandelkow, and E
Mandelkow. 1996. "Phosphorylation of Microtubule-Associated Proteins MAP2 and MAP4

by the Protein Kinase p110mark. Phosphorylation Sites and Regulation of Microtubule Dynamics.". *J Biol Chem* 271: 10834–43.

Drewes, G, A Ebneth, U Preuss, EM Mandelkow, and E Mandelkow. 1997. "MARK, a Novel Family of Protein Kinases That Phosphorylate Microtubule-Associated Proteins and Trigger Microtubule Disruption.". *Cell* 89: 297–308.

Hazel, J, K Krutkramelis, P Mooney, M Tomschik, K Gerow, J Oakey, and JC Gatlin. 2013. "Changes in Cytoplasmic Volume Are Sufficient to Drive Spindle Scaling.". *Science* 342: 853–56.

Good, MC, MD Vahey, A Skandarajah, DA Fletcher, and R Heald. 2013. "Cytoplasmic Volume Modulates Spindle Size during Embryogenesis.". *Science* 342: 856–60.

Rieckhoff, EM, K Ishihara, and J Brugués. 2019. "How to Tune Spindle Size Relative to Cell Size?". *Curr Opin Cell Biol* 60: 139–44.

Rieckhoff, Elisa Maria, Frederic Berndt, Stefan Golfier, Franziska Decker, Maria Elsner, Keisuke Ishihara, and Jan Brugués. 2020. "Spindle Scaling Is Governed by Cell Boundary Regulation of Microtubule Nucleation", June. https://doi.org/10.1101/2020.06.15.136937.

Kinoshita, K, I Arnal, A Desai, DN Drechsel, and AA Hyman. 2001. "Reconstitution of Physiological Microtubule Dynamics Using Purified Components.". *Science* 294: 1340–43.

Zanic, M, PO Widlund, AA Hyman, and J Howard. 2013. "Synergy between XMAP215 and EB1 Increases Microtubule Growth Rates to Physiological Levels.". *Nat Cell Biol* 15: 688–93.

Pelletier, James F., Christine M. Field, Sebastian Fürthauer, Matthew Sonnett, and Timothy J. Mitchison. 2020. "Co-Movement of Astral Microtubules Organelles and F-Actin Suggests Aster Positioning by Surface Forces in Frog Eggs", June. https://doi.org/10.1101/2020.06.17.154260.

Field, CM, JF Pelletier, and TJ Mitchison. 2017. "Xenopus Extract Approaches to Studying Microtubule Organization and Signaling in Cytokinesis.". *Methods Cell Biol* 137: 395–435.

Field, CM, PA Nguyen, K Ishihara, AC Groen, and TJ Mitchison. 2014. "Xenopus Egg Cytoplasm with Intact Actin.". *Methods Enzymol* 540: 399–415.

Tinevez, JY, N Perry, J Schindelin, GM Hoopes, GD Reynolds, E Laplantine, SY Bednarek, SL Shorte, and KW Eliceiri. 2017. "TrackMate: An Open and Extensible Platform for Single-Particle Tracking.". *Methods* 115: 80–90.

Ho, J, T Tumkaya, S Aryal, H Choi, and A Claridge-Chang. 2019. "Moving beyond P Values: Data Analysis with Estimation Graphics.". *Nat Methods* 16: 565–66.

Wühr, M, T Güttler, L Peshkin, GC McAlister, M Sonnett, K Ishihara, AC Groen, et al. 2015. "The Nuclear Proteome of a Vertebrate.". *Curr Biol* 25: 2663–71.