1 **Variation in transparency degree: response to light availability or part of**
2 **mimicry syndrome?**
3

4 Mónica Arias[1,2], Jérôme Barbut[2], Rodolphe Rougerie[2], Manon Dutry[1], Mireia Kohler[1],
5 Baptiste Laulan[1], Caroline Paillard [1], Serge Berthier[3], Christine Andraud[4], Marianne
6 Elias[2], Doris Gomez[1,3]
7

**Affiliations**
[1]CEFE, Univ Montpellier, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier,
France
[2]Institut de Systématique, Evolution, Biodiversité, CNRS, MNHN, Sorbonne
Université, EPHE, Université des Antilles, 45 rue Buffon CP50, France
[3]INSP, Sorbonne Université, CNRS, Paris, France
[4]CRC, CNRS, MNHN, Paris, France

15
16

17 **Corresponding author email**: moarias@gmail.com
18

**Abstract (247 words)**
Transparency is common in aquatic environments where it reduces detectability by
predators. Transparency degree (the proportion of transmitted light) varies under
water and is higher as light availability increases. Transparency is rare on land and
poorly studied. Recent studies have shown that Lepidoptera – a group in which
transparency has independently evolved multiple times – also display large variation
in transparency degree. The large diversity encompasses different types of clearwing
species, such as bee/wasp mimics and death leaf mimics. This study explores how
the evolution of different transparency degree could be related to habitat openness,
and to mimicry syndrome, through two complementary approaches. First, by
exposing artificial moth-like prey to wild avian predators in open and closed
habitats, our fieldwork experiment shows an increase in prey survival at higher
transparency degrees in open compared to closed habitats. Second, by analysing the
evolution of wing characteristics and ecological traits in 107 clearwing species,
comparative analyses show that instead, open habitat species show in average

similar or slightly lower transparency degree than closed habitat species. We also
find that bee/wasp mimics have highest and least varying transparency degrees and
are more often diurnal. Compared to bees/wasp mimics, leaf mimics have lower but
more varying transparency degree and are more often nocturnal. High transparency
degree likely reduces detectability in open habitats, as shown by field experiments
and comparative analyses of optical measurements on bee/wasp mimics. Our results
suggest that habitat openness, and species interactions play a crucial role in
determining transparency design.
42
43  **Keywords**: transparency, irradiance, mimicry, nocturnality, Lepidoptera
44

**Introduction**

Transparency is common in aquatic environments where it reduces detectability and works as an antipredatory defence, especially in the pelagic environment where there is nowhere to hide [1]. However, the amount of light that is transmitted by transparent tissues (hereafter referred to as transparency degree) varies between organisms. Based on the observation of 29 zooplankton species, Johnsen and Widder [2], found that transparency degree could range from 50 to 90%, with the most transparent species living in lighter, shallower places. In a theoretical approach, they show that efficiency at detecting transparent species depends on visual performance (for instance, minimum contrast that predator can detect, also known as contrast threshold), which varies with ambient light conditions: in shallow places with bright light conditions, high visual performance (low contrast threshold values) selects for high transparency degree, and only highly transparent organisms are effectively undetected. Instead, at deeper areas with dimmer light conditions, visual systems are less performing (high contrast threshold values) and select for low transparency degree, thus organisms can be poorly transparent and be effectively undetected [3, Figure S1]. On land, studies in primates have also shown higher visual performance at brighter light conditions as the retinal cells increase their neural response more strongly when mean luminance is higher [4,5]. Similarly, visual performance increases with ambient light luminance in other terrestrial vertebrates such as birds [6–8]

Transparency is rather rare in terrestrial environments and is almost restricted to insect wings. Transparency also reduces detectability of prey on land [9–11]. Recently, Gomez et al [12] reported that transparency has evolved multiple times independently in Lepidoptera, and that it is produced by a large diversity of structures and configurations with different transparent degrees. This study does not investigate the links between transparency and habitat, but suggests in several aspects an important role of visual predators in driving the evolution of transparency (for instance diurnal species were reported to have higher transparency degree than nocturnal species). Hence, we expect that terrestrial transparent species living in more open habitats - with brighter light conditions - should present higher transparency degree compared to species living in more closed habitats - with dimmer light conditions -. In addition, closed-habitat species

might show larger variation in transparency degrees as many degrees are similarly

visually protective, thus high transparency degrees are not strongly selected.

Predation risk can be reduced not only by reducing detectability, but also by

mimicking inedible elements such as leaves [masquerade, 13] or unprofitable prey

[batesian mimicry, 14], and transparency might help achieving such mimicry.

Several clearwing Lepidoptera exhibit brownish colorations that combined with

transparent surfaces resembling "holes" can mimic dead leaves (examples of

potential dead-leaf mimics in Fig. 1a). Other small clearwing moths and butterflies

exhibit yellow and dark stripes on their bodies and highly transparent wings

(examples in Fig. 1b), resembling harmful species for predators, such as bees and

wasps. Mimicry syndrome could concur in driving transparency degrees to have

different optima in bee/wasp and leaf mimics, but whichever factor (habitat light

availability versus mimicry) is driving the evolution is still unknown. Bees and

wasps often live in open habitats [15,16], and their mimics should do as well. High

transparency degree should be selected in bees, wasps and their mimics because

their body sizes are rather small, a trait associated to higher transparency degrees in

butterflies [12], and light conditions are brighter in open habitats. Instead, dead-leaf

mimics might show a larger variation in body size, may more often live in closed

habitats, where more leaves fall down and should exhibit a broader diversity of

transparency degrees, as all of them can contribute to mimic leaves in

decomposition and confer similar protection against predators. The expected

differences in transparency degrees should also be associated to variations in

morphology and coloration of wing scales, with absent or transparent hair-like

scales for bee/wasp mimics and classically shaped scales for leaf mimics.

Here, we studied whether differences in transparency degree found in

Lepidoptera clearwing species are related to light availability and/or mimicry using

two complementary approaches. First, we tested whether the efficiency of

transparency at reducing detectability changed according to light availability on

terrestrial environments, by carrying out a fieldwork experiment with artificial prey.

Second, using real butterflies, we tested whether differences in transparency degree

could be explained by species habitat and by mimicry, by conducting comparative

analyses at broad interspecific level. We took advantage of the 107-species dataset of

Lepidopteran species analysed by Gomez et al [12] to add species ecological traits

(habitat and mimicry) and relate them to clearwing optical and structural traits.
Together, these analyses help understanding the evolution of interspecific
differences in transparency degree on land.
114
**Material and Methods**
<u>Artificial butterfly Elaboration</u>
To test for the efficiency of transparency degree at reducing detectability in habitats
with different light availability, we elaborated plain grey artificial butterflies with
paper wings and a malleable edible body. Following the general methodology
described in Arias *et al* [9], butterflies did not replicate any real local butterfly, but
mimicked a general grey moth with closed wings (i. e., a triangular shape). The
rather cryptic butterfly colour was chosen based on 120 reflectance measures of
green oak *Quercus ilex* trunks. Reflectance measurements were taken using a
spectrophotometer (Starline Avaspec-2048 L, Avantes) and a deuterium halogen
lamp (Avalight DHS, Avantes) emitting in the 300-700 nm range, including UV, to
which some predators of butterfly and moths, such as birds, are sensitive [17].
Measurements were done relative to a white reference (lights turned on with no
sample) and a dark reference (light turned off with no sample). For reflectance
measurements, we used an optic probe (FC-UV200-2-1.5 x 100, Avantes) merging
illumination and collection angles. Grey wings (R=155, G=155, B=155) were printed
on sketch paper Canson using a HP Officejet Pro 6230 printer. Paper wings were
laminated with a Polyester Opale Mat 75μm poach. Four forms of artificial butterflies
we used that in order of transparency were:  completely opaque butterfly "C", poorly
transparent butterfly with 6 layers of transparent film "T6", highly transparent
butterfly with a single layer of transparent film "T1", and fully transparent butterfly
with no film in the transparent zones "T0". For the butterflies that included
transparent elements, triangular windows of 234 mm$^2$ were cut down from the grey
triangle and the remaining part was put on top of one or six layers of transparency
film 3M for inkjet printing. This transparent film was chosen as it is highly
transparent, even in the UV range of the spectrum, and going from 1 to 6 transparent
layers permitted a reduction of 50% of the transmittance between treatments (Fig.
S2). Such transparent layer was coated with a transparent mate varnish to reduce its
shininess. For transparency measurements, we measured specular transmittance

from 300 to 1100 nm, using a deuterium-halogen lamp (Avalight DHS, Avantes),

optical fibres (FC-UV200-2-1.5 x 100, Avantes) and a spectrometer (Starline

Avaspec-2048 L, Avantes). Fibres were separated, aligned and 5mm apart and the

wing sample was placed perpendicular between them at equal distance (light spot of

1mm diameter). Spectra were taken relative to a dark (light off) and to a white

reference (no sample between the fibres) measurement. On top of butterfly wings,

we added an artificial grey body. Butterfly bodies were prepared with flour (428 g),

lard (250 g) and water (36 g), as reported on Carrol & Sherratt [18]. Yellow, red and

blue food edible colourings were used to dye the pastry in grey, imitating artificial

wing colour. Such malleable mix permits to register and distinguish marks made by

bird beaks from insect jaws. Both paper wings and artificial bodies were measured

in spectrometry, similarly to how trunk reflectance was measured (Fig. S2). Bodies

could attract predator attention, but it has been previously shown that wingless

bodies had the same attractiveness as that of butterflies with large transparent

elements [9]. Artificial butterflies and bodies were pinned to green oak *Quercus ilex*

trunks. To avoid ant attacks, Vaseline and a piece of highly sticky double-faced

transparent tape were stuck in between the wings and the trunk.

<u>Experimental set up</u>

Predation experiments were performed in May and June 2018 at the zoological Park

of Montpellier (43.64°, 3.87°) and at La Rouviere forest (close to Montarnaud city

43.65°, 3.64°). This experiment overlapped with breeding season of local great

(*Parus major*) and blue tit (*Cyanistes caeruleus*) populations, when birds are looking

for food in an intensive way, what can maximise the number of attacks on our

artificial prey. Artificial prey items were pinned every 10 metres on oak trunks of at

least 10 cm of diameter and with few or no moose cover. Prey order was randomized

by blocks of eight items, two per treatment, before starting the experiment. Initially,

prey was evenly disposed in closed and open areas. However, throughout the

experiment a stronger predation rate was detected in open areas (see Results

section). Therefore, we increased the number of artificial prey released on closed

areas to compensate for the differential predation pressure and get more similar

numbers of attacked prey items in closed and open habitats. We analysed results for

both original (larger sample size in closed habitats) and reduced datasets (same

number of butterflies placed in both closed and open habitats). Prey were mostly
disposed facing north and as perpendicular to the floor as possible, to reduce direct
sunlight reflection on them. Prey was checked every 24h. After 96h all prey were
removed. When V- or U- marks were detected, or when the body was missing
without signals of invertebrate attacks (i.e. no body scraps left on the wings or
around the butterfly on the trunk) prey were considered as attacked and wings or
pins where removed. Instead, if invertebrate attacks were detected, or when prey
was missing, it was replaced. Non-attacked prey were considered as censored data.
Canopy cover was used to classify places in open and closed habitat. We took
pictures each 20 meters in the trails (at forest edges for the open habitat, inside the
forest for the closed habitat, examples Fig. S3) that were used for the experiment,
using a Canon EOS 550D camera and a fisheye lens. Canopy cover was further
quantified using Fiji software [19]. After verifying unimodality in canopy cover
distribution (Hartigans' dip test: D = 0.01, p-value = 0.6), we used a discrete
classification, assuming that places with cover larger than the mean where closed
habitats and places with lower covers where open habitats (see Fig. S3 for the
distribution of canopy cover in the experimental sites).

To test how different treatments survive along the experiment on both open and
closed habitats, Cox proportional hazard regression [20] were applied, first for all
data points including form, habitat openness and their interaction as explanatory
variables, and then, Cox regression were independently applied for each habitat
using the different treatments as factors, to further explore the effect of form within
each habitat. Overall significance was measured using a Wald test. Statistical
analyses were performed in R [21] using the *survival* package [22].

<u>Species selection and phylogeny</u>
We used a dataset of 107 clearwing species extracted from that used by Gomez *et al*
[12], and including those species for which information about the openness of their
habitat was available. These species spanned 31 of the 124 existing Lepidoptera
families and were selected based on the presence of transparent or translucent
zones in their wings. For each species, we selected one specimen from the
Lepidoptera collection of National Museum of Natural History (Paris, France). For
comparative analyses, we used the phylogenetic tree built by Gomez *et al* [12]. In

brief, 9 nuclear and mitochondrial DNA sequences were taken from GenBank or

BOLD for the clearwing species or a surrogate species from the same genus, tribe,

subfamily or family. 60 additional species were incorporated to improve tree

topology. After sequence alignment and partitioning, tree Bayesian inference using

BEAST 1.8.4 [23] by constraining family topology and node ages to follow [24] was

performed. After combining several runs with appropriate burn-in periods, the

maximum clade credibility tree (MCCT) was computed with median node ages,

which was then used for our analyses.

<u>Ecological data on species habitat, daytime activity and mimicry type</u>

Ecological information was gathered consulting papers and experts for a total of 107

species for activity rhythm (58% of species are diurnal species), habitat (56% of

species living in open habitats), and mimicry status (14 species were bee/wasp

mimics, 13 species were leaf mimics, while the remaining 80 did not fall in either of

those categories. Table S1 and Fig S4). We excluded from the analyses species that

were reported to live in open and closed habitats, or to fly at day and night. Bee and

wasp mimics were grouped together in the same mimicry category, as both exhibit

the striped pattern with black and yellow that advertise their unpalatability [25].

Butterflies and moths mimicking bees and wasps have forewings longer than their

hindwings, and often strong delimitations of their wing borders and veins by dark

coloration, as the hornet moth *Sesia apiformis* (*api* - bee, *formis* -shaped) or the

broad-bordered bee hawk-moth *Hemaris fuciformis* (Fig 1b). Fallen leaf mimics

include brownish moths and butterflies with small or large transparent surfaces and

that could touch or not the wing borders (Fig. 1a). *Rotschildia lebeau* has been

proposed as a leaf mimic [26] but such mimicry is controversial [27]. *Rotschildia*

*ericina* was therefore not considered as a leaf mimic for the analyses.

<u>Wing structural and optical measurements</u>

Museum specimens were photographed using a D800E Nikon camera equipped with

a 60mm lens, placed on a stand with an annular light. Photos were then analysed

using ImageJ [28] to extract descriptors of wing macrostructure: wing length (mm),

wing surface ($mm^2$), and clearwing area (the surface of transparent area in $mm^2$), for

the forewing and hindwing separately. We defined and computed proportion of

clearwing area as the ratio clearwing area/wing surface, i.e. the proportion of the
total wing area occupied by transparency.

For transparency measurements, we followed the same procedure described
above for measuring transparency in artificial prey. For each species and wing, we
took five measurements in the transparent zone. We analysed spectral shape using
Avicol v6 [29] to extract the mean transmittance over [300-700] nm, which
described the level of transparency.

<u>Physical properties of transparent wings</u>
Gomez et al [12] described a large diversity in transparent wing surface
characteristics. Using binocular imaging (Zeiss Stereo Discovery V20) and digital
microscopic imaging (Keyence VHX-5000), these authors found that transparency
could be achieved either with nude membranes or with membranes covered by
scales of various types (either classical scales, hair-like scales or both) that could be
coloured or transparent, and erected of flat according to their insertion on the wing
membrane. Therefore, we also explored the evolution of mimicry according to these
characteristics.

<u>Analyses of transparency degree differences between habitats and mimics</u>
To explore whether differences in transparency degree (percentage of transmitted
light) are related to variation in habitat and mimicry we fitted both 1) a linear mixed
model including 107 species for which we found data on habitat preference and 2) a
Bayesian phylogenetic mixed model, using the phylogenetic reconstructions
obtained by Gomez et al including 183 species from 31 families. Comparisons
between the first and the second analysis allowed us to assess the influence, if any, of
phylogenetic relationships on the observed trends. We included habitat
(open/closed) and mimicry (bee-wasp mimics/leaf mimics/no mimicry with the
contrasts bee/wasp mimics more transparent than all other species, and leaf mimics
less transparent than other types of species) as explanatory variables, together with
variables reported by Gomez et al (2020) as correlated to transparency degree: wing
length (as a proxy of butterfly size), proportion of clearwing surface, fore-/hindwing
and daytime activity (diurnal/nocturnal). Butterfly transparency, wing length and
clearwing proportion were obtained from Gomez *et al* [12]. As five measurements

per wing (fore- and hindwing), thus ten measurements per species were included in
our dataset, wing measures nested in species was considered as a random effect. We
fitted different models including different combinations of two- and three-factor
interactions and we compared them using the AIC criterion to select the best model.
We reported and compared the model without interactions and the best model
including interactions. Additionally, we fitted Bayesian phylogenetic mixed models
with Markov chain Monte Carlo analyses using the 'mulTree' R package [30]. Using
the best formulated linear mixed model, uninformative priors were chosen, with
variance and belief parameter set to 1 for random effects and 0.002 for residual
variances [31]. Models were run using two chains of 600,000 iterations with and a
thinning interval of 300 and a burn-in of 10,000. Fixed effects were considered
statistically significant when the probabilities in the 95% credible intervals did not
include zero.
Additionally, we performed several phylogenetic generalised least square (PGLS)
analyses (1 observation per species) to test for potential correlations between size,
proportion of clearwing surface, daytime activity, habitat and mimicry, and between
wing ratio (forewing length/hindwing length, usually high in bees and wasps) and
mimicry. Using BayesTraits [32] we additionally explored character coevolution
between mimicry/habitat, mimicry/daytime activity and habitat/daytime as well as
between mimicry and scale characteristics: absence/presence of scales, scale
insertion (flat or erected), scale coloration (coloured or transparent) as well as scale
type (hair-like scales and classic scales). When analysing physical wing properties
and mimicry evolution, independent analyses were performed for anterior and
posterior wings, as differences between wings have been reported by Gomez et al
[12]. We calculated the likelihood of the dependent and independent models of
evolution of each pair of binary characters, using a Markov-Chain Monte-Carlo
approach. We then estimated the likelihood ratio test between them and compared it
to a chi-square distribution with 4 degrees of freedom to obtain their statistical
significance [33]. As this approach can only be applied to binary characters, and
mimicry has three factors, models were run including the different possible pairwise
combinations to explore: a. evolution of leaf mimicry (including non-mimics and leaf
mimics); b. evolution of bee/wasp mimics (including non-mimics and bee/wasp
mimics) and c. differences in the evolution of leaf and bee/wasp- mimics (including

bee/wasp and leaf mimics). We calculated δ, a new metric inspired in the
phylogenetic analog of the Shannon entropy to measure phylogenetic signal in
categorical non-ordered data [34], for testing for phylogenetic signal in mimicry
evolution. δ is useful to detect whether the coevolution between mimicry and other
traits are homologies (mimicry and the other characteristics are shared within a
clade) or convergences (mimicry and the other characteristics evolved
independently and several time along the phylogeny).

Finally, we predicted more variance in transparency degree for those
conditions where different levels of transparency might be equally efficient as would
be the case for butterflies and moths that are active during the night, that live in
closed habitats and/or that are leaf mimics. We applied Fligner-Killeen tests of
homogeneity of variance in R to test for difference in variance between a) diurnal
and nocturnal species, b) species living in open and in close habitats and c)
bee/wasp mimics, leaf mimics and no mimic species. As Gomez et al (2020) reported
that transparency degree is closely related to wing size, we also tested for
differences in the variation of wing length between habitat, daytime activity and
mimicry using the test mentioned above.

<u>Results</u>

<u>Fieldwork experiment</u>

In total, 1149 artificial butterflies were used during the experiment and 243 prey
were attacked (21.14 %). When data from both open and closed habitats were
analysed together (same numbers of attacked prey items), survival was lower in
open than in closed areas. In closed areas 651 prey were released, 110 attacked, with
an attack rate of 16.89% while in open areas, 498 prey were released, 133 attacked,
with an attack rate of 26.71% (Wald test = 34.39, df = 7, p<0.001). Results are similar
when including the "reduced" dataset (a random subsample of the closed habitat set
and of the same sample size as the set released in open habitats): 15.66% of attacks
in closed habitats (92 attacks). Wald test = 24.67, df = 7, p<0.001). Additionally, in
open habitats the opaque form was more attacked than the other forms that had
transparency (z= 2.15, p=0.031), but when including the "reduced" dataset, such
difference could not be detected anymore (z= 1.52, p=0.127). When data was
independently analysed by habitat closeness, all prey treatments were similarly

attacked in closed areas (Fig. 2a, Wald test = 0.9, df = 3, p = 0.825, and for the
reduced dataset Wald test = 0.5, df = 3, p = 0.5), in contrast to open areas (Fig. 2b,
Wald test = 9.86, df = 3, p = 0.02). In open areas, opaque artificial butterflies were
more attacked than all the other treatments (z = 2.68, p = 0.007); artificial butterflies
with six layers of transparency were marginally more attacked than those with a
single transparency layer (z = 1.839, p = 0.066), while artificial butterflies without or
with a single transparent layer where similarly attacked and reported the lowest
number of attacks (z = -1.61, p = 0.11).
350
<u>Comparative analyses</u>
<u>Variations in transparency degree, and relationship to habitat and daytime activity</u>
Habitat explains a small part of the variation found in transparency degree. When
interactions between factors are excluded from the model, species have in general
the same level of transparency for closed and open habitats (t=-1.58, df=102, p
=0.117, Table 1). When interactions are included, species living in closed habitats
transmit slightly more light than species living in open habitats (t=-2.09, df=102, p
=0.04, Table 1, Fig. 3). In open habitats transmittance is slightly higher in most
species for the forewing and in species showing smaller hindwing sizes, but
transparency degree decreases steeply for hindwings with increase in wing size
(triple interaction wing, habitat and size: t=30,2, df=488, p=0.003, Table 1, Fig 3b.).
Regarding correlations between habitat and other physical characteristics of species
wings, species living in closed habitats are not different in size (PGLS forewing: t=-
1.8, p= 0.07; hindwing: t=-1.32, p= 0.19) or proportion of clearwing surface from
those living in open habitats (PGLS forewing: t=0.27, p = 0.78; hindwing: t=0.13,
p=0.9). No relationship was found between habitat and daytime activity (BayesTraits
LRT= 6.08, df=4, p=0.193, Table 2). Daytime activity has no significant contribution
to explaining variations in transparency degree in models including mimicry and
habitat as explanatory variables (Table 1). Diurnal species have larger hindwings
than nocturnal species (PGLS t=-2.33, p=0.02) but no differences in forewing sizes
were detected (PGLS t = -1.45, p =0.15). No differences in the proportion of
clearwing area were found between diurnal and nocturnal species (PGLS forewing: t
= -1.58, p =0.12; hindwing: t=0.16, p=0.87). No differences in the variance of
transparence degree were found between habitats (Fligner-Killeen test for forewing:

2.02, df=1, p=0.155; for hindwing: 1.48, df = 1, p=0.224). However, nocturnal species

vary more in transparency degree than diurnal species for their forewing (Fligner-

Killeen test for forewing: 15.47, df=1, p<0.001; for hindwing: 2.87, df = 1, p=0.09, Fig.

S8). Diurnal species vary more on their wing size than nocturnal species (Fligner-

Killeen test for all forewing: 8.79, df=1, p=0.003; for all hindwing: 30.51, df = 1,

p<0.001, Fligner-Killeen test for wings smaller than 60 mm, forewing: 22.22, df=1,

p<0.001; hindwing: 39.66, df=1, p<0.001, Fig. S8). No difference in wing size

variation was found for species from different habitat openness (Fligner-Killeen test

for forewing: 0.26, df=1, p=0.610; for hindwing: 0.196, df = 1, p=0.657).

<u>Variations in transparency degree, transparency structural basis and mimicry</u>

Mimicry explains a large portion of the variation found in transparency degree.

When fitting a model without interactions, bee/wasp mimics had a higher

transparency degree than other clearwing species (t=2, df = 102, p = 0.05, Table 1,

Fig. 4). When interactions were included, the best model additionally shows that

within bee/wasp mimics, small-sized species have higher transparency degrees than

large-sized species (t=-3.04, df=488, p=0.003), in contrast to leaf mimics that

transmit more light at larger wing sizes (t=-4, df=488, p<0.001, Fig. 4d). Additionally,

light transmittance differences between mimics are larger for hindwing than for

forewing (t=2.69, df=488, p=0.007, Figure 4a). Regarding the relationship between

mimicry and wing characteristics, leaf mimics have smaller hindwings than non-

mimics (PGLS forewing: t=1.66, p = 0.1; hindwing: t=1.99, p = 0.05, Figure 4b), and

the length ratio forewing/hindwing is larger for bee/wasp mimics than for other

clearwing species (PGLS t=4.51, p <0.001, Fig. 4e). Additionally, proportion of

clearwing surface is larger for no mimics and for leaf mimics (PGLS forewing:

t=4.005, p < 0.001; hindwing: t=2.86, p=0.005, Fig. 4c). Most bee/wasp mimics are

diurnal while most leaf mimics are nocturnal (BayesTraits LRT=13.56, df = 4,

p=0.009, Table 2, Fig S4 & Fig S6).

We found that transparence degree varies less in bee/wasp mimics than in

leaf-mimics (Fligner-Killeen test for forewing: 15.6, df= 1, p<0.001, for hindwing:

36.14, df = 1, p<0.001), or in other clearwing species (Fligner-Killeen test for

forewing: 16.38, df=1, p<0.001; for hindwing: 36.34, df = 1, p<0.001), but there is no

difference in the transparence degree variation in leaf mimics and non-mimic

species (Fligneer-Killeen test for forewing: 2.02, df=1, p= 0.155; for hindwing:
0.00005, df = 1, p=0.995). Likewise, wing length varies less in bee/wasp mimics than
in leaf mimics (Fligner-Killeen test for forewing: 18.91, df=1, p<0.001; for hindwing:
39.86, df = 1, p<0.001) or in other clearwing species (Fligner-Killeen test for
forewing: 39.06, df=1, p<0.001; for hindwing: 59.15, df = 1, p<0.001). Leaf and non-
mimics show a similar variation in wing length (Fligner-Killeen test for forewing:
0.58, df=1, p=0.447; for hindwing: 2.46, df = 1, p=0.117).

Some scale characteristics are correlated to mimicry evolution. Scales are
more often absent and less often coloured when present in hindwings of bee/wasp
mimics in comparison to non-mimics wings (BT LRT presence = 24.55, df=4,
p<0.001, Table S2 & Fig S7; BT LRT colour= 16.14, df=4, p=0.003, Table S2 & Fig S7).
Similarly, leaf mimics have less often coloured scales in comparison to non-mimics
both in forewing (BT LRT=11.62, df=4, p=0.02) and hindwing (LRT=12.87, df=4,
p=0.012, Table S2 & Fig S7). Moreover, leaf mimics only exhibit classic scales in
contrast to bee/wasp mimics and non-mimics (LRT=9.33, df=4, p=0.053, Table S2 &
Fig S7). More than evolutionary convergences, these similarities are likely to be
homologies as mimicry has a strong phylogenetic signal ($\delta$ =4.471).

Some characteristics were common for the general set of species that we
studied here. Transparency degree is higher on forewings (t=9.45, df=502, p<0.001),
wings with large proportion of clearwing areas (t=10.88, df=502, p<0.001, Fig. S5a)
and smaller species (t=-6.61, df=502, p<0.001, Fig. S5b). Regarding general wing
characteristics, smaller species have larger proportion of transparent surface in
hindwings than do larger species (PGLS forewing: t=-1.42, p = 0.16; hindwing: t=-
2.58, p = 0.01).

**Discussion**
Fieldwork experiments using artificial prey that differed on their transparency
degree suggest that variation in transparency can be detected only in brighter light
conditions, similar to what has been suggested for aquatic environments [2]. Poorly
transparent prey types survived less than highly transparent ones in open
environments, while they performed similarly in closed areas. Additionally, less
artificial moths were attacked in closed habitats, even after increasing the number of
artificial prey in closed habitats to compensate for the higher predation rate often

reported for open habitats such as forest edges [35]. By controlling for differences in

predation rate between habitats by increasing the number of prey placed in closed

habitat our results suggest that selective pressure towards an increase in

transmittance is higher in open than in closed habitats.

Our experimental results suggest that if detectability reduction is an

important trigger for the evolution of transparency degree variation, butterflies and

moths with poorer transparency degree should be mostly restricted to closed

habitats. However, we found that when excluding factor interactions, overall

butterflies and moths living in closed habitats have a similar transparency degree,

and that there is a slightly higher mean transmittance in species living in open

habitats, only revealed when interactions with other factors were included in the

analyses. Gomez et al [12] reported that poor light transmittance is associated to

small proportions of clearwing surfaces and large wing sizes, characteristics that

were similarly segregated in open and closed habitats (LMM estimates for the

interactions wing size: habitat = 0.31±0.05; and clearwing surface: habitat = -

0.28±0.14 , Fig. 3), probably explaining the small differences found in transparency

degree between different light conditions. However, we also found that forewings

and small hindwings transmitted more light in open than in closed habitat species, in

accordance to what would be selected according to the high visual performance of

visual predators under bright conditions [2]. This suggests that differences detected

in light transmittance between habitats are closely related to selective pressures

acting in size as well as in fore- and hindwing in an independent manner. As Gomez

*et al* [12] suggested, transparency might be highly effective at small body sizes, but

its efficiency likely decreases with size, as larger wings and bodies imply larger

opaque elements, such as the butterfly body itself. Our study confirms this trend and

sheds more light on the evolution of these morphological traits. For cryptic Erebidae

moths, Kang et al [36] have suggested that the increase of conspicuousness related

to larger body sizes might have triggered the evolution of hidden contrasting colour

signals on the hindwing that are probably involved in startle displays. Selective

pressure towards decreased detectability might promote different character

evolutions according to prey size, especially in open habitats. Species living in open

habitats incur in higher risks of tissue damage by higher exposure to UV light, more

wind or higher exposition to direct rainfall, conditions that have been reported as

important at shaping moth and butterfly segregation in different habitats [37]. In
case transparency entails costs related to one or several of these factors, they will
promote the presence of opaque scales in open habitats, as it might be the case for
large species, where additionally large opaque elements will be difficult to hide even
with the highest transparency degree.

Transparency can decrease prey detectability but it is also involved in
mimicry evolution, and differences in transparency degree seem to be intimately
associated to mimicry evolution. When mimicry is included in the analysis, we found
that bee/wasp mimics have higher transparency degrees and are more often
restricted to open habitats, in agreement to our fieldwork results and confirming
that high transparency in open habitats can be a visual optimum in terrestrial
environments. As this is not the case for all clearwing species, this highlights the
strong effect of biotic and abiotic interactions in shaping adaptive peaks, thus the
segregation of species under different conditions and the evolution of different
features.

Other characteristics of bee/wasp mimics include the absence of scales in
their wings, their diurnal activity [38], their smaller wing sizes and their smaller
hindwings in comparison to their forewings [39,40]. Insects that mimic bees or
wasps resemble their harmful counterparts in body colour, body shape and flight
pattern (for butterflies: [41]) but also in activity patterns (for diptera: [42]).
Convergence in high transparency degree along with reduction in variation in wing
size and increase in fore/wing ratio can contribute to such multicomponent mimicry,
that improves prey chances at deceiving a larger set of predators, in case several
traits are similarly salient for them [43] or in case different predator species find
different cues as salient [44]. The evolution of highly transparent wings in
pollinators is likely triggered by their higher mobility and the openness of the
habitat where the majority of nectar sources occurred [45]. Studies in other non-
insect transparent organisms have shown that transparent forms of *Hippolyte*
*obliquimanus* shrimps are more mobile and exploit a larger diversity of habitats [46].
Whether transparency is also related to higher mobility in insects remains to be
properly tested.

On the other hand, we found that leaf mimics are larger, are more often found
in closed habitats and in contrast to the trends for all clearwing species, have a

higher transparency degree at larger wing sizes and lower proportions of clearwing
surfaces. Some fully opaque fallen leaf mimic species, as the comma butterfly
(*Polygonia c-album*) exhibit a distinctive pale mark that might decrease their
predation [47,48], although it is controversial [49]. Either way, reduced and highly
transparent zones as those found in *Orthogonioptilum violascens* might produce a
similar appearance to those markings given their high transparency degree that
might be highly reflective and be perceived as a white mark, conferring thus the
potential benefits of the "comma mark" when seen for a given angle, as well as the
full crypsis, when no reflections can be perceived. Whether highly transparent
windows work as other highly visible elements displayed in cryptic species such as
eyespots that deter predators [50,51] and that are also important in mate choice
[52] remains to be tested.

Other fallen leaf mimics include large transparent surfaces that can be large
and can touch moth wings edges as for *Bertholdia* species or not touching them as
for *Pseudasellodes fenestraria*. In both cases, transparency surfaces are likely
disrupting prey shape (as in the first case) and/or prey surface (as in the second
example [53]), thus reducing their detectability. Disruptive coloration, especially
when touching prey borders has been broadly studied as it is common in fully
opaque cryptic species [54–57]. Therefore, in case transparent surfaces touch wing
edges, they might work as disruptive marks regardless of their transparency degree.

**Conclusion**
Similar to aquatic environments, differences in transparency degree keeping
butterfly size constant seems more easily detectable in lighter environments, and
higher transparency degrees can indeed decrease detectability of prey by predators,
as our field experiment suggests. In real species, transparency degree between
habitats is rather similar. Yet, bee/wasp mimics are more often diurnal and
associated to open habitats, and have higher and less varying transparency degrees,
while leaf-mimics have lower and more varying transparency degree. Both habitat
light availability and mimicry can concur to fostering variations in transparency
degree. Further studies exploring the eventual coast of transparency regarding
irradiance, wind and direct rainfall, will help to better understand the evolution and
the constraints imposed on transparency on land.

Acknowledgments

This work was funded by Clearwing ANR project (ANR-16-CE02-0012), HFSP project on transparency (RGP0014/2016) and a France-Berkeley fund grant (FBF #2015-58). We warmly thank Jacque Pierre for help with species choice, identification, and information on species ecology, Edgar Attivissimo for contributing to Keyence imaging, Thibaud Decaëns, Daniel Herbin and Claude Tautel for species selection and identification, Charles Perrier for fruitful discussions on statistical analyses and Adrian Hoskins and Tony Pittaway for permitting using their photos for Figure 1.

References

1. Johnsen S. 2014 Hide and seek in the open sea: pelagic camouflage and visual countermeasures. *Annu. Rev. Mar. Sci.* **6**, 369–392.

2. Johnsen S, Widder EA. 1998 Transparency and visibility of gelatinous zooplankton from the northwestern Atlantic and Gulf of Mexico. *Biol. Bull.* **195**, 337–348.

3. Ruxton GD, Allen, William L., Sherratt TN, Speed MP. 2018 *Avoiding attack: the evolutionary ecology of crypsis, warning signals and mimicry*. Second. Oxford University Press.

4. Ghim MM, Hodos W. 2006 Spatial contrast sensitivity of birds. *J. Comp. Physiol. A* **192**, 523–534.

5. Purpura K, Kaplan E, Shapley R. 1988 Background light and the contrast gain of primate P and M retinal ganglion cells. *Proc. Natl. Acad. Sci.* **85**, 4534–4537.

6. Hodos W, Leibowitz RW, Bonbright Jr JC. 1976 Near-field visual acuity of pigeons: Effects of head location and stimulus luminance. *J. Exp. Anal. Behav.* **25**, 129–141.

7. Kassarov L. 2003 Are birds the primary selective force leading to evolution of mimicry and aposematism in butterflies? An opposing point of view. *Behaviour* **140**, 433–451.

8. Lind O, Karlsson S, Kelber A. 2013 Brightness discrimination in budgerigars (Melopsittacus undulatus). *PLoS One* **8**, e54650.

9. Arias M, Elias M, Andraud C, Berthier S, Gomez D. 2020 Transparency improves concealment in cryptically coloured moths. *J. Evol. Biol.* **33**, 247–252.

10. Arias M, Mappes J, Desbois C, Gordon S, McClure M, Elias M, Nokelainen O, Gomez D. 2019 Transparency reduces predator detection in mimetic clearwing butterflies. *Funct. Ecol.* **33**, 1110–1119.

11. Mcclure M *et al.* 2019 Why has transparency evolved in aposematic butterflies?
Insights from the largest radiation of aposematic butterflies, the Ithomiini. *Proc.*
*R. Soc. B* **286**, 20182769.

12. Gomez D *et al.* 2020 Transparency in butterflies and moths: structural diversity,
optical properties and ecological relevance. *bioRxiv*

13. Skelhorn J, Rowland HM, Speed MP, Ruxton GD. 2010 Masquerade: camouflage
without crypsis. *Science* **327**, 51–51.

14. Bates HW. 1862 XXXII. Contributions to an Insect Fauna of the Amazon Valley.
Lepidoptera: Heliconidæ. *Trans. Linn. Soc. Lond.* **23**, 495–566.
(doi:10.1111/j.1096-3642.1860.tb00146.x)

15. Grundel R, Jean RP, Frohnapple KJ, Glowacki GA, Scott PE, Pavlovic NB. 2010
Floral and nesting resources, habitat structure, and fire influence bee
distribution across an open-forest gradient. *Ecol. Appl.* **20**, 1678–1692.

16. Yamaura Y, Royle JA, Shimada N, Asanuma S, Sato T, Taki H, Makino S. 2012
Biodiversity of man-made open habitats in an underused country: a class of
multispecies abundance models for count data. *Biodivers. Conserv.* **21**, 1365–
1380.

17. Chen D-M, Goldsmith TH. 1986 Four spectral classes of cone in the retinas of
birds. *J. Comp. Physiol. A* **159**, 473–479.

18. Carroll J, Sherratt T. 2013 A direct comparison of the effectiveness of two
anti-predator strategies under field conditions. *J. Zool.* **291**, 279–285.

19. Schindelin J *et al.* 2012 Fiji: an open-source platform for biological-image
analysis. *Nat. Methods* **9**, 676.

20. Cox DR. 1972 Models and life-tables regression. *JR Stat Soc Ser B* **34**, 187–220.

21. R Foundation for Statistical Computing RC. 2014 *R: A language and environment*
*for statistical computing.* Vienna, Austria.

22. Therneau T, Lumley T. 2009 Survival: Survival analysis, including penalised
likelihood. R package version 2.35-7. *R Found. Stat. Comput.*

23. Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012 Bayesian phylogenetics with
BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* **29**, 1969–1973.

24. Regier JC *et al.* 2013 A large-scale, higher-level, molecular phylogenetic study of
the insect order Lepidoptera (moths and butterflies). *PLoS One* **8**.

25. Plowright R, Owen RE. 1980 The evolutionary significance of bumble bee color
patterns: a mimetic interpretation. *Evolution* , 622–637.

26. Janzen DH. 1984 Weather-related color polymorphism of Rothschildia lebeau
(Saturniidae). *Bull. ESA* **30**, 16–21.

27. Hernández-Chavarría F, Hernández A, Sittenfeld A. 2004 The" windows", scales,
and bristles of the tropical moth Rothschildia lebeau (Lepidoptera: Saturniidae).
*Rev. Biol. Trop.* **52**, 919–926.

28. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* **9**, 671–675. (doi:10.1038/nmeth.2089)

29. Gomez D. 2006 AVICOL, a program to analyse spectrometric data. *Free Exec.*
*Available Httpsitesgooglecomsiteavicolprogram Author Dodogomezyahoofr* **Last**
**update october 2011**.

30. Guillerme T, Healy K. 2014 mulTree: a package for running MCMCglmm analysis
on multiple trees. *Zenododoi 105281zenodo 12902*

31. Hadfield JD. 2010 MCMC Methods for Multi-Response Generalized Linear Mixed
Models: The MCMCglmm *R* Package. *J. Stat. Softw.* **33**.
(doi:10.18637/jss.v033.i02)

32. Pagel M, Meade A. 2013 Bayes Traits V2. *Comput. Program Doc. Available*
*Httpwww Evol. Rdg Ac UkBayesTraits Html Accessed 12 July 2013*

33. Pagel M. 1994 Detecting correlated evolution on phylogenies: a general method
for the comparative analysis of discrete characters. *Proc. R. Soc. Lond. B Biol. Sci.*
**255**, 37–45.

34. Borges R, Machado JP, Gomes C, Rocha AP, Antunes A. 2019 Measuring
phylogenetic signal between categorical traits and phylogenies. *Bioinformatics*
**35**, 1862–1869.

35. Barbaro L, Giffard B, Charbonnier Y, van Halder I, Brockerhoff EG. 2014 Bird
functional diversity enhances insectivory at forest edges: a transcontinental
experiment. *Divers. Distrib.* **20**, 149–159.

36. Kang C, Zahiri R, Sherratt TN. 2017 Body size affects the evolution of hidden
colour signals in moths. *Proc. R. Soc. B Biol. Sci.* **284**, 20171287.
(doi:10.1098/rspb.2017.1287)

37. Brown Jr KS, Hutchings RW. 1997 Disturbance, fragmentation, and the dynamics
of diversity in Amazonian forest butterflies. *Trop. For. Remn. Ecol. Manag.*
*Conserv. Fragm. Communities Univ. Chic. Press Chic.* **632**.

38. Somanathan H, Kelber A, Borges RM, Wallén R, Warrant EJ. 2009 Visual ecology
of Indian carpenter bees II: adaptations of eyes and ocelli to nocturnal and
diurnal lifestyles. *J. Comp. Physiol. A* **195**, 571–583.

39. Lindauer M. 2019 Hymenopteran. *Encycl. Br.*

40. Wootton RJ. 1992 Functional morphology of insect wings. *Annu. Rev. Entomol.* **37**,
113–140.

41. Skowron Volponi MA, McLean DJ, Volponi P, Dudley R. 2018 Moving like a model:
mimicry of hymenopteran flight trajectories by clearwing moths of Southeast
Asian rainforests. *Biol. Lett.* **14**, 20180152.

42. Howarth B, Edmunds M, Gilbert F. 2004 Does the abundance of hoverfly
(Syrphidae) mimics depend on the numbers of their hymenopteran models?
*Evolution* **58**, 367–375.

43. Kazemi B, Gamberale-Stille G, Leimar O. 2015 Multi-trait mimicry and the
relative salience of individual traits. *Proc. R. Soc. B Biol. Sci.* **282**, 20152127.

44. Kikuchi DW, Mappes J, Sherratt TN, Valkonen JK. 2016 Selection for
multicomponent mimicry: equal feature salience and variation in preferred
traits. *Behav. Ecol.* **27**, 1515–1521.

45. van Halder I, Barbaro L, Jactel H. 2011 Conserving butterflies in fragmented
plantation forests: are edge and interior habitats equally important? *J. Insect
Conserv.* **15**, 591–601.

46. Duarte RC, Stevens M, Flores AA. 2016 Shape, colour plasticity, and habitat use
indicate morph-specific camouflage strategies in a marine shrimp. *BMC Evol. Biol.*
**16**, 218.

47. Dimitrova M, Stobbe N, Schaefer HM, Merilaita S. 2009 Concealed by
conspicuousness: distractive prey markings and backgrounds. *Proc. R. Soc. B Biol.
Sci.* **276**, 1905–1910.

48. Olofsson M, Dimitrova M, Wiklund C. 2013 The white 'comma'as a distractive
mark on the wings of comma butterflies. *Anim. Behav.* **86**, 1325–1331.

49. Troscianko J, Skelhorn J, Stevens M. 2018 Camouflage strategies interfere
differently with observer search images. *Proc. R. Soc. B Biol. Sci.* **285**, 20181386.

50. Stevens M, Stubbins CL, Hardman CJ. 2008 The anti-predator function of
'eyespots' on camouflaged and conspicuous prey. *Behav. Ecol. Sociobiol.* **62**,
1787–1793. (doi:10.1007/s00265-008-0607-3)

51. Stevens M, Cantor A, Graham J, I.S. W. 2009 The function of animal 'eyespots':
conspicuousness but not eye mimicry is key. *Curr. Zool.* **55**, 319–326.

52. Robertson KA, Monteiro A. 2005 Female Bicyclus anynana butterflies choose
males on the basis of their dorsal UV-reflective eyespot pupils. *Proc. R. Soc. Lond.
Ser. B-Biol. Sci.* **272**, 1541–1546. (doi:10.1098/rspb.2005.3142)

53. Costello LM, Scott-Samuel NE, Kjernsmo K, Cuthill IC. 2020 False holes as
camouflage. *Proc. R. Soc. B* **287**, 20200126.

54. Cuthill IC, Stevens M, Sheppard J, Maddocks T, Párraga CA, Troscianko TS. 2005
Disruptive coloration and background pattern matching. *Nature* **434**, 72.

55. Fraser S, Callahan A, Klassen D, Sherratt TN. 2007 Empirical tests of the role of
disruptive coloration in reducing detectability. *Proc. R. Soc. Lond. B Biol. Sci.* **274**,
1325–1331.

56. Schaefer HM, Stobbe N. 2006 Disruptive coloration provides camouflage
independent of background matching. *Proc. R. Soc. Lond. B Biol. Sci.* **273**, 2427–
2432.

57. Stevens M, Cuthill IC. 2006 Disruptive coloration, crypsis and edge detection in
early visual processing. *Proc. R. Soc. B Biol. Sci.* **273**, 2141.
(doi:10.1098/rspb.2006.3556)

**Figures and tables**

**Variation in transparency degree: response to light availability or part of mimicry syndrome?**

Mónica Arias[1,2], Jérôme Barbut[2], Rodolphe Rougerie[2], Manon Dutry[1], Mireia Kohler[1], Baptiste Laulan[1], Caroline Paillard [1], Serge Berthier[3], Christine Andraud[4], Marianne Elias[2], Doris Gomez[1,3]

**Affiliations**

[1]CEFE, Univ Montpellier, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

[2]Institut de Systématique, Evolution, Biodiversité, CNRS, MNHN, Sorbonne Université, EPHE, Université des Antilles, 45 rue Buffon CP50, France

[3]INSP, Sorbonne Université, CNRS, Paris, France

[4]CRC, CNRS, MNHN, Paris, France

a.                                              b.

Figure 1. Examples of a potential leaf mimic (a. *Pseudasellodes fenestraria,* Rio Kosnipata, Peru. Photo: © Adrian Hoskins) and a bee/wasp mimic (b. Adult female *Hemaris fuciformis fuciformis*, Catalonia, Spain. Photo: © Tony Pittaway). Forewing length is ~18mm in *Pseudasellodes fenestraria*, ~24mm in *Hemaris fuciformis fuciformis.*

Figure 2. Survival of artificial prey in closed (a.) and open (b.) areas, without any transparent area (C), and with transparent elements covered by 6 (T6), 1 (T1) or non (T0) transparent film layers. Prey placed on tree trunks were monitored for 'survival' every day for 4 days.

Figure 3: Variation between open (yellow) and closed habitats (dark blue) in a) transparency degree and b) the interaction between wings (forewing: fw, square/dashed and hindwing: hw, circle/plain), wing length and transparency degree. Values larger than 60 mm were not plotted for clarity reasons but were included in the analyses.

Figure 4: Variation between bee/wasp mimics (red), leaf mimics (purple) and no mimics (blue) in a) transparency degree b) wing length c) proportion of clearwing surface, d) Ratio between length of forewing and hindwing per mimicry group and e) in the interaction wing size and transparency.

Table 1. Variation of transparency degree (mean percentage of transmitted light) according to mimicry, habitat, daytime activity, wing size, wing and proportion of clearwing surface excluding (in the left) or including interactions between them (in the right).

| | Bayesian | Linear mixed model | | | | Bayesian | Linear mixed model | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | Estimate [CI95%] | Estimate ± se | DF | $t$ | | Estimate [CI95%] | Estimate ± se | DF | $t$ | |
| (Intercept) | 60.09[31.06;88.73] | 60.61±3.92 | 502 | 15.48 | *** | 62.24[30.79;93.17] | 64.94±5.6 | 488 | 11.59 | *** |
| Activity (D>N) | 10.34[21.66;-0.51] | 1.99±2 | 102 | 0.99 | | -11.43[-22.67;-0.12] | 2.98±2.01 | 102 | 1.48 | |
| Wing (F>H) | 7.92[9.57;6.28] | 3.67±0.39 | 502 | 9.45 | *** | -7.7[-12.43;-2.83] | 1.92±1.31 | 488 | 1.47 | |
| Clearwing proportion | 0.27[0.22;0.32] | 0.26±0.02 | 502 | 10.88 | *** | 0.04[-0.12;0.2] | -0.11±0.07 | 488 | -1.59 | |
| Mimicry:bw>all | 21.45[9.4;33.57] | 3.97±1.98 | 102 | 2 | * | 41.29[19.43;62.83] | 9.9±2.84 | 102 | 3.49 | *** |
| Mimicry:leaf<non-mimics | 5.57[-9.84;20.81] | -5.05±3.07 | 102 | -1.64 | | -3.17[-25.14;19.52] | 1.62±4.21 | 102 | 0.39 | |
| Habitat (Open>Closed) | -7.47[-17.14;1.92] | -5.96±3.77 | 102 | -1.58 | | -17.21[-31.38;-3.18] | -12.07±5.77 | 102 | -2.09 | * |
| Wing length | -0.5[-0.64;-0.36] | -0.43±0.07 | 502 | -6.61 | *** | -0.45[-0.71;-0.2] | -0.3±0.21 | 488 | -1.4 | |
| Activity (D>N):Wing Wing(F>H) | - | - | - | - | | 3.1[0.54;5.49] | 0.82±0.28 | 488 | 2.95 | |
| Mimicry (bw>all):Clearwing prop. | - | - | - | - | | 0.04[-0.15;0.22] | 0.09±0.03 | 488 | 2.99 | ** |
| Mimicry (leaf<n.m):Clearwing prop. | - | - | - | - | | -0.46[-0.68;-0.24] | 0.24±0.05 | 488 | 4.8 | *** |
| Habitat(O>C):Wing length | - | - | - | - | | -0.04[-0.31;0.24] | -0.28±0.14 | 488 | -2.05 | * |
| Habitat(O>C):Wing(F>H) | - | - | - | - | | 1.88[-3.24;7.02] | -0.64±1.17 | 488 | -0.55 | |
| Wing length:Wing (F>H) | - | - | - | - | | -0.02[-0.14;0.1] | 0±0.03 | 488 | -0.09 | |
| W.length:Clearwing prop. | - | - | - | - | | 0[0;0.01] | 0±0 | 488 | 0.89 | |
| Mimicry (bw>all):W.length | - | - | - | - | | -1.08[-1.99;-0.14] | -0.46±0.15 | 488 | -3.03 | ** |
| Mimicry (leaf<n.m):W.length | - | - | - | - | | 0.82[0.2;1.44] | -0.55±0.14 | 488 | -4 | *** |
| Clearwing prop:Habitat (O>C) | - | - | - | - | | 0.34[0.24;0.44] | 0.31±0.05 | 488 | 6.51 | *** |
| Clearwing prop:Wing (F>H) | - | - | - | - | | 0.03[-0.04;0.1] | -0.02±0.02 | 488 | -1.31 | |
| Mimicry (bw>all):Wing(F>H) | - | - | - | - | | -4.6[-14.17;5.06] | 1.02±0.79 | 488 | 1.28 | |
| Mimicry (leaf<n.m):Wing(F>H) | - | - | - | - | | 6.04[-1.19;13.2] | 2.19±0.81 | 488 | 2.69 | ** |
| Wing (F>H): Habitat (O>C):Wing length | - | - | - | - | | -0.26[-0.46;-0.06] | 0.14±0.05 | 488 | 3.02 | ** |
| Phylogenetic variance | 951.7[702.02;1276.09] | - | - | - | | 968.02[725.74;1287.29] | - | - | - | |

| Residual variance | 951.7[702.02;1276.09] | - | - | - | 67.24[61.24;73.71] | - | - | - |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |

Bayesian regression and linear mixed model to explain transparency degree include daytime activity (nocturnal/diurnal), wing length in mm, proportion of clearwing area (%Clearwing), type of habitat (open/closed), mimicry syndrome (bee/wasp -, leaf - or no mimics), wing (fore-/hindwing) and the interactions: mimicry syndrome and proportion of clearwing area, mimicry syndrome and wing size, proportion of clearwing area and habitat type, wing size and habitat type, wing size and proportion of clearing area, wing and wing length and proportion of clearwing area, wing length and habitat as explanatory variables. Wing measurements nested in species was considered as random effect. Linear mixed models take into account contrast between factors that are not considered by Bayesian estimations, explaining differences in the sign of the estimates between analyses. Bayesian estimates whose confidence interval did not include 0 and LMM p values below 0.05 are statistically significant. *** stands for p<0.001, ** for p<0.01 and * for p<0.05.

Table 2. Results of coevolution tests between habitat, mimicry and daytime activity

| Tested characters | Mimicry levels included | L.Dep. | L.Indep. | LRT | df | p | |
|---|---|---|---|---|---|---|---|
| Daytime activity/ mimicry | Bee/wasp & leaf mimics | -30 | -36.78 | 13.56 | 4 | 0.009 | ** |
| | No- & leaf mimics | -101.82 | -101.82 | 0.01 | 4 | 1 | |
| | No- & bee/wasp mimics | -97.75 | -97.88 | 0.26 | 4 | 0.993 | |
| Habitat / mimicry | bee/wasp & leaf mimics | -34.86 | -36.78 | 3.85 | 4 | 0.427 | |
| | No- & leaf mimics | -102.27 | -102.34 | 0.13 | 4 | 0.998 | |
| | No- and bee/wasp mimics | -99.73 | -101.26 | 3.05 | 4 | 0.549 | |
| Habitat/daytime activity | No-, bee/wasp- & leaf mimics | -144.8 | -141.76 | 6.08 | 4 | 0.193 | |

Likelihood of dependent (L. Dep. Coevolution, 4 parameters estimated) and independent (L. Indep. 8 parameters estimated) evolution between pairs of binary characters obtained from BayesTraits. As mimicry has three levels, subsets including two of them were used and reported in the second column. A significant p-value indicated that the dependent model did significantly better than the independent model, indicating a correlated evolution between the traits. ** stands for statistical significance below 0.01