Estimation of the SNP mutation rate in two vegetatively propagating species of duckweed

George Sandler[1], Magdalena Bartkowska[1], Aneil F. Agrawal[1,2]*, Stephen I. Wright[1,2]*

[1]Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street, Toronto, ON M5S 3B2, Canada; [2]Center for Analysis of Genome Evolution and Function, University of Toronto, 25 Willcocks Street, Toronto, ON M5S 3B2, Canada

Author for correspondence

George Sandler

Tel: +1 416 6172717

Email: george.sandler@mail.utoronto.ca

*These authors contributed equally to this work.

Key words: Mutation rate, vegetative reproduction, duckweed, asexual, effective population size

Total word count	6159	No. Figures	4
Abstract	128	No. Tables	0
Introduction	1007	No. Supporting information files	1
Materials and Methods	1926		
Results	1053		
Discussion	2173		
Acknowledgements	41		

Abstract

Mutation rate estimates for vegetatively reproducing organisms are rare, despite their frequent occurrence across the tree of life. Here we report mutation rate estimates in two vegetatively reproducing duckweed species, *Lemna minor* and *Spirodela polyrhiza*. We use a modified approach to estimating mutation rates by taking into account the reduction in mutation detection power that occurs when new individuals are produced from multiple cell lineages. We estimate an extremely low per generation mutation rate in both species of duckweed and note that allelic coverage at *de novo* mutation sites is very skewed. We also find no substantial difference in mutation rate between mutation accumulation lines

25   propagated under benign conditions and those grown under salt stress. We discuss the

26   implications of interpreting mutation rate estimates in vegetatively propagating organisms.

Introduction

Most research on the evolution of mutation rates has focused either on sexually

reproducing eukaryotes or unicellular organisms, both of which feature a single cell phase as

part of their life cycle. However, a diverse array of organisms reproduce either through

clonal budding, fission or vegetative growth, whereby a single cell phase is not imposed

every generation (Bell, 1982). This mode of reproduction potentially allows multiple cell

lineages to be transmitted from parent to offspring, complicating the process of genotyping

individuals. This happens because when individuals composed of a mosaic of cells are

sequenced, the mean number of sequencing reads supporting non-reference mutations is

no longer 50%. Such a skew in allelic coverage makes it harder to distinguish true mutations

from sequencing errors (Cibulskis *et al.*, 2013), complicating the assessment of power when

calculating per base pair mutation rates. Even if a cellular mutation rate can be calculated

for an organism with multiple cell lineages, it becomes more challenging to use this

parameter in population genetics analyses as mutations can potentially be lost within an

organism before truly contributing to population level genetic diversity. Previous theoretical

work modeling mutation load in organisms with multiple cell lineages has suggested that

cell lineage selection can significantly reduce mutation load by purging deleterious

mutations during somatic growth (Otto & Orive, 1995). As most new mutations are thought

to be deleterious (Sturtevant, 1937; Eyre-Walker & Keightley, 2007) this type of selection

might skew the level of genetic diversity observed in organisms with vegetative

reproduction compared to the level expected given their per base pair mutation rates.

Previous studies have investigated the rate of somatic mutations in plants where

multiple cell lineages can segregate within a generation (Watson *et al.*, 2016; Schmid-Siegert

et al., 2017; Plomion *et al.*, 2018; Wang *et al.*, 2019). While somatic mutations can be

transmitted from generation to generation in plants (Plomion *et al.*, 2018; Wang *et al.*,

2019), if somatic growth is followed by sexual reproduction, a single cell bottleneck is

nonetheless imposed on any segregating variation within the soma, removing the

persistence of multiple cell lineages across generations. This is however not the case for

organisms reproducing through vegetative growth, budding or fission. Despite their

frequency across the eukaryotic tree of life, almost no per-base-pair mutation rate

estimates exist for organisms procreating through such modes of reproduction. One recent

study in a vegetatively growing fairy-ring mushroom reported very low mutation rates per

mitotic cell division (Hiltunen *et al.*, 2019). The authors of this work used simulated

mutations to assess the level of power they had to detect low frequency *de novo* mutation

in this dataset, improving their estimate of the fairy ring mushroom mutation rate.

Here we report mutation rate estimates in two species of duckweed (*L. minor* and *S.*

polyrhiza). Both species are free-floating, facultatively sexual aquatic plants. While

duckweed can produce seed though sexual reproduction, most growth occurs vegetatively

via clonal budding from two pouches present in the duckweed frond (Landolt, 1986). As a

result, while these species are found all across the globe and likely have enormous census

population sizes, allozyme and genomic analyses have revealed low levels of genetic

diversity within local populations (Cole & Voskuil, 1996; Ho, 2018; Xu *et al.*, 2019; Ho *et al.*,

2019). Work by Xu et al (2019), has estimated the per base pair mutation rate in a genotype

of *S. polyrhiza* growing in the field, finding an extremely low rate of mutation. However,

their analysis did not take into account the fact that duckweed individuals are likely

composed of a mosaic of cell lineages during periods of asexual growth, potentially leading

to an underestimate of the true mutation rate.

Studying two duckweed species allows us to contribute to three other questions in

mutation rate evolution research. First, our mutation rate estimates provide another species

to add to the existing set of species with mutation rate estimates that, collectively, allow for

testing the theory that selection against mutators should be most efficient in species with

large effective population sizes (Sung *et al.*, 2012; Lynch *et al.*, 2016). Duckweeds are useful

additions to this set as previous work has suggested that the effective population size (N_e) of

S. polyrhiza is on the order of a million individuals (Ho et al. 2019; Xu et al. 2019). Second,

the inclusion of two facultatively sexual species that differ in their degree of sexuality allow

us to preliminarily investigate the effect of recombination on the evolution of mutation

rates. Genomic and allozyme patterns have suggested that *L. minor* undergoes bouts of

sexual reproduction more often than *S. polyrhiza*, a pattern that is in line with flowering

observations of these species in the field (Hicks, 1932; Landolt, 1986). Theoretical work has

shown that when recombination breaks apart associations between mutator alleles (that

elevate mutation rate) and the mutations they produce, mutation rates can evolve in

several ways. On one hand selection against mutators in more sexual populations may be

relaxed as they no longer remain linked to new deleterious mutations (Kimura, 1967; Leigh,

1970). Alternatively mutator alleles can spread when recombination is sufficiently low if

they hitch hike along with any beneficial mutations they produce (André & Godelle, 2006).

Finally, environmental stress is known to increase mutation rates in bacteria in a process

known as stress-induced mutagenesis (Foster, 2007). A few examples of stress increasing

mutation rates are known in eukaryotes (Matsuba et al. 2013; Jiang et al. 2014; Sharp and

Agrawal 2016; but see Saxena et al. 2019), however it is unclear how general this

phenomenon is. We performed our experiment both under a control and salt stress

treatment to test whether stress-induced mutagenesis is a common phenomenon in plants.

We estimated the mutation rate in 46 asexually propagated mutation accumulation

lines, including two genotypes of *S. polyrhiza* and one genotype of *L. minor*. We report an

exceptionally low rate of mutation in both species of duckweed and note a pattern of

skewed allelic counts at *de novo* sites that suggests the presence of multiple segregating cell

lineages in vegetatively reproducing duckweed.

Materials and Methods

<u>Mutation accumulation and DNA extraction</u>

MA lines were started for three genotypes in April 2014 and propagated for

approximately 60 generations. Two *Spirodela polyrhiza* genotypes were used: GP23 from

Grenadier Pond, Toronto, Canada and CC from Cowan Creek, Oklahoma, USA. A single

genotype of *L. minor* (GPL7) was also isolated from Grenadier Pond, Toronto, Canada. For

each genotype (CC, GP23 and GPL7), we established 16 MA lines. We generated each line

from a single maternal plant, which was started by isolating two fronds from each genotype

culture. Because daughter fronds are generated iteratively, we grew and isolated daughters

tracking pedigree until a minimum of 16 daughter fronds paired by generation were

available for each of the three genotypes (arising from a single starting maternal frond).

Daughters in each frond pairs (matched for generation relative to maternal frond) were

assigned to one of two growth medium treatments (salt stress and control). Daughters are

produced from two pockets of meristem tissue on either side of the maternal frond and

mature daughter fronds remain attached to the maternal plant via a stipule for a short time

(Landolt, 1986). To ensure that each generation was propagated with a daughter frond, we

separated the daughter from the maternal frond as soon as the daughter began producing

her own frond. Each line was checked for mature daughter fronds every two days. The first

daughter produced was used whenever possible. MA lines were propagated in 0.5X

Appenroth liquid growth medium (Appenroth *et al.*, 1996) at 24°C with 12 hours of light per

day. Salt stress lines were supplemented with 25mM of NaCl for *S. polyrhiza* and 50mM of

NaCl for *L. minor*. Prior to the start of the MA experiment, we performed growth assays to

establish stressful NaCl levels for both species. The chosen salinity levels caused duckweed

fronds to become patchy and thin but still allowed for continual asexual propagation.

After the termination of the MA experiment we allowed MA lines to continue

growing for several generations to obtain enough plant material to perform CTAB DNA

extractions.

<u>Sequencing and filtering</u>

We sequenced the MA lines at the McGill Innovation Centre. Illumina HISeq 2000

sequencing with 100bp paired end reads was used for both *S. polyrhiza* genotypes while

Illumina HISeq 2500 sequencing with 125bp paired end reads was used for the *L. minor*

genotype.

Paired end reads were mapped to the *S. polyrhiza* and *L. minor* reference genomes

(Van Hoeck *et al.*, 2015; Cao *et al.*, 2016) using the Burrows-Wheeler aligner (BWA) 0.717

using the BWA-MEM option (Li & Durbin, 2009). We then used Picard to remove duplicate

reads before calling indels using the HaploTypeCaller tool in GATK 3.7 (McKenna *et al.*,

2010). Next, we used the IndelRealigner tool in GATK to perform indel realignment. Finally,

we used BCFtools (1.6) (Li, 2011) to create mpileup files for the realigned output from GATK

and to call SNPs and short indels (indels no more than 10bp) .

We first filtered out sites with unusually high or low coverage. We did this by

eliminating sites that had coverage outside +/- 200x median coverage (summed across all

lines) in each genotype individually. We visualised relatedness between our lines using a

PCA plot created from heterozygous sites present in our MA lines in R (v5.3.5) (R Core Team,

2019) using the package SNPRelate (Zheng *et al.*, 2012). In doing so we discovered two

major outliers in one of our *S. polyrhiza* genotypes (CC) suggesting that these two lines were

cross contaminated. We subsequently removed these two lines from our analysis.

Our next round of filtering aimed to remove low quality regions of the genome that

contain unusually high amounts of in-phase heterozygous variants (e.g. Fig. S1). Such

regions likely represent collapsed duplications in the reference genome that map poorly to

an incorrect genomic coordinate. These variants are highly reference-biased in their allelic

coverage likely due to the poor mapping of reads that contain many differences relative to

the reference genome. To remove such regions, we first created a consensus genotype for

each set of lines; if more than one line in a given genotype supported the existence of a

heterozygote at a site, that site was designated as heterozygous in the consensus genotype.

We then performed a sliding window analysis on heterozygosity on each consensus

genotype. We used 1000bp windows with a 100bp step. After trying a variety of filtering

criteria, we decided to designate regions of the genome as callable if there existed no more

than 10 heterozygous calls in a 1000bp window in each *S. polyrhiza* genotype and no more

than 5 heterozygous calls per 1000bp window in the *L. minor* genotype. We used more

stringent criteria in *L. minor* due to the lower quality of the genome assembly. These cut-

offs represent a trade-off between eliminating problematic, variant-rich areas of the

genome and excluding well assembled genomic areas with higher than average diversity.

After filtering, around 100Mb of the genome was retained as callable in each of the three

genotypes. This filtering step greatly improved the allelic coverage of ancestral

heterozygous sites by removing suspected hidden duplications that map poorly to the

reference (Fig. S2).

Next, we called putative *de novo* mutations in the remaining callable regions. Within

each set of lines, we picked sites where one line had a heterozygous genotype, with at least

5 reads supporting the non-reference base, but all other lines supported a homozygous

genotype. We then extracted such sites from the mpileup file used to call genotypes. This

was done as the pileup file contains reads that are filtered out during genotype calling but

are useful in our case as they can lead to the elimination of false positive mutations. We

filtered putative *de novo* mutations using the mpileup file in two ways. First, if a line other

than the one which contained the *de novo* mutation had any reads which supported the *de*

novo base call we excluded the site. Second, if a site with a *de novo* base call contained

reads with more than two non-reference bases across all samples, we also excluded the site.

We did this to exclude sites where a high rate of sequencing errors might have occurred. We

used this cut-off based on the observation that at sites where all lines supported a

homozygous genotype, the vast majority of sites contain no more than one alternate base

call (again likely due to sequencing errors which can be observed in the mpileup file). The

remaining putative *de novo* mutations that passed these filtering criteria were visually

inspected in the Integrative Genomics Viewer (IGV) (Robinson *et al.*, 2011). We excluded a

few mutations which appeared on reads in complete linkage with other non-reference bases

(an indication of hidden genomic duplications) or on reads that looked like the product of

PCR or sequencing errors (see Fig. S3-5 for examples).

<u>Power analysis</u>

To calculate the mutation rate, we first needed to know how much power we had to

detect *de novo* mutations at our callable sites. To assess power, we first obtained a list of

sites where we knew we had non-zero power to detect mutations, this included sites where

all lines within a genotype supported a homozygous reference base call and no more than

one alternate base was present in the mpileup file (one less than in the case if a *de novo*

mutation was present). We then randomly sampled 500,000 such sites from each genotype

independently, and randomly chose a line where a mutation could have happened. We

randomly eliminated a third of the sites where one alternate base was present in the

mpileup file as our filtering criteria would eliminate true *de novo* mutations if another line

by chance contained a sequencing error which matched the *de novo* base call (i.e., we

assume the probability of this occurring is 1/3). Of the remaining sites, we assigned how

many reads would support the *de novo* base call by drawing from a binomial distribution

with a success rate of 50%, 34%, 26%, 20% and 10%. These different values were chosen to

represent a range of frequencies a mutation may be found due to the inheritance of

multiple cell lineages in asexual reproduction. This is similar to the approach taken by

Hiltunen et al. (2019) when calculating mutation rates in vegetatively growing fairy-ring

mushrooms.

Our estimate of power was the proportion of sites (out of the original 500,000) that

had at least 5 reads supporting the *de novo* mutation (for each possible binomial success

rate that we tested). We then multiplied our power estimates by the number of callable

sites in the genome. Then separately for each line, we multiplied the adjusted number of

callable sites by the number of MA generations and summed these values across all lines in

a given genotype (split by treatment). This provided us with a denominator for our mutation

rate calculation. Our final step was to divide the number of *de novo* mutations identified in

each genotype (split by treatment) by this denominator. The *de novo* mutation count was

adjusted for false positives identified during mutation validation.

Calling indels

We scanned our lines for *de novo* indels in the same way as we searched for point

mutations with one key modification. We used the same regions of the genome that we had

previously assessed as "callable"; however, we only considered *de novo* indels if they were

at least 2000 bp away from any other indel in any other line of the same genotype. This

filtering step was required to avoid false positive indel calls which appear due to spurious

mapping patterns in repetitive regions.

Mutation validation

We only had tissue from *S. polyrhiza* to perform validation on putative *de novo*

mutations. We extracted DNA from our MA lines using a Qiagen DNeasy kit. Afterwards, we

designed primers for 14 SNPs and one indel found in our two *S. polyrhiza* genotypes. We

performed PCR reactions using FroggaBio PCR mastermix which were then sent off for

Sanger sequencing at Eurofins Genomics. We inspected the Sanger sequencing

chromatograms in FinchTV 1.4.0 (Geospiza, Inc.; Seattle, WA, USA;

http://www.geospiza.com) to see if we could detect a peak suggesting the presence of a

mutant base in the focal MA line and checked that the base was not present in a different

MA line of the same genotype as a negative control. One line from the *S. polyrhiza* CC

genotype contained a massive overabundance of putative *de novo* mutations (of the 31

mutations found in 14 lines of the CC genotype, 9 were identified from this one line); all 3

variants that we attempted to Sanger sequence validate from this line failed. Ultimately, we

excluded this CC line from all the results presented here. Excluding that line, 11 total SNPs

that were checked with Sanger sequencing and used to correct for the false positive rate in

mutation detection in all our *S. polyrhiza* lines.

Statistical analysis

We calculated 95% confidence intervals for our mutation rates using the Agresti

Coull method implemented in the R package "binom" (Dorai-Raj, 2014). We tested for

significant differences between our salt and normal treatment lines using independent Chi-

square tests in each genotype. To test for differences between *L. minor* and *S. polyrhiza* we

merged the number of callable sites (scaled by power) and number of mutations in the two

S. polyrhiza genotypes (separately for each treatment), and also applied Chi-square tests to

test for significance in mutation rate variation between species and conditions.

Data availability

All sequencing data will be deposited to the Sequencing Read Archive at NCBI upon

acceptance of this manuscript for publication. Supplementary material has been uploaded

as a separate file.

Results

De novo mutations

In total, we identified 24 and 17 putative *de novo* mutations in *S. polyrhiza*

genotypes GP23 and CC, respectively and a further 29 mutations in *L. minor* (Supplemental

table 1). When inspecting the putative *de novo* mutations in IGV, we observed that most

mutant sites exhibited highly reference biased allelic counts (over 50% of reads support the

reference base call). On average, mutant sites contained 26%, 29%, and 34% mutant reads

in genotypes GP23, CC (*S. polyrhiza*) and GPL7 (*L. minor)* respectively. We implemented

several filtering and quality control steps to ensure these mapping patterns were not a

result of sequencing error or genome mis-assembly. First, we masked areas of the genome

that had odd coverage patterns or were enriched for heterozygous calls to avoid areas

containing hidden genomic duplications. The heterozygous variants that pass these filtering

criteria and are present in the ancestral genotypes (i.e. are shared by all MA lines within a

genotype) show relatively normal patterns of reference and non-reference allele coverage.

Second, we only considered *de novo* mutations at sites with minimal sequencing errors. This

step eliminated problematic areas of the genome prone to genotyping error. Finally, we

performed Sanger sequencing validation on 11 putative *de novo* mutations (in *S. polyrhiza*)

of which 6 were validated with both positive and negative controls. The pattern of reference

bias in our *de novo* mutations was also present in our Sanger sequencing chromatograms;

the reference base peak was generally much larger than the mutant peak. When we plotted

Illumina mutant base frequency vs. Sanger mutant peak height (standardized by reference

peak height), we observed strong concordance for our 6 validated mutations (R^2 = 0.56; Fig.

1) suggesting that these are not spurious mapping patterns or sequencing errors but rather

reflect a true bias in mutation abundance. When we inspected our PCR products with gel

electrophoresis we observed single, clean bands of the expected product size so we

consider it unlikely that multiple sites in the genome were amplified leading to odd mapping

patterns at our *de novo* mutation sites. One alternative is that reference biased mapping

patterns may be due to some bias in the PCR amplification process. We do not believe this is

likely however since our primers were designed for sequences that were flanking the site of

de novo mutations that should be identical whether a *de novo* mutations is present or not.

Fig. *1 Proportion of de novo bases from Illumina reads vs. relative peak height of mutant*

base in Sanger sequencing data in Spirodela polyrhiza. R^2 = 0.57.

The majority of mutations were C -> T transitions in both species of duckweeds (Fig.

2) in concordance with patterns found in previous mutation rate studies (Ossowski *et al.*,

2010; Thomas *et al.*, 2018). Transitions were more common than transversions, with the

average ti/tv ratio being 2.04 for *S. polyrhiza* and 4.5 for *L. minor* propagated in standard

medium. This is consistent with previous results reported in *S. polyrhiza* by Xu et al. (2019)

where three C -> T and one C -> A mutations were detected and validated.

Our analysis uncovered only one putatively *de novo* short indel that turned out to be

a false positive based on our Sanger sequencing validation analysis.

Fig. 2 *De novo mutation spectra in two species of duckweed (Spirodela polyrhiza, Lemna*
minor).

<u>Mutation rate comparisons between *S. polyrhiza* and *L. minor*</u>

Our estimate of the per base pair SNP mutation rate highly depends on our power to

detect mutations, which in turn depends on factors such as total read depth and the

number of reads that support a *de novo* mutation base-call (See Supplemental tables 2-5).

Heterozygous SNPs are expected to be supported by around 50% of reads in organisms with

a single cell phase. In our case, on average, 27% (*S. polyrhiza*) and 34% (*L. minor*) of reads

supported *de novo* heterozygous mutations. To improve our estimate of the mutation rate,

we assumed that we should only expect to find mutations that are on average supported by

27% (in *S. polyrhiza)* or 34% (in *L.* minor) of reads at each mutant site (see methods for more

details of power analysis). In this case, our point estimate of the mutation rate for plants

grown in standard medium is 8.33E-11 for *S. polyrhiza* and 8.66E-11 for *L. minor* (Fig. 3). For

plants grown in stressful salt medium our point estimates for the mutation rate are 9.20E-11

for *S. polyrhiza* and 16.9E-11 for *L. minor*.

While the estimated mutation rate is higher for *L. minor* in both conditions, the

difference between the two species is non-significant (chi-square test: p = 1 and p = 0.22 for

control and salt stressed conditions). The difference in mutation rate in the two species is,

however, very sensitive to the expected frequency of reads that support the mutant base.

For example, if our power estimate is based on the assumption that *de novo* mutations are

expected to constitute only 10% of reads, the mutation rate under normal conditions

increases to 4.93E-10 in *S. polyrhiza* and 16.8E-10 in *L. minor*. Our mutation rate estimates

were not significantly different between our two individual *S. polyrhiza* genotypes (chi-

square test: p = 1 and p = 0.29 for control and salt stressed conditions). Supplemental tables

3-4 give the inferred mutation rates with alternative assumptions for the expected

proportion of non-reference reads at sites harbouring *de novo mutations*.

Fig. 3 *Mutation rate estimates in two species of duckweeds duckweed (Spirodela polyrhiza,*

Lemna minor). Power adjusted to account for multiple cell lineages. Error bars show Agresti-

Coull 95% confidence intervals.

<u>Effect of salt stress on mutation rate</u>

There were no significant differences in the mutation rate between lines propagated

in normal and salt stress medium for either of our three genotypes (Fig. 4, chi-square test: p

= 0.81, p = 0.51, p = 0.29 for genotypes GP23, CC, GPL7 respectively). There was also no

consistent trend in the change of mutation rate with the addition of salt stress as genotype

GP23 appeared to have a decreased mutation rate while genotypes CC and GPL7 appeared

to have an increase in the mutation rate with the addition of salt stress.

Salt stress also did not have a significant impact on the ti/tv ratio. In two of the

genotypes (1.33 -> 2.67 *S. polyrhiza* CC and 4.5 -> 8 *L. minor*), salt stress increased the ti/tv

ration. In *S. polyrhiza* genotypes GP23 salt stress decreased the ti/tv ration (2.75 -> 2.00)

although the difference was not significant in either of the three genotypes (chi-square test:

p > 0.5 all cases). These results run counter to previous work in *A. thaliana* where salt stress

was found to increase the de-novo mutation rate ~two-fold and lower the ti/tv ratio (Jiang

et al., 2014).

Fig. 4 *Effect of salt stress on the mutation rate in two genotypes of Spirodela polyrhiza and*

one genotype of Lemna minor. Error bars show Agresti Coull 95% confidence intervals.

Discussion

Using whole genome sequencing on 36 MA lines we report the *de novo* SNP

mutation rate in two species of duckweeds under both control and stressful growth

conditions. Our results are in agreement with previous estimates of a low mutation rate in *S.*

polyrhiza (Xu *et al.*, 2019) while also extending this observation to another duckweed

species, *L. minor*. An important result in our study is that *de novo* mutations appear to have

considerably reference biased genomic coverage in both duckweed species. We believe that

this pattern is not indicative of sequencing or genome assembly errors but rather is a by-

product of vegetative reproduction for several reasons. First, upon our inspection of the

data in an independent study of mutation rates in *S. polyrhiza* by Xu et al.; we noticed

similar levels of reference bias in Illumina short-read sequencing data at validated *de novo*

mutations. Second, our own validation with Sanger sequencing showed that mutations with

a higher reference bias in the Illumina dataset tended to have higher reference base peaks

in their Sanger sequencing chromatograms and vice versa. Moreover, after filtering, most

ancestral heterozygous sites that were shared by all lines in the three clonal genotypes were

not reference biased in such a manner and those that were, were found in regions highly

enriched for non-reference base calls. None of our putative *de novo* mutations are found in

such regions; due to the low genetic diversity in duckweed, *de novo* mutations were

generally the only heterozygous variants present in the immediate genomic area. This

means that mapping bias due to divergence from the reference is unlikely to cause the

strong allelic coverage bias we observed. Finally, a study of mutations in a vegetatively

growing fairy-ring mushroom also reported reference biased allelic counts at *de novo*

mutations (Hiltunen *et al.*, 2019). Therefore, the reference bias in our mutations is most

likely a signature of the segregation of multiple cell lines from generation to generation in

our duckweed MA experiment. Given that vegetative reproduction through clonal budding

is the main form of reproduction in duckweed (Landolt, 1986), it seems reasonable that

duckweed might not undergo a single cell phase for prolonged periods of time.

Simple models can be helpful in clarifying how multi-cell descent during vegetative

reproduction may affect mutation rates and our estimation of them. Assume that clonal

reproduction occurs by n parental cells forming a diploid offspring (generation 1). If a

mutation occurs in one of these n cells, because of multi-cell descent, the offspring will be a

genetic mosaic (i.e., not all cells will be genetically identical). The mutation begins at a

frequency of $1/2n$ in the offspring. With conventional single-cell descent, where $n = 1$, a new

mutation is expected to be at 50% frequency but the frequency will be lower than 50%

whenever $n > 1$. The n cells multiply in some (unknown) growth pattern to produce the

mature offspring, which then itself reproduces. In the absence of detailed knowledge of

developmental growth trajectories, we do not know what the representation of the original

n cell lineages will be in the subsequent generation.

We can consider two simple scenarios: If only one of these original cell lineages gives

rise to the next set of n cells used to produce the next generation (generation 2), then the

mutation will either be completely lost from the lineage (if it was not in the chosen cell

lineage) or it will be present in heterozygous state in all cells of future generations (if it was

present in the chosen cell lineage). In this first scenario, the genetic mosaicism resulting

from multi-cell descent persists only a single generation. Thus, in an experiment such as

ours, only mutations occurring in the very last generation will be affected by this issue (i.e.,

multi-cell descent is a trivial issue in this case). A second scenario is that all cell lineages

grow at equal rates and, each generation, n cells are chosen at random to form the next

daughter. This is conceptually similar to the process of coalescence in a population (of cells

here rather than individuals) of size n. Thinking backwards in time, all cells must eventually

trace their coalescent history to a single cellular ancestor, but it may take many generations

for this to happen (i.e., $2n$ generations, on average, but with variance proportional to n^2).

Thinking forwards in time, a mutation occurring in one of the n cells forming the generation

1 offspring may be present for many future generations, possibly becoming quite common

within individuals, before eventually being present in all future cells (with chance $1/n$) or

none of them (with chance $(n - 1)/n$). With even modest values of n (e.g., $n = 8$-80), genetic

mosaicism can persist for many generations. Mutations that will eventually be present in all

lineages—as well as those that will eventually be eliminated—will thus be found below the

50% frequency expected in a "traditional" (i.e., non-mosaic) heterozygote. Although this

second scenario as formulated here is unrealistically simple (e.g., random and independent

choice of n cells for each generation), it illustrates how multi-cell descent can have

consequences across multiple generations. Though developmental growth trajectories in

duckweed are insufficient to formulate a more realistic model, we suspect that multiple cell

lineages persist across multiple generations in duckweed and is responsible for the clear

bias towards mutant SNPs being less than 50%. Recent work on segregating mutations in a

single *Zostera marina* seagrass clone has made similar arguments to this model (Yu et al.,

2020). In their study, Yu et al. uncovered a large class of reference biased SNPs that were

present in some but not all sampled *Z. marina* ramets. By reconstructing the genealogical

relationship of the sampled ramets, the authors demonstrated that such SNPs changed in

frequency during vegetative growth, with some reaching heterozygous fixation in specific

ramets. The authors argued that this data was well explained by a model of "somatic drift"

whereby *de novo* mutations arise at low frequency within a single cell lineage before

ultimately either reaching fixation or being lost.

Calculating mutation rates in organisms with multiple segregating cell lineages poses

a technical challenge due to the difficulty of assessing power when mutations are present in

only a subset of the cells of an individual. We implemented a method that takes into

account the fraction of reads we expect to support a *de novo* mutation by using observed

mapping patterns of putatively *de novo* mutations, similar to the approach used by Hiltunen

et al. (2019). This method considers the fact that we have reduced power to detect recently

arisen mutations that are at low frequency within their MA lines, giving a more accurate

estimate of the SNP mutation rate. This approach is an admittedly crude attempt to address

the problem. Rather than assuming the expected frequency of the mutant allele is 50%, we

simply choose a lower value based on the observed coverage at putative *de novo* mutations.

In reality, this lower value is unknown and will differ among mutations depending upon

when each mutation arose. Nonetheless, our approach is an improvement over completely

ignoring the issue. Moreover, the variation in mutation rate estimates inferred using

different values for the assumed expected mutant frequency provides some sense of the

sensitivity of these estimates to this assumption.

A natural consequence of the lack of a single cell phase is that from a population

genetics perspective, the mutation rate becomes a harder parameter to interpret. On one

hand, we may be interested in calculating the mutation rate that captures every new

mutation that has arisen in a clonal bud. Alternatively, from an evolutionary perspective, we

might be only be interested in mutations that will not only arise in a clonal bud but also

persist in a future clonal descendant such that they contribute to population level genetic

diversity. Aside from random inter-cell lineage "drift" one possible force that may bias which

mutations are eventually lost before fixing in a clonal lineage is cell lineage selection which

may purge deleterious *de novo* mutations (Otto & Orive, 1995). In principle, we might be

able to calculate an "evolutionarily relevant" mutation rate by performing long term

mutation accumulation experiments allowing the majority of *de novo* mutations to either be

lost or to have fixed within a clonal lineage such that all cells in any clone will be either fully

homozygous (in the case of loss) or heterozygous (in the case of fixation). However, from a

practical standpoint, it is hard to know *a priori* how many generations will be necessary for

this process to occur. In practice, we could attempt to estimate the frequency of *de novo*

mutations that have become sufficiently common such that they are likely to not be lost

before fixation leading to an estimate of an evolutionarily relevant mutation rate. For

example, we could assume that mutations found in at least 50% of cells (i.e. at a frequency

of at least 25% in a clonal bud) are more likely than not to eventually fix in their clonal

lineage, being found in 100% of cells in some future generation. In reality, some of these

mutations are still likely to be lost before they have a chance to fix while simultaneously

some mutations that are below 50% frequency might still reach fixation in the future. More

generally, the lower the frequency cut-off we use for mutations that are "likely" to fix, the

better the chance that we capture all of the mutations that will reach fixation with the

caveat that we will also be capturing more mutations that will eventually be lost.

Our estimate of the mutation rate in *S. polyrhiza* (9.20E-11 per bp per gen.) is similar

to the estimates reported by Xu et al. (7.92E-11 per bp per gen.), however there are several

important differences between our studies. First, our MA experiment was conducted only in

the lab (but under two growth conditions), while Xu et al. placed MA lines both in the lab

and in the field. While we observed multiple mutations in our lab MA lines, Xu et al. report

no *de novo* mutations in their lab MA lines, likely due to the lower number of MA

generations in their study. Thus, despite the similarity in our mutation rate estimates, they

come from very different environmental settings. Another critical difference is that Xu et al.

used ancestral heterozygous sites to estimate their power to detect *de novo* mutations

which are not strongly reference biased. This suggests that the estimate from Xu et al.,

while conducted in a more naturally realistic environment is likely an underestimate of the

mutation rate in the field.

The estimates of the mutation rate in *S. polyrhiza* and *L. minor* are among the lowest

mutation rates observed so far for multicellular eukaryotes, comparable to rates observed

in unicellular eukaryotes (Lynch *et al.*, 2016). A low mutation rate is consistent with efficient

selection against mutator alleles in highly asexual organisms such as duckweed (Kimura,

1967; Leigh, 1970). Our observation that the mutation rate in *L. minor* shows a trend

towards being marginally higher (though non-significantly) than that in *S. polyrhiza* is also

consistent with the role of linkage in favouring the evolution of lower mutation rates

(Kimura, 1967; Leigh, 1970) as genomic analyses and field observations suggest that *L. minor*

outcrosses more frequently than *S. polyrhiza* (Vasseur *et al.*, 1993; Ho, 2018; Xu *et al.*, 2019;

Ho *et al.*, 2019). This is in contrast to the theoretical prediction that strong linkage in

asexual genomes can allow mutators to fix in asexual populations if they hitchhike to

fixation with beneficial mutations they produce (André & Godelle, 2006). Population

genomic analyses in duckweed have shown that selection on protein coding genes is weak

as evidenced by elevated measures of π_N/π_S (Ho, 2018; Xu *et al.*, 2019; Ho *et al.*, 2019). This

suggests that strongly beneficial mutations might be too rare to allow mutators to be

selected for in duckweed.

Xu et al. inferred a global N_e for *S. polyrhiza* of ~1x10E-06 so it might also be possible

that the large effective population sizes of these species allow selection to be efficient

enough to lower the mutation rate more than in most multicellular eukaryotes (Sung *et al.*,

2012; Lynch *et al.*, 2016). This explanation, however, is also inconsistent with the fact that

measures for the efficacy of selection suggest that selection is weak in duckweed, likely due

to the predominance of asexual reproduction (Xu *et al.*, 2019; Ho *et al.*, 2019).

In conclusion, we report a very low SNP mutation rate in two species of duckweed

consistent with previous results in this group. We found that *de novo* mutations appear at

low frequencies within MA lines suggesting the presence of multiple segregating cell

lineages. We then used an approach that allows us to estimate the mutation rate when

multiple cell lineages are transmitted across generations. The low mutation rate of these

duckweeds is consistent with the idea that a higher degree of asexual reproduction leads to

strong selection for low mutation rates.

Acknowledgements

This work was supported by the Natural Sciences and Engineering Research Council of

Canada (AFA and SIW). We thank Mitch Cruzan for helpful suggestions with data analysis.

We thank Jade Lavallee and Victor Mollov for help with maintaining MA lines.

Author contributions

SIW and AFA conceived the project. AFA, SIW, and MB collected samples. MB performed the

MA experiment. GS did bioinformatics analyses. AFA, SIW, GS wrote the manuscript. All

authors contributed to manuscript editing. SIW and AFA contributed equally to this work.

Literature Cited:

André J-B, Godelle B. **2006**. The Evolution of Mutation Rate in Finite Asexual Populations. *Genetics* **172**: 611–626.

Appenroth K-J, Teller S, Horn M. **1996**. Photophysiology of turion formation and germination in *Spirodela polyrhiza*. *Biologia Plantarum* **38**: 95–106.

Bell G. **1982**. *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. University of California, USA: Academic Press.

Cao HX, Vu GTH, Wang W, Appenroth KJ, Messing J, Schubert I. **2016**. The map-based genome sequence of *Spirodela polyrhiza* aligned with its chromosomes, a reference for karyotype evolution. *New Phytologist* **209**: 354–363.

Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, Gabriel S, Meyerson M, Lander ES, Getz G. **2013**. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nature Biotechnology* **31**: 213–219.

Cole CT, Voskuil MI. **1996**. Population genetic structure in duckweed (*Lemna minor*, Lemnaceae). *Canadian Journal of Botany* **74**: 222–230.

Dorai-Raj S. **2014**. *binom: Binomial Confidence Intervals for Several Parameterizations. R package*. URL https://cran.r-project.org/web/packages/binom/index.html. [accessed October 5 2018]

Eyre-Walker A, Keightley PD. **2007**. The distribution of fitness effects of new mutations. *Nature Reviews Genetics* **8**: 610–618.

Foster PL. **2007**. Stress-induced mutagenesis in bacteria. *Critical Reviews in Biochemistry and Molecular Biology* **42**: 373–397.

Hicks LE. **1932**. Flower Production in the Lemnaceae. **32**: 115–132.

Hiltunen M, Grudzinska-Sterno M, Wallerman O, Ryberg M, Johannesson H. **2019**. Maintenance of High Genome Integrity over Vegetative Growth in the Fairy-Ring Mushroom *Marasmius oreades*. *Current Biology* **29**: 2758–2765.e6.

Ho KHE. **2018**. *The Effects of Asexuality and Selfing on Genetic Diversity, the Efficacy of Selection and Species Persistence.* PhD thesis, University of Toronto, Toronto, ON, Canada.

Ho E, Bartkowska M, Wright SI, Agrawal A. **2019**. Population genomics of the facultatively asexual duckweed *Spirodela polyrhiza*. *New Phytologist* **224:** 1361-1371.

Jiang C, Mithani A, Belfield EJ, Mott R, Hurst LD, Harberd NP. **2014**. Environmentally responsive genome-wide accumulation of de novo *Arabidopsis thaliana* mutations and epimutations. *Genome Research* **24**: 1821–1829.

Kimura M. **1967**. On the evolutionary adjustment of spontaneous mutation rates. *Genetics Research* **9**: 23–34.

Landolt E. 1986. *Biosystematic investigations in the family of duckweeds (Lemnaceae). Vol.2: The family of Lemnaceae: a monographic study*. Zurich, Switzerland. Geobotanischen Instutites der ETH.

Leigh EG. 1970. Natural Selection and Mutability. *The American Naturalist* **104**: 301–305.

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**: 2987–2993.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.

Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nature Reviews Genetics* **17**: 704–714.

Matsuba C, Ostrow DG, Salomon MP, Tolani A, Baer CF. 2013. Temperature, stress, and spontaneous mutation in *Caenorhabditis briggsae* and *Caenorhabditis elegans*. *Biology Letters* **9**: 20120334.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, *et al.* 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research* **20**: 1297–1303.

Ossowski S, Schneeberger K, Lucas-Lledó JI, Warthmann N, Clark RM, Shaw RG, Weigel D, Lynch M. 2010. The rate and molecular spectrum of spontaneous mutations in Arabidopsis thaliana. *Science* **327**: 92–94.

Otto SP, Orive ME. 1995. Evolutionary Consequences of Mutation and Selection within an Individual. *Genetics* **141**: 1173–1187.

Plomion C, Aury J-M, Amselem J, Leroy T, Murat F, Duplessis S, Faye S, Francillonne N, Labadie K, Provost GL, *et al.* 2018. Oak genome reveals facets of long lifespan. *Nature Plants* **4**: 440–452.

R Core Team. 2019. *R: A language and environment for statistical computing.* R Foundation for Statistical Computing, Vienna, Austria. URL https://www.r-project.org/ [Accessed September 1 2017]

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative Genomics Viewer. *Nature biotechnology* **29**: 24–26.

Saxena AS, Salomon MP, Matsuba C, Yeh S-D, Baer CF. 2019. Evolution of the Mutational Process under Relaxed Selection in *Caenorhabditis elegans*. *Molecular Biology and Evolution* **36**: 239–251.

Schmid-Siegert E, Sarkar N, Iseli C, Calderon S, Gouhier-Darimont C, Chrast J, Cattaneo P,
Schütz F, Farinelli L, Pagni M, *et al.* 2017. Low number of fixed somatic mutations in a long-
lived oak tree. *Nature Plants* **3**: 926–929.

Sharp NP, Agrawal AF. **2016**. Low Genetic Quality Alters Key Dimensions of the Mutational
Spectrum. *PLoS Biology* **14**.

Sturtevant AH. **1937**. Essays on Evolution. I. On the Effects of Selection on Mutation Rate.
The Quarterly Review of Biology **12**: 464–467.

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. **2012**. Drift-barrier hypothesis and
mutation-rate evolution. *Proceedings of the National Academy of Sciences* **109**: 18488–
18492.

Thomas GWC, Wang RJ, Puri A, Harris RA, Raveendran M, Hughes DST, Murali SC, Williams
LE, Doddapaneni H, Muzny DM, *et al.* 2018. Reproductive Longevity Predicts Mutation
Rates in Primates. *Current biology* **28**: 3193–3197.e5.

Van Hoeck A, Horemans N, Monsieurs P, Cao HX, Vandenhove H, Blust R. **2015**. The first
draft genome of the aquatic model plant Lemna minor opens the route for future stress
physiology research and biotechnological applications. *Biotechnology for Biofuels* **8**.

Vasseur L, Aarssen LW, Bennett T. **1993**. Allozymic Variation in Local Apomictic Populations
of *Lemna minor* (Lemnaceae). *American Journal of Botany* **80**: 974–979.

Wang L, Ji Y, Hu Y, Hu H, Jia X, Jiang M, Zhang X, Zhao L, Zhang Y, Jia Y, *et al.* 2019. The
architecture of intra-organism mutation rate variation in plants. *PLoS biology* **17**: e3000191.

Watson JM, Platzer A, Kazda A, Akimcheva S, Valuchova S, Nizhynska V, Nordborg M, Riha
K. **2016**. Germline replications and somatic mutation accumulation are independent of
vegetative life span in Arabidopsis. *Proceedings of the National Academy of Sciences of the*
United States of America **113**: 12226–12231.

Xu S, Stapley J, Gablenz S, Boyer J, Appenroth KJ, Sree KS, Gershenzon J, Widmer A, Huber
M. **2019**. Low genetic variation is associated with low mutation rate in the giant duckweed.
Nature Communications **10**: 1243.

Yu L, Boström C, Franzenburg S, Bayer T, Dagan T, Reusch TBH. 2020. Somatic genetic drift
and multilevel selection in a clonal seagrass. *Nature Ecology and Evolution*. *In press.*
619
620 **Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS**. **2012**. A high-performance
621 computing toolset for relatedness and principal component analysis of SNP data.
622 *Bioinformatics* **28**: 3326–3328.

Supporting Information

Fig. S1 Visualization of a low-quality genomic region that was filtered from our analyses.

Fig. S2 Allelic coverage at heterozygous sites before and after filtering.

Fig. S3 Visualization of a validated *de novo* mutation from both Illumina short read and Sanger sequencing data.

Fig. S4 Visualisation of a putative *de novo* mutation that was identified as a likely sequencing error

Fig. S5 Visualisation of a putative *de novo* mutation that was identified as a likely genome misassembly error

Table S1 List of *de novo* mutations identified in two species of duckweed

Table S2 Mutation rate estimates with power adjusted to reflect the observed proportion of reads that support *de novo* mutations

Table S3 Mutation rate estimates with power adjusted under several scenarios of allelic bias at *de novo* mutant sites in salt stressed mutation accumulation lines

Table S4 Mutation rate estimates with power adjusted under several scenarios of allelic bias at *de novo* mutant sites in control mutation accumulation lines

Table S5 Number and fraction of callable sites under several scenarios of allelic bias at *de novo* mutant sites