Estimation of the SNP mutation rate in two vegetatively

propagating species of duckweed

George Sandler[1], Magdalena Bartkowska[1], Aneil F. Agrawal[1,2]*, Stephen I. Wright[1,2]*

[1]Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks

Street, Toronto, ON M5S 3B2, Canada; [2]Center for Analysis of Genome Evolution and

Function, University of Toronto, 25 Willcocks Street, Toronto, ON M5S 3B2, Canada

Author for correspondence

George Sandler

Tel: +1 416 6172717

Email: george.sandler@mail.utoronto.ca

*These authors contributed equally to this work.

Key words: Mutation rate, vegetative reproduction, duckweed, asexual, effective

population size

Abstract

Mutation rate estimates for vegetatively reproducing organisms are rare, despite their

frequent occurrence across the tree of life. Here we report mutation rate estimates in two

vegetatively reproducing duckweed species, *Lemna minor* and *Spirodela polyrhiza*. We use a

modified approach to estimating mutation rates by taking into account the reduction in

mutation detection power that occurs when new individuals are produced from multiple cell

lineages. We estimate an extremely low per generation mutation rate in both species of

duckweed and note that allelic coverage at *de novo* mutation sites is very skewed. We also

find no substantial difference in mutation rate between mutation accumulation lines

propagated under benign conditions and those grown under salt stress. Finally, we discuss

the implications of interpreting mutation rate estimates in vegetatively propagating

organisms.

Introduction

Most research on the evolution of mutation rates has focused either on sexually

reproducing eukaryotes or unicellular organisms, both of which feature a single cell phase as

part of their life cycle. However, a diverse array of organisms reproduce either through

clonal budding, fission or vegetative growth, whereby a single cell phase is not imposed

every generation (Bell 1982). This mode of reproduction potentially allows multiple cell

lineages to be transmitted from parent to offspring, complicating the process of genotyping

individuals. This happens because when individuals composed of a mosaic of cells are

sequenced, the mean number of sequencing reads supporting non-reference mutations is

no longer 50%. Such a skew in allelic coverage makes it harder to distinguish true mutations

from sequencing errors (Cibulskis *et al.* 2013), complicating the assessment of power when

calculating per base pair mutation rates. Even if a cellular mutation rate can be calculated

for an organism with multiple cell lineages, it becomes more challenging to use this

parameter in population genetics analyses as mutations can potentially be lost within an

organism before truly contributing to population level genetic diversity. Previous theoretical

work modeling mutation load in organisms with multiple cell lineages has suggested that

cell lineage selection can significantly reduce mutation load by purging deleterious

mutations during somatic growth (Otto and Orive 1995). As most new mutations are

thought to be deleterious (Sturtevant 1937; Eyre-Walker and Keightley 2007) this type of

selection might skew the level of genetic diversity observed in organisms with vegetative

reproduction compared to the level expected given their per base pair mutation rates.

Previous studies have investigated the rate of somatic mutations in plants where

multiple cell lineages can segregate within a generation (Watson *et al.* 2016; Schmid-Siegert

et al. 2017; Plomion *et al.* 2018; Wang *et al.* 2019). While somatic mutations can be

transmitted from generation to generation in plants (Plomion *et al.* 2018; Wang *et al.* 2019),

if somatic growth is followed by sexual reproduction, a single cell bottleneck is nonetheless

imposed on any segregating variation within the soma, removing the persistence of multiple

cell lineages across generations. This is however not the case for organisms reproducing

through vegetative growth, budding or fission. Despite their frequency across the eukaryotic

tree of life, almost no per-base-pair mutation rate estimates exist for organisms procreating

through such modes of reproduction. One recent study in a vegetatively growing fairy-ring

mushroom reported very low mutation rates per mitotic cell division (Hiltunen *et al.* 2019).

The authors of this work used simulated mutations to assess the level of power they had to

detect low frequency *de novo* mutation in this dataset, improving their estimate of the fairy

ring mushroom mutation rate.

Here we report mutation rate estimates in two species of duckweed (*L. minor* and *S.*

polyrhiza). Both species are free-floating, facultatively sexual aquatic plants. While

duckweed can produce seed though sexual reproduction, most growth occurs vegetatively

via clonal budding from two pouches present in the duckweed frond (Landolt, 1986). While

these species are found all across the globe and likely have enormous census population

sizes, allozyme and genomic analyses have revealed low levels of genetic diversity within

local populations (Cole and Voskuil 1996; Ho 2018; Xu *et al.* 2019; Ho *et al.* 2019). Work by

Xu et al (2019), has estimated the per base pair mutation rate in a genotype of *S. polyrhiza*

grown in the field and the lab, finding an extremely low rate of mutation in both cases.

However, their analysis did not take into account the fact that duckweed individuals are

likely composed of a mosaic of cell lineages during periods of asexual growth, potentially

leading to an underestimate of the true mutation rate.

Studying two duckweed species allows us to contribute to three other questions in

mutation rate evolution research. First, our mutation rate estimates provide another species

to add to the existing set of species with mutation rate estimates that, collectively, allow for

testing the theory that selection against mutators should be most efficient in species with

large effective population sizes (Sung *et al.* 2012; Lynch *et al.* 2016). Duckweeds are useful

additions to this set as previous work has suggested that the effective population size (N_e) of

S. polyrhiza is on the order of a million individuals (Ho et al. 2019; Xu et al. 2019). Second,

the inclusion of two facultatively sexual species that differ in their degree of sexuality allow

us to preliminarily investigate the effect of recombination on the evolution of mutation

rates. Genomic and allozyme patterns have suggested that *L. minor* undergoes bouts of

sexual reproduction more often than *S. polyrhiza*, a pattern that is in line with flowering

observations of these species in the field (Hicks, 1932; Landolt, 1986). Theoretical work has

shown that when recombination breaks apart associations between mutator alleles (that

elevate mutation rate) and the mutations they produce, mutation rates can evolve in

several ways. On one hand selection against mutators in more sexual populations may be

relaxed as they no longer remain linked to new deleterious mutations (Kimura 1967; Leigh

1970). Alternatively mutator alleles can spread when recombination is sufficiently low if

they hitch hike along with any beneficial mutations they produce (André and Godelle 2006).

Finally, environmental stress is known to increase mutation rates in bacteria in a process

known as stress-induced mutagenesis (Foster 2007). A few examples of stress increasing

mutation rates are known in eukaryotes (Matsuba et al. 2013; Jiang et al. 2014; Sharp and

Agrawal 2016; but see Saxena et al. 2019), however it is unclear how general this

phenomenon is. We performed our experiment both under a control and salt stress

treatment to test whether stress-induced mutagenesis is a common phenomenon in plants.

We estimated the mutation rate in 46 asexually propagated mutation accumulation

lines, including two genotypes of *S. polyrhiza* and one genotype of *L. minor*. We report an

exceptionally low rate of mutation in both species of duckweed and note a pattern of

skewed allelic counts at *de novo* sites that suggests the presence of multiple segregating cell

lineages in vegetatively reproducing duckweed.

Materials and Methods

<u>Mutation accumulation and DNA extraction</u>

MA lines were started for three genotypes in April 2014 and propagated for

approximately 60 generations. Two *Spirodela polyrhiza* genotypes were used: GP23 from

Grenadier Pond, Toronto, Canada and CC from Cowan Creek, Oklahoma, USA. A single

genotype of *L. minor* (GPL7) was also isolated from Grenadier Pond, Toronto, Canada. For

each genotype (CC, GP23 and GPL7), we established 16 MA lines. We generated each line

from a single maternal plant, which was started by isolating two fronds from each genotype

culture. Because daughter fronds are generated iteratively, we grew and isolated daughters

tracking pedigree until a minimum of 16 daughter fronds paired by generation were

available for each of the three genotypes (arising from a single starting maternal frond).

Daughters in each frond pairs (matched for generation relative to maternal frond) were

assigned to one of two growth medium treatments (salt stress and control). Daughters are

produced from two pockets of meristem tissue on either side of the maternal frond and

mature daughter fronds remain attached to the maternal plant via a stipule for a short time

(Landolt, 1986). To ensure that each generation was propagated with a daughter frond, we

separated the daughter from the maternal frond as soon as the daughter began producing

her own frond. Each line was checked for mature daughter fronds every two days. The first

daughter produced was used whenever possible. MA lines were propagated in 0.5X

Appenroth liquid growth medium (Appenroth *et al.* 1996) at 24°C with 12 hours of light per

day. Generation times were similar in both species at ~2.9 and ~2.8 days under normal

conditions and 3.5 and ~3.3 days under salt stress for *S. polyrhiza* and *L. minor* respectively.

Salt stress lines were supplemented with 25mM of NaCl for *S. polyrhiza* and 50mM of NaCl

for *L. minor*. Prior to the start of the MA experiment, we performed growth assays to

establish stressful NaCl levels for both species. The chosen salinity levels caused duckweed

fronds to become patchy and thin but still allowed for continual asexual propagation.

After the termination of the MA experiment we allowed MA lines to continue

growing for several generations without removing any individuals to obtain enough plant

material to perform CTAB DNA extractions.

<u>Sequencing and filtering</u>

We sequenced the MA lines at the McGill Innovation Centre. Illumina HISeq 2000

sequencing with 100bp paired end reads was used for both *S. polyrhiza* genotypes while

Illumina HISeq 2500 sequencing with 125bp paired end reads was used for the *L. minor*

genotype.

Paired end reads were mapped to the *S. polyrhiza* and *L. minor* reference genomes

(Van Hoeck *et al.* 2015; Michael *et al.* 2017) using the Burrows-Wheeler aligner (BWA) 0.717

using the BWA-MEM option (Li and Durbin 2009). We then used Picard to remove duplicate

reads before calling indels using the HaploTypeCaller tool in GATK 3.7 (McKenna *et al.*

2010). Next, we used the IndelRealigner tool in GATK to perform indel realignment. Finally,

we used BCFtools (1.6) (Li 2011) to create mpileup files for the realigned output from GATK

and to call SNPs and short indels (indels no more than 10bp). After mapping, mean and

median coverage was 26, and 25 for individual *S. polyrhiza* lines, and 18, 17 for individual *L.*

minor lines. We also calculated total median coverage for each site within each genotype

(by summing across all individual lines), which was 436, 426, 280 for genotypes GP23, CC,

and GPL7 respectively.

We first filtered out sites with unusually high or low coverage. We did this by

eliminating sites that had coverage outside +/- 200x median coverage (summed across all

lines) in each *S. polyrhiza* genotype and +/- 100x median coverage in the *L. minor* genotype

due to the lower quality of the reference genome for this species. We visualised relatedness

between our lines using a PCA plot created from heterozygous sites present in our MA lines

in R (v5.3.5) (R Core Team 2019) using the package SNPRelate (Zheng *et al.* 2012). In doing

so we discovered two major outliers in one of our *S. polyrhiza* genotypes (CC) suggesting

that these two lines were cross contaminated. We subsequently removed these two lines

from our analysis.

Our next round of filtering aimed to remove low quality regions of the genome that

contain unusually high amounts of in-phase heterozygous variants (e.g. Figure S1). Such

regions likely represent collapsed duplications in the reference genome that map poorly to

an incorrect genomic coordinate. These variants are highly reference-biased in their allelic

coverage likely due to the poor mapping of reads that contain many differences relative to

the reference genome. To remove such regions, we first created a consensus genotype for

each set of lines; if more than one line in a given genotype supported the existence of a

heterozygote at a site, that site was designated as heterozygous in the consensus genotype.

We then performed a sliding window analysis on heterozygosity on each consensus

genotype. We used 1000bp windows with a 100bp step. After trying a variety of filtering

criteria, we decided to designate regions of the genome as callable if there existed no more

than 10 heterozygous calls in a 1000bp window in each *S. polyrhiza* genotype and no more

than 5 heterozygous calls per 1000bp window in the *L. minor* genotype. We used more

stringent criteria in *L. minor* due to the lower quality of the genome assembly. These cut-

offs represent a trade-off between eliminating problematic, variant-rich areas of the

genome and excluding well assembled genomic areas with higher than average diversity.

After filtering, around 100Mb of the genome was retained as callable in each of the three

genotypes. This filtering step greatly improved the allelic coverage of ancestral

heterozygous sites by removing suspected hidden duplications that map poorly to the

reference (Figure S2).

Next, we called putative *de novo* mutations in the remaining callable regions. Within

each set of lines, we picked sites where one line had a heterozygous genotype, with at least

5 reads supporting the non-reference base, but all other lines supported a homozygous

genotype. We then extracted such sites from the mpileup file used to call genotypes. This

was done as the pileup file contains reads that are filtered out during genotype calling but

are useful in our case as they can lead to the elimination of false positive mutations. We

filtered putative *de novo* mutations using the mpileup file in two ways. First, if a line other

than the one which contained the *de novo* mutation had any reads which supported the *de*

novo base call we excluded the site. Second, if a site with a *de novo* base call contained

reads with more than two non-reference bases across all samples, we also excluded the site.

We did this to exclude sites where a high rate of sequencing errors might have occurred. We

used this cut-off based on the observation that at sites where all lines supported a

homozygous genotype, the vast majority of sites contain no more than one alternate base

call (again likely due to sequencing errors which can be observed in the mpileup file). The

remaining putative *de novo* mutations that passed these filtering criteria were visually

inspected in the Integrative Genomics Viewer (IGV) (Robinson *et al.* 2011). We excluded a

few mutations which appeared on reads in complete linkage with other non-reference bases

(an indication of hidden genomic duplications) or on reads that looked like the product of

PCR or sequencing errors (see Figure S3-5 for examples).

<u>Power analysis</u>

To calculate the per generation mutation rate, we first needed to know how much

power we had to detect *de novo* mutations at our callable sites. To assess power, we first

obtained a list of sites where we knew we had non-zero power to detect mutations, this

included sites where all lines within a genotype supported a homozygous reference base call

and no more than one alternate base was present in the mpileup file (one less than in the

case if a *de novo* mutation was present). We then randomly sampled 500,000 such sites

from each genotype independently, and randomly chose a line where a mutation could have

happened. We randomly eliminated a third of the sites where one alternate base was

present in the mpileup file as our filtering criteria would eliminate true *de novo* mutations if

another line by chance contained a sequencing error which matched the *de novo* base call

(i.e., we assume the probability of this occurring is 1/3). Of the remaining sites, we assigned

how many reads would support the *de novo* base call by drawing from a binomial

distribution with a success rate of 50%, 34%, 28%, 20% and 10%. These different values

were chosen to represent a range of frequencies a mutation may be found due to the

inheritance of multiple cell lineages in asexual reproduction. This is similar to the approach

taken by Hiltunen et al. (2019) when calculating mutation rates in vegetatively growing

fairy-ring mushrooms.

Our estimate of power was the proportion of sites (out of the original 500,000) that

had at least 5 reads supporting the *de novo* mutation (for each possible binomial success

rate that we tested). We then multiplied our power estimates by the number of callable

sites in the genome. Then separately for each line, we multiplied the adjusted number of

callable sites by the number of MA generations and summed these values across all lines in

a given genotype (split by treatment). This provided us with a denominator for our mutation

rate calculation. Our final step was to divide the number of *de novo* mutations identified in

each genotype (split by treatment) by this denominator. The *de novo* mutation count was

adjusted for false positives identified during mutation validation.

<u>Calling indels</u>

We scanned our lines for *de novo* indels in the same way as we searched for point

mutations with one key modification. We used the same regions of the genome that we had

previously assessed as "callable"; however, we only considered *de novo* indels if they were

at least 2000 bp away from any other indel in any other line of the same genotype. This

filtering step was required to avoid false positive indel calls which appear due to spurious

mapping patterns in repetitive regions.

<u>Mutation validation</u>

We only had tissue from *S. polyrhiza* to perform validation on putative *de novo*

mutations as unlike *L. minor*, *S. polyrhiza* produces asexual resting stages called turions

which we were able utilize for long term refrigerated storage. After allowing turions to

germinate we extracted DNA from our MA lines using a Qiagen DNeasy kit. Afterwards, we

designed primers for 14 SNPs and one indel found in our two *S. polyrhiza* genotypes. We

performed PCR reactions using FroggaBio PCR mastermix which were then sent off for

Sanger sequencing at Eurofins Genomics. We inspected the Sanger sequencing

chromatograms in FinchTV 1.4.0 (Geospiza, Inc.; Seattle, WA, USA;

http://www.geospiza.com) to see if we could detect a peak suggesting the presence of a

mutant base in the focal MA line and checked that the base was not present in a different

MA line of the same genotype as a negative control. One line from the *S. polyrhiza* CC

genotype (line E) contained a massive overabundance of putative *de novo* mutations (of the

29 initial mutations found in 14 lines of the CC genotype, 9 were identified from this line); all

3 variants that we attempted to Sanger sequence validate from this line failed. Ultimately,

we excluded this CC line from all the results presented here. Excluding CC line E, 11 total

SNPs that were checked with Sanger sequencing and used to correct for the false positive

rate in mutation detection in all our *S. polyrhiza* lines. We adjusted our mutation rate for

false positive by excluding mutations that failed validation, summing all mutations that

passed validation, and multiplying the sum of unvalidated mutations by our estimate of the

false positive rate. Our formula for the true positive rate is as follows:

$$TP = \frac{n_{validated}}{n_{validated} + n_{failed}}$$

262   Where $n_{validated}$ refers to the total number of mutations successfully validated in *S.*

263   *polyrhiza*, and n_{failed} refers to the total number of mutations that failed validation.

265   Finally, the formula for our per generation, per base pair mutation rate estimates is

266   as follows:

$$\frac{n_{validated\ in\ lines} + TP \times n_{not-checked\ in\ lines}}{\sum(\#callable\ sites \times \#generations) \times power \times 2}$$

where $n_{validated\ in\ lines}$ refers to the number of validated mutations in the focal set of lines,

and $n_{not-checked\ in\ lines}$ refers to the number of mutations not tested with Sanger

sequencing in the focal set of lines. The number of mutations and callable sites was summed

across all lines within a genotype, within a treatment. The true positive rate, TP, was

estimated once using all Sanger-tested mutations from across the entire experiment.

<u>Statistical analysis</u>

We calculated 95% confidence intervals for our mutation rates using the Agresti

Coull method implemented in the R package "binom" (Dorai-Raj 2014). We tested for

significant differences between our salt and normal treatment lines using independent Chi-

square tests in each genotype. To test for differences between *L. minor* and *S. polyrhiza* we

merged the number of callable sites (scaled by power) and number of mutations in the two

S. polyrhiza genotypes (separately for each treatment), and also applied Chi-square tests to

test for significance in mutation rate variation between species and conditions.

Data availability
Raw sequencing data is available on the NCBI SRA database under accession PRJNA659313
for *S. polyrhiza* and PRJNA659264 for *L. minor*. Custom scripts and additional data are
available at https://github.com/gsan211/duckweed_MA. Supplementary material has been
uploaded as a separate file.

Results

De novo mutations

291   In total, we identified 23 and 19 putative *de novo* mutations in *S. polyrhiza*

genotypes GP23 and CC, respectively and a further 29 mutations in *L. minor* (Supplemental

table 1). When inspecting the putative *de novo* mutations in IGV, we observed that most

mutant sites exhibited highly reference biased allelic counts (over 50% of reads support the

reference base call). On average, mutant sites contained 26%, 30%, and 34% mutant reads

with SD 8.2%, 11.6%, and 9.1% in genotypes GP23, CC (*S. polyrhiza*) and GPL7 (*L. minor*)

respectively. We implemented several filtering and quality control steps to ensure these

mapping patterns were not a result of sequencing error or genome mis-assembly. First, we

masked areas of the genome that had odd coverage patterns or were enriched for

heterozygous calls to avoid areas containing hidden genomic duplications. The heterozygous

variants that pass these filtering criteria and are present in the ancestral genotypes (i.e. are

shared by all MA lines within a genotype) show relatively normal patterns of reference and

non-reference allele coverage. Second, we only considered *de novo* mutations at sites with

minimal sequencing errors. This step eliminated problematic areas of the genome prone to

genotyping error. Finally, we performed Sanger sequencing validation on 11 putative *de*

novo mutations (in *S. polyrhiza*) of which 6 were validated with both positive and negative

controls. The pattern of reference bias in our *de novo* mutations was also present in our

Sanger sequencing chromatograms; the reference base peak was generally much larger than

the mutant peak. When we plotted Illumina mutant base frequency vs. Sanger mutant peak

height (standardized by reference peak height), we observed strong concordance for our 6

validated mutations (R^2 = 0.56; Figure 1) suggesting that these are not spurious mapping

patterns or sequencing errors but rather reflect a true bias in mutation abundance. When

we inspected our PCR products with gel electrophoresis we observed single, clean bands of

the expected product size so we consider it unlikely that multiple sites in the genome were

amplified leading to odd mapping patterns at our *de novo* mutation sites. One alternative is

that reference biased mapping patterns may be due to some bias in the PCR amplification

process. We do not believe this is likely however since our primers were designed for

sequences that were flanking the site of *de novo* mutations that should be identical whether

a *de novo* mutations is present or not.

We report a high false positive rate for *de novo* mutation identification in *S. polyrhiza*

of ~45%. This is likely a consequence of having to distinguish between true mutations with

low allelic counts and sequencing errors or bioinformatic artifacts that can appear at

similarly low frequencies. Additionally, background noise in Sanger Sequencing

chromatograms can obscure variants with low allelic counts, posing a potential way in which

false positive rates may be artificially elevated. We attempted to avoid this issue by ensuring

that our chromatograms had relatively low levels of background noise before confidently

assigning mutations as failing or passing validation.

Figure *1 Proportion of de novo bases from Illumina reads vs. relative peak height of mutant*

base in Sanger sequencing data in Spirodela polyrhiza. R^2 = 0.57.

The majority of mutations were C -> T transitions in both species of duckweeds

(Figure 2) in concordance with patterns found in previous mutation rate studies (Ossowski

335 *et al.* 2010; Thomas *et al.* 2018). Transitions were more common than transversions, with

336 the average ti/tv ratio being 2.23 for *S. polyrhiza* and 4.5 for *L. minor*. This is consistent with

337 previous results reported in *S. polyrhiza* by Xu et al. (2019) where three C -> T and one C -> A

338 mutations were detected and validated. Approximately 50% of C -> T transitions occurred at

339 CpG sites in both species, a pattern that is consistent with previous evidence of elevated

340 mutation rates at such sites (Hodgkinson and Eyre-Walker 2011). We used SNPeff (Cingolani

341 *et al.* 2012) to annotate our putative de novo mutations with most SNPs being labelled as

342 intergenic (See supplemental table 1).

343  Our analysis uncovered only one putatively *de novo* short indel that turned out to be

344 a false positive based on our Sanger sequencing validation analysis.

347 **Figure 2** *De novo mutation spectra in two species of duckweed (Spirodela polyrhiza, Lemna*

348 *minor).*

351 <u>Mutation rate comparisons between *S. polyrhiza* and *L. minor*</u>

352  Our estimate of the per generation, per base pair SNP mutation rate highly depends

353 on our power to detect mutations, which in turn depends on factors such as total read

354 depth and the number of reads that support a *de novo* mutation base-call (See

355 Supplemental tables 2-5). Heterozygous SNPs are expected to be supported by around 50%

356 of reads in organisms with a single cell phase. In our case, on average, 28% (*S. polyrhiza*)

357 34% (*L. minor*) of reads supported *de novo* heterozygous mutations. To improve our

358 estimate of the mutation rate, we assumed that we should only expect to find mutations

359 that are on average supported by 28% (in *S. polyrhiza)* or 34% (in *L.* minor) of reads at each

360 mutant site (see methods for more details of power analysis). In this case, our point

estimate of the mutation rate for plants grown in standard medium is 8.39E-11 for *S.*

polyrhiza and 8.66E-11 for *L. minor* (Figure 3). For plants grown in stressful salt medium our

point estimates for the mutation rate are 9.91E-11 for *S. polyrhiza* and 16.9E-11 for *L. minor*.

While the estimated mutation rate is higher for *L. minor* in both conditions, the

difference between the two species is non-significant (chi-square test: p = 1 and p = 0.30 for

control and salt stressed conditions). The difference in mutation rate in the two species is,

however, very sensitive to the expected frequency of reads that support the mutant base.

For example, if our power estimate is based on the assumption that *de novo* mutations are

expected to constitute only 10% of reads, the mutation rate under normal conditions

increases to 5.13E-10 in *S. polyrhiza* and 16.8E-10 in *L. minor*. Our mutation rate estimates

were not significantly different between our two individual *S. polyrhiza* genotypes (chi-

square test: p = 0.50 and p = 0.77 for control and salt stressed conditions). Supplemental

tables 3-4 give the inferred mutation rates with alternative assumptions for the expected

proportion of non-reference reads at sites harbouring *de novo mutations*.

Figure 3 *Mutation rate estimates in two species of duckweeds duckweed (Spirodela*

polyrhiza, Lemna minor). Error bars show Agresti-Coull 95% confidence intervals.

Effect of salt stress on mutation rate

There were no significant differences in the mutation rate between lines propagated

in normal and salt stress medium for either of our three genotypes (Figure 4, chi-square

test: p = 1.00, p = 0.53, p = 0.29 for genotypes GP23, CC, GPL7 respectively). There was also

no consistent trend in the change of mutation rate with the addition of salt stress as

genotype GP23 appeared to have a decreased mutation rate while genotypes CC and GPL7

appeared to have an increase in the mutation rate with the addition of salt stress.

Salt stress also did not have a significant impact on the ti/tv ratio. In two of the

genotypes (1.33 -> 2 *S. polyrhiza* CC and 4.5 -> 8 *L. minor*), salt stress increased the ti/tv

ration. In *S. polyrhiza* genotypes GP23 salt stress decreased the ti/tv ration (3.67 -> 2.00)

although the difference was not significant in either of the three genotypes (chi-square test:

p > 0.5 all cases). These results run counter to previous work in *A. thaliana* where salt stress

was found to increase the de-novo mutation rate ~two-fold and lower the ti/tv ratio (Jiang

et al. 2014).

Figure 4 *Effect of salt stress on the mutation rate in two genotypes of Spirodela polyrhiza*

and one genotype of Lemna minor. Error bars show Agresti Coull 95% confidence intervals.

Discussion

Using whole genome sequencing on 46 MA lines, we report a low per base pair, per generation SNP mutation rate in two species of duckweeds under two growth conditions. An important result in our study is that *de novo* mutations appear to have considerably reference biased genomic coverage in both duckweed species. We believe that this pattern is not indicative of sequencing or genome assembly errors but rather is a by-product of vegetative reproduction for several reasons. First, upon our inspection of the data in an independent study of mutation rates in *S. polyrhiza* by Xu et al.; we noticed similar levels of reference bias in Illumina short-read sequencing data at validated *de novo* mutations. Second, our own validation with Sanger sequencing showed that mutations with a higher reference bias in the Illumina dataset tended to have higher reference base peaks in their Sanger sequencing chromatograms and vice versa. Moreover, after filtering, most ancestral heterozygous sites that were shared by all lines in the three clonal genotypes were not reference biased in such a manner and those that were, were found in regions highly enriched for non-reference base calls. None of our putative *de novo* mutations are found in such regions; due to the low genetic diversity in duckweed, *de novo* mutations were generally the only heterozygous variants present in the immediate genomic area. This means that mapping bias due to divergence from the reference is unlikely to cause the strong allelic coverage bias we observed. Finally, a study of mutations in a vegetatively growing fairy-ring mushroom also reported patterns of reference bias with an average allelic coverage of 44% across 111 *de novo* mutations (Hiltunen *et al.* 2019). Therefore, the reference bias in our mutations is most likely a signature of the segregation of multiple cell lines from generation to generation in our duckweed MA experiment. Given that vegetative reproduction through clonal budding is the main form of reproduction in duckweed (Landolt, 1986), it seems reasonable that duckweed might not undergo a single cell phase for prolonged periods of time.

Simple models can be helpful in clarifying how multi-cell descent during vegetative reproduction may affect mutation rates and our estimation of them. Assume that clonal reproduction occurs by n parental cells forming a diploid offspring (generation 1). If a mutation occurs in one of these n cells, because of multi-cell descent, the offspring will be a genetic mosaic (i.e., not all cells will be genetically identical). The mutation begins at a frequency of $1/2n$ in the offspring. With conventional single-cell descent, where $n = 1$, a new

431   mutation is expected to be at 50% frequency, but the frequency will be lower than 50%

432   whenever $n > 1$. The n cells multiply in some (unknown) growth pattern to produce the

433   mature offspring, which then itself reproduces. In the absence of detailed knowledge of

434   developmental growth trajectories, we do not know what the representation of the original

435   n cell lineages will be in the subsequent generation.

436   We can consider two simple scenarios: If only one of these original cell lineages gives

rise to the next set of n cells used to produce the next generation (generation 2), then the

mutation will either be completely lost from the lineage (if it was not in the chosen cell

lineage) or it will be present in heterozygous state in all cells of future generations (if it was

present in the chosen cell lineage). In this first scenario, the genetic mosaicism resulting

from multi-cell descent persists only a single generation. Thus, in an experiment such as

ours, only mutations occurring in the very last generation will be affected by this issue (i.e.,

multi-cell descent is a trivial issue in this case). A second scenario is that all cell lineages

grow at equal rates and, each generation, n cells are chosen at random to form the next

daughter. This is conceptually similar to the process of coalescence in a population (of cells

here rather than individuals) of size n. Thinking backwards in time, all cells must eventually

trace their coalescent history to a single cellular ancestor, but it may take many generations

for this to happen (i.e., $2n$ generations, on average, but with variance proportional to n^2).

Thinking forwards in time, a mutation occurring in one of the n cells forming the generation

1 offspring may be present for many future generations, possibly becoming quite common

within individuals, before eventually being present in all future cells (with chance $1/n$) or

none of them (with chance $(n - 1)/n$). With even modest values of n (e.g., $8 < n < 80$), genetic

mosaicism can persist for many generations. Mutations that will eventually be present in all

lineages—as well as those that will eventually be eliminated—will thus be found below the

50% frequency expected in a "traditional" (i.e., non-mosaic) heterozygote. Although this

second scenario as formulated here is unrealistically simple (e.g., random and independent

choice of n cells for each generation), it illustrates how multi-cell descent can have

consequences across multiple generations. Though developmental growth trajectories in

duckweed are insufficient to formulate a more realistic model, we suspect that multiple cell

lineages persist across multiple generations in duckweed and is responsible for the clear

bias towards mutant SNPs being less than 50%. Recent work on segregating mutations in a

single *Zostera marina* seagrass clone has made similar arguments to this model (Yu et al.,

2020). In their study, Yu et al. uncovered a large class of reference biased SNPs that were

present in some but not all sampled *Z. marina* ramets. By reconstructing the genealogical

relationship of the sampled ramets, the authors demonstrated that such SNPs changed in

frequency during vegetative growth, with some reaching heterozygous fixation in specific

ramets. The authors argued that this data was well explained by a model of "somatic drift"

whereby *de novo* mutations arise at low frequency within a single cell lineage before

ultimately either reaching fixation or being lost.

Calculating mutation rates in organisms with multiple segregating cell lineages poses

a technical challenge due to the difficulty of assessing power when mutations are present in

only a subset of the cells of an individual. We implemented a method that takes into

account the fraction of reads we expect to support a *de novo* mutation by using observed

mapping patterns of putatively *de novo* mutations, similar to the approach used by Hiltunen

et al. (2019). This method considers the fact that we have reduced power to detect recently

arisen mutations that are at low frequency within their MA lines, giving a more accurate

estimate of the SNP mutation rate. This approach is an admittedly crude attempt to address

the problem. Rather than assuming the expected frequency of the mutant allele is 50%, we

simply choose a lower value based on the observed coverage at putative *de novo* mutations.

In reality, this lower value is unknown and will differ among mutations depending upon

when each mutation arose, i.e., there is an unknown distribution of mutation frequencies at

the end of the MA experiment. Nonetheless, our approach is an improvement over

completely ignoring the issue. Moreover, the variation in mutation rate estimates inferred

using different values for the assumed expected mutant frequency provides some sense of

the sensitivity of these estimates to this assumption. However, the issue of the unknown

distribution of mutation frequencies adds a caveat to the between-species comparison.

Because of the moderate difference in coverage between species, which affects the power

to detect mutations segregating at different true frequencies, the estimates for the two

species may be somewhat differentially affected by any bias introduced by our method.

A natural consequence of the lack of a single cell phase is that from a population

genetics perspective, the mutation rate becomes a harder parameter to interpret. On one

hand, we may be interested in calculating the mutation rate that captures every new

mutation that has arisen in a clonal bud. Alternatively, from an evolutionary perspective, we

might be only be interested in mutations that will not only arise in a clonal bud but also

persist in a future clonal descendant such that they contribute to population level genetic

diversity. Aside from random inter-cell lineage "drift", cell lineage selection may bias which

de novo mutations are eventually lost in a clonal lineage (Otto and Orive 1995), although

this process likely has a minimal effect on mutation rate estimates in most mutation

accumulation studies. In principle, we might be able to calculate an "evolutionarily

relevant" mutation rate by performing long term mutation accumulation experiments

allowing the majority of *de novo* mutations to either be lost or to have fixed within a clonal

lineage such that all cells in any clone will be either fully homozygous (in the case of loss) or

heterozygous (in the case of fixation). However, from a practical standpoint, it is hard to

know *a priori* how many generations will be necessary for this process to occur. In practice,

we could attempt to estimate the frequency of *de novo* mutations that have become

sufficiently common such that they are likely to not be lost before fixation leading to an

estimate of an evolutionarily relevant mutation rate. For example, we could assume that

mutations found in at least 50% of cells (i.e. at a frequency of at least 25% in a clonal bud)

are more likely than not to eventually fix in their clonal lineage, being found in 100% of cells

in some future generation. In reality, some of these mutations are still likely to be lost

before they have a chance to fix while simultaneously some mutations that are below 50%

frequency might still reach fixation in the future. More generally, the lower the frequency

cut-off we use for mutations that are "likely" to fix, the better the chance that we capture

all of the mutations that will reach fixation with the caveat that we will also be capturing

more mutations that will eventually be lost. In our study we opted to use a read number

cut-off rather than a frequency cut-off as we were primarily concerned about differentiating

true *de novo* mutations from false positives, a task that is particularly challenging in

organisms with low mutation rates. In practice, our filtering criteria results in us mostly

identifying mutations with an allelic frequency of at least 20% (61/71 *de novo* mutation

reported in our study). As mentioned above however, some of these mutations may still be

lost prior to fixation within an organism meaning that the mutation rate we report here is

likely inflated compared to a true "evolutionarily relevant" rate. To some extent, this

upward bias is counter-balanced by mutations that could eventually go to fixation but are at

too low frequency at the time of sequencing to be either observed or even inferred by our

power calculation that is based on some threshold frequency (10-50%; Tables S2-5).

Our estimate of the mutation rate in *S. polyrhiza* (8.39E-11 per bp per gen.) is similar

to the estimates reported by Xu et al. (7.92E-11 per bp per gen.), however there are two

important differences between our studies. First, our MA experiment was conducted only in

the lab, while Xu et al. placed MA lines both in the lab and in the field, observing no

mutations in the lab setting, likely due to the smaller number of MA generations in their

study. Second, Xu et al. used ancestral heterozygous sites to estimate their power to detect

de novo mutations which are not strongly reference biased in a similar manner. This

suggests that the estimate from Xu et al., while conducted in a more naturally realistic

environment, may be an underestimate of the mutation rate in the field.

535
536

Figure 5 *Comparison of per base pair, per generation mutation rates between duckweed and*

other eukaryotic species. Duckweed estimated highlighted by red box. References: (Ossowski

et al. 2010; Lynch 2010; Denver et al. 2012; Schrider et al. 2013; Weller et al. 2014; Zhu et

al. 2014; Venn et al. 2014; Keightley et al. 2015; Uchimura et al. 2015; Yang et al. 2015;

Farlow et al. 2015; Ness et al. 2015; Smeds et al. 2016; Xie et al. 2016; Besenbacher et al.

2016 p.; Feng et al. 2017; Liu et al. 2017; Oppold and Pfenninger 2017; Flynn et al. 2017;

Krasovec et al. 2017, 2018, 2019; Hanlon et al. 2019; Orr et al. 2020)

The estimates of the per generation, per base pair mutation rate in *S. polyrhiza* and

L. minor are among the lowest so far for multicellular eukaryotes (Figure 5 and Table S6).

One potential explanation for why duckweeds have lower mutation rates than other plants

is the smaller number of cell divisions they likely go through compared to their larger, long-

lived relatives. Mutation rate studies in trees have indeed shown that while per generation

tree mutation rates are high (on the order of 1E-08 mutations per bp), mutation rates per

unit growth (a proxy for number of cell divisions), must be several orders of magnitude

lower (Xie *et al.* 2016; Hanlon *et al.* 2019; Orr *et al.* 2020). However, duckweeds do appear

to exhibit a low mutation rate compared to animals which have limited cell divisions

between meiotic events due to a segregated germline. Moreover, our per generation

duckweed mutation rate estimates fall on the lower end of values seen in green algae and

other unicellular eukaryotes, organisms which unlike duckweed only go through a single cell

division per generation (Figure 5). Overall, these patterns fit with previous work that has

suggested that number of cell divisions per generation alone is not enough to fully explain

variation in mutation rates (Lynch 2010).

The low mutation rate observed in our study is consistent with efficient selection

against mutator alleles in highly asexual organisms such as duckweed, bacteria and

unicellular eukaryotes (Kimura 1967; Leigh 1970). This explanation would also predict that

the mutation rate should be higher in *L. minor* than in *S. polyrhiza* as genomic analyses and

field observations suggest that *L. minor* outcrosses more frequently than *S. polyrhiza*

(Vasseur *et al.* 1993; Ho 2018; Xu *et al.* 2019; Ho *et al.* 2019). However, we did not observe a

significant difference in mutation rate between species in our study. This could either be

because we did not have the power to differentiate between such overall low mutation

rates, because the difference in rates of sexual reproduction is not large enough between

duckweed species, because a difference in rates of sex has arisen in recent history, or

because factors other than reproductive mode play a larger role in shaping mutation rate

evolution. The overall low mutation rate in both species is in contrast to the theoretical

prediction that strong linkage in asexual genomes can allow mutators to fix in asexual

populations if they hitchhike to fixation with beneficial mutations they produce (André and

Godelle 2006). Population genomic analyses in duckweed have shown that selection on

protein coding genes is weak as evidenced by elevated measures of π_N/π_S (Ho 2018; Xu *et al.*

2019; Ho *et al.* 2019) This suggests that strongly beneficial mutations might be too rare to

allow mutators to be selected for in duckweed.

Xu et al. inferred a global N_e for *S. polyrhiza* of ~1x10E-06 so it might also be possible

that the large effective population sizes of these species allow selection to be efficient

enough to lower the mutation rate more than in most multicellular eukaryotes (Sung *et al.*

2012; Lynch *et al.* 2016). This explanation, however, is also inconsistent with the fact that

measures for the efficacy of selection suggest that selection is weak in duckweed, likely due

to the predominance of asexual reproduction (Xu *et al.* 2019; Ho *et al.* 2019).

In conclusion, we report a very low SNP mutation rate in two species of duckweed

consistent with previous results in this group. We found that *de novo* mutations appear at

low frequencies within MA lines suggesting the presence of multiple segregating cell

lineages. We then used an approach that allows us to estimate the mutation rate when

multiple cell lineages are transmitted across generations. The low mutation rate of these

duckweeds is consistent with the idea that a higher degree of asexual reproduction leads to

strong selection for low mutation rates.

Acknowledgements

This work was supported by the Natural Sciences and Engineering Research Council of

Canada (AFA and SIW). We thank Mitch Cruzan for helpful suggestions with data analysis.

We thank Jade Lavallee and Victor Mollov for help with maintaining MA lines.

Author contributions

SIW and AFA conceived the project. AFA, SIW, and MB collected samples. MB performed the

MA experiment. GS did bioinformatics analyses. AFA, SIW, GS wrote the manuscript. All

authors contributed to manuscript editing. SIW and AFA contributed equally to this work.

Literature Cited:

André, J.-B., and B. Godelle, 2006 The evolution of mutation rate in finite asexual populations. Genetics 172: 611–626.

Appenroth, K.-J., S. Teller, and M. Horn, 1996 Photophysiology of turion formation and germination in Spirodela polyrhiza. Biol. Plant. 38: 95–106.

Bell, G., 1982 *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. University of California Press, Berkeley

Besenbacher, S., P. Sulem, A. Helgason, H. Helgason, H. Kristjansson *et al.*, 2016 Multi-nucleotide de novo mutations in humans. PLOS Genet. 12: e1006315.

Cibulskis, K., M. S. Lawrence, S. L. Carter, A. Sivachenko, D. Jaffe *et al.*, 2013 Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. Nat. Biotechnol. 31: 213–219.

Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen *et al.*, 2012 A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly (Austin) 6: 80–92.

Cole, C. T., and M. I. Voskuil, 1996 Population genetic structure in duckweed (Lemna minor, Lemnaceae). Can. J. Bot. 74: 222–230.

Denver, D. R., L. J. Wilhelm, D. K. Howe, K. Gafner, P. C. Dolan *et al.*, 2012 Variation in base-substitution mutation in experimental and natural lineages of Caenorhabditis nematodes. Genome Biol. Evol. 4: 513–522.

Dorai-Raj, S., 2014 binom: *Binomial confidence intervals for several parameterizations*. R package, version 1.1-1, URL https://cran.r-project.org/web/packages/binom/index.html

Eyre-Walker, A., and P. D. Keightley, 2007 The distribution of fitness effects of new mutations. Nat. Rev. Genet. 8: 610–618.

Farlow, A., H. Long, S. Arnoux, W. Sung, T. G. Doak *et al.*, 2015 The spontaneous mutation rate in the fission yeast Schizosaccharomyces pombe. Genetics 201: 737–744.

Feng, C., M. Pettersson, S. Lamichhaney, C.-J. Rubin, N. Rafati *et al.*, 2017 Moderate nucleotide diversity in the Atlantic herring is associated with a low mutation rate. eLife 6:.

Flynn, J. M., F. J. J. Chain, D. J. Schoen, and M. E. Cristescu, 2017 Spontaneous mutation accumulation in Daphnia pulex in selection-free vs. competitive environments. Mol. Biol. Evol. 34: 160–173.

Foster, P. L., 2007 Stress-induced mutagenesis in bacteria. Crit. Rev. Biochem. Mol. Biol. 42: 373–397.

Hanlon, V. C. T., S. P. Otto, and S. N. Aitken, 2019 Somatic mutations substantially increase the per-generation mutation rate in the conifer Picea sitchensis. Evol. Lett. 3: 348–358.

Hicks, L. E., 1932 Flower Production in the Lemnaceae. Ohio Journal of Science 32: 115–132.

Hiltunen, M., M. Grudzinska-Sterno, O. Wallerman, M. Ryberg, and H. Johannesson, 2019 Maintenance of high genome integrity over vegetative growth in the fairy-ring mushroom Marasmius oreades. Curr. Biol. 29: 2758-2765.e6.

Ho, K. H. E., 2018 The effects of asexuality and selfing on genetic diversity, the Efficacy of selection and species Persistence. [Thesis, University of Toronto].

Ho, E., M. Bartkowska, S. I. Wright, and A. Agrawal, 2019 Population genomics of the facultatively asexual duckweed Spirodela polyrhiza. New Phytol. 224: 1361–1371

Hodgkinson, A., and A. Eyre-Walker, 2011 Variation in the mutation rate across mammalian
genomes. Nat. Rev. Genet. 12: 756–766.
Jiang, C., A. Mithani, E. J. Belfield, R. Mott, L. D. Hurst *et al.*, 2014 Environmentally
responsive genome-wide accumulation of de novo Arabidopsis thaliana mutations
and epimutations. Genome Res. 24: 1821–1829.
Keightley, P. D., A. Pinharanda, R. W. Ness, F. Simpson, K. K. Dasmahapatra *et al.*, 2015
Estimation of the spontaneous mutation rate in Heliconius melpomene. Mol. Biol.
Evol. 32: 239–243.
Kimura, M., 1967 On the evolutionary adjustment of spontaneous mutation rates*. Genet.
Res. 9: 23–34.
Krasovec, M., M. Chester, K. Ridout, and D. A. Filatov, 2018 The mutation rate and the age of
the sex chromosomes in Silene latifolia. Curr. Biol. 28: 1832-1838.e4.
Krasovec, M., A. Eyre-Walker, S. Sanchez-Ferandin, and G. Piganeau, 2017 Spontaneous
mutation rate in the smallest photosynthetic eukaryotes. Mol. Biol. Evol. 34: 1770–
1779.
Krasovec, M., S. Sanchez-Brosseau, and G. Piganeau, 2019 First estimation of the
spontaneous mutation rate in diatoms. Genome Biol. Evol. 11: 1829–1837.
Landolt, E., 1986 *Biosystematic investigations in the family of duckweeds (Lemnaceae),*
volume 2. The family of Lemnaceae – a monographic study, volume 1.
Geobotanischen Institutes der ETH, Zurich.
Leigh, Egbert Giles, 1970 Natural selection and mutability. Am. Nat. 104: 301–305.
Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data.
Bioinformatics. 27: 2987–2993.
Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 25: 1754–1760.
Liu, H., Y. Jia, X. Sun, D. Tian, L. D. Hurst *et al.*, 2017 Direct determination of the mutation
rate in the bumblebee reveals evidence for weak recombination-associated
mutation and an approximate rate constancy in insects. Mol. Biol. Evol. 34: 119–130.
Lynch, M., 2010 Evolution of the mutation rate. Trends Genet. 26: 345–352.
Lynch, M., M. S. Ackerman, J.-F. Gout, H. Long, W. Sung *et al.*, 2016 Genetic drift, selection
and the evolution of the mutation rate. Nat. Rev. Genet. 17: 704–714.
Matsuba, C., D. G. Ostrow, M. P. Salomon, A. Tolani, and C. F. Baer, 2013 Temperature,
stress and spontaneous mutation in Caenorhabditis briggsae and Caenorhabditis
elegans. Biol. Lett. 9: 20120334.
McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 20: 1297–1303.
Michael, T. P., D. Bryant, R. Gutierrez, N. Borisjuk, P. Chu *et al.*, 2017 Comprehensive
definition of genome features in Spirodela polyrhiza by high-depth physical mapping
and short-read DNA sequencing strategies. Plant J. 89: 617–635.
Ness, R. W., A. D. Morgan, R. B. Vasanthakrishnan, N. Colegrave, and P. D. Keightley, 2015
Extensive de novo mutation rate variation between individuals and across the
genome of Chlamydomonas reinhardtii. Genome Res. 25: 1739–1749.
Oppold, A.-M., and M. Pfenninger, 2017 Direct estimation of the spontaneous mutation rate
by short-term mutation accumulation lines in Chironomus riparius. Evol. Lett. 1: 86–
92.

Orr, A. J., A. Padovan, D. Kainer, C. Külheim, L. Bromham *et al.*, 2020 A phylogenomic
approach reveals a low somatic mutation rate in a long-lived plant. Proc. R. Soc. B
Biol. Sci. 287: 20192364.
Ossowski, S., K. Schneeberger, J. I. Lucas-Lledó, N. Warthmann, R. M. Clark *et al.*, 2010 The
rate and molecular spectrum of spontaneous mutations in Arabidopsis thaliana.
Science 327: 92–94.
Otto, S. P., and M. E. Orive, 1995 Evolutionary consequences of mutation and selection
within an Individual. Genetics 141: 1173–1187.
Plomion, C., J.-M. Aury, J. Amselem, T. Leroy, F. Murat *et al.*, 2018 Oak genome reveals
facets of long lifespan. Nat. Plants 4: 440.
R Core Team, 2019 R: *A language and environment for statistical computing. R Foundation
for Statistical Computing*, Vienna, Austria. URL https://www.R-project.org/.
Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander *et al.*, 2011
Integrative Genomics Viewer. Nat. Biotechnol. 29: 24–26.
Saxena, A. S., M. P. Salomon, C. Matsuba, S.-D. Yeh, and C. F. Baer, 2019 Evolution of the
mutational process under relaxed selection in Caenorhabditis elegans. Mol. Biol.
Evol. 36: 239–251.
Schmid-Siegert, E., N. Sarkar, C. Iseli, S. Calderon, C. Gouhier-Darimont *et al.*, 2017 Low
number of fixed somatic mutations in a long-lived oak tree. Nat. Plants 3: 926–929.
Schrider, D. R., D. Houle, M. Lynch, and M. W. Hahn, 2013 Rates and genomic consequences
of spontaneous mutational events in Drosophila melanogaster. Genetics 194: 937–
954.
Sharp, N. P., and A. F. Agrawal, 2016 Low genetic quality alters key dimensions of the
mutational spectrum. PLoS Biol. 14:.
Smeds, L., A. Qvarnström, and H. Ellegren, 2016 Direct estimate of the rate of germline
mutation in a bird. Genome Res. 26: 1211–1218.
Sturtevant, A. H., 1937 Essays on Evolution. I. On the Effects of Selection on Mutation Rate.
Q. Rev. Biol. 12: 464–467.
Sung, W., M. S. Ackerman, S. F. Miller, T. G. Doak, and M. Lynch, 2012 Drift-barrier
hypothesis and mutation-rate evolution. Proc. Natl. Acad. Sci. 109: 18488–18492.
Thomas, G. W. C., R. J. Wang, A. Puri, R. A. Harris, M. Raveendran *et al.*, 2018 Reproductive
longevity predicts mutation rates in Primates. Curr. Biol. 28: 3193-3197.e5.
Uchimura, A., M. Higuchi, Y. Minakuchi, M. Ohno, A. Toyoda *et al.*, 2015 Germline mutation
rates and the long-term phenotypic effects of mutation accumulation in wild-type
laboratory mice and mutator mice. Genome Res. 25:1125-1134
Van Hoeck, A., N. Horemans, P. Monsieurs, H. X. Cao, H. Vandenhove *et al.*, 2015 The first
draft genome of the aquatic model plant Lemna minor opens the route for future
stress physiology research and biotechnological applications. Biotechnol. Biofuels 8:.
Vasseur, L., L. W. Aarssen, and T. Bennett, 1993 Allozymic variation in local apomictic
populations of Lemna minor (Lemnaceae). Am. J. Bot. 80: 974–979.
Venn, O., I. Turner, I. Mathieson, N. de Groot, R. Bontrop *et al.*, 2014 Strong male bias drives
germline mutation in chimpanzees. Science 344: 1272–1275.
Wang, L., Y. Ji, Y. Hu, H. Hu, X. Jia *et al.*, 2019 The architecture of intra-organism mutation
rate variation in plants. PLoS Biol. 17: e3000191.
Watson, J. M., A. Platzer, A. Kazda, S. Akimcheva, S. Valuchova *et al.*, 2016 Germline
replications and somatic mutation accumulation are independent of vegetative life
span in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. 113: 12226–12231.

Weller, A. M., C. Rödelsperger, G. Eberhardt, R. I. Molnar, and R. J. Sommer, 2014 Opposing
forces of A/T-biased mutations and G/C-biased gene conversions shape the genome
of the nematode Pristionchus pacificus. Genetics 196: 1145–1152.
Xie, Z., L. Wang, L. Wang, Z. Wang, Z. Lu *et al.*, 2016 Mutation rate analysis via parent–
progeny sequencing of the perennial peach. I. A low rate in woody perennials and a
higher mutagenicity in hybrids. Proc. R. Soc. B Biol. Sci. 283: 20161016
Xu, S., J. Stapley, S. Gablenz, J. Boyer, K. J. Appenroth *et al.*, 2019 Low genetic variation is
associated with low mutation rate in the giant duckweed. Nat. Commun. 10: 1243.
Yang, S., L. Wang, J. Huang, X. Zhang, Y. Yuan *et al.*, 2015 Parent-progeny sequencing
indicates higher mutation rates in heterozygotes. Nature 523: 463–467.
Zheng, X., D. Levine, J. Shen, S. M. Gogarten, C. Laurie *et al.*, 2012 A high-performance
computing toolset for relatedness and principal component analysis of SNP data.
Bioinforma. Oxf. Engl. 28: 3326–3328.
Zhu, Y. O., M. L. Siegal, D. W. Hall, and D. A. Petrov, 2014 Precise estimates of mutation rate
and spectrum in yeast. Proc. Natl. Acad. Sci. U. S. A. 111: E2310-2318.
757
758

Supporting Information

Figure S1 Visualization of a low-quality genomic region that was filtered from our analyses.

Figure S2 Allelic coverage at heterozygous sites before and after filtering.

Figure S3 Visualization of a validated *de novo* mutation from both Illumina short read and

Sanger sequencing data.

Figure S4 Visualisation of a putative *de novo* mutation that was identified as a likely

sequencing error

Figure S5 Visualisation of a putative *de novo* mutation that was identified as a likely genome

misassembly error

Table S1 List of *de novo* mutations identified in two species of duckweed

Table S2 Mutation rate estimates under the observed allelic bias at *de novo* mutations

Table S3 Mutation rate estimates under several scenarios of allelic bias at *de novo* mutant

sites in salt stressed mutation accumulation lines

Table S4 Mutation rate estimates under several scenarios of allelic bias at *de novo* mutant

sites in control mutation accumulation lines

Table S5 Number and fraction of callable sites under several scenarios of allelic bias at *de*

novo mutant sites

Table S6 Summary of mutation rate estimates retrieved from the literature