Phagocytosis and self-destruction execute dendrite degeneration of *Drosophila* sensory

neurons at distinct levels of NAD$^+$ reduction

Hui Ji[1,3], Maria L. Sapar[1,2,3], Ankita Sarkar[1], Bei Wang[1], and Chun Han[1*]

[1]Weill Institute for Cell and Molecular Biology and Department of Molecular Biology and

Genetics, Cornell University, Ithaca, NY 14853, USA

[2]Current address: New York Stem Cell Foundation, New York, NY 10019, USA

[3]These authors contributed equally to this work

*Correspondence and Lead Contact: chun.han@cornell.edu

RUNNING TITLE

NAD$^+$ reduction coordinates PS-mediated phagocytosis and self-destruction of dendrites

Keywords

phosphatidylserine, PS exposure, phagocytosis, phagocyte, self-destruction, Wallerian

degeneration, dendrite degeneration, NAD$^+$, Nmnat, Sarm, Axed, calcium, *Drosophila*, da

neuron

ABSTRACT

After injury, severed dendrites and axons expose the "eat-me" signal phosphatidylserine (PS) on their surface and degenerate by disassembly. While axon degeneration is controlled by a conserved "axon-death" pathway that is thought to activate self-destruction, how PS exposure is regulated by this pathway and whether PS-induced phagocytosis contributes to neurite breakdown *in vivo* remain unknown. Here we show that in *Drosophila* sensory dendrites, PS exposure and self-destruction are triggered by two distinct levels of NAD^+ reduction downstream of Sarm activation. Surprisingly, phagocytosis is the main driver of dendrite degeneration induced by both genetic NAD^+ disruptions and injury. Furthermore, the axon-death factor Axed is only partially required for self-destruction of injured dendrites, acting in parallel with PS-induced phagocytosis. Lastly, injured dendrites exhibit a unique rhythmic calcium flashing that correlates with self-destruction. Therefore, a special genetic program coordinates PS exposure and self-destruction in injury-induced dendrite degeneration *in vivo*.

INTRODUCTION

Physical insults to the nervous system often disrupt neuronal connectivity and function by damaging dendritic or axonal processes of neurons. Injured axons break down through a series of stereotypical events collectively called Wallerian degeneration (WD) (Coleman and Freeman, 2010; Waller, 1850). Dendrites undergo a similar program of degeneration after injury (Tao and Rolls, 2011). Before neurons can regenerate their processes and restore connections, the debris from damaged neurites has to be promptly cleared by phagocytes, which are cells that engulf dead cells or cell debris (Sapar and Han, 2019). Inefficient clearance can lead to neuroinflammation and further exacerbate the damage to the surrounding tissues (Davies et al., 2019; Galloway et al., 2019). Although WD is mainly considered to be a neurite-intrinsic, self-destructive process (Gerdts et al., 2016), whether phagocytosis actively contributes to degeneration of injured neurites *in vivo*, rather than merely passively removing neuronal debris, remains unclear.

WD is governed by an evolutionarily conserved pathway, which is also called "axon-death" pathway because it was discovered in studies focused primarily on axon degeneration in *Drosophila* and rodents (Freeman et al., 2003; Gerdts et al., 2016; Sapar and Han, 2019). This pathway is centered on nicotinamide adenine dinucleotide (NAD^+), an essential cellular

metabolite that is locally depleted after axons are injured. The signaling cascade starts with

injury-induced activation of the E3 ubiquitin ligase Highwire/Phr1 in severed axons (Babetto et

al., 2013; Xiong et al., 2012), which in turn causes degradation of nicotinamide mononucleotide

adenyltransferase (Nmnat), an enzyme required for the synthesis of NAD^+ (Zhai et al., 2009).

The resultant decrease of NAD^+ levels (Sasaki et al., 2016), together with an accumulation of the

NAD^+ precursor nicotinamide mononucleotide (NMN) (Di Stefano et al., 2015; Liu et al., 2018),

activates Sarm/SARM1, a sterile alpha/Armadillo/Toll-Interleukin receptor homology domain

protein (Bratkowski et al., 2020; Figley et al., 2021; Jiang et al., 2020; Osterloh et al., 2012;

Shen et al., 2021). Sarm/SARM1 carries NADase activity; thus its activation causes catastrophic

NAD^+ depletion, which drives axon breakdown through unclear mechanisms (Gerdts et al.,

2015; Wang et al., 2005).

Besides this core Highwire/Phr1-Nmnat-NAD^+-Sarm/SARM1 pathway, several other

factors have also been found to act in WD. Downstream of *Sarm*, *axundead* (*axed*) is required for

axon degeneration of olfactory receptor neurons (ORNs) and wing sensory neurons in

Drosophila (Neukomm et al., 2017). The loss of *axed* blocks axon degeneration even when Sarm

is dominantly activated, raising the possibility that Axed activation, rather than NAD^+ depletion,

is the key switch of WD (Neukomm et al., 2017). In addition, *pebbled* (*peb*) encodes a

Drosophila transcription factor required for axon degeneration of glutamatergic but not

cholinergic sensory neurons in the wing (Farley et al., 2018). Lastly, the dual leucine kinase

(DLK)/c-Jun N-terminal kinase (JNK) stress pathway also contributes to, but is not required for,

WD in both *Drosophila* and mice (Miller et al., 2009; Yang et al., 2015). Although how these

factors interact with the core pathway to promote axon degeneration is still mysterious, it is

generally believed that catastrophic NAD^+ depletion caused by Sarm activity initiates a neurite-

intrinsic self-destruction program that ultimately is responsible for WD of axons (Babetto et al.,

2013; Gerdts et al., 2015; Gerdts et al., 2016; Neukomm et al., 2017). While the WD pathway is

primarily characterized in axons, evidence suggests that NAD^+ reduction is also an essential step

in injury-induced dendrite degeneration (Sapar et al., 2018; Tao and Rolls, 2011). However,

which components of the WD pathway are conserved in dendrites remains unknown.

Neuronal debris is recognized by resident phagocytes of the nervous system through

specific "eat-me" signals exposed on the neuronal surface. A highly conserved eat-me signal is

phosphatidylserine (PS), a negatively charged phospholipid normally found in the inner leaflet of

the plasma membrane of healthy cells. During apoptosis, PS is externalized to the outer leaflet of

the plasma membrane to mark the cell for engulfment (Leventis and Grinstein, 2010). Genetic

analyses of certain PS-binding bridging molecules and cell membrane receptors in mice and

zebrafish suggest that PS recognition contributes to the phagocytosis of neurons (Fourgeaud et

al., 2016; Mazaheri et al., 2014; Nandrot et al., 2007). Similarly, clearance of injured axons and

dendrites in *Drosophila* requires Draper (Drpr), an engulfment receptor that binds to PS

(Freeman et al., 2003; Han et al., 2014; MacDonald et al., 2006; Tung et al., 2013). PS exposure

has also been directly observed on injured axons of mouse neurons *in vitro* (Shacham-Silverberg

et al., 2018; Sievers et al., 2003) and on injured dendrites of *Drosophila* sensory neurons *in vivo*

(Sapar et al., 2018). Although axonal PS exposure can be induced independent of axon

degeneration *in vitro* (Shacham-Silverberg et al., 2018), ectopically induced PS exposure

resulted in engulfment-dependent neurite reduction of otherwise healthy neurons in both the

central nervous system (CNS) and the peripheral nervous system (PNS) of *Drosophila* (Sapar et

al., 2018), pointing to a dominant effect of PS exposure in induced phagocytosis. Recent studies

revealed a link between neuronal PS exposure and the WD pathway. Overexpression of Wld[S], a

fusion protein containing the full-length murine Nmnat1 (Mack et al., 2001), in *Drosophila*

sensory neurons suppresses PS exposure of injured dendrites (Sapar et al., 2018). In addition,

Sarm1 ablation and NAD^+ supplementation in neuronal culture reduce PS exposure on injured

axons (Shacham-Silverberg et al., 2018). These observations raise two important questions: How

does the WD pathway regulate and coordinate both neurite PS exposure and self-destruction?

What are the relative contributions of PS-mediated phagocytosis and neurite self-destruction in

WD *in vivo*?

To address these questions, we utilized *Drosophila* class IV dendritic arborization (C4da)

neurons on the larval body wall, an established *in vivo* model of injury-induced dendrite

degeneration (Han et al., 2014). In this system, degenerating dendrites of C4da neurons are

phagocytosed by epidermal cells through the engulfment receptor Drpr. Here we show that PS

exposure and dendrite self-destruction are two steps activated at different levels of NAD^+

reduction, and that *in vivo*, phagocytosis is the main driving force of dendrite degeneration

induced by both genetic NAD^+ perturbations and injury. Interestingly, injury and genetic NAD^+

perturbations cause dendrite degeneration through different cellular mechanisms in membrane

disruption, dendrite calcium dynamics, and phagocytosis dependence. Furthermore, Axed

specifically promotes, but is not required for, dendrite self-destruction, suggesting different

mechanisms of injury-induced degeneration in axons and dendrites. Lastly, we report a

previously unknown rhythmic calcium flashing specifically in injured dendrites, which is

disrupted by Wld^S overexpression and *axed* loss, indicating a potential role of calcium flashing in

dendrite self-destruction.

RESULTS

***Nmnat* knockout induces spontaneous dendrite degeneration due to Sarm-mediated NAD$^+$**

reduction

NAD$^+$ reduction is required for PS exposure on injured dendrites (Sapar et al., 2018). To

determine if NAD$^+$ loss is also sufficient to cause neuronal PS exposure, we decided to first

investigate the impact of removing *Nmnat* from C4da neurons, as the loss of *Nmnat* is expected

to cause cell-autonomous NAD$^+$ reduction. *Nmnat* LOF is known to induce degeneration of eye

photoreceptors (Zhai et al., 2006) and wing sensory neurons (Neukomm et al., 2017). However, a

previous study found that C4da neurons mutant for *Nmnat* did not show dendrite degeneration,

despite displaying dendrite reduction and axon degeneration (Wen et al., 2011). To reexamine

the LOF phenotype of *Nmnat*, we used CRISPR-TRiM, a tissue-specific mutagenesis method we

previously developed (Poe et al., 2019), to knock out *Nmnat* in C4da neurons. In this method,

Nmnat is knocked out by C4da-specific *ppk-Cas9* and two ubiquitously expressed guide-RNAs

(gRNAs) targeting *Nmnat*. To distinguish dendrite reduction caused by degeneration from that

caused by growth defects, we used CD4-tdTomato (CD4-tdTom) to label C4da dendrites.

Because tdTom is stable in phagosomes, the presence of tdTom-labeled dendrite debris in

epidermal cells is an indication of dendrite breakdown and subsequent engulfment (Han et al.,

2014).

As expected, *Nmnat* knockout (KO) in C4da neurons (Figure 1B) caused strong dendritic

reduction (Figure 1G) in wandering 3rd instar larvae as compared to the control (Figures 1A). In

addition, dendrite debris was observed to spread in epidermal cells underneath and near the

dendrites of *Nmnat* KO neurons (Figures 1B and 1H), suggesting that some dendrites had

degenerated and were engulfed by epidermal cells. Unlike degenerating dendrites observed

during developmental pruning or after injury (Han et al., 2011; Han et al., 2014), the remaining

dendrites of *Nmnat* KO neurons did not show obvious blebbing or fragmentation, which may

explain why dendrite degeneration was not detected previously in *Nmnat* mutant C4da neurons,

considering that the membrane GFP marker used to label dendrites in the previous study is

rapidly degraded by epidermal cells once engulfed and thus cannot visualize phagosomes (Han et

al., 2014; Wen et al., 2011).

Nmnat protects neurons by both synthesizing NAD^+ and functioning as a chaperon

protein (Ali et al., 2013). To verify that the observed dendrite degeneration is due to the loss of

Nmnat enzymatic activity, we tried to rescue *Nmnat* KO neurons by overexpressing Wld^S, which

contains full NMNAT activity (Mack et al., 2001), and $Wld^{S\text{-}dead}$, a mutant version of Wld^S that

Figure 1: *Nmnat* knockout induces spontaneous dendrite degeneration due to Sarm-mediated NAD^+ reduction

(A-F) Partial dendritic fields of control (A), *Nmnat* KO (B), *Nmnat* KO + Wld^S OE (C), *Nmnat* KO + $Wld^{S\text{-}dead}$ OE (D), $Sarm^{4705/4621}$ (E), and *Nmnat* KO + $Sarm^{4705/4621}$ (F) ddaC C4da neurons. Neurons were labeled by *ppk>CD4-tdTom* (A-D) and *ppk-CD4-tdTom* (E and F). The C4da-specific KO was induced by *ppk-Cas9*. Scale bars, 25 μm.

(G) Quantification of dendrite length. n = number of neurons: control (n =11, 6 animals); *Nmnat* KO (n = 19, 10 animals); *Nmnat* KO + Wld^S OE (n = 17, 10 animals); *Nmnat* KO + $Wld^{S\text{-}dead}$ OE (n = 20, 11 animals); $Sarm^{4705/4621}$ (n = 12, 6 animals); *Nmnat* KO + $Sarm^{4705/4621}$ (n = 13, 8 animals). One-way ANOVA and Tukey's test.

(H) Quantification of debris coverage ratio, which is the percentage of debris area normalized by dendrite area ratio. Number of neurons: same as in (G). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method.

For all quantifications, ***p≤0.001; n.s., not significant. The significance level above each genotype is for comparison with the control. Black bar, mean; red bar, SD.

cannot synthesize NAD$^+$ but maintains the chaperon function (Avery et al., 2009). Wld^S

overexpression (OE) rescued the degeneration of *Nmnat* KO neurons and restored dendrite

morphology to the wildtype level (Figures 1C, 1G, and 1H), while Wld^{S-dead} OE did not change

dendrite length or debris level of *Nmnat* KO neurons (Figures 1D, 1G, and 1H). These results

suggest that NAD$^+$ reduction is responsible for the dendrite degeneration of *Nmnat* KO neurons.

We next asked whether Sarm plays a role in *Nmnat* KO-induced dendrite degeneration.

Indeed, *Sarm* LOF completely blocked dendrite degeneration of *Nmnat* KO neurons (Figures 1F,

1G, 1H), as evident in the absence of dendrite debris in epidermal cells. However, these neurons

showed more variable dendrite length (Figure 1G) compared to *Sarm* LOF alone (Figures 1E,

1G, and 1H), perhaps due to the loss of Nmnat chaperon function (Wen et al., 2011).

Importantly, these results demonstrate that *Sarm* is required for the dendrite degeneration of

Nmnat KO neurons. All the results above together suggest that *Nmnat* LOF in neurons cause

spontaneous dendrite degeneration through Sarm-mediated NAD$^+$ loss.

PS exposure-mediated phagocytosis causes dendrite degeneration of *Nmnat* KO neurons

To ask whether *Nmnat* KO also causes neuronal PS exposure, we used an established method to

visualize PS exposure on C4da dendrites. In this method, the fluorescent PS sensor GFP-Lact is

expressed by the larval fat body and secreted into the hemolymph (Sapar et al., 2018). As C4da

dendrites are largely exposed to the hemolymph, GFP-Lact labeling allows visualization of

dynamic PS exposure on dendrites in intact live animals (Sapar et al., 2018). We found that GFP-

Lact strongly labeled *Nmnat* KO neurons at distal branches that underwent degeneration (Figure

2B), while wildtype C4da neurons showed no labeling (Figure 2A). Interestingly, we also

observed weaker GFP-Lact labeling on dendrite segments that did not display obvious signs of

degeneration (outlined in Figure 2B), suggesting that PS exposure may precede dendrite

breakdown, instead of being merely a consequence of dendrite degeneration. This conclusion

was further corroborated by time-lapse imaging of *Nmnat* KO neurons (Video 1) and kymograph

analysis (Figure 2C): PS exposure (indicated by white dotted lines) occurred well ahead of

dendrite fragmentation (indicated by blue dotted lines).

As *Sarm* is required for dendrite degeneration of *Nmnat* KO neurons, we asked if Sarm

also regulates PS exposure in these neurons. *Nmnat* KO neurons showed no PS exposure in the

 Sarm mutant background (Figure 2D), suggesting that Sarm-mediated NAD$^+$ reduction is also

 responsible for inducing the observed PS exposure.

Figure 2: PS exposure-mediated phagocytosis causes dendrite degeneration of *Nmnat* KO neurons

(A) A control ddaC neuron showing the lack of GFP-Lact labeling. Insets show close-ups of an intact dendrite. Scale bar, 25 μm.

(B) A *Nmnat* KO ddaC neuron undergoing spontaneous degeneration and showing labeling by GFP-Lact. In the inset, the outlined dendrite segments do not show obvious blebbing or fragmentation but are weakly labeled by GFP-Lact. Scale bar, 25 μm.

(C) Kymographs of tdTom signal (dendrite) and GFP-Lact signal along a dendrite segment of a *Nmnat* KO ddaC neuron. White dotted lines indicate the timing of GFP-Lact initial binding to the dendrite; blue dotted lines indicate timing of the dendrite fragmentation based on the continuity of the dendrite signal. Scale bars, 10 μm horizontal, 60 min vertical. See also Video 1.

(D) Dendrites of a *Nmnat* KO + *Sarm$^{4705/4621}$* ddaC neuron showing the lack of GFP-Lact labeling. Insets show close-ups of an intact dendrite. Scale bar, 25 μm.

(E-I) Partial dendritic fields of control (E), *Nmnat* KO (F), *drpr$^{-/-}$* (G), *Nmnat* KO + *drpr$^{-/-}$* (H) and *Nmnat* KO + ATP8A OE (I) neurons. Scale bars, 25 μm.

(J) Quantification of dendrite length. n = number of neurons: control (n = 13, 7 animals); *Nmnat* KO (n = 13, 7 animals); *drpr$^{-/-}$* (n = 14, 8 animals); *Nmnat* KO + *drpr$^{-/-}$* (n = 22, 14 animals); ATP8A OE (n=17, 9 animals); *Nmnat* KO + ATP8A OE (n=23, 9 animals). One-way ANOVA and Tukey's test.

(K) Quantification of debris coverage ratio. Number of neurons: same as in (J). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method.

(L) A time series of *Nmnat* KO + *drpr$^{-/-}$* dendrites. Yellow arrowheads indicate growth of dendrites compared to the previous frame; blue arrows indicate retractions of dendrites compared to the previous frame. Scale bar, 10 μm. See also Video 2.

In all panels, neurons were labeled by *ppk-CD4-tdTom* (A-H) and *ppk-MApHS* (I). C4da-specific KO was induced by *ppk-Cas9*. For all quantifications, *p≤0.05, **p≤0.01, ***p≤0.001; n.s., not significant. The significance level above each genotype is for comparison with the control. Black bar, mean; red bar, SD.

The observation of PS exposure on *Nmnat* KO dendrites raises the question of whether

PS exposure contributes to dendrite breakdown by inducing phagocytic attacks from epidermal

cells. Because PS-mediated epidermal engulfment of dendrites requires Drpr (Sapar et al., 2018),

we examined *Nmnat* KO dendrites in *drpr* mutant larvae. Strikingly, *drpr* LOF completely

blocked dendrite degeneration of *Nmnat* KO neurons (Figure 2H as compared to Figures 2E-2G;

Figures 2J and 2K). These dendrites exhibited dynamic extension and retraction behaviors

(Figure 2L and Video 2), demonstrating that they were not fragmented dendrites that failed to be

cleared. Next, to examine whether dendrite breakdown induced by *Nmnat* KO depends on PS

exposure, we tried to rescue *Nmnat* KO neurons by overexpressing ATP8A. ATP8A is the

Drosophila ortholog of PS flippase responsible for keeping PS in the inner leaflet of the plasma

membrane (Raiders et al., 2021; Tanaka et al., 2011), whose LOF in da neurons causes dendritic

PS exposure (Sapar et al., 2018). Indeed, ATP8A OE completely blocked dendrite degeneration

of *Nmnat* KO neurons and resulted in even less dendrite debris than the wildtype control (Figures

1I and 1K).

Thus, these data demonstrate that Sarm-mediated NAD^+ reduction causes dendrite PS

exposure, which in turn induces phagocytic attack to drive dendrite degeneration of *Nmnat* KO

neurons.

Wld^S OE and *Sarm* KO suppress PS exposure and fragmentation of injured dendrites

Wld^S protects injured neurites from degenerating by maintaining the NAD^+ level in the neurites.

We previously found that overexpressing Wld^S in C4da neurons blocked fragmentation and PS

exposure of ablated dendrites at 10 hrs after injury (AI) (Sapar et al., 2018). To further

investigate the role of the WD pathway in neuronal PS exposure, we examined the effects of

overexpressing Wld^S and knocking out *Sarm* in C4da neurons at 24 hrs after laser-severing of

dendrites. Neuronal-specific KO was conducted using *SOP-Cas9*, which is active in precursor

cells of da neurons, to minimize potential gene perdurance (Poe et al., 2019). As expected, Wld^S

OE blocked degeneration and clearance of injured dendrites also at 24 hrs AI (Figure 3C, as

compared to the control in Figure 3B), even though the injured arbors were greatly simplified as

compared to uninjured dendrites (Figures 3A and 3E). Neuronal-specific *Sarm* KO showed

similar effects in injured dendrites (Figures 3D and 3E) to those of Wld^S OE.

We next examined the effects of Wld^S OE and *Sarm* KO on dendritic PS exposure 24 hrs

AI. Wld^S OE prevented PS exposure on severed dendrites, even on branches that showed

blebbing and breakage (Figures 3F and 3I). *Sarm* KO also effectively blocked PS exposure on

injured dendrites (Figures 3G and 3I), with occasionally strong PS labeling on fragmented distal

terminal branches (Figures 3H and 3I). The above data in dendrite injury together suggest that

Sarm-mediated NAD$^+$ reduction causes both PS exposure and degeneration of injured dendrites.

Figure 3: Wld^S OE and *Sarm* KO suppress PS exposure and fragmentation of injured dendrites

(A) Partial dendritic field of an uninjured ddaC control neuron.

(B-D) Partial dendritic fields of control (B), Wld^S OE (C) and *Sarm* KO (D) ddaC neurons 24 hrs after injury (AI). Blue asterisk, injury site. Yellow dots outline regions covered by injured dendrites.

(E) Quantification of dendrite density (1000 x dendritic length/area) in regions covered by injured dendrites. n = number of neurons: control no injury (NI)

(n = 10, 10 animals); control (n = 12, 6 animals); Wld^S OE (n = 13, 6 animals); *Sarm* KO (n = 19, 6 animals).

(F and G) Wld^S OE (F) and *Sarm* KO (G) ddaC neurons 24 hrs AI, showing unfragmented dendrites and lack of GFP-Lact binding to severed dendrites (yellow arrowheads).

(H) Dendrites of a *Sarm* KO ddaC neuron 24 hrs AI, showing infrequent GFP-Lact binding at fragmented terminal branches (yellow arrowheads).

(I) Quantification of GFP-Lact binding on injured dendrites. The GFP-Lact intensity is shown for both background epidermal regions and injured dendrites. The outliers in *Sarm* KO dendrite dataset correspond to infrequent degenerating terminal branches. Background: epidermal regions without dendrites. n = number of measurements: control background (n = 39) and control dendrite (n/a, no remaining dendrites), 7 animals; Wld^S OE background (n = 18) and Wld^S OE dendrite (n = 18), 4 animals; *Sarm* KO background (n = 48) and *Sarm* KO dendrite (n = 48), 9 animals.

In all panels, neurons were labeled by *ppk-MApHS* (A-C; only tdTom channel is shown), *ppk>CD4-tdTom* (D), and *ppk-CD4-tdTom* (F-H). Neuronal-specific KO was induced by *SOP-Cas9*. For all quantifications, One-way ANOVA and Tukey test; ***p≤0.001; n.s., not significant; n/a., not applicable. The significance level above each genotype in (E) is for comparison with the control. Black bar, mean; red bar, SD. Scale bars, 25 μm.

212 The PS exposure on fragmented terminal dendrites of *Sarm* KO neurons could be due to

213 relatively faster NAD$^+$ turnover in those local branches even in the absence of Sarm.

214 Ectopic PS exposure causes injured dendrites of Wld^S OE and *Sarm* KO neurons to

215 degenerate

The absence of PS exposure on injured dendrites of Wld^S OE and *Sarm* KO neurons raises the

question of whether the degeneration defects of these neurons are due to the lack of PS-mediated

epidermal attack on dendrites. To test this hypothesis, we ectopically induced neuronal PS

exposure by knocking out *CDC50*, which encodes a chaperone necessary for the function of

Drosophila flippases (Tanaka et al., 2011), and by overexpressing TMEM16F, a mammalian PS

scramblase (Segawa et al., 2011). These manipulations in C4da neurons cause mild but

appreciable phagocytosis-dependent dendrite loss (Sapar et al., 2018). *CDC50* KO or TMEM16F

OE alone in Wld^S OE neurons caused partial or complete degeneration of injured dendrites and

the accompanying epidermal engulfment (Figures 4A, 4B, and 4E). A much stronger effect was

observed when *CDC50* KO and TMEM16F OE were combined in Wld^S OE neurons:

Figure 4: Ectopic PS exposure causes injured dendrites of Wld^S OE and *Sarm* KO neurons to degenerate.

(A-D) Partial dendritic fields of Wld^S OE + TMEM16F OE (A), Wld^S OE + *CDC50* KO (B), Wld^S OE + TMEM16F OE + *CDC50* KO (C) and *Sarm* KO + TMEM16F OE + *CDC50* KO (D) ddaC neurons 24 hrs AI. Blue asterisk, injury site. Yellow dots outline regions covered by injured dendrites. Neurons were labeled by *ppk-MApHS* (A, B, and C) and *ppk-CD4-tdTom* (D). Neuronal-specific KO was induced by *ppk-Cas9* (B and C) and *SOP-Cas9* (D). Scale bars, 25 μm.

(E) Quantification of dendrite density. n = number of neurons: control (n = 12, 6 animals); Wld^S OE (n = 13, 6 animals); Wld^S OE + TMEM16F OE (n = 23, 8 animals); Wld^S OE + *CDC50* KO (n = 17, 8 animals); Wld^S OE + TMEM16F OE + *CDC50* KO (n = 16, 8 animals); *Sarm* KO (n = 19, 6 animals); *Sarm* KO + TMEM16F OE + *CDC50* KO (n = 13, 7 animals). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method. The dataset for Wld^S OE and *Sarm* KO is the same as in Figure 3E.

For all quantifications, *p≤0.05, ***p≤0.001; n.s., not significant. The significance level above each genotype is for comparison with the control. Black bar, mean; red bar, SD.

Degeneration and clearance of injured dendrites were restored to the wildtype level (Figures 4C

and 4E). Similarly, *CDC50* KO and TMEM16F OE caused injured dendrites of *Sarm* KO

neurons to completely degenerate (Figures 4D and 4E). Because injury induces a more rapid and

stronger PS exposure on dendrites than *CDC50* KO or TMEM16F OE (Sapar et al., 2018), our

data strongly suggest that injury-induced PS exposure is a major driver of dendrite breakdown by

activating phagocytic attack by epidermal cells.

Injury and NAD$^+$ depletion cause phagocytosis-independent dendrite self-destruction

Although PS-mediated phagocytosis is sufficient to break down injured dendrites, blocking

phagocytosis with *drpr* LOF failed to prevent the fragmentation of injured dendrites 20 hrs AI,

even though the dendrite debris was left unengulfed (Figures 5A-5C). A neurite self-destruction

Figure 5: Injury and NAD$^+$ depletion cause phagocytosis-independent dendrite self-destruction

(A-B) Partial dendritic fields of ddaC neurons in control (A) and *drpr$^{-/-}$* (B) animals 20 hrs AI. The dendrite debris lining up in the original dendritic pattern in (B) indicates dendrite fragmentation and the lack of epidermal engulfment. Blue asterisk, injury site. Yellow dots outline regions covered by injured dendrites.

(C) Quantification of dendrite density. n = number of neurons: control no-injury (NI) (n = 10, 10 animals); control AI (n = 12, 6 animals); *drpr$^{-/-}$* NI (n = 6, 5 animals); *drpr$^{-/-}$* AI (n = 13, 5 animals). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method.

(D-G) ddaC neurons in *Sarm^{GOF}* OE 24 hrs after egg laying (AEL) (D), *Sarm^{GOF}* OE + *drpr$^{-/-}$* 24 hrs AEL (E), *Sarm^{GOF}* OE 48 hrs AEL (F) and *Sarm^{GOF}* OE + *drpr$^{-/-}$* 48 hrs AEL (G).

(H-I) Partial dendritic fields of *Sarm^{GOF}* OE (H) and *Sarm^{GOF}* OE + *drpr$^{-/-}$* (I) ddaC neurons at the wandering stage.

(J) Quantification of dendrite degeneration showing percentages of neurons undergoing partial degeneration, complete degeneration, and no degeneration. n = number of neurons: *Sarm^{GOF}* OE 24 hrs AEL (n = 31, 7 animals); *Sarm^{GOF}* OE + *drpr$^{-/-}$* 24 hrs AEL (n = 49, 17 animals); *Sarm^{GOF}* OE 48 hrs AEL (n = 17, 7 animals); *Sarm^{GOF}* OE + *drpr$^{-/-}$* 48 hrs AEL (n = 33, 11 animals); *Sarm^{GOF}* OE wandering (n = 19, 14 animals); *Sarm^{GOF}* OE + *drpr$^{-/-}$* wandering (n = 10, 6 animals). Freeman-Halton extension of Fisher's exact test.

In image panels above, neurons were labeled by *ppk-MApHS* (A and B) and *ppk>CD4-tdTom* (D-I). C4da-specific KO was induced by *ppk-Cas9*. For all quantifications, **p≤0.01, ***p≤0.001; n.s., not significant. Black bar, mean; red bar, SD. Scale bars, 25 μm.

program triggered by NAD$^+$ depletion is known to drive WD of cultured neurons in the absence
of phagocytes (Gerdts et al., 2015). To test the possibility that a further NAD$^+$ reduction beyond
the level that triggers PS exposure causes phagocytosis-independent dendrite self-destruction, we
tested the effect of overexpressing a gain-of-function Sarm (Sarm^{GOF}) in C4da neurons, because
Sarm^{GOF} potently depletes NAD$^+$ and causes dominant neurodegeneration (Neukomm et al.,
2017). Indeed, Sarm^{GOF} OE caused complete dendrite degeneration in most neurons as early as
24 hrs after egg laying (AEL) (Figures 5D and 5J), while *Nmnat* KO did not cause dendrite
degeneration until 88-91 hrs AEL. Supporting the role of engulfment in NAD$^+$ reduction-induced
dendrite degeneration, *drpr* LOF strongly suppressed dendrite degeneration of Sarm^{GOF} OE
neurons at both 24 hrs AEL and 48 hrs AEL (Figures 5E, 5G, and 5J). However, *drpr* LOF failed
to prevent dendrite fragmentation of Sarm^{GOF} OE neurons at wandering 3rd instar (Figures 5H-
5J), suggesting that NAD$^+$ depletion is able to cause phagocytosis-independent dendrite self-
destruction. These results together support the hypothesis that dendrite self-destruction induced
by NAD$^+$ depletion is responsible for the fragmentation of injured dendrites when phagocytosis
is suppressed.

Axed promotes self-destruction of injured dendrites in parallel with PS-mediated
phagocytosis

To understand how other components of the WD pathway regulate dendrite degeneration, we
examined LOF of *axed* and *peb*. Tissue-specific KO of *axed* in Or22a olfactory neurons, an
established axon-injury model (MacDonald et al., 2006), effectively blocked degeneration of
severed axons 7 days after antenna ablation (Figure S1A-S1D, and S1G), demonstrating the
effectiveness of the gRNAs. In C4da neurons, both KO and knockdown (KD) of *axed* also
efficiently blocked injury-induced axon degeneration (Figures S1H-S1K), consistent with the
role of Axed in WD. To test if Axed is involved in PS exposure-induced dendrite degeneration
resulting from *Nmnat* LOF, we knocked out both *Nmnat* and *axed* in da neurons. Unlike *Sarm*
KO, which fully rescued *Nmnat* KO-induced degeneration (Figures S1M, S1O, S1P, and S1Q),
axed KO did not affect dendrite degeneration caused by *Nmant* KO (Figures S1L, S1N, S1P, and
S1Q), suggesting that *axed* is not required for the PS exposure induced by NAD$^+$ reduction.

Figure S1: Axed is not involved in PS-mediated phagocytosis

(A-F) Axons of control NI (A), control 7 days AI (B), *axed* KO NI (C), *axed* KO 7 days AI (D), *peb* KO NI (E), and *peb* KO 7 days AI (F) Or22a neurons in 14-day-old adult brains. Or22a neurons were labeled by *Or22a>CD8-GFP* and KO was induced by *ey-Cas9*. Scale bars, 25 μm.

(G) Quantification of GFP brightness of Or22a glomeruli. n = number of brains: control NI 14 days (n = 19); control 7 days AI 14 days (n = 15); *axed* KO NI 14 days (n = 16); *axed* KO 7 days AI 14 days (n = 14); *peb* KO NI 14 days (n = 18); *peb* KO 7 days AI 14 days (n = 20). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method. *p≤0.05, ***p<0.001, n.s., not significant. The significance level above each genotype is for comparison with the control. Statistical comparisons for *peb* KO glomeruli are not shown due to the variation within the group of *peb* KO NI 14 days. Black bar, mean; red bar, SD.

(H-J) Injured axons of wildtype 24 hrs AI (H), *axed* KO AI 24 hrs AI (I), and *axed* KD AI 24 hrs AI (J). Neurons were labeled by *ppk-MApHS*. Neuronal-specific KO was induced by *SOP-Cas9*, and neuronal-specific KD was driven by *21-7 Gal4*. Scale bars, 25 μm.

(K) Quantification of axon fragmentation showing percentages of neurons undergoing no fragmentation, and complete fragmentation of injured axons at 24 hrs AI. n = number of neurons: wildtype 24 hrs AI (n = 20, 12 animals); *axed* KO 24 hrs AI (n = 17, 9 animals); *axed* KD 24hrs AI (n = 20, 8 animals). Freeman-Halton extension of Fisher's exact test. For all quantifications, ***p≤0.001; n.s., not significant.

(L-O) Partial dendritic fields of *Nmnat* KO + *axed* KO (L and N) and *Nmnat* KO + *Sarm* KO (M and O) ddaC C4da neurons. Neurons were labeled by *ppk>CD4-tdTom* (L-O). The C4da-specific KO was induced by *ppk-Cas9* (L-M) or by *SOP-Cas9* (N-O). Scale bars, 25 μm.

(legends continued on the next page)

We next asked if Axed is involved in injury-induced dendrite degeneration. Injured

dendrites of wildtype neurons completely fragmented by 10 hrs AI (Figures 6A and 6E). *axed*

KO in neurons resulted in incomplete and no fragmentation of injured dendrites only in small

fractions (25% and 8%, respectively) of neurons (Figures 6B and 6E), suggesting that *axed* is not

absolutely required for the degeneration of injured dendrites. We next asked if Axed specifically

contributes to self-destruction of injured dendrites. Because PS-induced phagocytosis may be

sufficient to break down injured dendrites by 10 hrs AI, thereby masking dendrite protection by

axed KO, we examined the effect of *axed* KO in *drpr* mutant. *drpr* LOF alone caused incomplete

and no dendrite fragmentation in 36% and 23% injured neurons, respectively (Figures 6C and

6E), consistent with the idea that phagocytosis contributes significantly to dendrite breakdown at

this stage. In comparison, *axed* KO in *drpr* mutant strongly blocked dendrite degeneration at 10

hrs AI, with 56% incomplete fragmentation and 36% no fragmentation (Figures 6D and 6E),

suggesting that *axed* KO and *drpr* LOF have an additive effect in dendrite protection.

Surprisingly, this combined protection was significantly weakened by 24 hrs AI, with only 47%

incomplete fragmentation and 5% no fragmentation (Figures 6F-6J). These data suggest that

Axed promotes dendrite self-destruction in parallel with PS-induced phagocytosis; however,

other factors can also cause self-destruction in the absence of Axed.

In contrast to *axed*, *peb* KO in C4da neurons did not block dendrite fragmentation after

injury but altered dendrite morphology (Figures S1R-S1T). Unexpectedly, knocking out *peb*

from precursors of Or22a neurons caused loss of some or all of Or22a axons in uninjured adult

brains (Figure S1G), suggesting that *peb* plays a role in Or22a development or axon patterning.

The remaining axons of *peb* KO neurons did not show defects in axon degeneration after injury (Figures S1E-S1G). These results suggest that Peb may be a neuronal type-specific modulator of the WD pathway and is not required for dendrite degeneration of da neurons.

Figure 6: Axed drives dendrite self-destruction in parallel with PS-mediated phagocytosis

(A-D) Partial dendritic fields of wildtype (A), *axed* KO (B), *drpr⁻/⁻* (C), and *axed* KO + *drpr⁻/⁻* (D) ddaC neurons 10 hrs AI. (B) shows partial fragmentation; (C) and (D) show no fragmentation.

(E) Quantification of dendrite fragmentation showing percentages of neurons undergoing no fragmentation, partial fragmentation, and complete fragmentation of injured dendrites at 10 hrs AI. n = number of neurons: wildtype (n=18, 9 animals); *axed* KO (n = 24, 10 animals); *drpr⁻/⁻* (n = 22, 11 animals); *axed* KO + *drpr⁻/⁻* (n = 25, 11 animals). Freeman-Halton extension of Fisher's exact test.

(F-I) Partial dendritic fields of wildtype (F), *axed* KO (G), *drpr⁻/⁻* (H), and *axed* KO + *drpr⁻/⁻* (I) ddaC neurons 24 hrs AI. (I) shows partial fragmentation. In all image panels, blue asterisks indicate injury sites; yellow dots outline regions covered by injured dendrites. Neurons were labeled by *ppk>CD4-tdTom* (A, B, F and G) and *ppk-CD4-tdTom* (C, D, H, and I). Neuronal-specific KO was induced by *SOP-Cas9* (B, D, G, and I). Scale bars, 25 μm.

(J) Quantification of dendrite fragmentation showing percentages of neurons undergoing no fragmentation, partial fragmentation, and complete fragmentation of injured dendrites at 24 hrs AI. n = number of neurons: wildtype (n=21, 12 animals); *axed* KO (n = 23, 10 animals); *drpr⁻/⁻* (n = 12, 5 animals); *axed* KO + *drpr⁻/⁻* (n = 19, 8 animals). Freeman-Halton extension of Fisher's exact test. For all quantifications, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$; n.s., not significant. The significance level above each genotype is for comparison with the wildtype.

Injured dendrites undergo severe membrane rupture during dendrite fragmentation

To further understand how injury induces dendrite degeneration, we investigated the extent of membrane rupture during dendrite breakdown using a split GFP-based assay. In this membrane rupture assay (Figure 7A), neurons express myristoylated tdTom-GFP(1-10) and the fat body secretes GFP(11)x7 into the hemolymph. GFP(11)x7 will bind myr-tdTom-GFP(1-10) and reconstitute fluorescent GFP only when the dendrite membrane is ruptured to allow diffusion of GFP(11)x7 into the cytoplasm of neurons. While reconstituted GFP was not detected in uninjured wildtype dendrites (Figure S2A) or degenerating dendrites of *Nmnat* KO neurons

Figure S2: Membrane of uninjured dendrites does not rupture.

(A) Uninjured wildtype dendrites lacking reconstituted GFP in the membrane rupture assay. Scale bar, 25 μm.

(Figures 7B, open arrowheads, and 7D-7E), it was observed in severed wildtype dendrites that were undergoing blebbing and fragmentation (Figures 7C, arrowheads, and 7E), as well as on membrane pieces shed from injured dendrites (Figure 7C, arrows, and 7D). In time-lapse movies, GFP signals were visible at low levels in injured branches soon after laser ablation, likely due to GFP(11)$_{x7}$ entry through the injury site. The signals were kept at constantly low levels until injured dendrites fragmented, when GFP signals rapidly increased in large dendrite particles (Video 3). These results suggest that injured dendrites undergo severe membrane rupture during fragmentation. In contrast, *Nmnat* KO neurons experience much milder disruptions of membrane

Figure 7. Injured dendrites undergo severe membrane rupture during dendrite fragmentation

(A) A diagram for the membrane rupture assay. Extracellular GFP(11)$_{x7}$ is separated from intracellular myr-tdTom-GFP(1-10) attached to the inner membrane of dendrites. Fluorescent GFP is reconstituted only when the dendrite membrane is ruptured to allow diffusion of GFP(11)$_{x7}$ into the cytoplasm of neurons.

(B) *Nmnat* KO dendrites (open arrowheads) and debris (open arrows) lacking reconstituted GFP in the membrane rupture assay. Scale bar, 25 μm.

(C) Reconstituted GFP labeling of injured wildtype dendrites (arrowheads) and debris (arrows) in the membrane rupture assay. Scale bar, 25 μm. See also Video 3.

(D) Quantification of GFP-positive debris, showing the percentage of GFP-positive debris area in tdTom-positive debris area. Number of ROIs: wildtype NI (n= 15, 7 animals); *Nmnat* KO NI (n= 10, 6 animals) wildtype 6 hrs AI (n= 10, 4 animals). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method.

(E) Quantification of GFP-positive filament, showing the percentage of GFP-positive filament area in tdTom-positive filament area. Number of ROIs: same as in (D). Kruskal-Wallis (One-way ANOVA on ranks) and Dunn's test, p-values adjusted with the Benjamini-Hochberg method.

For all quantifications, **p≤0.01; ***p≤0.001; n.s., not significant. The significance level above each genotype is for comparison with the control. Black bar, mean; red bar, SD.

integrity, even though they are losing membranes due to the attack of epidermal cells, probably

because their dynamically growing branches are efficient in repairing membrane damages.

Injured dendrites exhibit calcium flashing that is suppressed by Wld^S and *axed* loss

The observed membrane rupture of injured dendrites is consistent with phagocytic attacks of

epidermal cells on PS-exposing dendrites (Figure 7C). To understand potential signaling events

that may lead to dendritic PS exposure and membrane rupture, we examined calcium dynamics

in injured dendrites using long-term time-lapse imaging (Ji and Han, 2020; Poe et al., 2017),

because calcium influx is essential for and correlates with degeneration of damaged axons in

neuronal culture and *in vivo* (George et al., 1995; Vargas et al., 2015; Williams et al., 2014). A

previous study in zebrafish identified an initial calcium influx at the time of axon severing and a

second calcium wave that coincides with axon fragmentation (Vargas et al., 2015). We recorded

calcium dynamics using a membrane-tethered GCaMP6s (myr-GCaMP6s) (Akbergenova et al.,

2018), which detected only occasional local rise of calcium in dendrites of wildtype (Video 4)

and *Nmnat* KO neurons (Video 5). In the injury model, we observed an initial calcium rise

immediately after laser injury in both detached dendrites and the rest of the neurons (Figure 8A,

2s AI). Interestingly, soon after calcium dropped to the baseline level, severed dendrites but not

those connected to the cell body entered a phase of continuous calcium flashes that lasted 1-5

hours (Figures 8A at 1h AI, 8D, and Video 6). Afterward, severed dendrites stayed relatively

quiescent for 1-3 hrs before calcium surged again at the time of dendrite blebbing and

fragmentation (Figures 8A at 6h AI, 8D, and Video 6). All severed dendrites but not those still

attached to cell bodies showed flashing, quiescence, and surge phases after injury, even though

the exact timing and duration of each phase varied from dendrite to dendrite (Figures S3A and

S3D).

It was reported that nanoscale ruptures of the axonal plasma membrane are responsible

for extracellular calcium influx in neuroinflammatory lesions (Witte et al., 2019) and after spinal

cord contusion (Williams et al., 2014). To determine whether the last calcium rise that coincides

with dendrite degeneration could be a result of phagocytosis-induced membrane breakage, we

imaged both calcium dynamics and PS exposure of injured dendrites using the PS sensor

Annexin V-mCardinal (AV-mCard) (Sapar et al., 2018). AV-mCard labeling of dendrites

appeared slightly ahead of the final calcium surge (Figures 8B and 8C, Video 7). Considering

that Annexin V-binding to PS and accumulation on dendrite surface are likely slower than

calcium activation of GCaMP6s, our data support the idea that PS-mediated phagocytosis causes

dendrite membrane rupture and the final calcium surge.

Figure 8: Calcium flashing precedes PS exposure and fragmentation of injured dendrites

(A) A ddaC neuron in a 3rd instar larva expressing myr-GCaMP6s before (BI) and after (2s AI, 1h AI, and 6h AI) laser injury. Blue arrows, uninjured dendrites or cell body; yellow arrowheads, injured dendrites; blue asterisk, injury site. Scale bar, 10 μm. See also Video 6.

(B) Labeling of injured ddaC dendrites expressing GCaMP6s by AV-mCard after (2 h, 4 h, and 5.5 h) injury. Scale bar, 10 μm. See also Video 7.

(C) Kymograph of GCaMP6s signal and AV-mCard binding on an injured dendrite shown in B. Yellow dotted lines, timing of AV-mCard binding; blue dotted lines, timing of the final calcium surge. Scale bars, 3 μm horizontal, 40 min vertical.

(D-F) Kymograph of GCaMP6s signal in wildtype (D), Wld^S-overexpressing (E), and *axed* KO (F) dendrites after injury with 3-min interval. Scale bar, 5 μm horizontal, 60 min vertical. See also Videos 6, 8 and 12.

(G-I) Kymograph of GCaMP6s signal in wildtype (G), Wld^S-overexpressing (H), and *axed* KO (I) dendrites after injury with 2-s interval. See also Videos 9-11 and 13. Scale bars, 5 μm horizontal, 45 sec vertical.

(J) Plot of GCaMP6s brightness in injured dendrites in (D), (E), and (F) over time with 3-m interval. Measurements in injured dendrites start from 0.5-1 hr AI.

(K) Plot of GCaMP6s brightness in injured dendrite in (G), (H), and (I) over time with 2-s interval. Measurements start from 0.5-1 hr AI.

(L) Quantification of average flashing amplitude (GCaMP6s brightness difference between maximum peak and neighboring minimum peak) captured in 2 s-interval time-lapse images. n=number of dendrite arbors: wildtype 0.5-1.5 hrs AI (n= 19, 5 animals); Wld^S-overexpressing 2 hrs AI (n=8, 4 animals); *axed* KO 0.5-1 hrs AI (n=7, 3 animals). One-way ANOVA and Tukey's test. For all quantifications, ***p≤0.001; n.s., not significant. The significance level above each genotype is for comparison with the control. Black bar, mean; red bar, SD.

Because the unique pattern of calcium flashing is absent in uninjured dendrites and

Nmnat KO neurons, we suspected that it may play an active role in promoting degeneration of

injured dendrites. If so, factors that can block dendrite degeneration may also alter the calcium

flashing. Indeed, Wld[S] OE dramatically reduced calcium fluctuations in injured dendrites and

eliminated the quiescent and surge phases for the entire duration of our time-lapse imaging (13

hrs) (Figures 8E, 8J, S3B, and Video 8). In addition, using time-lapse imaging at a higher

temporal resolution (2 s/frame), we found that wildtype injured dendrites displayed calcium

flashes at a frequency of 0.4-3/min (Figures 8G and 8K, and Video 9) while injured dendrites of

Figure S3: Wld[S] overexpression and loss of *axed* changed calcium flashes in injured dendrites.

(A-C) Representative plots of GCaMP6s brightness in injured dendrites of wildtype (A), Wld[S]-overexpressing (B), and *axed* KO (C) neurons over time with 3-min interval.

(D-F) Representative plots of GCaMP6s brightness in uninjured dendrites of wildtype (A), Wld[S]-overexpressing (B), and *axed* KO (C) neurons over time with 3-min interval.

Wld^S OE neurons either maintained a much milder calcium fluctuation or exhibited irregular and

infrequent calcium flashes (Figures 8H and 8K-8L; Videos 10 and 11). In contrast, the majority

of severed dendrites of *axed* KO neurons (6/9) maintained constant and high calcium levels

before the quiescence and surge phases (Figures 8F, 8J, S3C, and Video 12), which was further

confirmed by time-lapse imaging at the high temporal resolution (Figures 8I, 8K, 8L, S3C, and

Video 13). Overall, injured dendrites of Wld^S OE and *axed* KO neurons showed much smaller

average amplitudes of calcium fluctuations compared to the wildtype control (Figure 8L).

Because Wld^S OE and *axed* KO did not alter calcium dynamics in uninjured dendrites (Figures

S3E and S3F), these data suggest that these manipulations specifically modify calcium flashing

patterns in injured dendrites, consistent with the idea that calcium flashes may promote

degeneration of injured dendrites.

DISCUSSION

In this study, we investigate the mechanism of dendrite degenerations caused by genetic NAD^+

disruptions and injury in the *Drosophila* PNS. Although neurite phagocytosis has been observed

after neuronal injury both *in vivo* and *in vitro* (Han et al., 2014; MacDonald et al., 2006;

Nomura-Komoike et al., 2020; Sapar et al., 2018), WD is generally thought as a result of neurite

self-destruction triggered by NAD^+ depletion (Babetto et al., 2013; Gerdts et al., 2015; Gerdts et

al., 2016; Neukomm et al., 2017). However, our results using the engulfment-deficient *drpr*

mutant strongly suggest that PS-mediated phagocytosis is the main driving force for NAD^+

reduction-related dendrite degeneration *in vivo*. It is solely responsible for dendrite degeneration

of *Nmnat* KO neurons and greatly accelerates the degeneration of Sarm^{GOF} OE neurons. In

injury, phagocytosis is responsible for at least half of the dendrite fragmentation by 10 hrs AI

and may contribute more to dendrite breakdown than self-destruction at earlier stages.

Supporting this idea, ectopic PS exposure on injured dendrites is sufficient to revert the blockage

of dendrite fragmentation by Wld^S OE or *Sarm* KO, even though ectopically induced PS

exposure is much lower than natural PS exposure on injured dendrites (Sapar et al., 2018). Our

time-lapse analyses of PS exposure, the final calcium surge, and dendrite rupture also support

that PS-mediated phagocytosis breaks down injury dendrites at the time of dendrite

fragmentation. Therefore, at least in the context of dendrite injury, phagocytosis is the major

factor driving degeneration while self-
destruction acts as a secondary
mechanism to ensure complete
fragmentation.

Figure 9. Three levels of NAD$^+$ reduction elicit distinct responses in dendrites.

See Discussion for details.

NAD$^+$ reduction is known to be
essential for neuronal PS exposure and
neurite self-destruction during WD
(Gerdts et al., 2015; Sapar et al., 2018;
Shacham-Silverberg et al., 2018). How
does the same signaling input coordinate
the two different events? Our results show
that NAD$^+$ disruption controls PS
exposure and neurite self-destruction in
separate steps of WD. In the existing
model, Sarm activation is believed to cause catastrophic NAD$^+$ depletion that is sufficient to
initiate neurite self-destruction (Gerdts et al., 2015; Sasaki et al., 2016). However, we found that
downstream of Sarm activation and before the initiation of self-destruction, neurites first expose
PS to engage in phagocytosis-mediated non-autonomous degeneration. Therefore, in our revised
model, dendrites respond to at least three distinct, increasingly severe levels of NAD$^+$ reduction
by eliciting different molecular events (Figure 9). Between the NAD$^+$ level required for Sarm
activation (SA level) and the level that initiates self-destruction (SD level), Sarm activity lowers
NAD$^+$ to a level that causes neurons to expose PS on their surface (which we call the PSE level).
This PS exposure is sufficient to cause phagocytosis-mediated dendrite degeneration, which can
be completely prevented by blocking engulfment activity of phagocytes. However, below the SD
level, neurites spontaneously fragment even in the absence of phagocytosis.

Our results also suggest a direct correlation between the kinetics of NAD$^+$ reduction and
the severity of neurite degeneration. *Nmnat* KO is expected to cause slow NAD$^+$ reduction, due
to gene perdurance and the time required for natural NAD$^+$ turnover, and correspondingly causes
engulfment-dependent dendrite degeneration only in late 3rd instar larvae. In contrast, Sarm^{GOF}
OE should lead to a more rapid NAD$^+$ depletion and in fact causes engulfment-dependent
dendrite degeneration as early as the 1st instar and dendrite self-destruction by the 3rd instar.

Injury is known to cause even more rapid NAD^+ reduction in axons (Wang et al., 2005) and is

correlated with the fastest dendrite degeneration – initiation at around 4 hrs AI and completion

usually by 10 hrs AI.

How does NAD^+ reduction cause PS exposure? A direct consequence of NAD^+ loss is the

decline of neurite ATP levels due to the requirement of NAD^+ in glycolysis and oxidative

phosphorylation (Shacham-Silverberg et al., 2018; Wang et al., 2005). Consistent with ATP

reduction playing a role in inducing PS exposure, suppressing mitochondria ATP synthesis in

DRG culture caused gradual axonal PS exposure (Shacham-Silverberg et al., 2018). However,

how ATP reduction may induce PS exposure remain elusive. Although the maintenance of

membrane PS asymmetry by flippases requires ATP, flippase KO in C4da neurons causes a

much milder PS exposure than injury (Sapar et al., 2018), suggesting that mechanisms other than

flippase inhibition must be contributing to the rapid PS exposure seen after injury. Identifying

the PS transporters responsible for PS exposure on injured neurites will be a key step for

revealing the mechanisms of NAD^+ regulation of PS exposure.

Genetic analyses in *Drosophila* identified Axed as a key switch of WD, whose activity is

absolutely required for axon degeneration caused by injury and genetic depletion of NAD^+

(Neukomm et al., 2017). How does Axed regulate neurite degeneration? Our data suggest that

Axed is not required for PS-mediated phagocytosis but contributes to self-destruction of injured

dendrites, placing its activation below the SD level of NAD^+ (Figure 9). Surprisingly, Axed

seems to play a minor role in dendrite degeneration, as its LOF only slowed down but did not

block self-destruction, indicating the existence of other factors that promote self-destruction of

injured dendrites. Importantly, these results suggest that dendrites and axons may rely on

different mechanisms of self-destruction.

By exploring the different mechanisms employed in *Nmnat* KO- and injury-induced

dendrite degenerations, we discovered dynamic calcium activities only present in injured

dendrites, including an unreported calcium flashing pattern prior to any obvious degenerative

event and a final calcium surge that coincides with dendrite fragmentation. Calcium surge at the

time of neurite fragmentation is a shared feature between injured axons of zebrafish (Vargas et

al., 2015) and injured dendrites of *Drosophila* da neurons. Although calcium influx is required

for WD (George et al., 1995) and may activate calcium-dependent lipid scramblases (Suzuki et

al., 2010), our time-lapse analyses suggest that the final calcium surge is more likely a result of

phagocytosis-induced membrane rupture rather than the cause of fragmentation. In comparison,

the calcium flashing soon after the injury is unique to dendrites and may play an active role in

dendrite degeneration in ways similar to the compartmentalized calcium flashing that occurs

during developmental pruning of C4da neurons (Kanamori et al., 2013). Consistent with this

possibility, the calcium flashing is suppressed by Wld^S OE and *axed* KO. Interestingly, these two

manipulations block the calcium flashing in opposite ways, with Wld^S OE dampening the

calcium level and *axed* KO keeping the calcium level high. This distinction may be a useful clue

for understanding the regulation of PS exposure and self-destruction. For example, it is possible

that elevated calcium levels prepare dendrites for PS exposure and drastic changes of calcium

levels promote dendrite self-destruction.

Lastly, as impairment of NAD$^+$ metabolism is a general feature of neurodegenerative

disorders including Leber congenital amaurosis (LCA), Alzheimer's disease, Parkinson's

disease, and retinal degenerations (Ali et al., 2013; Canto et al., 2015; Fang et al., 2017; Lin et

al., 2016; Verdin, 2015), phagocytosis may play important roles in the pathogenesis of these

diseases through dysregulated neuronal PS exposure.

METHODS

Fly strains

The details of fly strains used in this study are listed in the key reagent table. For labeling of

C4da neurons, we used *ppk-CD4-tdTom*, *ppk-MApHS*, and *ppk-Gal4 UAS-CD4-tdTom*. For

labeling PS exposure on dendrites, we used *dcg-Gal4 UAS-GFP-LactC1C2*, *R16A03-LexA*

LexAop-GFP-LactC1C2, and *dcg-Gal4 UAS-AnnexinV-mCard*. To label Or22a axons, we used

Or22a-Gal4 UAS-mCD8-GFP. To visualize dendrite rupture, we used *dcg-Gal4 UAS-*

sGFP(11)x7 together with *ppk-LexA LexAop-myr-tdTom-GFP(1-10)*. To visualize calcium

activities in C4da dendrites, we used *ppk-LexA LexAop-myr-GCaMP6s*.

Molecular cloning and transgenic flies

ey-Cas9: Three tandem copies of a 211bp *ey* enhancer (corresponding to nucleotides 2577-2787

in GenBank accession number AJ131630) was inserted into XhoI/SalI sites of pENTR11

(Thermo Fisher Scientific). The resulting entry vector was combined with a Cas9 destination

vector which is similar to pDEST-APIC-Cas9 (Addgene 121657) but does not contain Inr, MTE, and DPE in the Hsp70 core promoter (Poe et al., 2019) to generate the pAPIC-ey-Cas9 expression vector through a Gateway LR reaction.

LexAop-myr-tdTom-GFP(1-10): pAPLO-CD4-tdTom (Poe et al., 2017) was digested by BglII/AscI, blunted, and religated to remove an XbaI site before 13xLexAop2. The resulting construct is called pAPLOm-CD4-tdTom. An myr-GFP fragment was isolated from pJFRC176-10XUAS-rox-dSTOP-rox-myr-GFP (Addgene 32147) by XhoI/XbaI digestion and ligated to XhoI/XbaI sites of pAPLOm-CD4-tdTom to make pAPLO-myr-GFP. pAPLO-myr-GFP was then digested by BamHI/XbaI and subsequently assembled with a tdTom PCR fragment and a GFP(1-10) gBlock fragment (synthesized by IDT) through NEBuilder DNA Assembly to generate pAPLO-myr-tdTom-GFP1-10.

UAS-sGFP(11)x7: A DNA fragment containing GFP(11)x7 was PCR amplified from a gBlock DNA fragment and cloned into Nhe/XbaI sites of pIHEU-sfGFP-LactC1C2 using NEBuilder DNA Assembly. The resulting construct pIHEU-sGFP11x7 inherits the signal peptide sequence from pIHEU-sfGFP-LactC1C2 but removes the sfGFP-LactC1C2 coding sequence.

LexAop-WldS: The WldS coding sequence was PCR amplified from UAS-WldS genomic DNA and cloned into XhoI/XbaI sites of pAPLOm-CD4-tdTom via restriction cloning to make pAPLO-WldS.

UAS-ATP8A: A fragment encoding a FLAG tag and the ER exit signal of Kir2.1 (Han et al., 2011) was cloned into the EcoRI and XbaI sites of pACU (Han et al., 2011) by ligating with annealed oligos. The ATP8A coding region that is shared by all isoforms was amplified by PCR from cDNA clone GH28327 (AA26 to AA1091, BDGP) and inserted into an NheI site before the FLAG tag.

gRNA expression vectors: For *Nmnat*, *Sarm*, and *axed*, gRNAs were cloned into pAC-U63-tgRNA-Rev as described (Poe et al., 2019). The first tRNA spacer in each final construct is tRNA[Gly] but the remaining tRNA spacers are tRNA[Gln] (Koreman et al., 2021). *gRNA-peb* was constructed in pAC-U63-QtgRNA2.1-BR. The gRNA target sequences are listed in the table below.

Gene	Target sequence 1	Target sequence 2

Sarm	GCATCTGTTCAAACACTCCG	CGATTCCAATATCAGCCCGG
Nmnat	GGAACCCACAGAGTGGTAGG	TAAGAGCCGCCGAATCAACG
axed	GTCGGTACTCGAGCGGAGCG	TGGACCCGATGGCCATCACG
peb	ATTTCGTCTGAATCGCTCGG	ACGCATGTGACGCACCAGGG

493   The above constructs were injected by Rainbow Transgenic Flies to transform flies
494   through φC31 integrase-mediated integration into attP docker sites. The sequences of constructs
495   will be provided upon request.

CRISPR-TRiM

The efficiency of transgenic gRNA lines was validated by the Cas9-LEThAL assay (Poe et al.,
2019). Homozygous males of each gRNA line were crossed to *Act-Cas9 w lig4* homozygous
females. *gRNA-Nmnat* crosses caused lethality of all progeny at the 2^{nd} instar larval stage; *gRNA-*
Sarm crosses yielded viable female progeny and male lethality between 3^{rd} instar larvae to
prepupae; *gRNA-axed* crosses caused lethality at the embryonic stage for all progeny; *gRNA-peb*
crosses caused lethality from embryonic stage to the 2^{nd} instar for all progeny. These results
suggest that all gRNAs are efficient.

C4da-specific gene knockout was carried out using *ppk-Cas9* (Poe et al., 2019). Tissue-
specific knockout in da neuron precursor cells were carried out with *SOP-Cas9* (Poe et al.,
2019). Gene knockout in the precursor cells of Or22a ORNs was carried out using *ey-Cas9* (this
study).

Live imaging

Animals were reared at 25°C in density-controlled vials (60-100 embryos/vial) on standard
yeast-glucose medium (doi:10.1101/pdb.rec10907). Larvae at 125 hours AEL (wandering stage)
or stages specified were mounted in 100% glycerol under coverslips with vacuum grease spacer
and imaged using a Leica SP8 microscope equipped with a 40X NA1.30 oil objective. Larvae
were lightly anesthetized with isoflurane before mounting. For consistency, we imaged dorsal
ddaC neurons from A1-A3 segments (2-3 neurons per animal) on one side of the larvae. Unless
stated otherwise, confocal images shown in all figures are maximum intensity projections of z

stacks encompassing the epidermal layer and the sensory neurons beneath, which are typically 8–

10 µm for 3rd instar larvae.

Injury assay

Injury assay at the larval stage was done as described previously (Sapar et al., 2018). Briefly,

larvae at 90 hr AEL were lightly anesthetized with isoflurane, mounted in a small amount of

halocarbon oil under coverslips with grease spacers. The laser ablation was performed on a Zeiss

LSM880 Confocal/Multiphoton Upright Microscope, using a 790 nm two-photon laser at

primary dendrites of ddaC neurons in A1 and A3 segments. Animals were recovered on grape

juice agar plates following lesion for appropriate times before imaging.

ORN axon injury assay was performed on 7-day-old male flies by removing the outer

segments of both antennae as described in (MacDonald et al., 2006). The injured males were

recovered for 7 days by transferring to fresh yeast-glucose medium every day. To examine

Or22a axon degeneration, brains were dissected in PBST (0.2% Triton-X in PBS), fixed in 4%

formaldehyde in PBS for 20 min, and then rinsed with PBST three times, 20 minutes each. Then

the brains were mounted in SlowFade® Diamond Antifade Mountant (Thermo Fisher Scientific)

and imaged using a Leica SP8 microscope with a 40x NA1.3 oil objective.

Long-term time-lapse imaging

Long-term time-lapse imaging at the larval stage was done as described previously (Ji and Han,

2020; Sapar et al., 2018). Briefly, a layer of double-sided tape was placed on the coverslip to

define the position of PDMS blocks. A small amount of UV glue was added to the groove of

PDMS and to the coverslip. Anesthetized larvae were placed on top of the UV glue on the

coverslip and then covered by PDMS blocks with the groove side contacting the larva. Glue was

then cured by 365nm UV light. The coverslip with attached PDMS and larvae was mounted on

an aluminum slide chamber that contained a piece of moisturized Kimwipes (Kimtech Science)

paper. Time-lapse imaging was performed on a Leica SP8 confocal equipped with a 40x NA1.3

oil objective and a resonant scanner at digital zoom 0.75 and a 3-min or 2-sec interval. For

imaging after ablation, larvae were pre-mounted in the imaging chamber and subjected to laser

injury. The larvae were then imaged 0.5-1 hours after ablation. For calcium imaging before and

immediately after ablation, images were captured on a Zeiss LSM880 Confocal/Multiphoton

Upright Microscope on which the ablation was performed.

Image analysis and quantification

Image processing and analyses were done in Fiji/ImageJ. Methods for tracing and measuring C4da neuron dendrite length have been previously described (Poe et al., 2017). Briefly, the images were segmented by Auto Local Threshold and reduced to single pixel skeletons before measurement of skeleton length by pixel distance. The dendrite debris measurement has been described previously (Sapar et al., 2018). Briefly, a dendrite mask was first generated from projected images by Auto Local Threshold in order to create a region of interest (ROI) by dilation to map areas within one-epidermal-cell diameter (40 μm) from dendrites. Dendrite debris within the ROI was converted to binary masks based on fixed thresholds. Different thresholds were used for *ppk-C4-tdTom* and *ppk-Gal4 UAS-CD-tdTom* as they have different brightness. The dendrite pixel area (ADen), debris pixel area (ADeb), and ROI area (AROI) were measured and dendrite coverage ratio was calculated based on the following formula: $100 \cdot Adeb \cdot AROI /(AROI- ADen) \cdot ADen$. For measuring Lact-GFP, two regions at empty epidermal regions were measured as background levels. TdTom signals on dendrites were used to generate dendrite masks for measurement of GFP within the masks. For kymographs, we used a custom macro based on the Straighten function to extract a strip of pixels centered at the selected dendrite branch. The maximum intensity pixel in the strip at each distance was used to generate a single-pixel line for each time frame. The final kymographs were displayed using the Fire lookup table (LUT).

Statistical Analysis

R was used to conduct statistical analyses and generate graphs. (*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$). Statistical significance was set at $p < 0.05$. Data acquisition and quantification were performed non-blinded. Acquisition was performed in ImageJ (batch processing for debris coverage ratio and fragmentation ratio, manually by hand for GFP-Lact binding) and Microsoft Excel. Statistical analyses were performed using R. We used the following R packages: car, stats, multcomp for statistical analysis and ggplot2 for generating graphs. Some graphs were made in Excel using its native plotting functions. For the statistical analysis we ran the following tests, ANOVA (followed by Tukey's HSD) when dependent variable was normally distributed and there was approximately equal variance across groups. When dependent variable was not normally distributed and variance was not equal across groups, we used Kruskal-Wallis

(followed by Dunn's test, p-values adjusted with Benjamini-Hochberg method) to test the null

hypothesis that assumes that the samples (groups) are from identical populations. To check

whether the data fit a normal distribution, we generated qqPlots to analyze whether the residuals

of the linear regression model is normally distributed. We used the Levene's test to check for

equal variance within groups. The state of neuronal degeneration or fragmentation was compared

using the Freeman-Halton extension of Fisher's exact test.

Replication

For all larval and adult imaging experiments, at least 3 biological replications were performed

for each genotype and/or condition.

AUTHOR CONTRIBUTIONS

Conceptualization, CH, HJ, MLS, AS; Methodology, CH, HJ, MLS, AS, BW; Investigation, HJ,

MLS, AS; Formal Analysis, HJ, MLS, AS; Resources, CH, MLS, HJ, BW, AS; Writing –

Original Draft, CH, MLS, HJ, AS; Writing – Review and Editing, CH, MLS, HJ, AS; Funding

Acquisition, CH.

ACKNOWLEDGMENTS

We thank Marc Freeman, Yang Xiang, Heather Broihier, and Bloomington Stock Center for fly

stocks; Addgene for plasmids; Cornell BRC Imaging facility for access to microscopes (funded

by NIH grant S10OD018516); Cornell CSCU for advice on statistics; Mike Goldberg and

Fenghua Hu for critical reading and suggestions on the manuscript. This work was supported by

a Cornell start-up fund and NIH grants (R01NS099125 and R21OD023824) awarded to C.H.

DECLARATION OF INTEREST

The authors declare no competing interests.

REFERENCE

Akbergenova, Y., Cunningham, K.L., Zhang, Y.V., Weiss, S., and Littleton, J.T. (2018).
Characterization of developmental and molecular factors underlying release heterogeneity at
Drosophila synapses. Elife *7*.
Ali, Y.O., Li-Kroeger, D., Bellen, H.J., Zhai, R.G., and Lu, H.C. (2013). NMNATs,
evolutionarily conserved neuronal maintenance factors. Trends Neurosci *36*, 632-640.
Avery, M.A., Sheehan, A.E., Kerr, K.S., Wang, J., and Freeman, M.R. (2009). Wld S requires
Nmnat1 enzymatic activity and N16-VCP interactions to suppress Wallerian degeneration. J Cell
Biol *184*, 501-513.

Babetto, E., Beirowski, B., Russler, E.V., Milbrandt, J., and DiAntonio, A. (2013). The Phr1
ubiquitin ligase promotes injury-induced axon self-destruction. Cell Rep *3*, 1422-1429.
Bratkowski, M., Xie, T., Thayer, D.A., Lad, S., Mathur, P., Yang, Y.S., Danko, G., Burdett,
T.C., Danao, J., Cantor, A*., et al.* (2020). Structural and Mechanistic Regulation of the Pro-
degenerative NAD Hydrolase SARM1. Cell Rep *32*, 107999.
Canto, C., Menzies, K.J., and Auwerx, J. (2015). NAD(+) Metabolism and the Control of Energy
Homeostasis: A Balancing Act between Mitochondria and the Nucleus. Cell Metab *22*, 31-53.
Coleman, M.P., and Freeman, M.R. (2010). Wallerian degeneration, wld(s), and nmnat. Annu
Rev Neurosci *33*, 245-267.
Davies, A.J., Kim, H.W., Gonzalez-Cano, R., Choi, J., Back, S.K., Roh, S.E., Johnson, E.,
Gabriac, M., Kim, M.S., Lee, J*., et al.* (2019). Natural Killer Cells Degenerate Intact Sensory
Afferents following Nerve Injury. Cell *176*, 716-728 e718.
Di Stefano, M., Nascimento-Ferreira, I., Orsomando, G., Mori, V., Gilley, J., Brown, R.,
Janeckova, L., Vargas, M.E., Worrell, L.A., Loreto, A*., et al.* (2015). A rise in NAD precursor
nicotinamide mononucleotide (NMN) after injury promotes axon degeneration. Cell Death Differ
22, 731-742.
Fang, E.F., Lautrup, S., Hou, Y., Demarest, T.G., Croteau, D.L., Mattson, M.P., and Bohr, V.A.
(2017). NAD(+) in Aging: Molecular Mechanisms and Translational Implications. Trends Mol
Med *23*, 899-916.
Farley, J.E., Burdett, T.C., Barria, R., Neukomm, L.J., Kenna, K.P., Landers, J.E., and Freeman,
M.R. (2018). Transcription factor Pebbled/RREB1 regulates injury-induced axon degeneration.
Proc Natl Acad Sci U S A *115*, 1358-1363.
Figley, M.D., Gu, W., Nanson, J.D., Shi, Y., Sasaki, Y., Cunnea, K., Malde, A.K., Jia, X., Luo,
Z., Saikot, F.K*., et al.* (2021). SARM1 is a metabolic sensor activated by an increased
NMN/NAD(+) ratio to trigger axon degeneration. Neuron *109*, 1118-1136 e1111.
Fourgeaud, L., Traves, P.G., Tufail, Y., Leal-Bailey, H., Lew, E.D., Burrola, P.G., Callaway, P.,
Zagorska, A., Rothlin, C.V., Nimmerjahn, A*., et al.* (2016). TAM receptors regulate multiple
features of microglial physiology. Nature *532*, 240-244.
Freeman, M.R., Delrow, J., Kim, J., Johnson, E., and Doe, C.Q. (2003). Unwrapping glial
biology: Gcm target genes regulating glial development, diversification, and function. Neuron
38, 567-580.
Galloway, D.A., Phillips, A.E.M., Owen, D.R.J., and Moore, C.S. (2019). Phagocytosis in the
Brain: Homeostasis and Disease. Front Immunol *10*, 790.
George, E., Glass, J., and Griffin, J. (1995). Axotomy-induced axonal degeneration is mediated
by calcium influx through ion-specific channels. The Journal of Neuroscience *15*, 6445-6452.
Gerdts, J., Brace, E.J., Sasaki, Y., DiAntonio, A., and Milbrandt, J. (2015). SARM1 activation
triggers axon degeneration locally via NAD(+) destruction. Science *348*, 453-457.
Gerdts, J., Summers, D.W., Milbrandt, J., and DiAntonio, A. (2016). Axon Self-Destruction:
New Links among SARM1, MAPKs, and NAD+ Metabolism. Neuron *89*, 449-460.
Han, C., Jan, L.Y., and Jan, Y.N. (2011). Enhancer-driven membrane markers for analysis of
nonautonomous mechanisms reveal neuron-glia interactions in Drosophila. Proc Natl Acad Sci U
S A *108*, 9673-9678.
Han, C., Song, Y., Xiao, H., Wang, D., Franc, N.C., Jan, L.Y., and Jan, Y.N. (2014). Epidermal
cells are the primary phagocytes in the fragmentation and clearance of degenerating dendrites in
Drosophila. Neuron *81*, 544-560.

Ji, H., and Han, C. (2020). LarvaSPA, A Method for Mounting Drosophila Larva for Long-Term
Time-Lapse Imaging. J Vis Exp.
Jiang, Y., Liu, T., Lee, C.H., Chang, Q., Yang, J., and Zhang, Z. (2020). The NAD(+)-mediated
self-inhibition mechanism of pro-neurodegenerative SARM1. Nature *588*, 658-663.
Kanamori, T., Kanai, M.I., Dairyo, Y., Yasunaga, K., Morikawa, R.K., and Emoto, K. (2013).
Compartmentalized calcium transients trigger dendrite pruning in Drosophila sensory neurons.
Science *340*, 1475-1478.
Koreman, G.T., Xu, Y., Hu, Q., Zhang, Z., Allen, S.E., Wolfner, M.F., Wang, B., and Han, C.
(2021). Upgraded CRISPR/Cas9 tools for tissue-specific mutagenesis in Drosophila. Proc Natl
Acad Sci U S A *118*.
Leventis, P.A., and Grinstein, S. (2010). The distribution and function of phosphatidylserine in
cellular membranes. Annu Rev Biophys *39*, 407-427.
Lin, J.B., Kubota, S., Ban, N., Yoshida, M., Santeford, A., Sene, A., Nakamura, R., Zapata, N.,
Kubota, M., Tsubota, K.*, et al.* (2016). NAMPT-Mediated NAD(+) Biosynthesis Is Essential for
Vision In Mice. Cell Rep *17*, 69-85.
Liu, H.W., Smith, C.B., Schmidt, M.S., Cambronne, X.A., Cohen, M.S., Migaud, M.E., Brenner,
C., and Goodman, R.H. (2018). Pharmacological bypass of NAD(+) salvage pathway protects
neurons from chemotherapy-induced degeneration. Proc Natl Acad Sci U S A.
MacDonald, J.M., Beach, M.G., Porpiglia, E., Sheehan, A.E., Watts, R.J., and Freeman, M.R.
(2006). The Drosophila cell corpse engulfment receptor Draper mediates glial clearance of
severed axons. Neuron *50*, 869-881.
Mack, T.G., Reiner, M., Beirowski, B., Mi, W., Emanuelli, M., Wagner, D., Thomson, D.,
Gillingwater, T., Court, F., Conforti, L.*, et al.* (2001). Wallerian degeneration of injured axons
and synapses is delayed by a Ube4b/Nmnat chimeric gene. Nat Neurosci *4*, 1199-1206.
Mazaheri, F., Breus, O., Durdu, S., Haas, P., Wittbrodt, J., Gilmour, D., and Peri, F. (2014).
Distinct roles for BAI1 and TIM-4 in the engulfment of dying neurons by microglia. Nat
Commun *5*, 4046.
Miller, B.R., Press, C., Daniels, R.W., Sasaki, Y., Milbrandt, J., and DiAntonio, A. (2009). A
dual leucine kinase-dependent axon self-destruction program promotes Wallerian degeneration.
Nat Neurosci *12*, 387-389.
Nandrot, E.F., Anand, M., Almeida, D., Atabai, K., Sheppard, D., and Finnemann, S.C. (2007).
Essential role for MFG-E8 as ligand for alphavbeta5 integrin in diurnal retinal phagocytosis.
Proc Natl Acad Sci U S A *104*, 12005-12010.
Neukomm, L.J., Burdett, T.C., Seeds, A.M., Hampel, S., Coutinho-Budd, J.C., Farley, J.E.,
Wong, J., Karadeniz, Y.B., Osterloh, J.M., Sheehan, A.E.*, et al.* (2017). Axon Death Pathways
Converge on Axundead to Promote Functional and Structural Axon Disassembly. Neuron *95*, 78-
91 e75.
Nomura-Komoike, K., Saitoh, F., and Fujieda, H. (2020). Phosphatidylserine recognition and
Rac1 activation are required for Muller glia proliferation, gliosis and phagocytosis after retinal
injury. Sci Rep *10*, 1488.
Osterloh, J.M., Yang, J., Rooney, T.M., Fox, A.N., Adalbert, R., Powell, E.H., Sheehan, A.E.,
Avery, M.A., Hackett, R., Logan, M.A.*, et al.* (2012). dSarm/Sarm1 is required for activation of
an injury-induced axon death pathway. Science *337*, 481-484.
Poe, A.R., Tang, L., Wang, B., Li, Y., Sapar, M.L., and Han, C. (2017). Dendritic space-filling
requires a neuronal type-specific extracellular permissive signal in Drosophila. Proc Natl Acad
Sci U S A *114*, E8062-E8071.

Poe, A.R., Wang, B., Sapar, M.L., Ji, H., Li, K., Onabajo, T., Fazliyeva, R., Gibbs, M., Qiu, Y.,
Hu, Y*., et al.* (2019). Robust CRISPR/Cas9-Mediated Tissue-Specific Mutagenesis Reveals
Gene Redundancy and Perdurance in Drosophila. Genetics *211*, 459-472.
Raiders, S., Black, E.C., Bae, A., MacFarlane, S., Klein, M., Shaham, S., and Singhvi, A. (2021).
Glia actively sculpt sensory neurons by controlled phagocytosis to tune animal behavior. Elife
10.
Sapar, M.L., and Han, C. (2019). Die in pieces: How Drosophila sheds light on neurite
degeneration and clearance. J Genet Genomics *46*, 187-199.
Sapar, M.L., Ji, H., Wang, B., Poe, A.R., Dubey, K., Ren, X., Ni, J.Q., and Han, C. (2018).
Phosphatidylserine Externalization Results from and Causes Neurite Degeneration in Drosophila.
Cell Rep *24*, 2273-2286.
Sasaki, Y., Nakagawa, T., Mao, X., DiAntonio, A., and Milbrandt, J. (2016). NMNAT1 inhibits
axon degeneration via blockade of SARM1-mediated NAD(+) depletion. Elife *5*.
Shacham-Silverberg, V., Sar Shalom, H., Goldner, R., Golan-Vaishenker, Y., Gurwicz, N.,
Gokhman, I., and Yaron, A. (2018). Phosphatidylserine is a marker for axonal debris engulfment
but its exposure can be decoupled from degeneration. Cell Death Dis *9*, 1116.
Shen, C., Vohra, M., Zhang, P., Mao, X., Figley, M.D., Zhu, J., Sasaki, Y., Wu, H., DiAntonio,
A., and Milbrandt, J. (2021). Multiple domain interfaces mediate SARM1 autoinhibition. Proc
Natl Acad Sci U S A *118*.
Sievers, C., Platt, N., Perry, V.H., Coleman, M.P., and Conforti, L. (2003). Neurites undergoing
Wallerian degeneration show an apoptotic-like process with Annexin V positive staining and loss
of mitochondrial membrane potential. Neurosci Res *46*, 161-169.
Suzuki, J., Umeda, M., Sims, P.J., and Nagata, S. (2010). Calcium-dependent phospholipid
scrambling by TMEM16F. Nature *468*, 834-838.
Tanaka, K., Fujimura-Kamada, K., and Yamamoto, T. (2011). Functions of phospholipid
flippases. J Biochem *149*, 131-143.
Tao, J., and Rolls, M.M. (2011). Dendrites have a rapid program of injury-induced degeneration
that is molecularly distinct from developmental pruning. J Neurosci *31*, 5398-5405.
Tung, T.T., Nagaosa, K., Fujita, Y., Kita, A., Mori, H., Okada, R., Nonaka, S., and Nakanishi, Y.
(2013). Phosphatidylserine recognition and induction of apoptotic cell clearance by Drosophila
engulfment receptor Draper. J Biochem *153*, 483-491.
Vargas, M.E., Yamagishi, Y., Tessier-Lavigne, M., and Sagasti, A. (2015). Live Imaging of
Calcium Dynamics during Axon Degeneration Reveals Two Functionally Distinct Phases of
Calcium Influx. J Neurosci *35*, 15026-15038.
Verdin, E. (2015). NAD(+) in aging, metabolism, and neurodegeneration. Science *350*, 1208-
1213.
Waller, A. (1850). Experiments on the Section of the Glossopharyngeal and Hypoglossal Nerves
of the Frog, and Observations of the Alterations Produced Thereby in the Structure of Their
Primitive Fibres. Philosophical Transactions of the Royal Society of London *140*, 423-429.
Wang, J., Zhai, Q., Chen, Y., Lin, E., Gu, W., McBurney, M.W., and He, Z. (2005). A local
mechanism mediates NAD-dependent protection of axon degeneration. J Cell Biol *170*, 349-355.
Wen, Y., Parrish, J.Z., He, R., Zhai, R.G., and Kim, M.D. (2011). Nmnat exerts neuroprotective
effects in dendrites and axons. Mol Cell Neurosci *48*, 1-8.
Williams, P.R., Marincu, B.N., Sorbara, C.D., Mahler, C.F., Schumacher, A.M., Griesbeck, O.,
Kerschensteiner, M., and Misgeld, T. (2014). A recoverable state of axon injury persists for
hours after spinal cord contusion in vivo. Nat Commun *5*, 5683.

Witte, M.E., Schumacher, A.M., Mahler, C.F., Bewersdorf, J.P., Lehmitz, J., Scheiter, A.,
Sanchez, P., Williams, P.R., Griesbeck, O., Naumann, R*., et al.* (2019). Calcium Influx through
Plasma-Membrane Nanoruptures Drives Axon Degeneration in a Model of Multiple Sclerosis.
Neuron *101*, 615-624 e615.
Xiong, X., Hao, Y., Sun, K., Li, J., Li, X., Mishra, B., Soppina, P., Wu, C., Hume, R.I., and
Collins, C.A. (2012). The Highwire ubiquitin ligase promotes axonal degeneration by tuning
levels of Nmnat protein. PLoS Biol *10*, e1001440.
Yang, J., Wu, Z., Renier, N., Simon, D.J., Uryu, K., Park, D.S., Greer, P.A., Tournier, C., Davis,
R.J., and Tessier-Lavigne, M. (2015). Pathological axonal death through a MAPK cascade that
triggers a local energy deficit. Cell *160*, 161-176.
Zhai, R.G., Cao, Y., Hiesinger, P.R., Zhou, Y., Mehta, S.Q., Schulze, K.L., Verstreken, P., and
Bellen, H.J. (2006). Drosophila NMNAT maintains neural integrity independent of its NAD
synthesis activity. PLoS Biol *4*, e416.
Zhai, R.G., Rizzi, M., and Garavaglia, S. (2009). Nicotinamide/nicotinic acid mononucleotide
adenylyltransferase, new insights into an ancient enzyme. Cell Mol Life Sci *66*, 2805-2818.
759

VIDEO LEGEND

Video 1. GFP-Lact labeling of degenerating *Nmnat* KO dendrites, related to Figure 2.

Time-lapse movie of GFP-Lact labeling on distal dendrites of a *Nmnat* KO C4da neuron before

and during degeneration. Imaging started around 96 hours AEL. *Dcg-Gal4* drives GFP-Lact

expression in both fat bodies (not shown) and hemocytes (mobile cells in the GFP-Lact channel).

Timestamp is relative to the first frame, with a 3-min interval between each frame.

Video 2. Dendrite dynamics of *Nmnat* KO neurons in *drpr* mutant larvae, related to Figure

2.

Time-lapse movie of dendrites of a *Nmnat* KO neuron exhibiting dynamic extension and

retraction behaviors in a *drpr* mutant larva. Imaging started around 120 hours AEL. Timestamp

is relative to the first frame, with a 3-min interval between each frame.

Video 3. Membrane rupture of injured dendrites, related to Figure 7.

Time-lapse movie of laser-injured C4da dendrites from 1 to 11 hours AI. Reconstituted GFP is

from extracellular GFP(11)$_{x7}$ entering the dendrites and myr-tdTom-GFP(1-10) attached to the

inner leaflet of dendritic membrane. Timestamp is relative to the first frame, with a 3-min

interval between each frame.

Video 4. Cytoplasmic calcium dynamics in uninjured wildtype dendrites, related to Figure

8.

Time-lapse movie of an uninjured wildtype C4da neuron showing low baseline GCaMP6s
signals and occasional local rises in dendrites. Timestamp is relative to the first frame, with a 3-
min interval between each frame.

Video 5. Cytoplasmic calcium dynamics in dendrites of *Nmnat* KO neurons, related to
Figure 8.

Time-lapse movie of a *Nmnat* KO C4da neuron showing calcium dynamics in uninjured
dendrites. Timestamp is relative to the first frame, with a 3-min interval between each frame.

Video 6. Calcium dynamics in injured wildtype dendrites, related to Figure 8.

Time-lapse movie of laser-injured C4da dendrites from 1 to 11 hrs AI showing calcium
dynamics in both severed dendrites and those attached to the soma. Timestamp is relative to the
first frame, with a 3-min interval between each frame.

Video 7. AV-mCard labeling and calcium dynamics of injured dendrites, related to Figure
8.

Time-lapse movie of laser-injured C4da dendrites from 1 to 6 hrs AI showing labeling of injured
dendrites by the PS sensor AV-mCard and GCaMP6s signals. Timestamp is relative to the first
frame, with a 3-min interval between each frame.

Video 8. Calcium dynamics in injured Wld^S OE dendrites, related to Figure 8.

Time-lapse movie of laser-injured Wld^S OE C4da dendrites from 1 to 13 hours AI showing
GCaMP6s signals in severed dendrites. Timestamp is relative to the first frame, with a 3-min
interval between each frame.

Video 9. Calcium dynamics in injured wildtype dendrites with a higher temporal
resolution, related to Figure 8.

High temporal-resolution time-lapse movie of laser-injured wildtype C4da dendrites around 1 hr
AI showing calcium dynamics. Timestamp is relative to the first frame, with a 2-sec interval
between each frame.

Video 10. Calcium dynamics in injured Wld^S OE dendrites at a higher temporal resolution,
related to Figure 8.

High temporal-resolution time-lapse movie of laser-injured Wld^S OE C4da dendrites around 2

hrs AI showing calcium dynamics. Timestamp is relative to the first frame, with a 2-sec interval

between each frame.

Video 11. Irregular and infrequent cytoplasmic calcium flashes in injured Wld^S OE

dendrites at a higher temporal resolution, related to Figure 8.

High temporal-resolution time-lapse movie of laser-injured Wld^S OE C4da dendrites around 2

hrs AI showing irregular and infrequent GCaMP6s signals in severed dendrites. Timestamp is

relative to the first frame, with a 2-sec interval between each frame.

Video 12. Calcium dynamics in injured *axed* KO dendrites, related to Figure 8.

Time-lapse movie of laser-injured *axed* KO C4da dendrites from 0.5 to 5.5 hours AI showing

GCaMP6s signals in severed dendrites. Timestamp is relative to the first frame, with a 3-min

interval between each frame.

Video 13. Calcium dynamics in injured *axed* KO dendrites at a higher temporal resolution,

related to Figure 8.

High temporal-resolution time-lapse movie of laser-injured *axed* KO C4da dendrites around 10

min AI showing calcium dynamics. Timestamp is relative to the first frame, with a 2-sec interval

between each frame.