Coupling between fast and slow oscillator circuits in *Cancer borealis* is temperature compensated

Powell, D.J.*[1] dpowell@bowdoin.edu https://orcid.org/0000-0002-9210-2201

Haddad, S.A.*[2] sara.haddad@mls.uzh.ch https://orcid.org/0000-0003-0807-0823

Gorur-Shandilya, S.* srinivasgs@brandeis.edu https://orcid.org/0000-0002-7429-457X

Marder, E. marder@brandeis.edu https://orcid.org/0000-0001-9632-5448

Biology Department and Volen Center, Brandeis University, Waltham, MA 02454

Corresponding author: Eve Marder marder@brandeis.edu

*Authors contributed equally to this work

[1] Present address:
Biology Department
Bowdoin College
255 Maine St.
Brunswick, ME 04011

[2] Present address:
Department of Molecular Life Sciences
University of Zürich, Irchel
Winterthurerstrasse 190
CH-8057 Zürich, Switzerland

Key Words: pyloric rhythm, gastric mill rhythm, stomatogastric ganglion, central pattern generator, integer coupling

Abbreviated title: Stable temperature coupling of fast and slow oscillators

34 Acknowledgments: We thank Dr. Michael P Nusbaum for insights relevant to the
35 operation of the gastric mill circuit.

37 Research supported by: 5T32-NS007292 (SGS); R35-NS097343 (EM)

Abstract

Coupled oscillatory circuits are ubiquitous in nervous systems. Given that most

biological processes are temperature sensitive, it is remarkable that the neuronal

circuits of poikilothermic animals can maintain coupling across a wide range of

temperatures. Within the stomatogastric ganglion (STG) of the crab, *Cancer borealis*,

the fast pyloric rhythm (~1Hz) and the slow gastric mill rhythm (~0.1Hz) are precisely

coordinated at ~11°C such that there is an integer number of pyloric cycles per gastric

mill cycle (integer coupling). Upon increasing temperature from 7-23°C, both oscillators

showed similar temperature-dependent increases in cycle frequency, and integer

coupling between the circuits was conserved. Thus, although both rhythms show

temperature dependent changes in rhythm frequency, the processes that couple these

circuits maintain their coordination over a wide range of temperature. Such robustness

to temperature changes could be part of a toolbox of processes that enables neural

circuits to maintain function despite global perturbations.

Introduction

Coupled oscillators are common in nervous systems. Although oscillatory circuits may have distinct frequencies and duty cycles, coordination between them is often necessary for proper function (Nadim et al., 1998; Bartos et al., 1999; Colgin, 2011; Gordon, 2011; Jacobson et al., 2013; Rojas-Líbano et al., 2014; Harris and Gordon, 2015; Karalis et al., 2016; Tamura et al., 2017). For example, theta frequency circuits coupled to those in the gamma range are thought to drive both sensory and behavioral processing (Lisman and Buzsáki, 2008; Fujisawa and Buzsáki, 2011; Gordon, 2011; Lisman and Jensen, 2013). Memory formation may rely on the phase coupling of discrete populations of neurons with distinct frequency bands in the hippocampus (Tass et al., 1998; Lisman, 2005; Montgomery and Buzsaki, 2007; Scheffer-Teixeira and Tort, 2016; Zheng et al., 2016). Network uncoupling or improper coupling may lead to disease states and circuit malfunction (Moran and Hong, 2011; Kirihara et al., 2012; de Hemptinne et al., 2013; Bahramisharif et al., 2016; Salimpour and Anderson, 2019). The coordination of oscillatory circuits, often with distinct temporal features, is therefore key to circuit function and information processing. Because fluctuations in a circuit's environment can impact function (Fonseca and Correia, 2007; Tang et al., 2010; Haddad and Marder, 2018; Haley et al., 2018; Kushinsky et al., 2019; He et al., 2020), it is important to know how robust this coordination is to externally or internally generated perturbations.

The stomatogastric ganglion (STG) of the Jonah crab, *Cancer borealis*, contains two well-characterized, coupled, oscillatory circuits with known connectivity that generate distinct rhythms with different cycle periods (Coleman et al., 1995; Nadim et

al., 1998; Marder and Bucher, 2007). The pyloric rhythm (~1 Hz) is driven by a three-neuron bursting pacemaker kernel composed of the single Anterior Burster (AB) neuron and two Pyloric Dilator (PD) neurons (Nadim et al., 1998; Marder and Bucher, 2007). The gastric mill rhythm (~0.1 Hz) is produced by activating a half-center oscillator (reciprocal inhibition that produces alternating bursts of activity) between the Lateral Gastric (LG) motor neuron and Interneuron 1 (Int1), and it depends on a complex set of mechanisms including electrical junctions, and modulatory input (Coleman and Nusbaum, 1994; Coleman et al., 1995; Bartos and Nusbaum, 1997; Nadim et al., 1998; Bartos et al., 1999; Marder and Bucher, 2007; Stein et al., 2007). Although the pyloric and gastric mill oscillators independently produce distinct rhythms (Bartos et al., 1999), the core generator neurons (AB and LG) are integer coupled to each other and influence one another (Coleman and Nusbaum, 1994; Marder et al., 1998; Nadim et al., 1998).

In the New England Atlantic Ocean, where *C. borealis* is found, intertidal temperatures normally range from 8-24°C during the summer and 3-18°C in the winter (Stehlik et al., 1991). Because temperature affects most biological processes differently, it poses an interesting challenge for the nervous systems of poikilothermic animals that experience substantial temperature changes (Hazel and Prosser, 1974; Zečević and Levitan, 1980; Prosser and Nelson, 1981; Burrows, 1989; Foster and Robertson, 1992; Xu and Robertson, 1994; Xu and Robertson, 1996; Franz and Ronacher, 2002; Garrity et al., 2010; Marder et al., 2015). For example, the timescales of activation and inactivation of ion channels can change $\sim 2 - 20$ fold across 10°C (Q_{10}s) (Zečević et al., 1985; Klockner et al., 1990; Moran et al., 2004; Cao and Oertel, 2005; Fohlmeister et al., 2010; Tang et al., 2010; Kang et al., 2012; Robertson and Money, 2012; Yang and Zheng,

2014). Because neuron burst generation depends on many interdependent processes, if

even a few key processes have Q_{10}s that are appreciably different, this could dramatically

disrupt burst generation and/or alter the Q_{10} of a neuron's cycle period. Although the

phase relationships between different pyloric neurons are stable between 7 and 23°C

(Tang et al., 2010; Haddad and Marder, 2018), little is known about temperature

compensation in the gastric mill oscillator (Städele et al., 2015) or how temperature

affects coupling between the pyloric and gastric mill rhythms.

A key feature of temperature compensation in the pyloric circuit is that the

pacemaker kernel of the pyloric rhythm maintains constant duty cycle across a range of

temperatures (Tang et al., 2010; Rinberg et al., 2013), and this facilitates temperature

compensation of neuron phase relationships in the rest of the circuit. Because the

pyloric and gastric mill rhythms arise from different processes, it was not clear that the

gastric mill rhythm would be phase compensated as a function of temperature, and

whether the relationships between the two rhythms, such as integer coupling, would be

maintained as temperature varies.

Results

Two rhythmic oscillators driven by different processes

There are two major rhythms produced by neurons in the STG. The pyloric rhythm is constitutively active, with a characteristic period of about 1 sec. The gastric mill rhythm can be evoked by stimulating projection neurons found within the commissural ganglia (CoGs), and has a characteristic period of 6-10 sec (Beenhakker et al., 2004). The modulatory commissural neuron 1 (MCN1) and commissural projection neuron 2 (CPN2) (**Fig. 1a**) evoke a gastric mill rhythm when stimulated or when they are spontaneously active (**Methods**) (Beenhakker et al., 2004).

The activity of both circuits can be monitored by extracellular recordings from the motor nerves exiting the STG. The gastric mill rhythm can be seen in the lateral gastric nerve (*lgn*), dorsal gastric nerve (*dgn*), and medial ventricular nerve (*mvn*) recordings in **Fig. 1b**. These recordings show alternating bursts of activity in the LG and Dorsal Gastric (DG) neurons, with the Gastric Mill (GM) neurons firing at the end of the LG burst. The VD (Ventricular Dilator) and IC (Inferior Cardiac) neurons, seen on the *mvn,* show an envelope of activity alternating with the LG neuron, but also fire in time with the faster pyloric rhythm seen on the lateral ventricular nerve (*lvn*) and pyloric dilator nerve (*pdn*) (**Fig. 1b**). Because the AB and PD neurons are strongly electrically coupled, the cycle period of the pyloric pacemaker can be monitored using the *pdn* (**Fig. 1b**), which contains only PD neuron action potentials. Similarly, the LG neuron indicates gastric mill rhythm cycle period (**Fig. 1b**).

Figure 1c is a connectivity diagram of neurons in the STG and shows how MCN1 and CPN2 activate the gastric mill neurons. While CPN2 drives the gastric mill circuit

via electrical synapses onto key motor neurons (LG and GM), MCN1 excites all neurons

in the STG, including the pyloric pacemaker and gastric mill half-center neurons (LG

and Int1) (**Fig. 1d**). Both pattern generating circuits provide feedback to these

projection neurons (Norris et al., 1994; Blitz and Nusbaum, 2008, 2012; Blitz, 2017).

The interactions between CPN2, MCN1, and LG shape the timing of the gastric mill

rhythm along with rhythmic inhibition of Int1 by AB that allows LG to be rhythmically

released from inhibition and burst (Coleman and Nusbaum, 1994; Coleman et al., 1995).

The cartoon voltage traces (**Fig. 1d**) demonstrate integer coupling between the pyloric

and gastric mill circuits, where rhythmic AB-inhibition of Int1 results in LG

disinhibition and firing following the peak of Int1 inhibition (Nadim et al., 1998; Bartos

et al., 1999) .

Figure 1. STG rhythmic motor patterns. (**a**) Schematic of the STNS showing the locations of salient ganglia, nerves, and projection neurons. Individual STG neurons are unambiguously identified on the basis of the motor nerves by which they project to specific stomach muscles. (**b**) Extracellular nerve recordings of both the gastric mill (*lgn, dgn, mvn*) and pyloric (*lvn, pdn*) rhythms. The gastric mill rhythm is seen as the alternating bursts of activity in the LG and DG neurons. The GM neurons are active late in the LG neuron burst. The IC and VD neurons are active in alternation with LG, but also show activity in time with the pyloric rhythm (*lvn)*, with the PD neurons shown on the *pdn*. (**c**) Wiring diagram of the gastric mill, pyloric, and gastropyloric neurons and the input they receive from MCN1 (tan) and CPN2 (brown) (Marder and Bucher, 2007; Blitz, 2017). LG (pink) and DG (green) are both gastric mill neurons and AB-PD (black) are pyloric neurons. Colors used for these neurons are consistent in all figures. The resistor symbols denote electrical synapses, diode symbol denotes a rectifying junction, chemical inhibitory synapses are shown as filled circles, and the T bars indicate chemical excitatory/modulatory synapses from MCN1. Neuron copy number is in parenthesis. (**d**) (Top) Simplified wiring diagram of MCN1 excitation of both pattern generating circuits and AB inhibition of Int1 (white) (Coleman and Nusbaum, 1994;

Bartos and Nusbaum, 1997). (Bottom) Cartoon voltage traces of AB, Int1, and LG as they would appear during a gastric mill rhythm showing the relative burst timing between these neurons. As a consequence of a slow excitation from MCN1 to LG, the envelope of peak disinhibition of LG (from Int1) increases slowly over several cycles, eventually resulting in LG crossing threshold and bursting in time with the AB neuron. Note the slight delays between the peak of AB depolarization, the trough of Int1 inhibition from AB, and the subsequent burst of LG as marked by the dashed line.

Spontaneous gastric mill rhythms span a physiological range of temperatures

While the circuit controlling the pyloric rhythm is temperature robust from 7°C – 25°C (Tang et al., 2010; Tang et al., 2012; Rinberg et al., 2013; Soofi et al., 2014; Haddad and Marder, 2018), it remains unclear if the gastric mill circuit could also operate across this same range (Städele et al., 2015). We inspected a dataset of extracellular recordings of *dgn* and *lgn* during temperature ramps (Haddad and Marder, 2018) and found spontaneous gastric mill activity at temperatures between 11°C and 30°C.

Figure 2a shows bursts of LG (*lgn*) and DG (*dgn*) neurons from a single preparation showing spontaneous gastric mill activity at these temperatures. While these recordings show relatively regular gastric mill rhythms at lower temperatures, they were less regular at higher temperatures. **Figure 2b** shows example raster plots of the LG neuron for two preparations with spontaneous gastric mill rhythms. Color indicates the temperature at which that rhythm was observed and can be seen in the raster plots from two preparations across a range of temperature. Preparation 1 intermittently displayed characteristic gastric mill rhythms at both high and low temperatures, but was much more irregular in the middle of the temperature range, whereas Preparation 2 was considerably less robust at high temperatures.

Figure 2. Spontaneous gastric mill rhythms over physiological temperature range. (**a**) Extracellular recordings from the *lgn* and *dgn* at the temperatures indicated using the color code. (**b**) Spike rasters of LG from two preparations at the indicated colors to show the activity of these preparations over extended periods of time at the indicated temperatures. Traces in (**a**) and rasters in Preparation 2 in (**b**) are from the same animal. Rasters with the same color are continuous recordings at the same temperature, and time wraps around the end of each raster and continues on the next line.

Although **Figure 2** only shows two examples of spontaneous gastric mill rhythm, there were a total of 9 preparations in which spontaneous rhythms were observed at a variety of different temperatures (**Fig. 2–figure supplement 1**). Together, these data

indicate that activating at least some combination of endogenous inputs to the gastric

mill circuit can generate rhythmic activity across a wide temperature range.

Nonetheless, because the cycle period of the gastric mill was often irregular in these

spontaneous rhythms, we were not able to assess whether the pyloric and gastric mill

circuits remained coupled. Consequently, we chose to reliably activate gastric rhythms

by stimulating a known sensory pathway (**Methods**), to get a more reproducible gastric

mill rhythm, of known mechanism, across temperature and across preparations.

Figure 2–figure supplement 1. Spontaneous gastric activity in seven additional preparations. Each panel shows rasters of the LG neuron from a single preparation at temperatures from 11-30°C. Figure shows seven different preparations separated by black lines. LG was spontaneously active at temperatures as high as 30°C. Color of each row of raster indicates temperatures.

Gastric mill rhythms speed up with increased temperature

We were able to evoke gastric rhythms across the same temperature range at

which the pyloric circuit is robust (Tang et al., 2010). **Figure 3** shows example

recordings of evoked gastric rhythms at 7°C (**Fig. 3a**) and 23°C (**Fig. 3b**). Raster plots

from this preparation for the entire temperature range are shown in **Figure 3c**. Similar

to the pyloric circuit, the gastric mill rhythm cycle period decreased with increased

temperature from 7°C to 23°C. Although we observed that some of the neuron phase

relationships (relative timing of the neurons in each cycle) were altered at 23°C (e.g.

between DG and LG in **Figure 3b**), at high temperatures, the gastric mill rhythm

generator still produced regular rhythmic activity following stimulation. We were able to

evoke gastric rhythms across a temperature range of 7–21°C in 9/10 preparations.

Figure 3. Gastric mill rhythms can be evoked by stimulation over physiological temperature range. (**a**) Extracellular recordings from a preparation at 7°C (blue) and (**b**) 23°C (orange). Neuron action potentials described by their relative amplitudes on each nerve: *lgn*: LG (large bursting unit); *dgn*: DG (large, bursting unit; out of phase with LG: only on 7°C trace, has ceased bursting at 23°C), AGR (tonic), GM (small bursting unit; in-phase with LG); *mvn*: IC (large bursting unit), VD (medium bursting unit, antiphase with LG, alternates with IC), GM (small bursting unit; in-phase with LG); *pdn*: PD. (**c**) Spike rasters showing the activity of the LG neuron over the entire temperature range. The first row corresponds to (**a**) and the last row corresponds to (**b**).

In some preparations (4/10), spontaneous gastric mill rhythms were observed

prior to evoking a rhythm. **Figure 3–figure supplement 1a** shows example

recordings of both a spontaneous (black) and evoked (blue) gastric mill rhythm from the

same preparation (11°C). Three of the four preparations that had spontaneous and

evoked rhythms shared similarities in LG spike and burst timing (**Fig. 3–figure**

supplement 1b) and rhythm cycle period (mean ± SD) (**Fig. 3–figure supplement**

1c). Because there are ten known processes of gastric mill rhythm generation and all but

two are distinct from one another (Heinzel et al., 1993; Coleman and Nusbaum, 1994;

Norris et al., 1994, 1996; Weimann et al., 1997; Blitz et al., 1999; Christie et al., 2004;

Saideman et al., 2007a; Saideman et al., 2007b; White and Nusbaum, 2011; Dickinson

et al., 2015; Blitz et al., 2019), similarities between spontaneous and evoked gastric mill

rhythms within a preparation suggested that these rhythms are driven by similar

processes.

Figure 3–figure supplement 1. Comparison of spontaneous and evoked gastric mill rhythms within the same preparation. In 4/10 preparations, spontaneous gastric mill rhythms were observed at 11°C. (**a**) Extracellular traces showing activity on *lgn, dgn, mvn* and *pdn* nerves during spontaneous gastric mill activity (black) and after stimulation (blue). (**b**) Inter-spike intervals (ISIs) of LG neuron in the four preparations that showed spontaneous gastric mill activity. Black dots indicate ISIs from spontaneous activity and blue dots from evoked activity. (**c**) Comparison of burst period of LG during spontaneous and evoked bursting. Raw traces in (**a**) correspond to Preparation 1 in (**b**).

Gastric mill and pyloric cycle periods are similarly temperature sensitive

The pyloric cycle period decreases ~2-fold with a 10°C increase in temperature (Tang et al., 2010). If the pyloric and gastric mill circuits are to be coordinated across temperature, we hypothesized that the gastric mill period might also show a similar temperature dependence. We measured the burst periods of PD neurons (**Fig 4, black**) and LG neurons (**Fig 4, magenta**) in all 10 preparations in which the gastric mill

rhythm was evoked by stimulation over a temperature range from 7–21°C. **Fig 4a** shows

that, in a single preparation, both LG and PD burst periods decrease with increasing

temperature, despite significant variability in the LG burst period, due to a slowing

down (**Figure 4 − figure supplement 1**) of the gastric mill rhythm with time since

stimulation (Beenhakker et al., 2004; Stein et al., 2007).

A coordinated decrease in both gastric mill and pyloric cycle periods was

observed in all ten preparations (**Fig. 4b**). We quantified the extent to which the cycle

period increased with temperature using the Q_{10} (**Methods**). **Figure 4c** shows that

most preparations (8 of 10) exhibited a cycle period Q_{10} of ~2 for both the pyloric and

gastric mill rhythms. Estimated Q_{10}s were not significantly different for LG and PD burst

periods (paired *t*-test, $p = 0.235, N = 10, t = 1.27$), suggesting that the periods of both

circuits were not differently influenced by increased temperature.

**Figure 4. Pyloric and Gastric mill periods are similarly temperature
sensitive.** (**a**) Burst period of PD (black) and LG (magenta) in a single preparation as a
function of temperature. Each dot corresponds to a single burst. (**b**) Burst period of PD
and LG for ten preparations. Each line is a single preparation. Mean of all preparations
shown in thick lines. (**c**) The apparent Q_{10} of the burst frequency of each preparation for
the PD and LG neurons, representing the pyloric and gastric mill rhythms. The
apparent Q_{10}s of the two rhythms are not significantly different (paired *t*-test, $p = 0.235, N = 10, t = 1.27$)

Figure 4 − figure supplement 1. Gastric mill rhythms slow down with time after stimulation. (**a**) Instantaneous LG burst period plotted against time for a single preparation. Each line shows LG burst period for each gastric mill rhythm. Lines are of varying length as gastric mill rhythm duration was different each time it is evoked (**Methods**). Colors indicate temperature (7 − 21°C). (**b**) Mean LG burst period for the first 600 seconds post-stimulation for all 10 preparations. (**c**) Linear fits for average LG burst period across all preparations at each temperature. (**d**) Slopes of the linear fits in (**c**) plotted vs. r^2 of each fit line.

Gastric mill rhythms can be evoked when the pyloric pacemaker is

suppressed

At 11°C the gastric mill half-center is capable of producing rhythmic activity even if the pyloric rhythm is turned off by hyperpolarization of the pacemakers (Bartos et al., 1999). Therefore we hypothesized that the gastric mill oscillator might be independently temperature robust in the absence of pyloric input. To test this, we evoked gastric mill rhythms at 11°C, 15°C, 19°C, and 21°C with the pyloric pacemaker both active and inactive (**Methods**).

Figure 4 – figure supplement 2a and b show example recordings from a single preparation in which, at each temperature, we evoked a "control" rhythm (AB/PD active) followed by an evoked rhythm with both PD neurons hyperpolarized (AB/PD inactive). **Figure 4 – figure supplement 2c** shows that the gastric mill rhythm cycle period steadily decreased with increased temperature regardless of pyloric pacemaker activity. **Figure 4 – figure supplement 2d** shows that in all 9 preparations, the gastric mill rhythm was activated at 11°C and 15°C, whereas at 19°C and 21°C only a subset of preparations produced gastric mill rhythms at these elevated temperatures.

Figure 4 – figure supplement 2. Evoked gastric mill rhythms persist when the pyloric pacemaker is suppressed. Extracellular recordings of the *lgn* and *pdn* at 11°C, 15°C, 19°C, and 21°C with the pyloric pacemaker kernel active (**a**) and inactive (**b**) in a single preparation. (**c**) Burst period of LG with the pacemaker active (magenta) and inactive (yellow) at each temperature step for the preparation shown in (**a**) and (**b**) (Mean ± SEM). (**d**) Number of preparations (out of 9 total) in which a gastric mill rhythm could be evoked with the pacemaker kernel active (bottom row) and with AB/PD hyperpolarized (top row) for each temperature.

Integer coupling is maintained across a physiological temperature range

Integer coupling between two interacting oscillators with different periods refers

to the phenomenon in which the slower rhythm (gastric mill) tends to have a period that

is an integer multiple of the period of the faster cycle (pyloric) (Nadim et al., 1998).

Integer coupling exists between the pyloric and gastric mill oscillators at 11°C (Bartos et

al., 1999; Hamood and Marder, 2015). It remains an open question if this form of

precise coordination is preserved across temperature, because both oscillators change

their cycle periods with temperature.

Extracellular recordings of *lgn* and *pdn* at 7°C and 23°C from the same

preparation are shown in **Fig. 5a**. PD bursts tended to terminate shortly before the

initiation of LG bursting at both temperatures (**Fig. 5a**, dotted vertical lines indicate

start of LG bursting). The ratio of LG to PD periods was close to an integer at both

temperatures (~10 at 7°C and ~11 at 23°C).

To visualize integer coupling between the PD and LG across the entire dataset, we

plotted the LG burst period as a function of the mean PD burst period during that LG

burst (**Fig. 5b**). Data tended to lie along lines with integer slope (gray lines), suggesting

integer coupling between LG and PD across temperature. DG burst periods, which were

similar to LG burst periods, did not tend to lie on lines with integer slope (**Fig. 5c**). To

quantify the degree of integer coupling, we computed the cumulative distribution of

significands (value after the decimal point) of the ratios of gastric mill to pyloric periods

(**Fig. 5d**, **Methods**). Perfect integer coupling would lead to a cumulative distribution

indicated by the dotted line, while randomly chosen significands would lie along the

diagonal. Cumulative distributions for LG were skewed towards perfect integer

coupling, while distributions for DG were closer to the randomly distributed

significands, suggesting that integer coupling was observed in LG but not DG.

To quantify how the degree of integer coupling varies across temperature, we

measured the area between the cumulative distributions and the diagonal in **Fig. 5d**.

Perfect integer coupling would yield an area of 0.25. Areas for LG/PD significands were

larger than areas for DG/PD significands at every temperature (**Fig. 5e**), suggesting

greater integer coupling. Areas for DG/PD significands overlapped with areas computed

after shuffling gastric mill neuron (both LG and DG) and PD neuron cycle periods (gray

lines, **Fig. 5e**). To determine if measured integer coupling was significantly different

from chance, we used a two-sample Kolmogorov-Smirnoff test (**Methods**). LG and PD

were significantly integer coupled, at temperatures from 7°C–21°C ($p < 10^{-5}, D > .24$),

while DG and PD were not significantly coupled at any temperature ($p > .02, D < .16$).

AB dictates the phase at which LG turns on because LG onset is timed with the peak of

Int1 inhibition (Nadim et al., 1998). Because all previous studies of integer coupling

were at ~11°C, it was not known if the process underlying integer coupling would shift

with temperature increase. To visualize the effect of temperature on the ratio of PD to

LG cycle period, we plotted the ratio of LG cycle period and PD cycle period as a

function of temperature (**Fig. 5f**). At low temperatures, when integer coupling was

particularly strong (as seen from the horizontal banding), these ratios were distributed

over a wide range from 5-25. Ratios of LG to PD cycle period were spread across a

similar range across temperature, even at higher temperatures when integer coupling

was less pronounced ($p = 0.88, n = 3799$ PD cycles pooled across all preparations,

Spearman test), suggesting that the mechanism of integer coupling was independent of

the number of PD cycles to LG cycle period.

Figure 5. Integer coupling is maintained over physiological range. (**a**) Extracellular recordings of *pdn* show bursting of LG and PD at 7°C (blue) and 23°C (orange). Black bars indicate burst period of LG and mean burst period of PD. LG (**b**) and DG (**c**) burst period as a function of mean PD period. In (**b-c**), gray lines have integer slopes, and are not fits. (**d**) Cumulative probability distributions (c.d.f.s) of significand of ratio of gastric to pyloric period for LG (magenta) and DG (green) for all temperatures. Shading indicates confidence estimates from bootstrapping the data. Dashed diagonal line indicates a uniform distribution, and the dotted line indicates

perfect integer coupling. (**e**) Area between significand c.d.f.s and diagonal (a measure of
how integer coupled the circuits are) as a function of temperature for all preparations.
Gray lines are computed from shuffled data. LG and PD were significantly integer
coupled between 7°C–21°C ($p < 10^{-6}$ two-sample Kolmogorov-Smirnoff test), while DG
and PD did not exhibit integer coupling. (**f**) Ratio of gastric to pyloric periods as a
function of temperature. Apparent banding along the temperature axis is a consequence
of the experimental protocol, where discrete temperature steps were used (Methods).
Ratios of LG to PD cycle period were similar across temperature ($p = 0.88$, Spearman
test).

LG preserves phase locking to PD across physiological temperatures

How do LG and PD maintain integer coupling in their burst periods across this

temperature range? One possibility is via temperature-invariant phase locking between

LG and PD. If LG bursts start at a particular phase in the PD cycle, then the burst

periods of LG and PD must be integer coupled (Nadim et al., 1998). We therefore set

out to study the fine structure of the phase coupling between LG and PD across

temperature.

To visualize the phase coupling between LG and PD, we plotted PD spikes aligned

to LG burst starts (**Fig. 6a**) normalizing time by the mean PD burst period in each row.

Doing this aligns all PD spikes, suggesting that LG burst starts were an excellent

predictor of PD phase. To quantify this effect across all preparations, we computed

probabilities of LG burst starts across the temperature range tested. LG burst start

probabilities were sharply peaked (**Fig. 6b**), and were significantly different from a

uniform distribution (**Fig. 6b**, dashed line, ($p < 4 \times 10^{-4}, 7.5 < z <$

45, Rayleigh test for non $-$ uniformity, **see Methods** for details). From 7-21°C, LG bursts

started at a phase of ~0.5 in the PD cycle (**Fig. 6b, inset**), and this phase offset did not

significantly trend across this temperature range ($p = 0.46$, $\rho = -.31$, Spearman test).

LG burst initiation being phase locked to PD suggests that faster dynamics in LG,

such as spiking, could also be affected by PD. Modulation of LG spiking by AB/PD has

been reported via feedback from AB to MCN1 and CPN2 (Blitz and Nusbaum, 2008) and

possibly Int1 feedback to CPN2 (Blitz, 2017). We measured the spike probability of LG

conditional on PD phase across this temperature range, and found a pronounced

decrease in the probability of spiking in LG at higher temperatures (**Fig. 6c**). This

decrease in LG spike probability occurred close to PD burst termination (**Fig. 6c**,

horizontal bars).

Although DG bursts on the same timescale as LG, it is not directly coupled to the

pyloric oscillator (**Fig. 1**) and does not show any signature of phase coupling with PD.

DG bursts starts do not align with PD spikes (**Fig. 6d**), DG burst starts are not sharply

peaked in PD phase, and they are not significantly different from a uniform distribution

for all preparations at all temperatures (**Fig. 6e**, $p \in [.001, .85], .15 < z < 13$, Rayleigh

test). DG spike probability also does not show a noticeable decrease close to the end of

PD bursting (**Fig. 6f,** cf. **Fig. 6c**).

Figure 6. LG, but not DG, bursts and spikes are phase locked to PD across temperature. (**a**) PD spike rasters aligned to LG burst starts (dashed line at 0) in a single preparation. Each row corresponds to a LG burst start, and time is normalized by PD burst period. Black histograms at the top show spike probability over all rasters. (**b**) LG burst start probability as a function of PD phase. Inset shows mean LG burst start phase a function of temperature (gray lines: individual preparations, black line: mean across all preparations). LG burst start probabilities were significantly different from a uniform distribution ($p < 4 \times 10^{-4}, 7.5 < z < 45$, Rayleigh test for non-uniformity). Phase offset between LG and PD was similar from 7 to 21°C (inset, $p = 0.46$, Spearman test). (**c**) LG spike probability as a function of PD phase at various temperatures. Bars indicate PD spiking. (**d-f**) Similar to (**a-c**), but for DG. DG burst starts are not always significantly non-uniformly distributed, and the effect size is much smaller ($p \in [.001, .85], .15 < z < 13$, Rayleigh test). Rasters in (**a,d**) are from the same preparation.

Discussion

Temperature sensitive circuits maintain coordinated activity

Coupling between rhythmic circuits is common in nervous systems and is often functionally important, although the strength and the extent of inter-circuit coupling is varied (Nadim et al., 1998; Bartos et al., 1999; Colgin, 2011; Gordon, 2011; Jacobson et al., 2013; Rojas-Líbano et al., 2014; Harris and Gordon, 2015; Karalis et al., 2016; Tamura et al., 2017). Temperature is a global perturbation that influences the properties of all ion channels and synapses in neuronal circuits (Zečević et al., 1985; Janssen, 1992; Xu and Robertson, 1996; Garrity et al., 2010; Tang et al., 2010; Kang et al., 2012; Robertson and Money, 2012; Marder et al., 2015). Because each cellular and molecular process is likely to have a different temperature dependence, or Q_{10}, maintaining the function of complex physiological processes as temperature is changed is non-trivial (Du et al., 2010; Caplan et al., 2014; Roemschied et al., 2014; O'Leary and Marder, 2016; Schleimer and Schreiber, 2016). Most often, changes in temperature affect neuronal and circuit behavior (Zečević and Levitan, 1980; Prosser and Nelson, 1981; Long and Fee, 2008; Ramot et al., 2008; Garrity et al., 2010; Robertson and Money, 2012). It then becomes critical to know whether the coupling between circuits is also altered (Stiebler and Narins, 1990), or whether the relative coordination between rhythmic behaviors can be maintained as temperature changes.

Here we show that two oscillator circuits can maintain coordinated output across a range of physiologically relevant temperatures. The cardiac ganglion, another circuit in the crab which governs heartbeat, is also temperature sensitive across a range of physiological temperature, again with a Q_{10} of ~2 (Kushinsky et al., 2019). In the pyloric,

gastric mill, and cardiac circuits, rhythmicity is maintained up to a critical temperature,

and this critical temperature approximates the upper range of native ambient

temperatures (Tang et al., 2010; Städele et al., 2015; Kushinsky et al., 2019).

Coupling of the gastric mill and pyloric circuits is likely to be important for the

crab. In the intact animal, muscle movements of the gastric mill enable ingested food to

be chewed and subsequently filtered and pumped into the hindgut by the pylorus

(Marder and Bucher, 2007). Coupling between these circuits coordinates the timing of

chewing and filtering enabling proper flow of food through the mid-gut (Meyrand et al.,

1994; Diehl et al., 2013). Furthermore, a subset of STG neurons participates in both

rhythms, and therefore coupling between the pyloric and gastric mill circuits enables

neurons with dual functions to maintain their appropriate participation in both rhythms

(Weimann et al., 1991; Weimann and Marder, 1994; Gutierrez et al., 2013; Blitz et al.,

2019).

Oscillator circuits with different periods are similarly temperature robust

Although the gastric mill and pyloric rhythms are similarly temperature sensitive,

they have an approximately 10-fold difference in cycle period. The pyloric rhythm is

temperature robust because the AB neuron maintains a relatively constant duty cycle

across this range (Tang et al., 2010; Rinberg et al., 2013). Temperature robustness in the

gastric mill rhythm arises from a different set of interdependent processes including

projection neuron input, coordination of the half-center oscillator neurons LG and Int1,

and feedback from LG to MCN1. The similarity between the apparent cycle period Q_{10}s

of the pyloric and gastric mill rhythms was unexpected as there are so many different

physiological processes generating the two rhythms. Moreover, a lock step variance in

cycle period is not necessary in order to maintain coupling as the relative number of

pyloric cycles between each gastric cycle can vary widely while circuits remain coupled

(Nadim et al., 1998; Bartos et al., 1999). Because the salient drivers of LG firing are

sufficient projection neuron excitation and disinhibition from Int1 (Bartos et al., 1999;

DeLong et al., 2009), the total number of pyloric cycles per gastric mill cycle can vary

greatly.

Spontaneous gastric mill rhythms were non-stationary and driven by

unknown processes

As spontaneous gastric mill activity was often irregular, it was difficult to

examine coupling between the two rhythm generating circuits. The observed irregularity

could be due to weak activation of one or more descending pathways and is likely to be

different across preparations, and potentially across temperatures. There are many

different processes that drive gastric mill rhythms including release of hormonal

peptides such as crustacean cardioactive peptide (CCAP) (Weimann et al., 1997),

FLRFamides (Heinzel et al., 1993; Weimann et al., 1993; Christie et al., 2004),

pyrokinins (Saideman et al., 2007b; Dickinson et al., 2015), and Gly[1]-SIFamide (Blitz et

al., 2019). Synaptically released peptides *C. borealis* tachykinin related peptide (CabTRP

Ia) (Blitz et al., 1999; Wood et al., 2000) and *C. borealis* pyrokinin peptide I & II

(CabPK) (Saideman et al., 2007a; Saideman et al., 2007b) generate similar gastric mill

rhythms, while proctolin (Blitz et al., 1999; Stein et al., 2007) excites all gastric mill

motor neurons.

Stimulation of certain projection neuron such as MCN1 (contains CabTRP,

proctolin, and GABA) (Coleman and Nusbaum, 1994; Coleman et al., 1995; Bartos and

Nusbaum, 1997), MCN5 (contains Gly^1-SIFamide) (Norris et al., 1996; Blitz et al., 2019),

MCN7 (contains proctolin) (Blitz et al., 1999), CPN2 (Norris et al., 1994), and IVN (the

inferior ventricular neurons) (contain FLRFamide and histamine) (Christie et al., 2004),

or stimulation of sensory pathways that activate groups of projection neurons (Katz and

Harris-Warrick, 1991; Beenhakker et al., 2004; Beenhakker and Nusbaum, 2004; Blitz

et al., 2004; Beenhakker et al., 2007; White and Nusbaum, 2011) generate gastric mill

rhythms. Because there are so many ways gastric mill rhythms can be generated, and

because each method of gastric mill rhythm generation could result in a gastric mill

rhythm via a different set of mechanisms, they could have different temperature

dependencies. Consequently, we chose to elicit gastric mill rhythms using a single,

known mechanism, and we used exogenous stimulation that would reliably produce

gastric mill rhythms, that were qualitatively similar to the spontaneous gastric mill

rhythms we had previously observed.

A previous study showed that one gastric mill rhythm variant is easily disrupted

by small temperature changes (Städele et al., 2015), while the one studied here

maintains rhythmicity following a large increase in temperature. Although the

functional significance of the diversity among gastric mill rhythms is unclear, the

possibility remains that the assortment of gastric mill rhythm generator mechanisms

bequeaths a robustness against environmental perturbation. Because there are other

versions of the gastric mill not considered here, it will be interesting in the future to

determine the extent to which they are temperature robust and whether, in turn, they

are coupled to the pyloric rhythm across temperature.

Temperature compensation is driven by multiple mechanisms

Previous work has shown that neuromodulatory substances can extend the

temperature range over which the gastric and pyloric rhythms maintain function

(Städele et al., 2015; Haddad and Marder, 2018). In *C. elegans* neuromodulation

enhances the robustness of thermosensory neurons and altering temperature dependent

behavior (Beverly et al., 2011). Therefore, it is possible that some modulators that act on

the STG, not studied here, may further stabilize these rhythms, and some of the

modulatory inputs to the STG may release transmitter in response to temperature stress

for precisely this reason.

The gastric mill oscillator is activated by both chemical and electrical synapses. In

crayfish, the rectifying electrical junction from the lateral giant fiber to the motor giant

fiber has reliable synaptic propagation between 20°C and 30°C (Heitler and Edwards,

1998). In a model of this crayfish circuit, Heitler and Edwards (1998) found that if the

electrical synapse had a Q_{10} of 11, when most other measured properties of the circuit

had Q_{10}s between 2 and 5, the model reliably reproduced physiological responses to

temperature increase. In this system, at high temperatures, this increased drive from the

electrical synapse offsets the decrease in input resistance of the postsynaptic neuron. In

the evoked gastric mill rhythms, CPN2 drives LG and GM via electrical synapses

(**Figure 1c**), and our spontaneous rhythms, indicative of CPN2 activation, were

observed up to 30°C. While we do not know the Q_{10} of electrical synapses in the STG,

several measured chemical synapses in the pyloric circuit have a Q_{10} of ~2.5 (Tang et al.,

2010).

Previous work showed that input resistance modestly decreases in both the

Lateral Pyloric (LP) and LG neurons with increased temperature (Tang et al., 2010;

Städele et al., 2015). The MCN1–LG electrical synapse may help offset the change in

input resistance in the gastric circuit. In the pyloric circuit, the IPSP amplitude in the LP

neuron is temperature invariant (Q_{10} ~1), but the IPSC amplitude increased (Q_{10} ~2.5)

across the same range of temperatures as those studied here (Tang et al., 2010). The

increase in synaptic current may partially compensate for the increase in leak as

temperature increases. This could aid circuits in maintaining rhythmicity across

temperature. Furthermore, many STG neurons, including both LP and LG, exhibit post-

inhibitory rebound (PIR). Both I_A and I_h (currents that promote bursting and PIR)

exhibit temperature dependent increases in activation rate and peak current (Q_{10}s ~2 −

4) (Tang et al., 2010).

Circuits are robust to global perturbation

Although it may not be uniformly true that all neuronal circuits maintain

coordinated function across a wide range of temperatures in cold blooded animals, here

we demonstrate that for two coupled circuits, where coupling is necessary for proper

function, coordination is maintained despite a global perturbation. Because each

circuit's oscillations arise from a different set of circuit and cellular mechanisms, it is

unlikely that the underlying processes endowing temperature compensation are the

same. Therefore, it was not necessarily expected that compensation would be similar for

both circuits.

The nervous systems of small homeotherms such as birds, rodents (Chaffee and

Roberts, 1971; Janssen, 1992), and even human infants (Mank et al., 2016; Barbi et al.,

2017) likely face similar challenges as a result of fluctuations in ambient temperature or

in fever response to illness. Although they have homeostatic processes that regulate

internal temperature shifts, these regulatory responses can be slow compared to

temperature change (Cheshire, 2016). Therefore, it is likely that their nervous systems

must be robust to some degree of temperature fluctuation to maintain function until a

homeostatic process can intervene to restore internal temperature. While mouse brain

neuron temperature sensitivities exist in a much narrower range than the crustaceans

studied here (Hori et al., 1999), the temperature range we use is well within the range

experienced by the crabs naturally.

In this study, we use temperature as a global perturbation. It is possible that

coupled neural circuits are similarly robust to other global perturbations, so that

coupling is maintained despite transient or chronic changes in ambient conditions. For

example, the pyloric oscillator is known to be robust to large changes in pH (Haley et al.,

2018) and to fluctuations in extracellular potassium (He et al., 2020). Collectively, these

data indicate that the crustacean STG (and likely many other neural circuits) are capable

of sustaining normal output when challenged with global perturbations. While the

processes that confer robustness may be different for different circuits and

perturbations, nervous systems have likely evolved to cope with a variety of

environmental changes.

Methods

Animals and dissection

Adult male Jonah crabs (*C. borealis*; $N = 19$ were obtained from Commercial Lobster (Boston, MA) and $N = 9$ from Moody's Seafood (Brunswick, ME)) and kept in tanks with circulating artificial sea water at 11°C for approximately one week prior to use. Preparations with spontaneous rhythms ($N = 9$), were from crabs caught in July of 2013 and studied soon thereafter. Preparations in which rhythms were evoked ($N = 10$) were from crabs caught July, Sept., and Dec. of 2017. Preparations in which the pyloric pacemaker was inhibited ($N = 9$), were from crabs caught Aug. and Sept. 2020. Thirty minutes prior to dissection, animals were chilled on ice. Dissections were carried out as previously described (Gutierrez and Grashow, 2009) and the dissected nervous systems were kept in chilled saline solution (440 mM NaCl, 11 mM KCl, 26 mM MgCl2, 13 mM CaCl2, 11 mM Trizma base, 5 mM maleic acid, pH 7.45 at 23°C) throughout the dissection. Each dissected stomatogastric nervous system was pinned out in a petri dish with a Sylgard-lined bottom and fresh saline was continuously superfused across the nervous system for the duration of the experiment (**Fig. 1a**).

Nerve Recordings

Motor neuron action potentials originating in the STG (**Fig. 1a**) were recorded extracellularly (**Fig. 1b**) from the nerves via stainless steel pin electrodes. Stretches of the *lgn*, *dgn*, *mvn*, *lvn*, and *pdn* were electrically isolated from the bath with Vaseline wells. Voltage signals were amplified using model 3500 amplifiers (A-M Systems) and digitized at 10 kHz using a Digidata 1440 A-D converter (Axon Instruments/ Molecular

621 Devices). Data were recorded using pClamp data acquisition software (Axon

622 Instruments/Molecular Devices, version 10.5).

Temperature Control

624 Saline temperature was monitored continuously using a temperature controller (model

625 CL-100, Warner Instruments) and altered during each experiment using an associated

626 Peltier device and thermocouple (SC-20 and TA-29, Warner Instruments). Saline inflow

627 to the nervous system was positioned within 1 cm of the STG so that the measured

628 temperature at the point of inflow was approximately that of the ganglion somata.

Gastric Mill Rhythm Stimulation

We bilaterally stimulated the dorsal posterior esophageal nerves (*dpons*) to elicit a

gastric mill rhythm (**Figure 1a & b**). Stainless steel pin electrodes were placed on either

side of each *dpon* and were sealed to the nerve using Vaseline. Stimuli were delivered

using a model 3800 stimulator (A-M Systems) via model 3820 stimulus isolation units

(A-M Systems). Ten, six-second long episodic stimulus trains (0.6 Hz) were used to

evoke gastric mill rhythms using a within-train stimulus rate of 15 Hz, as described in

Beenhakker and Nusbaum (2004). The *dpons* contain axons of the ventral cardiac

neurons (VCNs) that activate the MCN1 and CPN2, resulting in a VCN-gastric mill

rhythm that is often active for tens of minutes (Beenhakker et al., 2004; Beenhakker

and Nusbaum, 2004; Blitz and Nusbaum, 2008). The duration of each stretch of gastric

mill activity is not consistent either within an experiment at different temperatures or

across experiments at the same temperature. Thus, the time between each gastric mill

rhythm stimulation was different both between temperatures and across preparations.

Defining gastric mill rhythm variants

While evoked gastric mill rhythms are distinguished by the method used to drive them, in the case of spontaneous gastric mill rhythms, the phase relationships and timing of the gastric mill neurons are used to classify them (White and Nusbaum, 2011).

Recordings from the *mvn* show the Inferior Cardiac (IC) neuron and ventricular dilator (VD) neuron (**Fig. 1b**). Although in the absence of a gastric mill rhythm, the IC and VD neurons are typically active in time with the LP neuron and the Pyloric neurons (PY$_5$; there are approximately five PY neurons in each preparation) respectively, the initiation of a gastric mill rhythm will often shift the IC and VD neurons into gastric mill timing as shown in **Figure 1b**. Because the timing of the IC and VD neurons are state-dependent, they can aid in classifying spontaneous gastric mill rhythms.

In this study, spontaneous gastric mill rhythms resemble VCN-rhythms as IC and VD activity is restricted to the LG interburst interval, which is known to occur because of CPN2 and LG inhibition of the IC and VD neurons respectively. Additionally, the LG bursts in these spontaneous rhythms did not exhibit any obvious interruptions, as is the case for another gastric mill rhythm that is similar to a VCN-rhythm, save for this difference (White and Nusbaum, 2011).

Spontaneous and evoked gastric mill rhythm temperature ranges

In one set of experiments ($N = 9$), spontaneous gastric mill rhythms were observed. Here, temperature was increased from 11°C to 33°C in 2°C increments, and a given temperature step was held for at least 2 minutes. Time between steps was ~2 minutes.

For experiments in which gastric mill rhythms were evoked ($N = 10$), the temperature

was increased in either 2°C or 4°C steps starting at 7°C until a gastric mill rhythm could

no longer be evoked (typically 21°C or 23°C). Each temperature step was held for the

duration of evoked gastric mill activity, and only after the epoch of the gastric mill

rhythm terminated was the temperature advanced to the next step. The time between

temperature steps was ~2 minutes.

The VCN-gastric mill rhythms were difficult to generate consistently above 23°C, and in

some cases only persisted for several cycles. In some preparations, 19°C was the

warmest temperature that a gastric mill rhythm could be evoked. Because the

stimulation paradigm only activates a subset of projection neurons in the CoGs, these

manipulations may fail to characterize the full extent to which the gastric mill circuit can

maintain rhythmicity across temperature. Within the temperature range in which they

were readily evoked, gastric mill rhythms lasted for tens of minutes.

Evoked rhythms with AB/PD hyperpolarized

To study the effect of temperature on VCN-gastric mill rhythm alone, in a separate set of

experiments, gastric mill rhythms were evoked twice at 11°C, 15°C, 19°C, and 21°C with

the pyloric pacemaker both active and inactive at each temperature step ($N = 9$). To

inactivate the pyloric pacemaker, each PD neuron was impaled with a single bridge

balanced sharp electrode in a current clamp configuration. PD neurons were identified

by briefly hyperpolarizing the membrane potential and monitoring the number of spikes

recorded extracellularly on the *pdn*. Action potentials from both PD neurons can be

monitored on a single *pdn*. Because the electrical coupling between the AB and PD

neurons is strong (Maynard and Selverston, 1975; Shruti et al., 2014), it is not necessary

to hyperpolarize all three neurons. PD neurons were targeted because of their relatively

large size compared to the AB neuron. Because neurons swell with increased

temperature (Tang et al., 2010), it can be difficult to maintain intracellular recordings,

especially of small neurons, and not damage the cells.

Data analysis

Spike identification and sorting

To reliably identify spikes from extracellular traces, we wrote spike sorting software that

uses supervised machine learning to learn the shape of spikes from a small manually

labelled dataset. This software is freely available at https://github.com/sg-s/crabsort.

PD, LG and DG neuron spikes were identified on the *pdn, lgn* and *dgn* nerves recorded

extracellularly. In a small subset of the full data, spikes from these neurons were

identified using thresholding, dimensionality reduction of the full spike shape, and

manual labelling. Using this subset, we trained a fully connected neural network to

classify putative identified spikes in new data. The accuracy of the classification was

constantly monitored during sorting, and new annotations were used to continuously

re-train the network using an active learning framework. Classification accuracy always

exceeded 98%.

After the neural network classified putative spikes in a new dataset, to ensure high-

quality spike annotation: 1) we manually inspected the data in two-minute chunks to

ensure that no artifacts were accidentally labelled as spikes; 2) we used tools built-in to

the spike sorting software to visually inspect, at high-temporal resolution, spikes that

the neural network reported with low confidence; and 3) we used other built-in tools to

inspect spikes that differed statistically from other spikes in its class. These steps let us

generate high-quality spike annotations for PD, LG, and DG, even when these neurons

were partially obscured by other units or noise.

Detection of bursts and measurement of burst metrics

Bursts in PD, LG or DG were defined using inter-spike intervals (ISIs). ISIs longer than

1 second on LG or DG were defined to be an inter-burst interval and were used to

annotate the start of a burst. Spikes were considered to be part of a burst only if there

were at least n spikes per burst, with $n = 5$ for LG and DG and $n = 2$ for PD.

This method allowed us to identify all burst starts, burst stops, and therefore burst

durations and burst periods for every burst. We computed the duty cycle on a cycle-by-

cycle basis by dividing the burst duration by the burst period.

Estimation and comparison of Q_{10}s (Figure 4)

The Q_{10} of a process is defined as the factor by which it changes over a ten-degree

increment in temperature. We computed the Q_{10} of PD and LG burst periods at every

temperature using

$$Q_{10} = \langle \left(\frac{f_T}{f_{11}}\right)^{\frac{10}{T-11}} \rangle$$

where f_T is the mean burst frequency at temperature T and f_{11} is the mean burst

frequency at 11°C (the reference temperature).

To compare the Q_{10}s of the pyloric and gastric rhythms, we used a paired t-test and

compared the mean Q_{10}s averaged across each preparation. Each animal thus

contributed equally and we included all 10 preparations. t represents the test statistic.

Integer coupling analysis (Fig. 5)

To measure the integer coupling between PD burst periods and LG or DG burst periods

(**Fig. 5b, c**), we computed the mean burst period of all PD bursts within one LG/DG

burst period. Thus, every dot corresponds to one LG burst and all the PD bursts that

occur within that LG burst period. LG/DG burst periods are plotted on the vertical axis

and the mean PD burst period is plotted on the horizontal axis.

To determine if these burst periods are significantly integer coupled (**Fig. 5d**), we

computed the significand (defined here as the value after the decimal point) of the ratio

of the ordinate to the abscissa for each dot in **Fig. 5b,c**. For example, if an LG burst had

a period of 10.1 seconds, and the mean PD period during that LG burst was 1 second, the

significand would be 0.1. If the LG burst period was 9.9 seconds, the significand would

be 0.9. We sorted all significands for LG and DG and bootstrapped the data 1000 times

to estimate confidence intervals of cumulative density functions of the significands.

Shading in **Figure 5d** indicates these confidence intervals. Curves that deviated

substantially from the diagonal (the null distribution) corresponded to significant

integer coupling between the gastric mill and pyloric burst periods. To determine if any

set of significands was significantly different from the null distribution, we performed a

two-sample Kolmogorov-Smirnoff test comparing the significands from the ratios of

LG/DG to PD burst periods and significands from ratios of shuffled copies of LG/DG

and PD burst periods. The reason we used a two-sample Kolmogorov-Smirnoff test, and

not a one-sample test that compares to a uniform distribution directly, is that we cannot

expect significands to be distributed exactly uniformly, even for randomly distributed

data. We observed through numerical calculations that for certain combinations of

numerator (gastric cycle period) and denominator (pyloric cycle period) ranges, even

randomly sampled cycle periods can lead to non-uniform significand distributions. The

threshold for significance was divided by the number of comparisons (9 temperature

steps) to correct for multiple comparisons. Data in **Fig. 5d** are pooled across all

preparations and across all temperatures. Statistical tests were performed grouping by

temperature, but pooling across all preparations. D indicates the Kolmogorov-Smirnoff

test statistic.

The dotted and dashed lines in **Fig. 5d** indicate two extremes of the significand

distribution. If the gastric and pyloric burst periods were perfectly integer coupled, then

every significand of every ratio of periods would be 0 (or 1). Thus, the cumulative

probability of such an idealized distribution would be a step-like function shown by the

dotted line. On the other extreme, if the ratios of gastric to pyloric burst periods were

such that the significands were randomly and uniformly distributed across the unit line

(suggesting no integer coupling whatsoever), then the cumulative probability of such a

distribution would be close to a diagonal line with unit slope, shown by the dashed line.

These two extremes allow us to define a yardstick to measure the degree of integer

coupling for any observed spike train. We observe that all possible cumulative

distributions are bounded by the two extremal distributions (dotted line and dashed

line). Because the area of the square in **Fig. 5d** is 1, simple geometry tells us that the

area bounded by the maximal distributions is 0.25. We therefore quantified the degree

of integer coupling of the two burst periods (**Fig. 5e**) by measuring the area between the

null distribution (dashed diagonal line) and the significand c.d.f.

Generation of burst-aligned rasters of PD (Fig. 6a,d)

To visualize the interaction between pyloric and gastric mill rhythms, we generated

burst-aligned rasters of the PD neuron activity in which we aligned PD rasters to the

burst start in a gastric mill-timed neuron (LG or DG). For every burst start in LG/DG,

we plotted the raster of PD spikes centered around that burst start, showing spikes 3

seconds before the burst start in LG/DG and 3 seconds after.

Estimation of gastric mill spike and burst start probability in PD phase

(Fig. 6)

To estimate when the LG or DG neurons start bursting in the PD cycle (**Fig. 6**), we first

identified the burst start times of LG and DG. At these points, we estimated the PD

phase by dividing the time since the start of the previous PD burst by the burst period of

that PD cycle. This method of phase estimation assumes that the phase increases

uniformly and linearly over a PD cycle, following earlier work (Prinz et al., 2004; Bucher

et al., 2005). We used a similar procedure to estimate gastric mill spike probability in

PD phase.

Analysis of distribution of LG/DG burst starts in PD phase (Fig. 6 b,e)

LG burst starts in PD time follow circular statistics, since a LG burst start occurring at a PD phase of 1 is the same as a LG burst start at a PD phase of 0. We therefore used a circular statistics toolbox by Phillipp Berens (Berens, 2009) (http://www.eye-tuebingen.de/berenslab/technology-development/) to analyze this data. Means and standard deviations of LG and DG burst starts (**Fig. 6b** inset and **Fig. 6e** inset) were computed using this circular statistics toolbox. To test if LG and DG burst starts were significantly non-uniformly distributed, we used the Rayleigh test for non-uniformity. z is the Rayleigh test statistic. The distribution of burst starts of LG or DG in PD time for each preparation, at each temperature was tested for non-uniformity individually.

The number of LG/DG burst starts was not the same for every preparation at every temperature step. For this reason, and to avoid time-dependent confounds, we analyzed the first 40 burst starts for each preparation at each temperature step. Preparation and temperature combinations where there were fewer than 40 burst starts (~35% of data) were excluded. A total of n= 3816 LG burst starts and n=3228 DG burst starts across 9 temperature points and 10 preparations are plotted in **Fig. 6b, d**.

References

References

Bahramisharif A, Mazaheri A, Levar N, Richard Schuurman P, Figee M, Denys D (2016) Deep Brain
Stimulation Diminishes Cross-Frequency Coupling in Obsessive-Compulsive Disorder. Biol
Psychiatry 80:e57-58.

Barbi E, Marzuillo P, Neri E, Naviglio S, Krauss BS (2017) Fever in Children: Pearls and Pitfalls. Children
(Basel) 4:81.

Bartos M, Nusbaum MP (1997) Intercircuit control of motor pattern modulation by presynaptic
inhibition. J Neurosci 17:2247-2256.

Bartos M, Manor Y, Nadim F, Marder E, Nusbaum MP (1999) Coordination of fast and slow rhythmic
neuronal circuits. J Neurosci 19:6650-6660.

Beenhakker MP, Nusbaum MP (2004) Mechanosensory activation of a motor circuit by coactivation of
two projection neurons. J Neurosci 24:6741-6750.

Beenhakker MP, Blitz DM, Nusbaum MP (2004) Long-lasting activation of rhythmic neuronal activity by a
novel mechanosensory system in the crustacean stomatogastric nervous system. J Neurophysiol
91:78-91.

Beenhakker MP, Kirby MS, Nusbaum MP (2007) Mechanosensory gating of proprioceptor input to
modulatory projection neurons. J Neurosci 27:14308-14316.

Berens P (2009) CircStat: A MATLAB Toolbox for Circular Statistics. Journal of Statistical Software; Vol 1,
Issue 10 (2009).

Beverly M, Anbil S, Sengupta P (2011) Degeneracy and neuromodulation among thermosensory neurons
contribute to robust thermosensory behaviors in *Caenorhabditis elegans*. J Neurosci 31:11718-
11727.

Blitz DM (2017) Circuit feedback increases activity level of a circuit input through interactions with
intrinsic properties. J Neurophysiol 118:949-963.

Blitz DM, Nusbaum MP (2008) State-dependent presynaptic inhibition regulates central pattern
generator feedback to descending inputs. J Neurosci 28:9564-9574.

Blitz DM, Nusbaum MP (2012) Modulation of circuit feedback specifies motor circuit output. J Neurosci
32:9182-9193.

Blitz DM, Beenhakker MP, Nusbaum MP (2004) Different sensory systems share projection neurons but
elicit distinct motor patterns. J Neurosci 24:11381-11390.

Blitz DM, Christie AE, Cook AP, Dickinson PS, Nusbaum MP (2019) Similarities and differences in circuit
responses to applied Gly(1)-SIFamide and peptidergic (Gly(1)-SIFamide) neuron stimulation. J
Neurophysiol 121:950-972.

Blitz DM, Christie AE, Coleman MJ, Norris BJ, Marder E, Nusbaum MP (1999) Different proctolin neurons
elicit distinct motor patterns from a multifunctional neuronal network. J Neurosci 19:5449-5463.

Bucher D, Prinz AA, Marder E (2005) Animal-to-Animal Variability in Motor Pattern Production in Adults
and during Growth. The Journal of Neuroscience 25:1611-1619.

Burrows M (1989) Effects of temperature on a central synapse between identified motor neurons in the
locust. Journal of Comparative Physiology A 165:687-695.

Cao XJ, Oertel D (2005) Temperature affects voltage-sensitive conductances differentially in octopus
cells of the mammalian cochlear nucleus. J Neurophysiol 94:821-832.

Caplan JS, Williams AH, Marder E (2014) Many parameter sets in a multicompartment model oscillator
are robust to temperature perturbations. J Neurosci 34:4963-4975.

Chaffee RRJ, Roberts JC (1971) Temperature Acclimation in Birds and Mammals. Annual Review of
Physiology 33:155-202.

Cheshire WP (2016) Thermoregulatory disorders and illness related to heat and cold stress. Autonomic
Neuroscience 196:91-104.

Christie AE, Stein W, Quinlan JE, Beenhakker MP, Marder E, Nusbaum MP (2004) Actions of a
histaminergic/peptidergic projection neuron on rhythmic motor patterns in the stomatogastric
nervous system of the crab *Cancer borealis*. J Comp Neurol 469:153-169.
Coleman MJ, Nusbaum MP (1994) Functional consequences of compartmentalization of synaptic input. J
Neurosci 14:6544-6552.
Coleman MJ, Meyrand P, Nusbaum MP (1995) A switch between two modes of synaptic transmission
mediated by presynaptic inhibition. Nature 378:502-505.
Colgin LL (2011) Oscillations and hippocampal–prefrontal synchrony. Current Opinion in Neurobiology
21:467-474.
de Hemptinne C, Ryapolova-Webb ES, Air EL, Garcia PA, Miller KJ, Ojemann JG, Ostrem JL, Galifianakis
NB, Starr PA (2013) Exaggerated phase-amplitude coupling in the primary motor cortex in
Parkinson disease. Proc Natl Acad Sci U S A 110:4780-4785.
DeLong ND, Kirby MS, Blitz DM, Nusbaum MP (2009) Parallel regulation of a modulator-activated
current via distinct dynamics underlies comodulation of motor circuit output. J Neurosci
29:12355-12367.
Dickinson PS, Kurland SC, Qu X, Parker BO, Sreekrishnan A, Kwiatkowski MA, Williams AH, Ysasi AB,
Christie AE (2015) Distinct or shared actions of peptide family isoforms: II. Multiple pyrokinins
exert similar effects in the lobster stomatogastric nervous system. J Exp Biol 218:2905-2917.
Diehl F, White RS, Stein W, Nusbaum MP (2013) Motor circuit-specific burst patterns drive different
muscle and behavior patterns. J Neurosci 33:12013-12029.
Du Y, Lu Q, Wang R (2010) Using interspike intervals to quantify noise effects on spike trains in
temperature encoding neurons. Cognitive Neurodynamics 4:199-206.
Fohlmeister JF, Cohen ED, Newman EA (2010) Mechanisms and distribution of ion channels in retinal
ganglion cells: using temperature as an independent variable. J Neurophysiol 103:1357-1374.
Fonseca PJ, Correia T (2007) Effects of temperature on tuning of the auditory pathway in the cicada
Tettigetta josei (*Hemiptera, Tibicinidae*). J Exp Biol 210:1834-1845.
Foster JA, Robertson RM (1992) Temperature Dependency of Wing-Beat Frequency in Intact and
Deafferented Locusts. Journal of Experimental Biology 162:295-312.
Franz A, Ronacher B (2002) Temperature dependence of temporal resolution in an insect nervous
system. Journal of Comparative Physiology A 188:261-271.
Fujisawa S, Buzsáki G (2011) A 4 Hz oscillation adaptively synchronizes prefrontal, VTA, and hippocampal
activities. Neuron 72:153-165.
Garrity PA, Goodman MB, Samuel AD, Sengupta P (2010) Running hot and cold: behavioral strategies,
neural circuits, and the molecular machinery for thermotaxis in *C. elegans* and *Drosophila*.
Genes & Development 24:2365-2382.
Gordon JA (2011) Oscillations and hippocampal-prefrontal synchrony. Current opinion in neurobiology
21:486-491.
Gutierrez GJ, Grashow RG (2009) *Cancer borealis* stomatogastric nervous system dissection. J Vis
Exp:https://www.ncbi.nlm.nih.gov/pubmed/19308017.
Gutierrez GJ, O'Leary T, Marder E (2013) Multiple mechanisms switch an electrically coupled,
synaptically inhibited neuron between competing rhythmic oscillators. Neuron 77:845-858.
Haddad SA, Marder E (2018) Circuit Robustness to Temperature Perturbation Is Altered by
Neuromodulators. Neuron 100:609-623 e603.
Haley JA, Hampton D, Marder E (2018) Two central pattern generators from the crab, Cancer borealis,
respond robustly and differentially to extreme extracellular pH. Elife 7.
Hamood AW, Marder E (2015) Consequences of acute and long-term removal of neuromodulatory input
on the episodic gastric rhythm of the crab Cancer borealis. J Neurophysiol 114:1677-1692.
Harris AZ, Gordon JA (2015) Long-range neural synchrony in behavior. Annu Rev Neurosci 38:171-194.

Hazel JR, Prosser CL (1974) Molecular mechanisms of temperature compensation in poikilotherms.
Physiol Rev 54:620-677.
He LS, Rue MCP, Morozova EO, Powell DJ, James EJ, Kar M, Marder E (2020) Rapid adaptation to
elevated extracellular potassium in the pyloric circuit of the crab, *Cancer borealis*. Journal of
Neurophysiology 123:2075-2089.
Heinzel HG, Weimann JM, Marder E (1993) The behavioral repertoire of the gastric mill in the crab,
Cancer pagurus: an *in situ* endoscopic and electrophysiological examination. J Neurosci 13:1793-
1803.
Heitler WJ, Edwards DH (1998) Effect of temperature on a voltage-sensitive electrical synapse in
crayfish. The Journal of Experimental Biology 201:503.
Hori A, Minato K, Kobayashi S (1999) Warming-activated channels of warm-sensitive neurons in rat
hypothalamic slices. Neuroscience Letters 275:93-96.
Jacobson TK, Howe MD, Schmidt B, Hinman JR, Escabi MA, Markus EJ (2013) Hippocampal theta,
gamma, and theta-gamma coupling: effects of aging, environmental change, and cholinergic
activation. J Neurophysiol 109:1852-1865.
Janssen R (1992) Thermal influences on nervous system function. Neuroscience & Biobehavioral Reviews
16:399-413.
Kang K, Panzano VC, Chang EC, Ni L, Dainis AM, Jenkins AM, Regna K, Muskavitch MAT, Garrity PA (2012)
Modulation of TRPA1 thermal sensitivity enables sensory discrimination in *Drosophila*. Nature
481:76-80.
Karalis N, Dejean C, Chaudun F, Khoder S, Rozeske RR, Wurtz H, Bagur S, Benchenane K, Sirota A, Courtin
J, Herry C (2016) 4-Hz oscillations synchronize prefrontal–amygdala circuits during fear behavior.
Nature Neuroscience 19:605-612.
Katz PS, Harris-Warrick RM (1991) Recruitment of crab gastric mill neurons into the pyloric motor
pattern by mechanosensory afferent stimulation. J Neurophysiol 65:1442-1451.
Kirihara K, Rissling AJ, Swerdlow NR, Braff DL, Light GA (2012) Hierarchical organization of gamma and
theta oscillatory dynamics in schizophrenia. Biol Psychiatry 71:873-880.
Klockner U, Schiefer A, Isenberg G (1990) L-type Ca-channels: similar Q10 of Ca-, Ba- and Na-
conductance points to the importance of ion-channel interaction. Pflugers Arch 415:638-641.
Kushinsky D, Morozova EO, Marder E (2019) In vivo effects of temperature on the heart and pyloric
rhythms in the crab *Cancer borealis*. J Exp Biol 222.
Lisman J (2005) The theta/gamma discrete phase code occuring during the hippocampal phase
precession may be a more general brain coding scheme. Hippocampus 15:913-922.
Lisman J, Buzsáki G (2008) A Neural Coding Scheme Formed by the Combined Function of Gamma and
Theta Oscillations. Schizophrenia Bulletin 34:974-980.
Lisman JE, Jensen O (2013) The θ-γ neural code. Neuron 77:1002-1016.
Long MA, Fee MS (2008) Using temperature to analyse temporal dynamics in the songbird motor
pathway. Nature 456:189-194.
Mank A, van Zanten HA, Meyer MP, Pauws S, Lopriore E, Te Pas AB (2016) Hypothermia in Preterm
Infants in the First Hours after Birth: Occurrence, Course and Risk Factors. PLoS One
11:e0164817-e0164817.
Marder E, Bucher D (2007) Understanding circuit dynamics using the stomatogastric nervous system of
lobsters and crabs. Annu Rev Physiol 69:291-316.
Marder E, Manor Y, Nadim F, Bartos M, Nusbaum MP (1998) Frequency control of a slow oscillatory
network by a fast rhythmic input: pyloric to gastric mill interactions in the crab stomatogastric
nervous system. Ann N Y Acad Sci 860:226-238.

Marder E, Haddad SA, Goeritz ML, Rosenbaum P, Kispersky T (2015) How can motor systems retain
performance over a wide temperature range? Lessons from the crustacean stomatogastric
nervous system. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 201:851-856.
Maynard DM, Selverston AI (1975) Organization of the stomatogastric ganglion of the spiny lobster.
Journal of comparative physiology 100:161-182.
Meyrand P, Simmers J, Moulins M (1994) Dynamic construction of a neural network from multiple
pattern generators in the lobster stomatogastric nervous system. J Neurosci 14:630-644.
Montgomery SM, Buzsaki G (2007) Gamma oscillations dynamically couple hippocampal CA3 and CA1
regions during memory task performance. Proc Natl Acad Sci U S A 104:14495-14500.
Moran LV, Hong LE (2011) High vs low frequency neural oscillations in schizophrenia. Schizophr Bull
37:659-663.
Moran MM, Xu H, Clapham DE (2004) TRP ion channels in the nervous system. Curr Opin Neurobiol
14:362-369.
Nadim F, Manor Y, Nusbaum MP, Marder E (1998) Frequency regulation of a slow rhythm by a fast
periodic input. J Neurosci 18:5053-5067.
Norris BJ, Coleman MJ, Nusbaum MP (1994) Recruitment of a projection neuron determines gastric mill
motor pattern selection in the stomatogastric nervous system of the crab, *Cancer borealis*. J
Neurophysiol 72:1451-1463.
Norris BJ, Coleman MJ, Nusbaum MP (1996) Pyloric motor pattern modification by a newly identified
projection neuron in the crab stomatogastric nervous system. J Neurophysiol 75:97-108.
O'Leary T, Marder E (2016) Temperature-Robust Neural Function from Activity-Dependent Ion Channel
Regulation. Curr Biol 26:2935-2941.
Prinz AA, Bucher D, Marder E (2004) Similar network activity from disparate circuit parameters. Nat
Neurosci 7:1345-1352.
Prosser CL, Nelson DO (1981) The Role of Nervous Systems in Temperature Adaptation of Poikilotherms.
Annual Review of Physiology 43:281-300.
Ramot D, MacInnis BL, Goodman MB (2008) Bidirectional temperature-sensing by a single
thermosensory neuron in *C. elegans*. Nature Neuroscience 11:908-915.
Rinberg A, Taylor AL, Marder E (2013) The effects of temperature on the stability of a neuronal
oscillator. PLoS Comput Biol 9:e1002857.
Robertson RM, Money TGA (2012) Temperature and neuronal circuit function: compensation, tuning
and tolerance. Current Opinion in Neurobiology 22:724-734.
Roemschied FA, Eberhard MJB, Schleimer J-H, Ronacher B, Schreiber S (2014) Cell-intrinsic mechanisms
of temperature compensation in a grasshopper sensory receptor neuron. eLife 3:e02078.
Rojas-Líbano D, Frederick DE, Egaña JI, Kay LM (2014) The olfactory bulb theta rhythm follows all
frequencies of diaphragmatic respiration in the freely behaving rat. Front Behav Neurosci 8:214-
214.
Saideman SR, Blitz DM, Nusbaum MP (2007a) Convergent motor patterns from divergent circuits. J
Neurosci 27:6664-6674.
Saideman SR, Ma M, Kutz-Naber KK, Cook A, Torfs P, Schoofs L, Li L, Nusbaum MP (2007b) Modulation of
rhythmic motor activity by pyrokinin peptides. J Neurophysiol 97:579-595.
Salimpour Y, Anderson WS (2019) Cross-Frequency Coupling Based Neuromodulation for Treating
Neurological Disorders. Frontiers in Neuroscience 13.
Scheffer-Teixeira R, Tort ABL (2016) On cross-frequency phase-phase coupling between theta and
gamma oscillations in the hippocampus. eLife 5:e20515.
Schleimer J-H, Schreiber S (2016) Homeostasis: How Neurons Achieve Temperature Invariance. Current
Biology 26:R1141-R1143.

Shruti S, Schulz DJ, Lett KM, Marder E (2014) Electrical coupling and innexin expression in the
1000   stomatogastric ganglion of the crab *Cancer borealis*. Journal of Neurophysiology 112:2946-2958.
1001   Soofi W, Goeritz ML, Kispersky TJ, Prinz AA, Marder E, Stein W (2014) Phase maintenance in a rhythmic
1002   motor pattern during temperature changes *in vivo*. J Neurophysiol 111:2603-2613.
1003   Städele C, Heigele S, Stein W (2015) Neuromodulation to the Rescue: Compensation of Temperature-
1004   Induced Breakdown of Rhythmic Motor Patterns via Extrinsic Neuromodulatory Input. PLoS Biol
1005   13:e1002265.
1006   Stehlik LL, MacKenzie Jr. CL, Morse WW (1991) Distribution and abundance of four brachyuran crabs on
1007   the northwest atlantic shelf. Fishery Bulletin, US:473-492.
1008   Stein W, DeLong ND, Wood DE, Nusbaum MP (2007) Divergent co-transmitter actions underlie motor
1009   pattern activation by a modulatory projection neuron. Eur J Neurosci 26:1148-1165.
1010   Stiebler IB, Narins PM (1990) Temperature-dependence of auditory nerve response properties in the
1011   frog. Hearing Research 46:63-81.
1012   Tamura M, Spellman TJ, Rosen AM, Gogos JA, Gordon JA (2017) Hippocampal-prefrontal theta-gamma
1013   coupling during performance of a spatial working memory task. Nature Communications 8:2182.
1014   Tang LS, Taylor AL, Rinberg A, Marder E (2012) Robustness of a rhythmic circuit to short- and long-term
1015   temperature changes. J Neurosci 32:10075-10085.
1016   Tang LS, Goeritz ML, Caplan JS, Taylor AL, Fisek M, Marder E (2010) Precise Temperature Compensation
1017   of Phase in a Rhythmic Motor Pattern. PLOS Biology 8:e1000469.
1018   Tass P, Rosenblum MG, Weule J, Kurths J, Pikovsky A, Volkmann J, Schnitzler A, Freund HJ (1998)
1019   Detection of *n:m* Phase Locking from Noisy Data: Application to Magnetoencephalography.
1020   Physical Review Letters 81:3291-3294.
1021   Weimann JM, Marder E (1994) Switching neurons are integral members of multiple oscillatory networks.
1022   Curr Biol 4:896-902.
1023   Weimann JM, Meyrand P, Marder E (1991) Neurons that form multiple pattern generators:
1024   identification and multiple activity patterns of gastric/pyloric neurons in the crab stomatogastric
1025   system. J Neurophysiol 65:111-122.
1026   Weimann JM, Marder E, Evans B, Calabrese RL (1993) The effects of SDRNFLRFamide and
1027   TNRNFLRFamide on the motor patterns of the stomatogastric ganglion of the crab *Cancer
1028   borealis*. J Exp Biol 181:1-26.
1029   Weimann JM, Skiebe P, Heinzel HG, Soto C, Kopell N, Jorge-Rivera JC, Marder E (1997) Modulation of
1030   oscillator interactions in the crab stomatogastric ganglion by crustacean cardioactive peptide. J
1031   Neurosci 17:1748-1760.
1032   White RS, Nusbaum MP (2011) The same core rhythm generator underlies different rhythmic motor
1033   patterns. J Neurosci 31:11484-11494.
1034   Wood DE, Stein W, Nusbaum MP (2000) Projection neurons with shared cotransmitters elicit different
1035   motor patterns from the same neural circuit. J Neurosci 20:8943-8953.
1036   Xu H, Robertson RM (1994) Effects of temperature on properties of flight neurons in the locust. Journal
1037   of Comparative Physiology A 175:193-202.
1038   Xu H, Robertson RM (1996) Neural parameters contributing to temperature compensation in the flight
1039   CPG of the locust, *Locusta migratoria*. Brain Research 734:213-222.
1040   Yang F, Zheng J (2014) High temperature sensitivity is intrinsic to voltage-gated potassium channels.
1041   eLife 3:e03255.
1042   Zečević D, Levitan H (1980) Temperature acclimation: effects on membrane physiology of an identified
1043   snail neuron. American Journal of Physiology-Cell Physiology 239:C47-C57.
1044   Zečević D, Levitan H, Pašić M (1985) Temperature Acclimation: Influence on Transient Outward and
1045   Inward Currents in an Identified Neurone of *Helix pomatia*. Journal of Experimental Biology
1046   117:369.

1047   Zheng C, Bieri KW, Hsiao YT, Colgin LL (2016) Spatial Sequence Coding Differs during Slow and Fast
1048   Gamma Rhythms in the Hippocampus. Neuron 89:398-408.

1049