MAGIC: Mosaic Analysis by gRNA-Induced Crossing-over

Sarah E. Allen[1,3], Gabriel T. Koreman[1,2,3], Ankita Sarkar[1,2,3], Bei Wang[1,2], Mariana F. Wolfner[1,4],

and Chun Han[1,2,4]

[1]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA

[2]Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, USA

[3]These authors contributed equally to this work, listed alphabetically

[4]For correspondence: mariana.wolfner@cornell.edu (MFW) and chun.han@cornell.edu (CH)

Abstract:

Mosaic animals have provided the platform for many fundamental discoveries in developmental biology, cell biology, and other fields. Techniques to produce mosaic animals by mitotic recombination have been extensively developed in *Drosophila melanogaster* but are less common for other laboratory organisms. Here, we report mosaic analysis by gRNA-induced crossing-over (MAGIC), a new technique for generating mosaic animals based on DNA double-strand breaks produced by CRISPR/Cas9. MAGIC efficiently produces mosaic clones in both somatic tissues and the germline of *Drosophila*. Further, by developing a MAGIC toolkit for one chromosome arm, we demonstrate the method's application in characterizing gene function in neural development and in generating fluorescently marked clones in wild-derived *Drosophila* strains. Eliminating the need to introduce recombinase-recognition sites in the genome, this simple and versatile system simplifies mosaic analysis in *Drosophila* and can be applied in any organism that is compatible with CRISPR/Cas9.

Keywords: *MAGIC, mosaic analysis, CRISPR/Cas9, gRNA, clonal analysis, germline, imaginal disc, da neurons, Drosophila dominant female sterility,*

Introduction:

Mosaic animals contain genetically distinct populations of cells that have arisen from one zygote. Mosaic animals have historically played important roles in the study of pleiotropic genes, developmental timing, cell lineage, neural wiring, and other complex biological processes. Given its genetic tractability, *Drosophila* has been a major system for generating and studying such mosaics (Perrimon, 1998), which have led to important discoveries such as developmental compartments (Garcia-Bellido et al., 1976), cell autonomy (Heitzler and Simpson, 1991), and maternal effects of zygotic lethal genes (Perrimon et al., 1989). Mosaic (also called clonal) analysis is currently used to study tumor suppressors (Feng and Irvine, 2007), signaling pathways (Huang et al., 2014), sleep-wake behaviors (Chung et al., 2017), cell fates (Hu and Jasper, 2019), and neuronal lineages (Lee, 2014), among other biological processes.

The earliest mosaic analyses relied on spontaneous mitotic recombination (Stern, 1936), rare events in which a DNA double-strand break (DSB) during the G_2 phase of the cell cycle is repaired by homologous recombination, resulting in the reciprocal exchange of chromosomal arms between homologous chromosomes distal to the site of the DNA crossover (reviewed in Griffin et al., 2014). Ionizing radiation, such as X-rays (Muller, 1928), cause DSBs and thus were later used in *Drosophila* to increase the baseline level of mitotic recombination (Xu and Rubin, 2012). However, ionizing radiation breaks genomic DNA at random locations and is associated with a high degree of lethality.

To overcome these limitations, the yeast Flippase (Flp)/FRT system was introduced into *Drosophila* to mediate site-specific recombination at FRT sites (Golic and Lindquist, 1989; Xu and Rubin, 1993), enabling the development of an ever-expanding toolbox with enhanced power and flexibility for clonal analysis (Griffin et al., 2009; Harrison and Perrimon, 1993; Lee and Luo, 1999; Xu and Rubin, 1993; Yu et al., 2009). This system requires that both homologous chromosomes carry FRT sites at the same position proximal (relative to the centromere) to the gene of interest, and an independent marker on one of the homologs to allow

visualization of the genetically distinct clones (Xu and Rubin, 1993). For clonal analysis in the

Drosophila germline, a dominant female sterility (DFS) *ovo^{D1}* transgene was combined with

Flp/FRT methods, allowing production of, and selection for, germline clones homozygous for a

mutation of interest in a heterozygous mother (Chou and Perrimon, 1992; Perrimon and Gans,

1983). In this "Flp-DFS" technique, egg production from *ovo^{D1}*-containing heterozygous and

homozygous germline cells is blocked, resulting in progeny derived exclusively from germline

clones lacking *ovo^{D1}* that were generated by mitotic recombination (Chou et al., 1993). Clonal

analysis based on somatic recombination has also been achieved in mice using the Cre-LoxP

system and the reconstitution of fluorescent protein genes as markers (Henner et al., 2013;

Zong et al., 2005). Despite these successes, site-specific recombination systems have not been

widely used for mitotic recombination in model animals beyond *Drosophila melanogaster* due to

the challenging task of introducing recombinase-recognition sites into centromere-proximal

regions for every chromosome.

Given the power of mosaic animals in biological research, it would be useful to have a more

general approach for inducing interhomolog mitotic recombination in any organism,

circumventing the challenges just mentioned. The CRISPR/Cas9 system has great potential for

extending clonal analysis, because it can create targeted DSBs in the genomic DNA of a wide

array of organisms (Sander and Keith Joung, 2014). This binary system requires only the Cas9

endonuclease and a guide RNA (gRNA) that specifies the DNA target site (Jinek et al., 2012),

both of which can be introduced into the cell independently of the location of the target site.

CRISPR/Cas9-induced DSBs can be repaired either by non-homologous end joining (NHEJ) or

homology-directed repair (HDR). So far, most CRISPR/Cas9 applications in animals have been

focused on NHEJ-mediated mutagenesis and HDR-mediated gene replacement (Salsman and

Dellaire, 2017), and CRISPR/Cas9-induced mitotic recombination has not been demonstrated in

multicellular organisms. Here, we report mosaic analysis by gRNA-induced crossing-over

(MAGIC), a novel CRISPR/Cas9-based technique for clonal analysis. This method can be used

to generate mosaic clones in both the *Drosophila* soma and germline. Based on this method, we

built a convenient toolkit to generate and label mosaic clones for genes located on chromosome

arm 2L. We demonstrate the success of our toolkit for clonal analysis in the soma and the

germline and show its applications in analyzing gene functions in neuronal dendrite

development. Lastly, we also demonstrate that MAGIC can be used successfully with unmarked

wild-derived strains, indicating that this method can be extended to organisms beyond

Drosophila.

Results:

Rationale for MAGIC

MAGIC relies on the action of gRNA/Cas9 in a proliferating cell during G_2 phase to generate a

DSB at a specific position on one chromatid of a homologous pair (Figure 1A). The DSB can

induce a crossover between this chromatid and a chromatid from the homologous chromosome,

resulting in exchange of chromosome segments between the two chromatids at the location of

the DSB. During the subsequent mitotic segregation of chromosomes, a 50% chance exists for

identical distal chromosome segments to sort into the same daughter cells, generating "twin

spots", which contain two genetically distinct populations of cells homozygous for the

chromosome segment distal to the exchange.

Using CRISPR-induced crossover to generate clones in the Drosophila ***soma and***

germline

For our initial tests of the ability of MAGIC to generate mosaic clones in somatic tissues, we

used ubiquitously expressed gRNAs to induce DSBs at the *gnu* locus and a ubiquitous

fluorescent marker, *His2Av-GFP* (Clarkson and Saint, 1999), to trace clones (Figure 1B). Both

His2Av-GFP and *gnu* are located on the left arm of chromosome 3 (3L), and *His2Av-GFP* is

distal to *gnu*. We chose *gnu* as our gRNA target because we have already made an efficient

gRNA-gnu line for other purposes (to be published elsewhere); furthermore, this gene is only

required maternally for embryonic development (Freeman et al., 1966), so mutations of *gnu* are

not expected to affect the viability or growth of somatic cells. We induced clones using three

different Cas9 transgenes, each of which is expressed in the developing wing disc under the

control of a different enhancer. With all three Cas9s, we observed twin spots consisting of bright

His2Av-GFP homozygous clones abutting GFP-negative clones in the midst of *His2Av-GFP/+*

heterozygous cells (Figures 1C-1E) in every imaginal disc examined, demonstrating the

feasibility of MAGIC for generating somatic mosaics.

Given that CRISPR/Cas9 is active in both the soma and the germline of *Drosophila*

(Gratz et al., 2013; Port et al., 2014; Ren et al., 2013), we next tested for MAGIC clone induction

in the germline by using the DFS technique and the germline-specific *nos-Cas9* (Port et al.,

2014) (Figure S1A). We used an *ovo^{D1}* transgene located on chromosome arm 2R and induced

DSBs at the *Rab3* locus, which is located on the same arm proximal to the location of *ovo^{D1}*

(Figure 1F). Since *Rab3* is a non-essential gene expressed only in neurons (Chen et al., 2015),

its disruption in the female germline should affect neither egg production nor embryonic

development of the progeny. Due to the dominant effect of *ovo^{D1}*, restoration of egg production

can only result from mitotic recombination proximal to *ovo^{D1}* (e.g. at *Rab3* in this case), followed

by generation of *ovo^{D1}* negative clones. As expected, control females that contained *ovo^{D1}* and

nos-Cas9, but not *gRNA-Rab3*, did not produce any eggs. In contrast, most females carrying all

three components produced 20-90 eggs each (Figure 1G), many of which hatched into larvae

(Figure 1H), suggesting successful mitotic recombination.

The results above together show that, like the Flp/FRT system, MAGIC is an effective

approach for generating homozygous clones via mitotic recombination in both *Drosophila* soma

and germline.

A toolkit for generating labeled clones for genes on chromosome arm 2L

Towards making MAGIC a general approach for analyzing *Drosophila* genes, we built a toolkit

for genes located on chromosome arm 2L as a proof-of-principle. We designed transgenic

constructs that each integrate two features simultaneously: ubiquitously-expressed gRNAs that

target a pericentromeric region and a ubiquitously-expressed marker for labeling clones. The

constructs were inserted into a distal position of 2L. When used together with an unmodified 2nd

chromosome, they label clones homozygous for nearly the entirety of the unmodified arm either

negatively or positively. Negative labeling in the nMAGIC option is achieved by expressing a

nuclear blue fluorescent protein (nBFP) reporter such that clones homozygous for the

unmodified arm lose nBFP expression (Figure 2A). Positive labeling with pMAGIC utilizes a

Gal80 marker (Lee and Luo, 1999), which suppresses Gal4-driven expression of a fluorescent

reporter (Figure 2B). Therefore, only the cells that lose Gal80 transgene will be fluorescently

labeled, similarly to mosaic analysis with a repressible cell marker (MARCM) (Lee and Luo,

1999).

To identify appropriate gRNA target sites, we surveyed the pericentromeric sequences

of 2L for sequences that met three criteria: (1) being reasonably conserved so that DSBs can be

induced in most *Drosophila* strains; (2) not functionally critical and being distant from essential

sequences so that indel mutations in nearby regions would not disrupt important biological

processes; and (3) unique in the genome, so as to avoid off-target effects. Therefore, for each

MAGIC construct, we chose a pair of non-repeat gRNA target sequences in an intergenic region

to enhance the chance of DSBs. The two gRNA target sequences are closely-linked to reduce

the risk of large deletions (Figure 2C). In addition, we preferentially chose sequences that are

conserved among closely related *Drosophila* species (*D. melanogaster*, *D. simulans*, and *D.*

sechellia) but not in more distant species. Considering the varying efficiencies of different gRNA

target sequences, we selected three pairs of gRNAs targeting three chromosomal locations

(40D2, 40D4, and 40E1) and tested their ability to produce clones in wing discs, neurons, and

the germline.

Clones were induced in a specific tissue by a Cas9 transgene that is expressed in

precursor cells of that tissue. We used *hh-Cas9* (Poe et al., 2019) for nMAGIC in the wing

imaginal disc (Figure 2D), *SOP-Cas9* (Poe et al., 2019) for pMAGIC in larval class IV dendritic

arborization (C4da) sensory neurons (Figure 2F), and *nos-Cas9* for the female germline (Figure

S1B). gRNAs targeting 40D2 consistently performed the best in generating clones in wing discs

and C4da neurons (Figures 2E and 2G) and appear to be the most efficient in the germline,

even though the differences in the germline were not statistically significant (Figures 2H and 2I).

Although the overall efficiencies of *gRNA-40D2* and *gRNA-40D4* in inducing clones in da

neurons are similar (Figure 2G and Figure S2A), *gRNA-40D4* induced more clones in a different

type of neuron (class III). The fact that the three gRNA pairs showed similar trends in their ability

to induce clones across different tissues suggests that an efficient gRNA construct for one

tissue will likely perform well in other tissues also. These results indicate that we have created

an efficient MAGIC toolkit for genes located on chromosome arm 2L. Analogous toolkits could

easily be made for any other chromosome arm, using the same methods.

Clonal analysis of neuronal dendrite development

To evaluate the utility of our MAGIC toolkit for characterizing gene function at the single-cell

level, we combined the pMAGIC line *gRNA-40D2(Gal80)* with mutations on 2L that affect

dendrite morphogenesis in C4da neurons by disrupting vesicular trafficking. We first used two

genes, *Secretory 5* (*Sec5*) (Peng et al., 2015) and *Rab5* (Satoh et al., 2008), that have been

shown to be required for dendrite growth. We observed dendrite reduction in C4da clones

carrying homozygous mutations in these genes (Figures 3A and 3B), recapitulating previously

published results using MARCM with the same mutants (Peng et al., 2015; Satoh et al., 2008).

A third gene, *Syntaxin 5* (*Syx5*), was identified in our unpublished RNAi screens. Clones

carrying a null mutation of *Syx5* produced the most dramatic dendrite reduction, with almost all

terminal dendrites eliminated (Figure 3C), consistent with the expected role of Syx5 in ER to

Golgi vesicle trafficking (Hardwick and Pelham, 1992). Therefore, our MAGIC reagents for 2L

can be used to characterize gene functions in single cells with a power analogous to that of

MARCM but with a much simpler system.

Generation of clones by MAGIC in fly lines with wild-derived genomes

A substantial advantage of MAGIC compared to Flp/FRT-based mitotic recombination systems

is that MAGIC does not require prior genetic modification of the chromosome arm to be tested.

It therefore has the potential to be applied to fly strains with wild-derived genomes, and even

other organisms. To test the applicability of MAGIC to unmarked strains with wild-derived

genomes, we crossed *gRNA-40D2(nBFP); hh-Cas9* to five randomly chosen lines from the

Drosophila Genetic Reference Panel (DGRP) (MacKay et al., 2012), a set of standard strains

established from flies captured in the wild. In all cases, we observed efficient clone induction in

wing imaginal discs (Figures 4A-4E), demonstrating the potential of MAGIC for clonal analysis

of the function of natural alleles residing on wild-derived chromosomes.

Discussion:

We present here a new technique we name MAGIC (mosaic analysis by gRNA-induced

crossing-over) for clonal analysis based on CRISPR-induced mitotic recombination. We show

that MAGIC is capable of producing efficiently mosaic tissues in both the *Drosophila* soma and

germline, using gRNAs targeting various chromosomal locations. Integrated gRNA-marker

constructs enable both positive- and negative-labeling of homozygous clones. As demonstrated

by our 2L toolkit, MAGIC is simple and effective to use; similar MAGIC reagents can be

generated easily for all other chromosome arms to allow for genome-wide characterization of

gene functions.

Although conventional Flp/FRT-based techniques have been widely and successfully

used in *Drosophila* for similar clonal analyses (Griffin et al., 2014), MAGIC has two major

advantages. The first is the simplicity of MAGIC assays: This system eliminates the requirement

for genetic modification of the test chromosome; furthermore, integrating gRNAs and genetic

markers into one transgenic construct further reduces the number of necessary genetic

components. Therefore, gRNA-marker transgenes can be combined with existing mutant

libraries to perform MAGIC with very little additional effort. The second advantage of MAGIC is

that mitotic recombination is not limited by the available FRT insertion sites. While introducing

FRT to a specific pericentromeric region is very difficult and historically required labor-intensive

genetic screens, CRISPR/Cas9 can induce DNA DSBs and subsequent crossover at specific

pericentromeric sequences with ease. Therefore, MAGIC opens doors for clonal analysis of

genes that were previously impossible to study using existing FRT sites, such as those on the

fourth chromosome and the ones near centromeres. In addition, many *Drosophila* mutations are

associated with transgenic constructs containing FRT (Thibault et al., 2004), making analyses

complicated when using Flp/FRT-based techniques. In contrast, MAGIC should be compatible

with all of these gene disruption lines.

The unique mechanism of MAGIC requires three considerations for successful

applications. First, our results suggest that the gRNA target sequence strongly influences the

efficiency of clone induction, likely by affecting the frequency of DNA DSBs in premitotic cells.

Therefore, for clonal analysis of a specific chromosomal arm, it is beneficial to compare a few

candidate gRNA targets and select the most effective one. Second, because perfect DSB repair

will recreate the gRNA target site and allow for one more round of Cas9 cutting, most cells that

have expressed Cas9 in their lineages are expected to eventually harbor indel mutations that

disrupt the gRNA target site, regardless of whether or not the DSBs have led to mitotic

recombination. However, this caveat can be mitigated by choosing gRNA sites in non-critical

sequences, which can be validated by crossing gRNA lines to a ubiquitous Cas9 or by

comparing gRNA-induced control clones to wildtype cells. Lastly, since only DNA DSBs in the

G_2 phase can lead to clone generation, the timing of Cas9 action is expected to be critical for

MAGIC. For the cell type in question, an ideal Cas9 should be expressed in the precursor cells,

as too early expression can mutate gRNA target sites prematurely and too late expression will

lead to unproductive DSBs.

Perhaps the most exciting aspect of MAGIC is its potential for use with wild-derived

Drosophila strains and in organisms beyond *Drosophila*. DGRP wild-derived strains have played

important roles in identification of natural alleles that are associated with certain phenotypic

variations (Clark et al., 1994; Grenier et al., 2015; Mackay and Huang, 2018; Shorter et al.,

2015). However, it has been difficult to investigate the effect of homozygosity for alleles within

these strains without being able to use available genetic tools (e.g. Gal4 drivers and fluorescent

markers) in *Drosophila*. By combining MAGIC with the DGRP, it is now possible to validate

causal effects of specific natural alleles in cellular or developmental processes in a tissue-

specific manner. The DGRP can also be used in MAGIC-based genetic screens to identify

natural alleles that, when made homozygous, can cause or modify certain phenotypes.

Importantly, MAGIC can in theory be utilized in a wide array of organisms that are compatible

with CRISPR/Cas9 (Harrison et al., 2014). In model systems that allow for transgenesis of

gRNA-marker constructs, such as mouse, zebrafish, and *Xenopus*, Cas9 can be introduced by

injection or virus transduction to further simplify genetic manipulations. Therefore, the flexibility

and power of mosaic analysis that are familiar to the *Drosophila* research community are now in

reach of researchers who study organisms which have not, or have rarely, been amenable to

clonal analysis.

Materials and Methods:

Fly Stocks and Husbandry

See the Key Resource Table for details of fly stocks used in this study. Broadly, all fly lines were

either generated in the Han and Wolfner labs, or obtained from the Bloomington *Drosophila*

Stock Center or the *Drosophila* Genetic Reference Panel (MacKay et al., 2012). All flies were

grown on standard yeast-glucose medium, in a 12:12 light/dark cycle, at room temperature (22

$\pm$ 1°C, for the egg laying assay) or 25°C (for larval assays) unless otherwise noted. Virgin males

and females for mating experiments were aged for 3-5 days. Virgin females were aged on

yeasted food for germline clonal analysis.

To test germline clone induction, we combined *nos-Cas9* and *ovo^{D1}*, and then the gRNA

in two sequential crosses in schemes shown in Figure S1.

To visualize clones of C4da neurons, we used *ppk-Gal4 UAS-CD4-tdTom* (Figure 2) and

ppk-Gal4 UAS-MApHS (Figure 3, only the tdTom channel is shown).

Molecular Cloning

zk-Cas9: The entry vector pENTR221-ZK2 (Poe et al., 2017) and the destination vector pDEST-

APIC-Cas9 (Addgene 121657) were combined in a Gateway LR reaction to generate the

expression vector pAPIC2-ZK2-Cas9.

MAGIC gRNA-marker vectors: gRNA-marker vectors were constructed similarly to pAC-U63-

tgRNA-Rev (Addgene 112811, Poe et al., 2019) but have either a ubi-nBFP (in pAC-U63-

tgRNA-nlsBFP) or a ubi-Gal80 (in pAC-U63-tgRNA-Gal80) marker immediately after the U6 3'

flanking sequence. The markers contain a Ubi-p63E promoter, mTagBFP-NLS or Gal80 coding

sequence, and His2Av polyA sequence. The Ubi-p63E promoter was amplified from *Ubi-*

CasExpress genomic DNA using the oligonucleotides

TTAATGCGTATGCATTCTAGTggccatggcttgctgttcttcgcgttc and

TTGGATTATTctgcgggcagaaaatagagatgtggaaaattag. mTagBFP-NLS was synthesized as a

gBlock DNA fragment (Integrated DNA Technologies, Inc.). Gal80 coding sequence was PCR

amplified from pBPGAL80Uw-4 (Addgene 26235) using the oligonucleotides

aaaaaaaaatcaaaATGAGCGGTACCGATTACAACAAAAGGAGTAGTGTGAG and

GCCGACTGGCTTAGTTAattaattctagaTTAAAGCGAGTAGTGGGAGATGTTG. The His2Av

polyA sequence was PCR amplified from pDEST-APLO (Addgene 112805). DNA fragments

were assembled together using NEBuilder DNA Assembly (New England Biolabs Inc.).

gRNA expression vectors: For *gnu* and *Rab3*, gRNA target sequences were cloned into pAC-

U63-tgRNA-Rev as described (Poe et al., 2019). For gRNAs targeting 2L, gRNA target

sequences were cloned into pAC-U63-tgRNA-nlsBFP and pAC-U63-tgRNA-Gal80 using

NEBuilder DNA Assembly. In the gRNA-marker constructs, the tRNA between the first and

second gRNAs is a *Drosophila* glutamine tRNA

(cagcgcGGTTCCATGGTGTAATGGTTAGCACTCAGGACTCTGAATCCTGCGATCCGAGTTCA

AATCTCGGTGGAACCT) instead of a rice glycine tRNA.

Injections were carried out by Rainbow Transgenic Flies (Camarillo, CA 93012 USA) to

transform flies through φC31 integrase-mediated integration into attP docker sites.

pAPIC2-ZK2-Cas9 and gRNA-marker constructs were integrated into the *attP*^{VK00037} site on the

second chromosome and expression vectors containing gRNAs targeting *Rab3* or *gnu* were

integrated into the *attP*^{VK00027} site on the third chromosome. Transgenic insertions were

validated by genomic PCR or sequencing.

Identification of gRNA target sequence

gRNA target sequences for *Rab3* and *gnu* were identified as described previously (Poe et al.,

2019). Briefly, two gRNA prediction methods were used: sgRNA Scorer 2.0 (Chari et al., 2017)

(https://crispr.med.harvard.edu) and Benchling (www.benchling.com). Candidate target

sequences were those that obtained high on-target scores in both algorithms. CasFinder (Aach

et al., 2014) was used to identify and reject any sequences with more than one target site. Two

target sequences against coding exons for all splice isoforms were chosen for each targeted

gene. gRNA target sequences for 2L were identified by visually scanning through

pericentromeric sequences using UCSC Genome Browser (https://genome.ucsc.edu/) following

principles described in the Results section. The gRNA target sequences are listed in the table

below.

Gene	Target sequence 1	Target sequence 2
gnu	TTCGAATGTAAAAGCTTCGG	TTCCTGCCAACGCCTCCAGT
Rab3	GCCCACCGTGGAGACGAAGG	AGTGCGACATGGAGGACCAG
40D2	AGTCACCTGAAATAAGTCAG	GTTAGCCATCACAAGAACAG
40D4	GGGTTGTCTCCTGATATGGG	AACCGAACTGAACTCAACTG
40E1	CCGAATGATTTCATGTGAAG	TAGGGCAATTAAATATGTCA

Live Imaging of neurons

Live imaging was performed as previously described (Poe et al., 2017). Briefly, animals were reared at 25˚C in density-controlled vials for between 96 and 120 hours after egg-laying (to obtain third to late-third instar larvae). Larvae were mounted in glycerol and their C4da neurons at segments A1-A6 were imaged using a Leica SP8 confocal microscope with a 20x oil objective and a z-step size of 3.5 μm.

Imaginal disc imaging

Imaginal disc dissections were performed as described previously (Han et al., 2004). Briefly, wandering third instar larvae were dissected in a small petri dish filled with cold PBS. The anterior half of the larva was inverted and the trachea and gut were removed. The sample was then transferred to 4% formaldehyde in PBS and fixed for 15 minutes at room temperature. After washing with PBS, the imaginal discs were placed in SlowFade Diamond Antifade Mountant (Thermo Fisher Scientific) on a glass slide. A coverslip was lightly pressed on top. Imaginal discs were imaged using a Leica SP8 confocal microscope with a 20X oil objective.

Assays for Germline Clonal Analysis

To monitor mitotic recombination events resulting in germline clone generation, we performed egg-laying and egg hatchability assays as detailed in Hu and Wolfner (2019), with the exception of using Canton-S males in place of ORP2 males as wild-type mates. Hatchability was calculated only for females that laid eggs. Females that laid no eggs were eliminated from hatchability calculations to avoid inflation of false-zero values.

Image Analysis and Quantification

Counting of wing disc clones was completed manually in Fiji/ImageJ. Counting of neuronal clones was completed manually during the imaging process.

Statistical Analysis

Statistical analyses were performed in R. Student's t-test was conducted for egg-laying data using *Rab* and *kni* gRNAs. For egg-laying data using the 2L toolkit, we performed estimated

marginal means contrasts between gRNAs and *post hoc* one sample t-tests using a generalized

linear model with a negative binomial response. For hatchability data using the 2L toolkit, we

performed estimated marginal means contrasts between proportions of hatched/non-hatched

eggs for each gRNA using a generalized linear mixed-effects model with a binomial response.

For all contrasts, p-values were corrected for multiple comparisons using the Tukey method. For

wing disc and neuronal clone data, we performed Welch's analysis of variance (ANOVA)

followed by pairwise *post hoc* Welch's t-tests. p-values from the multiple *post hoc* Welch's t-

tests were corrected for multiple comparisons using the Bonferroni method.

Acknowledgments:

We thank Bloomington *Drosophila* Stock Center (NIH P40OD018537) and the *Drosophila*

Genetic Reference Panel for fly stocks; Cedric Feschotte, Andy Clark, Eric Alani, and Marcus

Smolka for advice on gRNA design; Cornell CSCU consultants Stephen Parry and Erika Mudrak

for advice on statistics; Michael Goldberg and John Schimenti for critical reading and

suggestions on the manuscript. This work was supported by a Cornell start-up fund and NIH

grants (R01NS099125 and R21OD023824) awarded to C.H., an NIH grant (R01-HD038921)

awarded to M.W., and an NIH training grant (T32-GM07273) awarded to the Cornell BMCB

graduate program that supported S.A..

References:

Aach, J., Mali, P., and Church, G.M. (2014). CasFinder: Flexible algorithm for identifying specific

Cas9 targets in genomes (Cold Spring Harbor Laboratory).

Chari, R., Yeo, N.C., Chavez, A., and Church, G.M. (2017). SgRNA Scorer 2.0: A Species-

Independent Model to Predict CRISPR/Cas9 Activity. ACS Synth. Biol. *6*, 902–904.

Chen, S., Gendelman, H.K., Roche, J.P., Alsharif, P., and Graf, E.R. (2015). Mutational Analysis

of Rab3 Function for Controlling Active Zone Protein Composition at the Drosophila

Neuromuscular Junction.

Chou, T.-B., and Perrimon, N. (1992). Use of a Yeast Sitespecific Recombinase to Produce

Female Germline Chimeras in Drosophila.

Chou, T.-B., Noll, E., and Perrimon, N. (1993). Autosomal P[ovoD1] dominant female-sterile

insertions in Drosophila and their use in generating germ-line chimeras.

Chung, B.Y., Ro, J., Hutter, S.A., Miller, K.M., Guduguntla, L.S., Kondo, S., and Pletcher, S.D.

(2017). Drosophila Neuropeptide F Signaling Independently Regulates Feeding and Sleep-

Wake Behavior. Cell Rep. *19*, 2441–2450.

Clark, A.G., Silveriat, S., Meyerst, W., and Langleyt, C.H. (1994). Nature screen: An efficient

method for screening natural populations of Drosophila for targeted P-element insertions

(transposon tagging/reverse genetics/PCR/mutagenesis).

Clarkson, M., and Saint, R. (1999). A His2AvDGFP Fusion Gene Complements a Lethal

His2AvD Mutant Allele and Provides an in Vivo Marker for Drosophila Chromosome Behavior

(M ary Ann Liebert, Inc. Pp).

Feng, Y., and Irvine, K.D. (2007). Fat and Expanded act in parallel to regulate growth through

Warts. Proc. Natl. Acad. Sci. U. S. A. *104*, 20362–20367.

Freeman, M., Niisslein-Volhard, C., and Glover', D.M. (1966). The Dissociation of Nuclear and

Centrosomal Division in gnu, a Mutation Causing Giant Nuclei in Drosophila.

Garcia-Bellido, A., Ripoll, P., and Morata, G. (1976). Developmental compartmentalization in the

dorsal mesothoracic disc of Drosophila. Dev. Biol. *48*, 132–147.

Golic, K.G., and Lindquist, S. (1989). The FLP recombinase of yeast catalyzes site-specific

recombination in the drosophila genome. Cell *59*, 499–509.

Gratz, S.J., Cummings, A.M., Nguyen, J.N., Hamm, D.C., Donohue, L.K., Harrison, M.M.,

Wildonger, J., and O'connor-Giles, K.M. (2013). Genome Engineering of Drosophila with the

CRISPR RNA-Guided Cas9 Nuclease.

Grenier, J.K., Arguello, J.R., Moreira, M.C., Gottipati, S., Mohammed, J., Hackett, S.R.,

Boughton, R., Greenberg, A.J., and Clark, A.G. (2015). Global Diversity Lines-A Five-Continent

Reference Panel of Sequenced Drosophila melanogaster Strains.

Griffin, R., Sustar, A., Bonvin, M., Binari, R., Del Valle Rodriguez, A., Hohl, A.M., Bateman, J.R.,

Villalta, C., Heffern, E., Grunwald, D., et al. (2009). The twin spot generator for differential

Drosophila lineage analysis. Nat. Methods *6*.

Griffin, R., Binari, R., and Perrimon, N. (2014). Genetic odyssey to generate marked clones in

Drosophila mosaics.

Han, C., Belenkaya, T.Y., Wang, B., and Lin, X. (2004). Drosophila glypicans control the cell-to-

cell movement of Hedgehog by a dynamin-independent process. Development.

Hardwick, K.G., and Pelham, H.R.B. (1992). SED5 encodes a 39-kD integral membrane protein

required for vesicular transport between the ER and the Golgi complex. J. Cell Biol. *119*, 513–

521.

Harrison, D.A., and Perrimon, N. (1993). Simple and efficient generation of marked clones in

Drosophila.

Harrison, M.M., Jenkins, B. V, O'connor-Giles, K.M., and Wildonger, J. (2014). A CRISPR view

of development.

Heitzler, P., and Simpson, P. (1991). The choice of cell fate in the epidermis of Drosophila. Cell

64, 1083–1092.

Henner, A., Ventura, P.B., Jiang, Y., and Zong, H. (2013). MADM-ML, a Mouse Genetic Mosaic

System with Increased Clonal Efficiency. PLoS One *8*, 77672.

Hu, D.J.-K., and Jasper, H. (2019). Control of Intestinal Cell Fate by Dynamic Mitotic Spindle

Repositioning Influences Epithelial Homeostasis and Longevity. Cell Rep. *28*, 2807-2823.e5.

Hu, Q., and Wolfner, M.F. (2019). The Drosophila Trpm channel mediates calcium influx during

egg activation.

Huang, X., Shi, L., Cao, J., He, F., Li, R., Zhang, Y., Miao, S., Jin, L., Qu, J., Li, Z., et al. (2014).

The Sterile 20-Like kinase tao controls tissue homeostasis by regulating the hippo pathway in

drosophila adult midgut. J. Genet. Genomics *41*, 429–438.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012). A

Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity.

Lee, T. (2014). Generating mosaics for lineage analysis in flies. Wiley Interdiscip. Rev. Dev.

Biol. *3*, 69–81.

Lee, T., and Luo, L. (1999). Mosaic Analysis with a Repressible Neurotechnique Cell Marker for

Studies of Gene Function in Neuronal Morphogenesis.

Mackay, T.F.C., and Huang, W. (2018). Charting the genotype-phenotype map: lessons from

the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip. Rev. Dev. Biol. *7*,

e289.

MacKay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., Casillas, S.,

Han, Y., Magwire, M.M., Cridland, J.M., et al. (2012). The Drosophila melanogaster Genetic

Reference Panel. Nature *482*, 173–178.

Muller, H.J. (1928). The Production of Mutations by X-Rays.

Peng, Y., Lee, J., Rowland, K., Wen, Y., Hua, H., Carlson, N., Lavania, S., Parrish, J.Z., and

Kim, M.D. (2015). Regulation of dendrite growth and maintenance by exocytosis.

Perrimon, N. (1998). Creating mosaics in Drosophila. Int. J. Dev. Biol. *42*, 243–247.

Perrimon, N., and Gans, M. (1983). Clonal analysis of the tissue specificity of recessive female-

sterile mutations of Drosophila melanogaster using a dominant female-sterile mutation

Fs(1)K1237. Dev. Biol. *100*, 365–373.

Perrimon, N., Engstromt, L., and Mahowald, A.P. (1989). Zygotic Lethals With Specific Maternal

Effect Phenotypes in Drosophila melanogaster. I. Loci on the X Chromosome.

Poe, A.R., Tang, L., Wang, B., Li, Y., Sapar, M.L., and Han, C. (2017). Dendritic space-filling

requires a neuronal type-specific extracellular permissive signal in Drosophila. Proc. Natl. Acad.

Sci.

Poe, A.R., Wang, B., Sapar, M.L., Ji, H., Li, K., Onabajo, T., and Fazliyeva, R. (2019).

Mutagenesis Reveals Gene Redundancy and Perdurance in Drosophila. Genetics *211*, 459–

472.

Port, F., Chen, H.-M., Lee, T., and Bullock, S.L. (2014). Optimized CRISPR/Cas tools for

efficient germline and somatic genome engineering in Drosophila. Proc. Natl. Acad. Sci. *111*,

E2967–E2976.

Ren, X., Sun, J., Housden, B.E., Hu, Y., Roesel, C., Lin, S., Liu, L.P., Yang, Z., Mao, D., Sun,

L., et al. (2013). Optimized gene editing technology for Drosophila melanogaster using germ

line-specific Cas9. Proc. Natl. Acad. Sci. U. S. A. *110*, 19012–19017.

Salsman, J., and Dellaire, G. (2017). Precision genome editing in the CRISPR era.

Sander, J.D., and Keith Joung, J. (2014). CRISPR-Cas systems for editing, regulating and

targeting genomes. Nat. Biotechnol. Vol. *32*.

Satoh, D., Sato, D., Tsuyama, T., Saito, M., Ohkura, H., Rolls, M.M., Ishikawa, F., and Uemura,

T. (2008). Spatial control of branching within dendritic arbors by dynein-dependent transport of

Rab5-endosomes.

Shorter, J., Couch, C., Huang, W., Carbone, M.A., Peiffer, J., Anholt, R.R.H., and Mackay,

T.F.C. (2015). Genetic architecture of natural variation in Drosophila melanogaster aggressive

behavior. Proc. Natl. Acad. Sci.

Stern, C. (1936). Somatic Crossing Over and Segregation In Drosophlia melanogaster.

Thibault, S.T., Singer, M.A., Miyazaki, W.Y., Milash, B., Dompe, N.A., Singh, C.M., Buchholz,

R., Demsky, M., Fawcett, R., Francis-Lang, H.L., et al. (2004). A complementary transposon

tool kit for Drosophila melanogaster using P and piggyBac.

Xu, T., and Rubin, G. (1993). Analysis of genetic mosaics in developing and adult Drosophila

tissues.

Xu, T., and Rubin, G. (2012). The effort to make mosaic analysis a household tool.

Development.

Yu, H.-H., Chen, C.-H., Shi, L., Huang, Y., and Lee, T. (2009). Twin-spot MARCM to reveal the

developmental origin and identity of neurons. Nat. Neurosci. Vol. *12*.

Zong, H., Espinosa, J.S., Su, H.H., Muzumdar, M.D., and Luo, L. (2005). Mosaic Analysis with

Double Markers in Mice. Cell.

Figures:

Figure 1. CRISPR-induced crossover generates somatic and germline clones in

Drosophila

(A) A diagram illustrating how a CRISPR/Cas9-induced DSB leads to generation of

homozygous clones in a heterozygous organism. The magenta bar indicates an allele that

becomes homozygous in the twin spots.

(B) Strategy for generating mosaic clones using the *His2Av-GFP* marker and a crossover at the

gnu locus.

(C-E) Mosaic clones in wing discs visualized by levels of *His2Av-GFP* expression, as described

in the text. A pair of arrows indicate *His2Av-GFP*$^{+/+}$ (yellow) and *His2Av-GFP*$^{-/-}$ (blue) cells in

each panel. Clones were induced by *Act5C-Cas9* (**C**), *hh-Cas9* (**D**) and *zk-Cas9* (**E**). Scale bars,

50 μm.

(F) Strategy for generating and detecting clones in the female germline using *ovo*^{D1}.

(G) Number of eggs produced by females carrying *nos-Cas9* and *ovo*^{D1}, with (MAGIC) or

without (no gRNA) *gRNA-Rab3*. ***p<0.001, Student's t test. n=number females: no gRNA

(n=18); MAGIC (n=18).

(H) Hatchability of eggs produced females carrying *nos-Cas9* and *ovo*^{D1}, with (MAGIC) or

without (no gRNA) *gRNA-Rab3*. n=number females: no gRNA (N/A); MAGIC (n=18).

For all quantifications, black bar, mean; red bar, SD. Asterisks in (B) and (F) indicate mutated

gRNA target sites.

Rab3

2L

Figure S1. Crossing scheme for germline clone induction

(A) Crossing scheme for germline clone induction using *gRNA-Rab3*, *ovo^{D1}*(2R), and *nos-Cas9*.

(B) Crossing scheme for testing gRNAs for 2L in germline clone induction. The gRNAs used in this test were *gRNA(BFP)* lines.

Figure 2. A toolkit for generating labeled clones for genes on chromosome arm 2L.

(A) Strategy for generating negatively labeled clones for genes on 2L using nuclear BFP.

(B) Strategy for generating positively labeled clones for genes on 2L using Gal80.

(C) A map for gRNA-(40D4) target sites derived from the UCSC Genome Browser. The height

of the conservation bar corresponds to the level of conservation among 23 *Drosophila* species.

(D) A wing imaginal disc showing clones generated by a negative labeling construct (nBFP)

paired with *hh-Cas9*.

(E) Number of clones in wing discs using three different gRNA(BFP) constructs. n=number of

discs: 40D2 (n=17); 40D4 (n=18); 40E1(n=18). Welch's t-tests, p-values corrected using

Bonferroni method; **p≤0.01, ***p≤0.001.

(F) A C4da neuron clone generated by a positive labeling construct (Gal80) paired with *SOP-*

Cas9 and *ppk>CD4-tdTom*.

(G) Quantification of C4da clones using three different gRNA(Gal80) constructs. n=number of

larvae: 40D2 (n=16); 40D4 (n=16); 40E1 (n=16). Clones were visualized by *ppk>CD4-tdTom.*

Welch's t-tests, p-values corrected using Bonferroni method; ***p≤0.001. The asterisk on 40D4

indicates frequent labeling of C3da clones.

(H) Number of eggs produced by using *ovo^{D1}*(2L), *nos-Cas9*, and three different gRNA(BFP)

constructs. The control (ctrl) has no gRNA. n=number of females: ctrl (n=13); 40D2 (n =12);

40D4 (n=12); 40E1 (n=14). Asterisks are from *post hoc* single sample t-tests that compare the

estimated marginal mean (EMM) of each gRNA against the control (μ = 0); ***p≤0.001. p-

values are from contrasts of EMMs of each gRNA (excluding control). EMMs are based on a

negative binomial model. All p-values were corrected using the Tukey method.

(I) Hatchability of eggs produced by females carrying *ovo^{D1}*(2L), *nos-Cas9*, and gRNA(BFP)

constructs. n=number female: 40D2 (n=9); 40D4 (n=7); 40E1(n=9). p-values are from contrasts

of EMMs of proportions of hatched/non-hatched eggs for each gRNA based on a binomial,

mixed-effects model and were corrected using the Tukey method.

For all quantifications: black bar, mean; red bar, SD. Scale bars, 50 μm.

Figure S2. Distribution of da neuron clones using gRNA(Gal80) for 2L

(A) Distribution of da neuron clones in each segment using gRNA(Gal80) for 40D2 and 40D4.

n=number of neurons: 40D2 (n=47); 40D4 (n=52);

Sec5^{E10}   *Rab5^2*   *Syx5^{AR113}*

Figure 3. Clonal analysis of example genes in neuronal dendrite development

(A-C) MAGIC clones of C4da neurons using *Sec5^{E10}* (A), *Rab5^2* (B) and *Syx5^{AR113}* (C),

visualized by *ppk>MApHS*. Scale bars, 50 µm.

DGRP 109   DGRP 223   DGRP 237   DGRP 280   DGRP 356

Figure 4. MAGIC generates clones with DGRP lines

(A-E) Clones in wing imaginal discs by pairing *gRNA-40D2(BFP); hh-Cas9* with DGRP line 109

(A), 223 (B), 237 (C), 280 (D) and 356 (E). Scale bars, 50 µm.

Key Resource Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER	ADDITIONAL INFORMATION
Experimental Models: Organisms/Strains			
ppk-Gal4	Han et al., 2012		*ppk-Gal4* [VK00037]
ppk-Gal4	Han et al., 2012		*ppk-Gal4* [1A]
UAS-CD4-tdTom	Han et al., 2011	RRID:BDSC_35841	*UAS-CD4-tdTom* [7M1]
UAS-MApHS	Han et al., 2014		*UAS-MApHS* [VK00019]
His2Av-GFP	Bloomington Drosophila Stock Center	RRID:BDSC_5941	*His2Av-GFP(S65T)* [62A]
Act5C-Cas9	Bloomington Drosophila Stock Center	RRID:BDSC_54590	*Act5C-Cas9.P*
hh-Cas9	Poe et al., 2019		*R28E04-Cas9* [6A]
SOP-Cas9	Poe et al., 2019		*[sc-E1]x8-Cas9* [3A]
zk-Cas9	This study		*zk-Cas9* [VK00037]
nos-Cas9	Bloomington Drosophila Stock Center	RRID:BDSC_54591	*nos-Cas9.P* [ZH-2A]
ovo [D1] *2L*	Bloomington Drosophila Stock Center	RRID:BDSC_2121	*ovoD1-18* [2La] *ovoD1-18* [2Lb]
ovo [D1] *2R*	Bloomington Drosophila Stock Center	RRID:BDSC_4434	*ovoD1-18* [2R]
gRNA-gnu	This study		*gRNA-gnu(U63)* [VK00027]
gRNA-Rab3	This study		*gRNA-Rab3* [VK00027]
gRNA-40D2(BFP)	This study		*w; gRNA-40D2(BFP)* [VK00037]
gRNA-40D4(BFP)	This study		*w; gRNA-40D4(BFP)* [VK00037]
gRNA-40E1(BFP)	This study		*w; gRNA-40E1(BFP)* [VK00037]
40D2(Gal80)	This study		*w; gRNA-40D2(Gal80)* [VK00037]
40D4(Gal80)	This study		*w; gRNA-40D4(Gal80)* [VK00037]
gRNA-40E1(Gal80)	This study		*w; gRNA-40E1(Gal80)* [VK00037]
gRNA-40D2(BFP)	This study		*w; gRNA-40D2(BFP)* [VK00037]
gRNA-40D4(BFP)	This study		*w; gRNA-40D4(BFP)* [VK00037]
Rab5 [2]	Bloomington Drosophila Stock Center	RRID:BDSC_42702	
Sec5 [E10]	Bloomington Drosophila Stock Center	RRID:BDSC_81044	
Syx5 [AR113]	Bloomington Drosophila Stock Center	RRID:BDSC_3645	
DGRP 109	Mackay et al., 2012	RRID:BDSC_28140	
DGRP 223	Mackay et al., 2012		
DGRP 237	Mackay et al., 2012		
DGRP 280	Mackay et al., 2012	RRID:BDSC_28164	
DGRP 356	Mackay et al., 2012	RRID:BDSC_28178	
Recombinant DNA			
pENTR221-ZK2	Poe et al., 2017		
pDEST-APIC-Cas9	Poe et al., 2019	RRID:Addgene_121657	
pAC-U63-tgRNA-Rev	Poe et al., 2019	RRID:Addgene_112811	
Ubi-CasExpress [attP40]	Bloomington Drosophila Stock Center	RRID:BDSC_65419	Genomic DNA used as a PCR template
pBPGAL80Uw-4	Addgene	RRID: Addgene_26235	
pDEST-APLO	Poe et al., 2017	RRID: Addgene_112805	

Key Resource Table

Software and Algorithms			
Fiji	https://fiji.sc/	RRID: SCR_002285	
R	https://www.r-project.org/	RRID: SCR_001905	
Adobe Photoshop	Adobe	RRID:SCR_014199	
Adobe Illustrator	Adobe	RRID:SCR_010279	
Other			
Gateway™ LR Clonase™ II Enzyme mix	Thermo Fisher Scientific,	#11791020	
NEBuilder® HiFi DNA Assembly Master Mix	New England Biolabs Inc.	#E2621	