Nfkbid is required for immunity and antibody responses to *Toxoplasma gondii*

Scott P. Souza[1,3], Samantha D. Splitt[1,3], Julia A. Alvarez[1,3], Juan C. Sanchez-Arcila[1], Jessica N. Wilson[1,3], Safuwra Wizzard[1], Zheng Luo[4], Nicole Baumgarth[4], Kirk D.C. Jensen[1, 2#]

[1]School of Natural Sciences, Department of Molecular and Cell Biology, University of California, Merced

[2]Health Science Research Institute, University of California, Merced

[3]Graduate Program in Quantitative and Systems Biology, University of California, Merced

[4]Center for Immunology & Infectious Diseases, and Department of Pathology, Microbiology and Immunology, University of California, Davis

[#]**Email:** kjensen5@ucmerced.edu

Keywords: B-1 cells, B-2 cells, isotype class switching, *Nfkbid*, IkBNS, *Toxoplasma gondii*, atypical strains, recombinant inbred mice, QTL, vaccines

SUMMARY (150 word limit)
Protective immunity to parasitic infections has been difficult to elicit by vaccines. Among

parasites that evade vaccine-induced immunity is *Toxoplasma gondii*, which causes lethal

secondary infections in chronically infected mice. To identify correlates of immunity,

mouse genetics were used to identify *Nfkbid*, a nuclear regulator of NF-κB that is required

for B cell activation and B-1 cell development. *Nfkbid*-null mice (bumble) did not generate

parasite-specific IgM and lacked robust parasite-specific IgG, which correlated with

defects in B-2 cell maturation and class-switch recombination. Though high-affinity

antibodies were B-2 derived, transfer of B-1 cells partially rescued the immunity defects

observed in bumble and bone marrow chimeric mice. Immunity in resistant mice correlated

with robust isotype class-switching in both B cell lineages, which can be fine-tuned by

Nfkbid gene expression. We propose a model whereby humoral immunity to *T. gondii* is

regulated by *Nfkbid* and requires B-1 and B-2 cells for full protection.

INTRODUCTION

38
The goal of vaccination is to induce immunological memory that can protect from natural

infection challenge. Depending on the pathogen, effective memory would need to protect

also against a wide variety of pathogen-specific strains encountered in nature. Such

protection is termed heterologous immunity and is effective against pathogen strains that

differ in virulence, immune evasion, or polymorphic antigens. Parasites represent a special

challenge to vaccine development. Indeed, an entirely protective vaccine has yet to be

achieved for any human parasite (Sacks, 2014). The apicomplexan parasite *Toxoplasmas*

gondii, provides an excellent system to explore requirements for heterologous immunity

to a parasitic pathogen. *T. gondii* is a globally spread intracellular protozoan parasite of

warm-blooded animals that exhibits great genetic diversity (Lorenzi et al., 2016). *T. gondii*

strains differ dramatically in primary infection virulence in laboratory mice (Howe and

Sibley, 1995) and in severity of human toxoplasmosis (Grigg et al., 2001; Khan et al.,

2006; McLeod et al., 2012). Such infections can be overcome by immunological memory

responses elicited by vaccination or natural infection. In particular, memory CD8 T cells

and induction of IFNγ are primarily responsible for protection against lethal secondary

infections with the widely studied type I RH strain, which has a lethal dose of one parasite

in naïve mice (Gazzinelli et al., 1991; Gigley et al., 2009; Suzuki, Y. and Remington, 1990).

CD4 T cells are required to help the formation of effector CD8 T cell (Casciotti et al., 2002)

and B cell responses (Johnson and Sayles, 2002), but the ability to adoptively transfer

vaccine-elicited cellular immunity to naïve recipients against the type I RH challenge is

unique to memory CD8 T cells (Gazzinelli et al., 1991; Gigley et al., 2009).

The role of B cells in *T. gondii* infections is less understood. Previous studies

showed that B cell deficient mice (muMT) are extremely susceptible to primary (Chen et

al., 2003), chronic (Kang et al., 2000) and secondary infections (Sayles et al., 2000),

despite unimpaired levels of IFNγ. Passive transfer of antibodies from immunized animals

into vaccinated muMT mice significantly prolongs their survival after challenge (Johnson

and Sayles, 2002; Sayles et al., 2000). IgM seems particularly suited for blocking cellular

invasion by *T. gondii* (Couper et al., 2005), while IgG can perform both neutralization

(Mineo et al., 1993) and opsonization functions (Joiner et al., 1990). Antibody responses

against *T. gondii* are dependent on CD4 T cells (Glatman Zaretsky et al., 2012; Johnson

and Sayles, 2002), and are regulated by cytokines that modulate T follicular helper cell

and germinal center B cell formation in secondary lymphoid organs (Stumhofer et al.,

2013), suggesting conventional "B-2" B cell responses provide antibody-mediated

immunity to *T. gondii.*

In addition, "B-1" cells are innate-like lymphocytes that are known for producing

self- and pathogen-reactive "natural" IgM. B-1 cells are the predominant B cell

compartment within the body cavities, including the peritoneal and pleural spaces and

contribute to antigen-specific responses to many pathogens. In mouse models of

secondary bacterial infections, including *Borrelia hermsii*, *Streptococcus pneumoniae* and

non-typhoid *Salmonella,* vaccination induces protective memory B-1 cells to T cell-

independent bacterial antigens (reviewed in (Smith and Baumgarth, 2019)). This memory

is often restricted to the B-1(b), or CD5- subset of B-1 cells (Alugupalli et al., 2004), but

not in all models (Yang et al., 2012). In the *T. gondii* model, one study suggested that

primed CD5+ B-1(a) cells can rescue B-cell-deficient mice during primary infection with a

low virulence strain (Chen et al., 2003). Memory B cells are also appreciated to secrete

pathogen-specific IgM (Pape et al., 2011), and generate somatically mutated IgM to

combat blood stage secondary infection with *Plasmodium* (Krishnamurty et al., 2016).

Whether IgM responses to *T. gondii* are B-2 or B-1 derived is unknown. Moreover, the role

of B-1 cells in promoting immunity to *T. gondii* during a secondary infection has yet to be

determined.

Particularly troubling for vaccine development for *T. gondii* is the lack of sterilizing

immunity achieved following infection (Jensen et al., 2015). Unlike the highly passaged

lab type I RH strain, for which most immunological memory studies have been performed,

the less passaged type I GT1 strain and atypical strains, many of which are endemic to

South America, cause lethal secondary infections in C57BL/6J mice and co-infect (i.e.

"superinfect") the brains of challenged survivors (Jensen et al., 2015). During secondary

infection memory CD8 T cells become exhausted, but checkpoint blockade fails to reverse

disease outcome (Splitt et al., 2018). The data suggest yet unknown mechanisms are

needed to provide heterologous immunity to highly virulent strains of *T. gondii*. Therefore,

we set out to address whether additional requirements are necessary for heterologous

immunity to *T. gondii*. Through use of forward and reverse genetics, we discovered a

previously unidentified essential role for *Nfkbid* in immunity and antibody responses to *T.*

gondii, and present evidence that both B-1 and B-2 cells assist resistance to secondary

infection with highly virulent parasite strains.

RESULTS

Non-MHC loci control resistance to secondary infection with *Toxoplasma gondii*

When mice are given a natural infection with a low virulent type III CEP strain and

allowed to progress to chronic infection for 35-42 days, the immunological memory that

develops is known to protect against secondary infections with the commonly studied lab

strain, type I RH. In contrast, highly virulent "atypical" strains such as those isolated from

South America (VAND, GUYDOS, GUYMAT, TgCATBr5) or France (MAS, GPHT, FOU)

and the clonal type I GT1 strain led to morbidity and mortality in C57BL/6J mice at varying

frequencies, depending on the strain type and their virulence factors (Jensen et al., 2015).

To explore whether host genetics influenced the ability to survive secondary infection, a

similar experiment was performed with A/J mice. In contrast to C57BL/6 mice, A/J mice

were resistant to secondary infection with all virulent atypical *T. gondii* strains analyzed

(Fig 1A) and cleared parasite burden as early as day 4 post challenge (Fig 1B). In contrast,

naïve A/J mice succumb to infections with atypical strains (not shown). These results

suggest that at least one or more genetic loci controls immunity to virulent challenge.

A/J mice are known to be resistant to primary infections with the intermediate

virulent type II strain (McLeod et al., 1989), and prevent cyst formation due to

polymorphisms in their MHC class I H-2 molecule, L^d (Brown et al., 1995). To test whether

the H-2 locus contributes to immunity we compared secondary infections in C57BL/10

(B10) and C57BL/10.A (B10.A) mouse strains, the latter carrying the MHC H-2 locus of

A/J (H-2a) in place of C57BL/6's MHC H-2 (H-2b). Compared to mice with the H-2b

haplotype, mice expressing H-2a had less cysts and weighed more during chronic

infection with type III strains (P<0.04; Fig S1). Despite their relative health at the time of

challenge, B10.A mice were highly susceptible (0%-30% survival) to secondary infection

with certain atypical strains (VAND, GUY-DOS, GPHT, FOU), but displayed varying

degrees of resistance to others (TgCATBr5, MAS, GUY-MAT; 60-100% survival) (Fig 1A).

In the case of MAS secondary infection, B10.A mice but not B10 mice were highly resistant

and exhibited reduced parasite burden by day 8 post challenge (Fig 1B). Together, the

data suggest that while the MHC H-2a locus is an important modifier of resistance to

certain *T. gondii* strains, this is not true for every challenge. Importantly, the A/J genetic

background encodes additional non-MHC-linked genes that control immunity to *T. gondii*.

Genetic mapping reveals four loci that correlate with immunity to *Toxoplasma*

gondii

To identify non-MHC loci that promote resistance to secondary infection, we first

analyzed the outcome of infection with the type I GT1 *T. gondii* strain, as this strain caused

lethal secondary infections in B10 and B10.A, but not in A/J or first filial generation A/J x

C57BL/6J mice ('F1') (Fig 1C). Following secondary infections, A/J and F1 mice showed

no overt symptoms of weight loss, dehydration, or lethargy (not shown). However, sterile

immunity was not achieved. For example, the GT1 strain was present in the brains of

these survivors (i.e. "superinfection"), and at greater frequencies in F1 compared to A/J

mice (Fig 1D). Superinfections were also detected at high frequencies in B10 and B10.A

surviving mice challenged with atypical strains. Overall, mice of the C57BL genetic

background were more prone to superinfection compared to A/J mice (Table S1). It is

unknown whether virulent strains of *T. gondii* have evolved to superinfect hosts with

immunological memory, as previously hypothesized(Jensen et al., 2015). Nonetheless,

our results underscore the difficulty in achieving sterile immunity to parasites in otherwise

genetically resistant hosts.

Then we performed secondary infection experiments with the type I GT1 strain

using 26 recombinant inbred (RI) mice (Table S2). The AxB:BxA RI mouse panel contains

an assortment of homozygous A/J and C57BL/6 alleles, which assist genetic mapping of

loci that contribute to various phenotypes, including those related to *T. gondii* infection

(Hassan et al., 2015; McLeod et al., 1989). Genetic mapping revealed four distinct

Quantitative Trait Loci (QTL) peaks with logarithm of the odds (LOD) scores greater than

3 on chromosomes 7, 10, 11 and 17 (Fig 2A). None of the QTLs bore evidence for epistatic

interactions (not shown), and only the chromosome 10 QTL surpassed genome-wide

permutation testing (n=1000, P<0.05). Nevertheless, an additive-QTL model including all

four QTLs best fit the data compared to any lesser combination of them (P<0.02, ANOVA).

The estimated effect on the phenotypic variance observed in the RI panel is 24%, 41%,

21% and 27% for the chromosome (chr) 7, 10, 11 and 17 QTLs, respectively. Consistent

with these estimates, complete phenotypic penetrance (i.e. allelic correlation at 100%)

was not observed for any locus (Fig 2B). Moreover, replacing chromosomes 7 or 10 of

C57BL/6J with those of A/J conferred no survival advantage to secondary infection in

consomic mice (Fig S2). Regardless, small effect QTLs controlling complex traits can still

lead to the identification of causal genes within a QTL region, as occurred for the

successful identification of MHC 1 L^d as the host resistance factor to chronic *T. gondii*

infection (Brown et al., 1995).

Nfkbid is one of the most polymorphic genes within the chr7 QTL region (Mb 30.3-

33.0) and sits between the genetic markers that flank the highest imputed LOD score

(Dataset S1). This gene encodes IκBNS, which is a member of atypical NF-κB inhibitors,

called nuclear IκBs, reflecting their restricted cellular localization. Unlike classical inhibitors

of NF-κB, atypical NF-κB inhibitors can modulate NF-κB to induce or repress transcription

(Schuster et al., 2013). Previous work has shown that *Nfkbid* null mice completely fail to

develop B-1 cells, lack circulating IgM and IgG3 antibodies, and cannot respond to T-

independent (T-I) type II antigens such as NP-ficoll (Arnold et al., 2012; Pedersen et al.,

2016; Touma et al., 2011). IκBNS also promotes early plasma blast differentiation

(Khoenkhoen et al., 2019) and IgG1 responses to model T-dependent antigens (Arnold et

al., 2012; Hosokawa et al., 2017; Touma et al., 2011), enhances T cell production of IL-2

and IFNγ (Touma et al., 2007), supports development of T regulatory cells (Schuster et

al., 2012) and suppresses TLR-induced cytokine expression in macrophages (Kuwata et

al., 2006). The tetraspanin, *Tspan8,* is within the chr10 QTL (Mb 115.8-116.2) and is the

most highly polymorphic gene between A/J and C57BL/6J mice in this region (Dataset

S1). *Tspan8* is 6-fold more highly expressed in spleens from A/J compared to C57BL/6

mice (immgen.org). *Tspan8* promotes cancer metastasis (Berthier-Vergnes et al., 2011)

and can impact leukocyte migration (Zhao et al., 2018), but its role in immunity is largely

unknown. Other polymorphic gene candidates within the four QTLs are listed in Dataset

S1.

Nfkbid on chromosome 7 is required for immunity and the generation of

Toxoplasma gondii-specific antibodies

Given the role of *Nfkbid* in several immune functions, degree of polymorphism and

central location within the chr7 QTL, the requirement for *Nfkbid* in immunity to *T. gondii*

was further explored. *Nfkbid* null "bumble" mice (C57BL/6) have previously been

described, which were derived from an ENU mutagenesis screen. These mice possess a

premature stop codon in *Nfkbid,* rendering them unable to support T-I antibody responses

and B-1 development (Arnold et al., 2012). Bumble mice survived primary infection with

the low virulent CEP strain at frequencies similar to wildtype C57BL/6J mice (Fig 2C) but succumbed to secondary infection three days earlier when challenged with the GT1 strain (Fig 2D). Since the C57BL genetic background is uniformly susceptible to GT1 secondary infections used in our genetic screen, susceptibility to challenge with the commonly studied type I RH strain was explored, which is normally controlled in vaccinated or chronically infected mice (Gazzinelli et al., 1991; Gigley et al., 2009; Suzuki, Y. and Remington, 1990). Importantly, bumble mice were entirely susceptible to secondary infection with the type I RH strain (Fig 2E), which exhibited greater parasite loads compared to wildtype mice (Fig 2F). Moreover, bumble mice failed to generate parasite specific-IgM, and were poor producers of parasite-specific IgG3, IgG2b and IgG2a antibody responses after chronic infection (Fig 3A). The remaining antibodies that were secreted exhibited defects in their ability to block parasite invasion of host cells (Fig 3B). Antibodies from naïve mice fail to bind *T. gondii*, thus natural antibodies do not recognize *T. gondii,* consistent with previous reports (Couper et al., 2005). Although *Nfkbid* promotes T cell production of IFNγ and IL-2 in in vitro stimulation assays (Touma et al., 2007) and promotes thymic development of FOXP3+ T regulatory cells (Schuster et al., 2012), no impairment of T cell cytokine production was observed, nor were frequencies of FOXP3+ CD25^{hi} CD4+ T regulatory cells altered in bumble compared to wildtype mice during secondary infection (Fig S3).

To determine where the breakdown in the B cell response occurred in bumble mice, immunophenotyping was employed. Consistent with previous reports, bumble mice have greatly reduced marginal zone B cells in naïve mice and did not increase in frequency during *T. gondii* infection (not shown). Atypical B cells (FCLR5+ CD80+ CD73+) which respond to *Plasmodium* infections in mice (Kim et al., 2019) were also reduced in bumble compared to B6 mice following *T. gondii* infection (not shown). The memory CD73+ B cell

compartment in bumble mice bore evidence for reduced class switching during *T. gondii*

infection, as they remain mainly IgM+IgD+ while C57BL/6J mice have higher frequencies

of IgM-IgD- cells (Fig 3C). Most pronounced, however, is a large accumulation of

transitional stage immature B-2 cells in bumble mice that occurs during chronic infection,

implicating that B cell responses to *T. gondii* may require reinforcement from recent bone

marrow derived emigrants that is blocked in the absence of *Nfkbid*. Hence, *Nfkbid* is not

only an important regulator of B-1 cell development through the transitional stage of

immature B cell development in the steady state (Pedersen et al., 2014), but also for B-2

cell maturation, differentiation and activation during *T. gondii* infection.

Defective B-1 and B-2 responses underlie bumble's defect in immunity

The impaired ability of bumble mice to generate parasite-specific antibodies,

combined with the documented collapse of the B-1 cell compartment in *Nfkbid*-deficient

mice (Arnold et al., 2012; Touma et al., 2011), prompted us to directly assess the role of

B-1 mediated immunity and humoral responses to *T. gondii*. First, total peritoneal exudate

cells (PerC) were adoptively transferred into bumble mice at day 2 of birth, which allows

optimal B-1 engraftment and self-renew for the life of the animal (Baumgarth et al., 2000).

Then, bumble mice that received total PerC transfers were infected with the avirulent type

III strain at 6-7 weeks of age and given a secondary infection 35 days later with the type I

RH strain (Fig 4A). Bumble mice receiving PerC partially reconstituted serum IgM to ~40%

of wildtype levels (Fig S4A), consistent with previous studies (Baumgarth et al., 2000),

and had a significantly delayed time to death relative to non-transferred littermates

following type I RH challenge (Fig 4A). Previous studies have shown B-1 cells respond to

infections and create both pathogen- (Alugupalli et al., 2004; Choi and Baumgarth, 2008)

and microbiota -specific antibodies (Kreuk et al., 2019). Antibody profiling of bumble mice

that received PerC transfer showed a trend of increased anti-*T. gondii* antibody generation

following chronic infection (Fig 4B), but antibody reactivity to the parasite did not reach

levels observed in wildtype mice, suggesting the B-1 compartment has a limited role in

generating high-affinity parasite-specific antibodies. The B cell compartment responsible

for generating parasite-specific antibody was further confirmed with Igh-allotype chimeric

mice which allow tracking of IgM responses of allotype-marked B-1 and B-2 cells (Lalor et

al., 1989). In this experimental setup, endogenous B cells of the C57BL/6 background

(IgH-b allotype) are depleted with allotype specific anti-IgM-b antibodies and replaced with

transferred PerC B-1 of the IgH-a allotype which are refractory to the depletion antibodies

and will engraft for the life of the animal. Following removal of the depleting antibodies,

the endogenous B-2 cell population reemerge and are marked with anti-IgM-b antibodies,

while the transferred B-1 cell are marked with anti-IgM-a antibodies (Fig 4C). Assessing

antibody responses generated in IgH-allotype chimeric mice during a *T. gondii* infection

revealed the presence of B-1 derived IgM-a antibodies that had low reactivity to *T. gondii*

at days 14 and 30 post-infection, but the majority of highly reactive parasite-specific IgM-

b was derived from the B-2 compartment (Fig 4D).

Attempts to explore the role of B-1 and B-2 cells utilizing B cell deficient muMT

mice were complicated by their high susceptible to primary infection with the CEP strain,

irrespective of whether they received PerC as neonates or splenic B-2 cells prior to the

primary infection (Fig S4C-E), thus underscoring the importance of B cells in resistance to

T. gondii infection (Kang et al., 2000). Instead, mixed bone marrow chimeras were

generated in which irradiated bumble or wildtype recipients were transferred wildtype or

bumble bone marrow to reconstitute B-2 cells and the rest of the hematopoietic

compartment. Since B-1a cells do not efficiently reconstitute irradiated recipients from

adult bone marrow (Kantor et al., 1992), some recipients received wildtype PerC to restore

the B-1 compartment in this setting (Choi and Baumgarth, 2008). These and other bone

marrow chimeras (not shown) all succumbed to primary CEP infections (Fig S5). To

bypass the susceptibility of irradiated bone marrow recipients to live *T. gondii* infections

(Yap and Sher, 1999), bone marrow chimeras were vaccinated with a replication deficient

uracil auxotroph strain (RH *Δompdc Δup*) improving overall survival (Fig S5). Whereas

vaccinated wildtype recipient mice that received *Nfkbid*-sufficient but not bumble bone

marrow were able to survive type I RH challenge (Fig 4E), complete immunity was

conferred only when they were transferred PerC. This study also suggests that *Nfkbid* is

important in the non-hematopoietic lineage, as complete immunity was restored only in

wildtype but not bumble recipients (Fig 4E). In summary, our reconstitution studies

emphasize the importance of *Nfkbid* sufficiency in multiple compartments and highlight B-

1 cells as an important contributor to *T. gondii* immunity.

Evidence for enhanced B-1 and B-2 cell activation in resistant A/J mice

Since *Nfkbid* has a profound effect on the maturation and activation of multiple B

cell populations in the C57BL/6J background, we extended our analysis of the humoral

response to the resistant A/J background. A/J mice were found to have a superior IgG

response to parasite lysate antigen during secondary infections (Fig 5A), suggesting

enhanced humoral responses in this background. Within the spleen, both susceptible

C57BL/6J and resistant A/J mice produced similar frequencies of memory B cell CD73+

FCRL5- and atypical memory CD73+ FCRL5+ CD80+ populations during infection (not

shown), however in A/J mice these compartments were drastically increased in their class-

switch recombination frequencies during secondary infection (Fig 5B). The enhanced

class switch potential of resistant mice was also observed in B-1 cells. Within the

peritoneal B-1 cell compartment (CD19+ B220^{int} CD11b+) increased percentages of CD5-

B-1b cells were observed during secondary infection in A/J mice, which appear to have

downregulated their BCR as evidenced by being IgM^{lo} (Fig 6A-B). To further validate these

findings, frequencies of class switched IgM-IgD- CD5+ (B-1a) or CD5- (B-1b) cells within

the CD43+ B-1 cell compartments of the peritoneum (not shown) and spleen were

determined, and a similar trend was observed (Fig 6C). Following infection, B-1 cells in

the spleens of A/J mice maintained high levels of BAFFR and TACI expression and

expressed higher levels of surface CD138 relative to C57BL/6J mice (Fig 6D-E), markers

known to be induced by *Nfkbid* and important for B cell activation and differentiation into

antibody secreting cells (Khoenkhoen et al., 2019). In addition to differences in humoral

immunity, an increase in peritoneal CD8 T cell in vitro recall production of IFNγ was noted

in A/J relative to C57BL/6J mice, however other parameters, including granzyme B and

IL-2 expression and CD4 T cell responses were similar (Fig S6). Collectively these data

suggest that while CD8 T cell production of IFNγ is enhanced in resistant mice, and likely

contributes to their resistance, B-2 and B-1 cells participate in a strikingly enhanced

humoral response in A/J mice, offering insight into how immunity against parasitic

infections may be achieved.

B-1 cells in resistant A/J mice have enhanced germline transcription of *Ighg*

constant regions, class switch recombination and different activation profiles

The diminished antibody response in bumble mice as well as the B cell phenotypic

differences noted in A/J mice prompted us to investigate further how *Nfkbid* may be

modulating the B cell compartment in both genetic backgrounds. A transcriptomic

approach was taken to define peritoneal B-1a, B-1b and B-2 cell response characteristics

in resistant and susceptible mice during naïve, chronic and secondary infection states

(Dataset S2). GO term enrichment analysis consistently found type I and II interferon

signaling and immune defense signatures as being enriched in the most differentially

regulated genes among B cells following infection irrespective of genetic background (Fig

S7A, not shown). Looking specifically at CD5- B-1b cells in A/J mice, which bore evidence

for enhanced activation (Fig 6), pathway enrichment analysis of genes differentially

upregulated on day 5 of secondary infection compared to the naïve state found additional

signatures of TLR-signaling, complement activation and somatic recombination (Fig S7A).

Germline transcription of *Ighg1*, *Ighg3*, *Ighg2b*, and *Ighg2a/c* was greatly enhanced in

CD5- B-1b cells from A/J compared to C57BL/6J mice on day 5 of secondary infection

(Fig 6F), suggesting heightened isotype class switching occurred in mice on the resistant

background. Consistent with their IgM-IgD- phenotype (Fig 6B 6C), both B-1a and B-1b

cells underwent significant IgG isotype class-switching as revealed by intracellular

staining, which was readily observed in the splenic environment of A/J mice (Fig 6 G-H)

where activated B-1 cells migrate to secrete antibody (Ha et al., 2006; Itakura et al., 2005;

Waffarn et al., 2015). Moreover, *Scimp* an adaptor for TLR4 signaling (Luo et al., 2017),

Semaphorin7 a noted inducer of inflammatory cytokines (Suzuki, K. et al., 2007), the

alarmins *S100a8* and *S100a9*, and genes associated with tissue tolerance including

Retnlg and *Slpi,* were specifically upregulated in CD5- B-1b cells from A/J compared to

C57BL/6 mice on day 5 of challenge (Fig S7B). Gene Set Enrichment Analysis (GSEA) of

CD5- B-1b transcriptional variation between mice on day 5 of challenge detected

correlation with gene sets that distinguish B cells following vaccination with different TLR-

stimulating adjuvants (MPL vs R848, GSE25677) (Fig S7C). In summary, the

transcriptomics data suggest in the resistant A/J background, but not in C57BL/6J mice,

B-1 cells undergo Ig class switch recombination, perhaps through enhanced TLR-

signaling.

Gene dosage of *Nfkbid* impacts parasite-specific IgG1 responses

Because A/J *Nfkbid* polymorphisms are largely found in non-coding regions

(Dataset S1), we hypothesized that B cell differences observed in our system could be

due to differences in *Nfkbid* expression levels. To investigate this possibility, we quantified

Nfkbid expression within our transcriptomic dataset as well as by qPCR. B cells from

C57BL/6J mice had greater expression of *Nfkbid* relative to A/J, particularly at the chronic

infection stage (Fig 7A). Because *Nfkbid* expression differences could reflect heightened

responsiveness to parasite load, we stimulated enriched peritoneal and splenic B cells

from A/J and C57BL/6J mice with LPS and noted that B cells from C57BL/6J mice had on

average 1.5- to 2-fold greater induction of *Nfkbid* transcripts relative to A/J (Fig 7B).

To explore how gene dosage of *Nfkbid* impacts antibody responses generated

against *T. gondii,* serum from chronically infected bumble heterozygotes (*Nfkbid+/-*),

C57BL/6J, and A/J mice were analyzed. Although parasite-specific IgM and IgG3 did not

differ between mouse strains (not shown), increased parasite-specific IgG1 serum

responses were observed in A/J relative to C57BL/6 mice, a response that was

phenocopied for IgG1 in bumble heterozygotes (Fig 7C-D). The increase in IgG1 was also

reflected in CD138+ plasma cell differentiation in the bumble heterozygotes (Fig 7E-F),

consistent with previous reports that *Nfkbid* regulates plasma blast and IgG1 responses

(Hosokawa et al., 2017; Khoenkhoen et al., 2019; Touma et al., 2011). Hence, while

Nfkbid is required for the generation of IgM and robust IgG responses against *T. gondii*,

gene dosage of *Nfkbid* further controls IgG1 isotype profiles, presenting *Nfkbid* as tunable

modulator of antibody IgG responses to parasites. Of note, when *Nfkbid+/-* mice were

given secondary infections with the GT1 strain, all mice succumbed to the challenge (not

shown), observations that are consistent with a multiple-QTL model for *T. gondii* immunity

and apparent need for additional modifiers on chromosomes 10 and 17 to survive highly

virulent challenges.

DISCUSSION

The findings of this study underscore the utility of using inbred mouse panels to

uncover novel determinants of immunity to parasites. An unbiased genetic screen

identified *Nfkbid*, a tunable regulator of humoral responses to parasites. Within the

C57BL/6J background, *Nfkbid* is required for maturation and class switching of B-2 cells

during chronic *T. gondii* infection. While B-2 responses dominate the antibody response

in this background, the lack of B-1 cells appear partially responsible for the immunity defect

observed in bumble and are required for immunity in bone marrow chimeric mice. In

contrast, survival against *T. gondii* infection in resistant mice correlates with a strong

layered humoral response: enhanced activation and class-switching in both B-1 and B-2

cells. In this context, B-1 cells may assist the B-2 response to provide full immunity to

challenge. The ability of B-1 cells to make parasite-specific antibodies, though of lower

affinity, potentially amplifies B-2 immune responses to *T. gondii* through internalization of

B-1 cell-derived antigen-antibody complexes (Nguyen et al., 2017), assisting MHCII

antigen presentation for CD4 T cell help (Blandino and Baumgarth, 2019). Moreover, the

effect of B cell-mediated immunity to *T. gondii* and other pathogens is likely

underestimated in murine models using the C57BL/6 background, as enhanced B-1 and

B-2 responses were primarily observed in A/J mice.

Though the exact pathway remains to be investigated, *Nfkbid* is downstream of

TLR signaling in B cells (Arnold et al., 2012; Khoenkhoen et al., 2019; Touma et al., 2011),

which in B-1a cells causes them to downregulate CD5 and facilitate differentiation into

antibody secreting cells (Kreuk et al., 2019; Savage et al., 2019). CD5 is a potent negative

regulator of antigen receptor signaling that renders B-1a cells unresponsive to B cell

receptor (BCR)-triggering. This inhibition is overcome by TLR-stimulation which causes

CD5 to dissociate with the BCR, thereby releasing repression of BCR-mediated signaling

and antibody secretion (Savage et al., 2019) against foreign- and self-antigen (Kreuk et

al., 2019). These data suggest CD5- B-1b cells may represent an activated state of B-1

cells, and calls into question a strict division of labor between these two subsets. This

supposition would fit several of the observations made in our system, including evidence

for enhanced class switch recombination and TLR-gene signatures observed in the CD5-

B-1 cells of the resistant background. In addition, BAFFR and TACI are known inducers

of class-switch recombination (Castigli et al., 2005), and increased expression of these

receptors may further lower the threshold of activation and differentiation into CD138+

plasmablasts/plasma cells, all of which are regulated by *Nfkbid* (Khoenkhoen et al., 2019)

and occurring with greater magnitude in B-1 cells of genetically resistant A/J mice.

Although class switch recombination occurs with much greater frequency in B-2 cells of

A/J mice, we found no evidence for enhanced expression of BAFFR and TACI in this

compartment following *T. gondii* infection (not shown). Further investigation of pathways

upstream of *Nfkbid* has the potential to elucidate key requirements for *T. gondii* immunity.

Nfkbid appears to regulate transitional development in B-2 cells, which is

analogous to previous findings in B-1 cells (Pedersen et al., 2014), but only evident

following *T. gondii* infection. As immature B cells migrate out of the bone marrow to the

spleen, there are several checkpoints which are controlled by NF-κB such as BCR- or

BAFF-mediated signaling (Rowland et al., 2010), both of which are regulated by *Nfkbid*

(Khoenkhoen et al., 2019). Our observation of an accumulation of transitional B cells

during *T. gondii* infection in bumble mice suggest *Nfkbid* plays a role in stabilizing

advancement out of these developmental checkpoints. *Nfkbid* could act as a negative

regulator of BCR signaling, enabling pathogen-reactive B-2 cells to develop beyond

negative selection that would otherwise occur as antigen accumulates in secondary

lymphoid organs over time. Alternatively, *Nfkbid* could be a positive regulator of NF-κB

signaling, increasing the strength of BCR signaling to enable transitional B cells to become

mature B cells. In both cases, a developmental defect likely restricts the pool of *T. gondii*-

reactive mature B cells, preventing replenishment of antibody secreting cells during

infection, culminating in the low parasite-specific antibody titers observed in bumble mice.

It is important to emphasize that multiple polymorphisms determine the complex

phenotype of secondary infection immunity to *T. gondii*. Our genetic screen revealed at

least 4 loci that each account for 20-40% of the overall heterologous immunity to *T. gondii*,

and that the H-2 locus can be an important modifier of resistance against certain parasite

strains. Perhaps not surprising, the *Nfkbid* polymorphism in a stand-alone fashion did not

fully restore immunity, as inferred from chromosome 7 consomic mice and by our attempts

to mimic the lower gene expression observed in resistant mice through heterozygous

expression. Whereas polymorphic *Nfkbid* contributes 21% to this phenotype, perhaps by

regulating plasma cell differentiation (Fig 7), this smaller effect QTL was instrumental in

identifying *Nfkbid*, where a more drastic gene inactivation revealed its role in multiple

compartments in bumble mice, notwithstanding its requirement for humoral immunity.

In summary, heterologous immunity to a parasitic pathogen should, at a minimum,

prevent disease against a wide variety of strains that differ in virulence or polymorphic

antigens. An ideal parasite vaccine would entirely protect against re-infection and induce

sterile immunity, thought possible since re-infection studies were first performed in mice

(Nussenzweig et al., 1967) and humans immunized with irradiated sporozoites and

Plasmodium sp. challenge (Hoffman et al., 2002). Yet, only one partially protective vaccine

is in use for any human parasitic pathogen, RTS,S/AS01, which has low efficacy for

malaria prevention (RTS, S Clinical Trials Partnership, 2015). Our findings highlight the

role of both innate and conventional B cells in humoral immunity to *T. gondii*, introducing

B-1 cells as a potential vaccine target along with B-2 cells to maximize humoral immunity

to parasitic infections. Moreover, we present a modulator of antibody responses against

parasitic infections, *Nfkbid*, a transcriptional regulator that can tune B cell responses to
provide an overall effective class-switched antibody response against parasites.
449

ACKNOWLEDGMENTS
This work was supported by NIH grants R01 AI137126, R15 AI131027, a Hellmann
Fellowship and an MCB departmental award to KJ; and by NIH grants U19-AI109962 and
R01 AI117890 to NB. The sequencing was carried out at the DNA Technologies and
Expression Analysis Cores at the UC Davis Genome Center, supported by NIH Shared
Instrumentation Grant 1S10OD010786-01. We would like to thank David Gravano and the
UC Merced Stem Cell Instrumentation Foundry for their assistance designing panels and
utilizing instruments, supported by the DoD Research and Education Program for
HBCU/MSI Instrumentation Grant W911NF1910529. The authors declare no conflict of
interest.

460
461 **Author Contributions:** S.P.S., S.D.S., N.B. and K.J. designed research; S.P.S., S.D.S.,
462 J.A., J.C.S.A., J.W., S.W., and K.J. performed research; Z.L. and N.B. contributed new
463 reagents/analytic tools; S.P.S., S.D.S., and K.J. analyzed data; and S.P.S., S.D.S., N.B.
464 and K.J. wrote the paper.
465 **Competing Interest Statement:** The authors disclose no competing interests.
466

FIGURES AND TABLES

Figure 1. MHC and non-MHC alleles promote immunity to virulent *Toxoplasma gondii* strains in A/J mice.

All mice were infected with 10^4 type III CEP *hxgprt-* avirulent *T. gondii* parasites and allowed to progress to chronic infection; then, 35 days later, mice were challenged with 5×10^4 of the indicated strains of *T. gondii*. A) Survival of A/J, C57BL/10J (B10), and C57BL/10J.A (B10.A) mice following secondary infection with atypical strains of *T. gondii*. Cumulative results are plotted from 1-2 experiments; n=4 to 12 mice per parasite strain and mouse genetic background. B) Bioluminescence imaging of individual mice challenged with luciferase expressing MAS strain on days 1, 4 and 8 of secondary infection; parasite burden is shown as a heat map depicting the relative number of photons (photons/sec/cm2/sr) detected over a 5 minute exposure. C) Survival of B10 (n=5), B10.A (n=5), A/J (n=12), and F1 (A/J x C57BL/6, n=9) mice following secondary infection with the type I GT1 strain; ***P<0.0001, Log-rank (Mantel-Cox) compared to A/J mice. Cumulative data from 1 to 2 experiments are plotted. D) Superinfection in surviving A/J

(n=12) and F1 (n=9) mice following 35 days of secondary infection with the type I GT1

strain. To evaluate superinfection, brain homogenate was grown in MPA-xanthine

selection medium, which selects for GT1 parasites expressing the endogenous *HXGPRT*

locus and against the *hxgprt-* type III CEP strain used to induce chronic infection. Plotted

is the fraction of mice for which the presence or absence of the GT1 strain was detected;

**** P< 0.0001, Fisher's exact test.

490
491
492
493

Figure 2. Genetic mapping reveals *Nfkbid* is required for immunity to *Toxoplasma gondii* secondary infections.

26 recombinant inbred (RI) mouse strains from the AxB;BxA panel were primed with 10^4 type III *T. gondii* CEP *hxgprt-* parasites; then, 35 days later, mice were challenged with 5×10^4 virulent type I GT1 *T. gondii* parasites (n=2 per RI line). A) LOD scores for each marker were calculated using Haley-Knott regression and the running LOD scores of primary (black) and secondary infection survival (blue) for each genetic marker is plotted. 1000 permutations were performed to obtain the genome wide threshold LOD values; P=0.10 and 0.05 thresholds are shown. B) Effect plots for the genetic markers closest to the maximal LOD scores calculated for the chromosome 7, 10, 11 and 17 QTLs are shown. Each dot indicates the percent survival of a unique RI line and whether it encodes an A/J (red) or C57BL/6 (blue) allele at the specified genetic marker. C) Cumulative survival of bumble (*Nfkbid*-/- C57BL/6) (n=71) and wildtype (C57BL/6J) (n=89) naïve mice infected with the avirulent type III CEP strain. D) Survival of chronically infected bumble (n=12) and wildtype (n=8) mice given a secondary infection with the type I GT1 strain. Cumulative survival from 3 separate experiments are shown; * P<0.02, Gehan-Breslow-Wilcoxon test. E) As in B, but survival to secondary infection with the type I RH strain is shown. Cumulative survival from 3 separate experiments is plotted (bumble n=17, C57BL/6J n=4); ***P<0.008, Mantel-Cox test. F) Frequency of GFP+ events in the peritoneal lavage 7 days post-secondary infection with GFP-expressing type 1 RH strain (RH 1-1). Each dot represents the result of one mouse, and cumulative results are shown from 3 separate experiments (bumble n=13, C57BL/6J n=17); **P<0.002, unpaired two-tailed t-test.

Figure 3. *Nfkbid* is required for humoral responses to *Toxoplasma gondii*.

A) Whole fixed GFP+ parasites were incubated with serum from chronically infected mice, stained with fluorescently labeled anti-isotype antibodies and assessed by flow cytometry. Quantification of *T. gondii*-specific antibody isotype binding (IgM, IgG1, IgG2a/c, IgG2b, and IgG3) over a range of serum concentrations is shown. Background staining in the absence of serum is indicated by the dotted line for each isotype. Plotted is the cumulative average +/-SD of the geometric mean fluorescence straining intensity (MFI) from 3 separate experiments (bumble n=8, C57BL/6J n=11). B) Neutralization of GFP+ parasites coated with serum over a range of concentrations from the indicated chronically infected mice. Parasites were incubated in serum for 20 minutes before infection of mouse embryonic fibroblasts and assessed by FACS 2h later. The fraction of infected host cells (GFP+ cells) is normalized to that of parasite infections without serum. Each dot represents the serum from an individual mouse and cumulative results from 3 separate experiments are shown (bumble n=8, C57BL/6J n=7, naïve n=6). For A and B, significance was assessed by unpaired t-tests with Holm-Sidak correction for multiple comparisons; *** P< 0.001, ** P<0.01, * P<0.01. C) Representative FACS plots of chronically infected bumble and B6 mice and scatter plot of the frequency of splenic B-2 cell compartments quantifying transitional B cells at naïve, d12 of primary infection, and chronic infection. Representative FACS plots of chronically infected bumble and B6 mice and scatter plot of the frequency of IgM- IgD- cells of the CD73+ conventional memory B population. Cumulative data from two experiments n=5-6 mice/condition. * P<0.05 by unpaired two-tailed t-test.

Figure 4. The contribution of B-1 and B-2 cells to *Toxoplasma gondii* immunity in C57BL/6J mice.

A) Schematic of the secondary infection experiment using PerC reconstituted bumble mice. 2-4 day old bumble neonates were transferred 5×10^6 total PerC and allowed to rest for 7 weeks before primary infection with the type III CEP strain. 5 weeks post-primary infection, mice were given a secondary infection with the type I strain RH. Survival of bumble mice given total PerC transfers (n=10) relative to littermate controls (n=12). Cumulative survival is shown from 3 separate experiments; *** P <0.001, Mantel-Cox test. B) Representative histograms of anti-isotype staining of parasites coated in serum (10^3 dilution for IgG, 10^2 for IgM) from chronically infected bumble, bumble given PerC transfer, and WT mice. C) Schematic of neonatal allotype chimera generation. C57BL/6J neonates were given anti-IgHb to deplete endogenous B cells at day 1 post birth and twice weekly after for 6 weeks, thereby depleting the endogenous B-1 pool for the life of the animal due to their restricted fetal/neonatal window of development. Neonates were given 5×10^6 total PerC from 6-8 week old IgH-a congenic C57BL/6 mice donors. These mice then rested for 6 weeks after the last depletion treatment to allow reemergence of the endogenous B-2 IgH-b cells. D) Representative histograms of serum derived anti-IgM-a (B-1 derived) or anti-IgM-b (B-2 derived) staining profiles of type I RH GFP+ parasites taken from the IgH allotype chimeras on day 14 and 30 following primary type III CEP infection. Staining controls with serum from chronically infected C57BL/6J IgH-b littermates, or IgH-a mice are shown. E) Irradiated bumble and WT recipients (45.1 or 45.2) were given WT or bumble bone marrow (BM) with or without total WT PerC (45.2). Mice were vaccinated with a replication deficient type I strain (RH *Δup Δompdc*) and 30d later challenged with

565   type I RH. Cumulative survival is shown from 2-3 experiments (n=4-9 per condition); *

566   P<0.05 by Mantel-Cox test.

Figure 5. Immunity in A/J mice correlates with enhanced class-switching in memory

B-2 cells and increased serum reactivity to parasite lysate antigen.

A) Serum obtained from A/J and C57BL/6J mice chronically infected (CEP) or at D5 of

secondary infection (GT1) were used to probe GT1 parasite lysate separated by SDS-

PAGE, western blots were detected with anti-mouse IgG. "Total Lane Signal" is the signal

obtained from the entire lane of C57BL/6J compared to that of A/J (=1); western blots were

developed in tandem and analyzed by Image J. Results are from 6 individual experiments;

** P < 0.01, * P < 0.05; unpaired two-tailed t-tests. B) Gating strategies for identifying

memory B cells. Memory B cells are identified as CD19+ B220+ CD23+ CD21^{mid} CD73+.

Conventional memory B cells are FCRL5- CD80- while atypical memory B cells are

identified as FCRL5+, CD80+. Representative FACS plots of memory compartments in

A/J and C57BL/6J mice on day 5 of secondary infection with the type I GT1 strain are

shown. The frequency of class-switched (IgM- IgD-) memory cells at the indicated infection

states were analyzed. Each dot represents the results from an individual mouse and the

cumulative averages +SD from 2 experiments are plotted. N=3-6 mice per infection state.

Significance was assessed with an unpaired two-tailed t-test; *** P<0.0001, ** P<0.01.

Figure 6. Evidence for enhanced B-1 cell activation in resistant A/J mice.

A) Gating strategies for identifying B-1 (CD19+ B220$^{int-neg}$) and B-2 (CD19+ B220^{hi}) B cells. The legend applies to panels C-H. B) Representative FACS plots of the CD11b+ peritoneal B-1 B cell compartment in A/J and C57BL/6J (B6) mice at naïve, chronic (Chr), and 5 days (D5) post-secondary infection with the GT1 strain. Numbers indicate the percent of cells that fall within the indicated gate. Representative histograms of IgM surface expression and percent of cells that fall within the IgM^{lo} gate of CD5- B-1b cells from A/J (red) and C57BL/6J (blue) at the indicated time points. C) Frequencies of splenic CD43+ B-1a (CD5+) or B-1b (CD5-) that are IgM+IgD- or IgM-IgD- in A/J and C57BL/6J mice at the indicated infection states. D) Representative FACS plots of splenic CD19+ B220$^{int-neg}$ B cells stained for CD43 and CD5 in A/J and C57BL/6J mice at D5 of secondary infection. Representative CD138 expression on CD5- CD43+ B-1b cells. E) Frequencies of CD43+ splenic B-1a and B-1b cells from A/J and C57BL/6J mice that express BAFFR+, TACI+, or CD138+ at the indicated infection states. F) Heatmap depicting the relative expression of all *Ighg* transcripts from the indicated B cell population, mouse strain and infection state. G) Representative FACS plots of intracellular IgG (H+L) of B-1b cells, and H) frequency of both peritoneal and splenic B-1a and B-1b cells of A/J and C57BL/6J mice at day 5 of secondary infection. For C, E and H, the cumulative average +SD from 2-4 experiments is plotted and each dot represents the result from an individual mouse; P values calculated by 2-way ANOVA with Tukey correction; **** $P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$.

A
Nfkbid expression

B
Relative Nfkbid expression A/J B cells
PerC   Spleen

C
Serum-coated *Toxoplasma gondii* (GFP+)

- C57BL/6J
- *Nfkbid*+/- (C57BL/6)
- A/J
- Naive
- - - No serum

D
IgG1

E
A/J   C57BL/6J   *Nfkbid*+/- (C57BL/6)

0.9   0.6   1.4

B220
CD138+

F
CD138+ B220-

A/J   C57BL/6J   *Nfkbid*+/-

Figure 7. Gene dosage of *Nfkbid* impacts parasite-specific IgG1 responses.

A) *Nfkbid* expression in CPM (Counts per million) of 3'-Tag RNA-seq reads of the indicated

B cell populations obtained from A/J and C57BL/6J mice that were either naïve, chronically

infected, or on D5 of secondary infection with the type I GT1 strain. B) Enriched B cells

from the PerC and spleen were stimulated with LPS for 2 hrs and *Nfkbid* transcripts were

quantified by qPCR; ** P< 0.01, paired t-test. C) Representative histograms display the

detection of parasite-specific IgG1 bound to formaldehyde fixed GFP+ type I GT1

parasites; diluted serums (10^3) from C57BL/6J, *Nfkbid*+/- (C57BL/6J x bumble F1), and

A/J mice chronically infected with the type III CEP strain were assayed. Anti-mouse IgG1

background staining in the absence of serum is shown. D) As in C, but quantification of

the parasite-specific IgG1 antibody isotype binding over a range of serum concentrations

is shown. Plotted is the cumulative average +/-SD of the MFI from 2-3 separate

experiments (C57BL/6J n=8; *Nfkbid*+/- n=7; A/J n=8); significance was assessed by

unpaired t-tests and Holm-Sidak corrections comparing A/J vs C57BL/6J (*) or *Nfkbid*+/-

vs C57BL/6J (#); **** P<0.0001, # P<0.05. IgG1 staining was not significantly different

between A/J and *Nfkbid*+/- serums. E) Representative FACS plots of the dump- CD19+

CD138+ plasmablast populations within A/J, C57BL/6J and *Nfkbid*+/- mice at day 5 of

secondary infection with type I GT1 parasites. F) Frequency of B220- CD138+

plasmablasts of total live dump- CD19+ cells. Plotted is the cumulative average +/-SD of

2 separate experiments (n=5 per mouse strain); significance was assessed by unpaired t-

tests; * P< 0.05.

STAR METHODS

Parasite strains and cell lines

Human foreskin fibroblasts (HFFs) monolayers were grown in DMEM (4.5 g/L D-glucose) (Life Technologies) supplemented with 2 mM L-glutamine, 20% fetal bovine serum (FBS) (Omega Scientific), 1% penicillin-streptomycin, and 0.2% gentamycin (Life Technologies). Mouse Embryonic Fibroblasts (MEFs) were grown in DMEM (4.5 g/L D-glucose) (Life Technologies) supplemented with 10% fetal bovine serum (FBS) (Omega Scientific), 20mM HEPES, 1% penicillin-streptomycin, and 0.2% gentamycin (Life Technologies). *Toxoplasma gondii* strains were passaged in HFFs in 'Toxo medium' (4.5 g/L D-glucose, L-glutamine in DMEM supplemented with 1% FBS and 1% penicillin-streptomycin). The following clonal strains were used (clonal types are indicated in parentheses): RH Δ*ku80* Δ*hxgprt* (type I), RH (1-1) *GFP::cLUC* (type I), GT1 (type I), GT1 *GFP::cLuc* (type I), and CEP *hxgprt-* (type III). The following atypical strains were used: MAS, MAS *GFP::cLuc* (2C8) (haplogroup 'HG' HG4), GUY-MAT (HG5), FOU (HG6), GPHT (HG6), TgCATBr5 (HG7), GUY-DOS (HG10), and VAND (HG10). The uracil auxotroph vaccine strain, RH Δ*up* Δ*ompdc* (Fox and Bzik, 2010), was passaged in HFFs in medium containing 250μM uracil.

Generation of GFP-expressing GT1 strains

GT1 parasites were transfected with linearized plasmids for parasite expression of GFP and click beetle luciferase (*GFP::cLUC*), parasites were grown on HFF monolayers in T-25 flasks in Toxo medium for 2 weeks. Parasites were removed from the flasks by scraping; the parasites were pelleted and washed with PBS and suspended in sterile FACS buffer (2% FBS in PBS). Fluorescent parasites where then sorted via fluorescence-activated cell sorting (FACS) into a 96-well plate with confluent HFF monolayers. To

ensure single plaque formation in at least one of the wells, the sort was titrated using the

following parasite numbers: 100, 50, 25, 12, 6, 3, 2, and 1 for each well per row of 8.

Mice and ethics statement

Female C57BL/6J (H-2b), A/J (H-2a), C57BL/10SnJ (H-2b), B10.A-*H2ᵃ H2-T18ᵃ*/SgSnJ

(H-2a), B6AF1/J (A/J x C57BL/6J F1 progeny), B6.129S2-*Ighm^{tm1Cgn}*/J (muMT), B6.SJL-

Ptprcᵃ Pepcᵇ/BoyJ (CD45.1 congenic), B6.Cg-*Gpi1ᵃ Thy1ᵃ Ighᵃ*/J (IgH-a triple congenic

mice), C57BL/6J-Chr7^{A/J}/NaJ and C57BL/6J-Chr10^{A/J}/NaJ (chromosome 7 and 10

consomic mice), and 26 (AxB;BxA) recombinant inbred (RI) mice derived from A/J and

C57BL/6 founders, were purchased from Jackson Laboratories. The bumble mouse line

used for this research project (Arnold et al., 2012), C57BL/6J-*Nfkbid*^{m1Btlr}/Mmmh,

RRID:MMRRC_036725-MU, was obtained from the Mutant Mouse Resource and

Research Center (MMRRC) at University of Missouri, an NIH-funded strain repository, and

was donated to the MMRRC by Bruce Beutler, M.D., University of Texas Southwestern

Medical Center. Bumble mice were crossed to C57BL/6J to generate F1 bumble

heterozygotes (*Nfkbid*+/-).

Mice were maintained under specific pathogen free conditions at UC Merced.

Every effort was made to ensure unnecessary stress on the animals was avoided. Mouse

work was performed in accordance with the National Institutes of Health Guide to the Care

and Use of Laboratory Animals. All protocols have been reviewed and approved by UC

Merced's Committee on Institutional Animal Care and Use Committee. UC Merced has an

Animal Welfare Assurance filed with OLAW (#A4561-01), is registered with USDA (93-R-

0518), and the UC Merced Animal Care Program is AAALAC accredited (001318).

Parasite infections

Parasite injections were prepared by scraping T-25 flasks containing vacuolated HFFs

and sequential syringe lysis first through a 25G needed followed by a 27G needle. The

parasites were spun at 400 rpm for 5 min to remove debris and the supernatant was

transferred, followed by a spin at 1700 rpm washing with PBS. For primary infections, mice

were infected intraperitoneally (i.p.) with 10^4 tachyzoites of type III CEP *hxgprt-*. For some

experiments, mice were vaccinated i.p. with 10^6 tachyzoites of RH *Δup Δompdc.* For

secondary infections, mice were infected I.P. with 5×10^5 type I parasites (RH or GT1).

Parasite viability of the inoculum was determined by plaque assay following i.p. infections.

In brief, 100 or 300 tachyzoites were plated in HFF monolayers grown in a 24-well plate

and 4-6 days later were counted by microscopy (4x objective).

Blood plasma isolation and assessment of seroconversion

All mice were assessed for sero-positivity to *T. gondii* 4-5 weeks post primary infection.

50µL of blood was isolated from mice in tubes containing 5µL of 0.5M EDTA on ice,

pelleted and the supernatant containing blood plasma was heat inactivated to denature

complement at 56°C for 20 minutes and then stored at -80°C. HFFs were grown on

coverslips and infected with GFP-expressing RH (1-1) overnight, fixed 18 hrs later with

3% formaldehyde (Polysciences) in PBS, washed, permeabilized and blocked with PBS

containing 3% bovine serum albumin Fraction V (Sigma), 0.2M Triton X-100, 0.01%

sodium azide, incubated with a 1:100 dilution of collected blood plasma for 2 hrs at room

temperature, washed with PBS, and detected with Alexa Fluor 594-labeled secondary

antibodies specific for mouse IgG (cat # A11032, Life Technologies). Seropositive

parasites were observed by immunofluorescence microscopy (Nikon Eclipse Ti-U).

Brain superinfection assays and cyst enumeration

Brains from chronically infected mice (CEP *hxgprt-*) that survived secondary challenge

were dissected, rinsed in PBS, passed through a 21G needle several times, pelleted and

suspended in 1mL of PBS. For rederivation, 100µL of the brain homogenate was used to

inoculate HFF monolayers in Toxo medium. One to two weeks later, infected HFFs were

syringe-lysed and plated on new HFF monolayers to encourage parasite growth. Once

HFFs were fully vacuolated, parasites were passaged in Toxo medium supplemented with

mycophenolic acid (MPA) and xanthine that selects for parasites encoding a functional

HXGPRT (i.e. the challenging strains) and against the chronically infecting type III *hxgprt-*

which lacks a functional *HXGPRT* gene. Outgrowth in MPA-xanthine was considered

evidence for superinfection.

For counting tissue cysts, 100µl of brain homogenate was fixed in 900µL of ice

cold methanol, incubated for 5 minutes in microtubes (MCT-175-C, Axygen), washed and

stained overnight in a 500µL PBS solution containing 1:150 dilution of FITC-conjugated

Dolichos biflorus agglutinin (Vector Laboratories) with slow rotation at 4°C. The stained

homogenate was further washed and suspended in 1mL of PBS, of which several 50µL

aliquots were counted by fluorescence microscopy, and the number of cysts per brain

were deduced.

Genetic linkage analysis

Quantitative trait loci (QTL) analysis was performed with the package r/QTL in R (version

3.6.1). LOD scores for each marker were calculated using the Haley-Knott regression

model with the function 'scanone', or for all possible combination of two markers (i.e.

epistatic interactions) using the function 'scantwo'. 1000 permutations were performed to

obtain the genome wide LOD threshold for a P value of ≤0.05, which was considered

statistically significant. Similar results were obtained with a linear mixed regression model.

To estimate the effect each QTL had on the overall phenotype, the function 'fitqtl' was first

used to fit the data to a multiple-QTL model. Statistical support was found for inclusion of

all four QTLs with LOD scores > 3 compared to any lesser combination of three-QTLs

(ANOVA P <0.02). Individual QTL effects were then calculated under the assumption of

the four-QTL model, which collectively accounts for 91% of the observed phenotypic

variance.

Cell isolation, in *vitro* recall infections, and FACS analysis

PECs were isolated by peritoneal lavage and splenocytes obtained, as described in (Splitt

et al., 2018). In brief, 4mL of FACS buffer (PBS with 1% FBS) and 3mL of air were injected

into the peritoneal cavity with a 27G needle. After agitation, the PEC wash was poured

into a conical tube. PEC washes were filtered through a 70μm cell strainer, pelleted, and

washed with FACS buffer before staining. Spleens were dissected and crushed through

70μm cell strainers, pelleted, incubated in ACK red blood cell RBC lysis buffer (0.15M

NH_4Cl, 10mM $KHCO_3$, 0.1mM EDTA) for 5 minutes at room temperature, then washed

with FACS buffer. To obtain peripheral blood leukocytes (PBLs), 50μL of blood was

isolated from mice in tubes containing 5μL of 0.5 M EDTA on ice, pelleted and incubated

in ACK lysis buffer, washed and peripheral blood leukocytes (PBLs) were suspended in

FACS buffer.

For FACS analysis, all preparations were done on ice, and cells were blocked in

FACS buffer containing Fc Block anti-CD16/32 (2.4G2) (BD Biosciences), 5% normal

hamster serum, and 5% normal rat serum (Jackson ImmunoResearch) for 20 minutes

prior to staining with fluorophore-conjugated monoclonal antibodies (mAbs). The following

mAbs (1:100 staining dilutions) were used: anti-CD1d-BV650 (1B1, BD Bioscience), anti-

CD11c-eFlour 450 (N418, eBioscience); anti-CD11c-eFlour 450 (N418, BD Bioscience);

anti-CD45.2-eFlour 450 (104, eBioscience); anti-CD4-eFlour 450 (GK1.5, eBioscience),

anti-CD4-PECy7 (GK1.5, eBioscience); anti-CD11b-FITC (M1/70, eBioScience), anti-

CD11b-BUV395 (M1/70, BD Bioscience), anti-CD11b-BV421 (M1/70, BD Bioscience),

anti-CD11b-Pacific Blue (M1/70, BioLegend); anti-IFNγ-PE (XMG1.2, BD Bioscience);

anti-CD8α-APC (53-6.7, eBioscience), anti-CD8α-BV510 (53-6.7, BioLegend), anti-Ly6G-

APC (1A8-Ly6g, eBioscience), anti-CD19-PerCP-Cy5.5 (ebio1D3, eBioscience), anti-CD19-PE (6D5, BioLegend), anti-CD19-BV785 (6D5, BioLegend), anti-CD3-eFlour 780 (17A2, BD Biosciences), anti-Ly6C-PECy7 (HK1.4, BioLegend), anti-CD23-Pacific Blue (B3B4, BioLegend), anti-CD23-AF700 (B3B4, BioLegend), anti-CD21/CD35-FITC (7E9, BioLegend), anti-CD21/CD35-PE (7E9, BioLegend), anti-CD5-APC (53-7.3, BioLegend), anti-CD43-BV510 (S7, BD Bioscience), anti-CD43-BUV737 (S7, BD Bioscience), anti-CD5-PerCP-Cy5.5 (53-7.3, BioLegend), anti-CD5- Cy7-APC (53-7.3, BioLegend), anti-CD45R/B220-Cy7-APC (RA3-6B2, anti-CD45R/B220-BUV-661 (RA3-6B2, BD Bioscience), anti-CD73-Cy7-PE (eBioTY/11.9, eBioscience), anti-CD80-BV711 (16-10A1, BioLegend), anti-FCRL5-af88 (polyclonal, R&D systems) anti-IgM-PECy7 (RMM-1, BioLegend), anti-IgM-BV605 (RMM-1, BioLegend), anti-IgD-FITC (11-26c.2a, BioLegend), anti-IgD-PEDazzle (11-26c.2a, BioLegend), anti-CD138/Syndecan-1-BV510 (281-2, BD Bioscience), anti-CD138/Syndecan-1-BV650 (281-2, BioLegend), anti-mouse-CD267/TACI-AlexaFlour-647 (8F10, BD Biosciences), and anti-CD268/BAFF-R-PE (7H22-E16, BioLegend). Other FACS reagents included the viability dye propidium iodide (Sigma) at a final concentration of 1μg/mL.

For *in vitro* recall, splenocytes and PerC were isolated from chronic and challenged mice (day 5 or 7 following secondary infection) and 6 x 10^5 cells per well (96-well plate) were plated in T cell medium (RPMI 1640 with GlutaMAX, 20% FBS, 1% Pen/Strep, 1mM NaPyruvate, 10mM HEPES, 1.75μl BME). Cells were infected with a type I strain (RH or GT1) strain at an MOI (multiplicity of infection) of 0.2 for 18 hr; 3μg/mL brefeldin A (eBioscience) was added for the last 5 hr of infection. 96-well plates were placed on ice, cells were harvested by pipetting and washed with FACS buffer, blocked, and stained for surface markers. Cells were fixed with BD Cytofix/Cytoperm and permeabilized with BD Perm/Wash solution (cat# 554714, BD Pharmingen), stained with anti-IFNγ-PE (XMG1.2,

BD Bioscience), anti-GZB (GB11, BioLegend) and anti-IL-2-APC (JES6-5H4, BioLegend) on ice for 1 hr or overnight. Cells were then washed once with BD Perm/Wash solution, once in FACS buffer, and analyzed by FACS.

For FoxP3 staining, peritoneal lavage was performed on chronic and challenged (day 7 following secondary infection) bumble and WT mice. Cells were washed and surface stained. Fixation and permeabilization was performed with the eBioscience Foxp3/Transcription Factor Staining Buffer Set (cat # 00-5523-00) before intracellular staining with anti-Foxp3-PE (MF-14, BioLegend) according to the manufacturer's recommendations. Flow cytometry was performed on a Beckman Coulter Cytoflex LX, LSR II (BD Biosciences), or the Bio-Rad ZE5 and analyzed with FlowJo™ software.

For intracellular staining of IgG H/L PerC and splenocytes were surface stained with anti-CD43-BV510 (S7, BD Bioscience), anti-CD19-PE (6D5, BioLegend), anti-IgD-FITC (11-26c.2a, Biolegend), anti-IgM-PE/Cy7 (RMM-1, BioLegend), anti-CD5-APC (53-7.3, BioLegend), anti-CD45R/B220-Cy7-APC (RA3-6B2). Cells were then fixed with BD Cytofix/Cytoperm and permeabilized with BD Perm/Wash solution (cat# 554714, BD Pharmingen), stained with anti-IgG(H+L)-a350 (Polyclonal, Invitrogen) for 30 minutes. Cells were then washed once with BD Perm/Wash solution, once in FACS buffer, and analyzed by FACS.

Analysis of isotype-specific antibody reactivity to *Toxoplasma gondii* by flow cytometry

For serum reactivity analysis, syringe-lysed GFP-expressing strains (RH1-1 and GT1-GFP) were fixed in 3% formaldehyde for 20 minutes, washed twice in PBS, and plated in 96 well micro-titer plates at 4×10^5 parasites/well. The parasites were then incubated with serum from chronically infected mice, at serum concentrations ranging from 10^{-2} to 10^{-6} diluted in FACS buffer, for 20 minutes at 37°C. Parasites were then washed with FACS

buffer and placed on ice for incubation with anti-isotype detection antibodies depending

on application: anti-IgG3-BV421 (R40-82, BD Bioscience), anti-IgM-PE/Cy7 (RMM-1,

BioLegend), anti-IgG1-FITC (RMG1-1, BioLegend), anti-IgG1-APC (RMG1-1,

BioLegend), anti-IgG2b-FITC (RMG2b-1, BioLegend), anti-IgG2b-PE (RMG2b-1,

BioLegend), anti-IgG2a-FITC (RMG2a-62, BioLegend), anti-IgG2a-PerCP/Cy5.5

(RMG2a-62, BioLegend), anti-IgM-a-PE (DS-1, BD Bioscience), Anti-IgM-b-PE (AF6-78,

BD Bioscience).

Parasite neutralization assay

Heat-inactivated serum was used to coat live parasites for 20 minutes at 37°C before

infecting $5x10^5$ mouse embryonic fibroblasts/well (MEFs) in 96 well plates. Immediately

following addition of parasite to MEF wells, plates were spun at 1200rpm for 3 minutes to

synchronize infection. 2 hrs after initiation of infection, cells were placed on ice and

harvested by scraping with pipette tips. Cells were washed twice in FACS buffer,

suspended in 1:1000 PI in FACS buffer, and then analyzed by flow cytometry.

SDS-PAGE and immunoblotting for parasite lysate antigen

To generate parasite lysate antigens *Toxoplasma gondii* was cultured in HFF and

expanded to approximately $2x10^8$ parasites. Parasites were syringe-lysed, washed with

sterile 1X PBS and the parasite pellet was lysed with (1mL) 0.1% TritonX-100 detergent

in 1X PBS. Solubilized parasites were centrifuged at 2,000 RCF for 20 minutes to remove

large debris. The supernatant was aliquoted and stored at -80°C. Parasite lysate was

reduced with β-mercaptoethanol (BME) and separated via SDS-PAGE in 4-20% Mini-

PROTEAN TGX pre-cast gels (cat # 4561096, Bio-Rad) before transfer to PVDF

membrane using a Trans-Blot Turbo Mini PVDF Transfer Pack (cat # 1704156, Bio-Rad)

via Bio-Rad Transblot Turbo (cat # 1704150, Bio-Rad). Membranes were blocked with

10% fortified bovine milk dissolved in Tris-Buffered Saline with 0.1% Tween (TBS-T 0.1%)

for 1-2 hrs at room temperature or overnight at 4°C. Blots were then probed with heat-

inactivated serum in block at either 1:1,000 dilution for serum IgM analysis or 1:5,000

dilution for serum IgG analysis overnight at 4°C. Membranes were washed with TBS-T

0.1% three times for 20 minutes per wash. Blots were then incubated for one hr at room

temperature with goat α-mouse horseradish peroxidase (HRP)-conjugated antibodies

(SouthernBiotec): anti-IgM secondary 1:1000 (cat# 1020-05) and total anti-IgG secondary

1:5000 (cat# 1030-05). Membranes were then washed with TBS-T 0.1% three times and

developed with Immobilon® Forte Western HRP Substrate (WBLUF0500). All blots were

imaged via chemiluminescence on a ChemiDoc Touch (cat# 12003153, Bio-Rad). Image

Lab 6.1 software (Bio-Rad) was used for analysis of bands and total lane signal. Western

blots comparing A/J to C57BL/6J were developed simultaneously and the band signal was

normalized to A/J.

RNA isolation and sequencing

Peritoneal B-1a (B220$^{int-neg}$ CD19^{high} CD11b+ CD5+ PI-), B-1b (B220$^{int-neg}$ CD19^{high}

CD11b+ CD5- PI-), or B-2 B cells (B220^{high} CD19+ CD11b- PI-) were sorted into 500ul

RNeasy lysis buffer using a FACS ARIA II cell sorter (BD Biosciences). RNA was purified

using the RNeasy mini kit (cat# 74134, Qiagen) according to the manufacturer's protocol.

RNA purity was tested by Qubit (ThermoFisher) and Agilent 2100 BioAnalyzer for total

RNA with picogram sensitivity. DNA libraries were generated with a Lexogen QuantSeq-

UMI FWD 3′ mRNA-Seq Library Prep Kit (cat# 015). Samples were sent to UC Davis for

QuantSeq 3' mRNA FWD-UMI sequencing.

Gene expression analysis and data availability

Raw reads were trimmed and mapped by the BlueBee genomic pipeline FWD-UMI Mouse

(GRCm38) Lexogen Quantseq 2.6.1 (Lexogen). In brief, reads were quality controlled with

'FASTQC', trimmed with 'Bbduk' to remove low quality tails, poly(A)read-through and

adapter contaminations, read alignments to the *Mus musculus* genome build GRCm38

were done with 'STAR', and gene reads were quantified with 'HTSeq-count'. Differentially

expressed gene (DEG) analysis was performed utilizing limma-voom in R version 3.6.3 in

RStudio with Bioconductor suite of packages. Heatmaps were generated with 'gplots'.

Pathway and GO term analysis was performed with MouseMine (mousemine.org), and

gene set enrichment analysis was performed with GSEA v4.0.3. RNA-sequencing data

generated in this study has been deposited in the NCBI Sequence Read Archive

(Bioproject accession number PRJNA637442). All other data that support the findings

herein are available from the corresponding author upon reasonable request.

LPS-stimulation of enriched B cells and quantitative PCR

PerC and splenocytes were isolated from naïve 6-8 week-old C57BL/6J and A/J mice and

enriched for B cells using the EasySep Mouse Pan B cell Isolation Kit (cat#19844,

StemTech). Bead enrichment for splenic samples had the addition of biotinylated anti-

CD43 antibodies (clone S7, BD Bioscience) to remove B-1 cells. Enriched samples were

plated in a 96-well plate at 400,000 cells per well and stimulated with 25ug/ml of LPS (cat#

L4391-1MG, Sigma). After 2 hrs, total RNA was isolated using the Rneasy Mini Kit (cat#

74134) and cDNA was synthesized using High Capacity cDNA Reverse Transcription Kit

(ThermoFisher, cat#4368814). Quantitative PCR was performed on synthesized cDNA

samples using ThermoFisher TaqMan Master Mix (cat# 4444556) and TaqMan probes:

Actb - Assay ID:Mm02619580_g1 (cat# 4331182), and *Nfkbid* - Assay ID:

Mm00549082_m1 (cat# 4331182), according to the manufacturer's protocol.

Normalization of *Nfkbid* expression in each sample was calculated in comparison to *Actb*

expression levels. Fold change in *Nfkbid* expression of AJ relative to that of C57BL/6J

cells was determined through the delta delta CT method ($2^{-\Delta\Delta CT}$).

PerC adoptive transfers and IgH allotype chimeric mouse generation

PerC was harvested by peritoneal lavage of 6-12 week old C57BL/6J donor mice as

described above and 5×10^6 total peritoneal exudate cells (total PerC)/60ul PBS dose were

transferred i.p. into 2-4 day old bumble neonates. Allotype chimeric mouse generation was

performed as previously described (Lalor et al., 1989). In brief, 1-day old C57BL/6J

neonates were treated with 0.1mg of anti-IgM-b (clone AF6-78) and twice weekly

thereafter treated with 0.2mg of anti-IgM-b for 6 weeks. On day 2 after birth the neonates

were given 5×10^6 total PerC from B6.Cg-*Gpi1^a Thy1^a Igh^a* delivered i.p. The mice were

then allowed to rest for 6 weeks after the last antibody treatment before infection with *T.*

gondii.

Bone marrow chimeric mice generation

B6.SJL-*Ptprc^a Pepc^b*/BoyJ, bumble, and C57BL/6J recipient mice were given 2 doses of

500cGy with an X-Rad320 (Precision X-Ray) with a 4 hr interval. Donor BM cells were

harvested from bumble and WT C57BL/6J mice, filtered with a 70um filter, incubated in

ACK red blood cell lysis buffer, washed with PBS and transplanted by retro-orbital injection

at a concentration of 10^7 cells/200ul PBS dose. For BM chimeras reconstituted with PerC,

$5*10^6$ total PerC from WT C57BL/6J mice were transferred I.P. Recipient mice were then

allowed to rest for 8 weeks. Reconstitution was assessed at 8 weeks by FACS analysis of

PBL.

ELISA

High-affinity protein binding microplates (Corning) were coated overnight goat anti-mouse

IgM (1mg/ml, Southern Biotech, cat# 1021-01) and blocked with coating buffer containing

with 1% BSA (w/v) and 2% goat serum (Omega Scientific) in PBS. Wells were washed

with ELISA wash buffer (1X PBS, 0.05% TweenTM-20). Mouse serum samples were

diluted 1:400 in coating buffer and incubated in the wells for 1 hr. Wells were washed 5

times and secondary goat anti-IgM-HRP (Southern Biotech, cat# 1021-08) at 1:5000

dilution was incubated for 1 hr in the wells. Wells were washed several times and

developed with TMB substrate (Invitrogen). Development was stopped after 20 minutes

with 1M H_3PO_4 stop solution. Absorbance was measured at 450nm on a BioTek Epoch

microplate spectrophotometer.

Statistics

Statistical analysis was performed with Graphpad Prism 8 software. Statistical significance

was defined as $P < 0.05$. P values were calculated using paired or unpaired two-tailed t-

tests, 2-way ANOVA with Tukey multiple comparison correction, and multiple t-tests with

the Holm-Sidak correction for multiple comparisons. Survival curve significance was

calculated using log-rank (Mantel-Cox) testing. Significance in time to death was

calculated using the Gehan-Breslow-Wilcoxon test. Differential gene expression analysis

statistics were calculated using the Limma-Voom R package, P values were adjusted with

the Benjamini-Hochberg correction for multiple comparisons. GO term and pathway

enrichment analysis statistics were performed at mousemine.org using the Holm-

Bonferroni test correction. Statistical methods used for each figure are indicated in the

figure legends.

SUPPLEMENTAL INFORMATION

Figure S1. Cyst burden and weight during chronic infection with the low virulent

type III CEP strain correlates with the murine H-2 locus.

A/J, C57BL/6J (B6), C57BL/10J (B10), and C57BL/10.AJ (B10.A) mice were injected with

the type III strain *CEP hxgprt-* and allowed to progress to chronic infection. Plotted (+/-

SEM) is the average cyst number (x 1000) in the brain vs. the average fraction of initial

weight, where 1 is the normalized weight on the day of type III injection; the regression

value (R^2) is indicated. Results are from 3 to 5 mice for cyst numbers (day 42 of chronic

infection), and 5-12 mice for weight measurements (day 30 of chronic infection) per mouse

strain.

Figure S2. Consomic mice succumb to type I GT1 secondary infection.

Consomic mice of the C57BL/6J background with A/J chromosomal substitutions for

chromosome 7 (C57BL/6J-Chr7$^{A/J}$/NaJ) or chromosome 10 (C57BL/6J-Chr10$^{A/J}$/NaJ)

were infected with the type III CEP *hxgprt-* strain and allowed to progress to chronic

infection. Mice were then given a secondary infection with the type I GT1 strain.

Cumulative survival is shown for 2 independent experiments (CSS7 n=5, CSS10 n=6);

n.s., Mantel-Cox.

Figure S3. Bumble mice have intact T cell responses during secondary infection.

A) Peritoneal CD4 and CD8 T cells from bumble and C57BL/6J mice were assessed at

day 7 of secondary infection with the type I RH strain by an in vitro recall assay and

assayed for intracellular IFNγ and IL-2. In brief, peritoneal cells were harvested and

infected with live type I RH parasites for 16 hrs. T cells were assessed for production of

granzyme IFNγ, and IL-2 by intracellular staining and FACS. B) Peritoneal T-regulatory

cells (CD4+ CD25+ Foxp3+) were quantified at day 7 of secondary infection with type I

RH strain. Each dot represents the result from one mouse, and plotted are cumulative

averages +SD from 3 experiments; no significant differences were observed between

bumble and C57BL/6J mice by unpaired t-tests with the Holm-Sidak correction for multiple

comparisons.

Figure S4. Assessing PerC reconstitutions by serum IgM ELISAs and flow cytometry, and survival of muMT mice.

A) Serum IgM from C57BL/6J, muMT, bumble, and A/J mice was measured by ELISA. Serum was harvested from mice either naïve, chronically infected with type III CEP strain, or on D5 post-secondary infection with type I GT1 *T. gondii* strain. PerC transfer (+) refers

to mice adoptively transferred 5×10^6 total PerC cells as a day 2 neonate. Each dot

represents the results from an individual mouse, and plotted is the average +/-SD of the

O.D. obtained at 450nm; *P<0.05, unpaired two-tailed t-test. B) Bumble reconstitution of

the peritoneal B-1 compartment after neonatal PerC adoptive transfer. Representative

FACS plots of peritoneal B-2 cells (B220[high] CD19+) and B-1 (B220[int-neg] CD19+) cells from

bumble mice with or without PerC adoptive transfer. Shown are mice on day 20 of primary

infection with the type III CEP strain. C) B cell deficient muMT mice (n=3), muMT given

WT PerC adoptive transfers as 2-day neonates then allowed to reconstitute for 6-7 weeks

into adulthood (n=2), and muMT given B cell enriched splenocytes (n=3) 1 day prior to

infection with the type III CEP strain were assessed for survival. D) muMT reconstitution

of B-2 cell compartment, WT and muMT with B cell enriched splenocytes (EasySep™

Mouse Pan-B Cell Isolation Kit, cat# 19844) adoptively transferred 24 hrs earlier E) muMT

reconstitution of peritoneal B cell compartment after neonatal adoptive transfer.

Representative FACS plots of peritoneal B-2 cells (B220high CD19+) and B-1 B cells

(B220int-neg CD19+) from WT, and muMT mice or muMT mice with neonatal PerC

adoptive transfer. For D and E, uninfected mice are 6-8 weeks of age and numbers

indicate the percent of cells that fall within the depicted gate.

Figure S5. Bone marrow chimeric mice fail to survive primary infection but exhibit improved survival to vaccine strains.

A) Survival of the indicated bone marrow (BM) chimeras infected with the type III CEP *T. gondii* strain are plotted from a single experiment (n=2 for bumble recipients per condition; n=1 for C57BL/6J recipients); n.s., Mantel-Cox. B) Survival of the indicated BM chimeras vaccinated (10^6 i.p.) with the uracil auxotroph strain, Rh *Δup Δompdc*. Results are cumulative from 2-3 separate transfers and vaccinations; (n=4-9 per condition).

Figure S6. CD8 T cell IFNγ frequencies are increased in resistant A/J mice.

A) Peritoneal and B) splenic cells were harvested from A/J and C57BL6/J mice chronically

infected with type III CEP *T. gondii* strain, and infected with live type I parasites for 16 hrs.

T cells were assessed for production of granzyme B (GZB), IFNγ, and IL-2 by intracellular

staining and FACS. The average frequency +/-SD of positive staining CD4+ or CD8+ T

cells (CD3+ CD19-) and cumulative results from 2-3 experiments (AJ n=8, C57BL6/J

n=11) are shown; * P<0.05, unpaired two-tailed t-test.

Figure S7. Genes uniquely induced in B-1b cells on day 5 of secondary infection in resistant A/J mice.

Transcriptomic analysis of peritoneal B-1a (CD19+ B220$^{int-neg}$ CD11b+ CD5+), B-1b (CD19+ B220$^{int-neg}$ CD11b+ CD5-) and B-2 (CD19+ B220^{hi} CD11b- CD5-) B cells from A/J and C57BL/6J mice was performed using 3'-Tag RNA sequencing. A) Genes that were differentially upregulated in B-1b cells on day 5 of secondary infection compared to naïve mice in the A/J genetic background. P values of differentially expressed genes were calculated using the Benjamini-Hochberg adjustment for false discovery rate, and only those genes that survived significance were included in the heatmap. For comparison, all B-1 compartments in A/J mice are shown for this gene set. A/J B-1 Pathway and GO term enrichment was assessed on the genes presented in the heatmap in A. P values for enrichment analysis were adjusted with the Holm-Bonferroni correction. B) A cluster of genes found to be differentially induced in B-1b cells in A/J compared to C57BL/6J mice on day 5 of secondary infection are plotted as a heat map. C) Gene set enrichment analysis of the rank-ordered list of differentially expressed genes between A/J and C57BL/6J B-1b cells at D5 of secondary infection. Gene set depicted was in the top 10 gene sets ranked by false discovery rate (FDR) after investigating MSigDB's C7: immunologic signatures collection. Enrichment score is the degree of overrepresentation of a gene set at the top or bottom of a ranked list. NES is the enrichment score after normalizing for gene set size.

Mouse Strain	Chronic Infection	Secondary Infection	[1]Heterologous Brain Coinfection?
C57BL/10	CEP *hxgprt-*	GUYMAT	1/1
Cumulative average C57BL/10			1/1 = 100%
C57BL/10.A	CEP *hxgprt-*	TgCATBr5	2/2
C57BL/10.A	CEP *hxgprt-*	GUYMAT	2/2
C57BL/10.A	CEP *hxgprt-*	MAS	1/2
C57BL/10.A	CEP *hxgprt-*	FOU	0/1
Cumulative average C57BL/10.A			5/7 = 71%
A/J	CEP *hxgprt-*	GUYMAT	0/4
A/J	CEP *hxgprt-*	TgCATBr5	0/5
A/J	CEP *hxgprt-*	GUYDOS	4/5
A/J	CEP *hxgprt-*	VAND	4/6
A/J	CEP *hxgprt-*	FOU	1/5
A/J	CEP *hxgprt-*	GPHT	0/5
A/J	CEP *hxgprt-*	MAS	0/5
Cumulative average A/J			9/35 = 26%

1022

Table S1. Virulent strains of *Toxoplasma gondii* superinfect genetically resistant hosts.

CEP *hxgprt-* chronically infected C57BL/10, C57BL/10.A, and A/J mice were given a secondary infection with the indicated atypical *T. gondii* strain; then, 35-45 days after secondary infection, brains from surviving mice were homogenized in PBS and used to inoculate HFF monolayers. The resultant parasite cultures were then tested for the presence of the secondary infection strain; numerator = number of mice tested positive for secondary infection strain in the brain; denominator = number of mice that exhibited parasite growth in the HFF monolayer prior to drug selection.

1032   [1]Parasite growth was observed in mycophenolic acid / xanthine medium, which selects for

1033   parasites encoding a functional *HXGPRT* gene (i.e. the challenging atypical strains) and

1034   kills the primary infection CEP *hxgprt-* strain.

1035
1036

Recombinant Inbred Strain	Type III CEP Primary Infection Survival	Type I GT1 Secondary Infection Survival	[1]Heterologous Brain Coinfection?
AXB1	100	0	N/A
AXB2	100	100	0/2
AXB4	0	N/A	N/A
AXB5	50	100	1/1
AXB6	0	N/A	N/A
AXB8	100	50	1/1
AXB10	0	N/A	N/A
AXB12	100	100	2/2
AXB13	0	N/A	N/A
AXB15	100	100	1/2
AXB19	100	0	N/A
AXB23	100	0	N/A
AXB24	100	0	N/A
BXA1	50	0	N/A
BXA2	100	0	N/A
BXA4	100	100	0/1
BXA7	100	0	N/A
BXA8	100	100	2/2
BXA11	100	0	N/A
BXA12	100	100	1/1
BXA13	100	100	0/1
BXA14	100	0	N/A
BXA16	100	0	N/A
BXA24	100	50	N/A
BXA25	100	100	1/2
BXA26	100	100	0/2
Cumulative Average	82.6%	50%	9/17 = 52.9%

1037

1038

Table S2. Primary and secondary infection survival of recombinant inbred mice

(AxB;BxA).

26 strains (n = 2) from the RI (AxB; BxA) panel were infected with 10^4 type III avirulent

CEP *hxgprt- T. gondii* parasites; then, 35 days later, mice were challenged with $5x10^4$

virulent type I GT1 *T. gondii* parasites. Primary and secondary infection survival

percentages are indicated. 35-45 days after secondary challenge, brains from the

surviving RI mice were homogenized in PBS and used to inoculate HFF monolayers. The

resultant parasite cultures were then tested for the presence of the challenging strain;

numerator = number of mice tested positive for secondary infection strain in the brain;

denominator = number of mice that exhibited parasite growth in the HFF monolayer prior

to drug selection. Some surviving mice failed to generate parasite positive cultures.

[1]Parasite growth was observed in mycophenolic acid / xanthine medium, which selects for

GT1 parasites (the challenging strain) encoding the endogenous *HXGPRT* gene and

against the primary infection CEP *hxgprt-* strain.

1053

1054

Dataset S1. Polymorphic genes between A/J and C57BL/6J encoded within the four QTLs that define immunity to *Toxoplasma gondii*.

For each of the four QTLs that define secondary infection immunity to *T. gondii*, a list of all genes that have a DNA polymorphism between A/J and C5BL/6J mice, and are encoded within the QTL boundaries defined by the maximal genetic marker (real or imputed markers inferred in r/QTL) and the two flanking markers on either side; LOD scores and position are indicated. For each gene, its location and unique identifiers are listed, as well as the number and class of small nucleotide polymorphism (SNP) between A/J and C57BL/6J are shown. Total number of SNPs for each gene were also tallied ('Sum mutation'), and in bold are genes who's SNPs are two standard deviations greater than the average SNP for that class within the QTL region. In these QTL regions, there were also 476 (chr7), 88 (chr10), 40 (chr11) and 375 SNPs (chr17) that did not associate with any gene in the QTL boundaries. UTR, untranslated region; 'Region SNP', SNPs that are +/- 2 kb of the gene boundaries. Data obtained from Mouse Genome Informatics (MGI) (informatics.jax.org), and each locus is in a separate tab.

Dataset S2. Transcriptomic analysis of B cells responding to *Toxoplasma gondii* infection.

Table of genes included (threshold of >=5 CPM in any sample) in differential gene expression analysis. For each gene the Ensembl ID, MGI ID, gene symbol and descriptions are listed. The CPM from 3 samples of peritoneal B-1a, B-1b, and B-2 at naïve (N), chronic (Ch), and day 5 (D5) of secondary infection in both A/J (AJ) and C57BL/6J (B6) mice are given. The average CPM was calculated for each set of samples and average counts less than 5 CPM were raised to 5 to reduce the effect of low expression genes in further analysis. The log2 fold change (FC) of gene expression was

calculated using the average CPM for each cell type (B-1a, B-1b, and B-2) of each strain

(A/J and C57BL/6J) as they progressed from naïve to chronic infection or chronic to D5

secondary infection. The Log2 FC was also calculated from naïve to D5 of secondary

infection. The Log2 FC between A/J and C57BL/6J was calculated for each cell type.

Genes were then ranked by the number of comparisons that exhibited greater than 4FC

by the indicated comparison.

1086

1087   **REFERENCES**

1088

Alugupalli, K.R., Leong, J.M., Woodland, R.T., Muramatsu, M., Honjo, T., and Gerstein,
R.M. (2004). B1b lymphocytes confer T cell-independent long-lasting immunity.
Immunity *21,* 379-390.

Arnold, C.N., Pirie, E., Dosenovic, P., McInerney, G.M., Xia, Y., Wang, N., Li, X., Siggs,
O.M., Karlsson Hedestam, G.B., and Beutler, B. (2012). A forward genetic screen
reveals roles for Nfkbid, Zeb1, and Ruvbl2 in humoral immunity. Proc. Natl. Acad. Sci. U.
S. A. *109,* 12286-12293.

Baumgarth, N., Jager, G.C., Herman, O.C., and Herzenberg, L.A. (2000). CD4+ T cells
derived from B cell-deficient mice inhibit the establishment of peripheral B cell pools.
Proc. Natl. Acad. Sci. U. S. A. *97,* 4766-4771.

Berthier-Vergnes, O., El Kharbili, M., de la Fouchardiere, A., Pointecouteau, T.,
Verrando, P., Wierinckx, A., Lachuer, J., Le Naour, F., and Lamartine, J. (2011). Gene
expression profiles of human melanoma cells with different invasive potential reveal
TSPAN8 as a novel mediator of invasion. Br. J. Cancer *104,* 155-165.

Blandino, R., and Baumgarth, N. (2019). Secreted IgM: New tricks for an old molecule.
J. Leukoc. Biol. *106,* 1021-1034.

Brown, C.R., Hunter, C.A., Estes, R.G., Beckmann, E., Forman, J., David, C.,
Remington, J.S., and McLeod, R. (1995). Definitive identification of a gene that confers
resistance against Toxoplasma cyst burden and encephalitis. Immunology *85,* 419-428.

Casciotti, L., Ely, K.H., Williams, M.E., and Khan, I.A. (2002). CD8(+)-T-cell immunity
against Toxoplasma gondii can be induced but not maintained in mice lacking
conventional CD4(+) T cells. Infect. Immun. *70,* 434-443.

Castigli, E., Wilson, S.A., Scott, S., Dedeoglu, F., Xu, S., Lam, K.P., Bram, R.J., Jabara,
H., and Geha, R.S. (2005). TACI and BAFF-R mediate isotype switching in B cells. J.
Exp. Med. *201,* 35-39.

Chen, M., Mun, H.S., Piao, L.X., Aosai, F., Norose, K., Mohamed, R.M., Belal, U.S.,
Fang, H., Ahmed, A.K., Kang, H.K*., et al.* (2003). Induction of protective immunity by

primed B-1 cells in Toxoplasma gondii -infected B cell-deficient mice. Microbiol. Immunol. *47,* 997-1003.

Choi, Y.S., and Baumgarth, N. (2008). Dual role for B-1a cells in immunity to influenza virus infection. J. Exp. Med. *205,* 3053-3064.

Couper, K.N., Roberts, C.W., Brombacher, F., Alexander, J., and Johnson, L.L. (2005). Toxoplasma gondii-specific immunoglobulin M limits parasite dissemination by preventing host cell invasion. Infect. Immun. *73,* 8060-8068.

Fox, B.A., and Bzik, D.J. (2010). Avirulent uracil auxotrophs based on disruption of orotidine-5'-monophosphate decarboxylase elicit protective immunity to Toxoplasma gondii. Infect. Immun. *78,* 3744-3752.

Gazzinelli, R.T., Hakim, F.T., Hieny, S., Shearer, G.M., and Sher, A. (1991). Synergistic role of CD4+ and CD8+ T lymphocytes in IFN-gamma production and protective immunity induced by an attenuated Toxoplasma gondii vaccine. J. Immunol. *146,* 286-292.

Gigley, J.P., Fox, B.A., and Bzik, D.J. (2009). Cell-mediated immunity to Toxoplasma gondii develops primarily by local Th1 host immune responses in the absence of parasite replication. J. Immunol. *182,* 1069-1078.

Glatman Zaretsky, A., Silver, J.S., Siwicki, M., Durham, A., Ware, C.F., and Hunter, C.A. (2012). Infection with Toxoplasma gondii alters lymphotoxin expression associated with changes in splenic architecture. Infect. Immun. *80,* 3602-3610.

Grigg, M.E., Ganatra, J., Boothroyd, J.C., and Margolis, T.P. (2001). Unusual Abundance of Atypical Strains Associated with Human Ocular Toxoplasmosis. J. Infect. Dis. *184,* 633 639.

Ha, S.A., Tsuji, M., Suzuki, K., Meek, B., Yasuda, N., Kaisho, T., and Fagarasan, S. (2006). Regulation of B1 cell migration by signals through Toll-like receptors. J. Exp. Med. *203,* 2541-2550.

Hassan, M.A., Jensen, K.D., Butty, V., Hu, K., Boedec, E., Prins, P., and Saeij, J.P. (2015). Transcriptional and Linkage Analyses Identify Loci that Mediate the Differential Macrophage Response to Inflammatory Stimuli and Infection. PLoS Genet. *11,* e1005619.

Hoffman, S.L., Goh, L.M., Luke, T.C., Schneider, I., Le, T.P., Doolan, D.L., Sacci, J., de la Vega, P., Dowler, M., Paul, C*., et al.* (2002). Protection of humans against malaria by immunization with radiation-attenuated Plasmodium falciparum sporozoites. J. Infect. Dis. *185,* 1155-1164.

Hosokawa, J., Suzuki, K., Meguro, K., Tanaka, S., Maezawa, Y., Suto, A., Fujimura, L., Sakamoto, A., Clevers, H., Ohara, O., and Nakajima, H. (2017). IkappaBNS enhances follicular helper T-cell differentiation and function downstream of ASCl2. J. Allergy Clin. Immunol. *140,* 288-291.e8.

Howe, D.K., and Sibley, L.D. (1995). Toxoplasma gondii comprises three clonal lineages: correlation of parasite genotype with human disease. J. Infect. Dis. *172,* 1561-1566.

Itakura, A., Szczepanik, M., Campos, R.A., Paliwal, V., Majewska, M., Matsuda, H., Takatsu, K., and Askenase, P.W. (2005). An hour after immunization peritoneal B-1 cells

are activated to migrate to lymphoid organs where within 1 day they produce IgM
antibodies that initiate elicitation of contact sensitivity. J. Immunol. *175,* 7170-7178.

Jensen, K.D., Camejo, A., Melo, M.B., Cordeiro, C., Julien, L., Grotenbreg, G.M., Frickel,
E.M., Ploegh, H.L., Young, L., and Saeij, J.P. (2015). Toxoplasma gondii superinfection
and virulence during secondary infection correlate with the exact ROP5/ROP18 allelic
combination. MBio *6,* 2280.

Johnson, L.L., and Sayles, P.C. (2002). Deficient humoral responses underlie
susceptibility to Toxoplasma gondii in CD4-deficient mice. Infect. Immun. *70,* 185-191.

Joiner, K.A., Fuhrman, S.A., Miettinen, H.M., Kasper, L.H., and Mellman, I. (1990).
Toxoplasma gondii: fusion competence of parasitophorous vacuoles in Fc receptor-
transfected fibroblasts. Science *249,* 641-646.

Kang, H., Remington, J.S., and Suzuki, Y. (2000). Decreased resistance of B cell-
deficient mice to infection with Toxoplasma gondii despite unimpaired expression of IFN-
gamma, TNF-alpha, and inducible nitric oxide synthase. J. Immunol. *164,* 2629-2634.

Kantor, A.B., Stall, A.M., Adams, S., Herzenberg, L.A., and Herzenberg, L.A. (1992).
Differential development of progenitor activity for three B-cell lineages. Proc. Natl. Acad.
Sci. U. S. A. *89,* 3320-3324.

Khan, A., Jordan, C., Muccioli, C., Vallochi, A.L., Rizzo, L.V., Belfort, R.,Jr, Vitor, R.W.,
Silveira, C., and Sibley, L.D. (2006). Genetic divergence of Toxoplasma gondii strains
associated with ocular toxoplasmosis, Brazil. Emerg. Infect. Dis. *12,* 942-949.

Khoenkhoen, S., Erikson, E., Adori, M., Stark, J.M., Scholz, J.L., Cancro, M.P.,
Pedersen, G.K., and Karlsson Hedestam, G.B. (2019). TACI expression and plasma cell
differentiation are impaired in the absence of functional IkappaBNS. Immunol. Cell Biol.
97, 485-497.

Kim, C.C., Baccarella, A.M., Bayat, A., Pepper, M., and Fontana, M.F. (2019). FCRL5(+)
Memory B Cells Exhibit Robust Recall Responses. Cell. Rep. *27,* 1446-1460.e4.

Kreuk, L.S., Koch, M.A., Slayden, L.C., Lind, N.A., Chu, S., Savage, H.P., Kantor, A.B.,
Baumgarth, N., and Barton, G.M. (2019). B cell receptor and Toll-like receptor signaling
coordinate to control distinct B-1 responses to both self and the microbiota. Elife *8,*
10.7554/eLife.47015.

Krishnamurty, A.T., Thouvenel, C.D., Portugal, S., Keitany, G.J., Kim, K.S., Holder, A.,
Crompton, P.D., Rawlings, D.J., and Pepper, M. (2016). Somatically Hypermutated
Plasmodium-Specific IgM(+) Memory B Cells Are Rapid, Plastic, Early Responders upon
Malaria Rechallenge. Immunity *45,* 402-414.

Kuwata, H., Matsumoto, M., Atarashi, K., Morishita, H., Hirotani, T., Koga, R., and
Takeda, K. (2006). IkappaBNS inhibits induction of a subset of Toll-like receptor-
dependent genes and limits inflammation. Immunity *24,* 41-51.

Lalor, P.A., Herzenberg, L.A., Adams, S., and Stall, A.M. (1989). Feedback regulation of
murine Ly-1 B cell development. Eur. J. Immunol. *19,* 507-513.

Lorenzi, H., Khan, A., Behnke, M.S., Namasivayam, S., Swapna, L.S., Hadjithomas, M.,
Karamycheva, S., Pinney, D., Brunk, B.P., Ajioka, J.W*., et al.* (2016). Local admixture of
amplified and diversified secreted pathogenesis determinants shapes mosaic
Toxoplasma gondii genomes. Nat. Commun. *7,* 10147.

Luo, L., Bokil, N.J., Wall, A.A., Kapetanovic, R., Lansdaal, N.M., Marceline, F., Burgess, B.J., Tong, S.J., Guo, Z., Alexandrov, K.*, et al.* (2017). SCIMP is a transmembrane non-TIR TLR adaptor that promotes proinflammatory cytokine production from macrophages. Nat. Commun. *8,* 14133.

McLeod, R., Boyer, K.M., Lee, D., Mui, E., Wroblewski, K., Karrison, T., Noble, A.G., Withers, S., Swisher, C.N., Heydemann, P.T.*, et al.* (2012). Prematurity and severity are associated with Toxoplasma gondii alleles (NCCCTS, 1981-2009). Clin. Infect. Dis. *54,* 1595-1605.

McLeod, R., Skamene, E., Brown, C.R., Eisenhauer, P.B., and Mack, D.G. (1989). Genetic regulation of early survival and cyst number after peroral Toxoplasma gondii infection of A x B/B x A recombinant inbred and B10 congenic mice. J. Immunol. *143,* 3031-3034.

Mineo, J.R., McLeod, R., Mack, D., Smith, J., Khan, I.A., Ely, K.H., and Kasper, L.H. (1993). Antibodies to Toxoplasma gondii major surface protein (SAG-1, P30) inhibit infection of host cells and are produced in murine intestine after peroral infection. J. Immunol. *150,* 3951-3964.

Nguyen, T.T., Klasener, K., Zurn, C., Castillo, P.A., Brust-Mascher, I., Imai, D.M., Bevins, C.L., Reardon, C., Reth, M., and Baumgarth, N. (2017). The IgM receptor FcmuR limits tonic BCR signaling by regulating expression of the IgM BCR. Nat. Immunol. *18,* 321-333.

Nussenzweig, R.S., Vanderberg, J., Most, H., and Orton, C. (1967). Protective immunity produced by the injection of x-irradiated sporozoites of plasmodium berghei. Nature *216,* 160-162.

Pape, K.A., Taylor, J.J., Maul, R.W., Gearhart, P.J., and Jenkins, M.K. (2011). Different B cell populations mediate early and late memory during an endogenous immune response. Science *331,* 1203-1207.

Pedersen, G.K., Adori, M., Khoenkhoen, S., Dosenovic, P., Beutler, B., and Karlsson Hedestam, G.B. (2014). B-1a transitional cells are phenotypically distinct and are lacking in mice deficient in IkappaBNS. Proc. Natl. Acad. Sci. U. S. A. *111,* 4119.

Pedersen, G.K., Adori, M., Stark, J.M., Khoenkhoen, S., Arnold, C., Beutler, B., and Karlsson Hedestam, G.B. (2016). Heterozygous Mutation in IkappaBNS Leads to Reduced Levels of Natural IgM Antibodies and Impaired Responses to T-Independent Type 2 Antigens. Front. Immunol. *7,* 65.

Rowland, S.L., Leahy, K.F., Halverson, R., Torres, R.M., and Pelanda, R. (2010). BAFF receptor signaling aids the differentiation of immature B cells into transitional B cells following tonic BCR signaling. J. Immunol. *185,* 4570-4581.

RTS, S Clinical Trials Partnership. (2015). Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial. Lancet *386,* 31-45.

Sacks, D.L. (2014). Vaccines against tropical parasitic diseases: a persisting answer to a persisting problem. Nat. Immunol. *15,* 403-405.

Savage, H.P., Klasener, K., Smith, F.L., Luo, Z., Reth, M., and Baumgarth, N. (2019).
TLR induces reorganization of the IgM-BCR complex regulating murine B-1 cell
responses to infections. Elife *8,* 10.7554/eLife.46997.

Sayles, P.C., Gibson, G.W., and Johnson, L.L. (2000). B cells are essential for
vaccination-induced resistance to virulent Toxoplasma gondii. Infect. Immun. *68,* 1026-
1033.

Schuster, M., Annemann, M., Plaza-Sirvent, C., and Schmitz, I. (2013). Atypical IkappaB
proteins - nuclear modulators of NF-kappaB signaling. Cell. Commun. Signal. *11,* 23-23.

Schuster, M., Glauben, R., Plaza-Sirvent, C., Schreiber, L., Annemann, M., Floess, S.,
Kuhl, A.A., Clayton, L.K., Sparwasser, T., Schulze-Osthoff, K.*, et al.* (2012).
IkappaB(NS) protein mediates regulatory T cell development via induction of the Foxp3
transcription factor. Immunity *37,* 998-1008.

Smith, F.L., and Baumgarth, N. (2019). B-1 cell responses to infections. Curr. Opin.
Immunol. *57,* 23-31.

Splitt, S.D., Souza, S.P., Valentine, K.M., Castellanos, B.E., Curd, A.B., Hoyer, K.K., and
Jensen, K.D.C. (2018). PD-L1, TIM-3, and CTLA-4 blockade fail to promote resistance to
secondary infection with virulent strains of Toxoplasma gondii. Infect. Immun.

Stumhofer, J.S., Silver, J.S., and Hunter, C.A. (2013). IL-21 is required for optimal
antibody production and T cell responses during chronic Toxoplasma gondii infection.
PLoS One *8,* e62889.

Suzuki, K., Okuno, T., Yamamoto, M., Pasterkamp, R.J., Takegahara, N., Takamatsu,
H., Kitao, T., Takagi, J., Rennert, P.D., Kolodkin, A.L., Kumanogoh, A., and Kikutani, H.
(2007). Semaphorin 7A initiates T-cell-mediated inflammatory responses through
alpha1beta1 integrin. Nature *446,* 680-684.

Suzuki, Y., and Remington, J.S. (1990). The effect of anti-IFN-gamma antibody on the
protective effect of Lyt-2+ immune T cells against toxoplasmosis in mice. J. Immunol.
144, 1954-1956.

Touma, M., Antonini, V., Kumar, M., Osborn, S.L., Bobenchik, A.M., Keskin, D.B.,
Connolly, J.E., Grusby, M.J., Reinherz, E.L., and Clayton, L.K. (2007). Functional role for
I kappa BNS in T cell cytokine regulation as revealed by targeted gene disruption. J.
Immunol. *179,* 1681-1692.

Touma, M., Keskin, D.B., Shiroki, F., Saito, I., Koyasu, S., Reinherz, E.L., and Clayton,
L.K. (2011). Impaired B cell development and function in the absence of IkappaBNS. J.
Immunol. *187,* 3942-3952.

Waffarn, E.E., Hastey, C.J., Dixit, N., Soo Choi, Y., Cherry, S., Kalinke, U., Simon, S.I.,
and Baumgarth, N. (2015). Infection-induced type I interferons activate CD11b on B-1
cells for subsequent lymph node accumulation. Nat. Commun. *6,* 8991.

Yang, Y., Ghosn, E.E., Cole, L.E., Obukhanych, T.V., Sadate-Ngatchou, P., Vogel, S.N.,
Herzenberg, L.A., and Herzenberg, L.A. (2012). Antigen-specific antibody responses in
B-1a and their relationship to natural immunity. Proc. Natl. Acad. Sci. U. S. A. *109,* 5382-
5387.

Yap, G.S., and Sher, A. (1999). Effector cells of both nonhemopoietic and hemopoietic
origin are required for interferon (IFN)-gamma- and tumor necrosis factor (TNF)-alpha-

dependent host resistance to the intracellular pathogen, Toxoplasma gondii. J. Exp.
Med. *189,* 1083-1092.

Zhao, K., Wang, Z., Hackert, T., Pitzer, C., and Zoller, M. (2018). Tspan8 and
Tspan8/CD151 knockout mice unravel the contribution of tumor and host exosomes to
tumor progression. J. Exp. Clin. Cancer Res. *37,* 312-6.

1291