Nfkbid* is required for immunity and antibody responses to *Toxoplasma gondii

Scott P Souza[1], Samantha D Splitt[1], Julia A Alvarez[1], Safuwra Wizzard[1], Jessica N Wilson[1],

Zheng Luo[2], Nicole Baumgarth[2], Kirk DC Jensen[1#]

[1]School of Natural Sciences, University of California, Merced

[2]Center for Immunology & Infectious Diseases, and Department of Pathology, Microbiology and

Immunology, University of California, Davis

Correspondence: Kirk D.C. Jensen, kjensen5@ucmerced.edu

ABSTRACT

No fully protective vaccine has yet been developed for any human parasite. We previously

reported that South American strains of *Toxoplasma gondii* are adept at immune evasion and

cause lethal secondary infections in chronically infected mice. Here we report that unlike

susceptible C57BL/6J mice, A/J mice were highly resistant to secondary infection. Mapping

immunity loci that segregated within a recombinant inbred panel of mice we identified *Nfkbid*, a

nuclear regulator of NF-κB that is required for B-1 cell development. *Nfkbid*-null mice (bumble)

were susceptible to secondary, but not primary infections and failed to generate parasite-

specific IgM and lacked robust parasite-specific IgG. In resistant mice, B-1 cells bore evidence

for *Nfkbid*-dependent responses, including *Ighg* expression, and were partially responsible for

the immunity defect in bumble mice. We propose a model where adaptive immunity to *T. gondii*

must be 'layered' with B-1 responses to promote protection to virulent challenge.

INTRODUCTION

The goal of vaccination is to induce immunological memory that can protect from natural infection challenge. Depending on the pathogen, effective memory would need to protect also against a wide variety of pathogen-specific strains encountered in nature. Such protection is termed heterologous immunity, and is effective against pathogen strains that differ in virulence, immune evasion or polymorphic antigens. Parasites represent a special challenge to vaccine development. Indeed, an entirely protective vaccine has yet to be achieved for any human parasite [1]. The apicomplexan parasite *Toxoplasmas gondii*, provides an excellent system to explore requirements for heterologous immunity to a parasitic pathogen. *T. gondii* is a globally spread intracellular protozoan parasite of warm-blooded animals that exhibits great genetic diversity [2]. *T. gondii* strains differ dramatically in primary infection virulence in laboratory mice [3] and in severity of human toxoplasmosis [4-7]. Such infections can be overcome by immunological memory responses elicited by vaccination or natural infection. In particular, memory CD8 T cells and induction of IFNγ are primarily responsible for protection against lethal secondary infections with the widely studied type I RH strain, which has a lethal dose of one parasite in naïve mice [8-13]. CD4 T cells are required to help the formation of effector CD8 T cell [14, 15] and B cell responses [16], but the ability to adoptively transfer vaccine-elicited cellular immunity to naïve recipients against the type I RH challenge is unique to memory CD8 T cells [10, 11].

The role of B cells in *T. gondii* infections is less understood. Previous studies showed that B cell deficient mice (muMT) are extremely susceptible to primary [17], chronic [18] and secondary infections [19], despite unimpaired levels of IFNγ. Passive transfer of antibodies from immunized animals into vaccinated muMT mice significantly prolongs their survival after challenge [16, 19]. IgM seems particularly suited for blocking cellular invasion by *T. gondii* [20], while IgG can perform both neutralization [21] and opsonization functions [22-24]. Antibody responses against *T. gondii* are dependent on CD4 T cells [16, 25], and are regulated by cytokines that modulate T follicular helper cell and germinal center B cell formation in secondary lymphoid

organs [26, 27], suggesting conventional "B-2" B cell responses provide antibody-mediated

immunity to *T. gondii.*

In addition, "B-1" cells are innate-like lymphocytes that are known for producing self- and

pathogen-reactive "natural" IgM. B-1 cells are the predominant B cell compartment within the

body cavities, including the peritoneal and plural spaces and contribute to antigen-specific

responses to many pathogens (reviewed in [28]). In mouse models of secondary bacterial

infections, including *Borrelia hermsii* [29], *Streptococcus pneumoniae* [30] and non-typhoid

Salmonella [31]*,* vaccination induces protective memory B-1 cells to T cell-independent bacterial

antigens. This memory is often restricted to the B-1(b), or CD5- subset of B-1 cells [32], but not in

all models [33]. In the *T. gondii* model, one study suggested that primed CD5+ B-1(a) cells can

rescue B-cell-deficient mice during primary infection with a low virulence strain [17]. Memory B

cells are also appreciated to secrete pathogen-specific IgM [34], and generate somatically

mutated IgM to combat blood stage secondary infection with *Plasmodium* [35]. Whether IgM

responses to *T. gondii* are B-2 or B-1 derived is unknown. Moreover, the role of B-1 cells in

promoting immunity to *T. gondii* during a secondary infection has yet to be determined.

Particularly troubling for vaccine development for *T. gondii* is the lack of sterilizing

immunity achieved following infection [36, 37]. Unlike the highly passaged lab type I RH strain, for

which most immunological memory studies have been performed, the less passaged type I GT1

strain and atypical strains, many of which are endemic to South America, cause lethal

secondary infections in C57BL/6J mice and co-infect (i.e. 'superinfect') the brains of challenged

survivors [37]. During secondary infection, memory CD8 T cells become exhausted but checkpoint

blockade fails to reverse disease outcome [38]. The data suggest that yet unknown mechanisms

are needed to provide heterologous immunity to highly virulent strains of *T. gondii.* Therefore,

we set out to address whether additional requirements are necessary for heterologous immunity

to *T. gondii.* Through use of forward and reverse genetics, we discovered a previously

unidentified, essential role for *Nfkbid* in immunity and antibody responses to *T. gondii*, and

present evidence that B-1 cells may assist effective immunity to secondary infection with a

highly virulent parasite strain.

RESULTS

Non-MHC loci control resistance to secondary infection with *Toxoplasma gondii*

When mice are given a natural infection with a low virulent type III CEP strain and

allowed to progress to chronic infection for 35-42 days, the immunological memory that

develops is known to protect against secondary infections with the commonly studied lab strain,

type I RH. In contrast, "atypical" strains such as those isolated from South America (VAND,

GUYDOS, GUYMAT, TgCATBr5) or France (MAS, GPHT, FOU) and the clonal type I GT1

strain led to morbidity and mortality in C57BL/6J mice at varying frequencies, depending on the

strain type and their virulence factors [37]. To explore whether host genetics influenced the ability

to survive secondary infection, a similar experiment was performed with A/J mice. In contrast to

C57BL/6 mice, A/J mice were resistant to secondary infection with all virulent atypical *T. gondii*

strains analyzed (Fig 1A) and cleared parasite burden as early as day 4 post challenge (Fig 1B).

These results suggest that at least one or more genetic loci controls resistance to secondary

infection with *T. gondii* strains.

A/J mice are known to be resistant to primary infections with the intermediate virulent

type II strain [39, 40], and prevent cyst formation due to polymorphisms in their MHC class I H-2

molecule, L^d [41, 42]. To determine the extent to which the H-2a locus may contribute to A/J mice's

resistance to secondary infection we compared infections in C57BL/10 (B10) and C57BL/10.A

(B10.A) mouse strains, the latter carrying the MHC H-2 locus of A/J (H-2a) in place of that of

C57BL/6 (H-2b). Compared to mice with the H-2b haplotype, mice expressing H-2a had less

cysts and weighed more during chronic infection with type III strains (P<0.04; Fig S1). Despite

their relative health at the time of challenge, B10.A mice were highly susceptible (0%-30%

survival) to secondary infection with certain atypical strains (VAND, GUY-DOS, GPHT, FOU),

but displayed varying degrees of resistance to others (TgCATBr5, MAS, GUY-MAT; 60-100%

survival) (Fig 1A). In the case of MAS secondary infection, B10.A mice but not B10 mice were

highly resistant and exhibited reduced parasite burden by day 8 post challenge (Fig 1B).

Together, the data suggest that while the MHC H-2a locus is an important modifier of resistance

to certain *T. gondii* strains, this is not true for every encounter. Importantly, however, the A/J

genetic background encodes additional non-MHC-linked genes that control immunity to *T.*

gondii.

Genetic mapping reveals four loci that correlate with immunity to *Toxoplasma gondii*

To identify non-MHC loci that promote resistance to secondary infection, we first

analyzed the outcome of infection with the type I GT1 *T. gondii* strain, as this strain caused a

lethal secondary infections in B10 and B10.A, but not in A/J or first filial generation A/J x

C57BL/6J mice ('F1') (Fig 1C). Following secondary infections A/J and F1 mice showed no overt

symptoms of weight loss, dehydration, or lethargy (not shown). However, sterile immunity was

not achieved. For example, the GT1 strain was present in the brains of these survivors (i.e.

'superinfection'), and at greater frequencies in F1 compared to A/J mice (Fig 1D).

Superinfections were also detected at high frequencies in B10 and B10.A surviving mice

challenged with atypical strains. Overall, mice of the C57BL genetic background were more

prone to superinfection compared to A/J mice (Table S1). It is unknown whether virulent strains

of *T. gondii* have evolved to superinfect hosts with immunological memory, as previously

hypothesized [37]. Nonetheless, our results underscore the difficulty in achieving sterile immunity

to parasites in otherwise genetically 'resistant' hosts.

Then we performed secondary infection experiments with the type I GT1 strain using 26

recombinant inbred (RI) mice (Table S2). The AxB:BxA RI mouse panel contains an assortment

of homozygous A/J and C57BL/6 alleles, which assist genetic mapping of loci that contribute to

various phenotypes, including those related to *T. gondii* infection [39, 43, 44]. Genetic mapping

revealed four distinct Quantitative Trait Loci (QTL) peaks with logarithm of the odds (LOD)

scores greater than 3 on chromosomes 7, 10, 11 and 17 (Fig 2A). None of the QTLs bore

evidence for epistatic interactions (not shown), and only the chromosome 10 QTL surpassed

genome-wide permutation testing (n=1000, P<0.05). Nevertheless, an additive-QTL model

including all four QTLs best fit the data compared to any lesser combination of them (P<0.02,

ANOVA). The estimated effect on the phenotypic variance observed in the RI panel is 24%,

41%, 21% and 27% for the chromosome (chr) 7, 10, 11 and 17 QTLs, respectively. Consistent

with these estimates, complete phenotypic penetrance (i.e. allelic correlation at 100%) was not

observed for any locus (Fig 2B). Regardless, small effect QTLs controlling complex traits can

still lead to the identification of causal variants within a QTL region, as occurred for the

successful identification of MHC 1 L^d as the host resistance factor to chronic *T. gondii* infection

[41].

Nfkbid is one of the most polymorphic genes within the chr7 QTL region (Mb 30.3-33.0)

and sits between the genetic markers that flank the highest imputed LOD score (Table S3). This

gene encodes IκBNS, which is a member of atypical NF-κB inhibitors, called nuclear IκBs,

reflecting their restricted cellular localization. Unlike classical inhibitors of NF-κB, atypical NF-κB

inhibitors can modulate NF-κB to induce or repress transcription [45]. Previous work has shown

that *Nfkbid* null mice completely fail to develop B-1 cells [46, 47], lack circulating IgM and IgG3

antibodies, and cannot respond to T-independent type II antigens such as NP-ficoll [46-48]. IκBNS

also promotes B-2 cell activation and isotype class switching [46], enhances T cell production of

IL-2 and IFNγ [49], supports development of T regulatory cells [50] and suppresses TLR-induced

cytokine expression in macrophages [51]. The tetraspanin, *Tspan8,* is within the chr10 QTL (Mb

115.8-116.2) and is the most highly polymorphic gene between A/J and C57BL/6J mice in this

region (Table S3). *Tspan8* is 6-fold more highly expressed in spleens from A/J compared to

C57BL/6 mice (immgen.org). *Tspan8* promotes cancer metastasis [52] and can impact leukocyte

migration [53], but its role in immunity is largely unknown. Other polymorphic gene candidates

within the four QTLs are listed in Table S3.

Nfkbid on chromosome 7 is required for immunity and the generation of *Toxoplasma*

***gondii*-specific antibodies**

Given the role of *Nfkbid* in several immune functions, polymorphism and central location

within the chr7 QTL, the requirement for *Nfkbid* in immunity to *T. gondii* was further explored.

Nfkbid null "bumble" mice (C57BL/6J) have previously been described, which were derived from

an ENU mutagenesis screen. These mice possess a premature stop codon in *Nfkbid,* rendering

them unable to support T-I antibody responses and B-1 development [47]. Bumble mice survived

primary infection with the low virulent CEP strain at frequencies similar to wildtype C57BL/6J

mice (Fig 3A), but succumbed to secondary infection three days earlier when challenged with

the GT1 strain (Fig 3B). Since the C57BL genetic background is uniformly susceptible to GT1

secondary infections used in our genetic screen, susceptibility to challenge with the commonly

studied type I RH strain was explored, which is normally controlled in vaccinated or chronically

infected mice [8-13]. Importantly, bumble mice were uniformly susceptible to secondary infection

with the type I RH strain (Fig 3C), which exhibited greater parasite loads compared to wildtype

mice (Fig 3D). Moreover, bumble mice generated strongly reduced parasite specific-IgM, IgG3,

IgG2b and IgG2a antibody responses after chronic infection (Fig 4A-4C). The remaining

antibodies that were secreted exhibited defects in their ability to block parasite invasion of host

cells (Fig 4D). Antibodies from naïve mice fail to bind *T. gondii* (Fig 4C), thus natural antibodies

do not recognize *T. gondii,* consistent with previous reports [20]. Although *Nfkbid* promotes T cell

production of IFNγ and IL-2 in in vitro stimulation assays [49] and promotes thymic development of

FOXP3+ T regulatory cells [50], no impairment of T cell cytokine production was observed, nor

were frequencies of FOXP3+ CD25^{hi} CD4+ T regulatory cells altered in bumble compared to

wildtype mice during secondary infection (Fig S2). Hence, defects in B cell-mediated immunity

appear to underlie the susceptibility of bumble mice to secondary infection with *T. gondii*.

Evidence for enhanced B-1 cell activation in resistant A/J mice

To generate a comprehensive view of immune response differences between A/J and

C57BL/6 mice, and to refine investigation of the chr7 QTL's contribution to *T. gondii* immunity,

immune-phenotyping was employed during chronic and secondary infections. Consistent with

the increased parasite burdens observed in susceptible mice (Fig 1B), greater frequencies of

neutrophils, inflammatory macrophages, and dendritic cells were all enhanced in the peritoneal

cavities and to a lesser extent within the spleens (not shown) of C57BL/6J compared to A/J

mice on day 5 of secondary infection (Fig S3). Phenotypes associated with *Nfkbid* were further

explored: T cell cytokine production (IL-2 and IFNγ) and the B-1 cell compartment. Although

CD4 and CD8 T cell frequencies were not significantly different between the resistant and

susceptible mouse backgrounds (Fig S3, not shown), an increase in peritoneal CD8 T cell in

vitro recall production of IFNγ was noted between A/J and C57BL/6J mice, but other

parameters, including granzyme B and IL-2 expression were similar (Fig S4). Importantly, within

the peritoneal B-1 cell compartment (CD19+, B220^{int}, CD11b+) increased percentages of CD5-

B-1b cells were observed during secondary infection in resistant A/J mice, which appear to have

downregulated their BCR as evidenced by being IgM^{lo} (Fig 5A, 5B). To further validate these

findings, frequencies of CD5+ (B-1a) and CD5- (B-1b) cells within the CD43+ B-1 cell

compartments of the peritoneum and spleen were determined and a similar trend was observed

(Fig 5C, 5D). In the spleen, both strains of mice had significant increased frequencies of IgM-

IgD- B-1a and B-1b cells in chronically infected compared to naïve animals, while in the

peritoneum this increase was observed following challenge (Fig 5E). In these situations,

resistant A/J mice often had two-fold greater percentages of IgM-IgD- B-1 cells relative to the

susceptible C57BL/6 mice. Following infection, B-1 cells in the spleens of A/J mice maintained

high levels of BAFFR and TACI expression and expressed higher levels of surface CD138

relative to C57BL/6J mice (Fig 5D), markers known to be induced by *Nfkbid* and important for B

cell activation and differentiation into antibody secreting cells [54]. Finally, the overall parasite-

specific serum antibody response to parasite lysate was assessed by western blot. Serum-

derived antibodies from the resistant A/J background had significantly greater lysate antigen

reactivity, in particular IgG during secondary infections, compared to C57BL/6J (Fig S4).

Altogether, the data suggest that CD8 T cell production of IFNγ is enhanced in resistant mice

and likely contributes to their resistance. However, the data further suggest that B-1 cells

participate in enhanced B cell responses in A/J mice, offering a new modality of immunity

against parasitic infections. Importantly, the data lends credence to the hypothesis that the

causal gene under the chr7 QTL is *Nfkbid.*

B-1 cells in resistant A/J mice have enhanced germline transcription of *Ighg* constant

regions and different activation profiles

Bumble mouse susceptibility and inability to generate parasite-specific antibodies,

prompted us to investigate further how *Nfkbid* may be modulating the B cell compartment. A

transcriptomic approach was taken to define peritoneal B-1a, B-1b and B-2 cell response

characteristics in resistant and susceptible mice during naïve, chronic and secondary infection

states. The top differentially expressed genes commonly induced in B-1 and B-2 cells following

infection are represented as a heatmap (Fig 6A), and the entire data set can be found in Table

S4. GO term enrichment analysis found type I and II interferon signaling and immune defense

signatures, including "response to protozoans", as being enriched in the most differentially

regulated genes among B cells following infection (Fig 6B), an observation consistent with the

robust T cell IFNγ responses induced by *T. gondii* (Fig S4). Looking specifically at CD5- B-1

cells in A/J mice, which bore evidence for enhanced activation (Fig 5), pathway enrichment

analysis of genes differentially upregulated on day 5 of secondary infection compared to the

naïve state found signatures of TLR-signaling, complement activation and somatic

recombination (Fig S6). Germline transcription of *Ighg1*, *Ighg3*, *Ighg2b*, and *Ighg2a/c* was

greatly enhanced in CD5- B-1 cells from A/J compared to C57BL/6J mice on day 5 of secondary

infection (Fig 6C), suggesting heightened isotype class switching occurred in mice on the

resistant background, consistent with their IgM-IgD- phenotype (Fig 5). Moreover *Cxcr3,* an

important chemokine receptor for promoting Th1 and CD8 T cell responses during *T. gondii*

infection [55-57], *Scimp* an adaptor for TLR4 signaling [58], *Semaphorin7* a noted inducer of

inflammatory cytokines [59], the alarmins *S100a8* and *S100a9*, and genes associated with tissue

tolerance including *Retnlg* and *Slpi,* were each specifically upregulated in CD5- B-1 cells from

A/J compared to C57BL/6 mice on day 5 of challenge (Fig S6). Gene Set Enrichment Analysis

(GSEA) of CD5- B-1 transcriptional variation between mice on day 5 of challenge detected

correlation with gene sets that distinguish B cells following vaccination with different TLR-

stimulating adjuvants (MPL vs R848), follicular vs early germinal center B cells, and tumor-

associated immuno-suppressive dendritic cells (Fig 6D). In summary, the transcriptomics data

suggest that the entire B cell compartment of both strains of mice experienced IFN-signaling

during *T. gondii* infection. In the resistant A/J background but not in C57BL/6J mice, B-1 cells

underwent Ig class switch recombination and were activated, perhaps through enhanced TLR-

signaling.

Gene dosage of *Nfkbid* impacts parasite-specific IgG1 responses

The altered antibody responses to secondary infection in the bumble mice as well as the

B cell phenotypic differences noted in A/J mice suggested that *Nfkbid* may control *T. gondii*-

specific antibody responses, and that this may be modulated by expression of *Nfkbid.* This is

supported by the observation that most *Nfkbid* mutations between A/J and C57BL/6J mice are

non-protein coding. This includes multiple mutations in the 3'UTR which can determine

transcript stability (Table S3). To investigate this possibility, we quantified *Nfkbid* expression

within our transcriptomic dataset as well as by qPCR. B cells from C57BL/6J mice had greater

expression of *Nfkbid* relative to A/J, particularly at the chronic infection stage (Fig 7A). Because

Nfkbid expression differences could reflect heightened responsiveness to parasite load, we

stimulated enriched peritoneal and splenic B cells from A/J and C57BL/6J mice with LPS and

noted that B cells from C57BL/6J mice had on average 1.5- to 2-fold greater induction of *Nfkbid*

transcripts relative to A/J (Fig 7B).

To explore how gene dosage of *Nfkbid* impacts antibody responses generated against *T.*

gondii, serum from chronically infected bumble heterozygotes (*Nfkbid*+/-), C57BL/6J, and A/J

mice were analyzed. Although parasite-specific IgM and IgG3 did not differ between mouse

strains, increased parasite-specific IgG1 and IgG2a/c serum responses were observed in A/J

relative to C57BL/6 mice, a response that was phenocopied in particular for IgG1 in bumble

heterozygotes (Fig 7C-D). In contrast, heterozygous expression of *Nfkbid* is sufficient to restore

wildtype levels of parasite-specific IgM, IgG3, and IgG2b in mice with a C57BL/6 background

when compared to bumble mice (Fig 4C, 7D). Hence, while *Nfkbid* is required for the generation

of IgM and robust IgG responses against *T. gondii*, gene dosage of *Nfkbid* further controls IgG1

isotype profiles, presenting *Nfkbid* as tunable modulator of antibody IgG responses to parasites.

Of note, when *Nfkbid+/-* mice were given secondary infections with the GT1 strain, all mice

succumbed to the challenge (not shown), observations that are likely consistent with a multiple-

QTL model for *T. gondii* immunity and apparent need for additional modifiers on chromosomes

10 and 17 to survive highly virulent challenges.

B-1 cells promote but do not fully restore immunity to *Toxoplasma gondii*

The impaired ability of bumble mice to generate parasite-specific antibodies, combined

with the documented collapse of the B-1 cell compartment in *Nfkbid*-deficient mice [46, 47],

prompted us to directly asses the role of B-1 mediated immunity to *T. gondii*. First*,* total

peritoneal exudate cells (PerC) were adoptively transferred into bumble mice at day 2 of birth,

which allows optimal B-1 engraftment and self-renew for the life of the animal [60]. Then, bumble

mice that received total PerC transfers were infected with the avirulent type III strain at 6-7

weeks of age and given a secondary infection 35 days later with the type I RH strain (Fig 8A).

Bumble mice receiving PerC partially reconstituted serum IgM to ~40% of wildtype levels (Fig

S7), consistent with previous studies [60], and had a significantly delayed time to death relative to

non-transferred littermates following type I RH challenge (Fig 8A). Previous studies have shown

B-1 cells respond to infections and create both pathogen- [29, 30, 32, 61] and microbiota -specific

antibodies [62]. Antibody profiling of bumble mice that received PerC transfer showed a trend of

increased anti-*T. gondii* antibody generation following chronic infection (Fig 8B), but antibody

reactivity to the parasite did not reach levels observed in wildtype mice.

These results were further confirmed with Igh-allotype chimeric mice which allow

tracking of IgM responses of allotype-marked B-1 and B-2 cells [63]. In this experimental setup,

endogenous B cells of the C57BL/6 background (IgH-b allotype) are depleted with allotype

specific anti-IgM-b antibodies and replaced with transferred PerC B-1 of the IgH-a allotype

which are refractory to the depletion antibodies and will engraft for the life of the animal.

Following removal of the depleting antibodies, the endogenous B-2 cell population reemerge

and are marked with anti-IgM-b antibodies, while the transferred B-1 cell are marked with anti-

IgM-a antibodies (Fig 8C). Assessing antibody responses generated in IgH-allotype chimeric

mice during a *T. gondii* infection revealed the presence of B-1 derived IgM-a antibodies that had

low reactivity to *T. gondii* at days 14 and 30 post-infection, but the majority of highly reactive

parasite-specific IgM-b was derived from the B-2 compartment (Fig 8D). Other attempts were

made to explore the role of B-1 and B-2 cells, utilizing B cell deficient muMT mice, but these

mice were highly susceptible to primary infection with the CEP strain irrespective of whether

they received PerC as neonates or splenic B-2 cells prior to the primary infection (Fig S8),

underscoring the importance of B cells in resistance to *T. gondii* infection [18]. Mixed bone marrow

chimeras where also attempted, in which CD45.1 recipients were given a 1:1 mixture of muMT

and wildtype or bumble adult bone marrow to generate mice in which the B-2 compartment

expressed or lacked the *Nfkbid* gene, respectively. B-1a cells do not efficiently reconstitute

irradiated recipients from adult bone marrow [64]. Though reconstitution occurred, the mice

succumbed to primary CEP infections (Fig S9).

DISCUSSION

Our data strongly implicate *Nfkbid* in B cell-mediated responses to *T. gondii* and

underscore the utility of using inbred mouse panels to uncover novel determinants of immunity

to parasites. While B-1 cells are partially responsible for the defect in immunity observed in

bumble mice, antigen-specific B-2 cells likely need assistance from their B-1 counterparts to

provide full immunity to challenge. We reason this because in none of the attempted

reconstitutions did recipients receive both *Nfkbid* sufficient B-1 *and* B-2 cells. The ability of B-1

cells to make parasite-specific antibodies, though of lower affinity, likely assist B-2 immune

responses to *T. gondii,* which internalize antigen-antibody complexes secreted from the B-1

compartment [65], assisting MHCII antigen presentation for CD4 T cell help [66]. It is also possible

the effect of B-1 cell-mediated immunity to *T. gondii,* and possibly to other pathogens, is

underestimated in murine models using the C57BL/6 background, as shown here for the

enhanced B-1 responses observed in A/J mice.

Though the exact pathway remains to be investigated, *Nfkbid* is downstream of TLR

signaling in B cells [46, 47, 54], which causes B-1a cells to downregulate CD5 and facilitate

differentiation into antibody secreting cells [62, 67]. CD5 is a potent negative regulator of antigen

receptor signaling that renders B-1a cells unresponsive to B cell receptor (BCR)-triggering [68].

This inhibition is overcome by TLR-stimulation which causes CD5 to dissociate with the BCR,

thereby releasing repression of BCR-mediated signaling and antibody secretion [67] against

foreign- and self-antigen [62]. These data suggest CD5- B-1b cells may actually represent an

activated state of B-1 cells, and calls into question a strict division of labor between these two

subsets [69]. This supposition would fit several of the observations made in our system, including

evidence for enhanced class switch recombination and TLR-gene signatures observed in the

CD5- B-1 cells of the resistant background. In addition, BAFFR and TACI are known inducers of

class-switch recombination [70], and increased expression of these receptors may further lower

the threshold of activation and differentiation into CD138+ plasmablasts/plasma cells, all of

which are regulated by *Nfkbid* [54] and occurring with greater magnitude in B-1 cells of genetically

resistant A/J mice. Further investigation of pathways upstream of *Nfkbid,* such as the TLR

pathway, has the potential to elucidate key requirements for *T. gondii* immunity.

Heterologous immunity to a parasitic pathogen should, at a minimum, prevent disease

against a wide variety of strains that differ in virulence or polymorphic antigens. An ideal

parasite vaccine would entirely protect against re-infection and induce sterile immunity, thought

possible since re-infection studies were first performed in mice [71] and humans immunized with

irradiated sporozoites and *Plasmodium sp.* challenge [72-75]. Yet, only one partially protective

vaccine is in use for any human parasitic pathogen, RTS,S/AS01, which has low efficacy for

malaria prevention [76]. Our results demonstrate that even in genetically resistant hosts, sterile

immunity was not always achieved. Other loci that provide heterologous immunity to *T. gondii*

include the MHC of A/J to certain strains like MAS and GUY-MAT. Whether *Tspan8* is the

causal gene within the chr10 QTL (Mb 115.8-116.2) is unknown. *Tspan8* is one of four

tetraspanins that encode four cysteines, including CD9 and CD81 known to influence B cell

activation [77], but besides one report on impacting leukocyte migration (in the B6 background) [53],

its role in immune function is unknown. *Tspan8* expression is highly expressed in epithelial cells,

and we found no evidence for its expression in B cells by transcriptomic analysis and qPCR (not

shown). Tetraspanins interact with integrins and play diverse roles in lymphocyte function [77]. It is

tempting to speculate that Tspan8 interacts with an unidentified lymphocyte integrin to mediate

migration to the site of infection.

The focus of this study was to use an unbiased genetic screen to find requirements for

heterologous immunity to the most virulent strains *T. gondii*. Our findings highlight the role of B-

1 cells in immunity to *T. gondii*, introducing this population of B cells as a potential target for

vaccines to induce antibody responses against parasitic infections. Moreover, we present a

modulator of antibody responses against parasitic infections, *Nfkbid*, a transcriptional regulator

that can tune B cell responses to provide an overall effective class-switched antibody response

against parasites.

ACKNOWLEDGEMENTS

This work was supported by NIH grants R01 AI137126, R15 AI131027, a Hellmann Fellowship

and an MCB departmental award to KJ; and by NIH grants U19-AI109962 and R01 AI117890 to

NB. The sequencing was carried out at the DNA Technologies and Expression Analysis Cores

at the UC Davis Genome Center, supported by NIH Shared Instrumentation Grant

1S10OD010786-01. The authors declare no conflict of interest.

MATERIAL AND METHODS

Parasite strains and cell lines

Human foreskin fibroblasts (HFFs) monolayers were grown in DMEM (4.5 g/L D-glucose) (Life

Technologies) supplemented with 2 mM L-glutamine, 20% fetal bovine serum (FBS) (Omega

Scientific), 1% penicillin-streptomycin, and 0.2% gentamycin (Life Technologies). Mouse

Embryonic Fibroblasts (MEFs) were grown in DMEM (4.5 g/L D-glucose) (Life Technologies)

supplemented with 10% fetal bovine serum (FBS) (Omega Scientific), 20mM HEPES, 1%

penicillin-streptomycin, and 0.2% gentamycin (Life Technologies). *Toxoplasma gondii* strains

were passaged in HFFs in 'Toxo medium' (4.5 g/L D-glucose, L-glutamine in DMEM

supplemented with 1% FBS and 1% penicillin-streptomycin). The following clonal strains were

used (clonal types are indicated in parentheses): RH Δ*ku80* Δ*hxgprt* (type I), RH (1-1)

GFP::cLUC (type I), GT1 (type I), GT1 *GFP::cLuc* (type I), and CEP *hxgprt-* (type III). The

following atypical strains were used: MAS, MAS *GFP::cLuc* (2C8) (haplogroup 'HG' HG4), GUY-

MAT (HG5), FOU (HG6), GPHT (HG6), TgCATBr5 (HG7), GUY-DOS (HG10), and VAND

(HG10).

Generation of GFP-expressing GT1 strains

GT1 parasites were transfected with linearized plasmids for parasite expression of GFP and

click beetle luciferase (*GFP::cLUC*), parasites were grown on HFF monolayers in T-25 flasks in

Toxo medium for 2 weeks. Parasites were removed from the flasks by scraping; the parasites

were pelleted and washed with PBS and suspended in sterile FACS buffer (2% FBS in PBS).

Fluorescent parasites where then sorted via fluorescence-activated cell sorting (FACS) into a

96-well plate with confluent HFF monolayers. To ensure single plaque formation in at least one

of the wells, the sort was titrated using the following parasite numbers: 100, 50, 25, 12, 6, 3, 2,

and 1 for each well per row of 8.

Mice and ethics statement

Female C57BL/6J (H-2b), A/J (H-2a), C57BL/10SnJ (H-2b), B10.A-*H2^a H2-T18^a*/SgSnJ (H-2a),

B6AF1/J (A/J x C57BL/6J F1 progeny), B6.129S2-*Ighm^{tm1Cgn}*/J (muMT), B6.SJL-*Ptprc^a*

Pepc^b/BoyJ (CD45.1 congenic), B6.Cg-*Gpi1^a Thy1^a Igh^a*/J (IgH-a triple congenic mice) and 26

(AxB;BxA) recombinant inbred (RI) mice derived from A/J and C57BL/6 founders, were

purchased from Jackson Laboratories. The bumble mouse line used for this research project [47],

C57BL/6J-*Nfkbid^{m1Btlr}*/Mmmh, RRID:MMRRC_036725-MU, was obtained from the Mutant

Mouse Resource and Research Center (MMRRC) at University of Missouri, an NIH-funded

strain repository, and was donated to the MMRRC by Bruce Beutler, M.D., University of Texas

Southwestern Medical Center. Bumble mice were crossed to C57BL/6J to generate F1 bumble

heterozygotes (*Nfkbid*+/-).

Mice were maintained under specific pathogen free conditions at UC Merced. Every

effort was made to ensure unnecessary stress on the animals was avoided. Mouse work was

performed in accordance with the National Institutes of Health Guide to the Care and Use of

Laboratory Animals. All protocols have been reviewed and approved by UC Merced's

Committee on Institutional Animal Care and Use Committee. UC Merced has an Animal Welfare

Assurance filed with OLAW (#A4561-01), is registered with USDA (93-R-0518), and the UC

Merced Animal Care Program is AAALAC accredited (001318).

Parasite infections

Parasite injections were prepared by scraping T-25 flasks containing vacuolated HFFs and

sequential syringe lysis first through a 25G needed followed by a 27G needle. The parasites

were spun at 400 rpm for 5 min to remove debris and the supernatant was transferred, followed

by a spin at 1700 rpm washing with PBS. For primary infections, mice were infected

intraperitoneally (i.p.) with 10^4 tachyzoites of type III CEP *hxgprt-.* For secondary infections mice

were infected I.P. with $5x10^5$ type I parasites (RH or GT1). Parasite viability of the inoculum was

determined by plaque assay following i.p. infections. In brief, 100 or 300 tachyzoites were plated

in HFF monolayers grown in a 24-well plate and 4-6 days later were counted by microscopy (4x

objective).

Blood plasma isolation and assessment of seroconversion

All mice were assessed for sero-positivity to *T. gondii* 4-5 weeks post primary infection. 50µL of

blood was isolated from mice in tubes containing 5µL of 0.5M EDTA on ice, pelleted and the

supernatant containing blood plasma was heat inactivated to denature complement at 56°C for

20 minutes and then stored at -80°C. HFFs were grown on coverslips and infected with GFP-

expressing RH (1-1) overnight, fixed 18 hrs later with 3% formaldehyde (Polysciences) in PBS,

washed, permeabilized and blocked with PBS containing 3% bovine serum albumin Fraction V

(Sigma), 0.2M Triton X-100, 0.01% sodium azide, incubated with a 1:100 dilution of collected

blood plasma for 2 hrs at room temperature, washed with PBS, and detected with Alexa Fluor

594-labeled secondary antibodies specific for mouse IgG (cat # A11032, Life Technologies).

Seropositive parasites were observed by immunofluorescence microscopy (Nikon Eclipse Ti-U).

Brain superinfection assays and cyst enumeration

Brains from chronically infected mice (CEP *hxgprt-*) that survived secondary challenge were dissected, rinsed in PBS, passed through a 21G needle several times, pelleted and suspended in 1mL of PBS. For rederivation, 100μL of the brain homogenate was used to inoculate HFF monolayers in Toxo medium. One to two weeks later, infected HFFs were syringe-lysed and plated on new HFF monolayers to encourage parasite growth. Once HFFs were fully vacuolated, parasites were passaged in Toxo medium supplemented with mycophenolic acid (MPA) and xanthine that selects for parasites encoding a functional *HXGPRT* (i.e. the challenging strains) and against the chronically infecting type III *hxgprt-* which lacks a functional *HXGPRT* gene. Outgrowth in MPA-xanthine was considered evidence for superinfection.

For counting tissue cysts, 100μl of brain homogenate was fixed in 900μL of ice cold methanol, incubated for 5 minutes in microtubes (MCT-175-C, Axygen), washed and stained overnight in a 500μL PBS solution containing 1:150 dilution of FITC-conjugated *Dolichos biflorus* agglutinin (Vector Laboratories) with slow rotation at 4°C. The stained homogenate was further washed and suspended in 1mL of PBS, of which several 50μL aliquots were counted by fluorescence microscopy, and the number of cysts per brain were deduced.

Genetic linkage analysis

Quantitative trait loci (QTL) analysis was performed with the package r/QTL in R (version 3.6.1). LOD scores for each marker were calculated using the Haley-Knott regression model with the function 'scanone', or for all possible combination of two markers (i.e. epistatic interactions) using the function 'scantwo'. 1000 permutations were performed to obtain the genome wide LOD threshold for a P value of ≤0.05, which was considered statistically significant. Similar results were obtained with a linear mixed regression model. To estimate the effect each QTL had on the overall phenotype, the function 'fitqtl' was first used to fit the data to a multiple-QTL model. Statistical support was found for inclusion of all four QTLs with LOD scores > 3 compared to any lesser combination of three-QTLs (ANOVA P <0.02). Individual QTL effects

were then calculated under the assumption of the four-QTL model, which collectively accounts

for 91% of the observed phenotypic variance.

Cell isolation, in *vitro* recall infections, and FACS analysis

PECs were isolated by peritoneal lavage and splenocytes obtained, as described in [38]. In brief,

4mL of FACS buffer (PBS with 1% FBS) and 3mL of air were injected into the peritoneal cavity

with a 27G needle. After agitation, the PEC wash was poured into a conical tube. PEC washes

were filtered through a 70µm cell strainer, pelleted, and washed with FACS buffer before

staining. Spleens were dissected and crushed through 70µm cell strainers, pelleted, incubated

in ACK red blood cell RBC lysis buffer (0.15M NH_4Cl, 10mM $KHCO_3$, 0.1mM EDTA) for 5

minutes at room temperature, then washed with FACS buffer. To obtain peripheral blood

leukocytes (PBLs), 50µL of blood was isolated from mice in tubes containing 5µL of 0.5 M

EDTA on ice, pelleted and incubated in ACK lysis buffer, washed and peripheral blood

leukocytes (PBLs) were suspended in FACS buffer.

For FACS analysis, all preparations were done on ice, and cells were blocked in FACS

buffer containing Fc Block anti-CD16/32 (2.4G2) (BD Biosciences), 5% normal hamster serum,

and 5% normal rat serum (Jackson ImmunoResearch) for 20 minutes prior to staining with

fluorophore-conjugated monoclonal antibodies (mAbs). The following mAbs (1:100 staining

dilutions) were used: anti-CD11c-eFlour 450 (N418, eBioscience); anti-CD45.2-eFlour 450 (104,

eBioscience); anti-CD4-eFlour 450 (GK1.5, eBioscience), anti-CD4-PECy7 (GK1.5,

eBioscience); anti-CD11b-FITC (M1/70, eBioScience), anti-CD11b-BUV395 (M1/70, BD

Bioscience), anti-CD11b-Pacific Blue (M1/70, BioLegend); anti-IFNγ-PE (XMG1.2, BD

Bioscience); anti-CD8α-APC (53-6.7, eBioscience), anti-CD8α-BV510 (53-6.7, BioLegend), anti-

Ly6G-APC (1A8-Ly6g, eBioscience), anti-CD19-PerCP-Cy5.5 (ebio1D3, eBioscience), anti-

CD19-PE (6D5, BioLegend), anti-CD19-BV785 (6D5, BioLegend), anti-CD3-eFlour 780 (17A2,

BD Biosciences), anti-Ly6C-PECy7 (HK1.4, BioLegend), anti-CD23-Pacific Blue (B3B4,

BioLegend), anti-CD21/CD35-FITC (7E9, BioLegend), anti-CD5-APC (53-7.3, BioLegend), anti-

CD5-PerCP-Cy5.5 (53-7.3, BioLegend), anti-CD45R/B220-Cy7-APC (RA3-6B2, BioLegend),

anti-IgM-PECy7 (RMM-1, BioLegend), anti-CD138/Syndecan-1-BV650 (281-2, BioLegend), anti-

mouse-CD267/TACI-AlexaFlour-647 (8F10, BD Biosciences), and anti-CD268/BAFF-R-PE

(7H22-E16, BioLegend). Other FACS reagents included the viability dye propidium iodide

(Sigma) at a final concentration of 1µg/mL.

For *in vitro* recall, splenocytes and PerC were isolated from chronic and challenged mice

(day 5 or 7 following secondary infection) and 6 x 10^5 cells per well (96-well plate) were plated in

T cell medium (RPMI 1640 with GlutaMAX, 20% FBS, 1% Pen/Strep, 1mM NaPyruvate, 10mM

HEPES, 1.75µl BME). Cells were infected with a type I strain (RH or GT1) strain at an MOI

(multiplicity of infection) of 0.2 for 18 hr; 3µg/mL brefeldin A (eBioscience) was added for the last

5 hr of infection. 96-well plates were placed on ice, cells were harvested by pipetting and

washed with FACS buffer, blocked, and stained for surface markers. Cells were fixed with BD

Cytofix/Cytoperm and permeabilized with BD Perm/Wash solution (cat# 554714, BD

Pharmingen), stained with anti-IFNγ-PE (XMG1.2, BD Bioscience), anti-GZB (GB11,

BioLegend) and anti-IL-2-APC (JES6-5H4, BioLegend) on ice for 1 hr or overnight. Cells were

then washed once with BD Perm/Wash solution, once in FACS buffer, and analyzed by FACS.

For FoxP3 staining, peritoneal lavage was performed on chronic and challenged (day 7

following secondary infection) bumble and WT mice. Cells were washed and surface stained.

Fixation and permeabilization was performed with the eBioscience Foxp3/Transcription Factor

Staining Buffer Set (cat # 00-5523-00) before intracellular staining with anti-Foxp3-PE (MF-14,

BioLegend) according to the manufacturer's recommendations. Flow cytometry was performed

on a Beckman Coulter Cytoflex LX or LSR II (BD Biosciences) and analyzed with FlowJo^{TM}

software.

Analysis of isotype-specific antibody reactivity to *Toxoplasma gondii* by FACS

For serum reactivity analysis, syringe-lysed GFP-expressing strains (RH1-1 and GT1-GFP)

were fixed in 3% formaldehyde for 20 minutes, washed twice in PBS, and plated in 96 well

micro-titer plates at 4×10^5 parasites/well. The parasites were then incubated with serum from

chronically infected mice, at serum concentrations ranging from 10^{-2} to 10^{-6} diluted in FACS

buffer, for 20 minutes at 37°C. Parasites were then washed with FACS buffer and placed on ice

for incubation with anti-isotype detection antibodies depending on application: anti-IgG3-BV421

(R40-82, BD Bioscience), anti-IgM-PE/Cy7 (RMM-1, BioLegend), anti-IgG1-FITC (RMG1-1,

BioLegend), anti-IgG1-APC (RMG1-1, BioLegend), anti-IgG2b-FITC (RMG2b-1, BioLegend),

anti-IgG2b-PE (RMG2b-1, BioLegend), anti-IgG2a-FITC (RMG2a-62, BioLegend), anti-IgG2a-

PerCP/Cy5.5 (RMG2a-62, BioLegend), anti-IgM-a-PE (DS-1, BD Bioscience), Anti-IgM-b-PE

(AF6-78, BD Bioscience).

Parasite neutralization assay

Heat-inactivated serum was used to coat live parasites for 20 minutes at 37°C before infecting

5×10^5 mouse embryonic fibroblasts/well (MEFs) in 96 well plates. Immediately following addition

of parasite to MEF wells, plates were spun at 1200rpm for 3 minutes to synchronize infection. 2

hrs after initiation of infection, cells were placed on ice and harvested by scraping with pipette

tips. Cells were washed twice in FACS buffer, suspended in 1:1000 PI in FACS buffer, and then

analyzed by flow cytometry.

SDS-PAGE and immunoblotting for parasite lysate antigen

To generate parasite lysate antigens *Toxoplasma gondii* was cultured in HFF and expanded to

approximately 2×10^8 parasites. Parasites were syringe-lysed, washed with sterile 1X PBS and

the parasite pellet was lysed with (1mL) 0.1% TritonX-100 detergent in 1X PBS. Solubilized

parasites were centrifuged at 2,000 RCF for 20 minutes to remove large debris. The

supernatant was aliquoted and stored at -80°C. Parasite lysate was reduced with β-

mercaptoethanol (BME) and separated via SDS-PAGE in 4-20% Mini-PROTEAN TGX pre-cast

gels (cat # 4561096, Bio-Rad) before transfer to PVDF membrane using a Trans-Blot Turbo

Mini PVDF Transfer Pack (cat # 1704156, Bio-Rad) via Bio-Rad Transblot Turbo (cat #

1704150, Bio-Rad). Membranes were blocked with 10% fortified bovine milk dissolved in Tris-

Buffered Saline with 0.1% Tween (TBS-T 0.1%) for 1-2 hrs at room temperature or overnight at

4°C. Blots were then probed with heat-inactivated serum in block at either 1:1,000 dilution for

serum IgM analysis or 1:5,000 dilution for serum IgG analysis overnight at 4°C. Membranes

were washed with TBS-T 0.1% three times for 20 minutes per wash. Blots were then incubated

for one hr at room temperature with goat α-mouse horseradish peroxidase (HRP)-conjugated

antibodies (SouthernBiotec): anti-IgM secondary 1:1000 (cat# 1020-05) and total anti-IgG

secondary 1:5000 (cat# 1030-05). Membranes were then washed with TBS-T 0.1% three times

and developed with Immobilon® Forte Western HRP Substrate (WBLUF0500). All blots were

imaged via chemiluminescence on a ChemiDoc Touch (cat# 12003153, Bio-Rad). Image Lab

6.1 software (Bio-Rad) was used for analysis of bands and total lane signal. Western blots

comparing A/J to C57BL/6J were developed simultaneously and the band signal was

normalized to A/J.

RNA isolation and sequencing

Peritoneal B-1a (B220$^{int-neg}$ CD19^{high} CD11b+ CD5+ PI-), B-1b (B220$^{int-neg}$ CD19^{high} CD11b+

CD5- PI-), or B-2 B cells (B220^{high} CD19+ CD11b- PI-) were sorted into 500ul RNeasy lysis

buffer using a FACS ARIA II cell sorter (BD Biosciences). RNA was purified using the RNeasy

mini kit (cat# 74134, Qiagen) according to the manufacturer's protocol. RNA purity was tested

by Qubit (ThermoFisher) and Agilent 2100 BioAnalyzer for total RNA with picogram sensitivity.

DNA libraries were generated with a Lexogen QuantSeq-UMI FWD 3′ mRNA-Seq Library Prep

Kit (cat# 015). Samples were sent to UC Davis for QuantSeq 3' mRNA FWD-UMI sequencing.

Gene expression analysis and data availability

Raw reads were trimmed and mapped by the BlueBee genomic pipeline FWD-UMI Mouse

(GRCm38) Lexogen Quantseq 2.6.1 (Lexogen). In brief, reads were quality controlled with

'FASTQC', trimmed with 'Bbduk' to remove low quality tails, poly(A)read-through and adapter

contaminations, read alignments to the *Mus musculus* genome build GRCm38 were done with

'STAR', and gene reads were quantified with 'HTSeq-count'. Differentially expressed gene

(DEG) analysis was performed utilizing limma-voom in R version 3.6.3 in RStudio with

Bioconductor suite of packages. Heatmaps were generated with 'gplots'. Pathway and GO term

analysis was performed with MouseMine (mousemine.org), and gene set enrichment analysis

was performed with GSEA v4.0.3. RNA-sequencing data generated in this study has been

deposited in the NCBI Sequence Read Archive (Bioproject accession number PRJNA637442).

All other data that support the findings herein are available from the corresponding author upon

reasonable request.

LPS-stimulation of enriched B cells and quantitative PCR

PerC and splenocytes were isolated from naïve 6-8 week-old C57BL/6J and A/J mice and

enriched for B cells using the EasySep Mouse Pan B cell Isolation Kit (cat#19844, StemTech).

Bead enrichment for splenic samples had the addition of biotinylated anti-CD43 antibodies

(clone S7, BD Bioscience) to remove B-1 cells. Enriched samples were plated in a 96-well plate

at 400,000 cells per well and stimulated with 25ug/ml of LPS (cat# L4391-1MG, Sigma). After 2

hrs, total RNA was isolated using the Rneasy Mini Kit (cat# 74134) and cDNA was synthesized

using High Capacity cDNA Reverse Transcription Kit (ThermoFisher, cat#4368814).

Quantitative PCR was performed on synthesized cDNA samples using ThermoFisher TaqMan

Master Mix (cat# 4444556) and TaqMan probes: *Actb* - Assay ID:Mm02619580_g1 (cat#

4331182), and *Nfkbid* - Assay ID: Mm00549082_m1 (cat# 4331182), according to the

manufacturer's protocol. Normalization of *Nfkbid* expression in each sample was calculated in

comparison to *Actb* expression levels. Fold change in *Nfkbid* expression of AJ relative to that of

C57BL/6J cells was determined through the delta delta CT method ($2^{-\Delta\Delta CT}$).

PerC adoptive transfers and IgH allotype chimeric mouse generation

PerC was harvested by peritoneal lavage of 6-12 week old C57BL/6J donor mice as described

above and 5×10^6 total peritoneal exudate cells (total PerC)/60ul PBS dose were transferred i.p.

into 2-4 day old bumble neonates. Allotype chimeric mouse generation was performed as

previously described [63]. In brief, 1-day old C57BL/6J neonates were treated with 0.1mg of anti-

IgM-b (clone AF6-78) and twice weekly thereafter treated with 0.2mg of anti-IgM-b for 6 weeks.

On day 2 after birth the neonates were given 5×10^6 total PerC from B6.Cg-*Gpi1^a Thy1^a Igh^a*

delivered i.p. The mice were then allowed to rest for 6 weeks after the last antibody treatment

before infection with *T. gondii.*

Bone marrow chimeric mice generation

CD45.1 B6.SJL-*Ptprc^a Pepc^b* /BoyJ recipient mice were given 2 doses of 500cGy with an X-

Rad320 (Precision X-Ray) with a 4 hr interval. Donor BM cells were harvested from muMT,

bumble and WT C57BL/6J mice, filtered with a 70um filter, incubated in ACK red blood cell lysis

buffer, washed with PBS and transplanted by retro-orbital injection of 1:1 (muMT:bumble, or

muMT:WT) mixed BM cells at a concentration of 10^7 cells/200ul PBS dose. Recipient mice were

then allowed to rest for 8 weeks. Reconstitution was assessed at 8 weeks by FACS analysis of

PBL.

ELISA

High-affinity protein binding microplates (Corning) were coated overnight goat anti-mouse IgM

(1mg/ml, Southern Biotech, cat# 1021-01) and blocked with coating buffer containing with 1%

BSA (w/v) and 2% goat serum (Omega Scientific) in PBS. Wells were washed with ELISA wash

buffer (1X PBS, 0.05% TweenTM-20). Mouse serum samples were diluted 1:400 in coating

buffer and incubated in the wells for 1 hr. Wells were washed 5 times and secondary goat anti-

IgM-HRP (Southern Biotech, cat# 1021-08) at 1:5000 dilution was incubated for 1 hr in the

wells. Wells were washed several times and developed with TMB substrate (Invitrogen).

Development was stopped after 20 minutes with 1M H_3PO_4 stop solution. Absorbance was

measured at 450nm on a BioTek Epoch microplate spectrophotometer.

Statistics

Statistical analysis was performed with Graphpad Prism 8 software. Statistical significance was

defined as $P < 0.05$. P values were calculated using paired or unpaired two-tailed t-tests, 2-way

ANOVA with Tukey multiple comparison correction, and multiple t-tests with the Holm-Sidak

correction for multiple comparisons. Survival curve significance was calculated using log-rank

(Mantel-Cox) testing. Significance in time to death was calculated using the Gehan-Breslow-

Wilcoxon test. Differential gene expression analysis statistics were calculated using the Limma-

Voom R package, P values were adjusted with the Benjamini-Hochberg correction for multiple

comparisons. GO term and pathway enrichment analysis statistics were performed at

mousemine.org using the Holm-Bonferroni test correction. Statistical methods used for each

figure are indicated in the figure legends.

FIGURE LEGENDS

Figure 1. MHC and non-MHC alleles promote immunity to virulent *Toxoplasma gondii*

strains in A/J mice.

All mice were primed with 10^4 type III avirulent *T. gondii* CEP *hxgprt*- parasites; then, 35 days

later, mice were challenged with $5x10^4$ of the indicated strain of *T. gondii* parasites. A) Survival

of A/J [VAND (n=7), GUYDOS (n=5), GPHT (n=5), FOU (n=5), TgCATBr5 (n=5), GUYMAT

(n=4), and MAS (n=7)],   B10 [VAND (n=4), GUYDOS (n=4), GPHT (n=4), FOU (n=4),

TgCATBr5 (n=4), GUYMAT (n=4), and MAS (n=12)], and B10.A [VAND (n=4), GUYDOS (n=4),

GPHT (n=4), FOU (n=4), TgCATBr5 (n=4), GUYMAT (n=5), and MAS (n=10)] mice following

secondary infection with virulent atypical *T. gondii* strains. Cumulative results are plotted from 1-

2 experiments. B) Bioluminescence imaging of individual mice challenged with luciferase

expressing MAS strain on days 1, 4 and 8 of secondary infection; parasite burden is shown as a

heat map depicting the relative number of photons (photons/sec/cm2/sr) detected over a 5

minute exposure. C) Survival of B10 (n=5), B10.A (n=5), A/J (n=12), and F1 (A/J x C57BL/6,

n=9) mice following secondary infection with the virulent *T. gondii* strain GT1; ***P<0.0001, Log-

rank (Mantel-Cox) compared to A/J mice. Cumulative data from 1 to 2 experiments are plotted.

D) Superinfection in surviving A/J (n=12) and F1 (A/J x C57BL/6, n=9) following 35 days of

secondary infection with the *T. gondii* type I GT1 strain. To evaluate superinfection, brains of A/J

(n=12) and F1 (A/J x C57BL/6, n=9) mice were taken from secondary challenge survivors,

emulsified, and grown in MPA-xanthine selection medium, which selects for parasites GT1

parasites expressing the endogenous *HXGPRT* locus against the *hxgprt-* type III CEP strain

used for chronic infection. The fraction of mice for which the presence of the secondary parasite

strain detected (i.e. brain co-infection) or not is plotted; **** P< 0.0001, Fisher's exact test.

Figure 2. Four genetic loci correlate with host survival to *Toxoplasma gondii* secondary

infections.

26 recombinant inbred (RI) mouse strains from the AxB;BxA panel were primed with $1x10^4$ type

III avirulent *T. gondii* CEP *hxgprt-* parasites; then, 35 days later, mice were challenged with

$5x10^4$ virulent type I GT1 *T. gondii* parasites (n=2 per RI line). A) LOD scores for each marker

were calculated using Haley-Knott regression and the running LOD scores of primary (black)

and secondary infection survival (blue) for each genetic marker is plotted. 1000 permutations

were performed to obtain the genome wide threshold LOD values; P=0.10 and 0.05 thresholds

are shown. QTL analysis was performed with the program r/QTL (Bowman). B) Effect plots for

the genetic markers closest to the maximal LOD scores calculated for the chromosome 7, 10,

11 and 17 QTLs are shown. Each dot indicates the percent survival of a unique RI line and

whether it encodes an A/J (red) or C57BL/6 (blue) allele at the specified genetic marker.

Figure 3. *Nfkbid* is required for immunity to secondary infection with *Toxoplasma gondii*.

A) Cumulative survival of bumble (*Nfkbid-/-* C57BL/6) (n=71) and wildtype (C57BL/6J) (n=89)

naïve mice infected with the avirulent type III CEP strain. B) Survival of chronically infected

bumble (n=12) and wildtype (n=8) mice given a secondary infection with the type I GT1 strain.

Cumulative survival from 3 separate experiments are shown; * P<0.02, Gehan-Breslow-

Wilcoxon test. C) As in B, but survival to secondary infection with the type I RH strain is shown.

Cumulative survival from 3 separate experiments is plotted (bumble n=17, C57BL/6J n=4);

***P<0.008, Mantel-Cox test. D) Frequency of GFP+ events in the peritoneal lavage 7 days

post-secondary infection with GFP-expressing type 1 RH strain (RH 1-1). Each dot represents

the result of one mouse, and cumulative results are shown from 3 separate experiments

(bumble n=13, C57BL/6J n=17); **P<0.002, unpaired two-tailed t-test.

Figure 4. *Nfkbid* is required for the generation of parasite-specific neutralizing

antibodies.

A) Schematic depicting the assay used to detect serum antibody reactivity to *T. gondii*. Whole

fixed GFP+ parasites are incubated with serum from infected mice which are then stained with

fluorescently labeled anti-isotype detection antibodies and assessed by FACS. B)

Representative histograms display the detection of parasite-specific IgM, IgG3 and IgG1 bound

to formaldehyde fixed GFP+ type I RH parasites; diluted serums (10^3) from naïve C57BL/6J

mice, or from type III CEP chronically infected C57BL/6J or bumble mice were assayed. C) As

in B, but quantification of the parasite-specific antibody isotype binding (IgM, IgG1, IgG2a/c,

IgG2b, and IgG3) over a range of serum concentrations is shown. Background staining in the

absence of serum is indicated for each isotype. Plotted is the cumulative average +/-SD of the

geometric mean fluorescence straining intensity (MFI) from 3 separate experiments (bumble

n=8, C57BL/6J n=11). D) Neutralization of GFP+ parasites coated with serum over a range of

concentrations from the indicated mice. Parasites were incubated in serum for 20 minutes

before infection of mouse embryonic fibroblasts and assessed by FACS 2h later. The fraction of

infected host cells (GFP+ cells) is normalized to that of parasite infections without serum. Each

dot represents the serum from an individual mouse and cumulative results from 3 separate

experiments are shown (bumble n=8, C57BL/6J n=7, naïve n=6). For C and D, significance was

assessed by unpaired t-tests with Holm-Sidak correction for multiple comparisons; *** P< 0.001,

** P<0.01.

Figure 5. Evidence for enhanced B-1 cell activation in resistant A/J mice.

A) Gating strategies for identifying B-1 B cells. B-1 B cells are identified as CD19+, B220^{int} and

CD11b+. B-1a B cells are CD5+ and B-1b B cells are CD5-. B) Representative FACS plots of

the peritoneal B-1 B cell compartment (CD19+ B220$^{int-neg}$ CD11b^{+}) in A/J and C57BL/6J (B6)

mice at naïve, chronic, and 5 days (D5) post-secondary infection with the GT1 strain. Numbers

indicate the percent of cells that fall within the indicated gate. Representative histograms of IgM

surface expression and percent of cells that fall within IgM^{lo} gate of B-1a and B-1b cells from A/J

(red) and C57BL/6J (blue) at the indicated time points. Boxed in red highlight the increased

frequencies of B-1b B cells that are IgM^{low} on day 5 post-secondary infection with *T. gondii*

observed in A/J relative to C57BL/6J mice. C) Representative FACS plots of splenic CD19+

B220$^{int-neg}$ B cells stained for CD43 and CD5 in A/J and C57BL/6J mice at day 5 of secondary

infection. The relative percentage of B-1a and B-1b among total CD43+ CD19+ B220$^{int-neg}$ B-1

cells is indicated. D) The frequency of B-1a (CD5+) or B-1b (CD5-) among CD43+ CD19+

B220$^{int-neg}$ B-1 cells from the spleen and peritoneum at the indicated infection states were

analyzed. Each dot represents the results from an individual mouse and the cumulative

averages +SD from 2-3 experiments are plotted. Significance was assessed with an unpaired

two-tailed t-test; *** P<0.001, ** P<0.01. E) Frequencies of peritoneal and splenic B-1a or B-1b

(CD19+ B220$^{int-neg}$ CD43+) that are IgM+IgD- or IgM-IgD- in A/J and C57BL/6J mice at the

indicated infection states. F) Frequencies of splenic B-1a or B-1b cells from A/J and C57BL/6J

mice that express BAFFR+, TACI+, or CD138+ at the indicated infection states. For E and F,

the cumulative average +SD from 2-4 experiments are plotted and each dot represents the

result from an individual mouse; P values calculated by 2-way ANOVA with Tukey correction;

**** P<0.0001, *** P<0.001, ** P<0.01, * P<0.05. The legend applies to panels D-F.

Figure 6. B-1 cells have enhanced germline transcription of *Ighg* constant regions and

different activation profiles in resistant A/J mice.

A) Transcriptomic analysis of peritoneal B-1a (CD19+ B220$^{int-neg}$ CD11b+ CD5+), B-1b (CD19+

B220$^{int-neg}$ CD11b+ CD5-) and B-2 (CD19+ B220^{hi} CD11b+ CD5+) B cells from A/J and

C57BL/6J mice, that were either naïve, chronically infected, or on day 5 of secondary infection

with the type I GT1 strain, was performed using 3'-Tag RNA sequencing. The heat map depicts

the relative expression of genes that were: i) at least fourfold differentially expressed in any of

the B cell compartments when comparing its expression between the naïve and either chronic

or secondary infection states; and ii) it made the cutoff in at least three separate comparisons

irrespective of genetic background. B) Pathway and GO term enrichment of the genes depicted

in A; P values were calculated with Holm-Bonferroni adjustments. C) Heatmap depicting the

relative expression of all *Ighg* transcripts from the indicated B cell population, mouse strain and

infection state. D) Gene set enrichment analysis of the rank-ordered list of differentially

expressed genes between A/J and C57BL/6J B-1b cells at day 5 of secondary infection. Gene

sets depicted (GSE28237, GSE25677 and GSE23502) were in the top 10 gene sets ranked by

false discovery rate (FDR) after investigating MSigDB's C7: immunologic signatures collection.

Enrichment score is the degree of overrepresentation of a gene set at the top or bottom of a

ranked list. NES is the enrichment score after normalizing for gene set size.

Figure 7. Gene dosage of *Nfkbid* impacts parasite-specific IgG1 responses.

A) *Nfkbid* expression in CPM (Counts per million) of 3'-Tag RNA-seq reads of the indicated B

cell populations obtained from A/J and C57BL/6J mice that were either naïve, chronically

infected, or on day 5 of secondary infection with the type I GT1 strain. B) Enriched B cells from

the PerC and spleen were stimulated with LPS for 2 hrs and *Nfkbid* transcripts were quantified

by qPCR; ** P< 0.01, paired t-test. C) Representative histograms display the detection of

parasite-specific IgG1 bound to formaldehyde fixed GFP+ type I GT1 parasites; diluted serums

(10^3) from C57BL/6J, *Nfkbid*+/- (C57BL/6J x bumble F1), and A/J mice chronically infected with

the type III CEP strain were assayed. Anti-mouse IgG1 background staining in the absence of

serum is shown. D) As in C, but quantification of the parasite-specific antibody isotype binding

(IgM, IgG1, IgG2a/c, IgG2b, and IgG3) over a range of serum concentrations is shown.

Background staining in the absence of serum is indicated for each isotype. Plotted is the

cumulative average +/-SD of the geometric mean fluorescence straining intensity (MFI) from 2-3

separate experiments (C57BL/6J n=8; *Nfkbid*+/- n=7; A/J n=8); significance was assessed by

unpaired t-tests and Holm-Sidak corrections comparing A/J vs C57BL/6J (*) or *Nfkbid*+/- vs

C57BL/6J (#); **** P< 0.0001, *** P< 0.001, *[/#] P<0.05. IgG1 staining was not significantly

different between A/J and *Nfkbid*+/- serums.

Figure 8. Contribution of B-1 cells to *Toxoplasma gondii* immunity.

A) Schematic of the secondary infection experiment using PerC reconstituted bumble mice. 2-4

day old bumble neonates were transferred $5x10^6$ total PerC and allowed to rest for 7 weeks

before primary infection with type III CEP strain. 5 weeks post-primary infection, mice were

given a secondary infection with the type I strain RH. Survival of bumble mice given total PerC

transfers (n=10) relative to littermate controls (n=12). Cumulative survival is shown from 3

separate experiments; *** P <0.001, Mantel-Cox test. B) Representative histogram of anti-

isotype (IgM, IgG1, IgG3, IgG2a/c, and IgG2b) MFI bound to parasites coated in serum from

chronically infected bumble, bumble given PerC transfer, and WT mice. C) Schematic of

neonatal allotype chimera generation. C57BL/6J neonates were given anti-IgHb to deplete

endogenous B cells at day 1 post birth and twice weekly after for 6 weeks, thereby depleting the

endogenous B-1 pool for the life of the animal due to their restricted fetal/neonatal window of

development. Neonates were given $5x10^6$ total PerC from 6-8 week old IgH-a congenic

C57BL/6 mice donors. These mice then rested for 6 weeks after the last depletion treatment to

allow reemergence of the endogenous B-2 IgH-b cells. D) Representative histograms of serum

derived anti-IgM-a (B-1 derived) or anti-IgM-b (B-2 derived) staining profiles of type I RH GFP+

parasites taken from the IgH allotype chimeras on day 14 and 30 following primary type III CEP

infection. Staining controls with serum from chronically infected C57BL/6J IgH-b littermates, or

IgH-a mice are shown.

Table S1. Virulent strains of *Toxoplasma gondii* superinfect genetically resistant hosts.

CEP *hxgprt-* chronically infected C57BL/10, C57BL/10.A, and A/J mice were given a secondary

infection with the indicated atypical *T. gondii* strain; then, 35-45 days after secondary infection,

brains from surviving mice were homogenized in PBS and used to inoculate HFF monolayers.

The resultant parasite cultures were then tested for the presence of the secondary infection

strain; numerator = number of mice tested positive for secondary infection strain in the brain;

denominator = number of mice that exhibited parasite growth in the HFF monolayer prior to drug

selection.

[1]Parasite growth was observed in mycophenolic acid / xanthine medium, which selects for

parasites encoding a functional *HXGPRT* gene (i.e. the challenging atypical strains) and kills the

primary infection CEP *hxgprt-* strain.

Table S2. Primary and secondary infection survival of recombinant inbred mice

(AxB;BxA)

26 strains (n = 2) from the RI (AxB; BxA) panel were primed with $1x10^4$ type III avirulent CEP

hxgprt- T. gondii parasites; then, 35 days later, mice were challenged with $5x10^4$ virulent type I

GT1 *T. gondii* parasites. Primary and secondary infection survival percentages are indicated.

35-45 days after secondary challenge, brains from the surviving RI mice were homogenized in

PBS and used to inoculate HFF monolayers. The resultant parasite cultures were then tested

for the presence of the challenging strain; numerator = number of mice tested positive for

secondary infection strain in the brain; denominator = number of mice that exhibited parasite

growth in the HFF monolayer prior to drug selection. Some surviving mice failed to generate

parasite positive cultures.

803 [1]Parasite growth was observed in mycophenolic acid / xanthine medium, which selects for GT1

804 parasites (the challenging strain) encoding the endogenous *HXGPRT* gene and against the

805 primary infection CEP *hxgprt-* strain.

807 **Table S3. Polymorphic genes between A/J and C57BL/6J encoded within the four QTLs**

808 **that define immunity to *Toxoplasma gondii*.**

809 For each of the four QTLs that define secondary infection immunity to *T. gondii*, a list of all

810 genes that have a DNA polymorphism between A/J and C5BL/6J mice, and are encoded within

811 the QTL boundaries defined by the maximal genetic marker (real or imputed markers inferred in

812 r/QTL) and the two flanking markers on either side; LOD scores and position are indicated. For

813 each gene, its location and unique identifiers are listed, as well as the number and class of

814 small nucleotide polymorphism (SNP) between A/J and C57BL/6J are shown. Total number of

815 SNPs for each gene were also tallied ('Sum mutation'), and in bold are genes who's SNPs are

816 two standard deviations greater than the average SNP for that class within the QTL region. In

817 these QTL regions, there were also 476 (chr7), 88 (chr10), 40 (chr11) and 375 SNPs (chr17)

818 that did not associate with any gene in the QTL boundaries. UTR, untranslated region; 'Region

819 SNP', SNPs that are +/- 2 kb of the gene boundaries. Data obtained from Mouse Genome

820 Informatics (MGI) (informatics.jax.org), and each locus is in a separate tab.

822 **Table S4. Transcriptomic analysis of B cells responding to *Toxoplasma gondii* infection.**

823 Table of genes included (threshold of >=5 CPM in any sample) in differential gene expression

824 analysis. For each gene the Ensembl ID, MGI ID, gene symbol and descriptions are listed. The

825 CPM from 3 samples of peritoneal B-1a, B-1b, and B-2 at naïve (N), chronic (Ch), and day 5

826 (D5) of secondary infection in both A/J (AJ) and C57BL/6J (B6) mice are given. The average

827 CPM was calculated for each set of samples and average counts less than 5 CPM were raised

828 to 5 to reduce the effect of low expression genes in further analysis. The log2 fold change (FC)

of gene expression was calculated using the average CPM for each cell type (B-1a, B-1b, and

B-2) of each strain (A/J and C57BL/6J) as they progressed from naïve to chronic infection or

chronic to D5 secondary infection. The Log2 FC was also calculated from naïve to D5 of

secondary infection. The Log2 FC between A/J and C57BL/6J was calculated for each cell type.

Genes were then ranked by the number of comparisons that exhibited greater than 4FC by the

indicated comparison.

Figure S1. Cyst burden and weight during chronic infection with the low virulent type III

CEP strain correlates with the murine H-2 locus.

A/J, C57BL/6J (B6), C57BL/10J (B10), and C57BL/10.AJ (B10.A) mice were injected with the

type III strain *CEP hxgprt-* and allowed to progress to chronic infection. Plotted (+/- SEM) is the

average cyst number (x 1000) in the brain vs. the average fraction of initial weight, where 1 is

the normalized weight on the day of type III injection; the regression value (R^2) is indicated.

Results are from 3 to 5 mice for cyst numbers (day 42 of chronic infection), and 5-12 mice for

weight measurements (day 30 of chronic infection) per mouse strain.

Figure S2. Bumble mice have intact T cell responses during secondary infection.

A) Peritoneal CD4 and CD8 T cells from bumble and C57BL/6J mice were assessed at day 7 of

secondary infection with the type I RH strain by an in vitro recall assay and assayed for

intracellular IFNγ and IL-2 as described in Figure S3. B) Peritoneal T-regulatory cells (CD4+

CD25+ Foxp3+) were quantified at day 7 of secondary infection with type I RH strain. Each dot

represents the result from one mouse, and plotted are cumulative averages +SD from 3

experiments; no significant differences were observed between bumble and C56BL/6 mice by

unpaired t-tests with the Holm-Sidak correction for multiple comparisons.

Figure S3. Enhanced myeloid cell frequencies in C57BL/6J mice following chronic and

secondary infections with *Toxoplasma gondii*.

A) Representative FACs plots of PerC characterizing immune cell populations within the

peritoneal compartment. Populations characterized include: B-2 B cells (CD19+, CD11b-), B-1 B

cells (CD19+ CD11b+), neutrophils (CD19- Ly6G+), T cells (CD19- CD3+), macrophages

(CD19- Ly6G- CD3- CD11b+), and dendritic cells (CD19- Ly6G- CD3- CD11c+ CD11b+/-

CD8+/-). B) Immune cell populations are plotted as a frequency of total leukocytes. PerC was

harvested from A/J and C57BL/6J mice chronically infected with type III CEP *T gondii* strain or 5

days post-secondary infection with type I GT1 strain; **** P<0.0001, *** P<0.001, * P<0.05,

unpaired two-tailed t-test.

Figure S4. CD8 T cell IFNγ frequencies are increased in resistant A/J mice.

A) Peritoneal and B) splenic cells were harvested from A/J and C57BL6/J mice chronically

infected with type III CEP *T. gondii* strain, and infected with live type I parasites for 16 hrs. T

cells were assessed for production of granzyme B (GZB), IFNγ, and IL-2 by intracellular staining

and FACS. The average frequency +/-SD of positive staining CD4+ or CD8+ T cells (CD3+

CD19-) and cumulative results from 2-3 experiments (AJ n=8, C57BL6/J n=11) are shown; *

P<0.05, unpaired two-tailed t-test.

Figure S5. Immunity in A/J mice correlates with heightened serum antibody reactivity to

***Toxoplasma gondii* lysate antigen.**

Western blot analysis of serum-derived antibody reactivity against type I GT1 parasite lysate

antigen. A) Blood plasma was harvested from A/J and C57BL/6J mice either chronically infected

with type III CEP *T. gondii* strain, or B) 5 days post-secondary infection with type I GT1 and

used to probe GT1 parasite lysate separated by SDS-PAGE then detected with anti-IgM-HRP or

anti-IgG-HRP. Western blots comparing A/J and C57BL/6J serum reactivity to GT1 lysate were

developed and imaged simultaneously to reduce variability in band intensity. Total lane signal is

the sum of the colorimetric intensity from lysate bands in a single lane quantified in Image Lab

(Bio-Rad). "Total Lane Signal" is normalized lane signal from C57BL/6J compared to A/J which

were developed in tandem. Results are from 6 individual experiments; ** P < 0.01, * P < 0.05;

unpaired two-tailed t-tests.

Figure S6. Genes uniquely induced in B-1b cells on day 5 of secondary infection in

resistant A/J mice.

A) Genes that were differentially upregulated in B-1b cells on day 5 of secondary infection

compared to naïve mice in the A/J genetic background. P values of differentially expressed

genes were calculated using the Benjamini-Hochberg adjustment for false discovery rate, and

only those genes that survived significance were included in the heatmap. For comparison, all

B-1 compartments in A/J mice are shown for this gene set. A/J B-1 Pathway and GO term

enrichment was assessed on the genes presented in the heatmap in A. P values for enrichment

analysis were adjusted with the Holm-Bonferroni correction. B) A cluster of genes found to be

differentially induced in B-1b cells in A/J compared to C57BL/6J mice on day 5 of secondary

infection are plotted as a heat map.

Figure S7. Assessing PerC reconstitutions by serum IgM ELISAs and flow cytometry.

A) Serum IgM from C57BL/6J, muMT, bumble, and A/J mice was measured by ELISA. Serum

was harvested from mice either naïve, chronically infected with type III CEP strain, or on day 5

post-secondary infection with type I GT1 *T. gondii* strain. PerC transfer (+) refers to mice

adoptively transferred 5×10^6 total PerC cells as a day 2 neonate. Each dot represents the

results from an individual mouse, and plotted is the average +/-SD of the O.D. obtained at

450nm; *P<0.05, unpaired two-tailed t-test. B) Bumble reconstitution of the peritoneal B-1

compartment after neonatal PerC adoptive transfer. Representative FACS plots of peritoneal B-

2 cells (B220^{high} CD19+) and B-1 (B220$^{int-neg}$ CD19+) cells from bumble mice and bumble mice

with PerC adoptive transfer. Shown are mice on day 20 of primary infection with the type III CEP

strain.

Figure S8. B cell deficient muMT mice are highly susceptible to primary infections

irrespective of B cell reconstitution.

A) B cell deficient muMT mice (n=3), muMT given WT PerC adoptive transfers as 2-day

neonates then allowed to reconstitute for 6-7 weeks into adulthood (n=2), and muMT given B

cell enriched splenocytes (n=3) 1 day prior to infection with the type III CEP strain were

assessed for survival. B) muMT reconstitution of peritoneal B cell compartment after neonatal

adoptive transfer. Representative FACS plots of peritoneal B-2 cells (B220^{high} CD19+) and B-1

B cells (B220$^{int-neg}$ CD19+) from WT, and muMT mice or muMT mice with neonatal PerC

adoptive transfer. C) As in B, but WT, and muMT with B cell enriched splenocytes (EasySep™

Mouse Pan-B Cell Isolation Kit, cat# 19844) adoptively transferred 24 hrs earlier. For B and C,

uninfected mice are 6-8 weeks of age and numbers indicate the percent of cells that fall within

the depicted gate.

Figure S9. Bone marrow chimeric mice fail to survive primary infection.

A) Schematic of bone marrow (BM) chimera generation. 1:1 mixtures of B-cell deficient muMT

BM and either WT (n=10) or bumble BM (n=10) were adoptively transferred into lethally

irradiated CD45.1 mice. In this way, the B cell compartment of the reconstituted chimeric mice

will either express or lack *Nfkbid,* respectively. B) FACS analysis of the peripheral blood 8

weeks after transfer. Reconstitution is measured by the frequency of transferred CD45.2+ cells

in the immune cell population analyzed. Each dot represents the result from an individual

mouse. C) Survival of BM chimeras given type III CEP *T. gondii* infection. Results are

cumulative from 2 independent transfers; n.s., Mantel-Cox.

REFERENCES

1. Sacks, D. L. Vaccines against tropical parasitic diseases: a persisting answer to a persisting problem. *Nat. Immunol.* **15**, 403-405 (2014).

2. Lorenzi, H. *et al.* Local admixture of amplified and diversified secreted pathogenesis determinants shapes mosaic Toxoplasma gondii genomes. *Nat. Commun.* **7**, 10147 (2016).

3. Howe, D. K. & Sibley, L. D. Toxoplasma gondii comprises three clonal lineages: correlation of parasite genotype with human disease. *J. Infect. Dis.* **172**, 1561-1566 (1995).

4. Khan, A. *et al.* Genetic divergence of Toxoplasma gondii strains associated with ocular toxoplasmosis, Brazil. *Emerg. Infect. Dis.* **12**, 942-949 (2006).

5. McLeod, R. *et al.* Prematurity and severity are associated with Toxoplasma gondii alleles (NCCCTS, 1981-2009). *Clin. Infect. Dis.* **54**, 1595-1605 (2012).

6. Grigg, M. E., Ganatra, J., Boothroyd, J. C. & Margolis, T. P. Unusual Abundance of Atypical Strains Associated with Human Ocular Toxoplasmosis. *J. Infect. Dis.* **184**, 633 639 (2001).

7. Campos, M. A. S. *et al.* Ocular Sequelae of Congenital Toxoplasmosis in Brazil Compared with Europe. *PLoS Neglected Tropical Diseases* **2**, e277 (2008).

8. Suzuki, Y. & Remington, J. S. Dual regulation of resistance against Toxoplasma gondii infection by Lyt-2+ and Lyt-1+, L3T4+ T cells in mice. *J. Immunol.* **140**, 3943-3946 (1988).

9. Suzuki, Y. & Remington, J. S. The effect of anti-IFN-gamma antibody on the protective effect of Lyt-2+ immune T cells against toxoplasmosis in mice. *J. Immunol.* **144**, 1954-1956 (1990).

10. Gazzinelli, R. T., Hakim, F. T., Hieny, S., Shearer, G. M. & Sher, A. Synergistic role of CD4+ and CD8+ T lymphocytes in IFN-gamma production and protective immunity induced by an attenuated Toxoplasma gondii vaccine. *J. Immunol.* **146**, 286-292 (1991).

11. Gigley, J. P., Fox, B. A. & Bzik, D. J. Cell-mediated immunity to Toxoplasma gondii develops primarily by local Th1 host immune responses in the absence of parasite replication. *J. Immunol.* **182**, 1069-1078 (2009).

12. Jordan, K. A. *et al.* Kinetics and phenotype of vaccine-induced CD8+ T-cell responses to Toxoplasma gondii. *Infect. Immun.* **77**, 3894-3901 (2009).

13. Gigley, J. P., Bhadra, R. & Khan, I. A. CD8 T Cells and Toxoplasma gondii: A New Paradigm. *J. Parasitol. Res.* **2011**, 243796 (2011).

14. Khan, I. A., Hwang, S. & Moretto, M. Toxoplasma gondii: CD8 T Cells Cry for CD4 Help. *Front. Cell. Infect. Microbiol.* **9**, 136 (2019).

15. Casciotti, L., Ely, K. H., Williams, M. E. & Khan, I. A. CD8(+)-T-cell immunity against
Toxoplasma gondii can be induced but not maintained in mice lacking conventional CD4(+) T
cells. *Infect. Immun.* **70**, 434-443 (2002).

16. Johnson, L. L. & Sayles, P. C. Deficient humoral responses underlie susceptibility to
Toxoplasma gondii in CD4-deficient mice. *Infect. Immun.* **70**, 185-191 (2002).

17. Chen, M. *et al*. Induction of protective immunity by primed B-1 cells in Toxoplasma gondii -
infected B cell-deficient mice. *Microbiol. Immunol.* **47**, 997-1003 (2003).

18. Kang, H., Remington, J. S. & Suzuki, Y. Decreased resistance of B cell-deficient mice to
infection with Toxoplasma gondii despite unimpaired expression of IFN-gamma, TNF-alpha, and
inducible nitric oxide synthase. *J. Immunol.* **164**, 2629-2634 (2000).

19. Sayles, P. C., Gibson, G. W. & Johnson, L. L. B cells are essential for vaccination-induced
resistance to virulent Toxoplasma gondii. *Infect. Immun.* **68**, 1026-1033 (2000).

20. Couper, K. N., Roberts, C. W., Brombacher, F., Alexander, J. & Johnson, L. L. Toxoplasma
gondii-specific immunoglobulin M limits parasite dissemination by preventing host cell invasion.
Infect. Immun. **73**, 8060-8068 (2005).

21. Mineo, J. R. *et al*. Antibodies to Toxoplasma gondii major surface protein (SAG-1, P30)
inhibit infection of host cells and are produced in murine intestine after peroral infection. *J.*
Immunol. **150**, 3951-3964 (1993).

22. Joiner, K. A., Fuhrman, S. A., Miettinen, H. M., Kasper, L. H. & Mellman, I. Toxoplasma
gondii: fusion competence of parasitophorous vacuoles in Fc receptor-transfected fibroblasts.
Science **249**, 641-646 (1990).

23. Mordue, D. G., Hakansson, S., Niesman, I. & Sibley, L. D. Toxoplasma gondii resides in a
vacuole that avoids fusion with host cell endocytic and exocytic vesicular trafficking pathways.
Exp. Parasitol. **92**, 87-99 (1999).

24. Zhao, Y., Marple, A. H., Ferguson, D. J., Bzik, D. J. & Yap, G. S. Avirulent strains of
Toxoplasma gondii infect macrophages by active invasion from the phagosome. *Proc. Natl.*
Acad. Sci. U. S. A. **111**, 6437-6442 (2014).

25. Glatman Zaretsky, A. *et al*. Infection with Toxoplasma gondii alters lymphotoxin expression
associated with changes in splenic architecture. *Infect. Immun.* **80**, 3602-3610 (2012).

26. Park, J. *et al*. Impact of Interleukin-27p28 on T and B Cell Responses during
Toxoplasmosis. *Infect. Immun.* **87**, 10.1128/IAI.00455-19. Print 2019 Dec (2019).

27. Stumhofer, J. S., Silver, J. S. & Hunter, C. A. IL-21 is required for optimal antibody
production and T cell responses during chronic Toxoplasma gondii infection. *PLoS One* **8**,
e62889 (2013).

28. Smith, F. L. & Baumgarth, N. B-1 cell responses to infections. *Curr. Opin. Immunol.* **57**, 23-
31 (2019).

29. Alugupalli, K. R. *et al*. B1b lymphocytes confer T cell-independent long-lasting immunity.
Immunity **21**, 379-390 (2004).

30. Haas, K. M., Poe, J. C., Steeber, D. A. & Tedder, T. F. B-1a and B-1b cells exhibit distinct
developmental requirements and have unique functional roles in innate and adaptive immunity
to S. pneumoniae. *Immunity* **23**, 7-18 (2005).

31. Gil-Cruz, C. *et al*. The porin OmpD from nontyphoidal Salmonella is a key target for a
protective B1b cell antibody response. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9803-9808 (2009).

32. Cunningham, A. F. *et al*. B1b cells recognize protective antigens after natural infection and
vaccination. *Front. Immunol.* **5**, 535 (2014).

33. Yang, Y. *et al*. Antigen-specific antibody responses in B-1a and their relationship to natural
immunity. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 5382-5387 (2012).

34. Pape, K. A., Taylor, J. J., Maul, R. W., Gearhart, P. J. & Jenkins, M. K. Different B cell
populations mediate early and late memory during an endogenous immune response. *Science*
331, 1203-1207 (2011).

35. Krishnamurty, A. T. *et al*. Somatically Hypermutated Plasmodium-Specific IgM(+) Memory B
Cells Are Rapid, Plastic, Early Responders upon Malaria Rechallenge. *Immunity* **45**, 402-414
(2016).

36. Elbez-Rubinstein, A. *et al*. Congenital toxoplasmosis and reinfection during pregnancy: case
report, strain characterization, experimental model of reinfection, and review. *J. Infect. Dis.* **199**,
280-285 (2009).

37. Jensen, K. D. *et al*. Toxoplasma gondii superinfection and virulence during secondary
infection correlate with the exact ROP5/ROP18 allelic combination. *MBio* **6**, 2280 (2015).

38. Splitt, S. D. *et al*. PD-L1, TIM-3, and CTLA-4 blockade fail to promote resistance to
secondary infection with virulent strains of Toxoplasma gondii. *Infect. Immun.* (2018).

39. McLeod, R., Skamene, E., Brown, C. R., Eisenhauer, P. B. & Mack, D. G. Genetic regulation
of early survival and cyst number after peroral Toxoplasma gondii infection of A x B/B x A
recombinant inbred and B10 congenic mice. *J. Immunol.* **143**, 3031-3034 (1989).

40. Jensen, K. D. *et al*. Toxoplasma gondii rhoptry 16 kinase promotes host resistance to oral
infection and intestinal inflammation only in the context of the dense granule protein GRA15.
Infect. Immun. **81**, 2156-2167 (2013).

41. Brown, C. R. *et al*. Definitive identification of a gene that confers resistance against
Toxoplasma cyst burden and encephalitis. *Immunology* **85**, 419-428 (1995).

42. Brown, C. R. & McLeod, R. Class I MHC genes and CD8+ T cells determine cyst number in
Toxoplasma gondii infection. *J. Immunol.* **145**, 3438-3441 (1990).

43. Hassan, M. A., Butty, V., Jensen, K. D. & Saeij, J. P. The genetic basis for individual
differences in mRNA splicing and APOBEC1 editing activity in murine macrophages. *Genome*
Res. **24**, 377-389 (2014).

44. Hassan, M. A. *et al*. Transcriptional and Linkage Analyses Identify Loci that Mediate the
Differential Macrophage Response to Inflammatory Stimuli and Infection. *PLoS Genet.* **11**,
e1005619 (2015).

45. Schuster, M., Annemann, M., Plaza-Sirvent, C. & Schmitz, I. Atypical IkappaB proteins -
nuclear modulators of NF-kappaB signaling. *Cell. Commun. Signal.* **11**, 23-23 (2013).

46. Touma, M. *et al*. Impaired B cell development and function in the absence of IkappaBNS. *J.*
Immunol. **187**, 3942-3952 (2011).

47. Arnold, C. N. *et al*. A forward genetic screen reveals roles for Nfkbid, Zeb1, and Ruvbl2 in
humoral immunity. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 12286-12293 (2012).

48. Pedersen, G. K. *et al*. Heterozygous Mutation in IkappaBNS Leads to Reduced Levels of
Natural IgM Antibodies and Impaired Responses to T-Independent Type 2 Antigens. *Front.*
Immunol. **7**, 65 (2016).

49. Touma, M. *et al*. Functional role for I kappa BNS in T cell cytokine regulation as revealed by
targeted gene disruption. *J. Immunol.* **179**, 1681-1692 (2007).

50. Schuster, M. *et al*. IkappaB(NS) protein mediates regulatory T cell development via
induction of the Foxp3 transcription factor. *Immunity* **37**, 998-1008 (2012).

51. Kuwata, H. *et al*. IkappaBNS inhibits induction of a subset of Toll-like receptor-dependent
genes and limits inflammation. *Immunity* **24**, 41-51 (2006).

52. Berthier-Vergnes, O. *et al*. Gene expression profiles of human melanoma cells with different
invasive potential reveal TSPAN8 as a novel mediator of invasion. *Br. J. Cancer* **104**, 155-165
(2011).

53. Zhao, K., Wang, Z., Hackert, T., Pitzer, C. & Zoller, M. Tspan8 and Tspan8/CD151 knockout
mice unravel the contribution of tumor and host exosomes to tumor progression. *J. Exp. Clin.*
Cancer Res. **37**, 312-6 (2018).

54. Khoenkhoen, S. *et al*. TACI expression and plasma cell differentiation are impaired in the
absence of functional IkappaBNS. *Immunol. Cell Biol.* **97**, 485-497 (2019).

55. Chu, H. H. *et al*. Continuous Effector CD8(+) T Cell Production in a Controlled Persistent
Infection Is Sustained by a Proliferative Intermediate Population. *Immunity* **45**, 159-171 (2016).

56. Cohen, S. B. *et al*. CXCR3-dependent CD4(+) T cells are required to activate inflammatory
monocytes for defense against intestinal infection. *PLoS Pathog.* **9**, e1003706 (2013).

57. Shah, S., Grotenbreg, G. M., Rivera, A. & Yap, G. S. An extrafollicular pathway for the
generation of effector CD8(+) T cells driven by the proinflammatory cytokine, IL-12. *Elife* **4**,
10.7554/eLife.09017 (2015).

58. Luo, L. *et al.* SCIMP is a transmembrane non-TIR TLR adaptor that promotes
proinflammatory cytokine production from macrophages. *Nat. Commun.* **8**, 14133 (2017).

59. Suzuki, K. *et al.* Semaphorin 7A initiates T-cell-mediated inflammatory responses through
alpha1beta1 integrin. *Nature* **446**, 680-684 (2007).

60. Baumgarth, N., Jager, G. C., Herman, O. C. & Herzenberg, L. A. CD4+ T cells derived from
B cell-deficient mice inhibit the establishment of peripheral B cell pools. *Proc. Natl. Acad. Sci. U.*
S. A. **97**, 4766-4771 (2000).

61. Choi, Y. S. & Baumgarth, N. Dual role for B-1a cells in immunity to influenza virus infection.
J. Exp. Med. **205**, 3053-3064 (2008).

62. Kreuk, L. S. *et al.* B cell receptor and Toll-like receptor signaling coordinate to control
distinct B-1 responses to both self and the microbiota. *Elife* **8**, 10.7554/eLife.47015 (2019).

63. Lalor, P. A., Herzenberg, L. A., Adams, S. & Stall, A. M. Feedback regulation of murine Ly-1
B cell development. *Eur. J. Immunol.* **19**, 507-513 (1989).

64. Kantor, A. B., Stall, A. M., Adams, S., Herzenberg, L. A. & Herzenberg, L. A. Differential
development of progenitor activity for three B-cell lineages. *Proc. Natl. Acad. Sci. U. S. A.* **89**,
3320-3324 (1992).

65. Nguyen, T. T. *et al.* The IgM receptor FcmuR limits tonic BCR signaling by regulating
expression of the IgM BCR. *Nat. Immunol.* **18**, 321-333 (2017).

66. Blandino, R. & Baumgarth, N. Secreted IgM: New tricks for an old molecule. *J. Leukoc. Biol.*
106, 1021-1034 (2019).

67. Savage, H. P. *et al.* TLR induces reorganization of the IgM-BCR complex regulating murine
B-1 cell responses to infections. *Elife* **8**, 10.7554/eLife.46997 (2019).

68. Bikah, G., Carey, J., Ciallella, J. R., Tarakhovsky, A. & Bondada, S. CD5-mediated negative
regulation of antigen receptor-induced growth signals in B-1 B cells. *Science* **274**, 1906-1909
(1996).

69. Baumgarth, N. B-1 Cell Heterogeneity and the Regulation of Natural and Antigen-Induced
IgM Production. *Front. Immunol.* **7**, 324 (2016).

70. Castigli, E. *et al.* TACI and BAFF-R mediate isotype switching in B cells. *J. Exp. Med.* **201**,
35-39 (2005).

71. Nussenzweig, R. S., Vanderberg, J., Most, H. & Orton, C. Protective immunity produced by
the injection of x-irradiated sporozoites of plasmodium berghei. *Nature* **216**, 160-162 (1967).

72. Rieckmann, K. H. Human immunization with attenuated sporozoites. *Bull. World Health*
Organ. **68 Suppl**, 13-16 (1990).

73. Hoffman, S. L. *et al*. Protection of humans against malaria by immunization with radiation-
attenuated Plasmodium falciparum sporozoites. *J. Infect. Dis.* **185**, 1155-1164 (2002).

74. Rieckmann, K. H., Carson, P. E., Beaudoin, R. L., Cassells, J. S. & Sell, K. W. Letter:
Sporozoite induced immunity in man against an Ethiopian strain of Plasmodium falciparum.
Trans. R. Soc. Trop. Med. Hyg. **68**, 258-259 (1974).

75. Clyde, D. F., McCarthy, V. C., Miller, R. M. & Hornick, R. B. Specificity of protection of man
immunized against sporozoite-induced falciparum malaria. *Am. J. Med. Sci.* **266**, 398-403
(1973).

76. RTS, S Clinical Trials Partnership. Efficacy and safety of RTS,S/AS01 malaria vaccine with
or without a booster dose in infants and children in Africa: final results of a phase 3, individually
randomised, controlled trial. *Lancet* **386**, 31-45 (2015).

77. Hemler, M. E. Tetraspanin functions and associated microdomains. *Nat. Rev. Mol. Cell Biol.*
6, 801-811 (2005).

1117

A

A/J

Percent survival

- VAND
- GUY-DOS
- GPHT
- FOU
- TgCATBr5
- GUY-MAT
- MAS

Day

B10.A

Percent survival

Day

B10

Percent survival

Day

B

AJ B10.A B10 B6

Day 1

Day 4

Day 8

Photons/sec/cm²/sr

MAS-luc secondary infection

C

GT1 secondary infection

Percent survival

- B10
- B10.A
- A/J
- F1

***P < 0.0001

Day

D

GT1 brain co-infection

Percent of mice

****

- A/J
- F1

Superinfected   Not Superinfected

A

WT PerC transfer — 1° infection — 2° infection

Birth — 2 day — 7 weeks — 12 weeks

Percent survival vs Days following 2° infection (RH)
Bumble mice

- No transfer
- PerC transfer

***

B

Parasite-specific antibody (after 1° infection)

- C57BL/6J
- Bumble +PerC
- Bumble
- no serum

IgM   IgG3

IgG1   IgG2a/c   IgG2b

C

C57BL/6
IgH-b

PerC IgH-a transfer — Deplete endogenous B cells with anti-IgH-b 2x weekly — Primary infection

Birth   2d — 6 weeks — 12 weeks

IgH-a B-1 cells engraft   IgH-b B-2 cells grow back

IgH chimeric mice

B-1: IgH-a
B-2: IgH-b

D

Day 14
1° infection

B-1   B-2

anti-IgM-a (B-1)   anti-IgM-b (B-2)

B-1 vs B-2

Infected IgH chimeric mice

Staining controls

Infected IgH-b mice
Infected IgH-a mice

Day 30
1° infection

B-1   B-2

anti-IgM-a (B-1)   anti-IgM-b (B-2)