Multispecies integrated population model reveals bottom-up dynamics in a seabird predator-prey

system

Maud Quéroué[1], Christophe Barbraud[2], Frédéric Barraquand[3], Daniel Turek[4], Karine Delord[2],

Nathan Pacoureau[5], Olivier Gimenez[1]

[1]CEFE UMR 5175, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3,

Montpellier, France

[2]CEBC UMR7372, CNRS – La Rochelle Université, 79360 Villiers en Bois, France

[3]Institute of Mathematics of Bordeaux, CNRS, Talence, France

[4]Department of Mathematics and Statistics, Williams College, Williamstown, MA, USA

[5]Department of Biological Sciences, Earth to Ocean Research Group, Simon Fraser University,

Burnaby, British Columbia, Canada

Correspondence author: Maud Quéroué, CEFE UMR 5175, Univ Montpellier, CNRS, EPHE,

IRD, Univ Paul Valéry Montpellier 3, Montpellier, France

E-mail: maud.queroue@gmail.com

Abstract

Assessing the effects of climate and interspecific relationships on communities is challenging

because of the complex interplay between species population dynamics, their interactions, and

the need to integrate information across several biological levels (individuals – populations –

communities). Usually used to quantify species interactions, integrated population models

(IPMs) have recently been extended to communities. These models allow fitting multispecies

matrix models to data from multiple sources while simultaneously accounting for various sources

of uncertainty in each data source. We used multispecies IPMs accommodating climate

conditions to quantify the relative contribution of climate vs. interspecific interactions on

demographic parameters, such as survival and breeding success, in the dynamics of a predator-

prey system. We considered a stage-structured predator–prey system combining 22 years of

capture–recapture data and population counts of two seabirds, the Brown Skua (*Catharacta*

lönnbergi) and its main prey the Blue Petrel (*Halobaena caerulea*) both breeding on the

Kerguelen Islands in the Southern Ocean. Our results showed that climate and predator-prey

interactions drive the demography of skuas and petrels in different ways. The breeding success

of skuas appeared to be largely driven by the number of petrels and to a lesser extent by

intraspecific density-dependence. In contrast, there was no evidence of predation effects on the

demographic parameters of petrels, which were affected by oceanographic factors (chlorophyll a

and sea surface temperature anomalies). We conclude that bottom-up mechanisms are the main

drivers of this skua-petrel system. We discuss the mechanisms by which climate variability and

predator-prey relationships may affect the demographic parameters of these seabirds. Taking into

account both species interactions and environmental covariates in the same analysis improved

our understanding of species dynamics.

Keywords: Bayesian inference, Demography, Environmental variations, Integrated Population

Model, Matrix population model, Nimble, Predator-Prey interactions

Introduction

The effects of climate changes on the diversity and the structure of communities have been

reported repeatedly (Walther et al. 2002, Parmesan 2006, Hoegh-Guldberg and Bruno 2010,

Miller et al. 2018). However, the underlying mechanisms remain poorly understood due to the

complex dynamics of interacting species: within species, between species and between species

and the environment (Godfray and May 2014). Following disturbance, such as changes in

environmental conditions, the abundance and distribution of species are expected to be modified

according to the position and extent of the species' niche (Thomas et al. 2004). Because the

effects of environmental variability on mortality, fecundity and dispersal may differ between

species (Grosbois et al. 2008, Jenouvrier 2013), changes in structure and diversity appear at the

community level. However, studying species-by-species responses to environmental changes

may overlook the role played by species interactions on those responses, and contribute to a

lesser extent to the larger understanding of species interactions that is required by community

ecology.

Population dynamics models have been used to understand the effect of interspecific interactions

and environment on species demography. However, these models are in general not

demographically structured (Stenseth et al. 2015, Pacoureau et al. 2019a, Stoessel et al. 2019) or

only partially (Millon et al. 2014, Saunders et al. 2018, Pacoureau et al. 2019b). Unstructured

approaches consider individuals as being equivalent but differences in size, age and ontogenic

stages exist within a population and may be of importance in the context of interspecific

interactions. As argued by Miller and Rudolf (2011), the consideration of the stage structure of

populations can lead to a better understanding of community structure and dynamics.

Interactions between species such as predation or competition do not necessarily have a

homogeneous impact on the different stages of the interacting species. For example, young

individuals might be predominantly preyed upon in carnivore–ungulate systems (Gervasi et al.

2015). Therefore, to detect and understand species interactions, we need to consider jointly the

demography of several stage-structured populations (Oken and Essington 2015).

Although well developed for single-species dynamics (Tuljapurkar and Caswell 1997, Caswell

2001), demographic stage-structured models have received little attention in community ecology

(but see Chu and Adler (2015) on a plant system). The difficulty is that multispecies demography

analysis requires integrating information across several biological levels (individual – population

– community) which, in turn, requires unifying all available data sources into a single

framework. Integrated population models (IPMs) have been recently developed to infer

population demography by making complete use of all available sources of information (see

Schaub and Abadi 2011, and Zipkin and Saunders 2018 for reviews). In their simplest form,

these models combine population counts and demographic data into a single framework, which

allows the estimation of demographic parameters while simultaneously accounting for various

sources of uncertainty in each data source (e.g. measurement error or parameter estimation)

(Besbeas et al. 2002). The IPM framework has been extended to multiple species (Péron and

Koons 2012) for competition/parasitism, and more recently for predator-prey interactions

(Barraquand and Gimenez 2019).

Here, our main objective was to quantify the relative contribution of environmental changes and

species interactions on demographic parameters of a predator and its prey. Therefore, we used a

multispecies IPM framework accommodating the effects of local and global climatic conditions

on demographic parameters, such as survival and breeding, while explicitly considering species

interactions. We applied our approach on a stage-structured predator–prey system combining 22

years of capture-recapture data and population counts on two seabirds, the Brown Skua

(*Catharacta lönnbergi*) and its main prey the Blue Petrel (*Halobaena caerulea*) ('skua' and

'petrel' hereafter) breeding on the Kerguelen Islands in the Southern Ocean.

Because seabirds often occupy high level positions in food-webs, bottom-up forcing which

implies population regulation through climate driven limitation in food availability, has long

been featured as the dominant paradigm to understand their dynamics (Lack 1967, Aebischer et

al. 1990, Stenseth et al. 2002). Seabird foraging behavior and demography reflect the influences

of climate variability which directly impacts biological processes in marine ecosystems and

cascade through food webs up to seabirds (Barbraud and Weimerskirch 2001, Jenouvrier et al.

2003). However, top-down pressures as predation at breeding colonies are also known to affect

the vital rates of seabirds (Hipfner et al. 2012). There is increasing evidence that bottom-up and

top-down processes often act in concert and differently affect demographic parameters (Suryan

et al. 2006, Horswill et al. 2014, 2016). For example, the effects of predation and resources

limitation caused breeding failure of Black-legged Kittiwakes (*Rissa tridactyla*) (Regehr and

Montevecchi 1997) and population declines of Arctic Skuas (*Stercorarius parasiticus*) (Perkins

et al. 2018). Therefore, quantifying the relative strength of environmental conditions and

predator-prey effects is essential for a better understanding of the drivers of population dynamics

in seabirds. This is all the more important as climate changes impact the physical properties of

the oceans, including the Southern Ocean (Gille 2002, Han et al. 2014) and, through the trophic

food web, affect demography and population dynamics of seabirds (Barbraud et al. 2012,

Sydeman et al. 2015), including some of the species studied here (Barbraud and Weimerskirch

2003).

Using a multispecies IPM, we assessed the relative contribution of environment and predator-

prey interactions on seabirds' demographic parameters. We estimated survival and adult

breeding success for the two interacting species, and assessed the impacts of climatic conditions

on these demographic parameters to understand the contribution of predator-prey interactions in

shaping population dynamics.

Materials and Methods

Study site and Species

Skuas and petrels were studied on Mayes Island (49°28'S, 69°57'E), a 2.8 km² uninhabited island

of the Kerguelen Islands in the Southern Ocean where the two species breed during the austral

summer (October-February).

The petrel is a small (150–250g) long-lived seabird belonging to the family of *Procellariiformes*.

At Kerguelen Islands, petrels feed on macrozooplankton and micronekton feeder, mainly

crustaceans and fishes (Cherel et al. 2002, 2014). Individuals from Mayes Island spend the

nonbreeding season (from mid-February to September) between the polar front and the northern

limit of the pack-ice (57-62°S) between longitudes 20°W and 90°E (Cherel et al. 2016). Birds

return to breeding colonies in early September (Quillfeldt et al. 2020). Mayes Island is covered

with dry soils and dense vegetation, providing suitable breeding sites for approximately 142,000

breeding pairs of these burrowing petrels (Barbraud and Delord 2006). In late October, a single

egg is laid in a burrow dug in peat soil under tall and dense vegetation. The incubation lasts 45-

49 days and the chick rearing period 43-60 days (Jouventin et al. 1985). The chick fledges in

early February. Both sexes participate in parental care by alternating foraging trips during the

incubation and fledging periods.

The skua is a medium sized (1.1 – 2.2 kg) long-lived seabird belonging to the family of

Charadriiformes. On Mayes Island between 80 and 120 pairs breed annually (Mougeot et al.

1998). Breeding pairs form in October with a high mate fidelity, and generally establish

themselves in the same territory each year (Parmelee and Pietz 1987), which they tenaciously

defend throughout the breeding season. Generally, two eggs are laid between October and

December. The incubation lasts 28-32 days and the chicks rearing period 40-50 days (Higgins

and Davied 1996). Skuas are extremely plastic in their foraging techniques and adapt their diet

depending on the local availability of prey (Carneiro et al. 2015). On Mayes Island, during the

breeding season, Blue Petrels represent 95% of the skua diet (Pacoureau et al. 2019c). Skuas

from Mayes Island overwinter in the southern hemisphere between 10°E and 150°E (Delord et

al. 2018).

During the breeding period on Mayes Island, the predation of petrels by skuas takes place mainly

at night, when petrels come out or arrive at their burrows (Mougeot and Bretagnolle 2000a).

Skuas mostly prey on petrels on the ground, but they can also catch petrels in flight (Mougeot et

al. 1998, Pacoureau et al. 2019c). Vocalizing petrels, especially those without partners, are more

easily detected by skuas during the courtship period (Mougeot and Bretagnolle 2000b). Skuas

may also prey on chicks during the fledging period.

Count and capture-recapture data

Data of both skuas and petrels were collected during the breeding seasons from 1996/1997 to

2017/2018. For convenience, breeding seasons are named from 1996 to 2017 hereafter. The time

interval used in our model starts before the wintering of species and ends at the end of the

breeding period. Two types of data were used: count data corresponding to the number of

burrows or territories occupied by seabirds and capture-recapture (CR) data of adult seabirds

found on the monitored area. Each year, adult individuals of both species were checked at

specific times following the species phenology to determine the breeding status of each bird. The

breeding status of marked birds was determined at the end of the breeding period. In the

following we describe how the data were collected for the two species. For clarity, all parameters

for skuas are indicated by S and by P for petrels.

Around 200 individually marked burrows of petrels were inspected each year from early-to-mid

November just after the egg-laying to check for eggs and to identify marked adults, and then in

late January just before fledging of the chicks. Each year since 1985 (see Barbraud and

Weimerskirch 2005), all fledglings as well as new individuals found in burrows were marked

with a stainless steel band (captured by hand, marked, and replaced in their burrow). Petrels

never observed with an egg or a chick during a given breeding season were considered as

nonbreeders (NB). Individuals were identified as breeders if they laid a single egg or raised a

chick and as successful breeders if their chick fledged (SB). Two categories of failed breeders

were used depending on the stage of failure: egg stage (FBE) or chick stage (FBC). Given that

the first sampling period occurred just after laying, it is very unlikely that nonbreeders were

failed breeders. These breeding statuses allowed the construction of the individual capture

histories (Ch_P) and constituted our CR data. The annual number of adult petrels (Y_P), *i.e.* count

data, was estimated as the number of occupied burrows. Each occupied burrow was considered

as being frequented by a pair of petrels. We considered that this count included all adult

individuals, both breeders and non-breeders.

For skuas, each year since 1991, the eastern side of Mayes Island was inspected to identify

territories of skuas. A territory was considered established when a pair strongly defended an area

against other skuas (Mougeot et al. 1998). Around 50 nesting territories were visited four to eight

times from mid-October (after egg-laying) to late-February (just before fledging of the chicks)

each year. Chicks just before fledging, as well as new adult individuals, were marked with a

metal ring and a plastic ring to facilitate individual identification using binoculars. Breeding

status was determined by checking the nest contents for the presence of eggs or young chicks.

Skuas never observed with an egg or a chick were considered as nonbreeders (NB). Individuals

were identified as breeders if they laid at least one egg or raised a chick. If the eggs did not hatch

or the chicks died, both members of the pair were considered as failed breeders (FB). Given that

the first sampling period occurred just after laying, it is very unlikely that nonbreeders

represented failed breeders. Successful breeders were defined as individuals that fledged one or

two chicks, and were denoted as $SB1$ or $SB2$, respectively. These breeding statuses allowed the

construction of the individual capture histories (Ch_S) and constituted our CR data. The annual

number of skuas (Y_S), *i.e.* count data, was estimated as the number of territories and each

territory was considered occupied by a pair of skuas. We considered that this count included all

adult individuals, both breeders and non-breeders.

For both species, individual breeding status could be considered as "uncertain" (C) in case of

difficulties to assign their breeding status (lack of information, missed checks, individuals never

re-observed). Only adult individuals that have bred at least once between the 1996 and the 2017

breeding seasons were kept in the data set for analysis to eliminate potential transient individuals

($n = 318$ for skuas and $n = 1210$ for petrels). Individual capture histories (Ch) started at their first

breeding attempt recorded. Based on the high probability of observing breeders in the study site,

we assumed that the first breeding attempt was correctly detected. New individuals found in

monitored burrows or territories are considered as immigrants to the study site (N_{im}).

The presence of chicks was used to assign a breeding status to adult individuals captured in the

breeding area. In order to maintain the independence of the data, we did not include information

on chicks in the model. Therefore, the fecundity was a fixed value. We considered one chick for

each pair of seabird, considered as successful breeders ($N_{SB,P}$) for petrels or successful breeders

with one chick ($N_{SB1,S}$) for skuas ($f_{SB,P}$ and $f_{SB1,S}$ are equal to 1, respectively). For skuas that

successfully fledged two chicks ($N_{SB2,S}$), we considered two chicks per pair of skuas ($f_{SB2,S}$ is

equal to 2). Since juveniles only return to the breeding sites as adults to attempt to breed for the

first time (from four year old or older), we did not have data on juvenile states.

Integrated Population Model

We built a two-species IPM that combines count and CR data and allows estimating abundances

and demographic rates (Besbeas et al. 2002, Schaub and Abadi 2011). More specifically, we

connected two IPMs, one for predatory skuas and one for petrels, their main prey, through

explicit predator-prey relationships (Barraquand and Gimenez 2019). We incorporated the

effects of predation within species-specific vital rates such as survival and breeding parameters.

This IPM is structured by states which represent life history states (Fig. 1). We built two

likelihoods, one for the CR data and the other for the count data which we combined into a joint

likelihood.

In the following, we detail the state process following a biological timeline and we explain the

different likelihood used. The structure was the same for the two species but states differed in

relation to species biology (Fig. 1). The two main differences were: (1) skuas could have up to

two chicks versus only one for petrels, (2) the failed-breeder stage in petrels could be split

further according to the timing of failure (failure at the incubation vs. chick-rearing stage). For

clarity, parameters are indexed by S (for skuas) or P (for petrels) when differences occur, or by X

(for S or P) when the structure is the same for both species. We used Poisson (Po) and binomial

(Bin) distributions to account for demographic stochasticity. Notations of all parameters and state

variables are detailed in Appendix S1: Table S1.

State process

Offspring production

The estimated number of skuas and petrels in their first year *i.e.* between 0 and 1 year old

($N_{J1,S,t}$) at year *t,* is modelled with a Poisson distribution:

$N_{J1,S,t} \sim Po(0.5 \times f_{SB1,S} \times N_{SB1,S,t-1} + 0.5 \times f_{SB2,S} \times N_{SB2,S,t-1})$ (1)

$N_{J1,P,t} \sim Po(0.5 \times f_{SB,P} \times N_{SB,P,t-1})$ (2)

with $N_{J1,S}$ the number of chicks produced by all successful skua breeders ($N_{SB1,S}$ and $N_{SB2,S}$)

according to their fecundity ($f_{SB1,S}$: 1 chick and $f_{SB2,S}$: 2 chicks per female skua, sex ratio: 0.5).

For petrels, $N_{J1,P}$ is also Poisson distributed but with only one chick ($f_{SB,P}$) per estimated

successful female breeder ($N_{SB,P}$ with a sex ratio of 0.5).

Juvenile survival

The number of juveniles between one and two years (N_{J2}), two and three years (N_{J3}), and three

and four years (N_{J4}), are modelled with binomial distributions:

$N_{J2,X,t} \sim Bin(\phi_{J1,X}, N_{J1,X,t-1})$ (3)

$N_{J3,X,t} \sim Bin(\phi_{J2,X}, N_{J2,X,t-1})$ (4)

$N_{J4,X,t} \sim Bin(\phi_{J3,X}, N_{J3,X,t-1})$ (5)

with the apparent survival between one and two years (ϕ_{J1}), between two and three years (ϕ_{J2})

and between three and four years (ϕ_{J3}) respectively. As we observed only adult breeding birds,

we had no information on the juvenile phase. We assumed that juvenile apparent survival

increased with age (Greig et al. 1983, Grande et al. 2009, Fay et al. 2015), as experienced birds

are on average more effective in foraging (Daunt et al. 2007), in competing with conspecifics or

in avoiding predators:

$$logit\left(\phi_{Jage\_i,X}\right) = \lambda_{1,X} + \lambda_{2,X} \times age\_i \tag{6}$$

where ϕ_J is the juvenile apparent survival, age_i the age of the juvenile state (from N_{J1} to N_{J4}),

λ_1 the intercept and λ_2 the slope which is constrained to be positive.

Juvenile first breeding attempt

The first breeding attempt in skuas and petrels could start from age four. Four years old

individuals and older that did not attempt to breed are in the state (N_{J4+}). The individuals that

attempted to breed for the first time with a first breeding attempt probability Pr are in the state

N_{J4B} and the individuals that did not attempt to breed are in the state N_{J4NB}:

$$N_{J4B,X,t} \sim Bin\left(Pr_{X,t}, \phi_{J4,X} \times N_{J4+,X,t-1}\right) \tag{7}$$

$$N_{J4NB,X,t} = \phi_{J4,X} \times N_{J4+,X,t-1} - N_{J4B,X,t} \tag{8}$$

260 with ϕ_{J4} the apparent survival for the N_{J4+} state. The N_{J4+} state includes individuals that did not

261 attempt to breed (N_{J4NB}) and individuals aged between three and four years (N_{J4}):

$$N_{J4+,X,t} = N_{J4,X,t} + N_{J4NB,X,t} \tag{9}$$

Adult survival

For the two species, we modelled the number of surviving adults (N_{alive}) at year t among the

total number of adult individuals (N_{adtot}) at year *t-1* with a binomial distribution, with ϕ the

adult apparent survival:

$$N_{alive,X,t} \sim Bin\left(\phi_{X,t-1}, N_{adtot,X,t-1}\right) \tag{10}$$

Breeding probability

The number of adult individuals that have bred or not bred among those that survived (N_{alive}) is

modelled as:

$$N_{Balive,X,t} \sim Bin\left(\beta_{X,t-1}, N_{alive,X,t}\right) \tag{11}$$

$$N_{NB,X,t} = N_{alive,X,t} - N_{Balive,X,t} \tag{12}$$

with β the probability of breeding, N_{Balive} the number of adult breeders that survived and N_{NB}

the number of adult nonbreeders. As capture histories started at their first breeding attempt

recorded, the immigrants, *i.e.* newly marked individuals (N_{im}) coming for the first time in the

colony, were considered as breeders. Then, the total number of breeders (N_B) corresponds to the

sum of the number of adult breeders that survived (N_{Balive}), the number of immigrants (N_{im})

and the number of juveniles attempting to breed for the first time (N_{J4B}):

$$N_{B,X,t} = N_{Balive,X,t} + N_{J4B,X,t} + N_{im,X,t} \tag{13}$$

The total number of adults (N_{adtot}) corresponds to the sum of nonbreeders (N_{NB}) and breeders

(N_B):

$N_{adtot,X,t} = N_{NB,X,t} + N_{B,X,t}$ (14)

Breeding success

Breeding success and failure are modelled differently for skuas and petrels. For skuas, the

numbers of failed breeders ($N_{FB,S}$) and successful breeders ($N_{SB,S}$) are modelled following a

binomial distribution:

$N_{SB,S,t} \sim Bin\left(\gamma_{S,t-1}, N_{B,S,t}\right)$ (15)

$N_{FB,S,t} = N_{B,S,t} - N_{SB,S,t}$ (16)

with γ_S the probability of a successful breeding. A successful breeder can then have one or two

chicks, respectively $N_{SB1,S}$ and $N_{SB2,S}$ and this is modelled following a binomial distribution:

$N_{SB2,S,t} \sim Bin\left(\delta_{S,t-1}, N_{SB,S,t}\right)$ (17)

$N_{SB1,S,t} = N_{SB,S,t} - N_{SB2,S,t}$ (18)

with δ_S the probability of producing two chicks rather than one among the successful breeders.

For petrels, there are two states for failed breeders: one with petrels that failed to hatch their egg

(named failed breeder at the egg stage $N_{FBE,P}$) and the second with petrels that failed to fledge

their chick (named failed breeder at the chick stage $N_{FBC,P}$). Hence, there is a parameter of

successful hatching (ω_P). The numbers of petrels with an egg that successfully hatched ($N_{SH,P}$)

and the failed breeders at the egg stage ($N_{FBE,P}$) were modelled following a binomial

distribution:

$N_{SH,P,t} \sim Bin\left(\omega_{P,t-1}, N_{B,P,t}\right)$ (19)

$N_{FBE,P,t} = N_{B,P,t} - N_{SH,P,t}$ (20)

with ω_P the probability of successful hatching. Successful breeders ($N_{SB,P}$) and failed breeders

at the chick stage ($N_{FBC,P}$) were modelled following a binomial distribution:

$N_{SB,P,t} \sim Bin\ (\gamma_{P,t-1}, N_{SH,P,t})$ (21)

$N_{FBC,P,t} = N_{SH,P,t} - N_{SB,P,t}$ (22)

with y_P the probability of successful breeding.

Count data

The observation equation links the observed adult population count (Y) (*i.e.* the number of

territories/burrows multiplied by two for a pair of seabird) with the true adult population size

(N_{adtot}), with an additional term for observation error:

$Y_{X,t} \sim N(N_{adtot,X,t}, \varepsilon_{X,t})$

$\varepsilon_{X,t} \sim N(0, \sigma^2_{Y,X})$ (23)

where ε is the error term and σ^2_Y its variance. As only the adult states were observed on the field,

we excluded the juvenile states from the observation equation. The likelihood for the population

count data is denoted as $L_{co,S}(Y_S | \phi_{J1,S}, \phi_{J2,S}, \phi_{J3,S}, \phi_{J4,S}, Pr_S, \phi_S, \beta_S, \gamma_S, \delta_S, N_{adtot,S})$ for skuas and

as $L_{co,P}(Y_P | \phi_{J1,P}, \phi_{J2,P}, \phi_{J3,P}, \phi_{J4,P}, Pr_P, \phi_P, \beta_P, \omega_P, \gamma_P, N_{adtot,P})$ for petrels.

Capture-recapture data

For adult CR data, we used multievent capture–recapture models to estimate the demographic

parameters (Pradel 2005). These models take into account the imperfect detectability of the

individuals as well as the uncertainty in the assignment of states to individuals (Gimenez et al.

2012).

For skuas, our multievent model includes five states: NB, FB, SB1, SB2, dead, and six events:

not seen, seen as NB, seen as FB, seen as SB1, seen as SB2, seen as C. For petrels, the five states

are: NB, FBE, FBC, SB, dead, and the six events are: not seen, seen as NB, seen as FBE, seen as

FBC, seen as SB, seen as C. The following demographic parameters were estimated for the two

species: the adult apparent survival probability (ϕ_X), the breeding probability (β_X), the

probability of successful breeding (γ_X). The probability of successful breeding with two

chicks (δ_S) was also estimated for skuas, as well as the probability of hatching (ω_P) for petrels.

Two additional parameters were also estimated: the detection probability (p_X) and the state

assignment probability of individuals with uncertain state (u_X). All parameters were time-

varying through a yearly random effect, except u (Table 1). State transitions were set to be state

dependent according to the breeding status in the previous breeding season (Table 1): Breeder

($\tilde{B}$) representing birds that attempted to breed the previous breeding season ($FB, SB1, SB2$ for

skuas or FBE, FBC, SB for petrels) or Nonbreeder ($\widetilde{NB}$) representing birds that already bred

previously but did not attempt to breed during the previous breeding season (NB). The detection

probability and the state assignment probability also depended on the breeding status (Table 1).

The likelihood for the CR data for skuas is denoted as $L_{cr,S}(Ch_S|\phi_S, \beta_S, \gamma_S, \delta_S, p_S, u_S)$

and $L_{cr,P}(Ch_P|\phi_P, \beta_P, \gamma_P, \delta_P, p_P, u_P)$ for petrels.

Joint likelihood

The joint likelihood of the skua IPM is the product of the likelihood for the count data ($L_{co,S}$)

and CR data ($L_{cr,S}$):

$L_{ipm,S}(Y_S, Ch_S | \phi_{J1,S}, \phi_{J2,S}, \phi_{J3,S}, \phi_{J4,S}, Pr_S, \phi_S, \beta_S, \gamma_S, \delta_S, N_{adtot,S}, p_S, u_S) =$

$L_{co,S}(Y_S | \phi_{J1,S}, \phi_{J2,S}, \phi_{J3,S}, \phi_{J4,S}, Pr_S, \phi_S, \beta_S, \gamma_S, \delta_S, N_{adtot,S})$

$\times L_{cr,S}(Ch_S | \phi_S, \beta_S, \gamma_S, \delta_S, p_S, u_S)$ (24)

For petrels, the product of the likelihood for the count data ($L_{co,P}$) and CR data ($L_{cr,P}$) is

denoted as: $L_{ipm,P}(Y_P, Ch_P | \phi_{J1,P}, \phi_{J2,P}, \phi_{J3,P}, \phi_{J4,P}, Pr_P, \phi_P, \beta_P, \omega_P, \gamma_P, N_{adtot,P}, p_P, u_P)$.

Interspecific relationships, intraspecific density-dependence, and environmental

covariates

We used different covariates to investigate their effects on the adult demographic parameters

estimated for the two species (Table 2). We focused only on the demographic parameters of adult

individuals because only adults were observed on the field. We tested interspecific predator-prey

relationships between skua and petrel, and intraspecific relationships with density-dependence

for both species. Moreover, we considered several climatic covariates that were suspected to

affect demographic parameters of skuas and petrels, the Southern Annular Mode (SAM) on a

large scale, and the Sea Surface Temperature anomalies (SSTa) and Chlorophyll a concentration

(Chla) on a local scale. In the following, we provide more details on covariates and how they

may affect the demography of skuas and petrels.

Predator-prey interactions

Multispecies IPMs allow us to explicitly include interspecific relationships between vital rates of one species and estimated population sizes of the other. Based on the high proportion of petrels in the diet of the skuas during the breeding season (Mougeot et al. 1998, Pacoureau et al. 2019c), we predicted that petrel adult apparent survival (ϕ_P) should decrease with the number of skuas. As skuas prey on adults and chicks during the fledging period, we predicted that the hatching success (ω_P) and fledging success (γ_P) would be impacted by the number of predators. Inversely, we predicted that a large number of petrels in the breeding colony would provide enough food resources for skua and then be favorable to their breeding success (γ_S) and breeding success with two chicks (δ_S).

Intraspecific density-dependence

We investigated the effect of intraspecific density-dependence on the demography of the two species as higher density of individuals on the breeding area can lead to and increasing competition for food resources or for territories. Skuas are highly territorial and defend their territories vigorously during the whole breeding season. The most violent fights may even lead to their death. Moreover, the limited number of territories could cause emigration of skuas without territory. Thus, we predicted that the apparent survival (ϕ_S), *i.e.* the joint estimation of the mortality and emigration, would be negatively impacted by the number of skuas. This limited number of territories could also lead to a negative density-dependence relationship between breeding probability (β_S) and population density. The energetic cost and the time spent in defending a territory throughout the breeding season may limit the time spent searching for food, potentially limiting energy investment in reproduction. We thus predicted a negative effect of

population density on the successful breeding parameter (γ_S) and the probability to have two

chicks rather than one for successful breeders (δ_S). For petrels, we also tested the effects of

intraspecific competition for food resources, which could affect their adult apparent survival

(ϕ_P) and their breeding parameters: breeding probability (β_P), hatching (ω_P) and fledging

success (γ_P).

Environmental covariates

Climate variability impacts biological processes in marine ecosystems, which cascade through

food webs and are integrated by seabirds (Barbraud and Weimerskirch 2001, Jenouvrier et al.

2003). Hence, we considered several covariates that are suspected to affect populations of petrels

and skuas through these bottom-up mechanisms. All covariates are used as proxies of food

availability at sea at different scales. In the following, we explain how environmental conditions

may impact the two species based on their diet and distribution.

Because skuas have broad wintering areas (Delord et al. 2018), we tested a large-scale

environmental covariate, the SAM. In contrast with their diet during the breeding season

specialized on the Blue Petrel, during winter skuas adopt a mixed diet composed of low trophic

level preys, such as macrozooplankton and crustaceans (Delord et al. 2018). We hypothesized

that availability of food resources at sea during the austral winter might have an effect on the

body condition of skuas and then affect the survival of skuas. Moreover, skuas may experience a

carry-over effect as the additional energy invested by individuals to maintain themselves during

poor wintering conditions may have repercussion on their ability to breed the next breeding

season (Harrison et al. 2011, Bogdanova et al. 2017).

For petrels, the wintering areas have been determined (Cherel et al. 2016) allowing us to test two

covariates used at local scale, the SSTa and the Chla, in addition of the SAM. As their diet is

mainly composed of crustaceans and fish feeding at low trophic levels (Cherel et al. 2002, 2014),

the food availability at sea may impact the survival of petrels. Moreover, during the breeding

season, male and female petrels take turns, one incubating the egg and fasting and the other

foraging at sea, which results in substantial variation in their body mass (Chaurand and

Weimerskirch 1994a, 1994b, Weimerskirch et al. 1994, Chastel et al. 1995). Therefore, high

food availability at sea may allow a good foraging success of the foraging partner that may return

to land after a short stay at sea, allowing a good synchronization of the breeding partners on the

nest. In contrast, poor conditions could increase the time spent at sea by the foraging partner,

which would increase desertion of the nest by the fasting partner and then, reduce the breeding

success. We thus predicted that conditions at sea during the breeding season would also affect

the breeding success of petrels.

Southern Annular Mode

The SAM is a large-scale climate index. SAM is the leading mode of climate variability over the

Southern Hemisphere. SAM is defined as the difference of atmospheric pressure between the

40°S and 65°S latitudes (Marshall 2003). SAM influences surface wind, sea surface temperature

(SST) and surface chlorophyll concentration. A large majority of the skuas from Mayes Island

overwinter north of the polar front (Delord et al. 2018). In the subtropical zone, SAM positive

phases induced warm SSTa, low surface chlorophyll concentration and easterly winds driving

Ekman transport (the 90° wind-driven net transport on the sea surface), while in the Subantarctic

zone there is a convergence of waters that increase downwelling and positive SSTa (Lovenduski

and Gruber 2005). We thus predicted that the positive phases of SAM, potentially leading to

poorer food availability in the areas used by skuas during the nonbreeding period, would have

negative impacts on skua survival and limit their ability to breed the next breeding season. South

of the polar front, where petrels spend the winter, positive phases of the SAM are associated with

westerly winds. This induces cold SSTa, increased equatorward Ekman transport and drives

increased upwelling (Lovenduski and Gruber 2005). Consequently, the biological productivity

and potential prey availability for petrels are higher during positive phases of the SAM. We thus

predicted that the positive phases of SAM would be favorable for petrel demographic

parameters. Data were obtained from the online database of the British Antarctic Survey

(http://www.nerc-bas.ac.uk/ icd/gjma/sam.html).

Sea Surface Temperature anomalies

SSTa reflect local oceanographic conditions that influence the whole marine trophic food web.

High SST generally reduces vertical mixing and provides poor growing conditions for

zooplankton communities which, through bottom-up mechanisms, induces reduced trophic

resources for seabirds (Barbraud et al. 2012, Sydeman et al. 2015). Consequently, year-to-year

variation of SST was previously found to be negatively correlated with petrel body condition

(Guinet et al. 1998). Therefore, we predicted that high SSTa would negatively affect overwinter

survival and breeding success of petrels. The SSTa data were downloaded from the National

Oceanic and Atmospheric Administration ("data: NOAA NCEP EMC CMB GLOBAL

Reyn_SmithOIv2 monthly ssta") from 1996 to 2018.

Chlorophyll a

Chlorophyll a lies at the bottom of the marine food web and provides resources for higher trophic

organisms up to seabirds. Because petrel diet is mainly composed of crustaceans and fish feeding

at low trophic levels (Cherel et al. 2002, 2014), we predicted that high concentrations of Chla

would be favorable to the survival and breeding success of petrels. The Chla data were

downloaded from the NASA Ocean Data with a 9km mapped concentration data of chlorophyll a

for the years 1997 to 2001 and from the Nasa Earth Observation (NEO AQUA/MODIS data)

monthly for the years 2002 to 2018.

Assessing the effect of environmental covariates and population densities

We fitted a single multispecies IPM including all the biologically relevant effects. Logit-linear

regressions were used to estimate the effect of environmental (SAM, SSTa and Chla) and inter-

and intra-specific interactions on demographic parameters (adult apparent survival, breeding

probability, hatching probability, breeding success) (Table 2). We used state variables

$N_{adtot,S}$ and $N_{adtot,P}$, respectively the number of adult skuas and petrels, to assess the effects of

inter- and intra-specific interactions. For example, we modelled the hatching probability for

petrels that bred the previous year ($\omega_{P,\tilde{B}}$) using a logit link:

$logit(\omega_{P,\tilde{B},t-1}) =$

$\mu_{\omega,P,\tilde{B}} + \alpha_{SAM,\omega,P,\tilde{B}} \times SAM_{\omega,P,t} + \alpha_{SSTa,\omega,P,\tilde{B}} \times SSTa_{\omega,P,t} + \alpha_{Chla,\omega,P,\tilde{B}} \times Chla_{\omega,P,t} +$

$\alpha_{DD,\omega,P,\tilde{B}} \times N_{adtot,P,t} + \alpha_{PP,\omega,P,\tilde{B}} \times N_{adtot,S,t} + \varepsilon_{\omega,P,\tilde{B},t}$

$\varepsilon_{\omega,P,\tilde{B},t} \sim N(0, \sigma^2{}_{\varepsilon,\omega,P,\tilde{B}})$ \hfill (25)

with $\mu_{\omega,P,\tilde{B}}$ the intercept, $\alpha_{SAM_{\omega,P,\tilde{B}}}$ the slope for the climatic covariate $SAM_{\omega,P}$, $\alpha_{SSTa,\omega,P,\tilde{B}}$ the

slope for the climatic covariate $SSTa_{\omega,P}$, $\alpha_{Chla,\omega,P,\tilde{B}}$ the slope for the climatic covariate $Chla_{\omega,P}$,

$\alpha_{DD,\omega,P,\tilde{B}}$ the slope indicating the strength of the intraspecific density-dependence with

$N_{adtot,P}$ the number of adult petrels, $\alpha_{PP,\omega,P,\tilde{B}}$ the slope indicating the strength of the predator-

prey relationship with $N_{adtot,S}$ the number of adult skuas, $\varepsilon_{\omega,P,\tilde{B}}$ is a yearly random effect and

$\sigma^2_{\varepsilon,\omega,P,\tilde{B}}$ its temporal variance. The descriptions of all logit-linear relationships used on

demographic parameters are available in Appendix S2.

For local covariates (SSTa and Chla), we calculated the average values of the covariates in the

areas in which petrels were located (Cherel et al. 2016) in a specific time period during which the

environment might affect the demographic parameter under investigation (Table 2). Each

environmental covariate was standardized to have zero mean and unit variance. However, the

inter- and intra-specific covariates were not standardized prior to the analyses because the

population sizes were estimated step by step each year. To compare the relative contribution of

the effects of each covariate, we calculated the standardized effect of population size (for inter-

and intra-specific relationship) posterior to the analyses by multiplying their slopes (α) by the

standard deviation of the estimated population sizes. Then, we compared the relative contribution

of each covariate using the regression estimate which we used as a measure of effect size.

We computed the 95% and 80% credible intervals (CRI) for the regression coefficients α. We

did not interpret uncertain effects (*i.e.* 80% CRI including zero) and focused particularly on clear

effects whose sign could be reliably assessed (*i.e.* 95% CRI excluding zero).

Model implementation

To fit the juvenile apparent survival parameters increasing with age, we modelled them as a positive linear function of age by assigning to the slope λ_2 a $U(0,1)$ prior, and by defining the intercept λ_1 with a normal $N(0,1)$ prior. The probability of the first breeding attempt (Pr) is time-dependent with a uniform prior: $Pr_t \sim U(0,1)$. Normal priors $N(0,10^4)$ were assigned to the regression coefficients (α) of the covariate effects. For the variance of the random year effects (σ_ε^2), we used a $U(0,10)$ vague prior. The state assignment probability of individuals with uncertain state parameter (u) was defined *a priori* with a $U(0,10)$ vague prior.

Bayesian posterior distributions were approximated *via* Markov chain Monte Carlo (MCMC) algorithms. Two independent MCMC chains of 200,000 iterations were used with a burn-in period of 100,000. One out of five iterations was kept and final inferences were derived from a sample of $2 \times 20,000$ iterations that resulted from merging the two chains. Gelman-Rubin convergence diagnostic (Brooks and Gelman 1998) was below 1.5 for each parameter and the mixing of the chains was satisfactory. We performed the analyses using Nimble (de Valpine et al. 2017 ; version 0.9.1) and program R (R Core Team 2020 ; R version 4.0.3). Code and data are available on GitHub at https://github.com/maudqueroue/MultispeciesIPM_SkuaPetrel.

Results

Predator-prey relationships

We estimated positive relationships between two breeding parameters of skuas and the number of adult petrels. The breeding success for at least one chick ($\gamma_{S,\tilde{B}}$) [slope mean ($\alpha_{PP_{\gamma,S,\tilde{B}}}$) = 0.66;

95% CRI (0.37, 1.05)] (Fig. 2a) and the breeding success with two chicks ($\delta_{S,\tilde{B}}$) [slope mean

($\alpha_{PP_{\delta,S,\tilde{B}}}$) = 1.24; 95% CRI (0.59, 2.12)] (Fig. 2b) for skuas that were breeders the previous

breeding season increased with an increasing number of prey. Even though the effects were less

clear (95% CRI including zero), the breeding success of petrels tended to be positively impacted

by the number of predators (Table 3). We detected a positive relationship between the number of

adult skuas and the hatching success of petrels that were breeders the previous breeding

season ($\omega_{P,\tilde{B}}$), and with the breeding success of petrels that were nonbreeders the previous

breeding season ($\gamma_{P,\widetilde{NB}}$). We found no other interspecific relationship on the other parameters

(Table 3).

Intraspecific density-dependence

The number of skuas had a clear effect on two demographic parameters, namely the breeding

success and the breeding success with two chicks for skuas that were breeders the previous

breeding season. We found negative density-dependence for the breeding success ($\gamma_{S,\tilde{B}}$) [slope

mean ($\alpha_{DD_{\gamma,S,\tilde{B}}}$) = -0.39; 95% CRI (-0.65, -0.12)] (Fig. 2c) and for the probability of producing

two chicks rather than one ($\delta_{S,\tilde{B}}$) [slope mean ($\alpha_{DD_{\delta,S,\tilde{B}}}$) = -0.53; 95% CRI (-1.02, -0.10)] (Fig.

2d). These two breeding parameters were also affected by interspecific relationships and we

observed that the predator-prey effects were stronger than intraspecific effects ($\left|\text{mean } \alpha_{PP_{\gamma,S,\tilde{B}}}\right|$ =

0.66; $\left|\text{mean } \alpha_{DD_{\gamma,S,\tilde{B}}}\right|$ = 0.39 respectively) for the breeding success and ($\left|\text{mean } \alpha_{PP_{\delta,S,\tilde{B}}}\right|$ = 1.24;

$\left|\text{mean } \alpha_{DD_{\delta,S,\tilde{B}}}\right|$ = 0.53 respectively) for the breeding success with two chicks (Table 3).

For petrels, we estimated a positive effect of increased number of adult petrels on the breeding

probability for individuals that were breeders the previous breeding season $(\beta_{P,\tilde{B}})$ [slope

mean $(\alpha_{DD_{\beta,P,\tilde{B}}}) = 0.51$; 95% CRI (0.03, 0.98)] (Fig. 2e). Moreover, the number of petrels tended

to negatively affect the breeding success of petrels that did not bred the previous breeding

season $(\gamma_{P,\widetilde{NB}})$ (Table 3).

Environmental covariates

We found ecologically relevant relationships between environmental covariates and

demographic parameters of the two species (Table 3). For petrels, we found positive

relationships between the two local environmental covariates (SSTa and Chla) and the breeding

probability for individuals that were nonbreeders the previous breeding season $(\beta_{P,\widetilde{NB}})$. The

effect of these environmental covariates on the breeding probability was stronger for the Chla

covariate than for the SSTa covariate ($\left| \text{mean } \alpha_{Chla_{\beta,P,\widetilde{NB}}} \right| = 0.63$; $\left| \text{mean } \alpha_{SST_{\beta,P,\widetilde{NB}}} \right| = 0.36$

respectively). We estimated a positive relationship between the SAM covariate and the apparent

survival of skuas that were nonbreeders the previous breeding season $(\phi_{S,\widetilde{NB}})$.

In addition to the results above, we also estimated the demographic parameters and the number

of individuals in each state for both species from 1996 to 2017 (see Appendix S3: Figs. S1– S6).

Discussion

In this paper, we provide the first application of a multispecies IPM in a predator-prey context.

Joint analysis of empirical data for two seabird species allowed us to estimate demographic

parameters and population size for both simultaneously. The key advantage of using a

multispecies IPM was that it enabled us to use the population sizes estimated by the model for

one species to analyze its effect on the demographic parameters of the other species while

propagating all sources of uncertainty. Hence, it allowed us to understand the contribution of

interspecific interactions on the demographic parameters while further taking into account the

effects of climatic conditions. Our results showed that the demography of the predatory skua was

mainly driven by the number of petrel prey during the breeding season and by the environment

during the nonbreeding season, whereas petrels were mostly impacted by the environment. This

suggests that this predator-prey system is mainly driven by bottom-up processes and density-

dependent processes.

Effects of predator-prey relationships

The number of prey is a determining factor in the breeding success of skuas according to our

results. Food availability is known to be positively related with breeding parameters in seabirds

(Cairns 1988, Piatt et al. 2007, Oro et al. 2014). As diet of skuas during the breeding period is

dominated by petrels (Mougeot et al. 1998, Pacoureau et al. 2019c), a large abundance of petrels

provides easier conditions for skuas to feed themselves and their chicks resulting in a higher

breeding success.

Interestingly, we did not find the opposite relationship in the prey dynamics. Our model provided

no evidence for a negative effect of the number of skuas on the demographic parameters of the

petrel. As skuas prey on both adults and juveniles during the breeding season, we expected a

negative effect of the number of skuas on the petrel breeding parameters. This lack of effect

could be explained by the large abundance of petrels compared to the skuas on Mayes Island.

Oro et al. (2006) reported that in another seabird predator-prey system, the highest breeding

success of the prey occurred when the prey/predator ratio was very high. On Mayes Island, the

breeding population of petrels is estimated at approximately 142,000 breeding pairs (Barbraud

and Delord 2006), and this does not include chicks (around 71,000 each year) and nonbreeders

(approximately 30% of the petrels). Hence, there are about 476,000 petrels during a breeding

season versus about 200 skuas, resulting in a very high prey/predator ratio. Moreover, Mougeot

et al. (1998) showed that skuas breeding at Mayes Island preyed on about 40,000 petrels each

breeding season. This corresponds to about 8% of the petrel population of the island. It is

therefore possible that skua predation is only a minor factor in shaping petrel demographics, and

this effect may be too weak to be detected by our model. Inversely, although the relationships

estimated were less clear, our results suggest that the density of skuas tended to increase slightly

with the hatching success and breeding success of the prey. However, it is unlikely that the

presence of predators increased the reproductive success of petrels. To explain these

relationships, we might rely on the other strong effects estimated by our model. Indeed, we found

that the number of petrels positively affected the breeding success of skuas and that skuas were

sensitive to intraspecific density-dependence. Therefore, years when preys experience a high

breeding success correspond to years with particularly abundant food resources for skuas and

this until the end of the breeding season. Since skuas are potentially less affected by intraspecific

density-dependence than by abundance of prey, they could consequently breed in higher

numbers in the breeding area.

Effects of intraspecific density-dependence

For skuas, we found negative density-dependent effects on breeding success and probability to fledge two chicks, in accordance with our predictions. Egg and chick predation by conspecifics has been reported in the Great Skua (*Catharacta skua*) (Hamer et al. 1991, Ratcliffe and Furness 1999). Hence, a higher abundance of skuas increases the risk of predation on eggs and chicks, resulting in higher breeding failure. To avoid predation by conspecifics, the skuas start defending their territories from conspecifics just a few days after arrival on a breeding site until the end of the season. This activity is energetically costly and may also limit the time spent searching for food, potentially limiting energy investment in reproduction. The heterogeneous habitat hypothesis already demonstrated in territorial birds (Dhondt et al. 1992, Krüger and Lindström 2001, Ferrer and Donazar 2015) could also explain the relationships we found. Indeed, when the population increases, some individuals may be forced to occupy poorer quality habitats, resulting in lower reproductive success. We did not find an effect of density-dependence on the breeding probability of skuas. As skuas are territorial with high site fidelity, we hypothesized that in years with a high abundance of skuas, the breeding probability would decrease, as all the skuas would not succeed in acquiring a territory. It is possible that we did not observe this effect because the logistic function used for density-dependence does not accurately model the territory acquisition dynamics by floaters (e.g. van de Pol et al. 2010, Barraquand et al. 2014).

We estimated that the breeding success of skuas was affected by both predator-prey relationships and intraspecific density-dependence. Predator-prey relationships had a higher contribution to the variability in breeding success of skuas than the density-dependent effect. Hamer et al. (1991) reported that, following a reduction of sandeel (*Ammodytes marinus*) abundance, great

skua increased their foraging effort reducing the adult territorial attendance. In turn, breeding

failure increased due to predation from adults of neighboring territories. We then may assume

that petrel abundance allowed a suitable territorial attendance for skuas reducing the negative

density-dependent effects such as chick predation by conspecifics.

For petrels, we found a negative relationship between the breeding success of petrel that did not

breed the previous year and the number of petrels on the colony. Combined effects of density-

dependence and climate have already been observed in petrels, with a lower winter survival

when density is high (Barbraud and Weimerskirch 2003), suggesting a mechanism of

competition between conspecifics for food resources. As nonbreeders are known to be in poorer

condition than breeders (Chastel et al. 1995), they are potentially more sensitive to the

competition for food resources explaining why this effect was only found on petrels that were

nonbreeders the previous years. Interestingly, we found a positive intraspecific density-

dependence relationship on the breeding probability of petrels that bred the previous year. This

suggests that years with a high abundance of petrels reflected a good return rate to the breeding

site because environmental conditions were favorable for breeding. This is in agreement with

studies showing that petrels might skip breeding and take sabbatical years when environmental

conditions are poor (Warham 1990, Chastel et al. 1995).

Effects of environmental conditions

Breeding probability of petrels tended to be impacted by two of the environmental covariates

tested, namely SSTa and Chla. This effect of environmental conditions on the breeding

probability is in accordance with previous research showing that the body condition of petrels

might impact their decision to attempt breeding (Warham 1990, Chastel et al. 1995). High Chla

increases resources availability for organisms at higher trophic levels (macrozooplankton,

fishes), which are consumed by petrels (Cherel et al. 2002). Consequently, high Chla may

increase abundance of petrel prey, with a positive effect on the breeding performances and body

condition of petrels. Unexpectedly, we detected a positive effect of SSTa on breeding probability

of petrels. This result is surprising as previous study showed that warm SST events negatively

affected the breeding performances and body condition of petrels at Kerguelen Islands (Guinet et

al. 1998). Indeed, high SST generally reduces vertical mixing and provides poor growing

conditions for zooplankton communities that in turn reduce trophic resources for seabirds

(Barbraud et al. 2012, Sydeman et al. 2015). However, it has been showed recently that during

the pre-laying period petrels use water masses situated at more northerly latitudes than during the

winter period or the breeding period (Quillfeldt et al. 2020), where relationships between SST

and primary productivity may differ. Indeed, the covariance between SST and Chla depends on

location and shows particularly complex patterns in the Southern Ocean (Dunstan et al. 2018).

Positive effects of SSTa have already been identified in other sub-antarctic seabirds (Pinaud and

Weimerskirch 2002, Nevoux et al. 2007, Horswill et al. 2014). Furthermore, we estimated that

Chla, at the bottom of the trophic food chain, had a higher effect on the breeding probability than

SSTa which reflect oceanographic conditions. This indicated that the effect size of

environmental covariates increased when the covariates approached the trophic level occupied

by the prey of the petrels, suggesting a bottom-up mechanism. This result is consistent with

many studies showing that climatic conditions affect seabirds through indirect processes by

influencing prey availability and resulting in changes in their dynamics (Frederiksen et al. 2006,

Barbraud et al. 2012, Jenouvrier 2013, Lauria et al. 2013).

We did not detect any relationship between the breeding parameters of the skua and the

environmental covariates. This lack of effect could be explained by an absence of a direct link

between skuas and the environmental covariates tested, as breeding skuas remain on their

territory to defend it or to forage. However, we found an effect of SAM on the apparent survival.

This effect was detected only for skuas that were nonbreeders during the previous season. It was

proposed that only seabirds attaining a threshold condition decide to breed (Weimerskirch 1992),

suggesting that nonbreeders are generally in poorer conditions than breeders (Chastel et al. 1995,

Cam et al. 1998) and thus more sensitive to environmental conditions. Nevertheless, we found a

positive relationship between survival and SAM whereas we expected a negative relationship.

Indeed, skuas mainly overwinter north of the polar front (Delord et al. 2018) where positive

phases of SAM induce warm SST, low surface Chla concentration (Lovenduski and Gruber

2005), and thus potentially poor feeding conditions. However, only breeding skuas were studied

in Delord et al. (2018) and nonbreeding individuals may use different wintering areas where the

relationships between SAM and oceanographic variables differ. Several studies reviewed in

Jenouvrier (2013) highlighted multifaceted effects of climatic conditions on the demography of

seabirds involving direct, time-lagged and non-linear effect, which we did not considered here.

Therefore, despite the important contribution of our approach in understanding the effect of the

environment in our predator-prey system, disentangling in details the complex mechanisms

between environmental covariates and their effects on the demography of the two seabirds

remain challenging.

A bottom-up dynamic in a predator-prey system

Overall, our study has highlighted the important role of bottom-up processes in the dynamics of this marine predator-prey system, *i.e.* the dynamics of these two seabirds was mostly driven by food availability. Petrel dynamics were more strongly affected by environmental covariates near to their trophic level and the number of petrels impacted the dynamics of skuas. The bottom-up control of demographic rates in oceanic predators have been largely assumed (Jenouvrier 2013). This is because the functioning of oceanic systems is controlled and structured by physical processes impacting nutrient fluxes (Behrenfeld et al. 2006) and then the whole trophic food web. We found no evidence of top-down processes, *i.e.* predation effects, in this system, although these two mechanisms have been found to jointly affect ecosystems (Hunter and Price 1992, Sinclair et al. 2003) including other seabird systems (Horswill et al. 2014, 2016, Perkins et al. 2018). Effects of skua predation on petrels were expected, based on their diet during the breeding season. However, given the very large number of petrels present on the island compared to the number of predators, the impact of predation may have been too small to be detected by our model.

Conclusion

This multispecies IPM framework allowed us to estimate demographic parameters and abundances for both skuas and petrels. Taking into account both species interactions and environmental covariates in the same analysis improved our understanding of species dynamics. We concluded that bottom-up mechanisms are the main drivers of this skua-petrel system. Generalizing such assessments of interspecific relationships and environmental conditions in a

690 single demographic framework may be essential to predict how contrasted climatic scenarios will

691 affect communities. A promising avenue of research in multispecies IPMs lies in fitting models

692 to data on a large number of species, which will much likely require further methodological

693 developments.

Acknowledgments

This study was made possible thanks to all the field workers involved in the monitoring

programs on Brown Skuas and Blue Petrels since 1985 at Mayes Island, Kerguelen Islands.

These monitoring programs were supported financially and logistically by the French Polar

Institute IPEV (program 109, resp. Henri Weimerskirch), the Zone Atelier Antarctique (CNRS-

INEE), Terres Australes et Antarctiques Françaises. All work was carried out in accordance with

the guidelines of the IPEV ethics committee. We thank Chloé R. Nater for constructive feedback

and helpful suggestions on the manuscript. We acknowledge Dominique Joubert for the

management of the demographic database. We thank Dave Koons and an anonymous reviewer

for useful comments that helped improved a previous version of the manuscript. This research

was funded by the French National Research Agency (grant ANR-16-CE02-0007).

Literature Cited

Aebischer, N. J., J. C. Coulson, and J. M. Colebrookl. 1990. Parallel long-term trends across four

marine trophic levels and weather. Nature 347:753–755.

Barbraud, C., and K. Delord. 2006. Population census of blue petrels *Halobaena caerulea* at

Mayes Island, Iles Kerguelen. Antarctic Science 18:199–204.

Barbraud, C., V. Rolland, S. Jenouvrier, M. Nevoux, K. Delord, and H. Weimerskirch. 2012.

Effects of climate change and fisheries bycatch on Southern Ocean seabirds: a review.

Marine Ecology Progress Series 454:285–307.

Barbraud, C., and H. Weimerskirch. 2001. Emperor penguins and climate change. Nature

411:183–186.

Barbraud, C., and H. Weimerskirch. 2003. Climate and density shape population dynamics of a

marine top predator. Proceedings of the Royal Society B: Biological Sciences 270:2111–

2116.

Barbraud, C., and H. Weimerskirch. 2005. Environmental Conditions and Breeding Experience

Affect Costs of Reproduction in Blue Petrels. Ecology 86:682–692.

Barraquand, F., and O. Gimenez. 2019. Integrating multiple data sources to fit matrix population

models for interacting species. Ecological Modelling 411.

Barraquand, F., T. T. Høye, J.-A. Henden, N. G. Yoccoz, O. Gilg, N. M. Schmidt, B. Sittler, and

R. A. Ims. 2014. Demographic responses of a site-faithful and territorial predator to its

fluctuating prey: long-tailed skuas and arctic lemmings. Journal of Animal Ecology 83:375–

387.

Behrenfeld, M. J., R. T. O'Malley, D. A. Siegel, C. R. McClain, J. L. Sarmiento, G. C. Feldman,

A. J. Milligan, P. G. Falkowski, R. M. Letelier, and E. S. Boss. 2006. Climate-driven trends

in contemporary ocean productivity. Nature 444:752–755.

Besbeas, P., S. N. Freeman, B. J. T. Morgan, and E. A. Catchpole. 2002. Integrating Mark-

Recapture-Recovery and Census Data to Estimate Animal Abundance and Demographic

Parameters. Biometrics 58:540–547.

Bogdanova, M. I., A. Butler, S. Wanless, B. Moe, T. Anker-Nilssen, M. Frederiksen, T.

Boulinier, L. S. Chivers, S. Christensen-Dalsgaard, S. Descamps, M. P. Harris, M. Newell,

B. Olsen, R. A. Phillips, D. Shaw, H. Steen, H. Strøm, T. L. Thórarinsson, and F. Daunt.

2017. Multi-colony tracking reveals spatio-Temporal variation in carry-over effects between

breeding success and winter movements in a pelagic seabird. Marine Ecology Progress

Series 578:167–181.

Brooks, S. P., and A. Gelman. 1998. General Methods for Monitoring Convergence of Iterative

Simulations. Journal of Computational and Graphical Statistics 7:434–455.

Cairns, D. K. 1988. Seabirds as Indicators of Marine Food Supplies. Biological Ocenaography

5:261–271.

Cam, E., J. E. Hines, J.-Y. Monnat, J. D. Nichols, and E. Danchin. 1998. Are Adult Nonbreeders

Prudent Parents? The Kittiwake Model. Ecology 79:2917.

Carneiro, A. P. B., A. Manica, W. Z. Trivelpiece, and R. A. Phillips. 2015. Flexibility in foraging

strategies of Brown Skuas in response to local and seasonal dietary constraints. Journal of

Ornithology 156:625–633.

Caswell, H. 2001. Matrix Population Models: Construction, Analysis, and Interpretation. Sinauer

Associates, Sunderland, Massachusetts, USA.

Chastel, O., H. Weimerskirch, and P. Jouventin. 1995. Influence of Body Condition on

Reproductive Decision and Reproductive Success in the Blue Petrel. The Auk 112:964–972.

Chaurand, T., and H. Weimerskirch. 1994a. The regular alternation of short and long foraging

trips in the blue petrel *Halobaena caerulea*: a previously undescribed strategy of food

provisioning in a pelagic seabird. The Journal of Animal Ecology 63:275–282.

Chaurand, T., and H. Weimerskirch. 1994b. Incubation routine, body mass regulation and egg

neglect in the Blue Petrel *Halobaena caerulea*. Ibis 136:285–290.

Cherel, Y., P. Bocher, C. Trouvé, and H. Weimerskirch. 2002. Diet and feeding ecology of blue

petrels *Halobaena caerulea* at Iles Kerguelen, Southern Indian Ocean. Marine Ecology

Progress Series 228:283–299.

Cherel, Y., M. Connan, A. Jaeger, and P. Richard. 2014. Seabird year-round and historical

feeding ecology: blood and feather δ13C and 1315N values document foraging plasticity of

small sympatric petrels. Marine Ecology Progress Series 505:267–280.

Cherel, Y., P. Quillfeldt, K. Delord, and H. Weimerskirch. 2016. Combination of At-Sea

Activity, Geolocation and Feather Stable Isotopes Documents Where and When Seabirds

Molt. Frontiers in Ecology and Evolution 4.

Chu, C., and P. B. Adler. 2015. Large niche differences emerge at the recruitment stage to

stabilize grassland coexistence. Ecological Monographs 85:373–392.

Daunt, F., V. Afanasyev, A. Adam, J. P. Croxall, and S. Wanless. 2007. From cradle to early

grave: Juvenile mortality in European shags *Phalacrocorax aristotelis* results from

inadequate development of foraging proficiency. Biology Letters 3:371–374.

Delord, K., Y. Cherel, C. Barbraud, O. Chastel, and H. Weimerskirch. 2018. High variability in

migration and wintering strategies of brown skuas (*Catharacta antarctica lonnbergi*) in the

Indian Ocean. Polar Biology 41:59–70.

Dhondt, A. A., B. Kempenaers, and F. Adriaensen. 1992. Density-dependent clutch size caused

by habitat heterogeneity. Journal of Animal Ecology 61:643–648.

Dunstan, P. K., S. D. Foster, E. King, J. Risbey, T. J. O'Kane, D. Monselesan, A. J. Hobday, J.

R. Hartog, and P. A. Thompson. 2018. Global patterns of change and variation in sea

surface temperature and chlorophyll a. Scientific Reports 8:1–9.

Fay, R., H. Weimerskirch, K. Delord, and C. Barbraud. 2015. Population density and climate

shape early-life survival and recruitment in a long-lived pelagic seabird. Journal of Animal

Ecology 84:1423–1433.

Ferrer, M., and J. A. Donazar. 2015. Density-Dependent Fecundity by Habitat Heterogeneity in

an Increasing Population of Spanish Imperial Eagles. Ecology 77:69–74.

Frederiksen, M., M. Edwards, A. J. Richardson, N. C. Halliday, and S. Wanless. 2006. From

plankton to top predators: bottom-up control of a marine food web across four trophic

levels. Journal of Animal Ecology 75:1259–1268.

Gervasi, V., E. B. Nilsen, and J. D. C. Linnell. 2015. Body mass relationships affect the age

structure of predation across carnivore-ungulate systems: A review and synthesis. Mammal

Review 45:253–266.

Gille, S. T. 2002. Warming of the Southern Ocean since the 1950s. Science 295:1275–1277.

Gimenez, O., J. D. Lebreton, J. M. Gaillard, R. Choquet, and R. Pradel. 2012. Estimating

demographic parameters using hidden process dynamic models. Theoretical Population

Biology 82:307–316.

Godfray, H. C. J., and R. M. May. 2014. Open questions: are the dynamics of ecological

communities predictable? BMC Biology 12.

Grande, J. M., D. Serrano, G. Tavecchia, M. Carrete, O. Ceballos, R. Díaz-Delgado, J. L. Tella,

and J. A. Donázar. 2009. Survival in a long-lived territorial migrant: effects of life-history

traits and ecological conditions in wintering and breeding areas. Oikos 118:580–590.

Greig, S. A., J. C. Coulson, and P. Monaghan. 1983. Age-Related differences in foraging success

in the herring gull (*Larus Argentatus*). Anim. Behav. 31:1237–1243.

Grosbois, V., O. Gimenez, J.-M. Gaillard, R. Pradel, C. Barbraud, J. Clobert, A. P. Møller, and

H. Weimerskirch. 2008. Assessing the impact of climate variation on survival in vertebrate

populations. Biological Reviews 83:357–399.

Guinet, C., O. Chastel, M. Koudil, J. P. Durbec, and P. Jouventin. 1998. Effects of warm sea-

surface temperature anomalies on the blue petrel at the Kerguelen Islands. Proceedings of

the Royal Society B: Biological Sciences 265:1001–1006.

Hamer, K. C., R. W. Furness, and R. W. G. Caldow. 1991. The effects of changes in food

availability on the breeding ecology of great skuas *Catharacta skua* in Shetland. Journal of

Zoology 223:175–188.

Han, W., J. Vialard, M. J. McPhaden, T. Lee, Y. Masumoto, M. Feng, and W. P. M. de Ruijter.

2014. Indian Ocean Decadal Variability: A Review. Bull. Amer. Meteor. Soc. 95:1679–

1703.

Harrison, X. A., J. D. Blount, R. Inger, D. R. Norris, and S. Bearhop. 2011. Carry-over effects as

drivers of fitness differences in animals. Journal of Animal Ecology 80:4–18.

Higgins, P. J., and S. J. Davied. 1996. Handbook of Australia, New Zealand & Antarctic Birds.

Vol. 3, Snipe to pigeons. Oxford University Press.

Hipfner, M. J., L. K. Blight, R. W. Lowe, S. I. Wilhelm, G. J. Robertson, R. T. Barrett, T. Anker-

Nilssen, and T. P. Good. 2012. Unintended consequences: How the recovery of sea eagle

Haliaeetus spp. populations in the northern hemisphere is affecting seabirds. Marine

Ornithology 40:39–52.

Hoegh-Guldberg, O., and J. F. Bruno. 2010. The Impact of Climate Change on the World's

Marine Ecosystems. Science 328:1523–1528.

Horswill, C., J. Matthiopoulos, J. A. Green, M. P. Meredith, J. Forcada, H. Peat, M. Preston, P.

N. Trathan, and N. Ratcliffe. 2014. Survival in macaroni penguins and the relative

importance of different drivers: Individual traits, predation pressure and environmental

variability. Journal of Animal Ecology 83:1057–1067.

Horswill, C., N. Ratcliffe, J. A. Green, R. A. Phillips, P. N. Trathan, and J. Matthiopoulos. 2016.

Unravelling the relative roles of top-down and bottom-up forces driving population change

in an oceanic predator. Ecology 97:1919–1928.

Hunter, M. D., and P. W. Price. 1992. Playing chutes and ladders: heterogeneity and the relative

roles of bottom-up and top-down forces in natural communities. Ecology 73:724–732.

Jenouvrier, S. 2013. Impacts of climate change on avian populations. Global Change Biology
19:2036–2057.

Jenouvrier, S., C. Barbraud, and H. Weimerskirch. 2003. Effects of climate variability on the
temporal population dynamics of southern fulmars. Journal of Animal Ecology 72:576–587.

Jouventin, P., J. L. Mougin, J.-C. Stahl, and H. Weimerskirch. 1985. Comparative biology of the
burrowing petrels of the Crozet Islands. Notornis 32:157–220.

Krüger, O., and J. Lindström. 2001. Habitat heterogeneity affects population growth in goshawk
Accipiter gentilis. Journal of Animal Ecology 70:173–181.

Lack, D. 1967. Interrelationships in breeding adaptations as shown by marine birds. In
Proceedings of the International Ornithological Congress. Vol. 14, pp. 3-42. Oxford. UK:
Blackwell Scientific Publications.

Lauria, V., M. J. Attrill, A. Brown, M. Edwards, and S. C. Votier. 2013. Regional variation in
the impact of climate change: Evidence that bottom-up regulation from plankton to seabirds
is weak in parts of the Northeast Atlantic. Marine Ecology Progress Series 488:11–22.

Lovenduski, N. S., and N. Gruber. 2005. Impact of the Southern Annular Mode on Southern
Ocean circulation and biology. Geophysical Research Letters 32.

Marshall, G. J. 2003. Trends in the Southern Annular Mode from Observations and Reanalyses.
Journal of climate 16:4134–4143.

Miller, D. A. W., E. H. C. Grant, E. Muths, S. M. Amburgey, M. J. Adams, M. B. Joseph, J. H.

Waddle, P. T. J. Johnson, M. E. Ryan, B. R. Schmidt, D. L. Calhoun, C. L. Davis, R. N. Fisher, D. M. Green, B. R. Hossack, T. A. G. Rittenhouse, S. C. Walls, L. L. Bailey, S. S. Cruickshank, G. M. Fellers, T. A. Gorman, C. A. Haas, W. Hughson, D. S. Pilliod, S. J. Price, A. M. Ray, W. Sadinski, D. Saenz, W. J. Barichivich, A. Brand, C. S. Brehme, R. Dagit, K. S. Delaney, B. M. Glorioso, L. B. Kats, P. M. Kleeman, C. A. Pearl, C. J. Rochester, S. P. D. Riley, M. Roth, and B. H. Sigafus. 2018. Quantifying climate sensitivity and climate-driven change in North American amphibian communities. Nature Communications 9.

Miller, T. E. X., and V. H. W. Rudolf. 2011. Thinking inside the box◻: community-level consequences of stage-structured populations. Trends in Ecology and Evolution 26:457–466.

Millon, A., S. J. Petty, B. Little, O. Gimenez, T. Cornulier, and X. Lambin. 2014. Dampening prey cycle overrides the impact of climate change on predator population dynamics: a long-term demographic study on tawny owls. Global Change Biology 20:1770–1781.

Mougeot, F., and V. Bretagnolle. 2000a. Predation risk and moonlight avoidance in nocturnal seabirds. Journal of Avian Biology 31:376–386.

Mougeot, F., and V. Bretagnolle. 2000b. Predation as a cost of sexual communication in nocturnal seabirds: an experimental approach using acoustic signals. Animal Behaviour 60:647–656.

Mougeot, F., F. Genevois, and V. Bretagnolle. 1998. Predation on burrowing petrels by the brown skua (*Catharacta skua lönnbergi*) at Mayes Island, Kerguelen. Journal of Zoology

244:429–438.

Nevoux, M., H. Weimerskirch, and C. Barbraud. 2007. Environmental variation and experience-

related differences in the demography of the long-lived black-browed albatross. Journal of

Animal Ecology 76:159–167.

Oken, K. L., and T. E. Essington. 2015. How detectable is predation in stage-structured

populations? Insights from a simulation-testing analysis. Journal of Animal Ecology 84:60–

70.

Oro, D., N. Hernández, L. Jover, and M. Genovart. 2014. From recruitment to senescence: food

shapes the age-dependent pattern of breeding performance in a long-lived bird. Ecology

95:446–457.

Oro, D., A. Martínez-Abraín, M. Paracuellos, J. C. Nevado, and M. Genovart. 2006. Influence of

density dependence on predator-prey seabird interactions at large spatio-temporal scales.

Proceedings of the Royal Society B: Biological Sciences 273:379–383.

Pacoureau, N., M. Authier, K. Delord, and C. Barbraud. 2019a. Population response of an apex

Antarctic consumer to its prey and climate fluctuations. Oecologia 189:279–291.

Pacoureau, N., K. Delord, S. Jenouvrier, and C. Barbraud. 2019b. Demographic and population

responses of an apex predator to climate and its prey: a long☐term study of South Polar

Skuas. Ecological Monographs 89.

Pacoureau, N., E. Gaget, K. Delord, and C. Barbraud. 2019c. Prey remains of brown skua is

evidence of the long-term decline in burrow occupancy of blue petrels and thin-billed prions

at Mayes Island, Kerguelen. Polar Biology 42:1873–1879.

Parmelee, D. F., and P. J. Pietz. 1987. Philopatry, Mate and Nest-Site Fidelity in the Brown

Skuas of Anvers Island, Antartica. The Condor 89:916–919.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual

Review of Ecology, Evolution, and Systematics 37:637–669.

Perkins, A., N. Ratcliffe, D. Suddaby, B. Ribbands, C. Smith, P. Ellis, E. Meek, and M. Bolton.

2018. Combined bottom-up and top-down pressures drive catastrophic population declines

of Arctic skuas in Scotland. Journal of Animal Ecology 87:1573–1586.

Péron, G., and D. N. Koons. 2012. Integrated modeling of communities: parasitism, competition,

and demographic synchrony in sympatric ducks. Ecology 93:2456–2464.

Piatt, J. F., A. M. A. Harding, M. Shultz, S. G. Speckman, T. I. van Pelt, G. S. Drew, and A. B.

Kettle. 2007. Seabirds as indicators of marine food supplies: Cairns revisited. Marine

Ecology Progress Series 352:221–234.

Pinaud, D., and H. Weimerskirch. 2002. Ultimate and proximate factors affecting the breeding

performance of a marine top-predator. Oikos 99:141–150.

van de Pol, M., Y. Vindenes, B.-E. Sæther, S. Engen, B. J. Ens, K. Oosterbeek, and J. M.

Tinbergen. 2010. Effects of climate change and variability on population dynamics in a

long-lived shorebird. Ecology 91:1192–1204.

Pradel, R. 2005. Multievent: An Extension of Multistate Capture-Recapture Models to Uncertain

States. Biometrics 61:442–447.

Quillfeldt, P., H. Weimerskirch, K. Delord, and Y. Cherel. 2020. Niche switching and leapfrog

foraging: movement ecology of sympatric petrels during the early breeding season.

Movement Ecology 8.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Ratcliffe, N., and R. W. Furness. 1999. The effect of parental age and experimentally

manipulated brood size on the foraging effort and breeding performance of great skuas

(*Catharacta skua*). Journal of Zoology 249:195–201.

Regehr, H. M., and W. A. Montevecchi. 1997. Interactive effects of food shortage and predation

on breeding failure of black-legged kittiwakes: Indirect effects of fisheries activities and

implications for indicator species. Marine Ecology Progress Series 155:249–260.

Saunders, S. P., F. J. Cuthbert, and E. F. Zipkin. 2018. Evaluating population viability and

efficacy of conservation management using integrated population models. Journal of

Applied Ecology 55:1380–1392.

Schaub, M., and F. Abadi. 2011. Integrated population models: a novel analysis framework for

deeper insights into population dynamics. Journal of Ornithology 152:227–237.

Sinclair, A. R. E., S. Mduma, and J. S. Brashares. 2003. Patterns of predation in a diverse

predator-prey system. Nature 425:288–290.

Stenseth, N. C., J. M. Durant, M. S. Fowler, E. Matthysen, F. Adriaensen, N. Jonzén, K. Chan,

H. Liu, J. De Laet, B. C. Sheldon, M. E. Visser, and A. A. Dhondt. 2015. Testing for effects

of climate change on competitive relationships and coexistence between two bird species.

Proceedings of the Royal Society B: Biological Sciences 282.

Stenseth, N. C., A. Mysterud, G. Ottersen, J. W. Hurell, K.-S. Chan, and M. Lima. 2002.

Ecological Effects of Climate Fluctuations. Science 297:1292–1296.

Stoessel, M., B. Elmhagen, M. Vinka, P. Hellström, and A. Angerbjörn. 2019. The fluctuating

world of a tundra predator guild: bottom-up constraints overrule top-down species

interactions in winter. Ecography 42:488–499.

Suryan, R. M., F. Sato, G. R. Balogh, K. David Hyrenbach, P. R. Sievert, and K. Ozaki. 2006.

Foraging destinations and marine habitat use of short-tailed albatrosses: A multi-scale

approach using first-passage time analysis. Deep-Sea Research Part II: Topical Studies in

Oceanography 53:370–386.

Sydeman, W. J., E. Poloczanska, T. E. Reed, and S. A. Thompson. 2015. Climate change and

marine vertebrates. Science 350:772–777.

Thomas, C. D., A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F.

N. Erasmus, M. Ferreira de Siquiera, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A. S.

van Jaarsveld, G. F. Midgley, L. Miles, M. A. Ortega-Huerta, A. Townsend Peterson, O. L.

Phillips, and S. E. Williams. 2004. Extinction risk from climate change. Nature 427:145–

148.

Tuljapurkar, S., and H. Caswell. 1997. Structured-Population Models in Marine, Terrestrial, and

Freshwater Systems. Chapman & Hall, New York.

de Valpine, P., D. Turek, C. J. Paciorek, C. Anderson-Bergman, D. T. Lang, and R. Bodik. 2017.

Programming With Models: Writing Statistical Algorithms for General Model Structures

With NIMBLE. Journal of Computational and Graphical Statistics 26:403–413.

Walther, G., E. Post, P. Convey, A. Menzel, C. Parmesan, T. J. C. Beebee, J. Fromentin, O.

Hoegh-Guldberg, and F. Bairlein. 2002. Ecological responses to recent climate change.

Nature 416:389–395.

Warham, J. 1990. The petrels: their ecology and breeding systems. Academic Press, London,

UK.

Weimerskirch, H. 1992. Reproductive effort in long-lived birds: age-specific patterns of

condition, reproduction and survival in the wandering albatross. Oikos 64:464–473.

Weimerskirch, H., O. Chastel, L. Ackermann, T. Chaurand, F. Cuenot-Chaillet, X. Hindermeyer,

and J. Judas. 1994. Alternate long and short foraging trips in pelagics seabird parents.

Anim. Behav. 47:472–476.

Zipkin, E. F., and S. P. Saunders. 2018. Synthesizing multiple data types for biological

conservation using integrated population models. Biological Conservation 217:240–250.

Tables

Table 1: Summary of the demographic parameters and their specificities (year random effect or state dependence) for the two species: the Brown Skua (top) and the Blue Petrel (bottom). Notations are $\widetilde{NB}$: Nonbreeder the previous year, $\widetilde{B}$: Breeder the previous year, NB: Nonbreeder, FB: Failed Breeder, SB1: Successful Breeder with one fledged chick, SB2: Successful Breeder with two fledged chicks, FBE: Failed Breeder at the Egg stage, FBC: Failed Breeder at the Chick stage and SB: Successful Breeder.

Species	Parameter	Year random effect	State dependence
Skua	Adult apparent survival ϕ_S	☑	$\widetilde{NB_S}\|\widetilde{B_S}$
	Breeding probability β_S	☑	$\widetilde{NB_S}\|\widetilde{B_S}$
	Breeding success γ_S	☑	$\widetilde{NB_S}\|\widetilde{B_S}$
	Breeding success 2 chicks δ_S	☑	$\widetilde{NB_S}\|\widetilde{B_S}$
	Detection probability p_S	☑	$\widetilde{NB_S}\|\widetilde{B_S}$
	Uncertain state assignment probability u_S	☑	$NB_S\|FB_S\|SB1_S\|SB2_S$
Petrel	Adult apparent survival ϕ_P	☑	$\widetilde{NB_P}\|\widetilde{B_P}$
	Breeding probability β_P	☑	$\widetilde{NB_P}\|\widetilde{B_P}$
	Hatching success ω_P	☑	$\widetilde{NB_P}\|\widetilde{B_P}$
	Breeding success γ_P	☑	$\widetilde{NB_P}\|\widetilde{B_P}$
	Detection probability p_P	☑	$\widetilde{NB_P}\|\widetilde{B_P}$
	Uncertain state assignment probability u_P	☑	$NB_P\|FBE_P\|FBC_P\|SB_P$

976   Table 2: Summary of the covariates tested on the demographic parameters of the two species –

977   the Brown Skua (top) and the Blue Petrel (bottom) – and the time period (in months) considered

978   for each demographic parameter. Notations are PP: Predator-Prey interactions, DD: intraspecific

979   Density-Dependence, SAM: Southern Annual Mode, SSTa: Sea Surface Temperature anomalies,

980   Chla: Chlorophyll a concentration.

Species	Parameter	Covariates tested	Time period
	Adult apparent survival ϕ_S	DD|SAM	Wintering (March-September)
Skua	Breeding probability β_S	DD|SAM	Pre-Breeding (July-November)
	Breeding success γ_S	PP|DD|SAM	Breeding (October-February)
	Breeding success 2 chicks δ_S	PP|DD|SAM	Breeding (October-February)
	Adult apparent survival ϕ_P	PP|DD|SAM|SSTa|Chla	Wintering (February-September)
Petrel	Breeding probability β_P	DD|SAM|SSTa|Chla	Pre-Breeding (August-October)
	Hatching success ω_P	PP|DD|SAM|SSTa|Chla	Breeding (October – December)
	Breeding success γ_P	PP|DD|SAM|SSTa|Chla	Breeding (December-January)

Table 3: Regression coefficients estimates for the relationships between covariates (DD:

intraspecific Density-Dependence, PP: Predator-Prey interactions, SAM: Southern Annular

Mode, SSTa: Sea Surface Temperature anomalies, Chla: Chlorophyll a concentration) and

demographic parameters (ϕ: adult apparent survival, β: breeding probability, γ: breeding

success, δ: breeding success with two chicks, ω: hatching success) for Brown Skuas (top) and

Blue Petrels (bottom), $\tilde{B}$: breeders or $\widetilde{NB}$: non breeders the previous years. 80% credible

intervals that do not include zero are in bold.

SKUA	DD				PP				SAM			
Parameters	slope	sd	10%	90%	slope	sd	10%	90%	slope	sd	10%	90%
$\phi_{S,\tilde{B}}$	-0.09	0.11	-0.22	0.06					-0.38	0.41	-0.88	0.12
$\phi_{S,\widetilde{NB}}$	-0.12	0.15	-0.28	0.09					**1.25**	**0.96**	**0.14**	**2.42**
$\beta_{S,\tilde{B}}$	0.23	0.22	-0.08	0.51					0.73	0.69	-0.13	1.61
$\beta_{S,\widetilde{NB}}$	0.11	0.19	-0.16	0.33					0.80	0.65	0.00	1.60
$\gamma_{S,\tilde{B}}$	**-0.39**	**0.14**	**-0.56**	**-0.22**	**0.66**	**0.17**	**0.45**	**0.89**	0.12	0.20	-0.14	0.37
$\gamma_{S,\widetilde{NB}}$	-0.23	0.28	-0.54	0.15	0.20	0.39	-0.26	0.65	0.21	0.54	-0.45	0.86
$\delta_{S,\tilde{B}}$	**-0.53**	**0.24**	**-0.85**	**-0.22**	**1.24**	**0.38**	**0.78**	**1.74**	0.02	0.41	-0.49	0.53
$\delta_{S,N\tilde{B}}$	-0.40	0.44	-0.95	0.16	0.71	0.96	-0.47	1.80	-1.30	2.33	-4.07	0.79

PETREL	DD				PP				SAM				SSTa				Chla			
Parameters	slope	sd	10%	90%	slope	sd	10%	90%	slope	sd	10%	90%	slope	sd	10%	90%	slope	sd	10%	90%
$\phi_{P,\tilde{B}}$	-0.14	0.30	-0.51	0.24	0.25	0.24	-0.06	0.56	-0.34	0.39	-0.83	0.16	-0.22	0.24	-0.53	0.07	-0.51	0.47	-1.08	0.05
$\phi_{P,\widetilde{NB}}$	-0.38	1.05	-1.67	1.04	0.43	0.49	-0.25	1.01	1.27	2.26	-1.11	4.51	0.01	0.96	-1.01	1.25	-0.56	1.70	-2.43	1.59
$\beta_{P,\tilde{B}}$	**0.51**	**0.25**	**0.19**	**0.86**					-0.34	0.39	-0.85	0.14	0.07	0.22	-0.22	0.34	0.34	0.38	-0.16	0.82
$\beta_{P,\widetilde{NB}}$	0.21	0.33	-0.22	0.67					0.40	0.38	-0.07	0.85	**0.36**	**0.22**	**0.09**	**0.63**	**0.63**	**0.40**	**0.12**	**1.13**
$\omega_{P,\tilde{B}}$	0.12	0.25	-0.18	0.43	**0.28**	**0.14**	**0.11**	**0.44**	-0.10	0.25	-0.41	0.20	0.08	0.16	-0.12	0.28	-0.35	0.29	-0.70	0.01
$\omega_{P,\widetilde{NB}}$	-0.47	0.52	-1.08	0.08	0.26	0.33	-0.16	0.66	-0.17	0.61	-0.87	0.53	-0.12	0.47	-0.56	0.37	-0.51	0.69	-1.30	0.28
$\gamma_{P,\tilde{B}}$	0.05	0.61	-0.68	0.81	-0.12	0.31	-0.49	0.27	0.32	0.46	-0.22	0.93	-0.44	0.42	-0.98	0.05	-0.13	0.64	-0.91	0.70
$\gamma_{P,\widetilde{NB}}$	**-0.79**	**0.59**	**-1.53**	**-0.12**	**0.46**	**0.33**	**0.01**	**0.89**	-0.43	0.45	-0.99	0.08	-0.34	0.51	-0.99	0.25	-0.85	0.79	-1.84	0.06

Figure Legends

Figure 1: Structure of the multispecies Integrated Population Model. Squares represent the state variables, circles represent the parameters. Data and fixed values are represented with a dark background, estimated state variables and parameters with a white background. Two types of data are used, capture histories (Ch) from capture-recapture data and count data (Y). Adult apparent survival (ϕ), breeding probability (β), hatching success (ω), breeding success (γ), breeding success with two chicks (δ), juvenile apparent survival for one to four years old and older (ϕ_{J1} to ϕ_{J4}), probability of first reproduction (Pr), state assignment probability (u) and detection probability (p) are parameters estimated in the model. Fecundity (f) is fixed. The number of adults (N_{adtot}), dead (N_{dead}), alive (N_{alive}), breeders (N_B), nonbreeders (N_{NB}), failed breeders (N_{FB}), failed breeders at the stage egg (N_{FBE}), breeders with an egg hatched (N_{SH}), failed breeders at the stage chick (N_{FBC}), successful breeders (N_{SB}), successful breeders with one chick (N_{SB1}) or with two chicks (N_{SB2}) and the number of juveniles of one year old to four years old and older (N_{J1} to N_{J4+}) are state variables estimated by the model. The number of immigrants (N_{im}) is a fixed vector. The blue part is for Blue Petrels and the brown part is for Brown Skuas. Interspecific relationships are represented with thick arrows.

Figure 2: Effects of predator-prey relationships (top panels) and intraspecific density-dependence (bottom panel) on adult demographic parameters for the two seabirds, the Brown Skua and the Blue Petrel. Solid lines represent the estimated relationship between the covariates and the demographic parameters. Shaded areas are the 50% and 95% credibility intervals. Points represent demographic parameter estimates each year (21 years) plotted against covariate. Error bars are standard deviation. Prey effect on (a) the estimated breeding success probability

$(\alpha_{PP_{\gamma,S,\tilde{B}}})$ and (b) breeding success with two chicks for skuas that bred the previous breeding

season $(\alpha_{PP_{\delta,S,\tilde{B}}})$. Intraspecific density-dependence effect on (c) the breeding success $(\alpha_{DD_{\gamma,S,\tilde{B}}})$

and on (d) breeding success of skuas that were breeders the previous breeding season $\left(\alpha_{DD_{\delta,S,\tilde{B}}}\right)$

and (e) on the breeding probability of petrels that bred the previous breeding season $(\alpha_{DD_{\beta,P,\tilde{B}}})$.

Figures

Figure 1

1020

1021

 Figure 2