Multispecies integrated population model reveals bottom-up dynamics in a seabird predator-prey

system

Maud Quéroué[1], Christophe Barbraud[2], Frédéric Barraquand[3], Daniel Turek[4], Karine Delord[2],

Nathan Pacoureau[5], Olivier Gimenez[1]

[1]CEFE UMR 5175, Univ. Montpellier, CNRS, EPHE, IRD, Univ. Paul Valéry Montpellier 3,

Montpellier, France

[2]CEBC UMR7372, CNRS – La Rochelle Université, 79360 Villiers en Bois, France

[3]Institute of Mathematics of Bordeaux, CNRS, Talence, France

[4]Department of Mathematics and Statistics, Williams College, Williamstown, MA, USA

[5]Department of Biological Sciences, Earth to Ocean Research Group, Simon Fraser University,

Burnaby, British Columbia, Canada

Correspondence author: Maud Quéroué, CEFE UMR 5175, Univ. Montpellier, CNRS, EPHE,

IRD, Univ. Paul Valéry Montpellier 3, Montpellier, France

E-mail: maud.queroue@gmail.com

Abstract

Assessing the effects of climate and interspecific relationships on communities is challenging because of the complex dynamic of species in interactions and the need to integrate information across several biological levels (individuals – populations – communities). To quantify species interactions, integrated population models (IPMs) have recently been extended to communities and allow fitting multispecies matrix models to data from multiple sources. However, abiotic factors are still rarely considered. Here, we used multispecies IPMs accommodating climate conditions to quantify the relative contribution of climate vs. interspecific interactions on demographic parameters, such as survival and breeding success, in the dynamics of a predator-prey system. We considered a stage-structured predator–prey system combining 22 years of capture—recapture data and population counts of two seabirds, namely the Brown skua (*Catharacta lönnbergi*) and its main prey the Blue petrel (*Halobaena caerulea*) both breeding on the Kerguelen Islands in the Southern Indian Ocean. Our results showed that the demography of the petrel and the skua were not driven in the same way by climate and prey-predator interactions. While the breeding success of skuas appears to be largely driven by the number of petrels, we did not find any link between climate fluctuations and the demography of skuas. In contrast, petrel breeding success was not explained by the number of skuas but mainly by the environmental conditions. We discuss the mechanisms by which climate variability and predator-prey relationships may affect the demographic parameters of these seabirds. We highlighted that bottom-up mechanisms drive this skua-petrel system. Taking into account both species interactions and environmental covariates in the same analysis improved our understanding of species dynamics in the face of environmental changes.

Keywords: Bayesian inference, Demography, Environmental variations, Integrated Population Model, Matrix population model, Nimble, Predator-Prey interactions

Introduction

The effects of climate changes on the diversity and the structure of communities have been reported repeatedly (Walther et al. 2002, Parmesan 2006, Hoegh-Guldberg and Bruno 2010, Miller et al. 2018). However, the underlying mechanisms remain poorly understood (Godfray and May 2014) due to the complex dynamics of interacting species (within species, between species and between species and the environment). Following disturbance, the abundance and distribution of species are expected to be modified according to the position and extent of the species' niche (Thomas et al. 2004). Because the effects of environmental variability on mortality, fecundity and dispersal may differ between species (Grosbois et al. 2008, Jenouvrier 2013), changes in structure and diversity appear at the community level. However, studying species-by-species responses to environmental changes may miss the role played by species interactions on those responses, and does not contribute to the larger understanding of species interactions that is required by community ecology.

Population dynamics models have been used to understand the effect of interspecific interactions and environment on species demography. However, these models are in general not demographically structured (Stenseth et al. 2015, Pacoureau et al. 2019a, Stoessel et al. 2019) or only partially (Millon et al. 2014, Pacoureau et al. 2019b). Unstructured approaches consider individuals as being equivalent while differences in size, age and ontogenic stages exist within a population and may be of importance to decipher interspecific interactions. As argued by Miller

& Rudolf (2011), the consideration of the structure of species in stages can lead to a better

understanding of community structure and dynamics. Interactions between species such as

predation or competition do not necessarily have a homogeneous impact on the different stages of

the interacting species. For example, young individuals might be predominantly preyed upon in

carnivore–ungulate systems (Gervasi et al. 2015). Therefore, to detect and understand species

interactions, we need to consider jointly the demography of several stage-structured populations

(Oken and Essington 2015).

Although well developed for single-species dynamics (Tuljapurkar and Caswell 1997, Caswell

2001), demographic stage-structured models have received little attention in community ecology

(but see Chu and Adler (2015) on plant system). The difficulty is that multispecies demography

analysis requires integrating information across several biological levels (individual – population

– community) which, in turn, requires unifying all available data sources into a single framework.

Integrated population models (IPMs) have been recently developed to infer population

demography by making complete use of all available sources of information (see Schaub and

Abadi 2011 and Saunders et al. 2018 for reviews). In their simplest form, these models combine

population counts and demographic data into a single analysis, which allows the estimation of

demographic parameters while simultaneously accounting for various sources of uncertainty in

each data source (e.g., measurement error or parameter estimation) (Besbeas et al. 2002). The IPM

framework has been extended to multiple species (Péron and Koons 2012) for

competition/parasitism, and more recently for predator-prey interactions (Barraquand and

Gimenez 2019).

Here, our main objective was to quantify the relative contribution of environmental changes and

species interactions on demographic parameters of a predator and its prey. Therefore, we used a

multispecies IPM framework accommodating the effects of local and global climatic conditions

on demographic parameters, such as survival and breeding, while explicitly considering species

interactions. We applied our approach on a stage-structured predator–prey system combining 22

years of capture-recapture data and population counts on two seabirds, the Brown skua

(*Catharacta lönnbergi*) and its main prey the Blue petrel (*Halobaena caerulea*) ('skua' and 'petrel'

hereafter) breeding on the Kerguelen Islands in the southern Indian Ocean. The Southern and

Indian Oceans have experienced a substantial increase in temperature (Gille 2002, Han et al. 2014),

making the Kerguelen Islands a critical location for studying the effects of climate change on

ecosystems. Seabirds are top predators that play an important role in ecosystem functioning and

are considered indicators of environmental changes in marine ecosystems (Hazen et al. 2019).

Their foraging behavior and demography reflect the influences of climate variability that directly

impacts biological processes in marine ecosystems, which cascade through food webs and are

integrated by seabirds (Barbraud and Weimerskirch 2001, Jenouvrier et al. 2003).

Using a multispecies IPM, we assessed the relative contribution of environment and predator-prey

interactions on seabirds' demographic parameters. We estimated survival and adult breeding

success for the two interacting species, and assessed the impacts of climatic conditions on these

demographic parameters to understand the contribution of predator-prey interactions in shaping

population dynamics.

Materials and Methods

Site and Species

Skua and Petrel were studied on Mayes Island (49°28'S, 69°57'E), a 5-km uninhabited island of the Kerguelen archipelago in the Southern Indian Ocean. We considered 22 breeding seasons from 1996/1997 to 2017/2018.

The petrel is a small (150–250g) long-lived Procellariiformes seabird. At Kerguelen, the petrel is a macrozooplankton and micronekton feeder, its diet being mainly composed of crustaceans and fishes (Cherel et al. 2002, 2014). Individuals from Mayes Island spend the nonbreeding season (from mid-February to September) between the polar front and the northern limit of the ice pack (57-62°S) between longitudes 20°W and 90°E (Cherel et al. 2016). Birds return to colonies in early September (Quillfeldt et al. 2020). On Mayes Island the breeding population is estimated at approximately 142,000 breeding pairs (Barbraud and Delord 2006). In late October, a single egg is laid in a burrow dug in peat soil under tall and dense vegetation. The incubation phase takes between 45 and 49 days and the chick rearing period between 43 and 60 days (Jouventin et al. 1985). The chick fledges in early February. Both sexes participate in parental care by alternating foraging trips during the incubation and fledging periods.

The skua is a medium sized (1.1 – 2.2 kg) long-lived Charadriiformes seabird. On Mayes Island between 80 and 120 pairs breed annually (Mougeot et al. 1998). Breeding pairs form in October with a high mate fidelity, and generally establish themselves in the same territory each year (Parmelee and Pietz 1987), which they tenaciously defend throughout the breeding season. Generally two eggs are laid between October and December. The incubation phase takes between

28 and 32 days and the chicks rearing period between 40 and 50 days (Higgins and Davied 1996).

With their different foraging techniques, skuas adapt their diet depending on the availability of

prey (Carneiro et al. 2015). On Mayes Island, during the breeding season, Blue petrels represent

95% of the skua diet (Pacoureau et al. 2019c). Skuas from Mayes Island overwinter in the southern

hemisphere between 10°E and 150°E (Delord et al. 2018).

During the breeding period on Mayes Island, the predation of petrels by skuas takes place mainly

at night, when the petrels come out or arrive at their burrows (Mougeot & Bretagnolle, 2000a).

Skuas mostly predate petrels on the ground, but they can also predate petrels in flight (Mougeot et

al. 1998, Pacoureau et al. 2019c). Vocalizing petrels, especially those without partners, are more

easily detected by skuas during the courtship period (Mougeot and Bretagnolle 2000b). Skuas may

also predate chicks during the fledging period.

Counts and capture-recapture data

Individuals of each species were checked at specific times following the species phenology to

determine the breeding status of each bird. The breeding status of marked birds was determined at

the end of the breeding period.

Around 200 individually marked burrows of petrels were inspected each year from early-to-mid

November just after laying to check for eggs and to record marked adults, and then in late January

just before fledging. Each year since 1985 (see Barbraud and Weimerskirch 2005), all fledglings

found in burrows were marked with a stainless steel band, as well as new individuals found in

monitored burrows (captured by hand, marked, and replaced in their burrow). We used capture-

recapture data from 1996 and considered new individuals found in monitored burrows as

immigrants to the study site. Petrels never observed with an egg or a chick during a given breeding

season were considered as nonbreeders (NB_p). Individuals were identified as breeders (B_p) if they

laid a single egg or raised a chick and as successful breeders if their chick fledged (SB_p). Two

categories of failed breeders were used depending on the stage of failure: egg stage (FBE_p) or

chick stage (FBC_p). Given that the first sampling period occurred just after laying, it is very

unlikely that nonbreeders were failed breeders. The annual number of petrels was estimated as the

number of occupied burrows.

For skuas, each year since 1991, the eastern side of Mayes Island was inspected to identify

territories of skuas. A territory was considered established when a pair strongly defended an area

against other skuas (Mougeot et al. 1998). Around 50 nesting territories were visited four to eight

times from mid-October (after laying) to late-March (just before fledging) each year. Chicks were

marked just before fledging, as well as new adult individuals with a metal ring and a plastic ring

to facilitate individual identification using binoculars. Breeding status was determined by checking

the nest contents for the presence of eggs or young chicks. Skuas never observed with an egg or a

chick were considered as nonbreeders (NB_s). Individuals were identified as breeders (B_s) if they

laid at least one egg or raised a chick. If the eggs did not hatch or the chicks died, the breeding pair

was considered as failed breeder (FB_s). Given that the first sampling period occurred just after

laying, it is very unlikely that nonbreeders represented failed breeders. Successful breeders were

able to fledge one or two chicks which are denoted as $SB1_s$ or $SB2_s$, respectively. The annual

number of skuas was estimated as the number of territories.

For both species, individuals could be considered as "uncertain" (C) in case of difficulties to assign

their breeding status (lack of information, missed checks, individuals never re-observed). Only

individuals that have bred at least once between the 1996 and the 2017 breeding seasons were kept

in the data set for analysis ($n = 320$ for skuas and $n = 1210$ for petrels). Individual capture histories

started at their first breeding attempt recorded. Based on the high probability of observing breeders

in the study site, we assumed that the first breeding attempt was correctly detected.

Integrated Population Model

We built a two-species IPM that combines counts and capture-recapture data and allows estimating

abundances and demographic rates (Besbeas et al. 2002, Schaub and Abadi 2011). More

specifically, we connected two IPMs, one for the skuas, the predators, and one for the petrels, the

preys, through explicit predator-prey relationships (Barraquand and Gimenez 2019). Counts and

capture-recapture data for both species were combined into a joint likelihood that also included

the effects of intra-species and inter-species density-dependence and climatic covariates on

demographic parameters.

The two IPMs have the same structure but states differ in relation to species biology. The two main

differences are: (1) skuas can have up to two chicks versus only one for petrels, (2) additional

information is available for petrels which can determine the precise stage of breeding in case of

failure (egg or chick stage). We explain the structure of the IPMs based on the skua model and

differences are noted when they occur. See Appendix S1 for a full description of the petrel model.

For clarity, all parameters for skuas are indicated by s and for petrels by p.

Count data

State process:

We use Poisson (Po) and binomial (Bin) distributions to account for demographic stochasticity.

Skuas from one year old to four years old are considered as juveniles. The number of skuas in their

first year ($J_{1_{s_t}}$) at year t is modelled with a Poisson distribution:

$J_{1_{s_t}} \sim Po(0.5 \times f_{SB1_s} \times SB1_{s_{t-1}} + 0.5 \times f_{SB2_s} \times SB2_{s_{t-1}})$ (1)

with J_{1_s} the number of chicks by successful breeder ($SB1_s$ and $SB2_s$) female (sex ratio: 0.5)

according to their fecundity (f_{SB1_s}: 1 chick and f_{SB2_s}: 2 chicks per skua). For petrels, J_{1_p} is also

Poisson distributed but with only one chick per successful breeder (Appendix S1). The number of

juveniles of two (J_{2_s}), three (J_{3_s}), and four years old (J_{4_s}) are modelled with binomial

distributions:

$J_{2_{s_t}} \sim Bin(\phi_{J_{1_s}}, J_{1_{s_{t-1}}})$ (2)

$J_{3_{s_t}} \sim Bin(\phi_{J_{2_s}}, J_{2_{s_{t-1}}})$ (3)

$J_{4_{s_t}} \sim Bin(\phi_{J_{3_s}}, J_{3_{s_{t-1}}})$ (4)

with the apparent survival between one and two years ($\phi_{J_{1_s}}$), between two and three years ($\phi_{J_{2_s}}$)

and between three and four years ($\phi_{J_{3_s}}$) respectively. As the first breeding attempt in skuas could

start from age four, the J_{4+_s} state was considered as the immature state with four years old skuas

(J_{4_s}) and older that have never reproduced ($J_{4+_s}^{NB}$):

$J_{4+_{s_t}}^{B} \sim Bin\left(Pr_{s_t}, \phi_{J_{4_s}} \times J_{4+_{s_{t-1}}}\right)$ (5)

$J_{4+_{s_t}}^{NB} = \phi_{J_{4_s}} \times J_{4+_{s_{t-1}}} - J_{4+_{s_t}}^{B}$ (6)

$$J_{4+_{s_t}} = J_{4_{s_t}} + J_{4+_{s_t}}^{NB} \qquad (7)$$

with $\phi_{J_{4_s}}$ the apparent survival for immatures skuas, Pr_s the probability of first breeding attempt

for skuas in the state J_{4+_s} that have not recruited yet and $J_{4+_s}^{B}$ the number of skuas that attempt to

breed for the first time.

Because we had no information on the juvenile phase, we assumed that juvenile apparent survival

increased with age (Greig et al. 1983, Grande et al. 2009, Fay et al. 2015), as experienced birds

are on average more effective in foraging (Daunt et al. 2007), in competing with conspecifics or

in avoiding predators:

$$logit\left(\phi_{j_{i_s}}\right) = \lambda_{1_s} + \lambda_{2_s} \times i \qquad (8)$$

where $\phi_{j_{i_s}}$ is the juvenile apparent survival, i the age of the juvenile state (from J_{1_s} to J_{4_s}), λ_{1_s}

the intercept and λ_{2_s} the slope which is constrained to be positive.

We use binomial distributions to account for stochasticity in the different adult states. We modelled

the number of surviving adult skuas (N_{alive_s}) with a binomial distribution:

$$N_{alive_{s_t}} \sim Bin\left(\phi_{ad_{s_{t-1}}}, N_{ad_{s_{t-1}}}\right) \qquad (9)$$

with ϕ_{ad_s} the adult apparent survival. The number of skuas that have not bred among those that

survived (NB_s) is modelled as:

$$B_{ad_{s_t}} \sim Bin\left(\beta_{s_{t-1}}, N_{alive_{s_t}}\right) \qquad (10)$$

$$NB_{s_t} = N_{alive_{s_t}} - B_{ad_{s_t}} \qquad (11)$$

with β_s the probability of breeding and B_{ad_s} the number of adult breeders. The total number of

breeders (B_s) correspond to the sum of the number of adult breeders (B_{ad_s}), the number of

immigrants (*i.e.,* newly marked) skuas (N_{im_s}) and the number of juveniles attempting to breed for

the first time $(J^B_{4+_{st}})$:

$B_{s_t} = B_{ad_{s_t}} + J^B_{4+_{st}} + N_{im_{s_t}}$ (12)

The number of failed breeders (FB_s) and successful breeders (SB_s) are modelled following a

binomial distribution:

$SB_{s_t} \sim Bin\left(\gamma_{s_{t-1}}, B_{s_t}\right)$ (13)

$FB_{s_t} = B_{s_t} - SB_{s_t}$ (14)

with γ_s the probability of a successful breeding. A successful breeder can then have one or two

chicks, respectively $SB1_s$ and $SB2_s$ and this is modelled following a binomial distribution:

$SB2_{s_t} \sim Bin\left(\delta_{s_{t-1}}, SB_{s_t}\right)$ (15)

$SB1_{s_t} = SB_{s_t} - SB2_{s_t}$ (16)

with δ_s the probability of successful breeding with two chicks among the successful breeders.

For the petrel, there are two states for failed breeders: one with petrels that failed to hatch their egg

(called failed breeder at the egg stage FBE_p) and the second with petrels that failed to fledge their

chick (named failed breeder at the chick stage FBC_p). Hence, there is a parameter of successful

hatching (ω_p). There is only one state for successful breeders (SB_p) defined with a parameter of

successful breeding (y_p) (Appendix S1).

Observation process:

The observation equation links the observed adult population count ($N_{s\,counts}$) (*i.e.,* the number of

territories multiplied by two for a pair of skuas) with the true adult population size (N_{ad_s}), with an

additional term for observation error.

$$246 \quad N_{s\,counts_t} = (0\ 0\ 0\ 0\ 1\ 1\ 1\ 1) \begin{pmatrix} J1_s \\ J2_s \\ J3_s \\ J4_s \\ NB_s \\ FB_s \\ SB1_s \\ SB2_s \end{pmatrix}_t + \varepsilon_{s_t}$$

$$247 \quad \varepsilon_{s_t} \sim N(0, \sigma^2_{N_{s\,counts}}) \qquad\qquad\qquad (17)$$

where ε_{s_t} is the error term and $\sigma^2_{N_{s\,counts}}$ its variance. As only the adult states were observed on the

field, we used a (0 0 0 0 1 1 1 1) vector, excluding the juvenile states from the observation equation.

The likelihood for the population count data is denoted

as $L_{co_s}(N_{s\,counts}|\phi_{J1_s}, \phi_{J2_s}, \phi_{J3_s}, \phi_{J4_s}, Pr_s, \phi_{ad_s}, \beta_s, \gamma_s, \delta_s, N_{ad_s})$.

Capture-recapture data

For adult capture-recapture data, we used multievent capture–recapture models to estimate the

demographic parameters (Pradel 2005). These models take into account the imperfect detectability

of the individuals as well as the uncertainty in the assignment of states to individuals (Gimenez et

al. 2012).

Our multievent model includes five states: Nonbreeder (NB_s), Failed breeder (FB_s), Successful

breeder with one chick ($SB1_s$), Successful breeder with two chicks ($SB2_s$) and the uncertain

individuals (C_s). The following demographic parameters were estimated: the adult apparent

survival probability (ϕ_{ad_s}), the breeding probability (β_s), the probability of successful

breeding (γ_s) and the probability of successful breeding with two chicks (δ_s). Two additional

parameters were also estimated: the detection probability (p_s) and the state assignment probability

of individuals with uncertain state (u_s). In the petrel model there are two states for failed breeders

(at egg stage FBE_p or at chick stage FBC_p) but one state for successful breeder (SB_s), and the

hatching success parameter (ω_p) is also estimated (for details see Appendix S1). All parameters

are time-varying through a yearly random effect, except u_s (Table 1). State transitions are set to

be state dependent according to the breeding status in the previous breeding season: Breeder ($\widetilde{B_s}$)

representing birds that try to breed the previous breeding season (FB_s , $SB1_s$, $SB2_s$) or Nonbreeder

($\widetilde{NB_s}$) representing birds that already bred previously but did not try to breed during the previous

breeding season (NB_s), immigrants (N_{im_s}) and immature birds (J_{4_s}) that attempt to breed for the

first time in the colony (Table 1). The detection probability and the state assignment probability

also depend on the breeding status (Table 1). The likelihood for the capture-recapture data is

denoted as $L_{cr_s}(ch_s|\phi_{ad_s}, \beta_s, \gamma_s, \delta_s, p_s, u_s)$.

Joint likelihood

The joint likelihood of this IPM is the product of the likelihood for the count data (L_{co_s}) and

capture-recapture data (L_{cr_s}):

$L_{ipm_s}(N_{s_{counts}}, ch_s | \phi_{J_{1_s}}, \phi_{J_{2_s}}, \phi_{J_{3_s}}, \phi_{J_{4_s}}, Pr_s, \phi_{ad_s}, \beta_s, \gamma_s, \delta_s, N_{adtot_s}, p_s, u_s)$

$= L_{co_s}(N_{s_{counts}} | \phi_{J_{1_s}}, \phi_{J_{2_s}}, \phi_{J_{3_s}}, \phi_{J_{4_s}}, Pr_s, \phi_{ad_s}, \beta_s, \gamma_s, \delta_s, N_{ad_s}) \times L_{cr_s}(ch_s | \phi_{ad_s}, \beta_s, \gamma_s, \delta_s, p_s, u_s)$

$$(18)$$

Interspecific relationship, intraspecific density-dependence, and environmental covariates

We used different covariates to quantify their contribution to the adult demographic parameters

estimated for the two species (Table 2). We tested predator-prey relationships between the skua,

the predator, and its prey, the petrel, and intraspecific relationships with density dependence for

both species. We considered several climatic covariates that were suspected to affect demographic

parameters of skuas and petrels, the Southern Annular Mode (SAM) on a large scale, and the Sea

Surface Temperature anomalies (SSTa) and Chlorophyll a concentration (Chla) on a local scale.

Predator-prey interactions

We combined the two IPMs for skuas and petrels in the same model, which allowed us to consider

interspecific relationships. Based on the high proportion of petrels in the diet of the skuas during

the breeding season (Pacoureau et al. 2019c), we predicted that petrel adult apparent survival

(ϕ_{ad_p}), hatching success (ω_p) and fledging success (γ_p), should decrease with the number of adult

skuas. Inversely, skua breeding success (γ_s) and breeding success with two chicks (δ_s) should

increase with the number of petrels.

Intraspecific density-dependence

As skuas are highly territorial and defend vigorously their territories, we expected a negative density-dependence relationship between breeding probability (β_s) and population density. Furthermore, the energetic cost and the time spent in defending a territory throughout the breeding season could jeopardized the breeding success and increase the adult mortality. We thus predicted a negative density-dependence relationship on the successful breeding parameter (γ_s), the probability to have two chicks for successful breeders (δ_s) and the adult apparent survival parameter (ϕ_{ad_s}). For petrels, we also tested the effects of intraspecific competition for food resources, which could affect their adult apparent survival (ϕ_{ad_p}) and their breeding parameters: breeding probability (β_p), hatching (ω_p) and fledging success (γ_p).

Southern Annular Mode

The Southern Annular Mode (SAM) is a large-scale climate index. SAM is the leading mode of climate variability over the Southern Hemisphere. SAM is defined as the difference of atmospheric pressure between the 40°S and 65°S latitudes (Marshall 2003). SAM influences surface wind, sea surface temperature (SST) and surface chlorophyll concentration. Data were obtained from the online database of the British Antarctic Survey (http://www.nerc-bas.ac.uk/ icd/gjma/sam.html).

South of the Polar Front, where petrels winter, positive phases of the SAM are associated with westerly winds. It induce cold SST anomalies and Ekman transport (the 90° wind-driven net transport on the sea surface) and drive increased upwelling (Lovenduski and Gruber 2005). Consequently, the biological productivity and potential prey availability for petrels are accentuated

during positive phases of the SAM. We thus predicted that the positive phases of SAM would be

favorable to petrel survival and breeding success.

Since skuas have broad wintering areas (Delord et al. 2018), we have chosen to test a large-scale

environmental covariate, the SAM. Contrary to their diet during the breeding season specialized

on the Blue petrel, during winter they adopt a mixed diet composed of low trophic level preys,

macrozooplankton and crustaceans (Delord et al. 2018). Availability of food resources at sea

during their wintering might have an effect on the condition of skuas. A large majority of the skuas

from Mayes Island overwinter north of the polar front (Delord et al. 2018). In the subtropical zone,

SAM positive phases induced warm SST anomalies, low surface chlorophyll concentration and

easterly winds driving Ekman transport, while in the Subantarctic zone there is a convergence of

waters that increase downwelling and positive SST anomalies (Lovenduski and Gruber 2005). We

thus predicted that the positive phases of SAM, leading to poorer food availability in the areas

used by skuas during the nonbreeding period, would have negative impacts on skua survival and

on their ability to reproduce in the next breeding season.

Sea Surface Temperature anomalies

We used the SSTa as a proxy of food availability because it is known to indirectly affect seabird

demography through its effects on marine food webs (Barbraud et al. 2012). Because the year-to-

year variation of SST was negatively correlated with the body condition of petrels (Guinet et al.

1998), we predicted that high SSTa would negatively affect its survival and breeding success. The

SSTa data were downloaded from NOAA ("data: NOAA NCEP EMC CMB GLOBAL

Reyn_SmithOIv2 monthly ssta") from 1996 to 2018.

Chlorophyll a

Because the petrel diet is mainly composed of crustaceans and fish feeding at low trophic levels (Cherel et al. 2002, 2014), we predicted that high concentrations of Chla would be favorable to survival and breeding success via bottom-up mechanisms. The Chlorophyll a concentration (Chla) data were downloaded from the NASA Ocean Data with a 9km mapped concentration data of chlorophyll a for the years 1997 to 2001 and from the Nasa Earth Observation (NEO AQUA/MODIS data) monthly for the years 2002 to 2018.

For local covariates (SSTa and Chla), we recovered the average values of the covariates in the areas in which petrels were located (Cherel et al. 2016) in a specific time period during which the environment might affect the demographic parameter under investigation (Table 2). Each environmental covariate was standardized to have zero mean and unit variance.

Assessing the effect of covariates and state variables

We fitted a single model including all the biologically relevant effects. Logit-linear regressions were used to estimate the effect of environmental (SAM, SSTa and Chla) and inter- and intra-specific interactions on demographic parameters (adult apparent survival, breeding probability, hatching probability, breeding success) (Table 2). We used state variables N_{ad_S} and N_{ad_p}, i.e. the number of adult skuas and petrels respectively, to assess the effects of inter- and intra-specific interactions. For example, we modelled the hatching probability for petrels that breed the previous year (ω_p^B) using a logit link:

$$logit\left(\omega_{p_t}^B\right) = \mu_{\omega_p^B} + \alpha_{SAM\omega_p^B} \times SAM_{\omega_{p_t}} + \alpha_{SST\omega_p^B} \times SST_{\omega_{p_t}} + \alpha_{Chla\omega_p^B} \times Chla_{\omega_{p_t}} +$$

$$\alpha_{DD\omega_p^B} \times Nad_{p_t} + \alpha_{PP\omega_p^B} \times Nad_{s_t} + \varepsilon_{\omega_{p_t}^B}$$

$\varepsilon_{\omega_{p_t}^B} \sim N(0, \sigma^2_{\varepsilon_{\omega_p^B}})$ (19)

with $\mu_{\omega_p^B}$ the intercept, $\alpha_{SAM\omega_p^B}$ the slope for the climatic covariate $SAM_{\omega_{p_t}}$, $\alpha_{SST\omega_p^B}$ the slope for

the climatic covariate $SST_{\omega_{p_t}}$, $\alpha_{Chla\omega_p^B}$ the slope for the climatic covariate $Chla_{\omega_{p_t}}$, $\alpha_{DD\omega_p^B}$ the

slope indicating the strength of the intra-specific density dependence with Nad_{p_t} the number of

adult petrels between year *t* and *t+1*, $\alpha_{PP\omega_p^B}$ the slope indicating the strength of the predator-prey

relationship with Nad_{s_t} the number of adult skuas between year *t* and *t+1*, $\varepsilon_{\omega_{p_t}^B}$ is a yearly random

effect and $\sigma^2_{\varepsilon_{\omega_p^B}}$ its temporal variance. The descriptions of all equations and parameter notations

are available in Appendix S2.

We computed the 95% credible intervals for the regression coefficients α. Effects were considered

statistically significant when credible intervals did not include 0 as described in Kéry and Schaub

(2011). We compared the relative contribution of each covariate using the regression estimate

which we used as a measure of effect size.

Model implementation

To fit the juvenile apparent survival parameters increasing with age, we modelled them as a

positive linear function of age by assigning to the slope λ_2 a $U(0,1)$ prior, and by defining the

intercept λ_1 with a normal $N(0,1)$ prior. The probability of the first breeding attempt (Pr) is time-

dependent with a non-informative prior: $Pr_t \sim U(0,1)$. We modelled covariate effects on the adult

demographic parameters as linear functions of time on the logit scale by assigning normal

priors $N(0,10^4)$ to the regression coefficients (α). For the variance of the random year effects (σ_ε^2),

we used a $U(0,10)$ non-informative prior. The state assignment probability of individuals with

uncertain state parameter (u_s) was defined *a priori* with a $U(0,10)$ non-informative prior.

Bayesian posterior distributions were approximated *via* Markov chain Monte Carlo (MCMC)

algorithms. Two independent MCMC chains of 220,000 iterations were used with a burn-in period

of 120,000. One out of five iterations was kept and final inferences were derived from a sample of

$2 \times 20,000$ iterations that resulted from merging the two chains. Gelman-Rubin convergence

diagnostic (Brooks and Gelman 1998) was below 1.5 for each parameter and the mixing of the

chains was satisfactory. We performed the analyses using Nimble (de Valpine et al. 2017) and

program R (R Core Team 2020) (R version 4.0.0). Code and data are available on GitHub at

https://github.com/maudqueroue/MultispeciesIPM_SkuaPetrel.

Results

For skuas, we found no significant effect of the SAM covariate on adult demographic parameters

(Table 3). We estimated that the number of skuas (for intra-specific relationship) and the number

of petrels (for inter-specific relationship) had a significant effect on two demographic parameters,

namely the breeding success and the breeding success with two chicks for skuas that were breeders

the previous breeding season. We found negative density-dependence for the breeding success

($\gamma_s B$) [mean $\alpha_{DD\ \gamma_s^B} = -0.56$; Credible intervals CRI $(-0.77, -0.34)$] (Fig. 1a) and for the

breeding success with two chicks ($\delta_s B$) [mean $\alpha_{DD\ \delta_s^B} = -0.78$; CRI $(-1.10, -0.38)$] (Fig. 1c).

We found positive predator-prey relationships for breeding success ($\gamma_s B$) and breeding success

with two chicks ($\delta_s B$) with an increase in these parameters with increasing number of petrels

[mean $\alpha_{PP\gamma_s^B} = 0.76$; CRI $(0.41, 1.14)$ and mean $\alpha_{PP\delta_s^B} = 1.14$; CRI $(0.57, 1.76)$] (Fig. 1b,

d). The predator-prey effects had stronger slopes than the intra-specific density-dependence effects

for the breeding success of skuas ($\left|\text{mean } \alpha_{PP\gamma_s^B}\right| = 0.76$; $\left|\text{mean } \alpha_{DD\gamma_s^B}\right| = 0.56$, respectively) or

the breeding success with two chicks ($\left|\text{mean } \alpha_{PP\delta_s^B}\right| = 1.14$; $\left|\text{mean } \alpha_{DD\delta_s^B}\right| = 0.78$,

respectively).

For petrels, we found significant relationships between the three environmental covariates (SAM,

SSTa and Chla) and hatching success for individuals that were nonbreeders the previous breeding

season ($\omega_p NB$). We found negative relationships with the SAM covariate [mean $\alpha_{SAM\omega_p^{NB}} =$

-1.03 ; CRI $(-1.93, -0.27)$] (Fig. 1e) and with the SSTa covariate [mean $\alpha_{SST\omega_p^{NB}} =$

-1.16 ; CRI $(-1.99, -0.47)$] (Fig. 1f). A positive relationship was estimated with the Chla

covariate [mean $\alpha_{Chla\omega_p^{NB}} = 1.19$; CRI $(0.47, 2.11)$] (Fig. 1g). The contributions of these

environmental covariates to hatching success increase in the order SAM → SSTa → Chla. There

was no significant effect of skua densities on petrel adult demographic parameters. We estimated

a positive effect of increased petrel adult number on the breeding probability for petrels that were

breeders the previous breeding season ($\beta_p B$) [mean $\alpha_{DD\beta_p^B} = 0.71$; CRI $(0.44, 0.90)$] (Fig. 1h).

In addition to the results above, we also estimated the demographic parameters and the number of

individuals in each state for both species from 1996 to 2017 (see Appendix S3: Figs. S1– S6).

Discussion

In this paper, we provide the first application of an IPM integrating predator-prey relationships to

empirical data. Our results showed that the demography of the prey, the petrel, and the predator,

the skua, was not driven in the same way by the different environmental variations. We found that

the petrel dynamic was mainly impacted by the environment whereas the skua dynamic was

primarily driven by the number of prey individuals, highlighting a bottom-up system.

Effects of predator-prey relationships

The number of prey is a determining factor in the breeding success of skuas. Food availability is

known to be positively related with breeding parameters in seabirds (Cairns 1988, Piatt et al. 2007,

Oro et al. 2014). As diet of skuas during the breeding period in Mayes Island is dominated by

petrels (Mougeot et al. 1998, Pacoureau et al. 2019c), a large abundance of petrels provides easier

conditions for skuas to feed themselves and their chicks resulting in a high breeding success.

Interestingly, we did not find the opposite relationship in the prey dynamic. Our model indicates

that the number of skuas had no effect on the demographic parameters of the petrel. However, it

has already been shown that in predator-prey terrestrial bird systems, the presence of a predator

might have consequences on the reproduction of the prey by increasing brood failures (Mueller et

al. 2016) or by reducing clutch sizes (Morosinotto et al. 2010). This lack of effect could be

explained by the large abundance of petrels compared to the skuas on Mayes Island. Oro et al.

(2006) reported that in another seabird predator-prey system, the highest breeding success of the

prey occurred when the prey/predator ratio was very high. On Mayes Island, the breeding

population of petrels is estimated at approximately 142,000 breeding pairs (Barbraud and Delord

2006), but this does not include the chicks (around 71,000 each year) and nonbreeders

(approximately 30% of the petrels). Hence, there are about 476,000 petrels during a breeding

season versus about 200 skuas showing a very high prey/predator ratio. Moreover, Mougeot et al.

(1998) showed that skuas of Mayes Island predated about 40,000 petrels each breeding season.

This corresponds to about 8% of the petrel population of the island. It is therefore possible that

skua predation did not affect petrel demographics or that the effect was not sufficient to be detected

in our model. In addition, as shown in our results and discussed later, it is likely that petrel

dynamics were mostly driven by climatic conditions.

Effects of intra-specific density-dependence

We found negative density-dependent effects on breeding success and probability to fledge two

chicks in the skua population. Egg and chick predation by conspecifics has been reported in the

Great skua *Catharacta skua* (Hamer et al. 1991, Ratcliffe and Furness 1999). Hence, a higher

abundance of skuas increases the risk of predation on eggs and chicks, resulting in higher breeding

failure. To avoid predation by conspecifics, the skuas start defending their territories from other

skuas just a few days after arrival on a breeding site until the end of the season. This activity is

energetically costly and may also limit the time spent searching for food, potentially limiting

energy investment in reproduction. The heterogeneous habitat hypothesis already demonstrated in

territorial birds (Dhondt et al. 1992, Krüger and Lindström 2001, Ferrer and Donazar 2015) could

also explain the relationships we found. Indeed, when the population increases, some individuals

are forced to occupy poorer quality habitats, resulting in lower reproductive success.

Predator-prey relationships had a higher contribution to the breeding success of skuas than the

density-dependent effect. Hamer et al. (1991) reported that, following a reduction of sandeel

abundance, Great skua increased their foraging effort reducing the adult territorial attendance. In

turn, breeding failure has increased due to predation from adults of neighboring territories. We

then may assume that petrel abundance allowed a suitable skua territorial attendance reducing the

negative density-dependent effects such as chick predation by conspecifics.

We did not find an effect of density-dependence on the breeding probability of skuas. As skuas

are territorial with a high degree of site fidelity, we assumed that in years with a high abundance

of skuas, the breeding probability would decrease as all the skuas would not have territories. It is

possible that we did not observe this effect because the logistic function used for density-

dependence does not accurately model the territory acquisition dynamics by floaters (e.g. van de

Pol et al. 2010, Barraquand et al. 2014).

Interestingly, we found a positive density-dependence relationship on the breeding probability of

petrels that bred the previous year. Combined effects of density-dependence and climate have

already been observed in the Blue petrel, with a lower winter survival when density is high

(Barbraud and Weimerskirch 2003), suggesting a mechanism of competition between conspecifics

for the resource. The inverse relationship we found rather suggests that years with a high

abundance of petrels reflect a good return rate to the breeding site because environmental

conditions were favorable for breeding. Indeed, petrels are known to skip breeding and take

sabbatical years when environmental conditions are poor (Warham, 1990; Chastel, Weimerskirch,

& Jouventin, 1995).

Effects of environmental conditions

Our results showed that the hatching success for the petrels that did not breed the year before was

impacted by the environmental covariates we tested, namely SAM, SST and Chla. Hatching

success is the only parameter that was impacted by the environment according to our model. This

result is in agreement with former studies showing that the hatching period is paramount during

the breeding period for Procellariiformes. The partners alternate on the nest, one incubating and

fasting and the other making foraging trips at sea, causing important variations in their body mass

(Chaurand and Weimerskirch 1994a, 1994b, Weimerskirch et al. 1994, Chastel et al. 1995). Blue

petrels can temporarily desert their burrow if they fall below a certain threshold mass (Chaurand

and Weimerskirch 1994b), even if the shift of the partner has not occurred. This desertion increases

the risk of hatching failure (Boersma and Wheelwright 1979, Chaurand and Weimerskirch 1994b).

It was shown that half of the failures during the breeding season occurred during the first

incubation shift, and were related to males in poor body condition (Chastel et al. 1995). Our model

might detect an effect on petrels that did not breed the previous year because nonbreeder petrels

are known to be in poorer condition than breeders (Chastel et al. 1995), potentially making them

more sensitive to environmental conditions.

High SST generally reduces vertical mixing and provides poor growing conditions for zooplankton

communities and then through bottom-up mechanisms induce reduced trophic resources for

seabirds (Barbraud et al. 2012, Sydeman et al. 2015). In contrast, high chlorophyll a concentration,

at the bottom of the trophic food chain, provides resources for higher trophic organisms including

seabirds. Hence high SSTs could decrease the foraging efficiency of petrels during incubation and

cause desertion of eggs whereas the high level concentration of chlorophyll a could have a positive

effect as shown in our results. Negative relationships have also be found between the SST during

the chick rearing period and the fledging success of petrels (Guinet et al. 1998). The negative

relationship detected in our analyses between the SAM and the hatching success contradicts our

expectations. Indeed, our assumption was that the positive phase of SAM would increase the

biological productivity and the prey availability in the petrel's feeding area south of the polar front.

A negative relationship was also found between SAM and juvenile survival in snow petrels

Pagodroma nivea (Sauser et al. 2018). These authors suggested that the extension of fast ice during

positive phases of SAM (Holland et al. 2017) decreases prey accessibility in the Antarctic zone at

the polar front, where the blue petrel feeds during the breeding season.

The effects of environmental covariates, used as a proxy of prey abundance, increase when the

covariates approach the trophic level occupied by the prey of the petrels (SAM $\rightarrow$ SST $\rightarrow$

Chlorophyll a). According to these results, the dynamic of petrels seems to be driven indirectly by

climate through bottom-up mechanisms. These mechanisms are consistent with many studies

showing that climatic conditions affect seabirds through indirect processes by influencing prey

availability and resulting in changes in their dynamics (Frederiksen et al. 2006, Barbraud et al.

2012, Jenouvrier 2013, Lauria et al. 2013).

We did not detect a significant relationship between the demographic parameters of the skua and

the environmental covariates. The lack of effect on breeding parameters could be explained by an

absence of a direct link between skuas and the environmental covariates tested as breeding skuas

remain on their territory to defend it or to forage. However, skuas are always at sea during their

wintering and their survival might be affected by environmental conditions. To our knowledge,

only few studies have attempted to link environmental covariates to skua survival and an

environmental effect was only detected on juvenile survival (Pacoureau et al. 2019b). We possibly

did not detect an effect of SAM because skuas showed high inter-individuals variations in

wintering area (Delord et al. 2018) making difficult to estimate the effects of SAM on

oceanographic conditions in such diversified areas. Nevertheless, even if skuas appear unaffected

by environmental conditions, we may assume an indirect effect of the environment through

bottom-up mechanisms in this skua-petrel system. Indeed, poor conditions at sea altered the

breeding success of petrels resulting in lower abundance of adults and chicks of the main prey of

skua. Since the breeding success of skuas depended on the abundance of petrels, as shown in our

results, it might be indirectly affected by the environmental conditions.

Bottom up processes affect breeding parameters

Overall, we highlighted a bottom-up effect in the dynamics of this marine predator-prey system,

i.e., the dynamics of these two seabirds was mostly driven by food availability. Petrel dynamics

were more strongly affected by environmental covariates near to their trophic level and the number

of petrels impacted the dynamics of skuas. We found no evidence of top-down impact, *i.e.,*

predation effects, in this system, although these two mechanisms have been found to jointly affect

ecosystems (Hairston et al. 1960, Hunter and Price 1992, Sinclair et al. 2003). Only breeding

parameters were affected by climate or intra- and interspecific relationships whereas the survival

did not seem to be impacted. In long-lived iteroparous species, environmental disturbance such as

a decreased in food availability is known to affect mainly the breeding processes whereas the

survival tends to remain constant (Pons and Migot 1995, Benton and Grant 1996, Oro et al. 1999).

Conclusion

Our multispecies IPM provided estimates of survival, breeding success and population size for two

seabird species. It had the advantage of using the population sizes estimated by the model for one

species to analyze its effect on the demographic of the other species while propagating all sources

of uncertainty. Hence, it allowed us to understand the contribution of interspecific interactions on

the demographic parameters while taking into account the effects of climatic conditions. The

estimated interspecific interactions revealed an effect of prey density on predator breeding

parameters, while predator density had no effect on prey parameters, showing bottom-up predator-

prey dynamics. Environmental effects were found on prey parameters but not on predator

parameters, although they may have indirect effects by propagating to top predator dynamics

through its prey. It highlights the crucial need of taking simultaneously into account both species

interactions and environmental covariates to understand species dynamics in the face of

environmental changes.

Acknowledgments

This study was made possible thanks to all the field workers involved in the monitoring programs

on Brown skuas and Blue petrels since 1985 at Mayes Island, Kerguelen Islands. These monitoring

programs were supported financially and logistically by the French Polar Institute IPEV (program

109, resp. H. Weimerskirch), the Zone Atelier Antarctique (CNRS-INEE), Terres Australes et

Antarctiques Françaises. All work was carried out in accordance with the guidelines of the IPEV

ethics committee. We acknowledge D. Joubert for the management of the demographic database.

This research was funded by the French National Research Agency (grant ANR-16-CE02-0007).

Literature Cited

Barbraud, C., and K. Delord. 2006. Population census of blue petrels Halobaena caerulea at Mayes

Island, Iles Kerguelen. Antarctic Science 18:199–204.

Barbraud, C., V. Rolland, S. Jenouvrier, M. Nevoux, K. Delord, and H. Weimerskirch. 2012.

Effects of climate change and fisheries bycatch on Southern Ocean seabirds: a review. Marine

Ecology Progress Series 454:285–307.

Barbraud, C., and H. Weimerskirch. 2001. Emperor penguins and climate change. Nature
411:183–186.

Barbraud, C., and H. Weimerskirch. 2003. Climate and density shape population dynamics of a
marine top predator. Proceedings of the Royal Society B: Biological Sciences 270:2111–
2116.

Barbraud, C., and H. Weimerskirch. 2005. Environmental Conditions and Breeding Experience
Affect Costs of Reproduction in Blue Petrels. Ecology 86:682–692.

Barraquand, F., and O. Gimenez. 2019. Integrating multiple data sources to fit matrix population
models for interacting species. Ecological Modelling 411.

Barraquand, F., T. T. Høye, J.-A. Henden, N. G. Yoccoz, O. Gilg, N. M. Schmidt, B. Sittler, and
R. A. Ims. 2014. Demographic responses of a site-faithful and territorial predator to its
fluctuating prey: long-tailed skuas and arctic lemmings. Journal of Animal Ecology 83:375–
387.

Benton, T. G., and A. Grant. 1996. How to keep fit in the real world: Elasticity analyses and
selection pressures on life histories in a variable environment. The American Naturalist
147:115–139.

Besbeas, P., S. N. Freeman, B. J. T. Morgan, and E. A. Catchpole. 2002. Integrating Mark-
Recapture-Recovery and Census Data to Estimate Animal Abundance and Demographic
Parameters. Biometrics 58:540–547.

Boersma, P. D., and N. T. Wheelwright. 1979. Egg Neglect in the Procellariiformes: Reproductive
Adaptations in the Fork-Tailed Storm-Petrel. The Condor 81:157–165.

Brooks, S. P., and A. Gelman. 1998. General Methods for Monitoring Convergence of Iterative
Simulations. Journal of Computational and Graphical Statistics 7:434–455.

Cairns, D. K. 1988. Seabirds as Indicators of Marine Food Supplies. Biological Ocenaography
5:261–271.

Carneiro, A. P. B., A. Manica, W. Z. Trivelpiece, and R. A. Phillips. 2015. Flexibility in foraging
strategies of Brown Skuas in response to local and seasonal dietary constraints. Journal of
Ornithology 156:625–633.

Caswell, H. 2001. Matrix Population Models: Construction, Analysis, and Interpretation. Sinauer
Associates, Sunderland, Massachusetts, USA.

Chastel, O., H. Weimerskirch, and P. Jouventin. 1995. Influence of Body Condition on
Reproductive Decision and Reproductive Success in the Blue Petrel. The Auk 112:964–972.

Chaurand, T., and H. Weimerskirch. 1994a. The regular alternation of short and long foraging trips
in the blue petrel Halobaena caerulea: a previously undescribed strategy of food provisioning
in a pelagic seabird. The Journal of Animal Ecology 63:275–282.

Chaurand, T., and H. Weimerskirch. 1994b. Incubation routine, body mass regulation and egg
neglect in the Blue Petrel Halobaena caerulea. Ibis 136:285–290.

Cherel, Y., P. Bocher, C. Trouvé, and H. Weimerskirch. 2002. Diet and feeding ecology of blue
petrels Halobaena caerulea at Iles Kerguelen, Southern Indian Ocean. Marine Ecology

Progress Series 228:283–299.

Cherel, Y., M. Connan, A. Jaeger, and P. Richard. 2014. Seabird year-round and historical feeding

ecology: blood and feather δ13C and 1315N values document foraging plasticity of small

sympatric petrels. Marine Ecology Progress Series 505:267–280.

Cherel, Y., P. Quillfeldt, K. Delord, and H. Weimerskirch. 2016. Combination of At-Sea Activity,

Geolocation and Feather Stable Isotopes Documents Where and When Seabirds Molt.

Frontiers in Ecology and Evolution 4.

Chu, C., and P. B. Adler. 2015. Large niche differences emerge at the recruitment stage to stabilize

grassland coexistence. Ecological Monographs 85:373–392.

Daunt, F., V. Afanasyev, A. Adam, J. P. Croxall, and S. Wanless. 2007. From cradle to early grave:

Juvenile mortality in European shags Phalacrocorax aristotelis results from inadequate

development of foraging proficiency. Biology Letters 3:371–374.

Delord, K., Y. Cherel, C. Barbraud, O. Chastel, and H. Weimerskirch. 2018. High variability in

migration and wintering strategies of brown skuas (Catharacta antarctica lonnbergi) in the

Indian Ocean. Polar Biology 41:59–70.

Dhondt, A. A., B. Kempenaers, and F. Adriaensen. 1992. Density-dependent clutch size caused by

habitat heterogeneity. Journal of Animal Ecology 61:643–648.

Fay, R., H. Weimerskirch, K. Delord, and C. Barbraud. 2015. Population density and climate shape

early-life survival and recruitment in a long-lived pelagic seabird. Journal of Animal Ecology

84:1423–1433.

Ferrer, M., and J. A. Donazar. 2015. Density-Dependent Fecundity by Habitat Heterogeneity in an

Increasing Population of Spanish Imperial Eagles. Ecology 77:69–74.

Frederiksen, M., M. Edwards, A. J. Richardson, N. C. Halliday, and S. Wanless. 2006. From

plankton to top predators: bottom-up control of a marine food web across four trophic levels.

Journal of Animal Ecology 75:1259–1268.

Gervasi, V., E. B. Nilsen, and J. D. C. Linnell. 2015. Body mass relationships affect the age

structure of predation across carnivore-ungulate systems: A review and synthesis. Mammal

Review 45:253–266.

Gille, S. T. 2002. Warming of the Southern Ocean since the 1950s. Science 295:1275–1277.

Gimenez, O., J. D. Lebreton, J. M. Gaillard, R. Choquet, and R. Pradel. 2012. Estimating

demographic parameters using hidden process dynamic models. Theoretical Population

Biology 82:307–316.

Godfray, H. C. J., and R. M. May. 2014. Open questions: are the dynamics of ecological

communities predictable? BMC Biology 12.

Grande, J. M., D. Serrano, G. Tavecchia, M. Carrete, O. Ceballos, R. Díaz-Delgado, J. L. Tella,

and J. A. Donázar. 2009. Survival in a long-lived territorial migrant: effects of life-history

traits and ecological conditions in wintering and breeding areas. Oikos 118:580–590.

Greig, S. A., J. C. Coulson, and P. Monaghan. 1983. Age-Related differences in foraging success

in the herring gull (Larus Argentatus). Anim. Behav. 31:1237–1243.

Grosbois, V., O. Gimenez, J.-M. Gaillard, R. Pradel, C. Barbraud, J. Clobert, A. P. Møller, and H.

Weimerskirch. 2008. Assessing the impact of climate variation on survival in vertebrate

populations. Biological Reviews 83:357–399.

Guinet, C., O. Chastel, M. Koudil, J. P. Durbec, and P. Jouventin. 1998. Effects of warm sea-

surface temperature anomalies on the blue petrel at the Kerguelen Islands. Proceedings of the

Royal Society B: Biological Sciences 265:1001–1006.

Hairston, N. G., F. E. Smith, and L. B. Slobodkin. 1960. Community Structure, Population Control,

and Competition. The American Naturalist 94:421–425.

Hamer, K. C., R. W. Furness, and R. W. G. Caldow. 1991. The effects of changes in food

availability on the breeding ecology of great skuas Catharacta skua in Shetland. Journal of

Zoology 223:175–188.

Han, W., J. Vialard, M. J. McPhaden, T. Lee, Y. Masumoto, M. Feng, and W. P. M. de Ruijter.

2014. Indian Ocean Decadal Variability: A Review. Bull. Amer. Meteor. Soc. 95:1679–1703.

Hazen, E. L., B. Abrahms, S. Brodie, G. Carroll, M. G. Jacox, M. S. Savoca, K. L. Scales, W. J.

Sydeman, and S. J. Bograd. 2019. Marine top predators as climate and ecosystem sentinels.

Frontiers in Ecology and the Environment 17:565–574.

Higgins, P. J., and S. J. J. F. Davied (editors). 1996. Handbook of Australia, New Zealand &

Antarctic Birds. Vol. 3, Snipe to pigeons. Oxford University Press.

Hoegh-Guldberg, O., and J. F. Bruno. 2010. The Impact of Climate Change on the World's Marine

Ecosystems. Science 328:1523–1528.

Holland, M. M., L. Landrum, Y. Kostov, and J. Marshall. 2017. Sensitivity of Antarctic sea ice to

665  the Southern Annular Mode in coupled climate models. Climate Dynamics 49:1813–1831.

666 Hunter, M. D., and P. W. Price. 1992. Playing chutes and ladders: heterogeneity and the relative

667  roles of bottom-up and top-down forces in natural communities. Ecology 73:724–732.

668 Jenouvrier, S. 2013. Impacts of climate change on avian populations. Global Change Biology

669  19:2036–2057.

670 Jenouvrier, S., C. Barbraud, and H. Weimerskirch. 2003. Effects of climate variability on the

671  temporal population dynamics of southern fulmars. Journal of Animal Ecology 72:576–587.

672 Jouventin, P., J. L. Mougin, J.-C. Stahl, and H. Weimerskirch. 1985. Comparative biology of the

673  burrowing petrels of the Crozet Islands. Notornis 32:157–220.

674 Kéry, M., and M. Schaub. 2011. Bayesian Population Analysis Using WinBUGS. Page Bayesian

675  Population Analysis Using WinBUGS. Academic Press.

676 Krüger, O., and J. Lindström. 2001. Habitat heterogeneity affects population growth in goshawk

677  Accipiter gentilis. Journal of Animal Ecology 70:173–181.

678 Lauria, V., M. J. Attrill, A. Brown, M. Edwards, and S. C. Votier. 2013. Regional variation in the

679  impact of climate change: Evidence that bottom-up regulation from plankton to seabirds is

680  weak in parts of the Northeast Atlantic. Marine Ecology Progress Series 488:11–22.

681 Lovenduski, N. S., and N. Gruber. 2005. Impact of the Southern Annular Mode on Southern Ocean

682  circulation and biology. Geophysical Research Letters 32.

683 Marshall, G. J. 2003. Trends in the Southern Annular Mode from Observations and Reanalyses.

684  Journal of climate 16:4134–4143.

Miller, D. A. W., E. H. C. Grant, E. Muths, S. M. Amburgey, M. J. Adams, M. B. Joseph, J. H.

Waddle, P. T. J. Johnson, M. E. Ryan, B. R. Schmidt, D. L. Calhoun, C. L. Davis, R. N.

Fisher, D. M. Green, B. R. Hossack, T. A. G. Rittenhouse, S. C. Walls, L. L. Bailey, S. S.

Cruickshank, G. M. Fellers, T. A. Gorman, C. A. Haas, W. Hughson, D. S. Pilliod, S. J. Price,

A. M. Ray, W. Sadinski, D. Saenz, W. J. Barichivich, A. Brand, C. S. Brehme, R. Dagit, K.

S. Delaney, B. M. Glorioso, L. B. Kats, P. M. Kleeman, C. A. Pearl, C. J. Rochester, S. P. D.

Riley, M. Roth, and B. H. Sigafus. 2018. Quantifying climate sensitivity and climate-driven

change in North American amphibian communities. Nature Communications 9.

Miller, T. E. X., and V. H. W. Rudolf. 2011. Thinking inside the box : community-level

consequences of stage-structured populations. Trends in Ecology and Evolution 26:457–466.

Millon, A., S. J. Petty, B. Little, O. Gimenez, T. Cornulier, and X. Lambin. 2014. Dampening prey

cycle overrides the impact of climate change on predator population dynamics: a long-term

demographic study on tawny owls. Global Change Biology 20:1770–1781.

Morosinotto, C., R. L. Thomson, and E. Korpimäki. 2010. Habitat selection as an antipredator

behaviour in a multi-predator landscape: All enemies are not equal. Journal of Animal

Ecology 79:327–333.

Mougeot, F., and V. Bretagnolle. 2000a. Predation risk and moonlight avoidance in nocturnal

seabirds. Journal of Avian Biology 31:376–386.

Mougeot, F., and V. Bretagnolle. 2000b. Predation as a cost of sexual communication in nocturnal

seabirds: an experimental approach using acoustic signals. Animal Behaviour 60:647–656.

Mougeot, F., F. Genevois, and V. Bretagnolle. 1998. Predation on burrowing petrels by the brown

skua (Catharacta skua lönnbergi) at Mayes Island, Kerguelen. Journal of Zoology 244:429–

438.

Mueller, A.-K., N. Chakarov, H. Heseker, and O. Krüger. 2016. Intraguild predation leads to

cascading effects on habitat choice, behaviour and reproductive performance. Journal of

Animal Ecology 85:774–784.

Oken, K. L., and T. E. Essington. 2015. How detectable is predation in stage-structured

populations? Insights from a simulation-testing analysis. Journal of Animal Ecology 84:60–

70.

Oro, D., N. Hernández, L. Jover, and M. Genovart. 2014. From recruitment to senescence: food

shapes the age-dependent pattern of breeding performance in a long-lived bird. Ecology

95:446–457.

Oro, D., A. Martínez-Abraín, M. Paracuellos, J. C. Nevado, and M. Genovart. 2006. Influence of

density dependence on predator-prey seabird interactions at large spatio-temporal scales.

Proceedings of the Royal Society B: Biological Sciences 273:379–383.

Oro, D., R. Pradel, and J.-D. Lebreton. 1999. Food availability and nest predation influence life

history traits in Audouin's gull, Larus audouinii. Oecologia 118:438–445.

Pacoureau, N., M. Authier, K. Delord, and C. Barbraud. 2019a. Population response of an apex

Antarctic consumer to its prey and climate fluctuations. Oecologia 189:279–291.

Pacoureau, N., K. Delord, S. Jenouvrier, and C. Barbraud. 2019b. Demographic and population

responses of an apex predator to climate and its prey: a long-term study of South Polar Skuas.

Ecological Monographs 89.

Pacoureau, N., E. Gaget, K. Delord, and C. Barbraud. 2019c. Prey remains of brown skua is

evidence of the long-term decline in burrow occupancy of blue petrels and thin-billed prions

at Mayes Island, Kerguelen. Polar Biology 42:1873–1879.

Parmelee, D. F., and P. J. Pietz. 1987. Philopatry, Mate and Nest-Site Fidelity in the Brown Skuas

of Anvers Island, Antartica. The Condor 89:916–919.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual

Review of Ecology, Evolution, and Systematics 37:637–669.

Péron, G., and D. N. Koons. 2012. Integrated modeling of communities: parasitism, competition,

and demographic synchrony in sympatric ducks. Ecology 93:2456–2464.

Piatt, J. F., A. M. A. Harding, M. Shultz, S. G. Speckman, T. I. van Pelt, G. S. Drew, and A. B.

Kettle. 2007. Seabirds as indicators of marine food supplies: Cairns revisited. Marine Ecology

Progress Series 352:221–234.

van de Pol, M., Y. Vindenes, B.-E. Sæther, S. Engen, B. J. Ens, K. Oosterbeek, and J. M.

Tinbergen. 2010. Effects of climate change and variability on population dynamics in a long-

lived shorebird. Ecology 91:1192–1204.

Pons, J.-M., and P. Migot. 1995. Life-history strategy of the herring gull: changes in survival and

fecundity in a population subjected to various feeding conditions. The Journal of Animal

Ecology 64:592–599.

Pradel, R. 2005. Multievent: An Extension of Multistate Capture-Recapture Models to Uncertain

States. Biometrics 61:442–447.

Quillfeldt, P., H. Weimerskirch, K. Delord, and Y. Cherel. 2020. Niche switching and leapfrog

foraging: movement ecology of sympatric petrels during the early breeding season.

Movement Ecology 8.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria.

Ratcliffe, N., and R. W. Furness. 1999. The effect of parental age and experimentally manipulated

brood size on the foraging effort and breeding performance of great skuas (Catharacta skua).

Journal of Zoology 249:195–201.

Saunders, S. P., F. J. Cuthbert, and E. F. Zipkin. 2018. Evaluating population viability and efficacy

of conservation management using integrated population models. Journal of Applied Ecology

55:1380–1392.

Sauser, C., K. Delord, and C. Barbraud. 2018. Increased sea ice concentration worsens fledging

condition and juvenile survival in a pagophilic seabird , the snow petrel. Biology Letters 14.

Schaub, M., and F. Abadi. 2011. Integrated population models: a novel analysis framework for

deeper insights into population dynamics. Journal of Ornithology 152:227–237.

Sinclair, A. R. E., S. Mduma, and J. S. Brashares. 2003. Patterns of predation in a diverse predator-

prey system. Nature 425:288–290.

Stenseth, N. C., J. M. Durant, M. S. Fowler, E. Matthysen, F. Adriaensen, N. Jonzén, K. Chan, H.

Liu, J. De Laet, B. C. Sheldon, M. E. Visser, and A. A. Dhondt. 2015. Testing for effects of

climate change on competitive relationships and coexistence between two bird species.

Proceedings of the Royal Society B: Biological Sciences 282.

Stoessel, M., B. Elmhagen, M. Vinka, P. Hellström, and A. Angerbjörn. 2019. The fluctuating

world of a tundra predator guild: bottom-up constraints overrule top-down species

interactions in winter. Ecography 42:488–499.

Sydeman, W. J., E. Poloczanska, T. E. Reed, and S. A. Thompson. 2015. Climate change and

marine vertebrates. Science 350:772–777.

Thomas, C. D., A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F.

N. Erasmus, M. Ferreira de Siquiera, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A. S.

van Jaarsveld, G. F. Midgley, L. Miles, M. A. Ortega-Huerta, A. Townsend Peterson, O. L.

Phillips, and S. E. Williams. 2004. Extinction risk from climate change. Nature 427:145–148.

Tuljapurkar, S., and H. Caswell. 1997. Structured-Population Models in Marine, Terrestrial, and

Freshwater Systems. Chapman & Hall, New York.

de Valpine, P., D. Turek, C. J. Paciorek, C. Anderson-Bergman, D. T. Lang, and R. Bodik. 2017.

Programming With Models: Writing Statistical Algorithms for General Model Structures

With NIMBLE. Journal of Computational and Graphical Statistics 26:403–413.

Walther, G., E. Post, P. Convey, A. Menzel, C. Parmesan, T. J. C. Beebee, J. Fromentin, O. Hoegh-

Guldberg, and F. Bairlein. 2002. Ecological responses to recent climate change. Nature

416:389–395.

Warham, J. 1990. The petrels: their ecology and breeding systems. Academic Press, London, UK.

Weimerskirch, H., O. Chastel, L. Ackermann, T. Chaurand, F. Cuenot-Chaillet, X. Hindermeyer,

and J. Judas. 1994. Alternate long and short foraging trips in pelagics seabird parents. Anim.

Behav. 47:472–476.

Tables

Table 1: Summary of the demographic parameters and their specificities (year random effect or state dependence) for the two species: the skua and the petrel. Notations are $\widetilde{NB}$: Nonbreeder the previous year, $\widetilde{B}$: Breeder the previous year, NB: Nonbreeder, FB: Failed Breeder, SB1: Successful Breeder with one fledged chick, SB2: Successful Breeder with two fledged chicks, FBE: Failed Breeder at the Egg stage, FBC: Failed Breeder at the Chick stage and SB: Successful Breeder.

Species	Parameter	Year random effect	State dependence
Skua	Adult apparent survival ϕ_{ad_s}	✓	$\widetilde{NB_s}\vert\,\widetilde{B_s}$
	Breeding probability β_s	✓	$\widetilde{NB_s}\vert\,\widetilde{B_s}$
	Breeding success γ_s	✓	$\widetilde{NB_s}\vert\,\widetilde{B_s}$
	Breeding success 2 chicks δ_s	✓	$\widetilde{NB_s}\vert\,\widetilde{B_s}$
	Detection probability p_s	✓	$\widetilde{NB_s}\vert\,\widetilde{B_s}$
	Uncertain state assignment probability u_s	✗	$NB_s\,\vert\,FB_s\,\vert\,SB1_s\vert\,SB2_s$
Petrel	Adult apparent survival ϕ_{ad_p}	✓	$\widetilde{NB_p}\vert\,\widetilde{B_p}$
	Breeding probability β_p	✓	$\widetilde{NB_p}\vert\,\widetilde{B_p}$
	Hatching success ω_p	✓	$\widetilde{NB_p}\vert\,\widetilde{B_p}$
	Breeding success γ_p	✓	$\widetilde{NB_p}\vert\,\widetilde{B_p}$
	Detection probability p_p	✓	$\widetilde{NB_p}\vert\,\widetilde{B_p}$
	Uncertain state assignment probability u_p	✗	$NB_p\vert\,FBE_p\vert\,FBC_p\vert\,SB_p$

798   Table 2: Summary of the covariates tested on the demographic parameters of the two species – the

799   skua and the petrel – and the time period (in months) considered for each demographic parameter.

800   Notation are PP: Predator-Prey interactions, DD: intraspecific density dependence, SAM:

801   Southern Annual Mode, SSTa: Sea Surface Temperature Anomalies, Chla: Chlorophyll a

802   concentration.

Seabird	Parameter	Covariates tested	Time period
Skua	Adult apparent survival ϕ_{ad_s}	DD|SAM	Wintering (March-September)
	Breeding probability β_s	DD|SAM	Pre-Breeding (July-November)
	Breeding success γ_s	PP|DD|SAM	Breeding (October-February)
	Breeding success 2 chicks δ_s	PP|DD|SAM	Breeding (October-February)
Petrel	Adult apparent survival ϕ_{ad_p}	PP|DD|SAM|SSTa|Chla	Wintering (February-September)
	Breeding probability β_p	DD|SAM|SSTa|Chla	Pre-Breeding (August-October)
	Hatching success ω_p	PP|DD|SAM|SSTa|Chla	Breeding (October – December)
	Breeding success γ_p	PP|DD|SAM|SSTa|Chla	Breeding (December-January)

Table 3: Regression coefficients estimates for the relationships between covariates (DD: density-
dependence, PP: predator-prey interaction, SAM: southern annular mode, SSTa: sea surface
temperature anomalies, Chla: chlorophyll a concentration) and demographic parameters (ϕad:
adult apparent survival, β: breeding probability, γ: breeding success, δ: breeding success with two
chicks, ω: hatching success) for skuas (top) and petrels (bottom), B: breeder or NB: non breeder
the previous years. 95% credible intervals that do not include 0 are in bold.

SKUA	DD				PP				SAM			
Parameters	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%
$\phi ad_s B$	-0.09	0.10	-0.24	0.12					0.31	0.31	-0.29	0.97
$\phi ad_s NB$	-0.07	0.17	-0.30	0.34					-1.19	0.78	-2.89	0.27
$\beta_s B$	-0.05	0.15	-0.25	0.35					-0.06	0.33	-0.72	0.61
$\beta_s NB$	-0.10	0.22	-0.47	0.32					-0.18	0.55	-1.23	0.94
$\gamma_s B$	**-0.56**	**0.11**	**-0.77**	**-0.34**	**0.76**	**0.18**	**0.41**	**1.14**	-0.05	0.23	-0.50	0.40
$\gamma_s NB$	-0.33	0.35	-0.95	0.46	0.67	0.53	-0.42	1.73	-0.24	0.68	-1.70	1.00
$\delta_s B$	**-0.78**	**0.19**	**-1.10**	**-0.38**	**1.14**	**0.31**	**0.57**	**1.76**	0.60	0.33	-0.04	1.26
$\delta_s NB$	-0.17	0.64	-1.08	1.03	-0.25	1.64	-3.53	2.38	4.09	3.99	-3.00	13.0

PETREL	DD				PP				SAM				SSTa				Chla			
Parameters	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%	slope	sd	2.5%	97.5%
$\phi ad_p B$	-0.30	0.28	-0.91	0.21	0.21	0.27	-0.21	0.75	-0.08	0.45	-0.95	0.86	-0.10	0.25	-0.65	0.36	-0.35	0.51	-1.40	0.62
$\phi ad_p NB$	-1.32	0.90	-2.62	0.70	0.70	0.35	-0.10	1.12	0.51	1.92	-2.65	5.00	0.13	0.96	-1.63	2.35	-0.36	1.47	-2.70	3.37
$\beta_p B$	**0.71**	**0.12**	**0.44**	**0.90**					0.11	0.25	-0.37	0.63	-0.23	0.22	-0.69	0.17	0.44	0.42	-0.36	1.30
$\beta_p NB$	0.21	0.31	-0.44	0.73					0.10	0.27	-0.40	0.66	-0.08	0.20	-0.47	0.32	0.71	0.44	-0.24	1.56
$\omega_p B$	0.40	0.30	-0.29	1.01	0.01	0.20	-0.40	0.39	-0.43	0.39	-1.24	0.30	-0.43	0.32	-1.10	0.16	0.47	0.37	-0.22	1.23
$\omega_p NB$	0.21	0.36	-0.40	1.14	-0.29	0.22	-0.72	0.16	**-1.03**	**0.42**	**-1.93**	**-0.27**	**-1.16**	**0.39**	**-1.99**	**-0.47**	**1.19**	**0.43**	**0.43**	**2.11**
$\gamma_p B$	-0.33	0.48	-1.43	0.49	0.03	0.28	-0.42	0.63	0.15	0.48	-0.73	1.25	-0.04	0.32	-0.67	0.65	-0.18	0.46	-1.17	0.67
$\gamma_p NB$	-0.07	0.49	-1.04	0.83	0.08	0.34	-0.56	0.83	-0.10	0.64	-1.30	1.00	-0.16	0.58	-1.31	0.84	0.70	0.80	-0.50	2.85

Figures

Figure 1: Structure of the skua IPM. Squares represent the state variables, circles represent the parameters. Two types of data are used, capture-recapture (CR) data and counts data (N_{counts_s}). Adult apparent survival (ϕ_{ad_s}), breeding probability (β_s), breeding success with at least one chick (γ_s), breeding success with two chicks (δ_s) for successful breeders, juvenile apparent survival for one to four years old and older (ϕ_{J1_s} to ϕ_{J4_s}), probability of first reproduction (Pr_s), state assignment probability (u_s) and detection probability (p_s) are parameters estimated in the model. Fecundity of breeder with one $\left(f_{SB_s}\right)$ or two chicks (f_{SB_s}) are fixed. The number of adults (N_{ad_s}), dead (N_{dead_s}), alive (N_{alive_s}), breeder (B_s), nonbreeder (NB_s), failed breeder (FB_s), successful breeder (SB_s), successful breeder with one chick ($SB1_s$) or with two chicks ($SB2_s$) and the number of juveniles of one year old to four years old and older ($J1_s$ to $J4_s$) are state variables estimated by the model. The number of immigrants (N_{im_s}) is a fixed vector.

Skua

Petrel

Figure 2: State variables and covariates effects on adult demographic parameters for the skua (top

panels) and the petrel (bottom panels). Solid lines represent the estimated relationship between the

covariates and the demographic parameters. Shaded areas are the 50% and 95% credibility

intervals. Points represent demographic parameter estimates each year (21 years) plotted with

respect to the covariate. Error bars are standard deviation. (a) Density-dependence effect and (b)

prey effect on the estimated breeding success probability for skuas that bred the previous breeding

season. (c) Density-dependence effect and (d) prey effect on the estimated breeding success

probability for two chicks for skuas that bred the previous breeding season. Effects of (e) Southern

Annular Mode, (f) Sea surface temperature anomalies and (g) Chlorophyll a concentration on the

estimated hatching success probability for petrels that were nonbreeders during the previous

breeding season. (h) Density-dependence effect on the breeding probability for petrels that bred

the previous breeding season.