1

Plant cell wall polysaccharides induce colony expansion of soil-derived

Flavobacterium spp.

Judith Kraut-Cohen[a], Orr H.Shapiro[b], Barak Dror[ac] Eddie Cytryn[a]

[a]Institute of Soil, Water and Environmental Sciences, Agricultural Research

Organization, Volcani Center, P.O Box 15159, Rishon LeZion 7505101, Israel

[b]Institute of Postharvest and Food Sciences, Agricultural Research Organization,

Volcani Center, P.O Box 15159, Rishon LeZion 7505101, Israel

[c]Department of Plant Pathology and Microbiology, The R.H. Smith Faculty of

Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot,

Israel

Running title: Pectin induces Flavobacteria colony expansion

#Address correspondence to Eddie Cytryn, eddie@volcani.agri.gov.il.

*Present address: Agricultural Research Organization - the Volcani Center,

68 HaMaccabim Road , P.O.B 15159 Rishon LeZion 7505101, Israel.

Summary

Flavobacterium is a genus, belonging to the *Bacteriodetes* phylum, characterized by a unique gliding motility. They are often abundant in root microbiomes of various plants, but the factors contributing to this high abundance are currently unknown. In this study, we evaluated the effect of various plant-associated poly- and mono-saccharides on colony expansion of two *Flavobacterium* strains. Both strains were able to grow on pectin and other polysaccharides such as microcrystalline cellulose. However, only pectin, a major component of plant cell walls, substantially enhanced colony expansion on solid surfaces in a dose- and substrate-dependent manner (but did not occur on pectin monomers). On pectin, flavobacteria exhibited a bi-phasic behavior, with an initial phase of rapid expansion, followed by growth within the colonized area. Proteomic and gene expression analyses revealed significant induction of carbohydrate metabolism related proteins when flavobacteria were grown on pectin, including selected SusC/D, TonB-dependent glycan transport operons. Our results suggest an unknown linkage between specific glycan associated operons and flavobacterial colony expansion. This may be associated with their capacity to rapidly glide along the root and metabolize plant cell wall carbohydrates, characteristics that are crucial to rhizosphere competence.

Keywords: Flavobacterium, Roots, Rhizosphere, Pectin, TonB

Originality-Significance Statement: This study reveals unique data linking plant glycan metabolism and bacterial motility, providing insight into bacterial-root associations and rhizosphere competence. Specifically, it explores mechanisms

associated with pectin-stimulated colony expansion in root-associated *Flavobacterium* strains. We determined that expansion of colonies on pectin was biphasic in nature, characterized by rapid proliferation followed by biomass accumulation. We demonstrate by proteomic and gene expression analyses that expansion of *Flavobacterium* on pectin strongly induces TonB related transporters, which seemingly play a role in motility in addition to the uptake and metabolism of plant-associated glycans.

Introduction

The complex interactions between plant-associated microorganisms and their hosts (collectively referred to as the "plant holobiont") are crucial for plant health and growth (1–3). Plants modulate the narrow region of soil that is influenced by root secretions, the rhizosphere, by exuding various small molecular weight compounds (rhizodeposites) such as sugars, amino acids and organic acids, by rhizodepositing root cap border cells, and by releasing various mono- and poly-saccharides in their mucilage (3–8). Organic compounds released from the roots transform the rhizosphere into a nutrient rich environment relative to the surrounding bulk soil, which facilitates colonization by soil microorganisms. Rapid growth, coupled to competition for root resources, results in less diverse rhizosphere and rhizoplane microbiomes relative to surrounding bulk soil, a phenomenon known as the "rhizosphere effect" (4, 9, 10). Rhizosphere and root colonizing bacteria have the capacity to outcompete other soil bacteria and thereby proliferate and thrive in the root ecosystem through a combination of specific traits collectively coined "rhizosphere competence", which include: motility, resistance to stress, ability to utilize plant-derived organic compounds, chemotaxis, and

the production of secondary metabolites (5). Some earlier studies demonstrated that

rhizosphere bacteria are attracted to the roots by plant-exuded organic acids such as

malic, citric or fumaric acid and various amino acids (11–15). Many of these sensed

chemo-attractants can also be consumed by the bacteria (16). Although rhizosphere

recruitment and colonization mechanisms of certain plant-growth-promoting

rhizobacteria (PGPR) have been identified and characterized, those responsible for

recruitment of the vast majority of rhizosphere and rhizoplane bacteria is currently an

enigma (17–19).

Flavobacterium is a Gram-negative genus of bacteria from the phylum *Bacteroidetes*

known to degrade complex organic compounds in freshwater, marine and soil

environments (20, 21). It is highly abundant in the rhizoplane of a wide array of plants,

in contrast to considerably lower abundance in bulk soil (21–28), suggesting that

members of this genus have specifically adapted to root ecosystems. Several root- and

soil-derived flavobacteria were found to antagonize various plant pathogens in different

crops (29–33), and recently it was discovered that entophytic *Flavobacterium* strains

play a fundamental role in antagonizing phytopathogenic fungi, through the production

chitinolytic enzymes and specific secondary metabolites (34). Furthermore, a recent

study that applied bacterial network analysis, indicated that they are potential drivers

of pathogen suppression in root ecosystems (35). In addition, selected members of this

genus have defined as plant growth promoting rhizobacteria (PGPR) of various crops

(23, 29, 30, 36–38).

Soil flavobacteria have specialized ability to decompose complex plant derived

polysaccharides, such as pectin and glucomannan and the ability to secret various

carbolytic enzymes via the *Bacteriodetes*-specific type IX secretion system (T9SS) (21,

39–43). Like other *Bacteroidetes*, they contain a myriad of genes that encode

Polysaccharide Utilization Loci (PULs) that are activated specifically to facilitate glycan capture and uptake (44). These PULs include outer membrane protein transducers involved in polysaccharide utilization, which are part of the TonB family, generally referred to as Starch Utilization System (SUS) proteins. Interestingly, comparative genomics revealed that genomes of soil and root-associated flavobacterial strains are significantly enriched with genes associated with plant polysaccharide degradation relative to aquatic strains from this genus, indicating that physiology of this genus is strongly influenced by its ecology (21).

Most terrestrial *Flavobacterium* strains possess a unique gliding mechanism that rapidly propels them over solid and semi-solid surfaces. The proteins that comprise this gliding system are molecularly intertwined with at least fifteen proteins that make up the T9SS, seven of whom are responsible for the secretion of SprB and RemA adhesins which are expressed on the cell surface and involved in gliding (45, 46). We previously demonstrated that this T9SS-gliding complex is crucial for root colonization by flavobacteria, and this colonization was positively linked to the induction of plant resistance to foliar pathogens (31).

Collectively, the above data strongly suggest that terrestrial flavobacterial strains have evolved means that enable them to interact with plant roots, and that these interactions are beneficial to plant health. Nonetheless, the specific mechanisms behind this phenomenon are currently unclear. In this study, we assessed the impact of various plant-derived sugars on the motility and growth dynamics of flavobacteria by coupling conventional petri dish assays and live-imaging fluorescent microscopy with proteomic and gene expression analyses. We demonstrate that pectin, a plant cell wall-associated polysaccharide, facilitates bi-phasic proliferation over solid surfaces through induction of specific TonB-associated glycan uptake operons. *In-planta* extrapolation of these

118 results, suggests that this link between pectin, motility and carbohydrate metabolism

119 may be fundamental to rhizosphere competence in flavobacteria.

Results

Growth of *Flavobacterium* strains on various carbon sources

The growth of flavobacteria was evaluated on minimal and rich media amended with

different carbon sources. *F. johnsoniae* was motile and grew on chitin, pectin and

microcrystalline cellulose but not on arabinose (**FigS1**). Subsequently, we evaluated

growth of flavobacteria on rich media (PY2 agar) coated with selected sugars (**Fig1A**)

to evaluate how specific carbon sources flavobacterial proliferation. Colony expansion

of *F. johnsoniae* on PY2 agar media coated with pectin was close to five times higher

than on the same media coated with other analyzed mono- and polysaccharides or with

double distilled water (DDW, control) ($p < 0.05$, Tukey-Kramer HSD test) (**Fig1B**).

Wild type (WT) and gliding/typeIX secretion system mutants (ΔgldJ) of *F. johnsoniae*

and the pepper root isolate *Flavobacterium* sp. F52 were inoculated in the center of

PY2 agar media amended with or without pectin (**Fig1C**). When grown on pectin, WT

colonies of both *Flavobacterium* strains significantly expanded after 48 hours of

incubation, while growth was reduced in the control without pectin (DDW). In contrast,

gliding mutant (ΔgldJ) colonies of both flavobacterial strains did not expand

(**Fig1C,D**), indicating that the gliding apparatus is a prerequisite for pectin-induced

colony expansion.

Dose-dependent pectin facilitated colony expansion

To determine whether expansion on pectin is dose dependent, *F. johnsoniae* and

Flavobacterium sp.F52 strains were inoculated at the center of PY2 agar media plates

streaked with pectin at final concentrations of 0.5, 1, 2 and 4%. For all the examined

pectin concentrations colonies radiated along the pectin streaks, but expansion was

more significant on 2% and 4% pectin ($p < 0.05$,Tukey-Kramer HSD test) (**FigS2**).

Since galacturonic acid and rhamnose are the two major components of pectins, we

examined the colony expansion of *F. johnsoniae* on PY2 agar coated with 2% titrated

galacturonic acid, 2% rhamnose, a combination of galacturonic acid and rhamnose, 2%

pectin and DDW. *F. johnsoniae* expansion on galacturonic acid, rhamnose (alone or

combined) did not facilitate significant colony expansion ($p < 0.05$,Tukey-Kramer HSD

test) in contrast to colonies that were grown on pectin (**Fig2A, B**).

Temporal dynamics of pectin induced *F. johnsoniae* colony expansion

The expansion of green fluorescent protein (GFP)-labeled *F. johnsoniae* on PY2 agar

coated with glucose, microcrystalline cellulose, glucomannan or pectin, was visualized

at a higher resolution using time-lapse microscopy. Colony morphology after 32 hours,

on each tested substance is presented in **Fig3A**.

Growth dynamics were clearly affected by the sugar type. Expansion on pectin was

characterized by a relatively long lag phase. However following this stage, colonies

rapidly expanded and after 18 hours, the fluorescent signal of colonies grown on pectin

surpassed both the control (without sugar amendment) and the other amended

compounds (according to Tukey-Kramer HSD test). Growth on pectin facilitated

multiple ring-like structures that resemble previously described "raft" structures (47)

(**Fig3B, Supplementary movie 1**). Conversely, expansion of colonies grown on

glucose and the control did not display this bi-phasic expansion (**Supplementary**

movie 1). Bacterial colonies grown on glucose expanded the least resulting in compact,

small colonies with less ringed structures, suggesting that glucose has an inhibitory

effect on *F. johnsoniae* motility and possibly growth as was previously demonstrated

(**Fig3C**) (48, 49). The most significant expansion observed between 6 to 20 hours was

on cellulose microcrystalline, but the colonies grown on glucomannan and pectin

proliferated at later times, and the colonies grown on pectin reached the greatest

intensity at 44 hours (**Fig3C**), thereby surpassing the colonies grown on other sugar

sources (p<0.05,Tukey-Kramer HSD test). The high sensitivity of the fluorescence

detection enabled us to visualize lateral colony expansion on cellulose microcrystalline,

which was less visible on agar plates (**Fig1A).**

Colonies grown on pectin were bi-phasic, with initial phase of rapid expansion,

followed by an enhanced growth phase where cells appeared to proliferate within the

expanded area (**Supplementary movie 1**). An initial peak in fluorescence occurred at

20 hours, and a second peak at 36 hours (**Fig3D**). After 20 hours of growth, total

fluorescence was highest in cells grown on pectin, glucomannan and microcrystalline

cellulose, and lowest on glucose (p<0.05,Tukey-Kramer HSD test).

Next, we estimated the velocity of colony expansion on the selected mono- and

polysaccharides by measuring the time it took the colonies to cross three radials (3mm,

6mm and 9mm), and subsequently calculating the mean velocity from circle to circle

(**Fig3E**). Initially, in the first 1.5mm radius the estimated colony expansion velocity

was similar on all substances, however, between 1.5-3mm, colony expansion on pectin

and glucomannan increased. Colony expansion velocity slowed slightly in the outer

circle with pectin, glucomannan still exhibiting the fastest expansion rates of all the

tested mono- and polysaccharides (**Fig3F**). Thus, *F. johnsoniae* expanded faster on

pectin than on any of other tested carbon sources (**Fig3G**). Collectively, we conclude

that growth of *F. johnsoniae* on pectin is bi-phasic. Following an initial lag phase, there was a rapid expansion phase, followed by a slower growth phase where cells appear to spread less and replicate more gaining biomass.

Specific TonB/Sus transducers are expressed in response to growth on pectin

In order to gain deeper insight into the molecular mechanisms associated with flavobacteria colony expansion on pectin, we conducted a proteomic assay in which we examined differential intracellular protein expression after 44 hr growth between *F. johnsoniae* growing on PY2 agar coated with pectin relative to the same rich media coated with DDW. Expression of eighty-three and forty-three proteins was higher and lower, respectively on pectin relative to the DDW control (**TableS1**). Genomic annotation with various tools revealed a high proportion of unassigned proteins, while a large fraction of the proteins (17%, 22% and 37% of KEGG, SEED and EggNog annotations, respectively) were associated with carbohydrate metabolism (**FigS3A-C**). Of the 25 most markedly pectin-induced proteins identified, 13 were involved in polysaccharide uptake, processing and metabolism, including four Sus C/D related proteins (**Fig4A**). Other pectin-induced proteins included a novel transcriptional regulator (12-fold higher on pectin) and a protein associated with auxin regulation (26-fold higher on pectin) (**Table S1**). Interestingly, none of the differentially synthesized proteins was gliding related (**Table S2**).

Of the 44 previously-described SusC and 42 SusD homologues identified in the *F. johnsoniae* genome (43), 27 SusC and 15 SusD proteins were detected in the proteomic analysis, in addition to 610 proteins encoding flanking genes surrounding these Sus proteins encoding genes that constitute PUL clusters seemingly associated with glycan metabolism. Of these, three SusC and six SusD proteins along with 31 associated PUL

genes encoding proteins (forming 4 gene clusters **Table S3**) were significantly induced

in response to growth on pectin in the proteomic analysis (**Table S4**).

In order to validate the proteomic results, *F. johnsoniae* cells were again grown on PY2

agar coated with pectin or DDW as described in the proteomic analysis, and the

expression of 8 genes (**Table S5**) were evaluated by quantitative real time PCR. Pectin

did not facilitate expression of remA (WP_012022896.1) encoding for an adhesion

protein involved in gliding motility (not evaluated in the proteomic analysis since it's

an extra-membranal protein), indicating that synthesis of this protein is not enhanced

on pectin at the tested time point. Expression of genes encoding for the novel

transcriptional regulator (Trans. Regul WP_012022876.1), pectate lyase (Pec lyase

WP_012026072.1), putative auxin-regulated annotated protein (Auxin

WP_012023111.1) and tonB 2144 (WP_012026069.1), were also not differentially

expressed on pectin-coated media despite the fact that they were significantly induced

on pectin in the proteomic analysis. Among the examined TonB/SusC related genes,

tonB 260 (WP_012026229.1) and sus73 (SHH12854.1) were significantly upregulated

(60-100 fold, $p < 0.05$ Tukey-Kramer HSD test) while TonB 445 (WP_012022294.1)

was substantially upregulated (10-20 fold) but not statistically significant (**Fig4B**).

tonB260 and sus73 are part of an 18-gene cluster, of which all of the encoded proteins

were upregulated in the presence of pectin in the proteomic analysis (**cluster 1-Table**

S3). Using two different prediction tools, the TonB260 and Sus73 encoding genes were

mapped to the same operon together with a gene encoding for the hydrolytic enzyme

polygalacturonase that cleaves the α (1-4) bonds between adjacent galacturonic acids

within the homogalacturonic acid backbone of pectin (**Table S6**).

Discussion

Roots modulate their microbiome by secreting exudates, mucilage and other components that can attract or deter bacteria (8, 47, 50), and specific root exudates were found to serve as chemoattractants for microbes that colonize roots (51). However, this seems to be the tip of the iceberg, and little is known about the plant-associated factors that recruit bacteria from the ecologically complex rhizosphere to the surface and endosphere of roots.

Bacterial colonization of plant roots is a multi-stage process that involves recognition, movement towards attractants and attachment (15). Once attached, persistence and propagation on plant roots is dependent on metabolic fitness (52, 53), resistance to plant protective mechanisms and inter-microbial interactions. The abundance of bacteria in rhizosphere soil is about 10^2 higher than in the rhizoplane and root endosphere (54, 55). This exemplifies the extreme selective pressure of this highly competitive environment and the fitness requirements of the root-colonizing microbiome.

Members of the genus *Flavobacterium* are highly abundant and substantially enriched (relative to the rhizosphere soil) on, and within roots of several plant species (25, 32, 33, 56, 57), and certain strains have been found to augment plant health and suppress pathogens in a variety of plants (29, 32, 33, 35). In this study, we explored the effect of plant-derived mono- and polysaccharides on colony expansion of flavobacteria, in attempt to elucidate mechanisms associated with plant root-flavobacterial interactions. It was previously shown that *F. johnsoniae* can effectively grow on polysaccharides such as pectin and glucomannan as single carbon sources (43). Collectively, results from this study demonstrate that flavobacteria not only metabolize complex plant cell wall-derived glycans such as pectin, glucomanan and microcrystalline cellulose, but

that these polysaccharides (particularly pectin) also facilitate rapid spreading of

flavobacteria over solid surfaces. We postulate that this plant cell wall glycan

recognition-uptake mechanism is critical for proliferation of flavobacteria on plant

roots.

Interestingly, our results demonstrated that even when carbon was not limited, plant

cell wall glycans and especially pectin, facilitated colony expansion of terrestrial

flavobacterial strains over solid surfaces. While the specific correlation between

expansion on pectin and the unique flavobacterial gliding motility mechanism was not

determined in this study, it is evident that the latter is required, because ΔgldJ mutants

did not expand on pectin. Propagation on pectin occurred in a concentration dependent

manner, and the response was specifically dependent on the mother compound, as the

response did not occur with bacteria grown on the pectin monomers D-galacturonic

acid and L-rhamnose. In contrast, growth on glucose was found to inhibit propagation

of flavobacteria as was previously demonstrated by others (48, 49). Specifically,

glucose was shown to inhibit colony spreading via MfsA, (49), which encodes a major

facilitator superfamily (MFS) transporter and creation of raft microstructures, but does

not inhibit gliding motility in general (47, 48).

Patterns and extent of *F. johnsoniae* colony expansion was dictated by the carbon

source supplemented to the growth media. The expansion of flavobacterial colonies on

rich media coated with plant-derived polysaccharides (pectin and glucomanan) was

rapid, patchy, non-uniform and sparse relative to colonies grown on rich media coated

with glucose or DDW. The rapid expansion of flavobacteria on pectin was strongly

linked to induction of TonB-related proteins, which undoubtedly assist in binding

(recognition), uptake and metabolism of this glucan. Therefore, we hypothesize that for

root-associated flavobacteria, pectin is not merely a carbon source but also an

environmental cue. To the best of our knowledge, this is the first time that a specific

root-associated factor was found to be specifically linked to flavobacterial spreading.

There are only a few examples of plant cell wall-associated polysaccharides that

facilitate root bacteria colonization and proliferation. Flexibacter FS-1 is not able to

glide on agarose (unless supplemented with cold water-soluble fraction of agar) but

can glide on agarose amended with 1% pectin, while galacturonic acid does not induce

a similar effect (58). Purified Arabidopsis polysaccharides (arabinogalactan, pectin, or

xylan) triggered biofilm and pellicle formation in *B. subtilis* when added to rich media

and induced root colonization (6). Similarly, addition of pectin and sucrose to the media

of the PGPR *Bacillus amyloliquefaciens* strain SQY 162, increased bacterial

abundance, induced biofilm formation and improved the ability of the amended

bacterium to suppress tobacco bacterial wilt disease (59). In the symbiotic nitrogen

fixing bacterium *Rhizobium leguminosarum*, glucomannan-mediated attachment was

important for legume infection and nodulation (12).

We did not observe any evidence of chemotactic behavior towards pectin, and therefore

postulate that propagation on pectin requires direct contact between the flavobacteria

and the glycan. This is logical given the large size and low solubility of this plant

polysaccaride. Based on proteomics and subsequent qPCR (quantitative real-time

PCR) gene expression validation, we believe that this phenomenon is dictated by the

induction of TonB-associated *F. johnsoniae* PULs, suggesting a link between niche

recognition, colony expansion and metabolic fitness. Similar induction of TonB and

PUL was observed in marine flavobacteria as response to phytoplankton blooms

characterized in decomposition of alga-derived organic matter (60).

Their high energetic cost and substrate specificity of TonB transducers explains why

the genes encoding them were induced by pectin and not constitutively expressed (61,

62). Nonetheless, the specific regulatory mechanism responsible for facilitating this

response and the subsequent linkage to the induced gliding motility in flavobacteria

remain an enigma. This is especially true in view of the fact that pectin did not induce

expression of remA, the lectin binding, flavobacterial cell surface adhesin involved in

gliding (63). The gliding machinery might be involved in the earlier phase of the

response to pectin, in which we observed intensive bacterial motility. Alternatively,

pectin might facilitate gliding in a protein expression-independent manner, or induce

other components of the gliding machinery not identified in the proteomic analyses.

Previous experiments showed that *Pseudomonas putida* mutants that lacked TonB,

were deficient in their capacity to uptake iron and displayed impaired seed colonization,

again linking TonB to metabolic and functional fitness (64). However, knock-out of

specific tonB genes might be less effective in *F. johnsoniae* due to the multitude of

predicted tonB genes in its genome, which suggests a high level of functional

redundancy (43). Similar functional redundancy was shown in proteomic analysis of

the soil bacterium *Cytophaga hutchinsonii* where multiple cellulolytic enzymes were

induced after growth on microcrystalline cellulose (65).

In *Xanthomonas campestris pv. Campestris*, pectate sensed by specific TonB-

dependent receptors triggered secretion of extracellular polygalacturonate. This

resulted in pectin degradation and generation of oligogalacturonides (OGA) that are

recognized as damage-associated molecular patterns (DAMPs), facilitating the

initiation of the plant defense mechanisms (66). In light of our results and the findings

in *X. campestris*, we speculate that sensing, uptake and degradation of pectin by TonB

and associated proteins, results in OGA release, and plant immune response activation.

This pathway may explain the biocontrol and improved plant disease resistance induced

by flavobacteria in various plant types. Further experiments will be needed to support

these assumptions.

Beside TonB related proteins, pectin induced additional protein targets. The up

regulation of auxin-regulated protein is especially interesting since auxin is a major

phytohormone responsible for plant growth and development demonstrating again a

possible connection between pectin sensing and flavobacterial-plant interactions. The

additional upregulation of the putative transcriptional regulator may suggest that pectin

induces a downstream response.

Gene expression (qPCR) analysis did not completely correlate with the proteomics

analysis, when comparing flavobacterial colonies growing on media amended with

pectin relative to the control without pectin. The discrepancy between mRNA and

protein expression might be explained by post-translational modifications or regulation

affecting protein stability, degradation and complex formation, as shown in similar

cases (67, 68).

Live-imaging fluorescent microscopy revealed that *F. johnsoniae* growth on PY2 media

coated with different mono- and polysaccarides was bi-phasic in nature, with an initial

phase of rapid expansion, followed by biomass production within the colonized area.

This bi-phasic growth pattern, which was most pronounced on pectin, resembles

previously described models in motile *E. coli*, which depicted an initial expansion

phase, where "pioneer" bacteria with high velocity advance in front of the colony,

followed by a second phase, where "settler" bacteria grow and replicate locally (16, 69).

Collectively, we hypothesize that plant-derived polysaccharides and especially pectin

may serve as signal that facilitates the expansion of "pioneer" cells, and later as carbon

sources that support growth of "settlers".

To summarize, we found that pectin, a prominent plant cell wall polysaccharide,

facilitates expansion of flavobacteria on solid surfaces, even in the presence of nutrient-

rich media. We postulate that pectin may be a "missing link" in elucidating the

mechanisms responsible for rhizosphere competence in flavobacteria, *i.e.* their capacity

to efficiently colonize and proliferate in the root niche. Thus, in the root environment,

plant cell wall polysaccharides, and specifically pectin, may not only serve as a nutrient

source for flavobacteria, but also as a potential cue for colonization and rapid expansion

along the root surface. The interaction between pectin (and potentially other root

glycans) and flavobacteria is mediated by induction of TonB/SusC operons and other

associated PULs that facilitate metabolism of pectin.

Experimental Procedures

Bacterial Strains and growth conditions

Flavobacterial strains targeted in this study included *F. johnsoniae* strain UW101

(ATCC17061), a gliding/secretion impaired (ΔgldJ) *F. johnsoniae* strain UW101,

Flavobacterium sp. F52, isolated from the roots of a greenhouse pepper (70, 71), and a

gliding/secretion impaired (ΔgldJ) *Flavobacterium* sp. F52 strain. For live imaging

microscopy (see below) the fluorescent strain *F. johnsoniae* UW101 WT+pAS43

(Flavo GFP) Flavo-ErytRFlavo-CefR, and the gliding/secretion impaired *F. johnsoniae*

UW102-48 (ΔgldJ)+pAS43 (Flavo GFP) Flavo-ErytR Flavo-CefR were used.

Erythromycin, 100 ug/ml was added to the media of the GFP labeled bacteria.

Flavobacteria strains were grown in CYE medium (Casitone, 10 mg/ml, yeast extract

at 5mg/ml, and 8 mM $MgSO_4$ in 10 mM Tris buffer [pH 7.6]) at 30°C, as previously

described (72). To observe colony-spreading, bacteria were grown on PY2 agar

medium (2 g peptone, 0.5 g yeast extract,10 mg/ml agar, pH 7.3) at 30°C (73).

Organic amendments to growth media

A suite of mono- and polysaccharides were amended to growth media in various

configurations (as described below), to evaluate colony spreading dynamics of the

selected flavobacterial strains. The following substances were used in this study: pectin

(Sigma P9135), arabinose (Sigma A9462), D(+)glucose (Fischer scientific 10373242),

microcrystalline cellulose (partially depolymerized cellulose synthesized from an α-

cellulose precursor, Merck 102331), D(-)arbinose (Acros 161451000), glucomannan

(Megazyme P-GLCML), chitin (Sigma C7170), L-rhamnose (Sigma 83650) and D(+)-

galacturonic acid monohydrate (Sigma 48280-5G-F). All substances were dissolved

and suspended to 2% final concentration in DDW. Glucose, arabinose, rhamnose and

galacturonic acid (titrated to pH 7) were dissolved and filtered through a 0.22-micron

filter. Pectin was dissolved in DDW heated at 80°C, and subsequently filtered through

0.45-micron filters. Chitin, microcrystalline cellulose, and glucomannan were mixed

with DDW heated to 50°C and then autoclaved for 20 min.

Flavobacterial growth on various sugars

The selected sugars were amended to the PY2 agar plates in two manners: (i) to test the

effect of specific carbon sources on the directional proliferation of flavobacteria, 10 µl

of the selected compounds were thinly applied along a line projecting outward from the

center of the petri dish using a pipetor (**Fig2S**); (ii) to test the effect of specific carbon

sources on the general proliferation of flavobacteria, 500µl were uniformly smeared

over the entire petri dish (**Fig1A, C, Fig2A**). Non-soluble substances such as

microcrystalline cellulose and chitin, were vigorously vortexed and dispensed using a

cut tip. Where two sugars were used (*i.e.* rhamnose+ galacturonic acid) we added 250µl

of each. In all cases, plates were left to dry overnight after adding organic amendments.

Flavobacteria were incubated on CYE media overnight, colonies were harvested and

subsequently diluted in 200ul saline to 0.6-1 OD ($5x10^9$-1.12 $x10^{10}$ cells), vortexed

well, and 2ul were spotted in the center of PY2 agar covered or streaked with the

selected sugars as indicated above. Plates were left to dry for 15min and then incubated

for 48 hours at 30^OC. The colony area or the length of expansion (in cm) were measured

using Fiji (74) and statistics was calculated using JMP®, Version Pro 14 (SAS Institute

Inc., Cary, NC, 1989-2019). Differences between length/area were considered as

significantly different when $p<0.05$ in Tukey HSD test unless indicated differently.

RNA extraction

To assess the expression of selected genes in the presence of pectin, total RNA was

extracted from *F. johnsoniae* cells grown for 48 hours at 30^OC on PY2 plates covered

with 500µl of DDW or 2% pectin as described above. For each plate, bacteria were

suspended in 1 ml cold (4^OC) PBS buffer, washed once in cold PBS, centrifuged for

2min at 18,000g, resuspended in TE buffer supplemented with 0.4mg/ml lysozyme and

then incubated for 10 min at room temperature. RNA was subsequently extracted from

cells using the TRIzol reagent (TRIzol® InvitrogenTM, #15596026), following the

manufacturer's instructions. Residual DNA was removed from the RNA samples by

digesting with RQ1 DNAse (Promega M6101A) at 37 °C for 40 min. For the real time

experiments, cDNA was synthesized using 50ng of DNAse treated RNA, with 1ul of

random primers (Promega C118A). Synthesis of single strand cDNA was achieved

using ImProm-IITM Reverse-Transcriptase (Promega, Madison, WI, United States).

The integrity and concentration of the extracted RNA and cDNA, was examined with a

QubitTM 3.0 Fluorometer (Thermo Fisher Scientific, United States) using reagents and

protocols supplied by the manufacturer, and by electrophoresis of samples on a 0.8%

agarose gel.

Live Imaging Fluorescent Microscopy Experiments.

To visualize the effect of different mono- and polysaccharides on *F. johnsoniae* colony

expansion, we filled the wells of a 24 well plate with 500µl of PY2 agar. Subsequently,

10ul of the mono- and polysaccharides (2%) or the DDW control were gently applied

to the wells in triplicates, plates were rotated for 1hour and dried overnight at room

temperature. A 30G needle (KDL 16-0310) was used to seed the bacteria in the center

of each well. Microscopic imaging was performed using a NIKON eclipse Ti

microscope (Nikon, Japan) equipped with a ProScan motorized XY stage, an HF110A

system (enabling rapid switching of emission filters)(Prior Scientific, MA, USA) and a

temperature-controlled stage incubator (LAUDA ECO RE 415, Korea). Bright field

illumination was provided by a cool LED pE-100A (Cool LED, UK). Excitation light

for epifluorescence microscopy was provided by a Spectra X light engine (Lumencor,

USA). Imaging was performed using a long working distance 40x objective (NA 0.6)

(Nikon, Japan). Images were captured at 2 hr intervals for 44 hr using an ANDOR zyla

5.5 MP sCMOS camera (Oxford Instruments, UK).

Images were processed using the NIS elements AR 4.6 (64 bit) software package

(Nikon, Japan) and Fiji (74). Fluorescence level was normalized to the initial measured

value (to avoid differences in the initial number of seeded bacteria) and to the maximal

fluorescence on PY2 media amended with DDW (to reduce variability of GFP

fluorescence levels between movies). The population growth dynamic of GFP- *F.*

johnsoniae on each substance computed using Fiji's time series analyzer plugin, and

average fluorescence density profiles of the expanding population over time was

quantified using JMP®, Version Pro 14 . (SAS Institute Inc., Cary, NC, 1989-2019).

Proteomics Sample preparation

Samples were subjected to in-solution tryptic digestion using a modified Filter Aided

Sample Preparation protocol (FASP). Sodium dodecyl sulfate buffer (SDS) included:

4% (w/v) SDS, 100mM Tris/HCl pH 7.6, 0.1M DTT. Urea buffer (UB): 8 M urea

(Sigma, U5128) in 0.1 M Tris/HCl pH 8.0 and 50mM Ammonium Bicarbonate.

Bacterial cell pellets were lysed in SDS buffer. Lysate was centrifuged at 16,000 g for

10min. 50 ug total protein were mixed with 2 mL UB, loaded onto 30 kDa molecular

weight cutoff filters (Sartorius VS15RH22) and centrifuged. 1.5 ml of UB were added

to the filter unit and centrifuged at 14,000 g for 40 min. Proteins were alkylated using

iodoacetamide (10 mM final concentration) and washed twice with Ammonium

Bicarbonate. Trypsin was then added (50:1 protein amount:trypsin) and samples

incubated at 37°C overnight, followed by a second trypsin digestion for 4 hours at 37°C.

Digested peptides were then collected in a clean tube by centrifugation, acidified with

trifloroacetic acid, desalted using HBL Oasis (Waters 094225), speed vaccumed to

dryness and stored in -80°C until analysis. All chemicals used were from Sigma

Aldrich, unless stated otherwise. Sample preparation and initial statistical analysis was

performed at the Nancy and Stephen Grand Israel National Center for Personalized

Medicine, The Weizmann Institute of Science, Rehovot, Israel.

Liquid chromatography and proteomics analysis

Each sample was loaded and analyzed using split-less nano-Ultra Performance Liquid

Chromatography (10 kpsi nanoAcquity; Waters, Milford, MA, USA). The mobile phase was:

A) H_2O + 0.1% formic acid and B) acetonitrile + 0.1% formic acid. Desalting of the samples

was performed online using a Symmetry C18 reversed-phase trapping column (180 µm internal

diameter, 20 mm length, 5 µm particle size; Waters). The peptides were then separated using a

T3 HSS nano-column (75 µm internal diameter, 250 mm length, 1.8 µm particle size; Waters)

at 0.35 µL/min. Peptides were eluted from the column into the mass spectrometer using the

following gradient: 4% to 20%B in 155 min, 20% to 90%B in 5 min, maintained at 90% for 5

min and then back to initial conditions. ULC/MS grade solvents were used for all

chromatographic steps.Each sample was analyzed on the instrument separately in a

random order in discovery mode. Raw data was searched against the *F. johnsoniae*

protein databases, to which a list of common lab contaminants was added. Database

search was done using the Mascot algorithm, and quantitative analysis was performed

using the Expressionist software from GeneData. As a rule of thumb we consider

significant differences to be >1 peptide per protein, fold change >2 or <0.5, and <0.05

p-value. Proteins were functionally annotated using RAST (Rapid Annotations using

Subsystems Technology) (75).

Twenty-five pectin-induced proteins were selected based on high fold change between

PY2 agar coated with pectin vs. PY2 agar coated with DDW and statistical significance

(proteins without annotation were removed from this analysis). Gene abundance across

the dataset was normalized to 100% for each gene, and a heatmap was created using

studio.plot.ly (https://plotly.com/).

Quantitative PCR Assessment of Gene Expression Levels

The expression of 10 genes encoding for proteins found to be significantly induced on

pectin in the proteomic analyses were analyzed using qPCR. Primers for qRT-PCR

experiments (**Table S5**) were constructed based on the *F. johnsoniae* genome sequence

and were pre-designed using the PrimerQuest® tool (Integrated DNA Technologies,

USA). Triplicate cDNA samples for each of the treatments (with or without pectin)

were diluted and 2ng was used in a 20 μl final reaction volume together with 10ul Fast

SYBR^{TM} green PCR master mix (Thermo Fisher scientific), 100 nM each of forward

and reverse primers, DDW and 1ul of template cDNA. Amplification was carried out

on a StepOnePlus real-time PCR thermocycler (Applied Biosystems, Foster City, CA,

United States) using the following program: heat activation of DNA polymerase at

95°C for 3 min and 40 cycles at 95°C for 5 sec for denaturation and primer annealing

and extension at 60°C for 30 sec. A melting curve was produced to confirm a single

gene-specific peak and to detect primer-dimer formation by heating the samples from

60 to 95°C in 0.3°C increments. Gene amplification was confirmed by detection of a

single peak in the melting curve analysis. For each gene, PCR gene amplification was

carried out using three independent biological replicates. Expression of each of the

targeted genes was normalized to that of three alternative housekeeping genes (16S

rRNA, DNA gyrase subunit B (*gyr*B, EC 5.99.1.3) , and the electron transfer

flavoprotein, alpha subunit (ETF) threonine synthase (EC 4.2.3.1). These genes were

selected because there was no detected difference in their expression when grown on

PY2 media amended with pectin *vs.* DDW in the proteomic analysis. The relative

abundance of each target gene relative to a reference gene was determined according to

the method described previously (Livak and Schmittgen2001). Concentrations and

ΔΔCT values were calculated and analyzed with the StepOne software v2.3 (Applied

Biosystems, Foster City, CA, United States). Concomitant "no-RT" reactions,

lacking reverse transcriptase, were performed for each sample and run to confirm

absence of DNA contamination, as well as no template controls (NTCs) to confirm lack

of contamination. Reaction efficiency was monitored by means of an in internal

standard curve using a 10-fold dilution of DNA ranging from 0.01-10ng of DNA per

reaction, in triplicates. Efficiency was between 92.1 and 97.2% for all primers, and R2-

values were greater than 0.99. Data analysis was conducted using the StepOne software

v2.3 (Applied Biosystems, Foster City, CA, United States).

Acknowledgments: We would like to thank Alla Usyskin-Tonne, for her help with the

proteomics functional annotation and Eduard Belausov for his help with the binocular

based imaging. We thank Prof. M. J. McBride from the University of Wisconsin–

Milwaukee for generously providing plasmids providing plasmids and bacterial *F.*

Johnsoniae strains.

References

1.   Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere
microbiome and plant health. Trends Plant Sci 17:478–486.
2.   Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P.
2013. Structure and Functions of the Bacterial Microbiota of Plants. Annu Rev
Plant Biol 64:807–838.
3.   Reinhold-Hurek B, Bünger W, Burbano CS, Sabale M, Hurek T. 2015. Roots
Shaping Their Microbiome: Global Hotspots for Microbial Activity. Annu Rev
Phytopathol 53:403–424.
4.   Dennis PG, Miller AJ, Hirsch PR. 2010. Are root exudates more important than
other sources of rhizodeposits in structuring rhizosphere bacterial
communities? FEMS Microbiol Ecol 72:313–327.
5.   Barret M, Morrissey JP, O'Gara F. 2011. Functional genomics analysis of plant
growth-promoting rhizobacterial traits involved in rhizosphere competence.
Biol Fertil Soils 47:729–743.
6.   Beauregard PB, Chai Y, Vlamakis H, Losick R, Kolter R. 2013. Bacillus
subtilis biofilm induction by plant polysaccharides. Proc Natl Acad Sci
110:E1621–E1630.
7.   Massalha H, Korenblum E, Malitsky S, Shapiro OH, Aharoni A. 2017. Live
imaging of root–bacteria interactions in a microfluidics setup. Proc Natl Acad
Sci 114:4549–4554.
8.   Sasse J, Martinoia E, Northen T. 2018. Feed Your Friends: Do Plant Exudates
Shape the Root Microbiome? Trends Plant Sci 23:25–41.
9.   Ofek M, Voronov-Goldman M, Hadar Y, Minz D. 2014. Host signature effect
on plant root-associated microbiomes revealed through analyses of resident vs .
active communities. Environ Microbiol 16:2157–2167.
10.   Rosenberg E., DeLong E.F., Lory S., Stackebrandt E. TF. 2014. The Family
Chitinophagaceae, p. 119–144. *In* E., R, E.F., D, S., L, E., S, F., T (eds.), The
Prokaryotes. Springer, Berlin, Heidelberg.
11.   Rudrappa T, Czymmek KJ, Paré PW, Bais HP. 2008. Root-Secreted Malic
Acid Recruits Beneficial Soil Bacteria. Plant Physiol 148:1547–1556.
12.   Williams A, Wilkinson A, Krehenbrink M, Russo DM, Zorreguieta A, Downie
JA. 2008. Glucomannan-mediated attachment of Rhizobium leguminosarum to
pea root hairs is required for competitive nodule infection. J Bacteriol
190:4706–4715.
13.   Oku S, Ayaka K, Takahisa TK, Junichi YN. 2012. Identification of Chemotaxis
Sensory Proteins for Amino Acids in Pseudomonas fluorescens Pf0-1 and Their
Involvement in Chemotaxis to Tomato Root Exudate and Root Colonization.
Microbes Env 27:462–469.
14.   Webb BA, Compton KK, del Campo JSM, Taylor D, Sobrado P, Scharf BE.
2017. Sinorhizobium meliloti Chemotaxis to Multiple Amino Acids Is
Mediated by the Chemoreceptor McpU. Mol Plant-Microbe Interact 30:770–
777.
15.   Feng H, Zhang N, Fu R, Liu Y, Krell T, Du W, Shao J, Shen Q, Zhang R.
2019. Recognition of dominant attractants by key chemoreceptors mediates
recruitment of plant growth-promoting rhizobacteria. Environ Microbiol
21:402–415.
16.   Cremer J, Honda T, Tang Y, Wong-Ng J, Vergassola M, Hwa T. 2019.

Chemotaxis as a navigation strategy to boost range expansion. Nature 575:658–
663.

17.   Lugtenberg BJJ, Dekkers LC. 1999. What makes Pseudomonas bacteria
598   rhizosphere competent? Environ Microbiol 1:9–13.

18.   Yan Y, Yang J, Dou Y, Chen M, Ping S, Peng J, Lu W, Zhang W, Yao Z, Li H,
600   Liu W, He S, Geng L, Zhang X, Yang F, Yu H, Zhan Y, Li D, Lin Z, Wang Y,
601   Elmerich C, Lin M, Jin Q. 2008. Nitrogen fixation island and rhizosphere
602   competence traits in the genome of root-associated Pseudomonas stutzeri
603   A1501. Proc Natl Acad Sci 105:7564–7569.

19.   Pieterse MJ, Zamioudis C, Berendsen RL, Weller DM, Wees SCM Van,
605   Bakker PAHM. 2014. Induced Systemic Resistance by Beneficial Microbes.

20.   McBride MJ, Liu W, Lu X, Zhu Y, Zhang W. 2014. The Family
607   Cytophagaceae, p. 577–593. *In* The Prokaryotes. Springer Berlin Heidelberg,
608   Berlin, Heidelberg.

21.   Kolton M, Sela N, Elad Y, Cytryn E. 2013. Comparative Genomic Analysis
610   Indicates that Niche Adaptation of Terrestrial Flavobacteria Is Strongly Linked
611   to Plant Glycan Metabolism. PLoS One 8:e76704.

22.   Janssen PH. 2006. Identifying the Dominant Soil Bacterial Taxa in Libraries of
613   16S rRNA and 16S rRNA Genes. Appl Environ Microbiol 72:1719–1728.

23.   Manter DK, Delgado JA, Holm DG, Stong RA. 2010. Pyrosequencing Reveals
615   a Highly Diverse and Cultivar-Specific Bacterial Endophyte Community in
616   Potato Roots. Microb Ecol 60:157–166.

24.   Johansen JE, Binnerup SJ, Lejbolle KB, Mascher F, Sorensen J, Keel C. 2002.
618   Impact of biocontrol strain Pseudomonas fluorescens CHA0 on rhizosphere
619   bacteria isolated from barley (Hordeum vulgare L.) with special reference to
620   Cytophaga-like bacteria. J Appl Microbiol 93:1065–1074.

25.   Kolton M, Meller Harel Y, Pasternak Z, Graber ER, Elad Y, Cytryn E. 2011.
622   Impact of Biochar Application to Soil on the Root-Associated Bacterial
623   Community Structure of Fully Developed Greenhouse Pepper Plants. Appl
624   Environ Microbiol 77:4924–4930.

26.   Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S,
626   Tremblay J, Engelbrektson A, Kunin V, Rio TG del, Edgar RC, Eickhorst T,
627   Ley RE, Hugenholtz P, Tringe SG, Dangl JL. 2012. Defining the core
628   Arabidopsis thaliana root microbiome. Nature 488:86–90.

27.   Bodenhausen N, Horton MW, Bergelson J. 2013. Bacterial Communities
630   Associated with the Leaves and the Roots of Arabidopsis thaliana. PLoS One
631   8:e56329.

28.   Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y,
633   McHardy AC, Schulze-Lefert P. 2015. Structure and Function of the Bacterial
634   Root Microbiota in Wild and Domesticated Barley. Cell Host Microbe 17:392–
635   403.

29.   Sang MK, Kim KD. 2012. The volatile-producing Flavobacterium johnsoniae
637   strain GSE09 shows biocontrol activity against Phytophthora capsici in pepper.
638   J Appl Microbiol 113:383–398.

30.   Gunasinghe RN, Ikiriwatte CJ, Karunaratne AM. 2004. The use of Pantoea
640   agglomerans and Flavobacterium sp . to control banana pathogens. J Hortic Sci
641   Biotechnol 79:1002–1006.

31.   Kolton M, Frenkel O, Elad Y, Cytryn E. 2014. Potential Role of Flavobacterial
Gliding-Motility and Type IX Secretion System Complex in Root Colonization
and Plant Defense. Mol Plant-Microbe Interact 27:1005–1013.

32. Xue C, Ryan Penton C, Shen Z, Zhang R, Huang Q, Li R, Ruan Y, Shen Q.
2015. Manipulating the banana rhizosphere microbiome for biological control
of Panama disease. Sci Rep 5:11124.

33. Kwak M-J, Kong HG, Choi K, Kwon S-K, Song JY, Lee J, Lee PA, Choi SY,
Seo M, Lee HJ, Jung EJ, Park H, Roy N, Kim H, Lee MM, Rubin EM, Lee S-
W, Kim JF. 2018. Rhizosphere microbiome structure alters to enable wilt
resistance in tomato. Nat Biotechnol 36:1100–1109.

34. Carrión VJ, Perez-Jaramillo J, Cordovez V, Tracanna V, de Hollander M, Ruiz-
Buck D, Mendes LW, van Ijcken WFJ, Gomez-Exposito R, Elsayed SS,
Mohanraju P, Arifah A, van der Oost J, Paulson JN, Mendes R, van Wezel GP,
Medema MH, Raaijmakers JM. 2019. Pathogen-induced activation of disease-
suppressive functions in the endophytic root microbiome. Science (80-)
366:606–612.

35. Wei Z, Gu Y, Friman V-P, Kowalchuk GA, Xu Y, Shen Q, Jousset A. 2019.
Initial soil microbiome composition and functioning predetermine future plant
health. Sci Adv 5:eaaw0759.

36. Hebbar P, Berge O, Heulin T, Singh SP. 1991. Bacterial antagonists of
Sunflower (Helianthus annuus L.) fungal pathogens. Plant Soil 133:131–140.

37. Sang MK, Kim J Do, Kim BS, Kim KD. 2011. Root Treatment with
Rhizobacteria Antagonistic to Phytophthora Blight Affects Anthracnose
Occurrence, Ripening, and Yield of Pepper Fruit in the Plastic House and Field.
Phytopathology 101:666–678.

38. Alexander BJR, Stewart A. 2001. Glasshouse screening for biological control
agents of Phytophthora cactorum on apple (Malus domestica). New Zeal J
Crop Hortic Sci 29:159–169.

39. McBride MJ, Nakane D. 2015. Flavobacterium gliding motility and the type IX
secretion system. Curr Opin Microbiol 28:72–77.

40. Kharade SS, McBride MJ. 2015. Flavobacterium johnsoniae PorV Is Required
for Secretion of a Subset of Proteins Targeted to the Type IX Secretion System.
J Bacteriol 197:147–158.

41. Lauber F, Deme JC, Lea SM, Berks BC. 2018. Type 9 secretion system
structures reveal a new protein transport mechanism. Nature 564:77–82.

42. Doug Hyatt, Gwo-Liang Chen, Philip F LoCascio, Miriam L Land, , Frank W
Larimer LJH. 2010. Integrated nr Database in Protein Annotation System and
Its Localization. Nat Commun 6:1–8.

43. McBride MJ, Xie G, Martens EC, Lapidus A, Henrissat B, Rhodes RG,
Goltsman E, Wang W, Xu J, Hunnicutt DW, Staroscik AM, Hoover TR, Cheng
Y-Q, Stein JL. 2009. Novel Features of the Polysaccharide-Digesting Gliding
Bacterium Flavobacterium johnsoniae as Revealed by Genome Sequence
Analysis. Appl Environ Microbiol 75:6864–6875.

44. Foley MH, Cockburn DW, Koropatkin NM. 2016. The Sus operon: a model
system for starch uptake by the human gut Bacteroidetes. Cell Mol Life Sci
73:2603–2617.

45. Shrivastava A, Johnston JJ, Van Baaren JM, McBride MJ. 2013.
Flavobacterium johnsoniae GldK, GldL, GldM, and SprA are required for
secretion of the cell surface gliding motility adhesins sprb and remA. J
Bacteriol 195:3201–3212.

46. Johnston JJ, Shrivastava A, McBride MJ. 2018. Untangling Flavobacterium
johnsoniae gliding motility and protein secretion. J Bacteriol 200.

47. Gorski L, Godchaux W, Leadbetter ER. 1993. Structural specificity of sugars

that inhibit gliding motility of Cytophaga johnsonae. Arch Microbiol 160:121–
125.

48. Wolkin RH, Pate JL. 1984. Translocation of motile cells of the gliding
bacterium Cytophaga johnsonae depends on a surface component that may be
modified by sugars. J Gen Microbiol 130:2651–2669.

49. Imamura K, Sato K, Narita Y, Kondo Y, Nakane D, Naito M, Fujiwara T,
Nakayama K. 2018. Identification of a major glucose transporter in
Flavobacterium johnsoniae : Inhibition of F. johnsoniae colony spreading by
glucose uptake. Microbiol Immunol 62:507–516.

50. Wood DC, Hayasaka SS. 1981. Chemotaxis of rhizoplane bacteria to amino
acids comprising eelgrass (Zostera marina L.) root exudate. J Exp Mar Bio
Ecol 50:153–161.

51. Raina J-B, Fernandez V, Lambert B, Stocker R, Seymour JR. 2019. The role of
microbial motility and chemotaxis in symbiosis. Nat Rev Microbiol 17:284–
294.

52. Walker TS, Bais HP, Grotewold E, Vivanco JM. 2003. Root Exudation and
Rhizosphere Biology: Fig. 1. Plant Physiol 132:44–51.

53. Rodrà-guez-Navarro DN, Dardanelli MS, RuÃ-z-SaÃ-nz JE. 2007.
Attachment of bacteria to the roots of higher plants. FEMS Microbiol Lett
272:127–136.

54. Curtis TP, Sloan W T. 2005. MICROBIOLOGY: Exploring Microbial
Diversity--A Vast Below. Science (80-) 309:1331–1333.

55. Lagos L, Maruyama F, Nannipieri P, Mora M., Ogram A, Jorquera M. 2015.
Current overview on the study of bacteria in the rhizosphere by modern
molecular techniques: a mini‒review. J soil Sci plant Nutr 15:0–0.

56. Bulgarelli D, Rott M, Schlaeppi K, Ver E, Themaat L Van, Ahmadinejad N,
Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner
FO, Amann R, Eickhorst T, Schulze-lefert P. 2012. Revealing structure and
assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature
488:91–95.

57. Bergelson J, Mittelstrass J, Horton MW. 2019. Characterizing both bacteria and
fungi improves understanding of the Arabidopsis root microbiome. Sci Rep
9:24.

58. Arlauskas J, Burchard RP. 1982. Substratum requirements for bacterial gliding
motility. Arch Microbiol 133:137–141.

59. Guan H, Pan B, Fang Z, Gong M, Shen B, Shen Q, Guo R, Wu K, Shi W, Yuan
S. 2015. Pectin Enhances Bio-Control Efficacy by Inducing Colonization and
Secretion of Secondary Metabolites by Bacillus amyloliquefaciens SQY 162 in
the Rhizosphere of Tobacco. PLoS One 10:e0127418.

60. Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM,
Kassabgy M, Huang S, Mann AJ, Waldmann J, Weber M, Klindworth A, Otto
A, Lange J, Bernhardt J, Reinsch C, Hecker M, Peplies J, Bockelmann FD,
Callies U, Gerdts G, Wichels A, Wiltshire KH, Glockner FO, Schweder T,
Amann R. 2012. Substrate-Controlled Succession of Marine Bacterioplankton
Populations Induced by a Phytoplankton Bloom. Science (80-) 336:608–611.

61. Postle K, Kadner RJ. 2003. Touch and go: Tying TonB to transport. Mol
Microbiol 49:869–882.

62. Postle K. 2007. TonB System, In Vivo Assays and Characterization, p. 245–
269. *In* Methods in Enzymology.

63. Shrivastava A, Rhodes RG, Pochiraju S, Nakane D, McBride MJ. 2012.

745   Flavobacterium johnsoniae RemA is a mobile cell surface lectin involved in
746   gliding. J Bacteriol 194:3678–3688.
747 64.   Molina MA, Godoy P, Ramos-Gonzalez MI, Munoz N, Ramos JL, Espinosa-
748   Urgel M. 2005. Role of iron and the TonB system in colonization of corn seeds
749   and roots by Pseudomonas putida KT2440. Environ Microbiol 7:443–449.
750 65.   Taillefer M, Arntzen MØ, Henrissat B, Pope PB, Larsbrink J. 2018. Proteomic
751   Dissection of the Cellulolytic Machineries Used by Soil-Dwelling
752   Bacteroidetes. mSystems 3:1–16.
753 66.   Vorhölter F-J, Wiggerich H-G, Scheidle H, Sidhu VK, Mrozek K, Küster H,
754   Pühler A, Niehaus K. 2012. Involvement of bacterial TonB-dependent
755   signaling in the generation of an oligogalacturonide damage-associated
756   molecular pattern from plant cell walls exposed to Xanthomonas campestris pv.
757   campestris pectate lyases. BMC Microbiol 12:239.
758 67.   Flory MR, Lee H, Bonneau R, Mallick P, Serikawa K, Morris DR, Aebersold
759   R. 2006. Quantitative proteomic analysis of the budding yeast cell cycle using
760   acid-cleavable isotope-coded affinity tag reagents. Proteomics 6:6146–6157.
761 68.   Liu Y, Beyer A, Aebersold R. 2016. On the Dependency of Cellular Protein
762   Levels on mRNA Abundance. Cell 165:535–550.
763 69.   Liu W, Cremer J, Li D, Hwa T, Liu C. 2019. An evolutionarily stable strategy
764   to colonize spatially extended habitats. Nature 575:664–668.
765 70.   Kolton M, Frenkel O, Elad Y, Cytryn E. 2014. Potential Role of Flavobacterial
766   Gliding-Motility and Type IX Secretion System Complex in Root Colonization
767   and Plant Defense. Mol Plant-Microbe Interact 27:1005–1013.
768 71.   Sela N, Cytryn E, Green SJ, Harel YM, Kolton M, Elad Y. 2012. Draft
769   Genome Sequence of Flavobacterium sp. Strain F52, Isolated from the
770   Rhizosphere of Bell Pepper (Capsicum annuum L. cv. Maccabi). J Bacteriol
771   194:5462–5463.
772 72.   Mcbride MJ, Baker SA. 1996. Development of techniques to genetically
773   manipulate members of the genera Cytophaga, Flavobacterium, Flexibacter,
774   and Sporocytophaga. Appl Environ Microbiol 62:3017–3022.
775 73.   Agarwal S., Hunnicutt DW, McBride MJ. 2001. Cloning and Characterization
776   of theFlavobacterium johnsoniae Gliding Motility GenesgldD and gldE. J
777   Bacteriol 183:4167–4175.
778 74.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
779   Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J, White DJ, Hartenstein
780   V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for
781   biological-image analysis. Nat Methods 9:676–682.
782 75.   Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA,
783   Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R.
784   2014. The SEED and the Rapid Annotation of microbial genomes using
785   Subsystems Technology (RAST). Nucleic Acids Res 42:D206–D214.
786

Fig1- Impact of sugars on proliferation of flavobacteria. Image (A) and quantification

(B) of colony area of *F. johnsoniae* colony expansion on PY2 agar amended with different

mono- and polysaccharides: arabinose, glucose, microcrystalline cellulose (M.Cellulose),

pectin and no sugar control, after 48hr incubation at 30°C (N=4). Image (C) and quantification

(D) of colony expansion on PY2 agar of wild type (WT) and gliding mutants (ΔgldJ) of *F.*

johnsoniae and *Flavobacterium* sp. F52, amended with DDW or 2% pectin after 48hr

incubation at 30°C (N=3). Colony area was measured using Fiji. Letters indicate statistical

significance calculated using ®JMP Pro14, was considered significant if p<0.05 by Tukey

 Fig2 - Impact of pectin precursors (galacturonic acid and rhamnose) on

 proliferation of flavobacteria. (A) *F. johnsoniae* colony expansion on PY2 agar

 amended with pectin, galacturonic acid, rhamnose, and galacturonic acid and rhamnose

 or no sugar control incubated at 30°C for 48 h (N=4). (B) Graphic description of colony

 area based on results from (A). Letters indicate statistical significance, calculated using

 JMP Pro14®, and differences between treatments were considered significant when

 $p < 0.05$ by Tukey HSD.

Fig3 - Temporal dynamics of flavobacterial proliferation on polysaccharides using live imaging microscopy. (A) Morphology of GFP-labeled *F. johnsoniae* colonies on selected sugars and polysaccharides. Bacteria were inoculated in the center of PY2 agar coated with the indicated sugars and polysaccharides (schematically described in the insert). Images show colony morphology after 32hr. (B) Enlarged image of GFP-*F. johnsoniae* colony morphology after 20hr of growth on the selected sugars and polysaccharides as indicated in (A). (C) Growth rates of GFP-*F. johnsoniae* colonies on the selected sugars and polysaccharides. Data was normalized as described in the materials and methods section. Differences in the average colony fluorescence intensity after 44 hr was compared and considered significant if $p < 0.05$ by Tukey HSD (indicated by letters). Data includes means and data from three biological replicates composed of three technical repeats in each. (D) Temporal dynamics of GFP-*F. johnsoniae* growth rates. Growth was compared at the peaks (20hr and 36hr) and considered significant if $p < 0.05$ by Tukey HSD. (E) Schematic diagram showing the three characterized regions of interest (ROI- 1.5, 3 and 4.5 mm radii) used to evaluate of bacterial expansion rates. (F) Estimated expansion rates of GFP-*F. johnsoniae* on the selected sugars and polysaccharides. Velocity was estimated by expansion time in hours taken to cross known ROIs as indicated in E. (G) Estimated expansion time relative to estimated growth of GFP-*F. johnsoniae* on the selected sugars and polysaccharides. Colored circles mark time (h) for bacteria to cross the 4.5mm radius on each substance as calculated in F. Differences between treatments were considered significant when $p < 0.05$ by Tukey HSD

Fig4 - Pectin induced flavobacterial genes and proteins. (A) Differential expression of the 25 most substantial pectin induced proteins based on proteomic analysis of *F. johnsoniae* colonies grown on PY2 medium amended with pectin relative to colonies grown on identical media amended without pectin (DDW). Heat map shows triplicates for each treatment. All described proteins are statistically significant ($p<0.05$). (B) The mRNA expression level of selected proteins (Sus73 (SHH12854.1), TonB260 (WP_012026229.1), TonB445 (WP_012022294.1), TonB 2144 (WP_012026069.1), Auxin regulator (WP_012023111.1), Transcriptional regulator (WP_012022876.1), Pectate lyase (WP_012026072.1) and RemA(WP_012022896.1) shown to be induced in the proteomic analysis described in (A), using quantitative real-time PCR (qPCR). Fold changes in mRNA levels of the target genes were normalized against the 16SrRNA gene (left), the Electron transfer Flavoprotein, alpha subunit (WP_012023552.1) (center), and the DNA gyrase subunit B (WP_012024321.1)(right). Change in target genes fold change RNA expression was calculated using the $2^{-\Delta\Delta CT}$ method and statistical significance ($p<0.05$) by Student T-test. Error bars represent standard errors of six independent experiments based on two independent RNA extraction.