1

Plant cell wall polysaccharides induce colony expansion of soil-derived

Flavobacterium spp.

Judith Kraut-Cohen[a], Orr H.Shapiro[b], Barak Dror[ac] Eddie Ctyryn[a]

[a]Institute of Soil, Water and Environmental Sciences, Agricultural Research

Organization, Volcani Center, Rishon LeZion 7505101, Israel

[b]Institute Postharvest and Food Sciences Agricultural Research Organization, Volcani

Center, Rishon LeZion 7505101, Israel

[c]Department of Plant Pathology and Microbiology, The R.H. Smith Faculty of

Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot,

Israel

Running Head: Pectin induce Flavobacteria colony expansion by TonB proteins

#Address correspondence to Eddie Cytryn, eddie@volcani.agri.gov.il.

*Present address: Agricultural Research Organization - the Volcani Center,

68 HaMaccabim Road , P.O.B 15159 Rishon LeZion 7505101, Israel.

Abstract

Flavobacterium is a genus of gram-negative bacteria, belonging to the *Bacteriodetes* phylum, characterized by a unique gliding motility. They are ubiquitous and often abundant in root microbiomes of various plants, but the factors contributing to this high abundance are currently unknown. In this study, we evaluate the effect of various plant-associated poly- and mono-saccharides on growth and colony expansion of two *Flavobacterium* strains (*F. Johnsoniae*, *Flavobacterium* sp. F52). Both strains were able to grow on pectin and other polysaccharides such as cellulose as a single carbon source. However, only pectin, a polysaccharide that is profuse in plant cell walls, enhanced colony expansion on solid surfaces even under high nutrient availability, suggesting a link between carbohydrate metabolism and gliding. Expansion on pectin was dose- and substrate-dependent, as it did not occur when bacteria were grown on the pectin monomers galacturonic acid and rhamnose. Using time-lapse microscopy, we demonstrated a bi-phasic expansion of *F. johnsoniae* on pectin: an initial phase of rapid expansion, followed by biomass production within the colonized area. Proteomics and gene expression analyses revealed significant induction of several carbohydrate metabolism related proteins when *F. johnsoniae* was grown on pectin, including selected operons of SusC/D, TonB-dependent glycan transport genes. Our results suggest a yet unknown linkage between specific glycan associated operons and flavobacterial motility. This may be associated with their capacity to rapidly glide along the root and metabolize plant cell wall carbohydrates, two characteristics that are crucial to rhizosphere competence.

Importance

The genus *Flavobacterium* is highly abundant and enriched (relative to surrounding bulk soil) in plant root microbiomes, where they may play a role in plant health and ecosystem functioning. However, little is known about genetic and physiological characteristics that enable flavobacteria to colonize and proliferate in this highly competitive environment. In this study, we found that plant cell wall-polysaccharides and specifically pectin stimulate flavobacteria colony development in a bi-phasic manner, initially characterized by rapid expansion followed by increased biomass production. This appears to be linked to pectin-facilitated induction of specific TonB-associated proteins evidentially involved in the detection and uptake of plant sugars. These findings suggest that the capacity to sense, expand on and metabolize pectin and other plant cell wall polysaccharides play a fundamental role in promoting rhizosphere competence in flavobacteria. This work sheds light on specific mechanisms that facilitate plant-microbe interactions, which are fundamental for promoting plant health and for understanding the microbial ecology of root ecosystems.

Keywords: Flavobacterium, Roots, Rhizosphere, Pectin, TonB

Introduction

The complex interactions between plant-associated microorganisms and their hosts (collectively referred to as the "plant holobiont") are crucial for plant health and growth ((1–3). Plants modulate the narrow region of soil that is influenced by root secretions, the rhizosphere, by exuding various small molecular weight compounds (rhizodeposites) such as sugars, amino acids and organic acids, by rhizodepositing root cap border cells, and by releasing various mono- and poly-saccharides in their mucilage (3–8). Organic compounds released from the roots transform the rhizosphere into a nutrient rich environment relative to the surrounding bulk soil, which facilitates colonization by soil microorganisms. Rapid growth, coupled to competition for root resources, results in less diverse rhizosphere and rhizoplane microbiomes relative to surrounding bulk soil, a phenomenon known as the "rhizosphere effect" (4, 9, 10). Rhizosphere and root colonizing bacteria have the capacity to outcompete other soil bacteria and thereby proliferate and thrive in the root ecosystem through a combination of specific traits collectively coined "rhizosphere competence", which include: motility, resistance to stress, ability to utilize plant-derived organic compounds, chemotaxis, and the production of secondary metabolites (5). Some earlier studies demonstrated that rhizosphere bacteria are attracted to the roots by plant-exuded organic acids such as malic, citric or fumaric acid and various amino acids (11–15). Many of these sensed chemo-attractants can also be consumed by the bacteria (16). Although rhizosphere recruitment and colonization mechanisms of certain plant-growth-promoting rhizobacteria (PGPR) have been identified and characterized, those responsible for recruitment of the vast majority of rhizosphere and rhizoplane bacteria is currently an enigma (17–19).

Flavobacterium is a Gram-negative genus of bacteria from the phylum *Bacteroidetes*

known to degrade complex organic compounds, which is abundant in freshwater,

marine and soil environments (20, 21)). It is highly abundant in the rhizoplane,

rhizosphere and phylosphere of a wide array of plants, in contrast to considerably lower

abundance in bulk soil (21–28), suggesting that members of this genus have specifically

adapted to root ecosystems. Several root- and soil-derived flavobacteria were found to

antagonize various plant pathogens in different crops (29–33), and recently it was

discovered that entophytic *Flavobacterium* strains play a fundamental role in

antagonizing phytopathogenic fungi, through the production chitinolytic enzymes and

specific secondary metabolites (34). Furthermore, they were recently identified as

potential driver taxa behind pathogen suppression in analysis of bacterial network in a

plant root microbiomes (35). In addition, selected members of this genus are linked to

increased plant biomass, and have therefore been defined as plant growth promoting

rhizobacteria (PGPR) of various crops (23, 29, 30, 36–38).

Soil flavobacteria have specialized ability to decompose complex plant derived

polysaccharides, such as pectin and cellulose and the ability to secret various carbolytic

enzymes via the *Bacteriodetes*-specific type IX secretion system (T9SS) (21, 39–43).

Like other *Bacteroidetes*, they contain a myriad of genes that encode Polysaccharide

Utilization Loci (PULs) that are activated specifically to facilitate glycan capture and

intake (44). These PULs include outer membrane protein transducers involved in

polysaccharide utilization, which are part of the TonB family, generally referred to as

Starch Utilization System (SUS) proteins. Interestingly, comparative genomics

revealed that genomes of soil and root-associated flavobacterial strains are significantly

enriched with genes associated with plant polysaccharide degradation relative to

aquatic strains from this genus, indicating that physiology of this genus is strongly influence by its ecology (21).

Most terrestrial *Flavobacterium* strains possess a unique gliding mechanism that rapidly propels them over solid and semi-solid surfaces. The proteins that comprise this gliding system are molecularly intertwined with at least fifteen proteins that make up the T9SS, seven of whom are responsible for the secretion of SprB and RemA adhesins which are expressed on the cell surface and involved in gliding (45, 46). We previously demonstrated that this T9SS-gliding complex is crucial for root colonization by flavobacteria, and this colonization was positively linked to the induction of plant resistance to foliar pathogens (31).

Collectively, the above data strongly suggest that terrestrial flavobacterial strains have evolved means that enable them to interact with plant roots, and that these interactions are beneficial to plant health. Nonetheless, the specific mechanisms behind this phenomenon are currently unclear.

In this study, we assessed the impact of various plant-derived sugars on the motility and growth dynamics of flavobacteria by coupling conventional petri dish assays and live-imaging fluorescent microscopy with proteomic and gene expression analyses. This study demonstrates that pectin, a plant cell wall-associated polysaccharide, facilitates bi-phasic proliferation over solid surfaces through induction of specific TonB-associated glycan uptake operons. *In-planta* extrapolation of these results, suggests that this link between pectin, motility and carbohydrate metabolism may be fundamental to rhizosphere competence in flavobacteria.

Results

Growth of *Flavobacterium* strains on various carbon sources

We previously demonstrated that genomes of terrestrial flavobacteria are enriched with

genes involved in plant carbohydrate metabolism (21). To better understand the

interaction of flavobacteria with sugars, growth was evaluated on minimal and rich

media amended with the selected mono- and polysaccharides described. *F. johnsoniae*

was motile and grew on chitin, pectin and cellulose but not on arabinose, as a single

carbon source, (**FigS1**). Subsequently, we evaluated growth of flavobacteria on rich

media (PY2 agar) coated with selected sugars (**Fig1A**). Colony expansion of *F.*

johnsoniae on pectin was found to be significantly greater than the other analyzed

sugars, and was five times higher than DDW ($p > 0.05$, Tukey-Kramer HSD test)

(**Fig1B**).

Plant cell wall polysaccharides expedite flavobacterial colony expansion

Wild type (WT) and gliding/typeIX secretion system mutants (ΔgldJ) of *F. johnsoniae*

and the pepper root isolate *Flavobacterium* sp. F52 were inoculated in the center of

PY2 agar media amended with DDW or pectin (**Fig1C**). After 48 hours of incubation,

significant colony expansion of wild type bacteria was observed in both

Flavobacterium strains when grown on pectin, while no similar growth was found on

DDW. In contrast, gliding mutants (ΔgldJ) of both flavobacterial strains did not

proliferate differentially on any of the plates (**Fig1C,D**), indicating that the gliding

apparatus is a prerequisite for pectin-induced colony expansion.

Dose-dependent pectin facilitated colony expansion

To determine whether expansion on pectin is dose dependent, *F. johnsoniae* and

Flavobacterium sp.F52 strains were inoculated at the center of PY2 agar media plates

streaked with pectin at final concentrations of 0.5, 1, 2 and 4% percent. For all the

examined pectin concentrations, colonies radiated along the pectin streaks, with 2% and

4% pectin concentrations inducing significantly more expansion than the lower

concentrations (p>0.05,Tukey-Kramer HSD test) (**FigS2**).

Since galactronic-acid and rhamnose are the two major components of pectins, we

examined the colony expansion of *F. johnsoniae* on rich media plates coated with

DDW, 2% titrated galactronic-acid, 2% rhamnose, and a combination of galactronic

acid and rhamnose. *F. johnsoniae* expansion on galactronic-acid, rhamnose or both of

them together did not result in significant colony expansion (p>0.05,Tukey-Kramer

HSD test), in contrast to bacteria grown on pectin (**Fig2A, B**).

Temporal dynamics of pectin induced *F. johnsoniae* colony expansion

The expansion of GFP labeled *F. johnsoniae* on PY2 agar coated with glucose,

cellulose, glucomannan or pectin, was visualized at a higher resolution using time-lapse

microscopy. Colony morphology after 32 hours, on each tested substance is presented

in **Fig3A**.

Growth dynamics were clearly affected by the sugar type. Expansion on pectin was

characterized by a relatively long lag phase. However following this stage, colonies

rapidly expanded and after 18 hours, the fluorescent signal of colonies grown on pectin

surpassed both the control (DDW) and the other amended compounds (according to

Tukey-Kramer HSD test). Growth on pectin facilitated multiple ring-like structures that

resemble previously described "raft" structures (47) (**Fig3B, Supplementary movie 1**).

Conversely, expansion of colonies grown on glucose or DDW did not display this bi-

phasic expansion (**Supplementary movie 1**). Bacteria growth on glucose expanded the

least from 8 hours and throughout the experiment, resulting in compact, small colonies

with less ringed structures, suggesting that glucose has an inhibitory effect on *F.*

johnsoniae motility and possibly growth (**Fig3C**). The most significant expansion

observed between 6 to 20 hours was on cellulose, but the colonies grown on

glucomannan and pectin proliferated at later times, and the colonies grown on pectin

reached the greatest intensity at 44 hours (**Fig3C**), thereby surpassing the colonies

grown on other sugar sources (p>0.05,Tukey-Kramer HSD test).

Colonies grown on pectin were characterized by a bi-phasic growth with initial

expansion, followed by more significant growth phase (**Supplementary movie 1**) with

an initial peak in fluorescence at 20 hours and a second peak at 36 hours (**Fig3D**). After

20 hours of growth, total fluorescence was highest in cells grown on pectin,

glucomannan and cellulose, and lowest on glucose (p>0.05,Tukey-Kramer HSD test).

Next, we estimated the velocity of colony expansion on the selected sugars by

measuring the time it took the colonies to cross three radials (3mm, 6mm and 9mm),

and subsequently calculating the mean velocity from circle to circle (**Fig3E**). Initially,

in the first 1.5mm radius the estimated velocity was similar on all substances. Between

1.5-3mm colony expansion on pectin and glucomannan increased. Colony expansion

velocity slowed slightly in the outer circle with pectin, glucomannan still exhibiting the

fastest expansion rate of all the tested sugars (**Fig3F**). Thus, *F. johnsoniae* expanded

faster on pectin then on any of other tested sugars (**Fig3G**). Collectively, we conclude

that growth of *F. johnsoniae* on pectin is bi-phasic. Following an initial lag phase, there

is a rapid expansion phase where the colonies proliferate, followed by a slower growth

phase where cells appear to proliferate less and replicate more and gain biomass. This

replication step is significantly more intense with pectin than with any other carbon

source.

Specific TonB/Sus transducers are expressed in response to growth on pectin

In order to gain deeper insight into the molecular mechanisms associated with flavobacteria colony expansion on pectin, we conducted a proteomic assay in which we examined differential intracellular protein expression after 44 hr growth between *F. johnsoniae* growing on rich media coated with pectin or DDW. Eighty-three proteins were found to be upregulated and forty-three were down regulated in the presence of pectin (**TableS1**). Genomic annotation with various tools revealed a high proportion of unassigned proteins, while a large fraction of the proteins (17%, 22% and 37% of KEGG, SEED and EggNog annotations, respectively) were associated with carbohydrate metabolism (**FigS3A-C**). Of the 25 most markedly pectin-induced proteins identified, 13 were involved in polysaccharide uptake, processing and metabolism, including four Sus C/D related proteins (**Fig4A**). Other pectin-induced proteins included a novel transcriptional regulator (upregulated 12-fold) and a protein associated with Auxin-regulation (upregulated 26-fold) (**Table S1**). Interestingly, none of the differentially expressed proteins were gliding related (**Table S2**).

Of the 44 previously-described SusC and 42 SusD homologues identified in the *F. johnsoniae* genome (48), 27 SusC and 15 SusD proteins were detected in the proteomic analysis, in addition to 610 flanking genes surrounding these Sus proteins that constitute PUL clusters seemingly associated with glycan metabolism. Of these, three SusC and six SusD proteins along with 31 associated PUL proteins (forming 4 gene clusters **Table S3**) were significantly induced in response to growth on pectin (**Table S4**).

In order to validate the proteomic results, *F. johnsoniae* cells were grown on pectin and DDW, and the expression of 8 genes (**Table S5**) was evaluated by quantitative real time PCR. Pectin did not facilitate expression of RemA involved in gliding adhesion (not evaluated in the proteomics since it's an extra-membranal protein), indicating that this protein is not associated with the induced colony expansion of flavobacteria on pectin

at the tested time point. Expression of genes encoding for the novel transcriptional regulator (Trans. Regul), Pectate lyase (Pec lyase), putative Auxin-regulated annotated protein (Auxin) and TonB 2144 was also not augmented by pectin despite the fact that they were significantly induced in the proteomic analysis. Among the examined TonB/SusC related genes, TonB 260 and Sus73 were significantly upregulated (60-100 fold, $p<0.05$ Tukey-Kramer HSD test) while TonB 445 was substantially upregulated (10-20 fold) but not statistically significant (**Fig4B**). TonB260 and Sus73 are part of an 18-gene cluster, of which all of the proteins were upregulated in the presence of pectin in the proteomic analysis (**cluster 1-Table S3**). Using two different prediction tools, the TonB260 and Sus73 encoding genes were mapped to the same operon together with a gene encoding for the hydrolytic enzyme polygalacturonase that cleaves the α (1-4) bonds between adjacent galacturonic acids within the homogalacturonic acid backbone of pectin (**Table S6**).

Discussion

Roots modulate their microbiome by secretion of exudates, mucilage and other components that can attract or deter bacteria (8, 47, 49). Specific root exudates were found to serve as chemoattractants for microbes colonize the roots (50). However, this seems to be the tip of the iceberg, and little is known about the plant-associated factors that recruit bacteria from the ecologically complex rhizosphere to the surface and endosphere of roots.

Bacterial plant root colonization is a multi-stage process that involves recognition, movement towards plant/seed attractants and attachment (15). Once attached, persistence and propagation on plant roots is dependent on metabolic fitness (51, 52)**,** resistance to plant protective mechanisms and inter-microbial interactions. The

abundance of bacteria in rhizosphere soil is about 10^2 higher than in the rhizoplane and

root endosphere (53, 54). This exemplifies the extreme selective pressure of this highly

competitive environment and the fitness requirements of the root-colonizing

microbiome.

Members of the genus *Flavobacterium* are highly abundant and substantially enriched

(relative to the rhizosphere soil) on, and within roots of several plant species (25, 32,

33, 55, 56), and certain strains have been found to augment plant health and suppress

pathogens in a variety of plants (29, 32, 33, 35). In this study, we explored the effect of

plant-derived sugars on surface expansion of flavobacteria, in attempt to elucidate

mechanisms associated with plant root-flavobacterial interactions. It was previously

shown that *F. johnsoniae* can effectively grow on polysaccharides such as pectin and

glucomannan as single carbon sources (48). Collectively, results from this study

demonstrate that flavobacteria not only metabolize complex plant cell wall-derived

glycans such as pectin, glucomanan and cellulose, but that these sugars (particularly

pectin) serve as cues for rapid spread of flavobacteria over solid surfaces. We postulate

that this plant cell wall glycan recognition-uptake mechanism is critical for colonization

and proliferation of flavobacteria on plant roots.

Interestingly, our results demonstrated that even when carbon was not limited, plant

cell wall glycans and especially pectin, facilitated expansion of terrestrial flavobacterial

strains over solid surfaces. While the specific correlation between expansion on pectin

and the unique flavobacterial gliding motility mechanism was not determined in this

study, it is evident that the latter is required, because ΔgldJ mutants did not expand on

pectin. Propagation on pectin occurred in a concentration dependent manner, and the

response was specifically dependent on the mother compound, as the response did not

occur with bacteria grown on the pectin monomers D-galacturonic acid and L-

rhamnose. In contrast, growth on glucose was found to inhibit propagation of flavobacteria. This is supported by previous studies, which demonstrated that specific sugars such as glucose, sucrose and lactose can reversibly inhibit *F. johnsoniae* growth on minimal media and inhibits creation of raft microstructures, despite previous studies showing that glucose can served as a sole carbon source, and that it does not prevent gliding (47).

Patterns of *F. johnsoniae* colony expansion was dictated by the carbon source supplemented to the growth media. Preferable expansion of Flavobacteria occurred on plant-derived polysaccharides (pectin and glucomanan) creating a patchy pattern without uniform and dense coverage. The rapid expansion of flavobacteria on pectin was strongly linked to upregulation of TonB-related proteins, which undoubtedly assist in binding (recognition), uptake and metabolism of this glucan. Therefore, we hypothesize that for root-associated flavobacteria, pectin is not merely a carbon source but also an environmental cue. To the best of our knowledge, this is the first time that a specific environmental factor was found to be specifically linked to flavobacteria colony growth.

There are only few examples of plant cell wall-associated polysaccharides that facilitate root bacteria colonization. Flexibacter FS-1 is not able to glide on agarose but can glide on 1% Pectin, while galactronic acid does not induce similar effect (57). Purified Arabidopsis polysaccharides (arabinogalactan, pectin, or xylan) triggered biofilm and pellicle formation in *B. subtilis* when added to rich media and induced root colonization (6). Similarly, amendment with the PGPR *Bacillus amyloliquefaciens* strain SQY 162 with pectin and sucrose, increased bacterial abundance, induced biofilm formation and improved the ability to suppress tobacco bacterial wilt disease (58). In the symbiotic

nitrogen fixing bacterium *Rhizobium leguminosarum*, glucomannan-mediated

attachment was important for legume infection and nodulation (12).

We did not observe any evidence of chemotactic behavior towards pectin, and therefore

postulate that propagation on pectin requires direct contact between the flavobacteria

and the glycan. Based on proteomics and subsequent gene expression validation using

qPCR, we believe that this phenomenon is dictated by the induction of TonB-associated

F. johnsoniae PULs, suggesting a link between niche recognition, colony expansion

and metabolic fitness. Similar induction of TonB and PUL was observed in marine

flavobacteria as response to phytoplankton blooms characterized in decomposition of

alga-derived organic matter (59).

Their high energetic cost and substrate specificity of TonB transducers explains why

they were induced by pectin and not constitutively expressed (60, 61). Nonetheless, the

specific regulatory mechanism responsible for facilitating this response and the

subsequent linkage to the induced gliding motility in flavobacteria remain an enigma.

This is especially true in view of the fact that pectin did not induce expression of RemA,

the lectin binding, flavobacterial cell surface adhesin that is a fundamental part of the

gliding machinery (62). The gliding machinery might be involved in the earlier phase

of the response to pectin, in which we observed intensive bacterial motility. This

assumption can be examined by analysis of earlier proteomic profiles in the future.

Previous experiments showed that *Pseudomonas putida* mutants that lacked TonB,

were deficient in their capacity to uptake iron and displayed impaired seed colonization,

again linking TonB to metabolic and functional fitness (63). However, knock-out of

specific TonB genes might be less effective in *F. johnsoniae* due to the multitude of

predicted TonB genes in its genome, which suggests a high level of functional

redundancy (43). Similar functional redundancy of putative cellulolytic activity was shown where multiple cellulolytic enzymes were induced in proteomic analysis of the soil bacterium *Cytophaga hutchinsonii* after growth on cellulose (64).

In *Xanthomonas campestris pv. Campestris* pectate sensed by specific TonB-dependent receptor triggered secretion of extracellular polygalacturonate, resulting in pectin degradation and generation of oligogalacturonides (OGA) that are recognized as damage-associated molecular patterns (DAMPs), resulting in the initiation of the plant defensive measures (65). In light of our results and the findings in *X. campestris*, we speculate that sensing, uptake and degradation of pectin by TonB and associated proteins, results in OGA release, and plant immune response activation. This pathway may explain the biocontrol and improved plant disease resistance induced by flavobacteria in various plant types. Further experiments will be needed to support these assumptions.

Beside TonB related proteins, pectin induced additional protein targets (**Table S1**). The up regulation of auxin-regulated protein is especially interesting since Auxin is a major phytohormone responsible for plant growth and development demonstrating again a possible connection between pectin sensing and flavobacterial-plant interactions. The additional upregulation of putative transcriptional regulator may suggest pectin induce down-stream response.

Gene expression (qPCR) analysis did not completely mimic the expression differences found in the proteomics analysis of pectin vs. DDW. The discrepancy between RNA and protein expression might be explained by post-translational modifications or regulation affecting protein stability, degradation and complex formation in bacteria grown on pectin vs. DDW as shown in similar cases (66, 67).

Live-imaging fluorescent microscopy revealed that *F. johnsoniae* growth on solid surfaces was primarily bi-phasic in nature, with an initial phase of rapid expansion, followed by biomass production within the colonized area. It appeared that pectin initially induced colony expansion at the expense of cell growth. This bi-phasic growth pattern resembles previously described models in motile *E. coli*, which depicted an initial expansion phase, where "pioneer" bacteria with high velocity advance in front of the colony, followed by a second phase, where "settler" bacteria grow and replicate locally (16, 68). Interestingly, this bi-phasic phenomenon was most pronounced on pectin, suggesting that it may serve as signal that facilitate the expansion of "pioneer" cells, and later as carbon sources that support proliferation of "settlers".

To summarize, we found that pectin, a prominent plant cell wall polysaccharide, facilitates expansion of flavobacteria on solid surfaces, even in the presence of nutrient-rich media. We postulate that pectin may be a "missing link" in elucidating the mechanisms responsible for rhizosphere competence in flavobacteria, *i.e.* their capacity to efficiently colonize and proliferate in the root niche. Thus, in the root environment, plant cell wall polysaccharides, and specifically pectin, not only serve as a nutrient source for flavobacteria, but also as a cue for colonization and rapid expansion along the root surface. The interaction between pectin (and potentially other root glycans) and flavobacteria is mediated by induction of TonB/SusC operons and other associated PULs that facilitate propagation and metabolism of pectin.

Materials and Methods

Bacterial Strains and growth conditions

Four flavobacterial strains were targeted in this study. *F. johnsoniae* strain UW101 (ATCC17061), a gliding/secretion impaired *F. johnsoniae* strain UW101 gldJ mutant,

Flavobacterium sp. strain F52, isolated from the roots of a greenhouse pepper (69, 70)a

gliding/secretion impaired *F52* strain. *F. johnsoniae* UW101 WT+pAS43 (Flavo GFP)

Flavo-ErytRFlavo-CefR, *F. johnsoniae* UW102-48 (ΔgldJ)+pAS43 (Flavo GFP)

Flavo-ErytR Flavo-CefR were used for microscopic analysis. Erythromycin, 100ug/ml

was added to the media of the GFP labeled bacteria.

Flavobacteria strains were grown in CYE medium (Casitone, 10 mg/ml, yeast extract

at 5mg/ml, and 8 mM $MgSO_4$ in 10 mM Tris buffer [pH 7.6]) at 30°C, as previously

described (71). To observe colony-spreading, bacteria were grown on PY2 agar

medium (2 g peptone, 0.5 g yeast extract,10 mg/ml agar, pH 7.3) at 30°C (72).

Organic amendments to growth media

A suite of sugars and other organic compounds were amended to growth media in

various configurations (as described below), to evaluate motility and growth dynamics

of the selected flavobacterial strains. The following substances were used in this study:

pectin (Sigma P9135), arabinose (Sigma A9462), D(+)glucose (Merck K13506942),

cellulose (Merck K239731), D(-)arbinose (Acros 161451000), , glucomannan

(Megazyme P-GLCML), chitin (Sigma C7170), L-rhamnoae (Sigma 83650), D(+)-

galacturonic acid monohydrate (Sigma 48280-5G-F),. All substances were dissolved to

2% final concentration in DDW. Glucose, arabinose, rhamnose and galactronic acid

(titrated to pH 7) were dissolved and filtered through a0.22-micron filter. Pectin was

dissolved in DDW heated at 80°C, and subsequently filtered on 0.45-micron filters.

Chitin, cellulose, and glucomannan were dissolved in DDW heated to 50°C and then

autoclaved for 20 min.

Flavobacteria growth on various sugars

The selected sugars were amended to the PY2 agar plates in two manners: (i) 10 µl of the selected compounds were thinly smeared along a line projecting outward from the center of the petri dish using a pipetor, to test the effect of specific carbon sources on the directional proliferation of flavobacteria (**Fig2S**) or (ii) 500µl were uniformly smeared over the entire petri dish to test the effect of specific carbon sources on the general proliferation of flavobacteria (**Fig1A, C, Fig2A**). Non-soluble substances such as cellulose and chitin, were vigorously vortexed and dispensed using a cut wide tip. Where two sugars were used (*i.e.* rhamnose+galactronic acid) we added 250µl of each. In all cases, plates were left to dry for overnight after adding organic amendments. Flavobacteria were thawed on CYE media overnight. Then bacteria were diluted in 200ul saline to 0.6-1 OD (5×10^9-1.12×10^{10} cells), vortexed well and 2ul were spotted in the center of PY2 plate covered or streaked with specific sugar/s as indicated above. Plate was left to dry 15min and then grown for 48 hours in 30^{O}C. The colony area or the length of expansion (in cm) were measured using Fiji (73) and statistics was calculated using JMP®, Version Pro 14 . (SAS Institute Inc., Cary, NC, 1989-2019). Differences between length/area were considered as significantly different when $p < 0.05$ in Tukey HSD test unless indicated differently.

RNA extraction

To assess the expression of selected genes in the presence of pectin, total RNA was extracted from *F. johnsoniae* cells grown for 48hours at 30^{O}C on PY2 plates covered with 500µl of DDW or 2% pectin as described above. For each plate, bacteria were suspended in 1 ml cold (4^{O}C) PBS buffer, washed once in cold PBS, centrifuged for 2min at 18,000g , re-suspended in TE buffer supplemented with 0.4mg/ml Lysozyme and then incubated in 10min in room temprature. RNA was subsequently extracted from cells using the TRIzol reagent (TRIzol® InvitrogenTM, #15596026), following the

manufacturer's instructions. Residual DNA was removed from the RNA samples by

digesting with RQ1 DNAse (Promega M6101A) at 37 °C for 40 min. For the real time

experiments, cDNA was synthesized using 50ng of DNAse treated RNA, with 1ul of

random prokaryotic primers (Promega C118A). Synthesis of single strand cDNA was

achieved using ImProm-IITM Reverse-Transcriptase (Promega, Madison, WI, United

States).

The integrity and concentration of the extracted RNA and cDNA, was examined with a

QubitTM 3.0 Fluorometer (Thermo Fisher Scientific, United States) using reagents and

protocols supplied by the manufacturer and electrophoresis of samples on a 0.8%

agarose gel.

Live Imaging Fluorescent Microscopy Experiments.

To visualize the effect of different plant derived sugars on *F. johnsoniae*, we prepare a

24 well plate with 500µl PY2 agar media in each well. 10ul of each examined 2%

substance was gently applied on the well in triplicates, and plate was rotated for

1hour and dried ON in RT. A 30G needle (KDL 16-0310) was used to seed the bacteria

in the center of each well. Microscopic imaging was performed using a NIKON eclipse

Ti microscope (Nikon, Japan) equipped with a ProScan motorized XY stage and an

HF110A system (enabling rapid switching of emission filters)(Prior Scientific, MA,

USA), and a temperature-controlled stage incubator (LAUDA ECO RE 415, Korea).

Bright field illumination was provided by a cool LED pE-100A (Cool LED, UK).

Excitation light for epifluorescence microscopy was provided by a Spectra X light

engine (Lumencor, USA). Imaging was performed using a long working distance 40x

objective (NA 0.6) (Nikon, Japan). Images were captured at 2 hours intervals for 44

hours using an ANDOR zyla 5.5 MP sCMOS camera (Oxford Instruments, UK). For

each frame, bright field (gray), phycocyanin (Ex: 555 nm/Em: 632 nm; pink) and

chlorophyll (Ex: 440 nm/Em: 697 nm; green) were captured separately. Images were

processed using the NIS elements AR 4.6 (64 bit) software package (Nikon, Japan) and

Fiji (73). Fluorescence level was normalized to the initial measured value (to avoid

differences in the initial number of seeded bacteria) and to the maximal fluorescence

on DDW (to reduce variability of GFP fluorescence levels between movies). The

population growth dynamic of GFP- *F. johnsoniae* on each substance computed using

Fiji's time series analyzer plugin, and average fluorescence density profiles of the

expanding population over time was quantified using JMP®, Version Pro 14 . (SAS

Institute Inc., Cary, NC, 1989-2019).

Proteomics analysis

The samples were subjected to in-solution tryptic digestion followed by a desalting

step. The resulting peptides were analyzed using nanoflow liquid chromatography

(nanoAcquity) coupled to high resolution, high mass accuracy mass spectrometry (Q

Exactive HF). Each sample was analyzed on the instrument separately in a random

order in discovery mode. Raw data was searched against the *F. johnsoniae* protein

databases, to which a list of common lab contaminants was added.

Database search was done using the Mascot algorithm, and quantitative analysis was

performed using the Expressionist software from GeneData. As a rule of thumb we

consider significant differences to be >1 peptide per protein, fold change >2 or <0.5,

and <0.05 p-value. Proteins were functionally annotated using RAST (Rapid

Annotations using Subsystems Technology) (74).

Twenty-five pectin-induced proteins were selected based on high fold change between

pectin vs. DDW and statistical significance (proteins without annotation were removed

from this analysis). Each gene abundance across the dataset were normalized to 100%

and a heatmap was created using studio.plot.ly (https://plotly.com/).

Quantitative PCR Assessment of Gene Expression Levels

The expression of 10 genes encoding for proteins found to be significantly induced on

pectin in the proteomic analyses were analyzed using quantitative realtime (qRT)-PCR.

For relative quantification of target genes, primers for qRT-PCR experiments (**Table**

S5) were based on the *F. johnsoniae* genome sequence and were predesigned using the

PrimerQuest® tool (Integrated DNA Technologies -IDT). Triplicate cDNA samples for

each of the treatments (Pectin/DDW) were diluted and 2ng was used in a 20 µl final

reaction volume together with 10ul Fast SYBR^{TM} (Thermo Fisher scientific) green PCR

master mix, 100 nM each forward and reverse primers, DDW and 1ul of template

cDNA. Amplification was carried out on a StepOnePlus real-time PCR thermocycler

using the following program: heat activation of DNA polymerase at 95°C for 3 min and

40 cycles at 95°C for 5 sec for denaturation and primer annealing and extension at 60°C

for 30 sec. A melt curve was produced to confirm a single gene-specific peak and to

detect primer-dimer formation by heating the samples from 60 to 95°C in 0.3°C

increments. Gene amplification was confirmed by detection of a single peak in the melt

curve analysis. No primer-dimer formation was detected. For each gene, PCR gene

amplification was carried out with three independent biological replicates. The relative

quantification of each target gene versus reference gene was determined according to

the method described (Livak and Schmittgen2001). Expression of each gene was

normalized to that of three alternative reference genes (16S rRNA, DNA gyrase subunit

B (EC 5.99.1.3) -*gyr*B (Housekeeping), and Electron transfer Flavoprotein, alpha

subunit, Threonine synthase (EC 4.2.3.1)-ETF), selected because there was no detected

difference in its expression between the pectin and DDW treatments in the proteomic

analysis. Concentrations and ΔΔCT values were calculated and analyzed with the

StepOne software v2.3 (Applied Biosystems, Foster City, CA, United States).

Concomitant "no-RT" reactions, lacking reverse transcriptase, were performed for each

sample and run to confirm absence of DNA contamination, as well as no template

controls (NTCs) to confirm lack of contamination of all used reagents. Efficiency of

reactions was monitored by means of an in internal standard curve using a 10-fold

dilution of DNA ranging from 0.01-10ng of DNA per reaction, done in triplicates.

Reported efficiency was between 92.1 and 97.2% for all primers, and R^2-values were

greater than 0.99. All runs were performed using a StepOnePlus real-time PCR system

(Applied Biosystems, Foster City, CA, United States) and data analysis was conducted

using the StepOne software v2.3 (Applied Biosystems, Foster City, CA, United States).

Acknowledgments: We would like to thank Alla Usyskin-Tonne, for her help with the proteomics functional annotation and Eduard Belausov for his help with the binocular based imaging.

References

1.   Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. Trends Plant Sci 17:478–486.

2.   Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. 2013. Structure and Functions of the Bacterial Microbiota of Plants. Annu Rev Plant Biol 64:807–838.

3.   Reinhold-Hurek B, Bünger W, Burbano CS, Sabale M, Hurek T. 2015. Roots

Shaping Their Microbiome: Global Hotspots for Microbial Activity. Annu Rev

Phytopathol 53:403–424.

4.   Dennis PG, Miller AJ, Hirsch PR. 2010. Are root exudates more important than

other sources of rhizodeposits in structuring rhizosphere bacterial

communities? FEMS Microbiol Ecol 72:313–327.

5.   Barret M, Morrissey JP, O'Gara F. 2011. Functional genomics analysis of plant

growth-promoting rhizobacterial traits involved in rhizosphere competence.

Biol Fertil Soils 47:729–743.

6.   Beauregard PB, Chai Y, Vlamakis H, Losick R, Kolter R. 2013. Bacillus

subtilis biofilm induction by plant polysaccharides. Proc Natl Acad Sci

110:E1621–E1630.

7.   Massalha H, Korenblum E, Malitsky S, Shapiro OH, Aharoni A. 2017. Live

imaging of root–bacteria interactions in a microfluidics setup. Proc Natl Acad

Sci 114:4549–4554.

8.   Sasse J, Martinoia E, Northen T. 2018. Feed Your Friends: Do Plant Exudates

Shape the Root Microbiome? Trends Plant Sci 23:25–41.

9.   Ofek M, Voronov-Goldman M, Hadar Y, Minz D. 2014. Host signature effect

on plant root-associated microbiomes revealed through analyses of resident vs .

active communities. Environ Microbiol 16:2157–2167.

10.   Rosenberg E., DeLong E.F., Lory S., Stackebrandt E. TF. 2014. The Family

Chitinophagaceae, p. 119–144. *In* E., R, E.F., D, S., L, E., S, F., T (eds.), The

Prokaryotes. Springer, Berlin, Heidelberg.

11.   Rudrappa T, Czymmek KJ, Paré PW, Bais HP. 2008. Root-Secreted Malic

Acid Recruits Beneficial Soil Bacteria. Plant Physiol 148:1547–1556.

12.   Williams A, Wilkinson A, Krehenbrink M, Russo DM, Zorreguieta A, Downie

JA. 2008. Glucomannan-mediated attachment of Rhizobium leguminosarum to

pea root hairs is required for competitive nodule infection. J Bacteriol

190:4706–4715.

13.   Oku S, Ayaka K, Takahisa TK, Junichi YN. 2012. Identification of Chemotaxis

Sensory Proteins for Amino Acids in Pseudomonas fluorescens Pf0-1 and Their

Involvement in Chemotaxis to Tomato Root Exudate and Root Colonization.

Microbes Env 27:462–469.

14.   Webb BA, Compton KK, del Campo JSM, Taylor D, Sobrado P, Scharf BE.

2017. Sinorhizobium meliloti Chemotaxis to Multiple Amino Acids Is

Mediated by the Chemoreceptor McpU. Mol Plant-Microbe Interact 30:770–

777.

15.   Feng H, Zhang N, Fu R, Liu Y, Krell T, Du W, Shao J, Shen Q, Zhang R.

2019. Recognition of dominant attractants by key chemoreceptors mediates

recruitment of plant growth-promoting rhizobacteria. Environ Microbiol

21:402–415.

16.   Cremer J, Honda T, Tang Y, Wong-Ng J, Vergassola M, Hwa T. 2019.

Chemotaxis as a navigation strategy to boost range expansion. Nature 575:658–

663.

17.   Lugtenberg BJJ, Dekkers LC. 1999. What makes Pseudomonas bacteria

rhizosphere competent? Environ Microbiol 1:9–13.

18.   Yan Y, Yang J, Dou Y, Chen M, Ping S, Peng J, Lu W, Zhang W, Yao Z, Li H,

Liu W, He S, Geng L, Zhang X, Yang F, Yu H, Zhan Y, Li D, Lin Z, Wang Y,

Elmerich C, Lin M, Jin Q. 2008. Nitrogen fixation island and rhizosphere

competence traits in the genome of root-associated Pseudomonas stutzeri

A1501. Proc Natl Acad Sci 105:7564–7569.

19.   Pieterse MJ, Zamioudis C, Berendsen RL, Weller DM, Wees SCM Van,

Bakker PAHM. 2014. Induced Systemic Resistance by Beneficial Microbes.

20.   McBride MJ, Liu W, Lu X, Zhu Y, Zhang W. 2014. The Family

Cytophagaceae, p. 577–593. *In* The Prokaryotes. Springer Berlin Heidelberg,

Berlin, Heidelberg.

21.   Kolton M, Sela N, Elad Y, Cytryn E. 2013. Comparative Genomic Analysis

Indicates that Niche Adaptation of Terrestrial Flavobacteria Is Strongly Linked

to Plant Glycan Metabolism. PLoS One 8:e76704.

22.   Janssen PH. 2006. Identifying the Dominant Soil Bacterial Taxa in Libraries of

16S rRNA and 16S rRNA Genes. Appl Environ Microbiol 72:1719–1728.

23.   Manter DK, Delgado JA, Holm DG, Stong RA. 2010. Pyrosequencing Reveals

a Highly Diverse and Cultivar-Specific Bacterial Endophyte Community in

Potato Roots. Microb Ecol 60:157–166.

24.   Johansen JE, Binnerup SJ, Lejbolle KB, Mascher F, Sorensen J, Keel C. 2002.

Impact of biocontrol strain Pseudomonas fluorescens CHA0 on rhizosphere

bacteria isolated from barley (Hordeum vulgare L.) with special reference to

Cytophaga-like bacteria. J Appl Microbiol 93:1065–1074.

25.   Kolton M, Meller Harel Y, Pasternak Z, Graber ER, Elad Y, Cytryn E. 2011.

Impact of Biochar Application to Soil on the Root-Associated Bacterial

Community Structure of Fully Developed Greenhouse Pepper Plants. Appl

Environ Microbiol 77:4924–4930.

26.   Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S,

Tremblay J, Engelbrektson A, Kunin V, Rio TG del, Edgar RC, Eickhorst T,

Ley RE, Hugenholtz P, Tringe SG, Dangl JL. 2012. Defining the core

Arabidopsis thaliana root microbiome. Nature 488:86–90.

27.   Bodenhausen N, Horton MW, Bergelson J. 2013. Bacterial Communities

Associated with the Leaves and the Roots of Arabidopsis thaliana. PLoS One

8:e56329.

28.   Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y,

McHardy AC, Schulze-Lefert P. 2015. Structure and Function of the Bacterial

Root Microbiota in Wild and Domesticated Barley. Cell Host Microbe 17:392–

403.

29.   Sang MK, Kim KD. 2012. The volatile-producing Flavobacterium johnsoniae

strain GSE09 shows biocontrol activity against Phytophthora capsici in pepper.

J Appl Microbiol 113:383–398.

30.   Gunasinghe RN, Ikiriwatte CJ, Karunaratne AM. 2004. The use of Pantoea

agglomerans and Flavobacterium sp . to control banana pathogens. J Hortic Sci

Biotechnol 79:1002–1006.

31.   Kolton M, Frenkel O, Elad Y, Cytryn E. 2014. Potential Role of Flavobacterial

Gliding-Motility and Type IX Secretion System Complex in Root Colonization

and Plant Defense. Mol Plant-Microbe Interact 27:1005–1013.

32.   Xue C, Ryan Penton C, Shen Z, Zhang R, Huang Q, Li R, Ruan Y, Shen Q.

2015. Manipulating the banana rhizosphere microbiome for biological control

of Panama disease. Sci Rep 5:11124.

33.   Kwak M-J, Kong HG, Choi K, Kwon S-K, Song JY, Lee J, Lee PA, Choi SY,

Seo M, Lee HJ, Jung EJ, Park H, Roy N, Kim H, Lee MM, Rubin EM, Lee S-

W, Kim JF. 2018. Rhizosphere microbiome structure alters to enable wilt

resistance in tomato. Nat Biotechnol 36:1100–1109.

34.   Carrión VJ, Perez-Jaramillo J, Cordovez V, Tracanna V, de Hollander M, Ruiz-

Buck D, Mendes LW, van Ijcken WFJ, Gomez-Exposito R, Elsayed SS,

Mohanraju P, Arifah A, van der Oost J, Paulson JN, Mendes R, van Wezel GP,

Medema MH, Raaijmakers JM. 2019. Pathogen-induced activation of disease-

suppressive functions in the endophytic root microbiome. Science (80-)

366:606–612.

35.   Wei Z, Gu Y, Friman V-P, Kowalchuk GA, Xu Y, Shen Q, Jousset A. 2019.

Initial soil microbiome composition and functioning predetermine future plant

health. Sci Adv 5:eaaw0759.

36.   Hebbar P, Berge O, Heulin T, Singh SP. 1991. Bacterial antagonists of

Sunflower (Helianthus annuus L.) fungal pathogens. Plant Soil 133:131–140.

37.   Sang MK, Kim J Do, Kim BS, Kim KD. 2011. Root Treatment with

Rhizobacteria Antagonistic to Phytophthora Blight Affects Anthracnose

Occurrence, Ripening, and Yield of Pepper Fruit in the Plastic House and Field.

Phytopathology 101:666–678.

38.   Alexander BJR, Stewart A. 2001. Glasshouse screening for biological control

agents of Phytophthora cactorum on apple (Malus domestica). New Zeal J

Crop Hortic Sci 29:159–169.

39.   McBride MJ, Nakane D. 2015. Flavobacterium gliding motility and the type IX

641   secretion system. Curr Opin Microbiol 28:72–77.

642   40.   Kharade SS, McBride MJ. 2015. Flavobacterium johnsoniae PorV Is Required

643   for Secretion of a Subset of Proteins Targeted to the Type IX Secretion System.

644   J Bacteriol 197:147–158.

645   41.   Lauber F, Deme JC, Lea SM, Berks BC. 2018. Type 9 secretion system

646   structures reveal a new protein transport mechanism. Nature 564:77–82.

647   42.   Doug Hyatt, Gwo-Liang Chen, Philip F LoCascio, Miriam L Land, , Frank W

648   Larimer LJH. 2010. Integrated nr Database in Protein Annotation System and

649   Its Localization. Nat Commun 6:1–8.

650   43.   McBride MJ, Xie G, Martens EC, Lapidus A, Henrissat B, Rhodes RG,

651   Goltsman E, Wang W, Xu J, Hunnicutt DW, Staroscik AM, Hoover TR, Cheng

652   Y-Q, Stein JL. 2009. Novel Features of the Polysaccharide-Digesting Gliding

653   Bacterium Flavobacterium johnsoniae as Revealed by Genome Sequence

654   Analysis. Appl Environ Microbiol 75:6864–6875.

655   44.   Foley MH, Cockburn DW, Koropatkin NM. 2016. The Sus operon: a model

656   system for starch uptake by the human gut Bacteroidetes. Cell Mol Life Sci

657   73:2603–2617.

658   45.   Shrivastava A, Johnston JJ, Van Baaren JM, McBride MJ. 2013.

659   Flavobacterium johnsoniae GldK, GldL, GldM, and SprA are required for

660   secretion of the cell surface gliding motility adhesins sprb and remA. J

661   Bacteriol 195:3201–3212.

46.   Johnston JJ, Shrivastava A, McBride MJ. 2018. Untangling Flavobacterium

johnsoniae gliding motility and protein secretion. J Bacteriol 200.

47. Gorski L, Godchaux W, Leadbetter ER. 1993. Structural specificity of sugars

that inhibit gliding motility of Cytophaga johnsonae. Arch Microbiol 160:121–

125.

48. McBride MJ, Xie G, Martens EC, Lapidus A, Henrissat B, Rhodes RG,

Goltsman E, Wang W, Xu J, Hunnicutt DW, Staroscik AM, Hoover TR, Cheng

YQ, Stein JL. 2009. Novel features of the polysaccharide-digesting gliding

bacterium Flavobacterium johnsoniae as revealed by genome sequence

analysis. Appl Environ Microbiol 75:6864–6875.

49. Wood DC, Hayasaka SS. 1981. Chemotaxis of rhizoplane bacteria to amino

acids comprising eelgrass (Zostera marina L.) root exudate. J Exp Mar Bio

Ecol 50:153–161.

50. Raina J-B, Fernandez V, Lambert B, Stocker R, Seymour JR. 2019. The role of

microbial motility and chemotaxis in symbiosis. Nat Rev Microbiol 17:284–

294.

51. Walker TS, Bais HP, Grotewold E, Vivanco JM. 2003. Root Exudation and

Rhizosphere Biology: Fig. 1. Plant Physiol 132:44–51.

52. RodrÃ-guez-Navarro DN, Dardanelli MS, RuÃ-z-SaÃ-nz JE. 2007.

Attachment of bacteria to the roots of higher plants. FEMS Microbiol Lett

272:127–136.

53. Curtis TP, Sloan W T. 2005. MICROBIOLOGY: Exploring Microbial

Diversity--A Vast Below. Science (80-) 309:1331–1333.

54. Lagos L, Maruyama F, Nannipieri P, Mora M., Ogram A, Jorquera M. 2015.

Current overview on the study of bacteria in the rhizosphere by modern

molecular techniques: a mini‒review. J soil Sci plant Nutr 15:0–0.

55. Bulgarelli D, Rott M, Schlaeppi K, Ver E, Themaat L Van, Ahmadinejad N,

Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner

FO, Amann R, Eickhorst T, Schulze-lefert P. 2012. Revealing structure and

assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature

488:91–95.

56. Bergelson J, Mittelstrass J, Horton MW. 2019. Characterizing both bacteria and

fungi improves understanding of the Arabidopsis root microbiome. Sci Rep

9:24.

57. Arlauskas J, Burchard RP. 1982. Substratum requirements for bacterial gliding

motility. Arch Microbiol 133:137–141.

58. Guan H, Pan B, Fang Z, Gong M, Shen B, Shen Q, Guo R, Wu K, Shi W, Yuan

S. 2015. Pectin Enhances Bio-Control Efficacy by Inducing Colonization and

Secretion of Secondary Metabolites by Bacillus amyloliquefaciens SQY 162 in

the Rhizosphere of Tobacco. PLoS One 10:e0127418.

59. Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM,

Kassabgy M, Huang S, Mann AJ, Waldmann J, Weber M, Klindworth A, Otto

A, Lange J, Bernhardt J, Reinsch C, Hecker M, Peplies J, Bockelmann FD,

Callies U, Gerdts G, Wichels A, Wiltshire KH, Glockner FO, Schweder T,

Amann R. 2012. Substrate-Controlled Succession of Marine Bacterioplankton

Populations Induced by a Phytoplankton Bloom. Science (80-) 336:608–611.

60. Postle K, Kadner RJ. 2003. Touch and go: Tying TonB to transport. Mol

Microbiol 49:869–882.

61.   Postle K. 2007. TonB System, In Vivo Assays and Characterization, p. 245–

269. *In* Methods in Enzymology.

62.   Shrivastava A, Rhodes RG, Pochiraju S, Nakane D, McBride MJ. 2012.

Flavobacterium johnsoniae RemA is a mobile cell surface lectin involved in

gliding. J Bacteriol 194:3678–3688.

63.   Molina MA, Godoy P, Ramos-Gonzalez MI, Munoz N, Ramos JL, Espinosa-

Urgel M. 2005. Role of iron and the TonB system in colonization of corn seeds

and roots by Pseudomonas putida KT2440. Environ Microbiol 7:443–449.

64.   Taillefer M, Arntzen MØ, Henrissat B, Pope PB, Larsbrink J. 2018. Proteomic

Dissection of the Cellulolytic Machineries Used by Soil-Dwelling

Bacteroidetes. mSystems 3:1–16.

65.   Vorhölter F-J, Wiggerich H-G, Scheidle H, Sidhu VK, Mrozek K, Küster H,

Pühler A, Niehaus K. 2012. Involvement of bacterial TonB-dependent

signaling in the generation of an oligogalacturonide damage-associated

molecular pattern from plant cell walls exposed to Xanthomonas campestris pv.

campestris pectate lyases. BMC Microbiol 12:239.

66.   Flory MR, Lee H, Bonneau R, Mallick P, Serikawa K, Morris DR, Aebersold

R. 2006. Quantitative proteomic analysis of the budding yeast cell cycle using

acid-cleavable isotope-coded affinity tag reagents. Proteomics 6:6146–6157.

67.   Liu Y, Beyer A, Aebersold R. 2016. On the Dependency of Cellular Protein

Levels on mRNA Abundance. Cell 165:535–550.

68.   Liu W, Cremer J, Li D, Hwa T, Liu C. 2019. An evolutionarily stable strategy

to colonize spatially extended habitats. Nature 575:664–668.

69.   Kolton M, Frenkel O, Elad Y, Cytryn E. 2014. Potential Role of Flavobacterial

Gliding-Motility and Type IX Secretion System Complex in Root Colonization

and Plant Defense. Mol Plant-Microbe Interact 27:1005–1013.

70.   Sela N, Cytryn E, Green SJ, Harel YM, Kolton M, Elad Y. 2012. Draft

Genome Sequence of Flavobacterium sp. Strain F52, Isolated from the

Rhizosphere of Bell Pepper (Capsicum annuum L. cv. Maccabi). J Bacteriol

194:5462–5463.

71.   Mcbride MJ, Baker SA. 1996. Development of techniques to genetically

manipulate members of the genera Cytophaga, Flavobacterium, Flexibacter,

and Sporocytophaga. Appl Environ Microbiol 62:3017–3022.

72.   Agarwal S., Hunnicutt DW, McBride MJ. 2001. Cloning and Characterization

of theFlavobacterium johnsoniae Gliding Motility GenesgldD and gldE. J

Bacteriol 183:4167–4175.

73.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J, White DJ, Hartenstein

V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for

biological-image analysis. Nat Methods 9:676–682.

74.   Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA,

Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R.

2014. The SEED and the Rapid Annotation of microbial genomes using

Subsystems Technology (RAST). Nucleic Acids Res 42:D206–D214.

75.   Buchfink B, Xie C, Huson DH. 2014. Fast and sensitive protein alignment

using DIAMOND. Nat Methods 12:59–60.

76.   Jensen LJ, Julien P, Kuhn M, von Mering C, Muller J, Doerks T, Bork P. 2007.

757   eggNOG: automated construction and annotation of orthologous groups of

758   genes. Nucleic Acids Res 36:D250–D254.

759   77.   Taboada B, Estrada K, Ciria R, Merino E. 2018. Operon-mapper: a web server

760   for precise operon identification in bacterial and archaeal genomes.

761   Bioinformatics 34:4118–4120.

762   78.   Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM,

763   Solovyev V V, Rubin EM, Rokhsar DS, Banfield JF. 2004. Community

764   structure and metabolism through reconstruction of microbial genomes from

765   the environment. Nature 428:37–43.

 Figures legend

 Fig1- Impact of sugars on proliferation of flavobacteria. Image (A) and quantification (B)

 of colony area of Flavobacteria colony expansion on PY2 agar amended on different mono-

 and polysaccharides after 48hr incubation at 30°C (N=4). Image (C) and quantification (D) of

 colony expansion on PY2 agar of wild type (WT) and gliding mutants (ΔgldJ) of *F. johnsoniae*

 and *Flavobacterium* sp. F52, amended with DDW or 2% pectin after 48hr incubation at 30°C

 (N=3). Colony area was measured using Fiji. Statistics significance calculated using ®JMP

 Pro14, was considered significant if p<0.05 by Tukey HSD.

 Fig2 - Impact of pectin precursors (galactronic acid and rhamnose) on

 proliferation of flavobacteria. (A) *F. johnsoniae* colony expansion on PY2 agar

 amended with DDW, pectin galactronic acid, rhamnose and galactronic acid and

 rhamnose incubated at 30°C for 48 h (N=4). (B) Graphic description of colony area

 based on results from (A). Statistics significance was calculated using ®JMP Pro14,

 and means was considered significant when p<0.05 by Tukey HSD

 Fig3 - Temporal dynamics of flavobacterial proliferation on different sugars using

 live imaging microscopy. (A) Morphology of GFP labeled *F. johnsoniae* colonies on

 selected sugars. Bacteria were inoculated in the center of PY2 agar coated with the

 indicated sugars (schematically described in the insert). Images show colony

 morphology after 32hr. (B) Enlarged image of GFP-*F. johnsoniae* colony morphology

 after 20hr of growth on the selected sugars as indicated in (A). (C) Growth rates of

 GFP-*F. johnsoniae* colonies on the selected sugars. Data was normalized as described

 in the materials and methods section. Differences in the average colony fluorescence

intensity after 44 hr was compared and considered significant if p<0.05 by Tukey HSD.

Data includes means and data from three biological replicates composed of three

technical repeats in each. (D) Temporal dynamics of GFP-*F. johnsoniae* growth rates.

Growth was compared at the peaks (20hr and 36hr) and considered significant if p<0.05

by Tukey HSD. (E) Schematic diagram showing the three characterized regions of

interest (ROI- 1.5, 3 and 4.5 mm radii) used to evaluate of bacterial expansion rates. (F)

Estimated expansion rates of GFP-*F. johnsoniae* on the selected sugars. Velocity of

bacterial movement was estimated by expansion time in hours taken to cross known

ROIs as indicated in E. Statistical significance is p<0.05 by Tukey HSD. (G) Estimated

expansion time relative to estimated growth of GFP-*F. johnsoniae* on the selected

sugars. Colored circles mark time (h) for bacteria to cross the 4.5mm radius on each

substance as calculated in F.

Fig4 - Pectin induced flavobacterial genes and proteins. (A) Differential expression

of the 25 most substantial pectin induced proteins based on proteomic analysis of *F.*

johnsoniae colonies grown on CYE medium amended with pectin relative to colonies

grown on identical media amended with DDW. Heat map shows triplicates for each

treatment. All described proteins are statistically significant (p<0.05). (B) Expression

of selected genes (sus73, tonB260, tonB445, tonB 2144, auxin regulator,

Transcriptional regulator, pectate lyase and remA) shown to be induced in the

proteomic analysis described in (A), using quantitative real-time PCR (qPCR). Fold

changes in mRNA levels of the target genes were normalized against the 16SrRNA

gene (left), the Electron transfer Flavoprotein, alpha subunit (center), and the DNA

gyrase subunit B (right). Change in target genes fold change RNA expression was

calculated using the $2^{-\Delta\Delta CT}$ method and statistical significance (p<0.05) by Student T-

test. Error bars represent standard errors of six independent experiments based on two

independent RNA extractions.

A.

DDW

Arabinose

Glucose

Cellulose

Pectin

1cm

B.

C.

DDW   Pectin

WT *F. johnsoniae*

ΔgldJ *F. johnsoniae*

WT *sp. F52*

ΔgldJ *sp. F52*

1cm

D.

■ F. johnsoniae WT
■ F. johnsoniae ΔgldJ
■ F.52 WT
□ F.52 ΔgldJ

DDW

Pectin

Galacturonic acid

Rhamnose

Rhamnose
+ Galacturonic acid

1 cm

A.

TonB related

Carbohydrate metabolism related

B.