Title: I-KCKT allows dissection-free RNA profiling of adult *Drosophila*

intestinal progenitor cells

Authors: Kasun Buddika*, Jingjing Xu*, Ishara Ariyapala, and Nicholas S. Sokol

Affiliation: Department of Biology, Indiana University, Bloomington, IN 47405,

USA

* Co-first authors.

Correspondence: nsokol@indiana.edu (email)

16   812-856-6812 (phone)

20 **Running Title:** I-KCKT RNA profiling

Abstract

The adult Drosophila intestinal epithelium is a model system for stem cell biology, but its utility is limited by current gene expression profiling methods, particularly as applied to RNA-binding proteins. Here, we describe a new profiling method that relies upon a novel intestinal progenitor Gal4 driver, termed intestinal kickout (I-KCKT)-Gal4, that is not expressed outside the gastrointestinal tract. This method uses UV cross-linked whole animal frozen powder as its starting material to specifically immunoprecipitate the RNA cargoes of transgenic epitope-tagged RNA binding proteins expressed under I-KCKT-Gal4 control. When applied to the general mRNA binder, polyA-binding protein (PABP), we show that RNA profiles obtained by this method significantly overlap with RNA-seq profiles of progenitor cells obtained via cell sorting and have very low background noise when compared to other datasets, even though we used whole animals as starting material. We further apply this approach to two specific RNA binders, LIN28 and FMRP, and identify intestinal targets for each. We expect that this approach will be broadly applicable to the field, and enable the RNA profiling of wildtype and mutant intestinal progenitor cells from intact flies subjected to a variety of environmental conditions.

Introduction

The adult *Drosophila* intestine has become a premier model for understanding the biology and behavior of resident stem cells in their native context [1]. One key approach has been gene expression profiling of intestinal cell types to characterize molecular profiles associated with cell identity, maintenance, and differentiation under both homeostatic as well as perturbed conditions (see, for example, refs [2, 3]). This approach has relied on the manual dissection of intestines. Manual dissection is not only laborious and time-consuming but also precludes certain types of analysis that require large amounts of intact, non-degraded starting material, such as immunoprecipitation (IP). Methods such as RiboTag, for example, that rely upon the sequencing of immunoprecipitates associated with RNA-binding proteins (RBPs) to analyze post-transcriptional control mechanisms, are not currently feasible in adult Drosophila intestinal stem cells but have been critical for characterizing identity and differentiation mechanisms in other stem cell lineages [4-6]. There is therefore a need for improved methods to profile this stem cell population in Drosophila [7].

Manual dissection of intestines has been required because of the lack of tools to exclusively label and manipulate individual intestinal cell types. Such tools would enable the use of the whole fly as starting material, allowing the rapid production of large amounts of starting material. Gal4 drivers have been described that label the various individual cell types in the intestinal stem cell lineage, including the intestinal stem cells (ISCs) themselves as well as their transient progenitor cell daughters, termed enteroblasts (EB), and two differentiated cell types, enterocytes (ECs) and enteroendocrine cells (EEs) [8-10]. Such Gal4 drivers can be used to express epitope-tagged transgenic proteins specifically in these various intestinal cell types that can be immunopurified. However, these Gal4 drivers are either known or likely to be active elsewhere in the adult [11].

Here, we describe a method that uses a new intestinal progenitor Gal4 driver that is not expressed outside the intestine to profile the general transcriptome as well as specific RBP cargoes expressed in this cell type. We reasoned that an intersectional

approach could be used to design such a Gal4 driver. Intersectional methods enable the

targeting of transgenic expression to the cells in which two different enhancers are both

active. In a recombinase-mediated intersectional method, for example, the recombinase

under the control of one enhancer activates a Gal4 driver under the control of a second

enhancer via recombinase-mediated removal of an intervening stop cassette (Fig 1A).

We reasoned that such a recombinase-based method involving a pan-intestinal

enhancer and a progenitor enhancer might limit expression to only intestinal progenitor

cells. We designed our transgenes with the KD recombinase (KDR), which was recently

shown to mediate the excision, or "kick-out", of sequences between KDR target (KDRT)

sites in *Drosophila* cells [12]. We chose to use KDR so that the resulting system could

be used in tandem with other recombinases, like FLP, for additional manipulation of

intestinal cells. Because this system is designed for intestine-specific activation of a

"kick-out" transgene, we referred to it as Intestinal-KiCK-ouT, or I-KCKT.

Results

I-KCKT-Gal4 labels most intestinal progenitor cells

While progenitor enhancer sequence has previously been identified in the *miranda* (*mira*) gene locus [13], the availability of a defined pan-intestinal enhancer was less clear. Defined enhancers for some intestine-specific genes have been reported (e.g. *mex1-gal4*, *npc1b-gal4*, etc. [14, 15]), but careful inspection indicated that these were either not expressed throughout the intestine or in all cell types. We therefore took a candidate gene approach to identify intestine-specific genes based on gene expression data reported in FlyAtlas [16], and then tested DNA fragments from those genes for intestinal activity. KDR transgenes containing putative enhancer sequences from either *βTrypsin* (*βTry*), *θTrypsin* (*θTry*), *κTrypsin* (*κTry*), *CG10116*, or *CG18404*, were tested for intestinal activity in flies also harboring two additional transgenes, a *tubulin-KDRT-stop-KDRT-Gal4::p65* "kickout" transgene and a *UAS-6XGFP* responder [17]. GFP expression was monitored in both the larval and adult intestine for all resulting strains except for *θTry* (Fig 1B-M); in its case, only larvae were analyzed because adults of the proper genotype failed to eclose. While all strains displayed some GFP expression in the larval intestine, most GFP patterns were patchy and non-uniform. In the adult intestine, however, two KDR lines, *βTry*-KDR and CG10116-KDR, displayed expression throughout the tissue (Fig 1H, M). Careful inspection of these two indicated that CG10116-KDR was expressed in most cells throughout the midgut, raising the possibility that the associated enhancer fragment was pan-intestinal-specific and could be used to exclusively label adult intestinal progenitor cells.

To test this possibility, we evaluated *CG10116-KDR* activity in the intestinal progenitor cells of flies harboring a *mira*-containing stop cassette transgene as well as a UAS-reporter that drove a nuclear-localized GFP, *UAS-stinger-GFP* [18]. For these experiments, we analyzed strains containing two different *mira*-transgenes, one containing an enhanced version of Gal4 with the p65 transcriptional activation domain (*I-KCKT^{p65}*) and one with a non-modified version of Gal4 (*I-KCKT*) that could be used in conjunction with the Gal80-dependent temporal and regional gene expression targeting (TARGET) system for conditional expression [19]. To analyze CG10116-KDR-mediated

expression in progenitor cells, we compared GFP expression with that of a previously

generated and validated progenitor reporter, *mira-His2A.mCherry.HA* [7]*,* in each of the

five intestinal regions (Fig 2A-J). Quantification of the percentage of mCherry+ cells that

were also GFP+ indicated that ~100% of progenitor cells were labeled in all regions of *I-*

KCKT strains (Fig 2P, Q). The only two exceptions to this trend were regions 1 and 3 of

I-KCKT^{p65} intestines, where only 58.7±15.5% (n=5) and 64.4±17.0% (n=5) of progenitor

cells were labeled, respectively. Importantly, we also found that no GFP+ cells were

mCherry- in *I-KCKT^{p65}* and *I-KCKT* strains indicating that no non-progenitor cells were

labeled in *I-KCKT* strains. To confirm this analysis, we also compared GFP expression

in *I-KCKT* to a second, validated progenitor reporter, *esg-LacZ* [9], and found similarly

that ~100% of LacZ+ cells were also GFP+ and that 0% of GFP+ cells were LacZ- (Fig

S1). The only exception was region 3, where ~69.9±9.0% of LacZ+ cells were GFP+,

indicating that the *miranda* enhancer labels a subset of progenitors in this region.

Collectively, this analysis indicated that almost all progenitor cells, but few if any other

cells, were labeled in the intestines of *I-KCKT* strains.

I-KCKT-Gal4 is not detected in non-intestinal tissue

134   To determine whether intestinal progenitor cells were exclusively labeled in *I-*

KCKT adults, we performed two analyses. First, we visually inspected *UAS-6XGFP-*

bearing adults for GFP expression. *I-KCKT* strains displayed prominent GFP

fluorescence in the intestine but little, if any, elsewhere (Fig 3A, B). For comparison, we

analyzed in parallel three other widely used drivers known to be expressed in some or

all intestinal progenitor cells: a *GawB* P-element insertion in the *escargot* (*esg*) locus

(*esg-Gal4*), a *GawB* P-element insertion in the *Delta* (*Dl*) locus (*Dl-Gal4*), and a *Notch-*

responsive Gal4 reporter that contains multiple, tandem binding sites for the Grainyhead

and Suppressor of Hairless transcription factors (*gbe-Gal4*) [9, 10]. All three were

detected in the intestine (Fig 3E-G); *Dl-Gal4* and *gbe-Gal4* also displayed prominent

non-intestinal expression while *esg-Gal4* appeared more similar to the *I-KCKT* with

regard to intestinal specificity. For a more rigorous analysis, we also performed

comparative Western Blot analysis for GFP expression on three different protein

extracts generated from each strain: whole animal extract (total), extract from dissected

gastrointestinal tracts (intestine), and extract from all remaining non-intestinal tissue

(carcass) (Fig 3H). For both *I-KCKT* strains, GFP was detected in intestinal but not

carcass extract, and the amount in the intestine was roughly similar to the amount in the

total extract. In contrast, GFP was detected in both the intestine and carcass of the

three comparison strains, *esg-Gal4*, *Dl-Gal4*, and *gbe-Gal4*. We also tested the

conditional expression of *I-KCKT* by generating an *I-KCKT^{TS}* strain that harbored a

ubiquitously expressed, temperature sensitive Gal80 transgene (*tub-Gal80^{TS}*) and then

performing both analyses on resulting *I-KCKT^{TS}* flies that had been incubated at the

non-permissive (18°) and permissive (30°) temperatures (Fig 3H-J). This analysis

detected no GFP expression at the non-permissive temperature, and clear expression

at the permissive temperature. We note that I-KCKT-driven GFP signal can also be

detected in the progenitor cells of the malphigian tubules (arrowheads in Fig 3J), which

are *esg$^+$* cells that are related to midgut progenitor cells [20]. We also note that, like

esg-Gal4 but to a lesser degree, *I-KCKT-Gal4* loses its progenitor-specificity in the

intestinal tissue of aged animals (Fig S2). Collectively, these results indicated that,

unlike other commonly used progenitor drivers, I-KCKT-based expression could not be

detected outside the intestine.

ISC-KCKT-Gal4 specifically labels adult intestinal stem cells

To further expand on the utility of the I-KCKT system, we investigated whether I-

KCKT expression could be limited to either the ISC or EB subsets of progenitor cells. To

do so, we prepared strains in which *I-KCKT^{TS}* was combined with a Gal4-silencing

Gal80 transgene that contained either the EB-specific *gbe* synthetic enhancer (*gbe-*

Gal80) or a fragment from the *Delta* locus reported to be active in ISCs (*Dl^{GMR24H06}*-

Gal80) [21, 22]. We referred to these strains as *ISC-KCKT^{TS}* and *EB-KCKT^{TS}*,

respectively. Like *I-KCKT^{TS}*, both *ISC-KCKT^{TS}* and *EB-KCKT^{TS}* activity were intestine-

specific based on visual and/or Western blot analysis (Fig 3C, D, H). To evaluate the

cell type specificity of this activity, we drove GFP expression in both strains and

compared it to a dual reporter combination that we verified effectively distinguishes

ISCs and EBs (Fig S3). These dual reporters are the progenitor marker, *mira-*

His2A.mCherry.HA, and the EB-specific marker, *3Xgbe-smGFP::V5::nls*. For this

analysis, we scored mCherry$^+$, V5$^-$ cells as ISCs and mCherry$^+$, V5$^+$ cells as EBs. Most

ISCs were specifically labeled in the *ISC-KCKT^{TS}* strain: 84.1±12.3% (n=5) of ISCs were

labeled whereas 11.2±10.0% (n=5) of EBs were labeled across all five regions of the

intestine (Fig 2K-O, R, S). In contrast, EB-specific labeling in *EB-KCKT^{TS}* was less

effective: 57±19.8% (n=5) % of ISCs were labeled whereas only 30.7±20.1% (n=5) % of

EBs were labeled across all five regions of the intestine (Fig S4). Various possibilities

could explain this *EB-KCKT^{TS}* result, including that the GMR24H06 enhancer fragment

was not active in most ISCs and/or that Gal80 activity perdured in EB cells.

Nevertheless, this analysis indicated that *ISC-KCKT^{TS}* specifically labeled most ISCs.

RIP-seq of PABP using I-KCKT-Gal4 identifies progenitor expressed genes

We next investigated whether these I-KCKT strains could be used to streamline

current methods to molecularly profile progenitor cells. Current RNA-seq profiling

methods that rely on *esg-Gal4*, for example, require labor-intensive dissection of

hundreds of intestines followed by digestion of this tissue to release labeled progenitor

cells for fluorescent activated cell sorting (FACS) [2, 23-25]. Given the specificity of I-

KCKT drivers for intestinal progenitors, we reasoned that progenitor cell RNAs could

alternatively be profiled from intact tissue without any disruption. To test this possibility,

we expressed a FLAG-tagged poly-A Binding Protein (PABP) specifically in progenitors

with I-KCKT, prepared whole animal extract, subjected this extract to UV crosslinking to

covalently link protein and RNA complexes, immunoprecipitated PABP using anti-FLAG

beads, recovered associated RNAs, and then prepared and analyzed libraries for deep-

sequencing (Fig. 4A). PABP is a general mRNA-binding protein and we verified that a

tagged, inducible version that had previously been used to profile mRNAs in *Drosophila*

photoreceptor cells was expressed in progenitor cells when crossed to I-KCKT strains

(Fig S5) [26]. Libraries were then prepared in duplicate from RNA extracted from total

animal lysates of this strain ("input") as well as the PABP immunoprecipitates (PABP

IP). We first comparted the differential gene expression between PABP IP and input

using DESeq2 [27] with a false discovery rate adjusted p-value (FDR) of < 0.05 and a

log$_2$ fold change greater than one, and identified 1,661 genes enriched in the PABP IP

samples and 3,293 enriched in input (Fig. 4B). Representative examples of these

respective classes are the progenitor-enriched *esg* gene and the ovary-enriched *oskar*

gene (*osk*) (Fig. 4C-D). To evaluate the 1,661-enriched genes, we cross-referenced this

set to two published datasets. First, we compared our results to transcriptome profiles

obtained by Doupé, who used a DamID based method to identify progenitor-enriched or

EC-enriched genes [28]. Twenty percent of the 1,661 were present in Doupé's

progenitor-enriched genes, whereas only 4% were present in the EC-enriched set (Fig.

4E-F). Next, we re-analyzed published RNA-seq datasets from FACS-isolated

progenitors, ECs, and EEs [2], then identified progenitor enriched genes compared to

either ECs or EEs (FDR < 0.05 and $\log_2$ fold change > 1), and found that 31%-32% of

these enriched genes overlapped with genes enriched in the PABP IP (Fig. 4G-H).

Thus, genes most highly represented in the PABP IP include a substantial number of

genes previously shown to be enriched in progenitor cells.

To further evaluate the accuracy of the PABP IP dataset, we reasoned that it

should overall bear significant similarity to the RNA-seq profile of FAC sorted progenitor

cells. We therefore analyzed in parallel the PABP IP dataset and four previously

published RNA-seq datasets from FAC-sorted progenitor cells [2, 23-25]. Read counts

for all five sets were normalized using the Trimmed Mean of M-values method (TMM)

from edgeR to allow for accurate comparisons between samples [29]. We then looked

for genes common to all of these datasets that had a normalized expression value > 10,

and found 71% (4,966 genes out of 7,000 genes) of expressed genes in the PABP IP

overlapped with genes identified by all the other FACS-based datasets (Fig. 4I). To

assess the correlation of these six datasets, Spearman correlation matrices were

generated and visualized as correlograms. As a negative control for this analysis, we

also prepared a sixth dataset of head tissue data from FlyAtlas [16]. As expected, the

PABP IP correlated well with all FACS based progenitor datasets while, furthermore, all

five of these datasets showed a similar, low correlation with the non-intestinal dataset

(Fig. 4J). We then used the GeneOverlap R package to perform Fisher's exact test to

evaluate the statistical significance of the overlap between different datasets. In addition

to computing the statistical significance, this package also computes both an odds ratio

and a Jaccard index that indicates the association and similarity between two datasets,

respectively. The PABP IP scored highly in both these indices when compared to each

of the four FAC-sorted datasets and significantly higher than either the total input or

head tissue datasets, further indicating the similarity between the PABP IP and FACS-

based datasets (Fig. 4K). We performed one additional analysis with the FACS-based

datasets: first, we selected progenitor-enriched genes by comparing FACS-based data

to two sources of whole animal datasets, total input of the PABP experiment or

reanalyzed whole animal dataset from FlyAtlas (FDR < 0.05 and $\log_2$ fold change > 1)

and then assessed the similarity of this set to the gene set enriched in the PABP IP

(FDR < 0.05 and $\log_2$ fold change > 1 compared to input). As expected, PABP IP

enriched genes showed significant low p-values and high odds ratio and Jaccard index

scores strengthening the positive relationship of PABP and FACS-based datasets (Fig.

S6A). Altogether, this bioinformatic analysis supported a conclusion that the PABP IP

dataset represented the gene expression profiles of progenitor cells.

Having established that the PABP IP dataset contained progenitor genes, we

also sought to evaluate how much non-specific background was represented in this

dataset. Such noise might come from the post-lysis association of transgenic PABP with

mRNAs from non-progenitor cells. Hypothesizing that the most abundant transcripts in

the whole animal lysate might be most prone to associate with PABP post-lysis, we first

identified the top 10% of genes in the input based on the TMM normalized expression

values and found 1049 genes. Only 207 of these genes were also identified in the

PABP IP enriched gene set, suggesting that transcripts abundantly expressed in the

starting lysate were not non-specifically recovered in the IP (Fig. 4L). Supporting this

observation, we identified 854 genes in the input that were expected not to be present in

the PABP IP (Fig. S6B), and confirmed that only 4 of these transcripts were present in

the PABP IP enriched gene list (Fig. 4M). Finally, we again used FlyAtlas data to

identify genes enriched in fat body, head, ovary and testis relative to midgut and then

compared those tissue-enriched genes to the genes either enriched in or depleted from

the PABP IP. We found that odds ratio and Jaccard index scores were low for PABP-

enriched genes, indicating minimal overlap, and significantly higher for PABP-depleted

genes (Fig. 4N). As a positive control for this analysis, we performed the same

comparisons with genes enriched in the Fly Atlas midgut dataset as well as genes
enriched in FACS-based progenitor dataset from Buddika et al., 2020 [30]. As expected,
PABP IP enriched genes show highest association with genes enriched in FACS-
isolated progenitors, and significant overlap was also observed with the midgut enriched
gene list (Fig. 4N). Collectively, this analysis indicated that the PABP dataset contained
minimal background noise from other tissues, likely due to the effectiveness of the
crosslinking step.
279

As an alternative to PABP, we also tested whether this same approach could be
used with HA-tagged eIF4E, a translation initiation factor known to bind the mRNA cap
structure [31]. Unlike PABP, overexpression of eIF4E had no effect on progenitor cell
number over time (Fig S5). First, we performed differential expression analysis between
eIF4E IP and the total input from the PABP experiment using DESeq2. Using a false
discovery rate corrected p-value threshold of < 0.05, and a $\log_2$ fold change threshold of
> 1, we found only 751 genes to be significantly enriched in the eIF4E IP, but
approximately 38% of these genes were significantly enriched in the PABP IP as well
(Fig. S6C, F). Similar to the PABP IP data, the eIF4E IP data showed a significant
depletion of non-intestinal tissue specific transcripts, providing more proof for the
specificity of the approach (Fig. S6C). The reduced number of enriched genes in the
eIF4E IP may partly be due to the use of PABP total input as the control, that only a
subset of transcripts are physically associated with eIF4E, or differences in IP
efficiencies due to antibodies or protein isoforms. Despite these potential shortcomings,
the eIF4E gene list with normalized expression values > 10 clearly overlapped well with
similar lists from FACS-based datasets or PABP IP (Fig. S6D-E). These data
collectively suggest that the progenitor specificity of I-KCKT drivers will simplify the
labor-intensive protocols of performing any form of a biochemical or RIP-based
experiments in intestinal progenitors.
299

Discussion

Here we describe a method to purify the RNA cargoes associated with intestinal progenitor RBPs using whole animal lysate as the starting material. Because it is Gal4-based, this method will have wide applicability for profiling the general PABP- or eIF4E-bound transcriptome of wildtype and, when coupled with UAS-RNAi transgenes, mutant progenitor cells. In addition, it opens the possibility of molecularly characterizing the progenitor targets of RBPs, a growing number of which have been shown to play critical roles in progenitor cells (e.g. LIN28, FMRP, SPEN, TIS11 [32-35]). Finally, this method can be used in conjunction with additional current and future Gal4-based mRNA profiling methods, such as TU-tagging and ribosome profiling [36, 37], as well as methods that target other classes of RNAs, including microRNAs [38]. While the method is currently limited to progenitor cells, its applicability could be further expanded by generating kickout transgenes that drive expression in additional intestinal cell types such as EEs and ECs. While we focus on RNA profiling in this study, we also expect that I-KCKT in combination with mass-spectrometry to identify and characterize protein complexes and protein-protein interactions.

The I-KCKT method offers a number of advantages over current progenitor cell profiling methods, which usually involve the FAC sorting of progenitor cells from dissected and then mechanically disrupted intestines. The use of whole animals rather than dissected intestines as starting material greatly facilitates sample preparation, enabling the analysis of larger numbers of samples that could be used to test additional conditions and manipulations. In addition, because sample preparation is fast, this approach may effectively capture labile RNA profiles. Along these lines, the elimination of the FAC sorting step will also likely improve the accuracy of results, since FAC sorting is known to affect gene expression profiles likely due to the time and mechanical disruption associated with this step [39, 40]. In addition, this method could make molecular profiling of progenitors more accessible, since cell sorting runs can be costly and the equipment is not always available.

A current key limitation of the I-KCKT method is its purification of epitope-tagged

UAS-based transgenic protein rather than endogenous protein. This raises the concern

that the transgenic protein might be expressed at higher-than-endogenous levels, which

might cause inappropriate interactions with non-target mRNAs. Because the TARGET

system controlling transgene expression is temperature sensitive, this concern can be

addressed by testing multiple temperatures in the permissive range to identify

conditions supporting physiological protein expression. As an alternative approach to

address this concern, our current effort is focused on modifying endogenous RBP loci to

contain FRT-flanked epitope tags. I-KCKT-based expression of FLP should then lead to

the production of epitope-tagged endogenous protein in progenitor cells that would be

available for eCLIP-based analysis. Thus, we expect that I-KCKT-based methods will

allow unprecedented analysis of RNA-based mechanisms of progenitor cells.

Materials and Methods

Drosophila strains and husbandry:

All fly strains were cultured on standard Bloomington *Drosophila* stock center media (https://bdsc.indiana.edu/information/recipes/bloomfood.html). Flies were reared in 18°C, 25°C and 29°C incubators set for a 12hr light/dark schedule and 65% humidity. The genotypes of all strains used in this study are listed in Table S1. Transgenesis to create new strains was performed by Rainbow Transgenic Flies (Camarillo, CA) using plasmid DNA described below. Additional stocks were obtained from the Bloomington Drosophila Stock Center ({20XUAS-6XGFP}attP2, P{UAS-Stinger}2, P{tubP-Gal80[ts]}20, P{UAS-dPF}D), FlyORF ({UAS-eIF4E::3xHA}ZH-86Fb (F000955)), Steven Hou (P{Su(H)GBE-Gal4, P{GawB}Dl[05151-G]}), the Kyoto Drosophila Stock Center (P{GawB}NP5130), and Steve Stowers ({20XUAS-DSCP 6XGFP}attP2). For GAL80[ts]-dependent conditional expression experiments, flies were reared at 18°C, collected over 2 days, and then shifted to 30°C for up to 10 days before analysis. The following strains have been deposited at the Bloomington Drosophila Stock Center: (**i**) *I-KCKT[p65]* (BDSC #. Full genotype: *mira-KDRT>-dSTOP-KDRT>-Gal4::p65[1126]}attP40 ; {CG10116-KD::PEST[1107]}attP2*), (**ii**) *I-KCKT* (BDSC #. Full genotype: *{mira -KDRT>-dSTOP-KDRT>-Gal4[1438]}attP40, P{tubP-Gal80[ts]}20 ; {CG10116-KD::PEST[1107]}attP2*), (**iii**) *ISC-KCK[TS]* (BDSC #. Full genotype*: {gbe-Gal80}ZH-2A ; {mira -KDRT>-dSTOP-KDRT>-Gal4[1438]}attP40, P{tubP-Gal80[ts]}20 ; {CG10116-KD::PEST[1107]}attP2*).

Transgenes:

KD enhancer transgenes: PCR products containing enhancer fragments amplified from genomic DNA were cloned into the HindIII and AatII sites of pJFRC161 (20XUAS-IVS-KD::PEST, a gift from Gerald Rubin, Addgene plasmid # 32140) using HiFi DNA Assembly Mix (E2621L, New England Biolabs). Oligo pairs used to amplify enhancer fragments from *βTry* (3437/3438), *θTry* (3439/3440), *κTry* (3446/3447), *CG18404* (3443/3444 for the 2.6 kb fragment, 3445/3443 for the 0.5kb fragment), and *CG10116* (3441/3442) are shown in parentheses, and oligo sequences are reported in Table S2. Junctions of resulting plasmids were verified by sequencing prior to the preparation of

plasmid DNA for transgenesis. KD enhancer transgenes were inserted into the attP2

landing site.

KDRT Gal4 transgenes: pJFRC164 (pJFRC164-21XUAS-KDRT>-dSTOP-KDRT>-

myr::RFP, a gift from Gerald Rubin, Addgene plasmid # 32141) was used as a

backbone for KDRT-flanked stop cassette plasmids. Four-way HiFi DNA Assembly

reactions were performed with (i) 7.2kb HindIII/XbaI-digested pJFRC164 plasmid

backbone, (ii) 1.4kb AatII/XhoI-digested pJFRC164 KDRT>-dSTOP-KDRT fragment, (iii)

PCR-amplified Gal4-containing fragments, and (iv) PCR amplified enhancer-containing

fragments. Gal4::p65 was amplified from pBPGAL4.2::p65Uw (a gift from Gerald Rubin,

Addgene plasmid # 26229) with oligo pair 3433/3434, while Gal4 sequence was

amplified from pBPGAL4.1Uw (a gift from Gerald Rubin, Addgene plasmid # 26226)

with oligo pair 4401/4402. Enhancer fragments include: (i) 2.6 kb *tubulin* fragment

amplified with oligo pair 3626/3627, (ii) 2.6 kb 5' and 1.6 kb 3' *miranda* fragments

amplified with oligo pairs 3489/3490 and 3491/3492, respectively, and (iii) 3.4 kb *delta*

fragment amplified with oligo pair 3431/3432. Plasmid junctions and open reading

frames were verified by sequencing prior to the preparation of DNA for transgenesis.

Oligo sequences are reported in Table S2, and complete plasmid sequences are

available upon request. KDRT transgenes were inserted into the attP40 landing site.

Gal80 transgenes: For DI^{R24H06}-Gal80, the previously defined R24H06 enhancer

fragment of the *delta* locus was PCR amplified from genomic DNA and subcloned

upstream of the Gal80 open reading frame in a pATTB-containing transformation

plasmid. 3xgbe-Gal80 was generated in a similar manner except using oligos encoding

three tandem copies of *gbe* sequence. Both of these Gal80 transgenes were inserted

into the ZH-2A landing site.

Dissections and immunostaining:

Adult female flies were dissected in ice cold 1xPBS and fixed in 4% paraformaldehyde

(Electron Microscopy Sciences, Cat. No. 15714) in PBS for 45 min. Resulting dissected

tissue was washed in 1xPBT (1xPBS, 0.1% Triton X-100) and then blocked (1xPBT,

0.5% Bovine Serum Albumin) for at least 45 min. For Delta staining, an alternative

sample preparation scheme, adapted from [41], was followed. Briefly, intestines were

dissected in ice-cold Grace's Insect Medium (Lonza Bioscience, 04-457F) and fixed in a

1:1 (V/V) mixture of heptane (Sigma, H2198) and 4% paraformaldehyde (Electron

Microscopy Sciences, Cat. No. 15714) in water for 15 min. The bottom aqueous

paraformaldehyde layer was removed, 500µl of ice-cold methanol was added, and

shook vigorously for 30 seconds. The methanol/heptane mixture was removed and

incubated with 1ml ice-cold methanol for 5 min. Next, samples were gradually

rehydrated with a series of 0.3% PBT (1xPBS, 0.3% Triton X-100) : Methanol (3:7, 1:1,

7:3) washes, washed with 0.3% PBT alone for another 5 min and blocked (0.3% PBT,

0.5% Bovine Serum Albumin) for at least 45 min. Samples were incubated at 4°C

overnight with primary antibodies, including rabbit anti-RFP (Clontech, 1:1000), mouse

anti-V5 (MCA1360GA, Bio-Rad, 1:250), chicken anti-LacZ (ab9361, Abcam, 1:2000),

mouse anti-FLAG (F3165, Sigma, 1:1000), mouse anti-Delta (C594.9B, Developmental

Studies Hybridoma Bank, 1:500), and rabbit anti-HA (3724S, Cell Signaling Technology,

1:1000). The following day, samples were washed in 1xPBT and incubated for 2-3

hours with secondary antibodies, including AlexaFluor-488 and -568-conjugated goat

anti-rabbit, -mouse, -rat and -chicken antibodies (Life Technologies, 1:1000). Finally,

samples were washed in 1xPBT, including one wash with DAPI (1:1000), and mounted

in Vectashield mounting medium (Vector Laboratories) or, for Delta staining, in ProLong

Diamond mounting medium (Invitrogen, P36970).

Microscopy and image processing:

Images of whole flies were collected on a Zeiss Axio Zoom microscope and images of

dissected intestines were collected on a Leica SP8 Scanning Confocal microscope.

Samples to be compared were collected under identical settings on the same day,

image files were adjusted simultaneously using Adobe Photoshop CC, and figures were

assembled using Adobe Illustrator CC.

Western blot analysis:

Female *Drosophila* flies were used for protein isolation. Extract was prepared from

either whole animals or separately from the intestines and remaining carcass tissue of

dissected animals. Tissues were lysed in protein lysis buffer (25mM Tris-HCl pH 7.5,

150mM KCl, 5mM $MgCl_2$, 1% NP-40, 0.5mM DTT, 1xprotease inhibitor cocktail), protein

extracts were resolved on a 4-20% gradient polyacrylamide gel (Bio-Rad, Cat. No. 456-

1093), transferred to Immobilon®-P membrane (Millipore, Cat. No. IPVH00010) and

probed with rabbit anti-GFP (ab290, Abcam, 1:10,000) or mouse anti-a-tubulin (12G10,

Developmental Studies Hybridoma Bank, 1:1000) antibodies. Subsequently, blots were

washed extensively with 1xTBST (1xTBS, 0.1% Tween-20) and incubated with anti-

rabbit or -mouse conjugated HRP secondary antibodies. After extensive secondary

washes with 1xTBST, blots were treated with ECL-detection reagents (Thermo

Scientific, Cat. No. 1859701 and 1859698) and finally exposed to chemiluminescence

films (GE Healthcare, Cat. No. 28906839).

RNA immunoprecipitation and Library construction:

Whole flies were collected and immediately snap frozen in liquid nitrogen and

stored at -80°C until sample preparation. Two hundred flies (100 male and 100 female)

of each sample were ground into a fine powder using a pre-cooled mortar and pestle on

dry ice. The powder was irradiated three times at 120 mJ per cm2 in a UV cross-linker,

with mixing between each irradiation cycle to maximize surface exposure. The fly

powder was transferred to a 2ml tube containing 1ml lysis buffer using a cold spatula,

and the remaining fly powder was washed off the spatula with an additional 1ml lysis

buffer. Tubes were incubated on ice for 15-30 min, with mixing every 5 min. Epitope-

tagged proteins were immunoprecipitated with either anti-Flag (Sigma) or anti-HA

(Pierce) magnetic beads and following the manufacturer's instructions. RNA was eluted

from the beads with Proteinase K and TRIzol LS Reagent was used to isolate

immunoprecipitated RNA. The Ovation SoLo RNA-Seq System was used to make

PABP and EIF4E libraries. eCLIP libraries were generated by following the protocol in

Nostrand et al., 2016 [42], with a minor modification that 8U of RNAse I was used per

sample.

RNA-seq data analysis

RNA-seq read files were processed and aligned to the Berkeley *Drosophila*

Genome Project (BDGP) assembly release 6.28 (Ensembl release 99) reference

genome using a python based in house pipeline (https://github.com/jkkbuddika/RNA-

Seq-Data-Analyzer). Briefly, the quality of raw sequencing files was assessed using

FastQC [43] version 0.11.9, low quality reads were removed using Cutadapt [44]

version 2.9, reads aligning to rRNAs were removed using TagDust2 [45] version 2.2,

remaining reads were mapped to the Berkeley *Drosophila* Genome Project (BDGP)

assembly release 6.28 (Ensembl release 99) reference genome using STAR [46]

genome aligner version 2.7.3a and deduplicated using SAMtools [47] version 1.10.

Subsequently, the aligned reads were counted to the nearest overlapping feature suing

the Subread [48] version 2.0.0 function *featureCounts*. Finally, bigWig files representing

RNA-seq coverage were generated using deepTools [49] version 3.4.2 with the settings

--normalizeUsing CPM --binSize 1. The TMM-normalized expression values were

computed using the Bioconductor package edgeR [29] version 3.28.1. Differential gene

expression analysis was performed with the Bioconductor package DESeq2 [27] version

1.26.0.

Enhanced-CLIP (eCLIP) data analysis

The eCLIP read file processing, alignment and processing was done using an

easy to use python based in house analysis pipeline, eCLIP data analyzer

(https://github.com/jkkbuddika/eCLIP-Data-Analyzer). While the eCLIP data analyzer

pipeline permits analysis in both paired and single-end modes, we performed all our

analyzes using the single-end mode, which uses only R2 read for analysis. The eCLIP

data analyzer pipeline performs the following steps: (1) read quality assessment using

FastQC [43], (2) trimming of universal eCLIP adaptors using Cutadapt [44], (3) addition

of UMI sequence (5'-NNNNNNNNNN in the R2 read) using UMI-tools [50] to read name

to facilitate PCR duplicate removal, (4) removal of reads aligning to rRNAs using

Tagdust2 [45], (5) alignment of remaining reads to the provided genome (i.e., BDGP

assembly release 6.28, Ensembl release 100) using STAR [46], (6) coordinating sort

and index alignment outputs using SAMtools [47], (7) removal of PCR duplicates using

UMI-tools [50], (8) generation of bigwig files representing RNA-seq coverage tracks

using deepTools [49], (9) quantification of nearest overlapping features using Subread

[48] function *featureCounts*, and (10) peak calling using PureCLIP [51]. All programs,

versions and dependencies required to execute the eCLIP data analyzer are described

in the user guide and can be installed using miniconda. Input normalized peak calling

output files were processed as described in [52]. Differential gene expression analysis

was performed with the Bioconductor package DESeq2 [27] version 1.26.0.

Statistical analysis:

All statistical analyses were performed using Prism (GraphPad, Version 7.0). Datasets

were tested for normality using D'Agostino-Pearson test. Comparisons of two datasets,

were performed with an Unpaired t-test. Unless otherwise noted, significance is

indicated as follows: n.s., not significant; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p <$

0.0001.

Acknowledgements

We thank Steven Hou, Gerald Rubin, Steve Stowers, the Kyoto Drosophila Stock

Center, the Bloomington Drosophila Stock Center (supported by grant

NIHP4OOD018537), the Drosophila Genome Resource Center (supported by grant

NIH2P40OD010949), and the Developmental Studies Hybridoma Bank for reagents; the

Light Microscopy Imaging Center (supported by grant NIH1S10OD024988-01) for

access to the SP8 confocal; the Center for Genomics and Bioinformatics for sequencing

results; and the National Institute of General Medical Sciences (Award R01GM124220)

for financial support. The authors declare no competing financial interests.

References:

1. Li, H. and H. Jasper, *Gastrointestinal stem cells in health and disease: from flies to humans.* Dis Model Mech, 2016. **9**(5): p. 487-99.
2. Dutta, D., et al., *Regional Cell-Specific Transcriptome Mapping Reveals Regulatory Complexity in the Adult Drosophila Midgut.* Cell Rep, 2015. **12**(2): p. 346-58.
3. Hung, R.J., et al., *A cell atlas of the adult Drosophila midgut.* Proc Natl Acad Sci U S A, 2020. **117**(3): p. 1514-1523.
4. Baser, A., et al., *Onset of differentiation is post-transcriptionally controlled in adult neural stem cells.* Nature, 2019. **566**(7742): p. 100-104.
5. Sanz, E., et al., *Cell-type-specific isolation of ribosome-associated mRNA from complex tissues.* Proc Natl Acad Sci U S A, 2009. **106**(33): p. 13939-44.
6. Tahmasebi, S., M. Amiri, and N. Sonenberg, *Translational Control in Stem Cells.* Front Genet, 2018. **9**: p. 709.
7. Buddika, K., et al., *Canonical nucleators are dispensable for stress granule assembly in Drosophila intestinal progenitors.* J Cell Sci, 2020. **133**(10).
8. Jiang, H. and B.A. Edgar, *EGFR signaling regulates the proliferation of Drosophila adult midgut progenitors.* Development, 2009. **136**(3): p. 483-93.
9. Micchelli, C.A. and N. Perrimon, *Evidence that stem cells reside in the adult Drosophila midgut epithelium.* Nature, 2006. **439**(7075): p. 475-9.
10. Zeng, X., C. Chauhan, and S.X. Hou, *Characterization of midgut stem cell- and enteroblast-specific Gal4 lines in drosophila.* Genesis, 2010. **48**(10): p. 607-11.
11. Biteau, B., et al., *Lifespan extension by preserving proliferative homeostasis in Drosophila.* PLoS Genet, 2010. **6**(10): p. e1001159.
12. Nern, A., et al., *Multiple new site-specific recombinases for use in manipulating animal genomes.* Proc Natl Acad Sci U S A, 2011. **108**(34): p. 14198-203.
13. Bardin, A.J., et al., *Transcriptional control of stem cell maintenance in the Drosophila intestine.* Development, 2010. **137**(5): p. 705-14.
14. Phillips, M.D. and G.H. Thomas, *Brush border spectrin is required for early endosome recycling in Drosophila.* J Cell Sci, 2006. **119**(Pt 7): p. 1361-70.
15. Voght, S.P., et al., *Drosophila NPC1b promotes an early step in sterol absorption from the midgut epithelium.* Cell Metab, 2007. **5**(3): p. 195-205.
16. Leader, D.P., et al., *FlyAtlas 2: a new version of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.* Nucleic Acids Res, 2018. **46**(D1): p. D809-D815.
17. Shearin, H.K., et al., *Hexameric GFP and mCherry reporters for the Drosophila GAL4, Q, and LexA transcription systems.* Genetics, 2014. **196**(4): p. 951-60.
18. Barolo, S., L.A. Carver, and J.W. Posakony, *GFP and beta-galactosidase transformation vectors for promoter/enhancer analysis in Drosophila.* Biotechniques, 2000. **29**(4): p. 726, 728, 730, 732.
19. McGuire, S.E., Z. Mao, and R.L. Davis, *Spatiotemporal gene expression targeting with the TARGET and gene-switch systems in Drosophila.* Sci STKE, 2004. **2004**(220): p. pl6.

565   20.   Singh, S.R., W. Liu, and S.X. Hou, *The adult Drosophila malpighian tubules are*
566   *maintained by multipotent stem cells.* Cell Stem Cell, 2007. **1**(2): p. 191-203.

567   21.   Guo, Z., I. Driver, and B. Ohlstein, *Injury-induced BMP signaling negatively*
568   *regulates Drosophila midgut homeostasis.* J Cell Biol, 2013. **201**(6): p. 945-61.

569   22.   Furriols, M. and S. Bray, *A model Notch response element detects Suppressor of*
570   *Hairless-dependent molecular switch.* Curr Biol, 2001. **11**(1): p. 60-4.

571   23.   Fast, D., et al., *Vibrio cholerae-Symbiont Interactions Inhibit Intestinal Repair in*
572   *Drosophila.* Cell Rep, 2020. **30**(4): p. 1088-1100 e5.

573   24.   Korzelius, J., et al., *The WT1-like transcription factor Klumpfuss maintains*
574   *lineage commitment of enterocyte progenitors in the Drosophila intestine.* Nat
575   Commun, 2019. **10**(1): p. 4123.

576   25.   Li, Q., et al., *Ingestion of Food Particles Regulates the Mechanosensing*
577   *Misshapen-Yorkie Pathway in Drosophila Intestinal Growth.* Dev Cell, 2018.
578   **45**(4): p. 433-449 e6.

579   26.   Yang, Z., H.J. Edenberg, and R.L. Davis, *Isolation of mRNA from specific tissues*
580   *of Drosophila by mRNA tagging.* Nucleic Acids Res, 2005. **33**(17): p. e148.

581   27.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
582   *dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

583   28.   Doupe, D.P., et al., *Drosophila intestinal stem and progenitor cells are major*
584   *sources and regulators of homeostatic niche signals.* Proc Natl Acad Sci U S A,
585   2018. **115**(48): p. 12218-12223.

586   29.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor*
587   *package for differential expression analysis of digital gene expression data.*
588   Bioinformatics, 2010. **26**(1): p. 139-40.

589   30.   Buddika, K., et al., *Intestinal progenitor P-bodies maintain stem cell identity by*
590   *suppressing pro-differentiation factors.* BioRxiV, 2020.

591   31.   Rissland, O.S., *The organization and regulation of mRNA-protein complexes.*
592   Wiley Interdiscip Rev RNA, 2017. **8**(1).

593   32.   Andriatsilavo, M., et al., *Spen limits intestinal stem cell self-renewal.* PLoS Genet,
594   2018. **14**(11): p. e1007773.

595   33.   Chen, C.H., A. Luhur, and N. Sokol, *Lin-28 promotes symmetric stem cell*
596   *division and drives adaptive growth in the adult Drosophila intestine.*
597   Development, 2015. **142**(20): p. 3478-87.

598   34.   Luhur, A., et al., *Opposing Post-transcriptional Control of InR by FMRP and LIN-*
599   *28 Adjusts Stem Cell-Based Tissue Growth.* Cell Rep, 2017. **21**(10): p. 2671-
600   2677.

601   35.   McClelland, L., H. Jasper, and B. Biteau, *Tis11 mediated mRNA decay promotes*
602   *the reacquisition of Drosophila intestinal stem cell quiescence.* Dev Biol, 2017.
603   **426**(1): p. 8-16.

604   36.   Chen, X. and D. Dickman, *Development of a tissue-specific ribosome profiling*
605   *approach in Drosophila enables genome-wide evaluation of translational*
606   *adaptations.* PLoS Genet, 2017. **13**(12): p. e1007117.

607   37.   Hida, N., et al., *EC-tagging allows cell type-specific RNA analysis.* Nucleic Acids
608   Res, 2017. **45**(15): p. e138.

609   38.   Luhur, A., G. Chawla, and N.S. Sokol, *MicroRNAs as components of systemic*
610   *signaling pathways in Drosophila melanogaster.* Curr Top Dev Biol, 2013. **105**: p.
611   97-123.
612   39.   Andra, I., et al., *An Evaluation of T-Cell Functionality After Flow Cytometry*
613   *Sorting Revealed p38 MAPK Activation.* Cytometry A, 2020. **97**(2): p. 171-183.
614   40.   Richardson, G.M., J. Lannigan, and I.G. Macara, *Does FACS perturb gene*
615   *expression?* Cytometry A, 2015. **87**(2): p. 166-75.
616   41.   Lin, G., N. Xu, and R. Xi, *Paracrine Wingless signalling controls self-renewal of*
617   *Drosophila intestinal stem cells.* Nature, 2008. **455**(7216): p. 1119-23.
618   42.   Van Nostrand, E.L., et al., *Robust transcriptome-wide discovery of RNA-binding*
619   *protein binding sites with enhanced CLIP (eCLIP).* Nat Methods, 2016. **13**(6): p.
620   508-14.
621   43.   Andrews, S., *FastQC: A quality control tool for high throughput sequence data.*
622   http://wwwbioinformaticsbabrahamacuk/projects/fastqc, 2010.
623   44.   Martin, M., *Cutadapt removes adapter sequences from high-throughput*
624   *sequencing reads.* 2011, 2011. **17**(1): p. 3.
625   45.   Lassmann, T., *TagDust2: a generic method to extract reads from sequencing*
626   *data.* BMC Bioinformatics, 2015. **16**: p. 24.
627   46.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013.
628   **29**(1): p. 15-21.
629   47.   Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics,
630   2009. **25**(16): p. 2078-9.
631   48.   Liao, Y., G.K. Smyth, and W. Shi, *The R package Rsubread is easier, faster,*
632   *cheaper and better for alignment and quantification of RNA sequencing reads.*
633   Nucleic Acids Res, 2019. **47**(8): p. e47.
634   49.   Ramirez, F., et al., *deepTools2: a next generation web server for deep-*
635   *sequencing data analysis.* Nucleic Acids Res, 2016. **44**(W1): p. W160-5.
636   50.   Smith, T., A. Heger, and I. Sudbery, *UMI-tools: modeling sequencing errors in*
637   *Unique Molecular Identifiers to improve quantification accuracy.* Genome Res,
638   2017. **27**(3): p. 491-499.
639   51.   Krakau, S., H. Richard, and A. Marsico, *PureCLIP: capturing target-specific*
640   *protein-RNA interaction footprints from single-nucleotide CLIP-seq data.* Genome
641   Biol, 2017. **18**(1): p. 240.
642   52.   Busch, A., et al., *iCLIP data analysis: A complete pipeline from sequencing reads*
643   *to RBP binding sites.* Methods, 2020. **178**: p. 49-62.
644
645

Figure Legends:

Figure 1: KD driver analysis identifies a pan-intestinal enhancer in adults. (A)
Schematic of the I-KCKT system, which involves three transgenes: (i) a KD-expressing
transgene under the control of an intestinal enhancer, (ii) a transgene in which a KDRT-
flanked stop-cassette separates Gal4 from a progenitor enhancer, and (iii) a UAS
responder, in this case, controlling GFP expression. (B-M) KD recombinase-mediated
labeling of (B-G) larval and (H-M) adult intestinal cells. KD lines contain 3.4kb *βTry*
(B,H), 1.6kb *θTry* (C, I), 0.6kb *κTry* (D, J), 2.6kb *CG18404* (E, K), 0.5kb *CG18404* (F, L),
or 1kb *CG10116* (G, M) enhancer sequences. KD expression pattern was detected
based on KDRT-mediated activation of a *tubulin*-based Gal4::p65 driver in combination
with a *UAS-6XGFP* responder. Full genotypes are listed in Table S1.

Figure 2: I-KCKT[P65] and I-KCKT labels most intestinal progenitor cells and ISC-KCKT[TS] labels most ISCs. (A-O) Representative micrographs of the five regions (R1-R5) from intestines stained for I-KCKT[P65]>UAS-stinger-GFP (A-E), I-KCKT>UAS-stinger-GFP (F-J), or ISC-KCKT>UAS-stinger-GFP (K-O) in green and counterstained for intestinal progenitor marker *mira-His2A.mCherry.HA* in red and the DAPI DNA marker in blue. (P-Q) Histograms showing percentage of *mira-His2A.mCherry.HA-*

labeled intestinal progenitor (IP) cells that are also labeled with I-KCKT^{P65}>UAS-stinger-

GFP (F) or I-KCKT>UAS-stinger-GFP (L) per intestinal region (R1-R5). (R-S)

Histograms showing the percentage of ISCs (R) and EBs (S) labeled with ISC-KCKT^{TS}.

Full genotypes are listed in Table S1.

Figure 3: I-KCKT drivers are intestine-specific. (A-G) Pictures of whole adult females

showing 6XGFP expression patterns driven by I-KCKT^{P65} (A), I-KCKT^{TS} (B), EB-KCKT^{TS}

(C), ISC-KCKT^{TS} (D), esg$^{P\{GawB\}NP5130}$-Gal4 (E), Dl$^{05151-G}$-Gal4 (F), or Su(H)GBE-Gal4 (G)

Insets are enlargements of abdominal regions displaying intestinal GFP expression. (H)

Western blot analysis of tissue extracts from whole animals (total), dissected intestines,

or dissected carcasses showing 6XGFP levels driven by I-KCKT^{P65}, I-KCKT, I-KCKT^{TS}

at 18°C, I-KCKT^{TS} at 30°C, ISC-KCKT^{TS}, esg$^{P\{GawB\}NP5130}$-Gal4, Dl$^{05151-G}$-Gal4, or

Su(H)GBE-Gal4. (I, J) 6XGFP expression in dissected intestines of I-KCKT^{TS}>6GFP

adults kept at 18°C (I) or 30°C (J) and counterstained for the DAPI DNA marker in blue.

Yellow arrowheads indicate GFP expression at the base of the malphigian tubules. Full

genotypes are listed in Table S1.

Figure 4: RIP-seq of PABP using I-KCKT identifies progenitor expressed genes.
(A) Schematic representation of the conventional FACS (left) and I-KCKT-based
intestinal progenitor RNA-sequencing (right). (B) Volcano plot of differentially expressed
genes from comparison of PABP IP to total input. Each dot represents a single gene.
Yellow indicates a false discovery rate adjusted p-value (FDR) < 0.05 and a Log_2 fold
change > 1 or < -1. Green indicates Log_2 fold change > 1 or < -1 and an FDR >= 0.05.
Red indicates an FDR < 0.05 and log_2 fold change > 1 or < -1. A selected set of
significantly changed genes have been labeled. (C-D) Genome browser tracks of
normalized total input and PABP IP at the (C) *esg* or (D) *osk* locus of the *Drosophila*
genome. Note that the two replicates of the total input (green and yellow) and the two
replicates of the PABP IP (red and blue) have been overlayed for simplicity. (E-H) Venn
diagram of genes significantly enriched in PABP-RIP compared to total input (FDR <
0.05 and log_2 fold change > 1) and (E) in intestinal progenitors based on progenitor
DamID, or (F) in ECs based on EC DamID based on [28], or (G) in intestinal progenitors
compared to ECs (FDR < 0.05 and log_2 fold change > 1) or (H) in intestinal progenitors
compared to EEs (FDR < 0.05 and log_2 fold change > 1) based on reanalyzed data from
[2]. (I) Upset plot showing the overlap of moderately expressed genes identified by
PABP RIP-seq versus previously published FACS based methods. Only a selected set
of meaningful overlaps are shown for simplicity. Numbers above each bar shows the
size of each intersection. (J) Correlogram of Spearman's *rho* values for pairwise
comparisons of progenitor expressed genes based on PABP RIP-seq and FACS-based
progenitor RNA-seq as well as *Drosophila* female head expressed genes based on
RNA-seq of dissected heads. (K) Heatmaps of odds ratio (in green) and Jaccard index
(in orange) values for pairwise comparisons of PABP RIP-seq, RNA-seq of total input or
RNA-seq of dissected female heads against FACS-based progenitor RNA-seq.
Numbers on each colored box show the p-value for each overlap based on Fisher's
exact test. For I-K, PABP IP indicates data from I-KCKT-based PABP RIP-seq. Buddika,
Dutta, Fast and Korzelius indicate data from RNA-seq of FACS isolated progenitors
from [2, 23, 24, 30], respectively. For J-K, Head indicates reanalyzed female head RNA-
seq data from Leader *et al.,* 2017. (L-M) Venn diagram of genes significantly enriched in
PABP-RIP compared to total input (FDR < 0.05 and log_2 fold change > 1) and (L) top

10% genes with highest TMM normalized expression values or (M) genes uniquely

present in input, but not in any of the FACS-based RNA-seq gene lists. (N) Heatmaps of

odds ratio (in blue) and Jaccard index (in green) values for pairwise comparisons of

PABP IP enriched (FDR < 0.05 and $\log_2$ fold change > 1 vs input) or PABP IP depleted

(FDR < 0.05 and $\log_2$ fold change < -1 vs input) genes against fat body, head, ovary or

testis enriched genes compared to midgut (FDR < 0.05 and $\log_2$ fold change > 1) or

genes enriched in midgut or FACS-based progenitor dataset from Buddika *et al.,* 2020

[30] compared to whole animal input (FDR < 0.05 and $\log_2$ fold change > 1). Numbers

on each colored box show the p-value for each overlap based on Fisher's exact test.

Note that fat body, head, ovary, testis, midgut and whole animal datasets represent

reanalyzed data from [16].

Supplementary Figure Legends:

Figure S1: I-KCKT and esg-LacZ label the same cells. (A-E) Representative

micrographs of the five regions (R1-R5) from the intestines of 5-day old females stained

for *I-KCKT^{TS}>UAS-stinger-GFP* in green, *esg-LacZ* in red, and the DAPI DNA marker in

blue. (F) Histogram of the percentage of *esg-LacZ*-positive cells that also expressed *I-*

KCKT^{TS}>UAS-stinger-GFP in each region.

Figure S2: Comparison of I-KCKT-Gal4 and esg expression in aged flies.

Representative micrographs of the five regions (R1-R5) from the intestines from young

(5 day) and older (30-day) females stained for *esg>UAS-stinger-GFP* (top) or *I-*

KCKT^P65^>UAS-stinger-GFP (bottom) in green and counterstained for the DAPI DNA

marker in blue.

Figure S3: Verification that dual reporters *mira-His2A.mCherry.HA* and *3Xgbe-

smGFP::V5::nls **effectively distinguish ISCs and EBs.** (A-E) Representative

micrographs of the five regions (R1-R5) from the intestines of 5-day old females stained

for ISCs (anti-Delta, green cytoplasmic staining), EEs (anti-Prospero, green nuclear

staining), *mira-His2A.mCherry.HA* (red), *3Xgbe-smGFP::V5::nls (white),* and the DAPI

DNA marker (blue). (F) Quantification of the percent of ISCs and EBs labeled by the

dual reporters *mira-His2A.mCherry.HA* and *3Xgbe-smGFP::V5::nls* in each of the five

intestinal regions. ISCs were defined as the number of Delta+ cells per field of view.

EBs were defined as the number of Delta-, mCherry+ cells per field of view. For

quantification, field of view images were taken and quantified from defined portions of

each intestinal region in 6-7 intestines. We note that we detected a few (1-5) cells that

were both Delta+ and *3Xgbe-smGFP::V5::nls*+ in regions of most intestines. These

were not scored as either ISCs or EBs, and could represent cells transitioning between

these two cell fates.

Figure S4: ISC-KCKT^TS labels many ISCs. (A-E) Representative micrographs of the

five regions (R1-R5) from the intestines of 5-day old females stained for *I-*

KCKT^TS>UAS-stinger-GFP in green, *mira-His2A.mCherry.HA* in red, *3Xgbe-*

smGFP::V5::nls in white, and the DAPI DNA marker in blue. ISCs were scored as cells

positive for *mira-His2A.mCherry.HA* and negative for *3Xgbe-smGFP::V5::nls*, while EBs

were scored as cells positive for both markers. (F) Histogram of the percentage of EBs that

expressed *I-KCKT^TS>UAS-stinger-GFP* in each region. (G) Histogram of the percentage

of ISCs that expressed *I-KCKT^TS>UAS-stinger-GFP* in each region.

775
776

Figure S5: I-KCKT^{TS}-dependent expression of tagged RBPs in intestinal progenitor cells. (A-J) Micrographs of intestines showing I-KCKT^{TS}-dependent expression of UAS-GFP::FLAG (A, F), UAS-EIF4E::HA (D, I), UAS-PABP::FLAG (E, J), UAS-LIN28::FLAG::HA (B, G), and UAS-FMR1::GFP::FLAG (C, H) after two (A-E) and ten (F-J) days at the permissive temperature. Expressed proteins were detected with either anti-FLAG or -HA antibodies (shown in red) and intestines were counterstained with the DAPI DNA marker in blue. Full genotypes are listed in Table S1. (K, L) Histograms of the percentage of progenitor cells after two and ten days of I-KCKT^{TS}-dependent expression, based on quantification of the numbers of HA/FLAG$^+$ and DAPI$^+$ cells.

Figure S6: I-KCKT based RIPs identifies genes expressed in intestinal

progenitors. (A) Heatmaps of odds ratio (in blue) and Jaccard index (in green) values

for pairwise comparisons of PABP IP enriched or PABP IP depleted genes against input

normalized FACS-based progenitor RNA-seq. Two sources of input data have been

used for FACS data normalization: Total input of the PABP experiment, or female whole

animal dataset from Leader *et al.,* 2017. Numbers on each colored box show the p-

value for each overlap based on Fisher's exact test. N.S.: Not significant. (B) Upset plot

showing the overlap of moderately expressed genes identified by FACS based methods

and total PABP IP input. Only a select set of meaningful overlaps are shown for

simplicity. The number above each bar shows the precise size of each intersection.

Note that 854 genes were uniquely present in the input, but not in any of the FACS

based datasets. (C) Volcano plot of differentially expressed genes from comparison of

eIF4E IP to total input from the PABP experiment. Each dot represents a single gene.

Yellow indicates a false discovery rate adjusted p-value (FDR) < 0.05 and a Log_2 fold

change > 1 or < -1. Green indicates Log_2 fold change > 1 or < -1 and an FDR >= 0.05.

Red indicates an FDR < 0.05 and log_2 fold change > 1 or < -1. A selected set of

significantly changed genes have been labeled. (D) Upset plot showing the overlap of

moderately expressed genes identified by eIF4E RIP-seq or FACS based methods.

Only a selected set of overlaps are shown for simplicity. Number above each bar shows

the precise size of each intersection. (E-F) Venn diagram of (E) moderately expressed

genes identified by eIF4E or PABP RIP-seq methods or (F) PABP or eIF4E IP enriched

genes compared to total input (FDR < 0.05 and log_2 fold change > 1). Only moderately

expressed genes were considered for the analyses shown in B, D and E. A gene is

defined as moderately expressed, if the TMM normalized expression value is greater

than 10 in all replicates. eIF4E IP indicates data from I-KCKT-based eIF4E RIP-seq.

Buddika, Dutta, Fast and Korzelius indicate data from RNA-seq of FACS isolated

progenitors from [2, 23, 24, 30], respectively.

Supplementary Tables

 ### **Table S1: Fly strains used in this study**

Figure	Panels	Genotypes of flies used in each panel
Fig. 1	B, H	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {βTry-KD::PEST^{1105}}attP2
	C, I	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {θTry-KD::PEST^{1106}}attP2 / +
	D, J	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {κTry-KD::PEST^{1110}}attP2 / +
	E, K	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {CG18404-2.6kb-KD::PEST^{1108}}attP2 / +
	F, L	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {CG18404-0.5kb-KD::PEST^{1109}}attP2 / +
	G, M	{tub-KDRT>-dSTOP-KDRT>-Gal4::p65^{1156}}attP40 / + ; {20XUAS-6XGFP}attP2 ; {CG10116-KD::PEST^{1107}}attP2 / +
Fig 2	A-E,P	{3Xgbe-smGFP::V5::nls}ZH2A, M{mira-His2A.mCherry.HA}Su(hw)attP18 / + ; {mira-KDRT>-dSTOP-KDRT>-Gal4::p65^{1126}}attP40 / P{UAS-Stinger}2 ; {CG10116-KD::PEST^{1107}}attP2 / +
	F-J,Q	{3Xgbe-smGFP::V5::nls}ZH2A, M{mira-His2A.mCherry.HA}Su(hw)attP18 / + ; {mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / P{UAS-Stinger}2 ; {CG10116-KD::PEST^{1107}}attP2 / +
	K-O, R,S	{3Xgbe-smGFP::V5::nls}ZH2A, M{mira-His2A.mCherry.HA} Su(hw)attP18 / {3Xgbe-Gal80}ZH-2A ; {mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; P{UAS-Stinger}2; {CG10116-KD::PEST^{1107}}attP2 / +
Fig. 3	A, H	{mira-KDRT>-dSTOP-KDRT>-Gal4::p65^{1126}}attP40 / + ; {20XUAS-DSCP 6XGFP}attP2 / {CG10116-KD::PEST^{1107}}attP2
	B, H, I, J	{mira -KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {20XUAS-DSCP-6XGFP} attP2 / {CG10116-KD::PEST^{1107}}attP2
	C, H	{Dl^{R24H06}-Gal80}ZH-2A / + ; {mira -KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {20XUAS-DSCP-6XGFP} attP2 / {CG10116-KD::PEST^{1107}}attP2
	D, H	{3xgbe-Gal80}ZH-2A / + ; {mira -KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {20XUAS-DSCP-6XGFP} attP2 / {CG10116-KD::PEST^{1107}}attP2
	E, H	P{GawB}NP5130 / + ; {20XUAS-DSCP 6XGFP}attP2 / +
	F, H	P{GawB}Dl$^{05151-G}$/ {20XUAS-DSCP 6XGFP}attP2
	G, H	P{Su(H)GBE-Gal4} / + ; {20XUAS-DSCP 6XGFP}attP2 / +
Fig. 4	A-N	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {CG10116-KD::PEST^{1107}}attP2 / P{UAS-dPF}D
Fig S1	A-F	{mira -KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / P{lacW}esg^{k00606} ; {20XUAS-DSCP-6XGFP} attP2 / {CG10116-KD::PEST^{1107}}attP2
Fig S2	Top	P{GawB}NP5130 / P{UAS-Stinger}2 ;
	Bottom	{mira-KDRT>-dSTOP-KDRT>-Gal4::p65^{1126}}attP40 / P{UAS-Stinger}2 ; {CG10116-KD::PEST^{1107}}attP2 / +
Fig S3	A-F	{3Xgbe-smGFP::V5::nls}ZH2A, M{mira-His2A.mCherry.HA} Su(hw)attP18 /
Fig S4	A-O	{Dl^{R24H06}-Gal80}ZH-2A / + ; {mira -KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / P{UAS-Stinger}2 ; + / {CG10116-KD::PEST^{1107}}attP2
Fig. S5	A,F,K,L	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / {UAS-FLAG::TEV::GFP^{1237}}attp40 ; {CG10116-KD::PEST^{1107}}attP2 / +

Fig. S5	B,G,K,L	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {CG10116-KD::PEST^{1107}}attP2 / {UAS-eIF4E::3xHA}ZH-86Fb (F000955*)*
	C,H,K,L	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {CG10116-KD::PEST^{1107}}attP2 / P{UAS-dPF}D
	D,I,K,L	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / {20XUAS-Lin-28::FLAG-HA}attP40 ; {CG10116-KD::PEST^{1107}}attP2 / +
	E,J,K,L	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / {20X UAS-FMRP::GFP::TEV::FLAG}attP40 ; {CG10116-KD::PEST^{1107}}attP2 /
Fig S6	A,E,F	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {CG10116-KD::PEST^{1107}}attP2 / {UAS-eIF4E::3xHA}ZH-86Fb (F000955*)*
	C, D, E,F	{mira-KDRT>-dSTOP-KDRT>-Gal4^{1438}}attP40, P{tubP-Gal80[ts]}20 / + ; {CG10116-KD::PEST^{1107}}attP2 / P{UAS-dPF}D

Table S2: Oligos used in this study

Note: Oligos used in Gibson Assembly are 50-60 nucleotides (nts) in length with the 5'

nts encoding plasmid sequence and the 3' nts (**in Bold**) encoding amplicon sequence.

No.	Sequence	Additional Information
3431	ATGCGCACCCGTGGCCAGGGCCGCAAGCTT**CGTCTAAGTGTGACTGATCCTGAAAGGGTT**	With 3432, amplifies 3.4kb *delta* enhancer
3432	GCCTCTATTTATACTCCGGCGCTCGACGTC**CGGAGACATCTGCTGGTGGAGGACATTATG**	With 3431, amplifies 3.4kb *delta* enhancer
3433	CATTATCAGACAAATGTCTGATATCGTTTCTCGAG**AATCAAAATGAAGCTGCTGAGTAGT**	With 3434, amplifies Gal4::p65
3434	AAGTAAGGTTCCTTCACAAAGATCCTCTAGA**TTAGGAGCTTATCTGGCTCAGCAAAGCGG**	With 3433, amplifies Gal4::p65
3437	ATGCGCACCCGTGGCCAGGGCCGCA**TGCCAGCACAGGCGGAACCCACAGCCAGGC**	With 3438, amplifies 3.4kb *βTry* enhancer
3438	CTATTTATACTCCGGCGCTCGACGT**GCGGTACTTCTTTCCGTTTTGGCCTGCGCC**	With 3437, amplifies 3.4kb *βTry* enhancer
3439	ATGCGCACCCGTGGCCAGGGCCGCA**CTAACAGCATCTAAGCTGGTACCCAGTAGG**	With 3440, amplifies 1.6kb *θTry* enhancer
3440	CTATTTATACTCCGGCGCTCGACGT**GCAGACATTTATAACTTTCCATCAGGCGCG**	With 3439, amplifies 1.6kb *θTry* enhancer
3441	ATGCGCACCCGTGGCCAGGGCCGCA**CCGCCCCAATTCGTCCGAATACCTAAGCTG**	With 3442, amplifies 1kb *CG10116* enhancer
3442	CTATTTATACTCCGGCGCTCGACGT**TATTCGTCTGAGGCATTCCTTAAGAGCTCG**	With 3441, amplifies 1kb *CG10116* enhancer
3443	ATGCGCACCCGTGGCCAGGGCCGCA**GGCAGGCGATTGTTGCCCAATGGAATGCCC**	With 3444, amplifies 2.6kb *CG18404* enhancer
3444	CTATTTATACTCCGGCGCTCGACGT**GCAATTCTTATCGTTTGGCTTATGTGGGAG**	With 3443, amplifies 2.6kb *CG18404* enhancer
3445	ATGCGCACCCGTGGCCAGGGCCGCA**CGTAGCTGGAGATGTAGCTAATGCTGCTGG**	With 3444, amplifies 0.5kb *CG18404* enhancer
3446	ATGCGCACCCGTGGCCAGGGCCGCA**TGGCGTACCTGAGGCCTTGGATCGATGCCG**	With 3447, amplifies 0.6kb *κTry* enhancer
3447	CTATTTATACTCCGGCGCTCGACGT**GGATTTCAGTTTAGACTTAGCTAGGTTGCG**	With 3446, amplifies 0.6kb *κTry* enhancer
3489	ATGCGCACCCGTGGCCAGGGCCGCA**GGTGAGTACCACGCTCTGCCATCAC**	With 3490, amplifies 2.6kb *mira* enhancer
3490	AGACAAATGTCTGATATCGTTTA**TTTCGAATTTGTTTTTTGGACTTTCGG**	With 3489, amplifies 2.6kb *mira* enhancer
3491	TTGCTGAGCCAGATAAGCTCCTAAT**GCTATTCCTTTGTAACAACTACTCT**	With 3492, amplifies 2.6kb *mira* enhancer
3492	ATCACGAGGCCCTTTCGTCTTCAAG**TGAATCTTAACGTCTGCGCAATTGC**	With 3491, amplifies 2.6kb *mira* enhancer
3626	ATGCGCACCCGTGGCCAGGGCCGCAAGCTT**AATTCGATATCAAGCTTGCACAGGTCCTGT**	With 3627, amplifies 2.6kb *tubulin* enhancer
3627	GCCTCTATTTATACTCCGGCGCTCGACGTC**GCGGATCTGATACCCCGGGCGGGTACCTTC**	With 3626, amplifies 2.6kb *tubulin* enhancer
4401	TGTCTGATATCGTTTCTCGAGAAGCTTAAC**GATCCGGGATCCCGTAAGCAAATAAACAAG**	With 4402, amplifies Gal4
4402	CGATAGAGTAGTTGTTACAAAGGAATAGCA**CCTACTCCTTCTTTGGGTTCGGTGGAGTAT**	With 4401, amplifies Gal4