Activation of the Crtc2/Creb1 transcriptional network in skeletal muscle enhances weight loss during intermittent fasting

Nelson E. Bruno[1]*, Jerome C. Nwachukwu[1], Sathish Srinivasan[1], Richard Hawkins, David Sturgill[3], Gordon L. Hager[3], Stephen Hurst[4], Shey-Shing Sheu[4], Michael D. Conkright[5], Kendall W. Nettles[1]*

[1]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

[2]Department of Cancer Biology, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

[3]Laboratory of Receptor Biology and Gene Expression, National Cancer Institute, NIH, Bethesda, MD 20892, USA

[4]Center for Translational Medicine, Department of Medicine, Thomas Jefferson University, Philadelphia, PA, 19107, USA.

[5]Department of Shared Services, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

Correspondence: Nelson E Bruno, nbruno@scripps.edu; Kendall Nettles, knettles@scripps.edu

Conflict of interest statement: The authors declare no conflicts of interest.

Abstract

Exercise is a behavior modification indispensable for long-term weight loss. Exercise activates the Creb-Regulated Transcriptional Coactivator (Crtc) family of transcriptional coregulators to drive Creb1-mediated anabolic transcriptional programs in skeletal muscle. Here, we show that induced overexpression of a skeletal muscle specific Crtc2 transgene in aged mice leads to greater weight loss during alternate day fasting, and selective loss of fat rather than lean mass. Transcriptional profiling revealed that fasting and weight loss downregulated most of the mitochondrial electron transport genes and other regulators of mitochondrial function that were substantially reversed in the Crtc2 mice, which maintained higher energy expenditure during fasting. The Crtc2 mice displayed greater mitochondrial activity, metabolic flux capacity for both carbohydrates and fats, improved glucose tolerance and insulin sensitivity, and increased oxidative capacity before the fast, suggesting muscle-intrinsic mechanisms in support of improved weight loss. This work reveals that Crtc2/Creb1-mediated signaling coordinates metabolic adaptations in skeletal muscle that explain how Crtc2/Creb contribute to the effects of exercise on metabolism and weight loss.

Introduction

Long-term dieting is usually not effective in maintaining a lower body weight due to increased hunger and a slower metabolic rate (1). In humans, loss of >10% of body weight also increases metabolic efficiency, such that ~500 less calories per day are required to maintain the same energy expenditure (2). Weight loss reduces the metabolic rate due to a number of changes including increased circulating leptin and decreased thyroid hormone and sympathetic nervous system (SNS) activity (2-4). People who maintain long-term weight loss almost invariably exercise (5), and both preclinical and clinical data suggest that this is through raising metabolic rate (6-8), decreased energy efficiency (9), and increased energy expenditure from the increased physical activity (10).

Creb1-regulated transcription is stimulated by GPCRs that activate adenylyl cyclases to convert ATP to the second messenger cAMP, which in turn liberates the catalytic subunit of protein kinase A (PKA) (11). Crtc1–3 family members are primary coactivators for Creb1 that are also highly conserved in the cAMP transcriptional response. Crtc1–3 are rendered cytoplasmic and inactive by phosphorylation (12). In contrast, they are fully activated through synergistic dephosphorylation by phosphatases downstream of PKA and calcium flux in polarizable tissues including the skeletal muscle (13), hippocampus (14), cortex (15), islet cells (12), and the renal proximal tubule (16). In skeletal muscle, Crtc/Creb1 is activated by exercise downstream of the β-adrenergic receptor and calcium flux, stimulating anabolic transcriptional programs, including induction of PGC-1α (13, 17) and in vitro enhancement of oxidative capacity (17). These training adaptations lead to significant enhancement of high intensity exercise performance along with greater glycogen and intramuscular triglyceride stores (13). While Crtc/Creb1 coordinates metabolic transcriptional programs in other tissues such as adipocytes, liver, and pancreas, its role in skeletal muscle metabolism has not been studied in vivo.

To investigate the effects of Crtc2/Creb1 activation in adult skeletal muscle, we generated

doxycycline (dox)-inducible transgenic mice that show skeletal muscle-specific overexpression of

Crtc2 (FLAG-Crtc2 S171A/S275A/K628R mutant that prevents cytoplasmic localization) under

the control of a tetracycline response element (13). Given the primary role of exercise adaptations

in regulating whole-body metabolism, we examined here the role of the Crtc2/Creb1

transcriptional network in skeletal muscle metabolism during alternate day fasting (ADF),

mimicking a common dieting scheme in humans. Activation of Creb1-regulated transcriptional

responses during fasting enabled maintenance of a lower body weight due to the stimulation of

higher energy expenditure and other metabolic changes, enabling selective loss of fat mass and

retention of lean mass, as observed in humans who exercise (18, 19). Crtc2 mice displayed the

metabolic hallmarks of an exercised phenotype, including enhanced mitochondrial activity,

improved lipid and carbohydrate flux capacity and storage, and insulin sensitization. These data

support a model where lower metabolic activity during fasting and weight loss contributes to

weight regain, which can be reversed by exercise-induced activation of Crtc2/Creb1-regulated

transcriptional programs.

Results

Crtc2 overexpression in skeletal muscle facilitates weight loss

We studied aged Crtc2 (rtTA/Crtc2^{tm}) or WT (rtTA/WT) mice, which in the C57BL/6 background

naturally gain fat mass as they age (18 weeks, 30–39 g, **Figure 1A–B**). During dox treatment,

Crtc2 mice gained weight similarly to control WT mice (**Figure S1A**). On the ADF protocol, three

separate cohorts of Crtc2 mice displayed significantly greater and sustained weight loss than WT

mice, with Crtc2 mice losing about twice as much weight in each case, 9-14% of body weight

(**Figure 1C, Figure S1B**).

After 3 weeks of ad libitum refeeding, the Crtc2 mice re-gained more weight to catch up

with the WT mice (**Figure 1D**), which may be due to the anabolic effects of higher insulin

sensitivity, described below. We found that locomotion was similarly reduced in both groups

during the refeeding period (**Figure S1C**), the number of meals doubled in both groups, while

meal duration and number of bouts per meal declined similarly (**Figure 1E**). We observed a

reduction in kilocalories consumed in the Crtc2 relative to the WT group, which planned

comparisons demonstrated was significant only during the second ADF cycle, when the WT mice

increased consumption at a faster rate than the Crtc2 mice (**Figure 1E**). Increases in skeletal muscle

storage of glycogen and triglycerides associated with the anabolic phenotype of Crtc2/Creb1

activation (13) likely contributed to the early response to fasting, perhaps explaining the initially

slower rate of increased food consumption in the Crtc2 mice. These differences in caloric

consumption are consistent with the differences in body weight between cohorts and disappear

after normalization for body weight. Control mice lost more lean mass, while Crtc2 mice

conserved lean mass and burned the excess fat accumulated during aging (**Figure 1F**), similar to

the effects of exercise (20). To verify that Crtc2 helps maintain lower body weight in a different

model, we placed mice on a high fat diet, and found that Crtc2 mice also maintained lower body

weight than WT mice in this context (**Figure 1G**). Thus, the activation of Crtc2/Creb1 signaling

in skeletal muscle significantly modulated body composition and weight loss, mimicking the

effects of exercise training including preferential loss of fat mass.

The transcriptional response to fasting and weight loss and its regulation by Crtc2

To understand the skeletal muscle intrinsic effects of Crtc2 on dieting and weight loss, we performed

transcriptional profiling of mice tibialis anterior (TA) muscles electroporated with control and Crtc2

expression plasmids in contralateral legs. One cohort of electroporated mice was subject to 3 cycles

of ADF, while the other cohort was fed ad libitum. TA muscle tissues were collected from both

cohorts at the end of the third 24-hr fast. The transduction increased *Crtc2* mRNA levels by ~2.5-

fold across groups (**Figure S2A**). In response to ADF, we identified 1,880 differentially expressed

genes (DEGs) in the control-transduced muscle and 3,197 DEGs in Crtc2-transduced muscle, with

1,380 genes in common between the two groups (**Figure S2B, Supplementary Dataset_1**).

Analysis of canonical signaling pathways and gene ontology (GO) annotations revealed that ADF

modulated the transcriptional programs for lipid and carbohydrate metabolism, fatty acid beta-

oxidation, and insulin signaling (**Figure S2C-D**). ADF also regulated transcriptional programs that

control protein balance, including autophagy and ubiquitin-dependent proteasomal degradation,

RNA splicing, and transcription (**Figure S2C-D, Supplementary Dataset_1**). There were also over

200 DEGs regulated by ADF that were annotated as transcription factors and transcriptional

coregulators along with > 50 transcription factor and coregulators altered by Crtc2 over-expression

(**Figure S2E-G, Supplementary Dataset_1)**.

Crtc2 reverses fasting- and weight loss-induced effects on mitochondria and energy
expenditure

124   In addition to lower metabolic rate, fasting and weight loss render humans and other mammals more

125   energetically efficient, such that the same amount of work is performed with less ATP production,

126   suggesting a remodeling of glycolytic or oxidative capacity (2, 21). ADF modulated the expression

127   of >200 genes whose proteins localize in mitochondria according to MitoCarta (22) (**Figure 2A**).

128   ADF downregulated most of the genes that encode the electron transport chain (complexes I–IV),

the ATP synthase (complex V), the mitoribosome, and the cytosolic ribosome (**Figure 2A-C**, **Figure S3A**), but significantly upregulated genes such as *Sdha*, *Uqcrc1,* and *Atp5a1* (**Figure 2B**, **Figure S3A**).

We observed a significant ADF-induced expression of genes encoding ROS scavenger proteins such as thioredoxin (*Txnrd1*) and catalase (*Cat*), and genes regulating mitochondrial stability*, Tomm7, Timm9,* and the AAA protease *Spg,7* known to be active during the removal of damaged mitochondrial proteins (**Figure 2D**). Among the ATP synthase genes, there was a more pronounced loss of gene expression in the lateral stalk genes (**Figure 2B**), which may decrease the conductance of mitochondrial permeability transition pore (mPTP) and thus protect mitochondria from detrimental swelling and oxidative injury (23). ADF also induced increases in mitochondrial fusion (*Mfn2* and *Opa1)*, and decreased fission (*Fis)* gene expression, consistent with the protective effects of fusion during nutrient deprivation (24). These adaptations could be advantageous for weight loss as they prepare the mitochondria for the reduced metabolic rate and increased reliance on lipid metabolism during fasting, without causing permanent damage to the mitochondrial pool within the cell.

In mice subjected to ADF, Crtc2 transduction altered expression of 80 genes whose proteins localize in the mitochondria. Crtc2 expression strongly attenuated the ADF-dependent repression of electron transport chain genes and the mitoribosome (**Figure 2A–C**, **Figure S3A**), and generally upregulated the other genes encoding proteins that localize to mitochondria. In addition, Crtc2 enhanced expression of ribosomal genes that facilitate mitoribosome assembly, and genes such as *Spg7, Pink1, RhoT2*, and *Chchd3* that control the quality of mitochondrial proteins, but blocked ADF-dependent downregulation of *Minos1*, a central component of the mitochondrial inner membrane organizing system (**Figure 2D**). Crtc2 also upregulated expression of *Tefm*, the

mitochondrial transcriptional elongation factor, and *Bola3*, which participates in electron transport

chain assembly (25).

To understand how ADF affects mitochondria, we noted that glucocorticoids (GCs) are

one of the primary drivers of catabolic changes during fasting, and in response to ADF and Crtc2

overexpression, we observed over a thousand DEGs that were annotated as "glucocorticoid

response" genes (**Figure S2C**). In the context of an overnight fast, treatment with Dex more than

doubled loss of lean mass (**Figure 2E**). Using high-content imaging, we observed that Dex reduced

the diameter and mitochondrial potential of nutrient-deprived primary mouse skeletal myotubes

(**Figure 2F**). GCs can also activate the master energy sensor, AMP-activated protein kinase

(AMPK) (26), which can act as a glucocorticoid receptor (GR) coactivator (27). AMPK is

activated by a lower ATP/ADP ratio during fasting to inhibit anabolism and stimulate catabolic

processes such as autophagy and mitophagy. In primary myotubes, we observed a Dex-dependent

activating phosphorylation of AMPK that increased over time, consistent with an underlying

transcriptional mechanism (**Figure 2G**). Activation of AMPK with both Dex and the AMP analog,

AICAR, had an additive detrimental effect on myotube diameter (**Figure 2H**), suggesting that GR

collaborates with AMPK to drive skeletal muscle atrophy.

We also observed that ADF modulated the expression of genes encoding three of the

AMPK subunits. Changes in subunit composition are known to alter the regulation, substrate

preferences, and cellular location of AMPK. The α2 and β2 AMPK isoforms were significantly

upregulated by ADF (**Figure 2I**, SI Appendix). The α2 subunit is activated by low glycogen and

mediating nuclear translocation (28), while the β2 subunit is required for maintenance of muscle

ATP levels during fasting (29). In contrast, ADF significantly downregulated the γ3 AMPK

isoform (**Figure 2I**), which is required for glycogen synthesis (30), a process inhibited during

fasting. Thus, a combination of GR and AMPK signaling drives skeletal muscle atrophy during

nutrient deprivation and a loss of mitochondrial potential.

We further examined whether ADF- and Crtc2-induced changes in mitochondria—the

powerhouse of cellular respiration—are associated with changes in global energy expenditure. As

expected, ad libitum oxygen consumption cycled with activity in the mice, which increased at

night and decreased during the day (**Figure 2J, Figure S3B**). Fasting reduced oxygen consumption

at every cycle, and this effect was more pronounced during the day as the fast was extended

(**Figure S3B**). However, Crtc2 mice consistently showed higher oxygen consumption,

independent of fasting (**Figure 2J**).

When fed ad libitum, Crtc2 and control mice display similar rates of energy expenditure

(EE) (**Figure 2K**). We calculated EE from indirect calorimetry measurements of VO_2 and VCO_2

and performed ANCOVA analysis using body weight and NMR data as covariates to adjust the

EE to total body weight and body composition (**Figure S3C**). While fasting, the EE of WT mice

dropped significantly. In contrast, during fasting EE in Crtc2 mice did not drop below the ad

libitum levels (**Figure 2K**). Respiratory quotient was also reduced during fasting, indicating

greater utilization of fat for fuel, but the differences between WT and Crtc2 mice were not

significant (**Figure S3D–E**). These data suggest potential mechanisms to explain how physical

activity—through Crtc2/Creb1-regulated transcriptional programs—can help maintain a higher

metabolic rate and a lower body weight by upregulating expression of mitochondrial respiratory

chain subunits and modulating expression of other mitochondria-localized gene products, consistent

with the well-known beneficial effects of exercise on mitochondrial biogenesis and energy

expenditure.

Crtc2 stimulates mitochondrial activity and improves oxidative capacity

Prolonged fasting promotes lipolysis, insulin resistance, and mobilization of fatty acids to skeletal muscle through the combined effects of glucagon and glucocorticoids on adipocytes. These actions augment the rate of fatty acid oxidation by mitochondria and the ectopic accumulation of IMTGs. As expected, ADF upregulated a panoply of genes involved in the transport, oxidation, and esterification of fatty acids (**Figure S4A**).

Crtc2 mice displayed higher oxygen consumption before the fast; therefore, we examined them for improved mitochondrial function. After one week of dox treatment, Crtc2 mice showed increased mitochondrial succinate dehydrogenase activity in the gastrocnemius muscle (**Figure 3A, Figure S4B–C**) and increased expression of *Cytochrome C (Cycs)* (**Figure 3B**). Overexpression of PGC-1α increases mitochondrial density and exercise performance, but without the increase in metabolic rate that can occur from exercise training (31-34), highlighting the requirement for additional exercised-induced signals. In addition to the previously reported direct upregulation of *Ppargc1a/*PGC-1α by Crtc2 (13, 17), Crtc2 mice showed increased expression of *Nr4a3/*NOR-1 nuclear receptor (**Figure 3C**) which induces mitochondrial biogenesis and increases oxidative capacity (35, 36), and is induced by high intensity exercise (13). Crtc2 also upregulated the *Dgat1* gene (**Figure 3C**), which is a direct Creb1 target gene (13), the product of which catalyzes a rate-limiting step that increases IMTG stores. These mitochondria-related genes were also upregulated in primary myotubes overexpressing Crtc2, indicating a skeletal muscle-intrinsic effect of Crtc2 that mimics the effects of exercise (**Figure 3C**). To show that Creb1 is directly regulating atrophy, we transduced primary myocytes with the dominant negative A-Creb, which reduced myotube diameter (**Figure 3D**). We also observed a doubling of mitochondrial DNA after in vivo electroporation of Crtc2 (**Figure 3E**). The increased mitochondrial mass was associated with

increased oxidative capacity, as myotubes overexpressing Crtc2 or Crtc3 oxidized more palmitate

than control myotubes (**Figure 3F**), as previously reported (17).

Crtc2 improved glucose tolerance and carbohydrate metabolism before and during weight loss

ADF altered regulation of a number of carbohydrate metabolism and insulin signaling genes (**Figure**

S2D, Figure S5A, Supplementary Dataset_1). To probe for Crtc2-dependent changes in

carbohydrate metabolism, we performed glucose tolerance tests comparing Crtc2 and WT mice

following ADF or normal diet. The Crtc2 mice displayed significantly improved glucose disposal

after ADF, including a dramatic improvement in their ability to reduce glycemia (**Figure 4A**), and

required a lower insulin excursion (**Figure 4B**). Crtc2 overexpression also improved insulin

tolerance after fasting and weight loss (**Figure 4C**). These effects were also apparent after 1 week

of dox treatment in Crtc2 mice fed ad libitum (**Figure S5B–D**). In primary myotubes, expression of

Crtc2 was sufficient to significantly increase the uptake of 2-Deoxy-D-glucose (**Figure 4D**).

Crtc2 mice and Crtc2/3-overexpressing myotubes upregulated a number of carbohydrate

signaling genes in normal fed conditions at both the RNA and protein levels (**Figure 4E–H**),

including the glucose transporter, *Glut4*, whose insulin-dependent translocation to the membrane

drives glucose uptake into skeletal muscle, accounting for 80% of glucose disposal. The *Nr4a1* gene

encodes Nur77, a nuclear receptor family transcription factor that regulates transcriptional programs

for carbohydrate metabolism, lipolysis, and mitochondrial biogenesis (37), and is upregulated by

high intensity exercise or adrenergic signaling (13). *Irs2*, *Akt1*, and *Akt2* encode core components of

the insulin receptor signaling pathway that can coordinate anabolic signaling and improve

carbohydrate metabolism. We also identified a significant inhibition of myogenin expression in the

normal fed Crtc2 mice (**Figure 5E**), a primary driver of muscle atrophy during fasting. To validate

on target mechanism of action, we generated stable cell lines overexpressing Crtc3, dominant

negative A-Creb, or dominant negative Crtc2 expressing only the Creb binding domain (12). These

results demonstrate that A-Creb and dn-Crtc2 blocked basal expression of PGC-1α, Irs2, and Akt2,

and prevented phosphorylation of Akt1/2 at Thr308 (**Figure 4G–H**).

To test for pre-fasting changes in the central control of feeding and energy expenditure, we

completed a leptin infusion study, as leptin secretion from adipocytes both suppresses hunger and

stimulates the SNS. The WT and Crtc2 mice displayed identical patterns of weight loss following

leptin infusion (**Figure 4I**), and there were also no differences in plasma levels of feeding hormones,

cholesterol, or triglycerides (**Figure S5E–F**). Lastly, we considered that Crtc2 might be regulating

secreted proteins that modulate the response to dieting, using the Metazoan Secretome

Knowledgebase (38). Among 2,500 genes encoding likely secreted proteins, 171 DEGs were

regulated by Crtc2 during ADF. Using the Reactome.org peer-reviewed pathway database, we

detected significant enrichment of pathways regulating extracellular matrix, inter-cell signaling, and

immune function, among others (**Figure S5G**), but no changes in myokines that would drive weight

loss. However, Crtc2 signaling in skeletal muscle would be expected to have effects on other tissues

just from changes in metabolic flux, as Crtc2 mice displayed pre-fast increases in skeletal muscle

oxidative flux capacity.

This work reveals mechanisms through which diet and exercise regulate substrate mobilization

and metabolism and highlights how these lifestyle modifications can alter mitochondrial function,

energy expenditure, and the different dramatic metabolic rewiring associated with fasting versus

Crtc/Creb1 signals that are downstream of exercise. Reduced energy expenditure and increased

hunger during fasting are partly mediated by a lowering of the satiety hormone leptin, which acts

centrally to stimulate the SNS (2-4). Leptin is also secreted basally from adipose tissue in proportion

to fat mass, so it is chronically lower during fasting, which lowers sympathetic tone and energy

expenditure, along with reductions in circulating thyroid hormone and stimulating the HPA axis

(**Figure 5A**). Within skeletal muscle, increased glucocorticoids and AMPK activity are sufficient to

reduce mitochondrial potential (**Figure 2**), which along with likely contributions from fasting-

induced lowering of thyroid hormone, βAR, and insulin receptor activity, induce mitochondrial

dynamics, mitophagy, and the metabolic rewiring to conserve glucose for the brain (**Figure 5A**).

These energetic effects are reversed by exercise (**Figure 5B**), which we now show can be

recapitulated by enforced expression of Crtc2/Creb1 and upregulation of mitochondrial biogenesis

master regulatory genes, *Essrra/*ERRα (**Figure S2E**), *Nr4a3* (NOR1) (**Figure 3B–C**),

*Ppargc1a/*PGC-1α (**Figure 3B**), and *Nr4a1* (NUR77) (**Figure 4E–F**). The combination of βAR and

Ca^{2+} signaling are required to maximally stimulate nuclear translocation of Crtc2 in skeletal muscle

(13). In this model, increased GC and AMPK activity during exercise promotes mitochondrial

dynamics, which is then coupled with mitochondrial biogenesis to increase mitochondrial potential

and mass, in coordination with other metabolic changes (**Figures 3–4**). Overexpression of Crtc2

circumvented the neuronal activation of Creb1, as Crtc2 mice had most of the hallmarks of exercise

adaptations before fasting and weight loss (**Figures 3–5**) (13).

Materials and Methods

Statistical analyses

Changes in body weight and body were analyzed with 2-way analysis of variance. For VO_2,

respiratory quotient, and ambulation data, the ad libitum and fasted periods were analyzed

separately with 2-way ANOVA, and the refeeding data analyzed with Student's T-test. For the

energy consumption data, the fasted and re-fed data were analyzed together with 2-way ANOVA.

Gene expression data were analyzed with 1-way ANOVA, with Dunnett's test for multiple

comparisons of each gene to its corresponding control. Glycemia and insulin changes over time

were analyzed with 2-way ANOVA, and the AUC assessed with Student's T-test. The behavioral

data for meal consumption were analyzed with Student's T-tests.

295   Crtc2 transgenic mice and animal care

All animal procedures were approved by the Scripps Research Institute ICACUC. TRE-Crtc2[tm]

mice were generated by cloning FLAG-Crtc2 (S171A, S275A, K628R) into pTRE-Tight. Oocyte

injections were conducted into C57Bl6 at the University of Cincinnati. Double transgenic mice

(DTg) were generated by crossing C57BL/6-Tg(TRE-Crtc2tm)1Mdc/j mice with 1256[3Emut]

MCK-rtTA transgenic mice, which expresses the reverse tetracycline transactivator (rtTA)

regulated by the 1256[3Emut] MCK promoter specifically in skeletal muscle (Grill *et al*, 2003).

Wild type and TRE littermates were used as controls and treated with DOX in the same manner as

DTg mice. High fat diet (Research Diet D12450J) contained 60 kcal% fat and matched control low

fat diet (D12492) contained 10kCal% fat and 7% sucrose.

306   Glucose and insulin tolerance tests

307   Glucose tolerance testing was performed after an overnight fast. Blood glucose was quantified at

308   15, 30, 60, 90 and 120 min post-i.p. administration of 20% D-glucose (1 g/kg fasted body weight)

309   with an Accu-check glucose monitor as described (Heikkinen *et al*, 2007). Insulin tolerance testing

310   was performed on 6 hr-fasted mice injected i.p. with human insulin (1.5 U/kg fasted body weight).

311   Blood glucose levels were determined immediately before, and 15 and 30 min after, injection.

In vivo electroporation

Plasmid DNA (50 mg) was transfected into mouse tibialis anterior (TA) muscles by electroporation as described previously (13). 1 hr before electroporation mice were anesthetized, and a small incision was made through the skin covering the TA muscle and then injected with 30 Al of 0.5 U/Al hyaluronidase and then injected with plasmid DNA. *Crtc2* and *GFP* expression plasmids were purified from DH5a *Escherichia coli* cultures using an EndoFree plasmid kit (Qiagen, Valencia, CA, USA), and resuspended in 71 mM sterile PBS. After the injections, electric pulses were administered by using non-invasive platinum-coated tweezertrodes. Eight 100-ms square-wave electric pulses at a frequency of 1 Hz and at 200-ms intervals were delivered with an ECM 830 electroporation unit (BTX-Harvard Apparatus, Holliston, MA, USA) with a field strength of 40 V/cm. After the electroporation procedure, the incision was closed with Vetbond surgical glue.

Indirect calorimetry and energy expenditure

Mice were individually housed in a sixteen chamber Oxymax comprehensive lab animal monitoring system (CLAMS; Columbus Instruments, Columbus OH) at 26° C. Animals were allowed to acclimate for 1 day followed by VO_2 and VCO_2 were measured at 13 min intervals over a continuous duration of 60 hr. Energy expenditure (EE) was calculated using the Lusk equation, kcal/h = $(3.815 + 1.232 \times RQ) \times VO_2$ in l/h, where RQ is the ratio of VCO_2 to VO_2. Opto-Varimetrix-3 sensor system in the x-, y-, and z-axes of each chamber recorded sterotype, ambulatory and z-axis activity. Consecutive adjacent infrared beam breaks in either the x- or y-axes were scored as ambulatory activity.

Body composition

Body composition was determined using time domain nuclear magnetic resonance (TD-NMR) in a Bruker minispec.

Automated food intake monitoring

Feeding behavior was assessed using a BioDAQ episodic Food Intake Monitor (BioDAQ, Research Diets, Inc) as described (Stengel *et al*, 2010). Feeding parameters were calculated with BioDAQ Monitoring Software 2.2.02. Feeding bouts were defined as a change in stable weight of food. Meals consisted of two or more bouts within 5 s with a minimum meal amount of 0.02 g. Clusters of bouts occurring greater than 600 seconds apart were considered separate meals. Water or water containing Dox was provided *ad libitum* from regular water bottles. Mice habituated to the new environment within 1 day and showed normal food intake and regular body weight gain.

mRNA-seq and bioinformatics

Total RNA was extracted from tissues using TRIzol™ reagent (Invitrogen), and cleaned up using the RNeasy kit with on-column DNA digestion (QIAGEN). Three mRNA-seq libraries per condition were prepared using the NEBNext Ultra II directional RNA library Prep kit for Illumina (New England Biolabs), and read in paired-end 40-cycle sequencing runs on the NextSeq 500 system (Illumina). Sequences were aligned to the Ensembl *Mus musculus* GRCm38.83.chr.gtf genome (mm10) assembly using HISAT2 v2.0.5 (PMID: 25751142), sorted and indexed using SAMtools v1.7 (39) and quantified for differential expression analysis using Subread v1.6.1 featureCounts (40). Differential expression analysis was performed using DESeq2 v1.18.1 (41, 42). Enriched canonical pathways were identified using the Ingenuity pathways analysis software

(QIAGEN). Gene Ontology (GO) analysis excluding inferred electronic annotations, was

performed using the Mouse Genome Informatics GO Browser and GO Term Mapper

(http://www.informatics.jax.org). Genes encoding mouse nuclear hormone receptors were

obtained from the Nuclear Receptor Signaling Atlas (NURSA) (43). Genes encoding

mitochondrial proteins were obtained from the mouse Mitocarta2.0 (22, 44). Genes encoding

transcriptional coregulators were obtained from the EpiFactors database (45). Glucocorticoid-

responsive genes were obtained by combining previously published datasets (46) reanalyzed with

GEO2R, with other dexamethasone-regulated genes that we identified in C2C12 myotubes, GEO

accession no. GSE149453. Heatmaps were prepared using Pheatmap v1.0.8, RColorBrewer v1.1–

2, and other software packages available in R. Other data visualizations were performed using

GraphPad Prism7.

Glucose uptake

371 Insulin mediated 2-deoxy-D-[^{14}C] glucose uptake was determined by incubating primary

372 myotubes with PBS or PBS containing 100 nM insulin for 30 min followed by addition of 0.1 mM

373 of cold 2-deoxy-D-glucose and 0.1μCi/well of 2-deoxy-[^{14}C] D-glucose for 5 min prior to

374 solubilization. Nonspecific deoxyglucose uptake was measured in the presence of 20 μM

375 cytochalasin B and subtracted from the total glucose uptake.

Primary myoblast isolation and culture

Primary myoblasts were isolated from P1 to P3 (1–3 day old) C57BL/6J pups as described

(Springer *et al*, 1997) and cultured on collagen coated plates in media consisting in 1:1 F-

10/DMEM supplemented with 20% FCS and 25 μg/ml bFGF. Subconfluent myoblasts were

differentiated into multinucleated myotubes using a 1:1 F-10/5 mM glucose DMEM supplemented

with 4% heat inactivated horse serum (Hyclone) for up to five days. For steroid-free culture

conditions, cells were grown in charcoal:dextran-stripped FBS (Gemini Bio-products, cat no. 100-

119).

Adenoviruses

Control GFP-expressing and Crtc2 expressing adenoviruses have been described (Koo *et al*, 2005;

Wu *et al*, 2006). For all experiments myotubes were infected after 72 hr of differentiation with

4×10^8 viral particles per ml per well for 24–48 hr.

Immunoblotting analysis

Immunoblot analysis was performed as described (Conkright *et al*, 2003a). Whole cell extracts

were prepared in HEPES whole cell lysis buffer (20 mM HEPES, pH 7.4, 1% TX-100,1x

phosphoSTOP complete inhibition cocktail tablets [Roche], and 1X Complete-EDTA Free

protease inhibitor cocktail [Roche]). Frozen muscle tissue was homogenized in lysis buffer T-

PER® Tissue Protein Extraction reagent (Thermo Scientific) by a polytron homogenizer and

sonication. The protein concentration was determined using the protein assay reagent (Bio-Rad).

60–80 μg of protein were separated by SDS-PAGE and transferred to nitrocellulose membranes

for immunoblotting. Nur77/Nr4a1 #2518 Rabbit polyclonal antibody was from Novus Biological.

CRTC2 #A300-638A Rabbit polyclonal antibody was from Bethyl Laboratories. Irs2 (L1326)

#3089S Rabbit polyclonal antibody was Cell Signaling Technology, Inc. Tubulin #T9026-.2ML

Mouse Monoclonal antibody was from Sigma-Aldrich. pAKT(Thr308) (244f1)Rb and

pAKT(Ser473) (971s) were from Cell Signaling Technology. PGC1α ab54481 Rb was from

Abcam.

Quantitative RT-PCR (qPCR)

Total RNA from tissue or cells was harvested using Rneasy RNA Kit (Qiagen) and cDNA prepared

using Transcriptor High Fidelity cDNA Synthesis Kit (Roche). Relative abundance of cDNAs was

determined a Roche LightCycler 480, and the resulting data were normalized to *Rpl-23* ribosomal

RNA (IDT) as described (Conkright *et al*, 2003b). The primer pairs are as follows: *Nr4a1* (5′-

ttctgctcaggcctggtact′, 5′-gattctgcagctcttccacc-3′); *Nr4a3* (5′-tcagccttttttggagctgtt′, 5′-

tgaagtcgatgcaggacaag-3′); *Irs2* (5′-acaacctatcgtggcacctc-3′, 5′-gacggtggtggtagaggaaa-3′); Idh3a

(5′-gtgacaagaggttttgctggt-3′, 5′-tgaaatttctgggccaattc-3′); *Cytc* (5′-gatgccaacaagaacaaaggt-3′, 5′-

tgggattttccaaatactccat-3′); *Glut4* (5′-tgtggctgtgccatcttg-3′, 5′-cagggccaatctcaaagaag-3′).

Mitochondrial DNA (mt-DNA) analysis

Total DNA was extracted using the GenElute™ mammalian genomic DNA miniprep kit (Sigma-

Aldrich). Triplicate real-time PCR reactions were performed using 50 ng of total DNA, *CoxI* (5′-

agcattcccacgaataaataaca-3′, 5′-agcattcccacgaataaataacat-3′) or *CoxII* (5′-tttcaacttggcttacaagacg-3′,

5′-tttcaacttggcttacaagacg-3′) mitochondrial DNA primers, and a control genomic DNA primer,

Gapdh (5'-gaaaaggagattgctacg-3', 5'-gcaagaggctaggggc-3').

Histology

Dissected muscle tissue was mounted in OCT medium (TissueTek) and froze in liquid N_2–cooled isopentane. Tissue sections were stained for succinate dehydrogenase activity to distinguish between oxidative and nonoxidative fibers using standard methods.

Fatty acid metabolism

Fatty acid oxidation was assayed by incubating $(9,10(n)-^3H)$ palmitic acid (60 Ci/mmol) bound to fatty-acid free albumin (final concentration:100 μM, palmitate:albumin 2:1) and 1 mM carnitine with primary myocytes for 2 hr. Tritiated water released was collected and quantitated.

High-content imaging and analysis

Primary or C2C12 myotubes in black 96- or 384-well tissue culture plates with clear bases (Greiner Bio-One, North America, Inc.) were stained for 15 min with 200 nM MitoTracker® Orange CM-H_2TMRos dye (Invitrogen™ by ThermoFisher Scientific). The myotubes were rinsed with steroid-free DMEM to remove un-incorporated dye, and then incubated in the differentiation media for 45 min to reach peak fluorescence. Myotubes were fixed in 4% formaldehyde for 20 min, stained with 300 nM DAPI for 5 min, permeabilized in PBS containing 0.1% Triton X-100 for 20 min, blocked for 1 h with 1x TBS containing 0.1% Tween-20 (TBS-T) and 2.5% normal goat serum, and incubated at 4°C overnight with an AlexaFluor® 488-conjugated anti-skeletal muscle myosin (F59) antibody (Santa Cruz Biotechnology, Inc. cat no. sc-32732 AF488). The next day, the myotubes were washed 4 times with TBS-T to remove unbound antibodies and rinsed twice with PBS. The stained myotubes were imaged at 10–20X magnification on the IN Cell Analyzer 6000 platform (GE Healthcare). For each treatment condition, a stack of 24–27 images containing an average of

50–100 myotubes per image were analyzed using the IN Cell Developer Toolbox image analysis

software, with a customized segmentation protocol for myotubes. The average (mean) diameter

and mitochondrial potential (i.e. MitoTracker staining density x area) of myotubes in these images

were then calculated.

Data Availability

The raw and processed RNA-seq dataset is freely available to the public at the National Center for

Biotechnology   Information   (NCBI)   Gene   Expression   Omnibus   (GEO)

https://www.ncbi.nlm.nih.gov/geo/ and can be retrieved with the accession number GSE149150,

which will be released upon publication of this manuscript. The processed RNA-seq dataset is also

provided in SI Appendix, **Dataset S1**. Relevant protocols and raw data not detailed in the Methods

or Supplementary Information are available upon request.

Acknowledgements. N.E.B. was supported by the BallenIsles Men's Golf Association. J.C.N. was

supported by the Frenchman's Creek Women for Cancer Research.

Author contributions Conceptualization, N.E.B., M.D.C., K.W.N.; Methodology, N.E.B., M.D.C.;

Investigation, N.E.B., J.C.N., S.S., R.H., D.S.; Writing – Original Draft, K.W.N., N.E.B., M.D.C.,

S.H., S.S.S.; Writing – Revision, K.W.N, J.C.N., N.E.B.; Supervision, M.D.C, G.L.H., K.W.N.,

S.S.S., N.E.B.

References

1. C. L. Melby, H. L. Paris, R. M. Foright, J. Peth, Attenuating the Biologic Drive for Weight Regain Following Weight Loss: Must What Goes Down Always Go Back Up? *Nutrients* **9** (2017).

2. M. Rosenbaum *et al.*, Effects of experimental weight perturbation on skeletal muscle work efficiency in human subjects. *Am J Physiol Regul Integr Comp Physiol* **285**, R183-192 (2003).

3. L. J. Arone, R. Mackintosh, M. Rosenbaum, R. L. Leibel, J. Hirsch, Autonomic nervous system activity in weight gain and weight loss. *Am J Physiol* **269**, R222-225 (1995).

4. M. Rosenbaum *et al.*, Low-dose leptin reverses skeletal muscle, autonomic, and neuroendocrine adaptations to maintenance of reduced weight. *J Clin Invest* **115**, 3579-3586 (2005).

5. R. M. Foright *et al.*, Is regular exercise an effective strategy for weight loss maintenance? *Physiol Behav* **188**, 86-93 (2018).

6. P. S. MacLean *et al.*, Regular exercise attenuates the metabolic drive to regain weight after long-term weight loss. *Am J Physiol Regul Integr Comp Physiol* **297**, R793-802 (2009).

7. Y. Turk *et al.*, High intensity training in obesity: a Meta-analysis. *Obesity science & practice* **3**, 258-271 (2017).

8. A. Mourier *et al.*, Mobilization of visceral adipose tissue related to the improvement in insulin sensitivity in response to physical training in NIDDM. Effects of branched-chain amino acid supplements. *Diabetes care* **20**, 385-391 (1997).

9. M. Rosenbaum *et al.*, Resistance Training Reduces Skeletal Muscle Work Efficiency in Weight-Reduced and Non-Weight-Reduced Subjects. *Obesity (Silver Spring)* **26**, 1576-1583 (2018).

10. D. M. Ostendorf *et al.*, Physical Activity Energy Expenditure and Total Daily Energy Expenditure in Successful Weight Loss Maintainers. *Obesity (Silver Spring)* **27**, 496-504 (2019).

11. D. A. Walsh, J. P. Perkins, E. G. Krebs, An adenosine 3',5'-monophosphate-dependant protein kinase from rabbit skeletal muscle. *The Journal of biological chemistry* **243**, 3763-3765 (1968).

12. R. A. Screaton *et al.*, The CREB coactivator TORC2 functions as a calcium- and cAMP-sensitive coincidence detector. *Cell* **119**, 61-74 (2004).

13. N. E. Bruno *et al.*, Creb coactivators direct anabolic responses and enhance performance of skeletal muscle. *The EMBO journal* **33**, 1027-1043 (2014).

14. T. H. Ch'ng *et al.*, Activity-dependent transport of the transcriptional coactivator CRTC1 from synapse to nucleus. *Cell* **150**, 207-221 (2012).

15. P. S. Baxter, M. A. Martel, A. McMahon, P. C. Kind, G. E. Hardingham, Pituitary adenylate cyclase-activating peptide induces long-lasting neuroprotection through the induction of activity-dependent signaling via the cyclic AMP response element-binding protein-regulated transcription co-activator 1. *J Neurochem* **118**, 365-378 (2011).

16. M. Taub, S. Garimella, D. Kim, T. Rajkhowa, F. Cutuli, Renal proximal tubule Na,K-ATPase is controlled by CREB-regulated transcriptional coactivators as well as salt-inducible kinase 1. *Cell Signal* **27**, 2568-2578 (2015).

17.   Z. Wu *et al.*, Transducer of regulated CREB-binding proteins (TORCs) induce PGC-1alpha
transcription and mitochondrial biogenesis in muscle cells. *Proc Natl Acad Sci U S A* **103**,
14379-14384 (2006).

18.   N. D. Knuth *et al.*, Metabolic adaptation following massive weight loss is related to the
degree of energy imbalance and changes in circulating leptin. *Obesity (Silver Spring)* **22**,
2563-2569 (2014).

19.   J. C. Kerns *et al.*, Increased Physical Activity Associated with Less Weight Regain Six
Years After "The Biggest Loser" Competition. *Obesity (Silver Spring)* **25**, 1838-1843
(2017).

20.   R. J. Verheggen *et al.*, A systematic review and meta-analysis on the effects of exercise
training versus hypocaloric diet: distinct effects on body weight and visceral adipose tissue.
Obes Rev **17**, 664-690 (2016).

21.   R. Goldsmith *et al.*, Effects of experimental weight perturbation on skeletal muscle work
efficiency, fuel utilization, and biochemistry in human subjects. *Am J Physiol Regul Integr*
Comp Physiol **298**, R79-88 (2010).

22.   S. E. Calvo, K. R. Clauser, V. K. Mootha, MitoCarta2.0: an updated inventory of
mammalian mitochondrial proteins. *Nucleic Acids Res* **44**, D1251-1257 (2016).

23.   V. Giorgio *et al.*, Dimers of mitochondrial ATP synthase form the permeability transition
pore. *Proc Natl Acad Sci U S A* **110**, 5887-5892 (2013).

24.   A. S. Rambold, B. Kostelecky, N. Elia, J. Lippincott-Schwartz, Tubular network formation
protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc*
Natl Acad Sci U S A **108**, 10190-10195 (2011).

25.   V. Nasta, D. Suraci, S. Gourdoupis, S. Ciofi-Baffoni, L. Banci, A pathway for assembling
[4Fe-4S](2+) clusters in mitochondrial iron-sulfur protein biogenesis. *FEBS J*
10.1111/febs.15140 (2019).

26.   J. Liu *et al.*, Mitochondrial Dysfunction Launches Dexamethasone-Induced Skeletal
Muscle Atrophy via AMPK/FOXO3 Signaling. *Mol Pharm* **13**, 73-84 (2016).

27.   D. Ratman *et al.*, Chromatin recruitment of activated AMPK drives fasting response genes
co-controlled by GR and PPARalpha. *Nucleic Acids Res* **44**, 10539-10553 (2016).

28.   G. R. Steinberg *et al.*, Reduced glycogen availability is associated with increased
AMPKalpha2 activity, nuclear AMPKalpha2 protein abundance, and GLUT4 mRNA
expression in contracting human skeletal muscle. *Appl Physiol Nutr Metab* **31**, 302-312
(2006).

29.   B. Dasgupta *et al.*, The AMPK beta2 subunit is required for energy homeostasis during
metabolic stress. *Mol Cell Biol* **32**, 2837-2848 (2012).

30.   B. R. Barnes *et al.*, The 5'-AMP-activated protein kinase gamma3 isoform has a key role
in carbohydrate and lipid metabolism in glycolytic skeletal muscle. *The Journal of*
biological chemistry **279**, 38441-38447 (2004).

31.   C. S. Choi *et al.*, Paradoxical effects of increased expression of PGC-1alpha on muscle
mitochondrial function and insulin-stimulated muscle glucose metabolism. *Proc Natl Acad*
Sci U S A **105**, 19926-19931 (2008).

32.   M. Tadaishi *et al.*, Skeletal muscle-specific expression of PGC-1alpha-b, an exercise-
responsive isoform, increases exercise capacity and peak oxygen uptake. *PLoS One* **6**,
e28290 (2011).

33.   R. P. Shook *et al.*, Moderate cardiorespiratory fitness is positively associated with resting
metabolic rate in young adults. *Mayo Clin Proc* **89**, 763-771 (2014).

34.   M. Gilliat-Wimberly, M. M. Manore, K. Woolf, P. D. Swan, S. S. Carroll, Effects of
habitual physical activity on the resting metabolic rates and body compositions of women
aged 35 to 50 years. *J Am Diet Assoc* **101**, 1181-1188 (2001).
35.   J. C. Drake, R. J. Wilson, Z. Yan, Molecular mechanisms for mitochondrial adaptation to
exercise training in skeletal muscle. *FASEB J* **30**, 13-22 (2016).
36.   M. A. Pearen, G. E. Muscat, Minireview: Nuclear hormone receptor 4A signaling:
implications for metabolic disease. *Mol Endocrinol* **24**, 1891-1903 (2010).
37.   M. A. Maxwell *et al.*, Nur77 regulates lipolysis in skeletal muscle cells. Evidence for cross-
talk between the beta-adrenergic and an orphan nuclear hormone receptor pathway. *The*
Journal of biological chemistry **280**, 12573-12584 (2005).
38.   J. Meinken, G. Walker, C. R. Cooper, X. J. Min, MetazSecKB: the human and animal
secretome and subcellular proteome knowledgebase. *Database (Oxford)* **2015** (2015).
39.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009).
40.   Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
41.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
42.   S. Anders, W. Huber, Differential expression analysis for sequence count data. *Genome*
Biol **11**, R106 (2010).
43.   R. N. Margolis, R. M. Evans, B. W. O'Malley, N. A. Consortium, The Nuclear Receptor
Signaling Atlas: development of a functional atlas of nuclear receptors. *Mol Endocrinol*
19, 2433-2436 (2005).
44.   D. J. Pagliarini *et al.*, A mitochondrial protein compendium elucidates complex I disease
biology. *Cell* **134**, 112-123 (2008).
45.   Y. A. Medvedeva *et al.*, EpiFactors: a comprehensive database of human epigenetic factors
and complexes. *Database (Oxford)* **2015**, bav067 (2015).
46.   T. Kuo *et al.*, Genome-wide analysis of glucocorticoid receptor-binding sites in myotubes
identifies gene networks modulating insulin signaling. *Proc Natl Acad Sci U S A* **109**,
11160-11165 (2012).
586
587

FIGURES

Figure 1. Crtc2/Creb1 signaling enables greater loss of weight and adiposity during alternate day fasting (ADF).

(**A**) Summary of previous work showing that Crtc2 overexpression produced an exercised phenotype (13).

(**B**) 18-week old C57BL/6 male control (WT) and Crtc2 mice were treated with dox for 35 days and weighed. Data are shown as box-and-whisker plot, indicating the middle quartiles and range. $N = 6$ (WT) or 5 (Crtc2) mice per group.

(**C-F**) Mice from (**B**) were subject to alternate day fasting (ADF).

(**C**) Changes in the body weight during ADF.

(**D**) Weight regain after the ADF treatment and 3 weeks of ad libitum feeding. *significant in
Sidak's test for planned comparisons.

(**E**) Behaviors were assayed with Biodak monitoring system, including number of meals, bouts per
meal, meal duration, and kilocalories consumed over the 12-hr re-feeding period.

(**F**) Adiposity and lean mass determined by whole-body NMR measurements taken before and
after 8 cycles of ADF.

(**G**) 8-week old C57BL/6 male control and Crtc2 mice were treated with dox and placed on a high
fat diet (HFD). $N = 9$ (WT) or 7 (Crtc2) mice per group

Data are shown as mean ± SEM.

Figure 2. Electron transport chain gene expression and whole-body energy expenditure regulation by weight loss and fasting is reversed by Crtc2 expression.

(A–D) Gene expression profiles in control and Crtc2-transduced TA muscles of mice subjected to ADF relative to the ad libitum fed mice (columns 1–2), and effect of Crtc2 transduction relative to control during ADF (column 3). The profiles of all genes encoding **A)** mitochondrial proteins, including subunits of electron transport chain (ETC) complexes, **B)** subunits of the mitochondrial ATPase (complex V), **C)** ribosomal subunits, and **D)** highlighted mitochondrial proteins, are shown. ADF-regulated genes in control muscle appear

in **bold**. *Crtc2-regulated genes in mice subjected to ADF. *ADF-regulated genes in Crtc2-transduced muscle but not control. FC, fold change.

(**E**) Effects of Dex on lean body mass. Mice injected with vehicle or 10 mg/kg Dex were fasted overnight. Lean mass was measured by whole-body NMR before and after treatment. $N = 5$ mice per group.

(**F**) Diameter and mitochondrial potential of steroid-deprived primary myotubes treated for 2 days with vehicle or 100 nM Dex were compared by confocal high-content imaging. $N = 24$–27 images x 50–100 myotubes per image.

(**G**) Steroid-deprived C2C12 myotubes were stimulated with 100 nM Dex for 0–12 h. Levels of phosphorylated and total AMPK in whole lysates were compared by western blot.

(**H**) Diameter of steroid deprived C2C12 myotubes treated for 48 h with vehicle, 100 nM Dex, or 1 uM AICAR were compared by confocal high content imaging. $N = 24$–27 images x 50–100 myotubes per image.

(**I**) Expression levels of genes encoding AMPK subunits in mice described in **Figure 2**.

(**J**) VO_2 and VCO_2 were measured continuously (**Figure S2A**) for 72 hrs. in a CLAMS animal monitoring system. Shown are the 12-hr averages. $N = 8$ mice per group.

(**K**) Total energy expenditure was calculated from VO_2 and VCO_2 values using the Lusk equation.

Data are shown as mean ± SEM.

Figure 3. Crtc2 expression in skeletal muscle enhances mitochondrial biogenesis.

(**A**) Expression profiles of genes that regulate lipid metabolism in mice described in **Figure 2.**

(**B**) Histological analysis of succinate dehydrogenase in myofibers from gastrocnemius muscle sections after dox treatment. Scale bar = 500 μm.

(**C**) Gene induction calculated relative to WT mice. $N = 8$ mice per group.

(**D**) Gene expression in primary myocytes transduced with adenovirus expressing Crtc2 or GFP control 72 hr post-transduction.

(**E**) C2C12 myocytes were transduced with adenovirus GFP, Crtc2, or A-Creb genes and analyzed by confocal high content imaging. $N = 24$–27 images x 50–100 myotubes per image.

(**F**) Crtc2 expression plasmid was electroporated into the tibialis anterior muscle and GFP plasmid in the contralateral leg. After 7 days, DNA was extracted for assessment of mitochondrial DNA (mt-DNA) for Crtc2/GFP control. $N = 5$ mice per group.

649   **(G)** Oxidation of ^{3}H-Palmitic acid in the supernatant of primary myocytes transduced with control,

650   Crtc2, or Crtc3-expressing adenovirus.

651   * $p < 0.05$. Data are shown as mean + SEM.

Figure 4. Crtc2 transgenic mice maintain enhanced glucose disposal and insulin tolerance

(**A**) Expression profiles of genes that regulate carbohydrate metabolism in mice described in
Figure 2.

(**B**) Glucose tolerance test (GTT) after alternate day fasting. Right panel, area under the curve. *N*
= 10 mice per group

(**C**) Plasma insulin levels during GTT described in panel **B**.

(**D**) Insulin tolerance tests after 8 cycles of ADF. *N* = 10 mice per group

(**E**) Insulin-mediated uptake of 2-deoxy-D-glucose in primary myocytes transduced with
adenovirus expressing Crtc2 or GFP control. *N* = 10 per group. Student's t-test.

(**F**) Crtc2 mice were treated +/- dox for 1 week and the gene induction compared by qPCR relative

to the no dox groups. N= 10 per group.

(**G**) Gene expression in primary myocytes transduced with adenovirus expressing Crtc2 or GFP

control were compared by qPCR. *N*=4 per group.

(**H**) Western blot of whole lysates from primary myocytes transduced with the indicated plasmids.

(**I–J**) Myocytes stably expressing GFP, Crtc3 (lanes 2 and 3), dominant negative Creb (A-CREB),

or dominant negative Crtc2 (dn-Crtc) were compared by western blots.

(**K**) WT or Crtc2 mice were injected twice daily with 0.5 mg/kg leptin and weight was monitored.

N=8 mice per group.

Data are mean ± SEM.

Figure 5. Hormonal and neural modulation of energy expenditure in skeletal muscle

(**A**) Fasting-induced regulation of mitochondrial dynamics and mitophagy.

(**B**) Exercise-induced regulation of mitochondrial dynamics and mitochondrial biogenesis, which can be constitutively activated by overexpression of Crtc2 in skeletal muscle.

bAR, b-adrenergic receptor; GR, glucocorticoid receptor; HPA, hypothalamic-pituitary-adrenal axis; HPT, hypothalamic-pituitary-thyroid axis; InsR, insulin receptor; nAchR, nicotinic acetylcholine receptor; SNS, sympathetic nervous system; TH, thyroid hormone