Activation of the Crtc2/Creb1 transcriptional network in skeletal muscle enhances weight

loss during intermittent fasting

Nelson E. Bruno[1]*, Jerome C. Nwachukwu[1], Sathish Srinivasan[1], Richard Hawkins, David Sturgill[3], Gordon L. Hager[3], Stephen Hurst[4], Shey-Shing Sheu[4], Michael D. Conkright[5], Kendall W. Nettles[1]*

[1]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

[2]Department of Cancer Biology, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

[3]Laboratory of Receptor Biology and Gene Expression, National Cancer Institute, NIH, Bethesda, MD 20892, USA

[4]Center for Translational Medicine, Department of Medicine, Thomas Jefferson University, Philadelphia, PA, 19107, USA.

[5]Department of Shared Services, The Scripps Research Institute, 130 Scripps Way, Jupiter FL 33458, USA

Correspondence: Nelson E Bruno, nbruno@scripps.edu; Kendall Nettles, knettles@scripps.edu

Competing interest statement: The authors declare no competing interests.

Abstract

Exercise is a behavior modification indispensable for long-term weight loss. Exercise activates the Creb-Regulated Transcriptional Coactivator (Crtc) family of transcriptional coregulators to drive Creb1-mediated anabolic transcriptional programs in skeletal muscle. Here, we show that induced overexpression of a skeletal muscle specific Crtc2 transgene in aged mice leads to greater weight loss during alternate day fasting, and selective loss of fat rather than lean mass. Transcriptional profiling revealed that fasting and weight loss downregulated most of the mitochondrial electron transport genes and other regulators of mitochondrial function that were substantially reversed in the Crtc2 mice, which maintained higher energy expenditure during fasting. The Crtc2 mice displayed greater mitochondrial activity, metabolic flux capacity for both carbohydrates and fats, improved glucose tolerance and insulin sensitivity, and increased oxidative capacity before the fast, suggesting muscle-intrinsic mechanisms in support of improved weight loss. This work reveals that Crtc2/Creb1-mediated signaling coordinates metabolic adaptations in skeletal muscle that explain how Crtc2/Creb contribute to the effects of exercise on metabolism and weight loss.

Introduction

Long-term dieting is usually not effective in maintaining a lower body weight due to increased

hunger and a slower metabolic rate (1). In humans, loss of >10% of body weight also increases

metabolic efficiency, such that ~500 less calories per day are required to maintain the same energy

expenditure (2). Weight loss reduces the metabolic rate due to a number of changes including

increased circulating leptin and decreased thyroid hormone and sympathetic nervous system (SNS)

activity (2-4). People who maintain long-term weight loss almost invariably exercise (5), and both

preclinical and clinical data suggest that this is through raising metabolic rate (6-8), decreased

energy efficiency (9), and increased energy expenditure from the increased physical activity (10).

Creb1-regulated transcription is stimulated by GPCRs that activate adenylyl cyclases to

convert ATP to the second messenger cAMP, which in turn liberates the catalytic subunit of protein

kinase A (PKA) (11). Crtc1–3 family members are primary coactivators for Creb1 that are also

highly conserved in the cAMP transcriptional response. Crtc1–3 are rendered cytoplasmic and

inactive by phosphorylation (12). In contrast, they are fully activated through synergistic

dephosphorylation by phosphatases downstream of PKA and calcium flux in polarizable tissues

including the skeletal muscle (13), hippocampus (14), cortex (15), islet cells (12), and the renal

proximal tubule (16).

To study the effects of Crtc2/Creb1 activation in adult skeletal muscle, we generated

doxycycline (dox)-inducible transgenic mice that show skeletal muscle-specific overexpression of

Crtc2 under the control of a tetracycline response element (13). C57BL/6-Tg(TRE-

Crtc2tm)1Mdc/j mice express the FLAG-Crtc2 S171A/S275A/K628R mutant attenuating the

phosphorylation-driven negative feedback loop that exports Crtc2 from the nucleus to the

cytoplasm. These mice were crossed with the C57BL/6-Tg 1256[3Emut] MCK-rtTA mice that

express the reverse tetracycline transactivator (rtTA) specifically in skeletal muscle to control spatial expression limited to the skeletal muscle compartment (Grill *et al*, 2003). We identified a double transgenic line with expression that was tightly regulated by dox and highly selective for skeletal muscle, that we refer to as Crtc2 mice. These mice showed an exercised phenotype including induction of *Ppargc1a*/PGC-1α and other anabolic genes, increased glycogen and intramuscular triglycerides (IMTGs), increased muscle cross sectional area, and a remarkable increase in high intensity exercise performance after less than two weeks of transgene induction (13). Activation of Crtc2/Creb1 after exercise has now been demonstrated a number of times in humans, but may depend on the type, duration, and intensity of exercise to upregulate anabolic genes (17-20).

Given the primary role of exercise adaptations in regulating whole-body metabolism, we examined here the role of the Crtc2/Creb1 transcriptional network in skeletal muscle metabolism during alternate day fasting (ADF), mimicking a common dieting scheme in humans. Activation of Creb1-regulated transcriptional responses during fasting enabled maintenance of a lower body weight due to the stimulation of higher energy expenditure and other metabolic changes, enabling selective loss of fat mass and retention of lean mass, as observed in humans who exercise (21, 22). Transcriptional profiling revealed that ADF reduced the expression of most mitochondrial respiratory chain subunits and lowered energy expenditure, which was prevented by modest Crtc2 overexpression. Crtc2 mice displayed the metabolic hallmarks of an exercised phenotype, including enhanced mitochondrial activity, improved lipid and carbohydrate flux capacity and storage, and insulin sensitization. These data support a model where lower metabolic activity during fasting and weight loss contributes to weight regain, which can be reversed by exercise-induced activation of Crtc2/Creb1-regulated transcriptional programs.

Results

Crtc2 overexpression in skeletal muscle facilitates weight loss

We studied aged Crtc2 (rtTA/Crtc2^{tm}) or WT (rtTA/WT) mice, which in the C57BL/6 background naturally gain fat mass as they age (18 weeks, 30–39 g, **Figure 1A–B**). During dox treatment, Crtc2 mice gained weight similarly to control WT mice (SI Appendix, **Figure S1A**). On the ADF protocol, three separate cohorts of Crtc2 mice displayed significantly greater and sustained weight loss than WT mice, with Crtc2 mice losing about twice as much weight in each case, 9-14% of body weight (**Figure 1C, Figure S1B**).

After 3 weeks of ad libitum refeeding, the Crtc2 mice re-gained more weight to catch up with the WT mice (**Figure 1D**), which may be due to the anabolic effects of higher insulin sensitivity, described below. We also completed behavioral profiling studies and found that locomotion was similarly reduced in both groups during the refeeding period (**Figure S1C**). During the refeeding period the number of meals doubled in both groups, while meal duration and number of bouts per meal declined (**Figure 1E**). We observed a reduction in kilocalories consumed in the Crtc2 relative to the WT group, which planned comparisons demonstrated was significant only during the second ADF cycle, when the WT mice increased consumption at a faster rate than the Crtc2 mice (**Figure 1E**). These differences in caloric consumption are consistent with the differences in body weight between cohorts and disappear after normalization for body weight. Control mice lost more lean mass, while Crtc2 mice conserved lean mass and burned the excess fat accumulated during aging (**Figure 1F**). To verify that Crtc2 helps maintain lower body weight in a different model, we placed mice on a high fat diet, and found that Crtc2 mice also maintained lower body weight than WT mice in this context (**Figure 1G**). Thus, the activation of Crtc2/Creb1

signaling in skeletal muscle significantly modulated body composition and weight loss.

The transcriptional response to fasting and weight loss and its regulation by Crtc2

To understand the skeletal muscle intrinsic effects of Crtc2 on dieting and weight loss, we performed

transcriptional profiling of mice tibialis anterior (TA) muscles electroporated with control and Crtc2

expression plasmids in contralateral legs. One cohort of electroporated mice was subject to 3 cycles

of ADF, while the other cohort was fed ad libitum. TA muscle tissues were collected from both

cohorts at the end of the third 24-hr fast. The transduction increased *Crtc2* mRNA levels by ~2.5-

fold across groups (**Figure 2A**). In response to ADF, we identified 1,880 differentially expressed

genes (DEGs) in the control-transduced muscle and 3,197 DEGs in Crtc2-transduced muscle, with

1,380 genes in common between the two groups (**Figure 2B**). Analysis of canonical signaling

pathways and gene ontology (GO) annotations revealed that ADF modulated the transcriptional

programs for lipid and carbohydrate metabolism, fatty acid beta-oxidation, and insulin signaling

(**Figure 2C-D**). ADF also regulated transcriptional programs that control protein balance, including

autophagy and ubiquitin-dependent proteasomal degradation which were upregulated, RNA

splicing, and transcription (**Figure 2C-D**).

To understand other transcriptional control mechanisms underlying the skeletal muscle

response to fasting and weight loss, we examined genes annotated as transcription factors or

transcriptional coregulators and found over 200 DEGs regulated by ADF. These include the

upregulation of a cohort of nuclear receptors that control energy expenditure, lipid metabolism and

mitochondrial function (*Thra*, *Nr4a3, PPARs, RXRα, RORα*) (**Figure 2E**) (23). Mice lacking *Nr2c2*

(TR4) display abnormal mitochondrial composition and function (24), but its role in the fasting and

weight loss is not known. As expected by their roles in coordinating the skeletal muscle atrophy, we

observed upregulation of the *Foxo3* and *Foxo4* transcription factor genes following ADF (**Figure 2F**). We also observed upregulation of the *Foxn3* gene which regulates hepatic glucose homeostasis, but whose role in skeletal muscle is unknown (25). We identified a number of transcription factors implicated in myoblast differentiation and regeneration (*Foxk1*, *Nfix*, *Klf9*, *Vgll2*, *Peg3*, *Sox7*), suggesting they may play additional roles in the skeletal muscle response to fasting and weight loss.

The 85 transcriptional coregulators with ADF-dependent changes in expression, include histone acetyltransferases, corepressors, chromodomain proteins, protein methyltransferases and demethylases (**Figure 2G**). For example, Carm1 is a methyltransferase required for the effects of Foxo3 on atrophy (26), and the histone deacetylases Hdac2, Hdac3, and Hdac7, have broad roles in skeletal muscle physiology (27, 28). *Prkcb*/PKCβ was upregulated by ADF, and in addition to its role as a transcriptional coregulator, PKCβ phosphorylates and inhibits the insulin receptor to inhibit its activity (29), and activates the generator of mitochondrial ROS, p66shc (30).

Overexpression of Crtc2 in the context of ADF altered expression of 42 transcription factors and 15 coregulators. These include the nuclear receptor, ERRα, which regulates mitochondrial biogenesis, and RXRβ. RXRs are the obligate heterodimer partner for TR and PPARs. *E2F6*, which causes cardiac hypertrophy upon overexpression (31), was upregulated by ADF only in the Crtc2 condition. However, 74% of Crtc2-dependent DEGs that encode transcription factors were inhibited, including myogenin, a primary driver of atrophy, and *Irf1*, *Irf7*, *Notch2*, and *Stat1,* which control immune and inflammatory responses. Crtc2 also inhibited many other genes implicated in the inflammatory response including the nuclear factor-kappa B (NF-κB) signaling pathway. These findings demonstrate that ADF and its associated weight loss profoundly changes the expression of transcriptional regulators in skeletal muscle, effects significantly modulated by Crtc2.

Crtc2 reverses fasting- and weight loss-induced effects on mitochondria and energy expenditure

In addition to lower metabolic rate, fasting and weight loss render humans and other mammals more energetically efficient, such that the same amount of work is performed with less ATP production, suggesting a remodeling of glycolytic or oxidative capacity (2, 32). ADF modulated the expression of >200 genes whose proteins localize in mitochondria according to MitoCarta (33) (**Figure 3A**). ADF downregulated most of the genes that encode the electron transport chain (complexes I–IV), the ATP synthase (complex V), the mitoribosome, and the cytosolic ribosome (**Figure 3A-C**, **Figure S3A**), but significantly upregulated genes such as *Sdha*, *Uqcrc1*, and *Atp5a1* (**Figure 3B**, **Figure S3A**).

We observed a significant ADF-induced expression of genes encoding ROS scavenger proteins such as thioredoxin (*Txnrd1*) and catalase (*Cat*), and genes regulating mitochondrial stability*, Tomm7, Timm9,* and the AAA protease *Spg,7* known to be active during the removal of damaged mitochondrial proteins (**Figure 3D**). Among the ATP synthase genes, there was a more pronounced loss of gene expression in the lateral stalk genes (**Figure 3B**), which may decrease the conductance of mitochondrial permeability transition pore (mPTP) and thus protect mitochondria from detrimental swelling and oxidative injury (34). ADF also induced increases in mitochondrial fusion (*Mfn2* and *Opa1)*, and decreased fission (*Fis)* gene expression, consistent with the protective effects of fusion during nutrient deprivation (35). These adaptations could be advantageous for weight loss as it prepares the mitochondria for the reduced metabolic rate and increased reliance on lipid metabolism during fasting, without causing permanent damage to the mitochondrial pool within the cell.

In mice subjected to ADF, Crtc2 transduction altered expression of 80 genes whose proteins

localize in the mitochondria. Crtc2 expression strongly attenuated the ADF-dependent repression of

electron transport chain genes and the mitoribosome (**Figure 3A–C**, **Figure S3A**), and generally

upregulated the other genes encoding proteins that localize to mitochondria. In addition, Crtc2

enhanced expression of ribosomal genes that facilitate mitoribosome assembly, and genes such as

Spg7, Pink1, RhoT2, and *Chchd3* that control the quality of mitochondrial proteins, but blocked

ADF-dependent downregulation of *Minos1*, a central component of the mitochondrial inner

membrane organizing system (**Figure 3D**). Crtc2 also upregulated expression of *Tefm*, the

mitochondrial transcriptional elongation factor, and *Bola3*, which participates in electron transport

chain assembly (36).

To understand how ADF affects mitochondria, we noted that glucocorticoids (GCs) are

one of the primary drivers of catabolic changes during fasting, and in response to ADF and Crtc2

overexpression, we observed over a thousand DEGs that were annotated as "glucocorticoid

response" genes (**Figure 2C**). In the context of an overnight fast, treatment with Dex more than

doubled loss of lean mass (**Figure 3E**). Using high-content imaging, we observed that Dex reduced

the diameter and mitochondrial potential of nutrient-deprived primary mouse skeletal myotubes

(**Figure 3F**). GCs can also activate the master energy sensor, AMP-activated protein kinase

(AMPK) (37), which can act as a glucocorticoid receptor (GR) coactivator (38). AMPK is

activated by a lower ATP/ADP ratio during fasting to inhibit anabolism and stimulate catabolic

processes such as autophagy and mitophagy. In primary myotubes, we observed a Dex-dependent

activating phosphorylation of AMPK that increased over time, consistent with an underlying

transcriptional mechanism (**Figure 3G**). Activation of AMPK with both Dex and the AMP analog,

AICAR, had an additive detrimental effect on myotube diameter (**Figure 4H**), suggesting that GR

collaborates with AMPK to drive skeletal muscle atrophy.

We also observed that ADF modulated the expression of genes encoding three of the

AMPK subunits. Changes in subunit composition are known to alter the regulation, substrate

preferences, and cellular location of AMPK. The α2 and β2 AMPK isoforms were significantly

upregulated by ADF (SI Appendix, **Figure 3I**). The α2 subunit is activated by low glycogen and

mediating nuclear translocation (39), while the β2 subunit is required for maintenance of muscle

ATP levels during fasting (40). In contrast, ADF downregulated the γ3 AMPK isoform (**Figure**

3I), which is required for glycogen synthesis (41), a process inhibited during fasting. Thus, a

combination of GR and AMPK signaling drives skeletal muscle atrophy during nutrient

deprivation and a loss of mitochondrial potential.

We further examined whether ADF- and Crtc2-induced changes in mitochondria—the

powerhouse of cellular respiration—are associated with changes in global energy expenditure. As

expected, ad libitum oxygen consumption cycled with activity in the mice, which increased at

night and decreased during the day (**Figure 3J, Figure S3B**). Fasting reduced oxygen consumption

at every cycle, and this effect was more pronounced during the day as the fast was extended

(**Figure S3B**). However, Crtc2 mice consistently showed higher oxygen consumption,

independent of fasting (**Figure 2J**).

When fed ad libitum, Crtc2 and control mice display similar rates of energy expenditure

(EE) (**Figure 3JK**). We calculated EE from indirect calorimetry measurements of VO_2 and VCO_2

and performed ANCOVA analysis using body weight and NMR data as covariates to adjust the

EE to total body weight and body composition (**Figure S3C**). While fasting, the EE of WT mice

dropped significantly. In contrast, during fasting EE in Crtc2 mice did not drop below the ad

libitum levels (**Figure 3I**). Respiratory quotient was also reduced during fasting, indicating greater

utilization of fat for fuel, but the differences between WT and Crtc2 mice were not significant

(**Figure S3D–E**). These data suggest potential mechanisms to explain how physical activity—

through Crtc2/Creb1-regulated transcriptional programs—can help maintain a higher metabolic

rate and a lower body weight by upregulating expression of mitochondrial respiratory chain subunits

and modulating expression of other mitochondria-localized gene products, consistent with the well-

known beneficial effects of exercise on mitochondrial biogenesis and quality control.

Crtc2 stimulates mitochondrial activity and improves oxidative capacity

Prolonged fasting promotes lipolysis, insulin resistance, and mobilization of fatty acids to skeletal

muscle through the combined effects of glucagon and glucocorticoids on adipocytes. These actions

augment the rate of fatty acid oxidation by mitochondria and the ectopic accumulation of IMTGs.

As expected, ADF upregulated a panoply of genes involved in the transport, oxidation, and

esterification of fatty acids. Key among these are: *Ppard* and *Ppara,* the chief regulators of fat

metabolism in muscle; *Cd36* that regulates plasma membrane fatty acid transport and utilization

to spare glucose; *Ucp3,* which is implicated in β-oxidation as well as protection from triglyceride

accumulation in muscle (42); *Pdk2* and *Pdk4,* which promote fatty acid oxidation over glycolysis

(43); and multiple enzymes involved in Acyl-CoA metabolism (**Figure 4A**). *Acox1* is a

peroxisomal Acyl-CoA oxidase, while *Acad10*, *Acadl*, *Acadm* are members of the acyl-coenzyme

A dehydrogenase family that are critical for effective beta-oxidation in the mitochondria. *Acsl1*,

Acot1, *Acot2*, and *Acot4* regulate the generation and hydrolysis of Acyl-CoA. Differential

expression of *Dgat2*, *Daglb*, *Plin5* and other genes that regulate lipid storage and accumulation

highlights their importance in lipid homeostasis during fasting. Notably, the lipoprotein lipase

(*Lpl*) gene, which controls a rate limiting step in lipoprotein metabolism, was upregulated by ADF

more robustly in Crtc2-transduced legs than in control legs, suggesting an improved post prandial

metabolism of triglyceride-rich lipid particles. *Sirt4*, a mitochondrial sirtuin that regulates the rate

of fatty acid oxidation in muscle, was also substantially suppressed during fasting but significantly

less so in the Crtc2-overexpressing legs which may favor the higher lipid accumulation that we

previously described in muscle overexpressing Crtc2 (13).

Crtc2 mice displayed higher oxygen consumption before the fast; therefore, we examined

them for improved mitochondrial function. After one week of dox treatment, Crtc2 mice showed

increased mitochondrial succinate dehydrogenase activity in the gastrocnemius muscle (**Figure 4B,**

Figure S4A–B) and increased expression of *Cytochrome C (Cycs)* (**Figure 4C**). In addition to the

previously reported upregulation of *Ppargc1a/*PGC-1α (13), Crtc2 mice showed increased

expression of *Nr4a3/*NOR-1, a nuclear receptor family transcriptional factor (**Figure 4C**). PGC-

1α and NOR-1 induce mitochondrial biogenesis and increase oxidative capacity (44, 45). Crtc2

also upregulated the *Dgat1* gene (**Figure 4C**), which is a direct Creb1 target gene (13), the product

of which catalyzes a rate-limiting step that increases IMTG stores. These mitochondria-related

genes were also upregulated in primary myotubes overexpressing Crtc2, indicating a skeletal

muscle-intrinsic effect of Crtc2 that mimics the effects of exercise (**Figure 4D**). To show that

Creb1 is directly regulating atrophy, we transduced primary myocytes with the dominant negative

A-Creb, which reduced myotube diameter (**Figure 4E**). We also observed a doubling of

mitochondrial DNA after in vivo electroporation of Crtc2 (**Figure 4F**). The increased

mitochondrial mass was associated with increased oxidative capacity, as myotubes overexpressing

Crtc2 or Crtc3 oxidized more palmitate than control myotubes (**Figure 4G**), as previously reported

(46).

Crtc2 improved glucose tolerance and carbohydrate metabolism before and during weight loss

During fasting, the rise of non-esterified fatty acid levels in plasma leads to a suppression of insulin

signaling and glucose metabolism in skeletal muscle, where glycogen levels can drop by ~40% (47).

After 3 cycles of ADF, we found a significant activation of genes involved in glycogen metabolic

processes, including synthesis (*Gys1*, *Fbp2*, *Ugp2*, *Epm2a*, and *Ppp1cb*), branching (*Gbe1*), and

breakdown (*Gaa*, *Agl*, *Gsk3b*) of glycogen, perhaps in preparation for feeding (**Figure 5A**).

Likewise, a number of genes involved in different aspects of insulin signaling were also activated

in both Crtc2-transduced and control muscle, including *Irs2*, *Sorbs1,* and *Insr*. Many genes

necessary for glycolysis and fructose metabolism were suppressed by ADF, including *Pfkfb1,*

Pfkfb2, and *Fuk*. *Ogt*, an important inducer of insulin resistance in skeletal muscle via the inhibition

of AKT phosphorylation was also downregulated.

To probe for Crtc2-dependent changes in carbohydrate metabolism, we performed glucose

tolerance tests comparing Crtc2 and WT mice following ADF or normal diet. The Crtc2 mice

displayed significantly improved glucose disposal after ADF, including a dramatic improvement in

their ability to reduce glycemia (**Figure 5B**), and required a lower insulin excursion (**Figure 5C**).

Crtc2 overexpression also improved insulin tolerance after fasting and weight loss (**Figure 5D**).

These effects were also apparent after 1 week of dox treatment in Crtc2 mice fed ad libitum (**Figure**

S5A–C). In primary myotubes, expression of Crtc2 was sufficient to significantly increase the uptake

of 2DG-glucose (**Figure 5E**).

Crtc2 mice and Crtc2/3-overexpressing myotubes upregulated a number of carbohydrate

signaling genes in normal fed conditions at both the RNA and protein levels (**Figure 5F–J**),

including the glucose transporter, *Glut4*, whose insulin-dependent translocation to the membrane

drives glucose uptake into skeletal muscle, accounting for 80% of glucose disposal. The *Nr4a1* gene

encodes Nur77, a nuclear receptor family transcription factor that regulates transcriptional programs

for carbohydrate metabolism, lipolysis, and mitochondrial biogenesis (48). *Irs2*, *Akt1*, and *Akt2*

encode core components of the insulin receptor signaling pathway that can coordinate anabolic

signaling and improve carbohydrate metabolism. We also validated a significant inhibition of

myogenin expression in the normal fed Crtc2 mice, a primary driver of muscle atrophy during

fasting. To validate on target mechanism of action, we generated stable cell lines overexpressing

Crtc3, dominant negative A-Creb, or dominant negative Crtc2 expressing only the Creb binding

domain (12). These results demonstrate that A-Creb and dn-Crtc3 blocked basal expression of Pgc-

1α, Irs2, and Akt2, and prevented phosphorylation of Akt1/2 at Thr308 (**Figure 5I–J**).

To test for pre-fasting changes in the central control of feeding and energy expenditure, we

completed a leptin infusion study, as leptin secretion from adipocytes both suppresses hunger and

stimulates the SNS. The WT and Crtc2 mice displayed identical patterns of weight loss following

leptin infusion (**Figure 5K**), and there were also no differences in plasma levels of feeding hormones,

cholesterol, or triglycerides (**Figure S5D–E**). Lastly, we considered that Crtc2 might be regulating

secreted proteins that modulate the response to dieting, using the Metazoan Secretome

Knowledgebase (49). Among 2,500 genes encoding likely secreted proteins, 171 DEGs were

regulated by Crtc2 during ADF. Using the Reactome.org peer-reviewed pathway database, we

detected significant enrichment of pathways regulating extracellular matrix, inter-cell signaling, and

immune function, among others (**Figure S5F**), but no changes in myokines that would drive weight

loss. However, Crtc2 signaling in skeletal muscle would be expected to have effects on other tissues

just from changes in metabolic flux, as Crtc2 mice displayed pre-fast increases in skeletal muscle

oxidative flux capacity.

Discussion

Skeletal muscle can be viewed as having semi-autonomous flux engines responding to a variety of

homeostatic stresses including nutrient flux, hormones, and exercise. We demonstrated here, and in

our previous study, that Crtc2/Creb1 signaling in skeletal muscle globally coordinates many of the

anabolic and metabolic benefits of exercise through the combined regulation of a number of these

flux engines. Our transcriptional and metabolic profiling revealed a profound effect of Crtc2 activity

that prevented the ADF-dependent loss of mitochondrial electron transport chain gene expression

and maintained energy expenditure (**Figures 2–3**). Pre-fast, Crtc2 stimulated mitochondrial

biogenesis and activity, upregulated β-oxidation genes, increased lipid flux capacity, and improved

insulin sensitivity and glucose disposal before and during fasting (**Figures 4–5**). While altered

mitochondrial function is the most likely mechanism explaining increased oxygen consumption and

maintenance of energy expenditure during fast, we did not examine whether Crtc2 regulated aerobic

glycolysis. While PGC-1α is sufficient to upregulate mitochondrial biogenesis, it does not replicate

the effects of exercise on energy expenditure Additionally, the previously reported increases in

skeletal muscle storage of glycogen and triglycerides also likely contributed to the early response,

perhaps explaining the initially slower rate of increased food consumption in the Crtc2 mice, while

the WT mice rapidly increased food intake to maintain weight. (**Figure 1E**). Changes in muscle

metabolic capacity may favor energy mobilization from tissues other than skeletal muscle. However,

when we looked at the obvious signaling pathways recruited by fasting, like leptin signaling, we

found no difference in leptin sensitivity.

It is not clear whether high- or moderate-intensity exercise is better for weight loss or insulin

sensitivity (50, 51). However, there is a clear effect of exercise dose on the amount of fat and weight

loss and cardiorespiratory health (52, 53), highlighting the importance of regular exercise for human

health. Exercise also improves the utilization of fat for fuel, which we observed here. A recent NIH

working group on obesity and exercise concluded that exercise intervention trials may have failed to

show an effect of exercise on long-term weight loss due to compliance issues, but it is very clear that

people who effectively maintain long-term weight loss exercise regularly (5, 54). These

transcriptional and metabolic adaptations may also explain how exercise increases energy

expenditure beyond that required during and immediately after completion of the exercise and alter

substrate utilization. Exercised calorie-restricted (CR) mice respond to refeeding by preferentially

oxidizing lipids, while using additional energy to refill fat stores by *de novo* lipogenesis (55). In

contrast, sedentary CR mice store fat more efficiently for rapid weight regain (55). This parallels the

effects of Crtc2 overexpression on the preferential loss of fat mass during ADF, similar to the effects

seen in humans who exercise during weight loss (21).

In skeletal muscle, adenylate cyclase/cAMP signaling is activated by several exercise-

regulated receptors, including GPCRs such as the β-adrenergic receptor (βAR) which is activated

by the SNS, the cholinergic receptor activated at the neuromuscular junction to induce calcium-

mediated muscle contraction, and receptors for adenosine and calcitonin gene-related peptide

(CGRP), which cause vasodilation during exercise (56, 57). We previously demonstrated in mice

that exercise-induced expression of the Creb1 target genes, *Ppargc1a*/PGC-1α and *Sik1*, and that

this transcriptional output was reversed by βAR blocker administration (13).

This work reveals mechanisms through which diet and exercise regulate substrate mobilization

and metabolism and highlights how these lifestyle modifications can alter mitochondrial function

and energy expenditure. Reduced energy expenditure and increased hunger during fasting are partly

mediated by a lowering of the satiety hormone leptin, which acts centrally to stimulate the SNS (2-

4). Leptin also secreted basally from adipose tissue in proportion to fat mass, so it is chronically

lower during fasting, which lowers sympathetic tone and energy expenditure, along with reductions

in circulating thyroid hormone and stimulating the HPA axis (**Figure 6A**). Within skeletal muscle,

increased glucocorticoids and AMPK activity are sufficient to reduce mitochondrial potential

(**Figure 3)**, which along with likely contributions from fasting-induced lowering of thyroid hormone,

βAR, and insulin receptor activity, induce mitochondrial dynamics and mitophagy (**Figure 6A**).

These energetic effects are reversed by exercise (**Figure 6B**), which we now show can be

recapitulated by enforced expression of Crtc2/Creb1 and upregulation of mitochondrial biogenesis

master regulatory genes, *Nr4a1*, *Nr4a3*, *Essrra/*ERRα, and *Ppargc1a/*PGC-1α (**Figure 5**).

overexpression of PGC-1α increases mitochondrial density and exercise performance, but without

the increase in metabolic rate that can occur from exercise training (58-61), highlighting the

coordination with other exercised-induced signaling pathways. The combination of βAR and Ca^{2+}

signaling are required to maximally stimulate nuclear translocation of Crtc2 in skeletal muscle (13).

In this model, increased GC and AMPK activity during exercise promotes mitochondrial dynamics,

which is then coupled with mitochondrial biogenesis to increase mitochondrial potential and mass,

in coordination with other metabolic changes (**Figures 4–5**). Overexpression of Crtc2 circumvented

the neuronal activation of Creb1, as Crtc2 mice had most of the hallmarks of exercise adaptations

before fasting and weight loss (**Figures 4–5, Figure S5B–C**) (13).

376
377

Acknowledgements. N.E.B. was supported by the BallenIsles Men's Golf Association. J.C.N. was

supported by the Frenchman's Creek Women for Cancer Research.

Author contributions Conceptualization, N.E.B., M.D.C., K.W.N.; Methodology, N.E.B., M.D.C.;

Investigation, N.E.B., J.C.N., S.S., R.H., D.S.; Writing – Original Draft, K.W.N., N.E.B., M.D.C.,

S.H., S.S.S.; Writing – Revision, K.W.N, J.C.N. ,N.E.B.   Supervision, M.D.C, G.L.H., K.W.N.,

S.S.S., N.E.B.

References

1. C. L. Melby, H. L. Paris, R. M. Foright, J. Peth, Attenuating the Biologic Drive for Weight Regain Following Weight Loss: Must What Goes Down Always Go Back Up? *Nutrients* **9** (2017).

2. M. Rosenbaum *et al.*, Effects of experimental weight perturbation on skeletal muscle work efficiency in human subjects. *Am J Physiol Regul Integr Comp Physiol* **285**, R183-192 (2003).

3. L. J. Arone, R. Mackintosh, M. Rosenbaum, R. L. Leibel, J. Hirsch, Autonomic nervous system activity in weight gain and weight loss. *Am J Physiol* **269**, R222-225 (1995).

4. M. Rosenbaum *et al.*, Low-dose leptin reverses skeletal muscle, autonomic, and neuroendocrine adaptations to maintenance of reduced weight. *J Clin Invest* **115**, 3579-3586 (2005).

5. R. M. Foright *et al.*, Is regular exercise an effective strategy for weight loss maintenance? *Physiol Behav* **188**, 86-93 (2018).

6. P. S. MacLean *et al.*, Regular exercise attenuates the metabolic drive to regain weight after long-term weight loss. *Am J Physiol Regul Integr Comp Physiol* **297**, R793-802 (2009).

7. Y. Turk *et al.*, High intensity training in obesity: a Meta-analysis. *Obesity science & practice* **3**, 258-271 (2017).

8. A. Mourier *et al.*, Mobilization of visceral adipose tissue related to the improvement in insulin sensitivity in response to physical training in NIDDM. Effects of branched-chain amino acid supplements. *Diabetes care* **20**, 385-391 (1997).

9. M. Rosenbaum *et al.*, Resistance Training Reduces Skeletal Muscle Work Efficiency in Weight-Reduced and Non-Weight-Reduced Subjects. *Obesity (Silver Spring)* **26**, 1576-1583 (2018).

10. D. M. Ostendorf *et al.*, Physical Activity Energy Expenditure and Total Daily Energy Expenditure in Successful Weight Loss Maintainers. *Obesity (Silver Spring)* **27**, 496-504 (2019).

11. D. A. Walsh, J. P. Perkins, E. G. Krebs, An adenosine 3',5'-monophosphate-dependant protein kinase from rabbit skeletal muscle. *The Journal of biological chemistry* **243**, 3763-3765 (1968).

12. R. A. Screaton *et al.*, The CREB coactivator TORC2 functions as a calcium- and cAMP-sensitive coincidence detector. *Cell* **119**, 61-74 (2004).

13. N. E. Bruno *et al.*, Creb coactivators direct anabolic responses and enhance performance of skeletal muscle. *The EMBO journal* **33**, 1027-1043 (2014).

14. T. H. Ch'ng *et al.*, Activity-dependent transport of the transcriptional coactivator CRTC1 from synapse to nucleus. *Cell* **150**, 207-221 (2012).

15. P. S. Baxter, M. A. Martel, A. McMahon, P. C. Kind, G. E. Hardingham, Pituitary adenylate cyclase-activating peptide induces long-lasting neuroprotection through the induction of activity-dependent signaling via the cyclic AMP response element-binding protein-regulated transcription co-activator 1. *J Neurochem* **118**, 365-378 (2011).

16. M. Taub, S. Garimella, D. Kim, T. Rajkhowa, F. Cutuli, Renal proximal tubule Na,K-ATPase is controlled by CREB-regulated transcriptional coactivators as well as salt-inducible kinase 1. *Cell Signal* **27**, 2568-2578 (2015).

17.   D. V. Popov, E. A. Lysenko, P. A. Makhnovskii, N. S. Kurochkina, O. L. Vinogradova,
Regulation of PPARGC1A gene expression in trained and untrained human skeletal
muscle. *Physiological reports* **5** (2017).

18.   D. V. Popov *et al.*, Contractile activity-specific transcriptome response to acute endurance
exercise and training in human skeletal muscle. *Am J Physiol Endocrinol Metab* **316**, E605-
E614 (2019).

19.   B. Egan *et al.*, Exercise intensity-dependent regulation of peroxisome proliferator-
activated receptor coactivator-1 mRNA abundance is associated with differential activation
of upstream signalling kinases in human skeletal muscle. *J Physiol* **588**, 1779-1790 (2010).

20.   N. Brandt *et al.*, Impact of adrenaline and metabolic stress on exercise-induced intracellular
signaling and PGC-1alpha mRNA response in human skeletal muscle. *Physiological*
reports **4** (2016).

21.   N. D. Knuth *et al.*, Metabolic adaptation following massive weight loss is related to the
degree of energy imbalance and changes in circulating leptin. *Obesity (Silver Spring)* **22**,
2563-2569 (2014).

22.   J. C. Kerns *et al.*, Increased Physical Activity Associated with Less Weight Regain Six
Years After "The Biggest Loser" Competition. *Obesity (Silver Spring)* **25**, 1838-1843
(2017).

23.   L. B. Becnel *et al.*, Nuclear Receptor Signaling Atlas: Opening Access to the Biology of
Nuclear Receptor Signaling Pathways. *PLoS One* **10**, e0135615 (2015).

24.   S. Liu *et al.*, Mice lacking TR4 nuclear receptor develop mitochondrial myopathy with
deficiency in complex I. *Mol Endocrinol* **25**, 1301-1310 (2011).

25.   S. Karanth, E. K. Zinkhan, J. T. Hill, H. J. Yost, A. Schlegel, FOXN3 Regulates Hepatic
Glucose Utilization. *Cell Rep* **15**, 2745-2755 (2016).

26.   Y. Liu, J. Li, Y. Shang, Y. Guo, Z. Li, CARM1 contributes to skeletal muscle wasting by
mediating FoxO3 activity and promoting myofiber autophagy. *Exp Cell Res* **374**, 198-209
(2019).

27.   A. Galmozzi *et al.*, Inhibition of class I histone deacetylases unveils a mitochondrial
signature and enhances oxidative metabolism in skeletal muscle and adipose tissue.
Diabetes **62**, 732-742 (2013).

28.   U. Dressel *et al.*, A dynamic role for HDAC7 in MEF2-mediated muscle differentiation.
The Journal of biological chemistry **276**, 17007-17013 (2001).

29.   J. Boucher, A. Kleinridders, C. R. Kahn, Insulin receptor signaling in normal and insulin-
resistant states. *Cold Spring Harb Perspect Biol* **6** (2014).

30.   P. Pinton, R. Rizzuto, p66Shc, oxidative stress and aging: importing a lifespan determinant
into mitochondria. *Cell Cycle* **7**, 304-308 (2008).

31.   J. L. Major, M. Salih, B. S. Tuana, Interplay between the E2F pathway and beta-adrenergic
signaling in the pathological hypertrophic response of myocardium. *J Mol Cell Cardiol* **84**,
179-190 (2015).

32.   R. Goldsmith *et al.*, Effects of experimental weight perturbation on skeletal muscle work
efficiency, fuel utilization, and biochemistry in human subjects. *Am J Physiol Regul Integr*
Comp Physiol **298**, R79-88 (2010).

33.   S. E. Calvo, K. R. Clauser, V. K. Mootha, MitoCarta2.0: an updated inventory of
mammalian mitochondrial proteins. *Nucleic Acids Res* **44**, D1251-1257 (2016).

34.   V. Giorgio *et al.*, Dimers of mitochondrial ATP synthase form the permeability transition
pore. *Proc Natl Acad Sci U S A* **110**, 5887-5892 (2013).

35. A. S. Rambold, B. Kostelecky, N. Elia, J. Lippincott-Schwartz, Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc Natl Acad Sci U S A* **108**, 10190-10195 (2011).

36. V. Nasta, D. Suraci, S. Gourdoupis, S. Ciofi-Baffoni, L. Banci, A pathway for assembling [4Fe-4S](2+) clusters in mitochondrial iron-sulfur protein biogenesis. *FEBS J* 10.1111/febs.15140 (2019).

37. J. Liu *et al.*, Mitochondrial Dysfunction Launches Dexamethasone-Induced Skeletal Muscle Atrophy via AMPK/FOXO3 Signaling. *Mol Pharm* **13**, 73-84 (2016).

38. D. Ratman *et al.*, Chromatin recruitment of activated AMPK drives fasting response genes co-controlled by GR and PPARalpha. *Nucleic Acids Res* **44**, 10539-10553 (2016).

39. G. R. Steinberg *et al.*, Reduced glycogen availability is associated with increased AMPKalpha2 activity, nuclear AMPKalpha2 protein abundance, and GLUT4 mRNA expression in contracting human skeletal muscle. *Appl Physiol Nutr Metab* **31**, 302-312 (2006).

40. B. Dasgupta *et al.*, The AMPK beta2 subunit is required for energy homeostasis during metabolic stress. *Mol Cell Biol* **32**, 2837-2848 (2012).

41. B. R. Barnes *et al.*, The 5'-AMP-activated protein kinase gamma3 isoform has a key role in carbohydrate and lipid metabolism in glycolytic skeletal muscle. *The Journal of biological chemistry* **279**, 38441-38447 (2004).

42. C. Aguer *et al.*, Muscle uncoupling protein 3 overexpression mimics endurance training and reduces circulating biomarkers of incomplete beta-oxidation. *FASEB J* **27**, 4213-4225 (2013).

43. S. Zhang, M. W. Hulver, R. P. McMillan, M. A. Cline, E. R. Gilbert, The pivotal role of pyruvate dehydrogenase kinases in metabolic flexibility. *Nutr Metab (Lond)* **11**, 10 (2014).

44. J. C. Drake, R. J. Wilson, Z. Yan, Molecular mechanisms for mitochondrial adaptation to exercise training in skeletal muscle. *FASEB J* **30**, 13-22 (2016).

45. M. A. Pearen, G. E. Muscat, Minireview: Nuclear hormone receptor 4A signaling: implications for metabolic disease. *Mol Endocrinol* **24**, 1891-1903 (2010).

46. Z. Wu *et al.*, Transducer of regulated CREB-binding proteins (TORCs) induce PGC-1alpha transcription and mitochondrial biogenesis in muscle cells. *Proc Natl Acad Sci U S A* **103**, 14379-14384 (2006).

47. R. K. Conlee, M. J. Rennie, W. W. Winder, Skeletal muscle glycogen content: diurnal variation and effects of fasting. *Am J Physiol* **231**, 614-618 (1976).

48. M. A. Maxwell *et al.*, Nur77 regulates lipolysis in skeletal muscle cells. Evidence for cross-talk between the beta-adrenergic and an orphan nuclear hormone receptor pathway. *The Journal of biological chemistry* **280**, 12573-12584 (2005).

49. J. Meinken, G. Walker, C. R. Cooper, X. J. Min, MetazSecKB: the human and animal secretome and subcellular proteome knowledgebase. *Database (Oxford)* **2015** (2015).

50. S. E. Keating, N. A. Johnson, G. I. Mielke, J. S. Coombes, A systematic review and meta-analysis of interval training versus moderate-intensity continuous training on body adiposity. *Obes Rev* **18**, 943-964 (2017).

51. R. W. McGarrah, C. A. Slentz, W. E. Kraus, The Effect of Vigorous- Versus Moderate-Intensity Aerobic Exercise on Insulin Action. *Curr Cardiol Rep* **18**, 117 (2016).

52. T. S. Church, C. P. Earnest, J. S. Skinner, S. N. Blair, Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal

521 women with elevated blood pressure: a randomized controlled trial. *JAMA* **297**, 2081-2091
522 (2007).

53. C. A. Slentz *et al.*, Effects of the amount of exercise on body weight, body composition, and measures of central obesity: STRRIDE--a randomized controlled study. *Arch Intern Med* **164**, 31-39 (2004).

54. P. S. MacLean *et al.*, NIH working group report: Innovative research to improve maintenance of weight loss. *Obesity (Silver Spring)* **23**, 7-15 (2015).

55. A. J. Steig *et al.*, Exercise reduces appetite and traffics excess nutrients away from energetically efficient pathways of lipid deposition during the early stages of weight regain. *Am J Physiol Regul Integr Comp Physiol* **301**, R656-667 (2011).

56. R. Berdeaux, R. Stewart, cAMP signaling in skeletal muscle adaptation: hypertrophy, metabolism, and regeneration. *Am J Physiol Endocrinol Metab* **303**, E1-17 (2012).

57. R. Rudolf *et al.*, Alterations of cAMP-dependent signaling in dystrophic skeletal muscle. *Front Physiol* **4**, 290 (2013).

58. C. S. Choi *et al.*, Paradoxical effects of increased expression of PGC-1alpha on muscle mitochondrial function and insulin-stimulated muscle glucose metabolism. *Proc Natl Acad Sci U S A* **105**, 19926-19931 (2008).

59. M. Tadaishi *et al.*, Skeletal muscle-specific expression of PGC-1alpha-b, an exercise-responsive isoform, increases exercise capacity and peak oxygen uptake. *PLoS One* **6**, e28290 (2011).

60. R. P. Shook *et al.*, Moderate cardiorespiratory fitness is positively associated with resting metabolic rate in young adults. *Mayo Clin Proc* **89**, 763-771 (2014).

61. M. Gilliat-Wimberly, M. M. Manore, K. Woolf, P. D. Swan, S. S. Carroll, Effects of habitual physical activity on the resting metabolic rates and body compositions of women aged 35 to 50 years. *J Am Diet Assoc* **101**, 1181-1188 (2001).

Figure Legends

Figure 1. Crtc2/Creb1 signaling enables greater loss of weight and adiposity during alternate day fasting (ADF).

(**A**) Summary of previous work showing that Crtc2 overexpression produced an exercised phenotype (13).

(**B**) 18-week old C57BL/6 male control (WT) and Crtc2 mice were treated with dox for 35 days and weighed. Data are shown as box-and-whisker plot, indicating the middle quartiles and range. $N = 6$ (WT) or 5 (Crtc2) mice per group.

(**C-F**) Mice from (**B**) were subject to alternate day fasting (ADF).

(**C**) Changes in the body weight during ADF.

(**D**) Weight regain after the ADF treatment and 3 weeks of ad libitum feeding. *significant in Sidak's test for planned comparisons.

(**E**) Behaviors were assayed with Biodak monitoring system, including number of meals, bouts per meal, meal duration, and kilocalories consumed over the 12-hr re-feeding period.

(**F**) Adiposity and lean mass determined by whole-body NMR measurements taken before and after 8 cycles of ADF.

(**G**) 8-week old C57BL/6 male control and Crtc2 mice were treated with dox and placed on a high fat diet (HFD). $N = 9$ (WT) or 7 (Crtc2) mice per group

Data are shown as mean ± SEM. See also **Figure S1**

Figure 2. Transcriptional control of fasting and weight loss and its regulation by Crtc2.

(**A**) Mice TA muscles were transduced with GFP control or Crtc2 expression vector. Crtc2 mRNA levels in the TA of mice fed ad libitum (Fed), or subject to ADF (Fast) were compared by qPCR.

N = 3 mice per group.

(**B**) Venn diagram showing the numbers of DEGs identified by mRNA-seq comparing the effects

of ADF in control versus Crtc2-transduced TA muscles.

(**C**) Biological processes and functions regulated by ADF and Crtc2. Gene ontology (GO) analysis

suggests that ADF regulates several processes/ functions in a Crtc2-sensitive manner. The numbers

of DEGs involved in each process are shown. See SI Appendix, **Dataset S1** for a complete list of

represented GO annotations.

(**D**) Examples of ADF-regulated transcriptional programs and mRNAs in control TA muscles.

(**E–G**) Gene expression profiles in control and Crtc2-transduced TA muscles of mice subjected to

ADF relative to the ad libitum fed mice (columns 1–2), and effect of Crtc2 transduction relative to

control during ADF (column 3). Expression profiles of genes that encode **E**) nuclear receptors, **F**)

other transcription factors, and **G**) transcriptional coregulators. ADF-regulated genes in control

muscle appear in **bold**. *Crtc2-regulated genes in mice subjected to ADF. *ADF-regulated genes

in Crtc2-transduced muscle but not control. FC, fold change. Also see SI Appendix, **Dataset S1**

Figure 3. Electron transport chain gene expression and whole-body energy expenditure

regulation by weight loss and fasting is reversed by Crtc2 expression.

(**A–D**) Gene expression profiles in control and Crtc2-transduced TA muscles of mice subjected to

ADF relative to the ad libitum fed mice (columns 1–2), and effect of Crtc2 transduction relative to

control during ADF (column 3). The profiles of all genes encoding **A**) mitochondrial proteins,

including subunits of electron transport chain (ETC) complexes, **B**) subunits of the mitochondrial

ATPase (complex V), **C**) ribosomal subunits, and **D**) highlighted mitochondrial proteins, are

shown. ADF-regulated genes in control muscle appear in **bold**. *Crtc2-regulated genes in mice

subjected to ADF. *ADF-regulated genes in Crtc2-transduced muscle but not control. FC, fold

change.

(**E**) Effects of Dex on lean body mass. Mice injected with vehicle or 10 mg/kg Dex were fasted

overnight. Lean mass was measured by whole-body NMR before and after treatment. $N = 5$ mice

per group.

(**F**) Diameter and mitochondrial potential of steroid-deprived primary myotubes treated for 2 days

with vehicle or 100 nM Dex were compared by confocal high-content imaging. $N = 24–27$ images

x 50–100 myotubes per image.

(**G**) Steroid-deprived C2C12 myotubes were stimulated with 100 nM Dex for 0–12 h. Levels of

phosphorylated and total AMPK in whole lysates were compared by western blot.

(**H**) Diameter of steroid-deprived C2C12 myotubes treated for 48 h with vehicle, 100 nM Dex, or

1 uM AICAR were compared by confocal high content imaging. $N = 24–27$ images x 50–100

myotubes per image.

(**I**) Expression levels of genes encoding AMPK subunits in mice described in **Figure 2**.

(**J**) VO_2 and VCO_2 were measured continuously (**Figure S2A**) for 72 hrs. in a CLAMS animal

monitoring system. Shown are the 12-hr averages. $N = 8$ mice per group.

(**K**) Total energy expenditure was calculated from VO_2 and VCO_2 values using the Lusk equation.

Data are shown as mean ± SEM. Also see **Figure S3**

Figure 4. Crtc2 expression in skeletal muscle enhances mitochondrial biogenesis.

(**A**) Expression profiles of genes that regulate lipid metabolism in mice described in **Figure 2.**

(**B**) Histological analysis of succinate dehydrogenase in myofibers from gastrocnemius muscle

sections after dox treatment. Scale bar = 500 μm.

(**C**) Gene induction calculated relative to WT mice. $N = 8$ mice per group.

(**D**) Gene expression in primary myocytes transduced with adenovirus expressing Crtc2 or GFP

control 72 hr post-transduction.

(**E**) C2C12 myocytes were transduced with adenovirus GFP, Crtc2, or A-Creb genes and analyzed

by confocal high content imaging. $N = 24$–27 images x 50–100 myotubes per image.

(**F**) Crtc2 expression plasmid was electroporated into the tibialis anterior muscle and GFP plasmid

in the contralateral leg. After 7 days, DNA was extracted for assessment of mitochondrial DNA

(mt-DNA) for Crtc2/GFP control. $N = 5$ mice per group.

(**G**) Oxidation of ^{3}H-Palmitic acid in the supernatant of primary myocytes transduced with control,

Crtc2, or Crtc3-expressing adenovirus.

* $p < 0.05$. Data are shown as mean + SEM. **See also Figure S4**

Figure 5. Crtc2 transgenic mice maintain enhanced glucose disposal and insulin tolerance

(**A**) Expression profiles of genes that regulate carbohydrate metabolism in mice described in

Figure 2.

(**B**) Glucose tolerance test (GTT) after alternate day fasting. Right panel, area under the curve. N

$= 10$ mice per group

(**C**) Plasma insulin levels during GTT described in panel **B**.

(**D**) Insulin tolerance tests after 8 cycles of ADF. $N = 10$ mice per group

(**E**) Insulin-mediated uptake of 2-deoxy-D-glucose in primary myocytes transduced with

adenovirus expressing Crtc2 or GFP control. $N = 10$ per group. Student's t-test.

(**F**) Crtc2 mice were treated +/- dox for 1 week and the gene induction compared by qPCR relative

to the no dox groups. N= 10 per group.

(**G**) Gene expression in primary myocytes transduced with adenovirus expressing Crtc2 or GFP

control were compared by qPCR. *N*=4 per group.

(**H**) Western blot of whole lysates from primary myocytes transduced with the indicated plasmids.

(**I–J**) Myocytes stably expressing GFP, Crtc3 (lanes 2 and 3), dominant negative Creb (A-CREB),

or dominant negative Crtc2 (dn-Crtc) were compared by western blots.

(**K**) WT or Crtc2 mice were injected twice daily with 0.5 mg/kg leptin and weight was monitored.

N=8 mice per group.

Data are mean ± SEM. **See also Figure S5.**

Figure 6. Hormonal and neural modulation of energy expenditure in skeletal muscle

(**A**) Fasting-induced regulation of mitochondrial dynamics and mitophagy.

(**B**) Exercise-induced regulation of mitochondrial dynamics and mitochondrial biogenesis, which

can be constitutively activated by overexpression of Crtc2 in skeletal muscle.

bAR, b-adrenergic receptor; GR, glucocorticoid receptor; HPA, hypothalamic-pituitary-adrenal axis;

HPT, hypothalamic-pituitary-thyroid axis; InsR, insulin receptor; nAchR, nicotinic acetylcholine

receptor; SNS, sympathetic nervous system; TH, thyroid hormone

Materials and Methods

Statistical analyses

Changes in body weight and body were analyzed with 2-way analysis of variance. For VO_2, respiratory quotient, and ambulation data, the ad libitum and fasted periods were analyzed separately with 2-way ANOVA, and the refeeding data analyzed with Student's T-test. For the energy consumption data, the fasted and re-fed data were analyzed together with 2-way ANOVA. Gene expression data were analyzed with 1-way ANOVA, with Dunnett's test for multiple comparisons of each gene to its corresponding control. Glycemia and insulin changes over time were analyzed with 2-way ANOVA, and the AUC assessed with Student's T-test. The behavioral data for meal consumption were analyzed with Student's T-tests.

Crtc2 transgenic mice and animal care

All animal procedures were approved by the Scripps Research Institute ICACUC. TRE-Crtc2[tm] mice were generated by cloning FLAG-Crtc2 (S171A, S275A, K628R) into pTRE-Tight. Oocyte injections were conducted into C57Bl6 at the University of Cincinnati. Double transgenic mice (DTg) were generated by crossing C57BL/6-Tg(TRE-Crtc2tm)1Mdc/j mice with 1256[3Emut] MCK-rtTA transgenic mice, which expresses the reverse tetracycline transactivator (rtTA) regulated by the 1256[3Emut] MCK promoter specifically in skeletal muscle (Grill *et al*, 2003). Wild type and TRE littermates were used as controls and treated with DOX in the same manner as DTg mice. High fat diet (Research Diet D12450J) contained 60 kcal% fat and matched control low fat diet (D12492) contained 10kCal% fat and 7% sucrose.

Data Availability

The raw and processed RNA-seq dataset is freely available to the public at the National Center for

Biotechnology   Information   (NCBI)   Gene   Expression   Omnibus   (GEO)

https://www.ncbi.nlm.nih.gov/geo/ and can be retrieved with the accession number GSE149150,

which will be released upon publication of this manuscript. The processed RNA-seq dataset is also

provided in SI Appendix, **Dataset S1**. Relevant protocols and raw data not detailed in the Methods

or Supplementary Information are available upon request.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6