1 H3K27me3 is dispensable for early differentiation but required to maintain differentiated cell
2 identity
3

Sara A. Miller[1,2], Manashree Damle[1], Robert E. Kingston[1,2,*]

AFFILIATIONS:

[1]Department of Molecular Biology, Massachusetts General Hospital Research Institute,

Massachusetts General Hospital, Boston, Massachusetts 02114, USA

[2]Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA

*Correspondence: kingston@molbio.mgh.harvard.edu

11
12
13 **Abstract**
14

Polycomb repressive complex 2 (PRC2) catalyzes trimethylation of histone H3 on lysine 27 and
is required for normal development of complex eukaryotes. The requirement for H3K27me3 in
various aspects of mammalian differentiation is not clear. Though associated with repressed
genes, the modification is not sufficient to induce gene repression, and in some instances is not
required. To examine the role of the modification in mammalian differentiation, we blocked
trimethylation of H3K27 with both a small molecule inhibitor, GSK343, and by introducing a
point mutation into EZH2, the catalytic subunit of PRC2. We found that cells with substantively
decreased H3K27 tri-methylation were able to differentiate, which contrasts with EZH2 null
cells. Different PRC2 targets had varied requirements for H3K27me3 in repressive regulation
with a subset that maintained normal levels of repression in the absence of methylation. The
primary cellular phenotype when H3K27 tri-methylation was blocked was an inability of the
altered cells to maintain a differentiated state when challenged. This phenotype was
determined by H3K27me3 deposition both in embryonic stem cells and in the first four days of
differentiation. H3K27 tri-methylation therefore was not necessary for formation of
differentiated cell states but was required to maintain a stable differentiated state.

30
31 **Introduction**
32
Polycomb repressive complex 2 (PRC2) is a highly conserved protein complex that is
required for proper axial patterning of vertebrates. It is comprised of the core subunits EZH2,
SUZ12, EED and RBAP48, with additional subunits some of which are cell-type or developmental
stage specific (Healy et al., 2019; Margueron & Reinberg, 2011; Shen et al., 2009; van Mierlo et
al., 2019). The complex is crucial for differentiation, but it is not required for the self-renewing
phenotype associated with embryonic stem (ES) cells (Aloia et al., 2013; Chamberlain et al.,

2008). At a molecular level, PRC2 family complexes are the only histone 3 lysine 27 (H3K27)
methyltransferases identified in mammals. EZH2 is the primary catalytic component of these
complexes, with its paralog EZH1 also contributing to catalytic activity in some instances
(Margueron et al., 2008; Shen et al., 2008; Wassef et al., 2019; Xu et al., 2015). Lysine residues
can be mono- di- or tri- methylated, with PRC2 able to catalyze all levels of methylation.
Increased methylation is generally associated with gene repression, and H3K27 tri-methylation
(H3K27me3) is a key marker of facultative heterochromatin. However, the tri-methylation of
H3K27 is neither required nor sufficient to induce gene repression (Ahmed et al., 2018;
Chamberlain et al., 2008; Ferguson et al., 2018; O'Geen et al., 2017). Dissecting the role of tri-
methylated H3K27 in gene regulation is important for understanding how genes become
repressed during development and retain their repression in differentiated cells.
50
Expression levels of Polycomb complexes have developmental and disease effects. Both
hyper-activating and inactivating mutations in PRC2 have been identified from a variety of
human cancers (Basheer et al., 2019; Cyrus et al., 2019; Jain & Di Croce, 2016). Malignancies
with both types of alterations to PRC2 function have been linked to cancer progression and
poor prognosis in a wide variety of tissue (Abdel Raouf et al., 2019; Basheer et al., 2019; Bohm
et al., 2019; Bremer et al., 2019; Deng et al., 2019; Dou et al., 2019; Karlowee et al., 2019; Krill
et al., 2020; Matsubara et al., 2019; Mechaal et al., 2019; Shi et al., 2019; Tian et al., 2019;
Wasenang et al., 2019; M. J. Zhang et al., 2019; Q. Zhang et al., 2019). These findings are
consistent with the initial identification of Polycomb-Group genes in Drosophila where
haploinsufficiency yielded developmental phenotypes (Lewis & Mislove, 1947; Schuettengruber
et al., 2017). Inserting these disease mutations into cells alters the modification profiles of
H3K27 and can change their developmental potential. Increased methylation can often push
cells toward a neuro progenitor phenotype (Juan et al., 2016; Pasini et al., 2007; Thornton et
al., 2014). In cancer patients, increased levels of expression of PRC2 components and especially
EZH2 leads to hypermethylation and is associated with poor prognosis (Tian et al., 2019; Wu et
al., 2019). Consequently, there is significant interest in targeting this complex pharmaceutically.
Small molecules that inhibit PRC2 methyltransferase activity have been approved for clinical
trials (Fioravanti et al., 2018; Kondo, 2014; Liu et al., 2015; Lue & Amengual, 2018; Qi et al.,
2012; Shi et al., 2019; Xu et al., 2015; Yamagishi & Uchimaru, 2017; Yang et al., 2019). It
remains, however, unclear how far this activity can be manipulated before risking adverse
effects from having too little of the modification present. The ambiguity of the role of
H3K27me3 in disease progression increases the importance of understanding the mechanisms
by which PRC2 methylation activities regulate gene expression during development.
74
Previous work on PRC2 has shown that the complex is dispensable for the propagation
of self-renewing, undifferentiated cells that are phenotypically indistinguishable from WT ES
cells. These studies have also shown that PRC2 deficient cells do not differentiate properly
(Chamberlain et al., 2008; Lavarone et al., 2019; Pasini et al., 2007; Pasini et al., 2004; Shen et
al., 2008). Since the entire complex is disrupted when any of the core components are knocked
out, it is not clear whether it is solely the lack of methylation that is causing these
developmental defects, or whether there are other mechanistic roles PRC2 plays that are
distinct from methylation(Collinson et al., 2016; Rai et al., 2013; Shan et al., 2017; Yu et al.,

2017). A recent study examined the impact of a mutation in *Ezh2* that blocks all levels of
methylation. When these cells were differentiated into embryoid bodies they showed visual
phenotypic differences (Lavarone et al., 2019), demonstrating that complete removal of the
methyltransferase activity of PRC2 impairs normal differentiation. We focus here on the
developmental phenotypes caused by specific loss of the H3K27me3 modification, as opposed
to loss of all methylation.
89
To determine the contribution that H3K27me3 makes to gene repression, we examined
cells with a point mutation within the SET domain of *Ezh2* that generates a hypomorph that is
predominately defective in tri-methylation. We also analyzed cells treated with a small
molecule inhibitor that blocks tri-methylation more efficiently than di-methylation. The
expression level of PRC2 has known effects on gene regulation, so we used strategies that do
not impact the protein level or complex integrity, thus allowing a focus on methyltransferase
activity. There are multiple small molecules that interrupt PRC2 methyltransferase activity both
for research and current clinical trials; we used GSK343 (Bradley et al., 2014; Fraineau et al.,
2017; Liu et al., 2015; Xu et al., 2019; Yang et al., 2019). Mutations in the catalytic SET domain
of *Ezh2* have been identified from a variety of cancers; we created cell lines containing one of
the inactivating mutations for our experiments (Antonysamy et al., 2013). We found that this
mutation had a strong impact on H3K27me3 levels, but not on H3K27me2 levels. When these
cells were differentiated in an undirected fashion they did not show the significant visual
phenotypes seen with *Ezh2* knockout cells. While expression of some genes was altered, there
were a number of PRC2 target genes that did not rely upon H3K27me3 for their regulation. The
most dramatic phenotype observed was when cells were challenged to maintain their
differentiated identity. Inhibitor treated and point mutated cells reverted readily to an ES
phenotype when placed back into conditions that support ES cell growth. Thus, H3K27me3 was
required for cells to maintain their identity rather than for the initial differentiation.
109
Results
PRC2 catalyzes di- and tri-methylation of histone H3 and is associated with gene
repression. Previous studies have shown that deletion of PRC2 components in mouse ES cells
has minimal effects on the undifferentiated, self-renewing state (Juan et al., 2016; Lavarone et
al., 2019; Shen et al., 2008; Wassef et al., 2019). Yet deletion of EZH2, the catalytic subunit of
PRC2, results in cells that do not differentiate and embryos that are reabsorbed by day E10.5
(Pasini et al., 2004). However, the requirements for specific methylation functions of PRC2 in
differentiation and in the accompanying changes in gene expression have not been explored in
detail. We examined the role for H3K27me3 in regulation by looking at stochastic
differentiation of ES cells into embryoid bodies. Embryoid body (EB) formation is a well-studied
method of undirected differentiation with predictable changes in gene expression (Fig 1A and
(Behringer et al., 2016; Dang et al., 2002). Cells where PRC2 core components haven been
deleted fail to form normal embryoid bodies, presumably because the cells apoptose when
pushed to differentiate. Since PRC2 has been shown to play regulatory roles independent of
H3K27 methylation, the exact requirement for the H3K27me3 modification at individual genes
during differentiation as well as its role in maintaining cell identity remains unclear (Ahmed et
al., 2018; Ai et al., 2017; Pereira et al., 2010).

127
Inhibiting methyltransferase activity with the small molecule GSK343 does not block
differentiation potential

130
As a starting point to understand the role of H3K27 methylation by PRC2 during ES cell
differentiation, we treated cells with the small molecule inhibitor GSK343 which blocks the
methyltransferase activity of EZH2 (Fioravanti et al., 2018; Lue & Amengual, 2018; Yang et al.,
2019). Treatment with DMSO was used as a control (Fig 1B). Unlike PRC2 knockouts, GSK343
treated cells form embryoid bodies that have a similar morphology to their DMSO treated
counterparts (Fig 1C.) Embryoid bodies formed after either DMSO or GSK343 treatment are
phenotypically similar to those made by WT cells. In contrast, PRC2 knockout cells tested in
parallel failed to form embryoid bodies ((Lavarone et al., 2019) and Fig. 2E). Thus, the inhibition
of methyltransferase activity can be phenotypically separated from PRC2 knockout cells.

140
To determine the molecular effects of blocking methyltransferase activity, we
performed RNA-seq experiments to compare changes in gene expression between cells treated
with the inhibitor and control over the differentiation time course. We examined PRC2 target
genes and found examples of genes whose expression pattern across differentiation was the
same in DMSO and GSK343 treated cells (e.g. Tbx3 or Sox3, Fig. 1D) as well as genes whose
expression was altered in GSK343 treated cells (e.g., Wnt3 and Itgb7, Fig 1D). The trends
observed with these two classes of genes were found to be mirrored in large sets of genes. We
classified PRC2 target genes into two sets, those that required H3K27me3 for their regulation
and those that do not need this modification to preserve normal gene expression patterns
during embryoid body formation. Genes were defined as independent of the H3K27me3
modification if their expression remained within 1.2 fold of the WT expression at all time points
(Fig 1E). We conclude that inhibiting methylation activity of PRC2 does not block differentiation
but does alter gene expression at a subset of genes though not at others.

154
Point mutation to the SET domain inhibits methyltransferase activity

156
We sought to extend and refine the findings made using GSK343 by generating a
hypomorphic mutation in *Ezh2* that had a defined impact on H3K27 methylation. We generated
mutant cells that allowed us to examine the contribution of H3K27 tri-methylation to gene
regulation and differentiation during EB formation. Inactivating point mutations in the SET
domain of EZH2 have been identified in several human cancers. We chose one of those
mutations, human R635C, which is analogous to mouse R681C, for analysis in mouse cells. This
residue is located in a region that is important for coordinating the methyl donor (Fig 2A and
(Antonysamy et al., 2013). We purified PRC2 complexes comprised of the core components
RBAP46/48, EED, SUZ12, EZH2 (WT or mutant) and the accessory protein AEBP2 (Supp. Fig. 2).
We confirmed that this residue is important for tri-methylation of H3K27 by PRC2 using an in
vitro methyltransferase assay. We compared it to WT EZH2 and another SET domain mutation
(722D) that has been previously shown to block methylation (Lavarone et al., 2019), Fig 2B).
The complexes that contained WT EZH2 methylated both core histones and purified H3 in a
time and concentration dependent manner (Supp. Fig. 2). In contrast, mutant complexes show

drastically reduced levels of activity. Using two separate purifications, residual activity of the
mutant complex was always under 2% of the purified WT complexes (Fig. 2B and Supp. Fig. 2).
We conclude that the *Ezh2* 681C mutation reduces the methyltransferase activity of the
complex by at least 50-fold.
175
We introduced the R681C mutation into the CJ7 mouse ES cell line using the CRISPR-
Cas9 system. We isolated two independent clones in the mouse ES cells line CJ7 with this
homozygous point mutation in the endogenous *Ezh2* gene (called CJ7 Ezh2[681C-99] and CJ7
Ezh2[681C-102]). The phenotypes that we describe below were consistent between these two
independent lines. The mutant cells look phenotypically similar to WT cells and self-renew (Fig
2C). This was the anticipated result since knockout cells in the PRC2 core components EZH2 and
EED are also phenotypically similar to WT cells and can self-renew. We examined whether the
point mutation would lead to significant reduction of H3K27me3 in cells as anticipated from the
enzymatic defect seen in vitro. Lysine residues can be mono, di or tri-methylated. Mono-
methylation is spread broadly throughout the genome and di-methylation is loosely linked with
repressed genes, though this is poorly defined in most cell types. Tri-methylation of H3K27 is
closely associated with repressed genes, presumably due to binding of the CBX family of
proteins contained in the PRC1 complex to K27me3 and subsequent repression by this family of
complexes (Bernstein et al., 2006). We performed western blots on whole cell lysates from WT,
knockout and the point mutant cells and probed them with antibodies to di- and tri- H3K27
methylation and EZH2 (Fig 2D and Supp. Fig. 3). H3K27me3 was reduced to levels we could not
detect in the mutant cells, as was seen the knockout cells, but the levels of di-methylation were
largely unaffected in the R681C mutant cells. We conclude that in these cells the major effect of
the point mutation was to reduce the tri-methylation of H3K27. This is similar to the results
found in other studies of *Ezh2* point mutations (Lavarone et al., 2019). The R681C mutation
therefore offers an opportunity to examine the contribution of tri-methylated H3K27 to gene
regulation during differentiation.
198
We examined whether the mutant cell lines would form embryoid bodies similar to
those formed by WT cells, would apoptose like *Ezh2* or *Eed* knockout cells, or would take an
alternate path. We formed embryoid bodies by the hanging drop procedure and compared the
resultant phenotypes of WT, the point mutant and knockout cells. We found that the mutant
cells formed embryoid bodies, making them phenotypically separate from the knockout cells
(Fig 2E). This revealed that the dramatic reduction in H3K27me3 we see in the point mutant
cells does not stop the cells from differentiating, at least at a gross level. Another method for
testing the developmental potential of embryonic stem cells is assessing the formation of
beating clusters of cardiac cells. For this assay cells are differentiated in hanging drops and then
plated in individual wells to form a monolayer in differentiation media. After 10-15 days of
differentiation each well is visually examined for the presence of pulsing cells. Beating clusters
formed from both WT and mutant cells. There was some variation in the proportion of EBs that
could form beating clusters between the two 681C mutant strains, but in all cases the mutant
cells could form beating clusters (Fig. 2F). These data bolster the conclusion that the reduction
in H3K27me3 does not prevent cells from differentiating. This phenotype raised the issue of
whether there are molecular changes caused by the point mutation.

215
Molecular phenotype of *Ezh2* point mutant cells

217
To determine whether there were molecular phenotypes associated with the
hypomorphic mutation of *Ezh2*, and the resultant loss of H3K27me3, we determined the target
genes of PRC2 in our cell lines and measured their gene expression levels. We performed
CUT&RUN analysis using antibodies to H3K27me3, H3K27Ac, RING1b, EZH2 and SUZ12 in WT
and point mutant cells as embryonic stem cells (ESC) and as Embryoid bodies after
differentiation for eight days (D8EB). There have only been a small number of studies of PRC2
localization in differentiating cells, but of the approximately seven thousand target genes we
identified as having an H3K27me3 peak within 1kb of their transcription start site in WT cells,
just over 70% overlapped with previously published data (Fig. 3A and (Juan et al., 2016)). We
note that peaks called from our data set that did not overlap with previous called peaks showed
signal in the published data that was below the threshold, indicating further agreement
between the analysis done here and previous work. There were far more targets called using
the H3K27me3 than with the individual components of either polycomb complex, though the
targets are largely overlapping (Fig. 3B). This is likely due to differences in the strength of
individual antibodies. Therefore, we used the targets identified with H3K27me3 for further
analysis. In keeping with the western blot from whole cell lysates, H3K27me3 was reduced on
target genes in mutant cells (Fig. 3C). Levels of H3K27me3 were significantly lower in mutant
cells at day 4 of differentiation and then were increased at day 8 of differentiation but to levels
well below WT levels. The residual tri-methylation at day 8 might be due to EZH1 function
(Lavarone et al., 2019; Margueron et al., 2008; Shen et al., 2008). We focus below on the events
that happen during these first four days of differentiation and the impact of the lack of
H3K27me3 during this time frame. We conclude that the ability of the mutant cells to
differentiate was not due to retention of normal tri-methylation levels specifically on target
genes.

The experiments described above using small molecule inhibitors revealed that there
are genes that require H3K27me3 for their regulation and those that do not require H3K27me3
to maintain proper regulation. We investigated whether the point mutant cells showed these
same two gene classes. We performed RNA-seq analysis from ES cells and both D4 and D8EBs
from WT, CJ7 Ezh2$^{681C-99}$ and CJ7 Ezh2$^{681C-102}$ cells. The data for both mutant cell lines are nearly
overlapping and so for clarity we present the data from the CJ7 Ezh2$^{681C-99}$ (Supp. Fig. 4). For
this analysis we defined PRC2 target genes as those that had a statistically significant peak of
H3K27me3 from CUT&RUN at any time point in the EB formation protocol. PRC2 target genes
were identified using wild-type H3K27me3 peaks called by Homer using a p-value threshold of
0.001, a length of at least 1500 bp, 1rpkm in at least two time points and signal overlapping
Refseq annotated genes (TSS+-5kb). As with the cells treated with the small molecule inhibitor,
we could separate genes that rely on H3K27me3 from those that do not need a high level of the
modification for their regulation (Fig 3D). The gene expression patterns from the mutant cells
and the GSK343 treated cells clustered based upon day of differentiation rather than by
whether they were drug treated or mutant (Supp. Fig. 5), although the correlation was not as
strong as that seen between the two mutant cell lines. We detected both activated and

repressed genes that fell into methyltransferase-dependent and –independent categories. The
350 genes that are normally repressed during differentiation in WT cells, and were either
similarly repressed or were not repressed in mutant cells, showed more consistent patterns
than the genes that were normally activated in WT cells. These patterns were well established
by D4 of differentiation, thus we used repressed genes to further analyze any characteristics
specific to the methyltransferase dependent or to the independent genes.
265
We examined whether the level of H3K27me3 normally found on the genes in WT cells
or the amount of signal remaining in the mutant cells could predict whether a gene would be
dependent on the modification for its regulation. However, there was not a difference in the
levels of tri-methylation based on whether the genes require this modification for their
regulation (Fig. 3E). We then examined several characteristics of 350 genes that are normally
repressed during differentiation of WT ES cells including other histone modifications, CpG
methylation, and further sub dividing genes by their dependence on H3K27me3 (Supp. Fig. 6).
None of these characteristics showed any significant differences between genes who repression
was dependent upon methyltransferase activity and genes whose repression was not
dependent on methyltransferase activity. We conclude that there is a variable reliance on
H3K27me3 for gene regulation and that H3K27me3 is not the only driver of PRC2 target gene
repression during EB formation, just as full levels are not needed for differentiation.
278
279   **Methyltransferase point mutants cannot maintain differentiated cell identity**
280
A major issue in developmental gene expression concerns the interplay between
establishment and maintenance of gene expression profiles during differentiation. The
Polycomb group (PcG) system, including PRC2, plays a role in both aspects of gene regulation in
flies and in mammals. Substantial early work on mutant Drosophila highlighted a role for the
PcG, including gene products now known to compose PRC2, in maintenance. Given that we saw
limited impact of the ablation of H3K27me3 on establishment of the differentiated EB
phenotype, we tested whether the mutant or drug treated mouse cells were able to maintain a
differentiated state.
289
Under normal conditions, cells differentiated into embryoid bodies cannot revert to
ESCs without major manipulation such as introducing Yamanaka transcription factors
(Nakagawa et al., 2008; Takahashi & Yamanaka, 2006). To determine if the 681C point mutant
cells stably committed to a differentiated state, we formed day 8 EBs with WT and mutant cells,
dissociated the embryoid bodies into single cells and re-plated them in ESC conditions without
any additional manipulation. The cells were cultured in the ESC media for five days and then
stained for alkaline phosphatase (AP) activity. After incubation with an appropriate
colorometric substrate, ESCs become bright pink due to their alkaline phosphatase activity,
while feeder cells or any other differentiated cells remain unstained. We found a ten-fold
increase in the number of AP-positive colonies generated by the mutant cells over the WT (Fig.
4A and 4B). We asked whether this lack of commitment was maintained after longer periods of
differentiation and found that after 14 days there remained more cells that could revert to ES
cells, but to a considerably lesser extent than seen after 8 days (Figs. 4A, 4B). We conclude that

the mutant cells are not stably committed to the more defined lineages but can switch back to
an undifferentiated state, but that this flexibility decreases after two weeks of differentiation.
305
We used the small molecule inhibitor GSK343 to examine the time period during which
this flexibility is established. There were three possibilities: a) full levels of H3K27me3 might be
needed continuously to maintain the differentiated state; b) they might be needed at specific
times as the cells are differentiating; c) they might be needed only when cells are challenged
with external stimuli that allow reversion to embryonic stem cells. Examination of the mutant
cells addressed blocking full levels of methylation throughout the experimental time course,
but did not address the time period when that lack of activity was most important. To separate
the possibilities, we added or removed GSK343 during the differentiation time-course (Fig. 5A
shows the experimental design). To mimic the WT and point mutant conditions a subset of cells
were treated continuously with DMSO or GSK343 respectively. These treatments served as both
a baseline for altered times of GSK343 application. They also served to validate that the results
observed with re-plating of mutant cells were not caused by off target effects of the CRISPR
genetic manipulation or by mutations acquired during selection of these cells. Cells
continuously treated with GSK343 had higher numbers of AP-positive cells following re-plating
at 8 days, similar to the mutant cells; in contrast to the mutant cells there even more AP-
positive cells following re-plating after 14 days of differentiation in GSK343. GSK343 treatment
affects both di- and tri-methylation which might account for the increased plasticity seen at
later time points in drug treated cells (Supp. Fig. 7).
324
In addition to continual treatment with inhibitor, we varied the timing of GSK343
addition as depicted in Fig. 5A. Briefly, we treated ES cells for three days with either DMSO or
GSK343 before starting the differentiation. When embryoid body formation was initiated or at
day 4 of differentiation we changed the treatment from DMSO to GSK343 (or vice versa) for
half of the cells. We also switched the treatment at the time of re-plating into the ES conditions
at day 8. Finally, we allowed cells to differentiate to day 14 and switched treatment regimen
(Fig. 5A). The treatment type was switched just once in each scheme so we could determine
whether blocking the methylation of H3K27 at early or later stages of differentiation had the
greatest effect. These experiments allow us to determine whether reducing H3K27me3 has the
greatest effect on cell identity when the cells are differentiating, when they are challenged by
re-plating or throughout the differentiating time-course.
336
We found that the crucial window for treatment with the inhibitor was in the first four
days of embryoid body formation (Fig. 5A, B, C). Increased staining by alkaline phosphatase
activity was seen in all cells that had initially been treated with inhibitor, regardless of when it
was removed (Fig. 5C). Notably, treating with GSK343 for three days prior to inducing
differentiation, then removing the inhibitor when differentiation was initiated, led to a
significant increase in ES cells following re-plating at day 8. The only case where we saw
increased staining in re-plated cells that were initially treated with DMSO was when the
inhibitor was added at the onset of differentiation (Fig 5B, C). We conclude that the inhibition
of methyltransferase activity either immediately prior to differentiation or during the first four

days of differentiation allows the cells to remain in a more plastic state such that they can
revert to a stem cell-like phenotype when placed in the proper growth conditions.
348
When we examined the ability of treated cells to form beating colonies after treatment
with the small molecule we found no difference in the proportion of colonies that
spontaneously start beating between any of the treatment groups (Fig. 5D). This is consistent
with the data from the point mutant cells where both WT and mutant cells could form beating
colonies and further indicates that there is no deficit in the ability of the cells to differentiate
when H3K27 methylation is inhibited. Thus, as was seen with the comparison of WT and point
mutant cells, the obvious difference between DMSO and small molecule treated cells occurred
when cells were challenged to grow in the ESC culture conditions (Figs 5B,C). We conclude that
H3K27me3 methylation is more important in maintaining the differentiated state than in
generating that state, and that the critical time window occurs early in the differentiation
process.
360
The GSK343 treated cells and the mutant cells both appeared to revert to a pluripotent
state. To verify that these cells retained developmental potential we re-differentiated these
cells and determined whether they display the characteristic ability of ES cells to develop more
committed cells. We used the re-plated cells from both 681C mutant cells and those treated
with the inhibitor and attempted to make embryoid bodies. Both sets of re-plated cells were
able to make embryoid bodies, which confirms that they have developmental potential (Data
not shown).
368
To examine whether the reversion to a pluripotent state involved substantive changes in
gene expression, as opposed to gene expression changes in a few key genes, we examined
genome-wide gene expression pattern of the reverted cells and compared those to ESCs. To
examine the molecular phenotype of the re-plated cells, we isolated and sequenced RNA from
drug treated day 14 cells that were re-plated and had reverted to ES cell phenotype.
Unsupervised clustering showed that the average expression profiles of the cells that had been
treated with GSK343, and therefore had much higher levels of reversion, were more like WT ES
cells than to the DMSO treated and re-plated control cells (Fig. 5E.) We conclude that the cells
we see staining with alkaline phosphatase in our re-plating assays are reverting to an ES
phenotype. We examined the subset of PRC2 target genes that are normally repressed during
differentiation and found that many are reactivated when cells treated with GSK343 are re-
plated. This differs from the genes that are reactivated in the cells initially treated with DMSO.
The changes in gene expression observed following re-plating of cells treated with GSK343 were
significantly different from the patterns observed after re-plating of cells treated with DMSO.
(Supp. Fig. 7C). These expression pattern changes show the same response to the time of
treatment with GSK343 as seen above; reversion to the WT pattern requires GSK343 treatment
early in differentiation. From these data, we conclude that though all cells are placed under
developmental stress when re-plated into ES conditions, only those where H3K27 methylation
has been blocked revert to an ES cell phenotype. This underlines the importance of K3K27me3
in establishing the heritable gene expression profile of differentiated lineages.
389

Discussion

These studies offer two advances in understanding the role for tri-methylation of H3K27 during differentiation of ES cells into embryoid bodies. First, many PRC2 targets continue to be regulated in a normal manner during the first four days of differentiation despite significantly reduced H3K27me3 levels on these targets (Fig. 3.) Thus, H3K27me3 is not necessary for repression of a significant set of PRC2 targets, indicating compensating mechanisms for repression of these genes. Second, while many PRC2 targets are dysregulated when H3K27me3 is blocked, we did not observe a significant impact on differentiation. In contrast, there was a large enhancement of the ability of differentiated cells to revert to a pluripotent phenotype when placed into embryonic stem cell culture conditions. We conclude that the primary role for H3K27me3 during early differentiation is maintenance of the differentiated state.

Differentiated WT cells are not generally capable of reverting to an ES phenotype when their growth conditions are altered by a change in media. In normal cases, it takes the reactivation of key transcription factors to allow cells to return to that state. In contrast, *Ezh2* point mutant cells and those that that have been treated with a small molecule inhibitor readily revert to an ES phenotype when placed into media that supports that type of growth. This was seen at both the cellular and molecular level. When we varied the time windows where inhibitor was present, we found that the crucial window for normal H3K27me3 levels was between the start of differentiation through the first four days of embryoid body formation. Not having the ability to add the tri-methylation modification at those early stages of differentiation sets the stage for the cells to be able to revert to an ES phenotype when challenged, even if H3K27me3 is restored later during embryoid body formation. In the setting of the early stages of ES differentiation into embryoid bodies, the H3K27me3 modification is acting analogously to a stopper on a swinging door. When the modification is present the door will only open one way and the cells cannot go backward, but removal of the modification enables the door to swing both ways allowing cells to go back and forth between the differentiated and pluripotent state in response to external signals.

There are potential implications for these data in terms of the use of small molecule inhibitors in therapeutic situations. Multiple adult cell types have different requirements for PRC2 during their differentiation. Most relevant to the inhibition of PRC2 is the development of blood cells. PRC2 is required for the differentiation of blood stem cells and is the site of some of the highest levels of PRC2 expression in healthy adult tissues. If the phenotypes that we have observed during embryoid body formation occur in a similar manner in blood stem cells, treating patients with the small molecule inhibitors might alter the stability of commitment of healthy stem cells, raising the possibility of novel cancers arising from cells that cannot stably differentiate. Indeed, at least one clinical trial was temporarily suspended because patients had developed novel cancers (Fioravanti et al., 2018; Harris, 2018; Italiano et al., 2018). Determining the effect of blocking H3K27me3 in other differentiating cell lineages might be important to clinical intervention by expanding the knowledge of the potential side effects.

We note that it is difficult to completely eliminate H3K27 methylation with mutations
(Lavarone et al., 2019), and that the mutation we generated in *Ezh2* specifically impacts
H3K27me3, especially early in differentiation, but does not eliminate all methylation of H3K27.
This is both a limitation and an advantage; the significant impact on H3K27me3 early in
differentiation allowed us to show that loss of tri-methylation has a potent phenotype
(unstable commitment) at this stage yet does not have a discernible differentiation phenotype.
These data demonstrate a striking difference in the dependency of these two key phenotypes
on H3K27me3. This *Ezh2* mutation also has a molecular phenotype when gene regulation is
examined, raising the possibility that the network of genes that require H3K27me3 for
appropriate regulation at this stage are primarily responsible for driving stable commitment to
the differentiated state.

Methods

Cell Culture

ES cells: CJ7 WT, CJ7 *Ezh2*-/- and CJ7 *Eed*-/- cells were a generous gift from the laboratory of
Stuart Orkin. We cultured all embryonic stem cells on a monolayer of feeder MEFs in ES media
(DMEM (Gibco 11995-506) supplemented with 20%FBS, NEAA, pen/strep, glutamax, and LIF) on
tissue culture treated flasks coated with 0.2% geletin. Media was changed daily and cells were
split every 2-3 days.

EZH2 point mutant Cells: Point mutations were introduced into CJ7 WT using CRISPR RNP
transfected into cells with the Amaxa mouse ES kit (Lonza VPH-1001). $1x10^6$ cells were
transfected with the RNP containing two separate guide RNAs a single stranded donor oligo and
a linearized puromycin resistance gene. Cells were plated into three wells of a six well plate
with puroR MEFs and regular ES media. The next day puromycin was added to the wells in a
range of concentrations and kept on the cells for the next two days. Following selection,
resistant cells were expanded, individual colonies were then re-plated into twenty-four well
plates and checked for the presence of the mutation by restriction enzyme digestion followed
by confirmation with sequencing.

EB formation: Embryoid bodies were formed using the hanging drop method(Behringer et al.,
2016; Dang et al., 2002). Briefly, ES cells were trypsinized, de-MEFed and then resuspended in
differentiation media (IMDM (Gibco 12440-053) supplemented with 20% FBS, pen/strep, and
glutamax. No LIF). Droplets containing approximately 180 cells were incubated for four days so
that spheres of differentiating cells could form. After the four days EBs were collected and
grown in suspension in non-adherent plates or used for subsequent experiments.

Beating heart assay: Beating clusters were differentiated from day 4 embryoid bodies (Boheler
et al., 2002; Hescheler et al., 1997). Individual EBs were transferred into individual wells of a
gelatinized 12 well plate. Cells were maintained in differentiation media throughout the
experiment. Wells were monitored for beating colonies for the next 15 days and were scored as
positive if there were any beating cells in that time period.

Western Blot
The antibodies that were used probe the Western blots were from Cell Signaling Technology
H3K27me3 (9733S) and H3K27me2 (9728S) as well as from EMD Millipore EZH2 (07-689).

RNA-seq
RNA was isolated from whole cells using the Nucleospin RNA kit from Macherey Nagel
(740955.50). We then depleted rRNA using the Ribozero Gold kit from Epicentre (RZG1224).
cDNA was synthesized from the purified RNA using the Superscript Vilo cDNA sequencing kit
from ThermoFisher (11754050) the libraries were assembled as previously described (Bowman
et al., 2013). Three replicates were completed for all experimental conditions except the Day 14
re-plated cells where two replicates were completed.

Methyltransferase Assay
Purified protein complexes were incubated at room temperature with a histone substrate at a
final concentration of 500nM and approximately 500nM radioactive SAM (Adenosyl-L-
methionine S-methyl^{3}H from PerkinElmer (NET155H250UC)) in methyltransferase buffer (10%
glycerol, 25mM HEPES PH 7.9, 2mM MgCl with 1mM DTT added fresh). Unless otherwise
specified all reactions ran for one hour before being stopped with the addition of 6x SDS buffer.
Samples were then run on an SDS page gel, coomassie stained and incubated in AMPLIFY
Amersham/GE (NAMP100) for 20 minutes. The gel was dried and exposed to film for a
minimum of 24 hours before developing.

Protein purification
The ORF for each member of the core PRC2 complex were cloned into the pFastBac1
baculovirus with Suz12 tagged with 3xFlag. Sf9 transfection with bacmid DNA and virus
amplification were performed essentially as described for the Bac-to-Bac Baculovirus
Expression System (ThermoFisher Scientific). Sf9 cells were maintained in Hyclone CCM3
(CCM3 liquid medium with L-glutamine, GE Healthcare Life Sciences SH30065.02)
supplemented with 50 U/ml Penicillin-Streptomycin (ThermoFisher Scientific, 15140-122). For
protein expression, 2x10^6 Sf9 cells/ml were infected at an MOI of approximately 10. Cells were
harvested 66 hours post infection by centrifugation at 5000xG for 15 minutes. Cells were lysed,
treated with DNAse1 then the complex was bound to M2 and the complex was eluted with flag
peptide.

CUT&RUN
CUT&RUN experiments were performed as described (Skene & Henikoff, 2017). The antibodies
we used were from Cell Signaling Technology H3K27me3 (9733S), EZH2 (5246S), SUZ12 (3737S)
and Bethyl laboratories RING1b (A302-869A). Libraries were constructed as described in the
RNA-seq section. Two replicates were completed and representative experiments are shown.

Alkaline Phosphotase Assay
Cells were stained according to the instructions in the Stemgent AP Staining Kit II from
Reprocell (00-0055). Images of stained wells were quantified using ImageJ FIJI software.

Bioinformatics Methods

RNA-seq data processing: All RNA sequencing reads were aligned to the mm10 genome using STAR v2.5.3 (Dobin et al., 2013). Gene annotations were obtained from Ensembl (Hunt et al., 2018) Genome browser tracks were generated using Homer v4.10.3 (Heinz et al., 2010) and visualized in IGV (Robinson et al., 2011). Reads in exons of Refseq annotated genes were counted using featureCounts v1.6.1(Liao et al., 2014). edgeR (Robinson et al., 2010) was used to normalize reads and calculate FDR using triplicates. All further calculations and figures were made using R v3.3.2 (Dessau & Pipper, 2008). For all heatmaps, standard z-scores calculations were made using average RPKMs of triplicates for each gene across all conditions. Genes were narrowed down to only PRC2 targets using H3K27me3 CUT&RUN data. Only genes changing in untreated or wild-type cells over the time-course were used for further analysis. The genes were filtered using cut-offs of at least 0.5 RPKM average expression at any timepoint, at least 1.5 fold-change and maximum 0.05 FDR while comparing Day 8 and Day 4 or Day 4 and Day 0. Genes were then clustered based on up or down-regulation in control cells and up or down-regulation or unchanged gene expression in mutant or treated cells.

For Fig. 5E, unsupervised clustering was performed using R's heatmap.2 function. CpG and GC levels were calculated using Homer's annotatePeaks function.

CUT&RUN data processing: All CUT&RUN reads were aligned to the mm10 genome using bowtie2 and filtered using samtools (Li et al., 2009) to keep uniquely aligned reads. Genome browser tracks were generated using Homer v4.10.3 and visualized in IGV. Peaks were called using Homer's findPeaks function for broad peaks. PRC2 target genes were identified using wild-type H3K27me3 peaks at least 1500 bp in length, 1rpkm in at least two time points and overlapping Refseq annotated genes (TSS+-5kb). Deeptools v3.3.0(Ramírez et al., 2014) was used to make average profile plots for Fig. 3E.

Published ChIP-seq data sources and processing: Published data for other canonical and non-canonical PRC2 components was downloaded from GEO as follows : H3K27me3 (GSM2282188, GSM2282191, GSM2282192) (Juan et al., 2016), EPOP (GSM2098943) (Beringer et al., 2016), Jarid2 (GSM491760) (Li et al., 2010), H3K4me1 (GSM1180178), H3K4me3 (GSM1180179), H3K9me3 (GSM1180180), H3K36me3 (GSM1180183) (Hon et al., 2014). FASTQ files were downloaded from GEO using sratools v2.9.1 (https://ncbi.github.io/sra-tools/). All reads were aligned to the mm10 genome using bowtie2 and filtered using samtools to keep uniquely aligned reads. Genome browser tracks were generated using Homer v4.10.3. Deeptools v3.3.0 was used to make average profile plots for Fig. 3C and E.

Figure 1.

Figure 2.

A.

B.

Methyltransferase Assay

PRC2 Ezh2^{WT}	PRC2 Ezh2^{681C}	PRC2 Ezh2^{722D}
1hr 6hr	1hr 6hr	1hr 6hr

Recombinant H3

C.

Western Blot

αH3K27me3

αH3K27me2

αEzh2

CJ7 Ezh2^{WT} CJ7 Eed$^{-/-}$ CJ7 Ezh2$^{681C-99}$

D.

ESCs

CJ7 *Ezh2*^{WT} CJ7 *Eed*$^{-/-}$ CJ7 *Ezh2*$^{681C-99}$ CJ7 *Ezh2*$^{681C-102}$

E.

Day 4 EBs

CJ7 *Ezh2*^{WT} CJ7 *Eed*$^{-/-}$ CJ7 *Ezh2*$^{681C-99}$ CJ7 *Ezh2*$^{681C-102}$

F.

Beating Heart Assay

p=0.74

p=0.008   p=0.02

Beating
Not Beating

561
562

Figure 3.

Figure 4.

A. Alkaline Phosphatase Images

CJ7 *Ezh2*WT CJ7 *Ezh2*681C-99 CJ7 *Ezh2*681C-102

Re-plated after 8 days
EB differentiation

Re-plated after 14 days
EB differentiation

B. Alkaline Phosphatase Quantification

Figure 5.

A.

B.

Alkaline Phosphatase Images

C.

Alkaline Phosphatase Quantification

D.

Beating Heart Assay

E.

14 Day re-plating RNA-seq samples
PRC2 target genes average expression

568   Figure Legends:
569
Figure 1
(A) Diagram of embryoid body (EB) formation time-course. (B) Western blot of cells treated
with GSK343 EZH2 inhibitor or DMSO control probed with H3K27me3 and EZH2 antibodies. (C)
54x magnification of day 4 EBs from cells treated with either DMSO control or GSK343. (D)
Screen shots from RNA-seq over the EB differentiation time-course from Mtase independent
(Sox3 and Tbx3) and Mtase dependent (Itgb7 and Wnt3) genes from cells treated with DMSO
(blue) or GSK343 (red). (E) Heatmap showing expression patterns of Mtase dependent and
independent genes in untreated (DMSO) and treated (GSK343) ES cells over development time-
course. Z-scores of average RPKMs of duplicates are shown for each gene across all samples.
Genes are separated based on if they were activated (top) or repressed (bottom) in untreated
cells over time as well as if they had a similar expression pattern in treated cells (Mtase
dependent genes) (left) or a different expression pattern (Mtase independent genes) (right).
The four groups are further clustered based on fold-changes over time in untreated cells.
583
Figure 2
(A) Diagram of *Ezh2* SET domain structure with the 681 residue circled and an alignment of SET
domains with the analogous residue highlighted in red adapted from (Antonysamy et al., 2013).
It is highly conserved across species and methyltransferases. (B) In vitro methyltransferase
assay with PRC2 comprised of WT or mutant EZH2, EED, SUZ12, and AEBP2. Recombinant H3 is
the substrate and a radioactive SAM was the methyl donor. Reactions progressed for 1 or 6
hours. (C) Western blot from WT, mutant and PRC2 knockout cells probed with H3K27me and
EZH2 antibodies. (D) 54x magnification of embryonic stem cells (ESCs) WT, 681C-99, 681C-102
and *Eed*[-/-] cells. (E) 54x magnification of embryoid bodies that have differentiated for 4 days
from WT, 681C-99, 681C-102 and *Eed*[-/-] cells. (F) Quantification of beating heart assay from WT,
681C-99 and 681C-102 cells that were differentiated as EBs for 4 days and then individually
plated in differentiation media. Blue shows the number of EBs that gave rise to beating cells
and Orange shows those that did not.
597
598
599
Figure 3
(A) Venn diagram showing overlap of H3K27me3 cut-and-run and published ChIP-seq data
peaks. (B) Venn diagrams showing overlap of target genes of H3K27me3 and other PRC2
components by cut-and-run. (C) Average profiles showing H3K27me3 cut-and-run versus IgG
control signal at Day0, 4 and 8 in wild-type and mutant cells. (D) Heatmap showing expression
patterns of Mtase dependent and independent genes in wild-type and mutant ES cells over
developmental time-course. Z-scores of average RPKMs of triplicates are shown for each gene
across all samples. Genes are separated based on if they were activated (left two panels) or
repressed (right two panels) in wild-type cells over time as well as if they had a similar
expression pattern in mutant cells (Mtase dependent genes) or a different expression pattern
(Mtase independent genes). The four groups are further clustered based on fold-changes over
time in untreated cells. (E) Average profiles showing H3K27me3 binding in wild-type and

mutant cells by cut-and-run at Mtase dependent or independent and developmentally
repressed PRC2 target genes over the time course.
614
Figure 4
(A) Alkaline phosphatase staining from WT, 681C-99 and 681C-102 cells that had been
differentiated for 8 or 14 days and then transferred into ESC conditions for five days before
staining. ESCs stain bright pink in this assay. (B) Quantification of alkaline phosphatase staining
from re-plated cells with standard deviation error bars. Samples with significant difference
(p<0.05) from WT cells marked with an asterisk.
621
622
Figure 5
(A) Diagram showing the drug treatment plan for differentiation and re-plating of cells. Red
circles indicate the cells used for the beating heart assay (D), Purple are shown in images in (B)
as well as quantified in (C) and blue circles highlight samples only quantified in (C). (B) Alkaline
phosphatase staining from treated cells re-plated after 8 or 14 days of EB differentiation. Only
cells that start in GSK343 treatment is stain pink indicating that the cells have reverted to an ES
phenotype. (C) Quantification of alkaline phosphatase staining from cells treated with DMSO or
GSK343. Time of the treatment switch is indicated in the legend. Samples with significant
differences from those continuously treated with DMSO (p<0.05) are marked with an asterisk.
(D) Quantification of beating heart assay with GSK343 and DMSO treated cells. Cells were
differentiated as EBs for 4 days and then plated into differentiation media. Blue indicates the
number of EBs that formed beating cells and orange indicates the number that did not. (E)
Unsupervised clustering of RNA-seq average expression of cells re-plated for 5 days after 14
days of EB formation. Cells that were treated with GSK343 have gene expression profiles that
cluster with average expression from WT ESCs.
638

Supp. Fig. 1

A.

GSK343 Titration 3 day treatment

Western Blot

αH3K27me3

αEzh2

0 100nM 300nM 1µM 2µM 4µM

GSK343

Supp. Fig. 2

A.

In vitro methyltransferase assay

204   42   Pixel intensity

0.5   5   2.5   1.25 0.625

WT   681C

C.

Methyltransferase Assay

31K
24K
17K
12K

WT 681A 681C   WT 681A 681C

Recombinant H3   Core Histone

B.

In vitro methyltransferase assay

112   12.5 Pixel intensity

0.1   1   1   2   4   7

WT   681C

D.

Methyltransferase Assay

15.5 17 21 20 21 22 27.5 41   Pixel intensity

0.5ug
WT complex

15   30   45   60   75   90 105 120

Reaction time (min)

Supp. Fig. 3

A.

Western Blot

αEzh2

αH3K27me3

CJ7 Ezh2^wt

CJ7 Ezh2^681C-99

CJ7 Ezh2^-/-

CJ7 Eed^-/-

B.

Ezh2^681C-102 mutant

αEzh2

αH3K27me3

αH3K27me2

Supp. Fig. 4

A.

B.

Supp. Fig. 5

Log2 fold-change as compared to WT comparisons between drug treated and mutant cells

A. Log2 fold-change over WT ESC

B. Log2 fold-change over WT ESC

Supp. Fig. 6.

A. PRC2 developmentally repressed genes

ChIP-seq at PRC2 developmentally repressed genes

Jarid2 | EPOP | H3K9me3

H3K36me3 | H3K4me1 | H3K4me3

— Mtase independent
— Mtase dependent

B. PRC2 developmentally repressed genes

ESC expression level | Gene size

CpG percentage | GC percentage

C. PRC2 developmentally repressed genes

Mtase independent | Mtase dependent

CJ7 Ezh2^{WT} CJ7 Ezh2$^{681C-99}$
D0 D4 D8 D0 D4 D8

Similar starting ESC expression

59 | 45

Different starting ESC expression

123 | 121

z-scores of RPKM per

D. H3K27me3 Cut and Run D8EB

Mtase independent | Mtase dependent

CJ7 Ezh2^{WT}

CJ7 Ezh2$^{681C-99}$

— Similar ESC expression
— Different ESC expression

Supp. Fig. 7.

A.
Western Blot

αEzh2

αH3K27me3

αH3K27me2

DMSO D0 GSK343 D0 DMSO D8 GSK343 D8

B.   14 Day re-plating RNA-seq samples
PRC2 target genes

-2   0   2
z-scores
of RPKM

GSK/DMSO/1
GSK/DMSO/2
GSK/GSK/1
DMSO/DMSO/1
GSK/GSK/2
WT-ESC/3
WT-ESC/1
WT-ESC/2
DMSO/GSK/2
GSK/DMSO/3
DMSO/GSK/1
DMSO/DMSO/3
DMSO/GSK/3
DMSO/DMSO/2

C.   14 Day re-plating RNA-seq samples
PRC2 developmentally repressed genes

Mtase independent   Mtase dependent

-2   0   2
z-scores
of RPKM

DMSO-DMSO
DMSO-GSK343
GSK343-DMSO
GSK343-GSK343

DMSO-DMSO
DMSO-GSK343
GSK343-DMSO
GSK343-GSK343

644
645

646 Supplemental Figure Legends
647
648 Supp. Fig. 1
649 (A) Western blot of WT CJ7 cells treated for three days with an increasing concentration of
650 GSK343 probed with antibodies to H3K27me3 and EZH2. The 4uM concentration was non-toxic
651 to cells and yielded a loss of H3K27me3 signal and so this concentration was used for all
652 experiments.
653
Supp. Fig. 2
(A,B) In vitro methyltransferase assay comparing EZH2-WT and EZH2-681C. Concentration of
the PRC2 complex is indicated below the image with pixel intensity indicated above. (C) In vitro
methyltransferase assay with WT, 681A and 681C complexes with both recombinant histones
that do not have any covalent modifications and core histones that do as the substrate. With
both substrates, the mutant complexes did not show signal. (D) In vitro methyltransferase assay
with WT complex over time. All other methyltransferase assays shown here were done for 60
minutes which is well within the linear range.
662
663 Supp. Fig. 3
664 (A) Western blot showing the EZH2 and H3K27me3 signal for WT, EZH2 mutant and PRC2 KO
665 cells. (B) Western blot showing the EZH2, H3K27me3 and H3K27me2 signal of EZH2$^{618C-102}$
666 mutant cells.
667
668 Supp. Fig. 4
669 (A) Comparisons between Ezh2$^{618C-99}$ and Ezh2$^{681C-102}$ RNA-seq over the differentiation time-
670 course. The two mutants are nearly identical at all time points. (B) Unsupervised clustering of
671 the mutant RNA-seq showing that the day of differentiation is more relevant predictor of
672 similarity rather than the mutant strain.
673
674 Supp. Fig. 5
675 (A) Unsupervised clustering showing the fold change over WT ESC of GSK343 treated or
676 Ezh2681C-99 cells. The day of differentiation is the stronger predictor of clustering. (B) Direct
677 comparison of mutant and drug treated cells RNA-seq fold change over WT gene expression.
678
Supp. Fig. 6
(A) Average profiles showing binding of other PRC2 components and histone marks in wild-type
and mutant cells at Mtase dependent or independent and developmentally repressed PRC2
target genes over the time course. (B) Boxplots showing comparisons of different
characteristics of Mtase dependent and independent, developmentally repressed PRC2 target
genes in wild-type and mutant cells. (C) Heatmaps showing developmentally repressed genes
from Fig 3(D), separated based on expression levels of wild-type and mutant ES cells at Day 0 of
the time-course. Genes with similar expression levels at Day 0 in wild-type and mutant (within
1.2 fold) are on the top and those with different starting expression levels (more than 1.2 fold)
are at the bottom. (D) Average profiles showing H3K27me3 binding in wild-type and mutant
cells by cut-and-run at Mtase dependent or independent and developmentally repressed PRC2

target genes that have either different (more than 1.2 fold) or similar (within 1.2 fold)
expression over the time course.
692
Supp. Fig. 7
(A) Western blot of CJ7 cells treated with DMSO or GSK343 showing the (levels of Ezh2,
H3K27me3 and H3K27me2 at Day 0 and Day 8 or EB formation. B) Unsupervised clustering of
PRC2 target genes in re-plated cells and wild-type ES-cells. Untreated or drug treated cells were
replated at 14 days in DMSO or GSK343. Z-scores of average RPKMs of triplicates (duplicates for
GSK-GSK) are shown for each gene across all samples. (C) Heatmaps showing comparison of
treated and untreated cells, re-plated in DMSO or GSK343 for previously identified Mtase
dependent and independent genes. Z-scores of average RPKMs of triplicates (duplicates for
GSK-GSK) are shown for each gene across all samples.
702
703
704

705   Abdel Raouf, S. M., Ibrahim, T. R., Abdelaziz, L. A., Farid, M. I., & Mohamed, S. Y. (2019, Dec 10).
706   Prognostic Value of TWIST1 and EZH2 Expression in Colon Cancer. *J Gastrointest Cancer*.
707   https://doi.org/10.1007/s12029-019-00344-4
708
709   Ahmed, A., Wang, T., & Delgado-Olguin, P. (2018). Ezh2 is not required for cardiac regeneration
710   in neonatal mice. *PLoS One, 13*(2), e0192238.
711   https://doi.org/10.1371/journal.pone.0192238
712
713   Ai, S., Peng, Y., Li, C., Gu, F., Yu, X., Yue, Y., Ma, Q., Chen, J., Lin, Z., Zhou, P., Xie, H., Prendiville,
714   T. W., Zheng, W., Liu, Y., Orkin, S. H., Wang, D. Z., Yu, J., Pu, W. T., & He, A. (2017, Apr
715   10). EED orchestration of heart maturation through interaction with HDACs is
716   H3K27me3-independent. *Elife, 6*. https://doi.org/10.7554/eLife.24570
717
718   Aloia, L., Di Stefano, B., & Di Croce, L. (2013, Jun). Polycomb complexes in stem cells and
719   embryonic development. *Development, 140*(12), 2525-2534.
720   https://doi.org/10.1242/dev.091553
721
722   Antonysamy, S., Condon, B., Druzina, Z., Bonanno, J. B., Gheyi, T., Zhang, F., MacEwan, I., Zhang,
723   A., Ashok, S., Rodgers, L., Russell, M., & Gately Luz, J. (2013). Structural context of
724   disease-associated mutations and putative mechanism of autoinhibition revealed by X-
725   ray crystallographic analysis of the EZH2-SET domain. *PLoS One, 8*(12), e84147.
726   https://doi.org/10.1371/journal.pone.0084147
727
Basheer, F., Giotopoulos, G., Meduri, E., Yun, H., Mazan, M., Sasca, D., Gallipoli, P., Marando, L.,
Gozdecka, M., Asby, R., Sheppard, O., Dudek, M., Bullinger, L., Dohner, H., Dillon, R.,
Freeman, S., Ottmann, O., Burnett, A., Russell, N., Papaemmanuil, E., Hills, R., Campbell,
P., Vassiliou, G. S., & Huntly, B. J. P. (2019, Apr 1). Contrasting requirements during
disease evolution identify EZH2 as a therapeutic target in AML. *J Exp Med, 216*(4), 966-
981. https://doi.org/10.1084/jem.20181276

Behringer, R., Gertsenstein, M., Nagy, K. V., & Nagy, A. (2016, Dec 1). Differentiating Mouse
Embryonic Stem Cells into Embryoid Bodies by Hanging-Drop Cultures. *Cold Spring Harb*
Protoc, 2016(12). https://doi.org/10.1101/pdb.prot092429
738

739 Beringer, M., Pisano, P., Di Carlo, V., Blanco, E., Chammas, P., Vizán, P., Gutiérrez, A., Aranda, S.,
740   Payer, B., Wierer, M., & Di Croce, L. (2016, Nov 17). EPOP Functionally Links Elongin and
741   Polycomb in Pluripotent Stem Cells. *Mol Cell, 64*(4), 645-658.
742   https://doi.org/10.1016/j.molcel.2016.10.018
743

744 Bernstein, E., Duncan, E. M., Masui, O., Gil, J., Heard, E., & Allis, C. D. (2006, Apr). Mouse
745   polycomb proteins bind differentially to methylated histone H3 and RNA and are
746   enriched in facultative heterochromatin. *Mol Cell Biol, 26*(7), 2560-2569.
747   https://doi.org/10.1128/mcb.26.7.2560-2569.2006
748

749 Boheler, K. R., Czyz, J., Tweedie, D., Yang, H. T., Anisimov, S. V., & Wobus, A. M. (2002, Aug 9).
750   Differentiation of pluripotent embryonic stem cells into cardiomyocytes. *Circ Res, 91*(3),
751   189-201. https://doi.org/10.1161/01.res.0000027865.61704.32
752

753 Bohm, J., Muenzner, J. K., Caliskan, A., Ndreshkjana, B., Erlenbach-Wunsch, K., Merkel, S.,
754   Croner, R., Rau, T. T., Geppert, C. I., Hartmann, A., Roehe, A. V., & Schneider-Stock, R.
755   (2019, Sep). Loss of enhancer of zeste homologue 2 (EZH2) at tumor invasion front is
756   correlated with higher aggressiveness in colorectal cancer cells. *J Cancer Res Clin Oncol,*
757   *145*(9), 2227-2240. https://doi.org/10.1007/s00432-019-02977-1
758

759 Bowman, S. K., Simon, M. D., Deaton, A. M., Tolstorukov, M., Borowsky, M. L., & Kingston, R. E.
760   (2013, Jul 9). Multiplexed Illumina sequencing libraries from picogram quantities of
761   DNA. *BMC Genomics, 14*, 466. https://doi.org/10.1186/1471-2164-14-466
762

Bradley, W. D., Arora, S., Busby, J., Balasubramanian, S., Gehling, V. S., Nasveschuk, C. G.,
Vaswani, R. G., Yuan, C. C., Hatton, C., Zhao, F., Williamson, K. E., Iyer, P., Méndez, J.,
Campbell, R., Cantone, N., Garapaty-Rao, S., Audia, J. E., Cook, A. S., Dakin, L. A.,
Albrecht, B. K., Harmange, J. C., Daniels, D. L., Cummings, R. T., Bryant, B. M., Normant,
E., & Trojer, P. (2014, Nov 20). EZH2 inhibitor efficacy in non-Hodgkin's lymphoma does
not require suppression of H3K27 monomethylation. *Chem Biol, 21*(11), 1463-1475.
https://doi.org/10.1016/j.chembiol.2014.09.017
770

771 Bremer, S. C. B., Conradi, L. C., Mechie, N. C., Amanzada, A., Mavropoulou, E., Kitz, J., Ghadimi,
772   M., Ellenrieder, V., Strobel, P., Hessmann, E., Gaedcke, J., & Bohnenberger, H. (2019,
773   Nov 6). Enhancer of Zeste Homolog 2 in Colorectal Cancer Development and
774   Progression. *Digestion*, 1-9. https://doi.org/10.1159/000504093
775

776 Chamberlain, S. J., Yee, D., & Magnuson, T. (2008, Jun). Polycomb repressive complex 2 is
777   dispensable for maintenance of embryonic stem cell pluripotency. *Stem Cells, 26*(6),
778   1496-1505. https://doi.org/10.1634/stemcells.2008-0102
779

780 Collinson, A., Collier, A. J., Morgan, N. P., Sienerth, A. R., Chandra, T., Andrews, S., & Rugg-Gunn,
781   P. J. (2016, Dec 6). Deletion of the Polycomb-Group Protein EZH2 Leads to Compromised
782   Self-Renewal and Differentiation Defects in Human Embryonic Stem Cells. *Cell Rep,*
783   *17*(10), 2700-2714. https://doi.org/10.1016/j.celrep.2016.11.032
784

785 Cyrus, S., Burkardt, D., Weaver, D. D., & Gibson, W. T. (2019, Dec). PRC2-complex related
786   dysfunction in overgrowth syndromes: A review of EZH2, EED, and SUZ12 and their
787   syndromic phenotypes. *Am J Med Genet C Semin Med Genet, 181*(4), 519-531.
788   https://doi.org/10.1002/ajmg.c.31754
789

790 Dang, S. M., Kyba, M., Perlingeiro, R., Daley, G. Q., & Zandstra, P. W. (2002, May 20). Efficiency
791   of embryoid body formation and hematopoietic development from embryonic stem
792   cells in different culture systems. *Biotechnol Bioeng, 78*(4), 442-453.
793   https://doi.org/10.1002/bit.10220
794

795 Deng, Y., Chen, X., Huang, C., Chen, G., Chen, F., Lu, J., Shi, X., He, C., Zeng, Z., Qiu, Y., Chen, J.,
796   Lin, R., Chen, Y., & Chen, J. (2019). EZH2/Bcl-2 Coexpression Predicts Worse Survival in
797   Diffuse Large B-cell Lymphomas and Demonstrates Poor Efficacy to Rituximab in
798   Localized Lesions. *J Cancer, 10*(9), 2006-2017. https://doi.org/10.7150/jca.29807
799

800 Dessau, R. B., & Pipper, C. B. (2008, Jan 28). ["R"--project for statistical computing]. *Ugeskr*
801   *Laeger, 170*(5), 328-330. (R--en programpakke til statistisk databehandling og grafik.)
802

803 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &
804   Gingeras, T. R. (2013, Jan 1). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics,*
805   *29*(1), 15-21. https://doi.org/10.1093/bioinformatics/bts635
806

807 Dou, D., Ge, X., Wang, X., Xu, X., Zhang, Z., Seng, J., Cao, Z., Gu, Y., & Han, M. (2019). EZH2
808   Contributes To Cisplatin Resistance In Breast Cancer By Epigenetically Suppressing miR-
809   381 Expression. *Onco Targets Ther, 12*, 9627-9637. https://doi.org/10.2147/ott.S214104
810

811 Ferguson, J., Devarajan, M., DiNuoscio, G., Saiakhova, A., Liu, C. F., Lefebvre, V., Scacheri, P. C.,
812   & Atit, R. P. (2018, Feb 2). PRC2 Is Dispensable in Vivo for β-Catenin-Mediated
813   Repression of Chondrogenesis in the Mouse Embryonic Cranial Mesenchyme. *G3*
814   *(Bethesda), 8*(2), 491-503. https://doi.org/10.1534/g3.117.300311
815

816 Fioravanti, R., Stazi, G., Zwergel, C., Valente, S., & Mai, A. (2018, Dec). Six Years (2012-2018) of
817   Researches on Catalytic EZH2 Inhibitors: The Boom of the 2-Pyridone Compounds. *Chem*
818   *Rec, 18*(12), 1818-1832. https://doi.org/10.1002/tcr.201800091
819

Fraineau, S., Palii, C. G., McNeill, B., Ritso, M., Shelley, W. C., Prasain, N., Chu, A., Vion, E., Rieck,
K., Nilufar, S., Perkins, T. J., Rudnicki, M. A., Allan, D. S., Yoder, M. C., Suuronen, E. J., &
Brand, M. (2017, Nov 14). Epigenetic Activation of Pro-angiogenic Signaling Pathways in
Human Endothelial Progenitors Increases Vasculogenesis. *Stem Cell Reports, 9*(5), 1573-
1587. https://doi.org/10.1016/j.stemcr.2017.09.009

Harris, J. (2018). Partial Clinical Hold on Tazemetostat Trials Lifted by FDA [News article].
Targeted Oncology.

Healy, E., Mucha, M., Glancy, E., Fitzpatrick, D. J., Conway, E., Neikes, H. K., Monger, C., Van
Mierlo, G., Baltissen, M. P., Koseki, Y., Vermeulen, M., Koseki, H., & Bracken, A. P. (2019,
Nov 7). PRC2.1 and PRC2.2 Synergize to Coordinate H3K27 Trimethylation. *Mol Cell,*
76(3), 437-452.e436. https://doi.org/10.1016/j.molcel.2019.08.012

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh,
H., & Glass, C. K. (2010, May 28). Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. *Mol Cell, 38*(4), 576-589. https://doi.org/10.1016/j.molcel.2010.05.004

Hescheler, J., Fleischmann, B. K., Lentini, S., Maltsev, V. A., Rohwedel, J., Wobus, A. M., &
Addicks, K. (1997, Nov). Embryonic stem cells: a model to study structural and functional
properties in cardiomyogenesis. *Cardiovasc Res, 36*(2), 149-162.
https://doi.org/10.1016/s0008-6363(97)00193-4

Hon, G. C., Song, C. X., Du, T., Jin, F., Selvaraj, S., Lee, A. Y., Yen, C. A., Ye, Z., Mao, S. Q., Wang,
B. A., Kuan, S., Edsall, L. E., Zhao, B. S., Xu, G. L., He, C., & Ren, B. (2014, Oct 23). 5mC
oxidation by Tet2 modulates enhancer activity and timing of transcriptome
reprogramming during differentiation. *Mol Cell, 56*(2), 286-297.
https://doi.org/10.1016/j.molcel.2014.08.026

Hunt, S. E., McLaren, W., Gil, L., Thormann, A., Schuilenburg, H., Sheppard, D., Parton, A.,
Armean, I. M., Trevanion, S. J., Flicek, P., & Cunningham, F. (2018, Jan 1). Ensembl
variation resources. *Database (Oxford), 2018*. https://doi.org/10.1093/database/bay119

Italiano, A., Soria, J. C., Toulmonde, M., Michot, J. M., Lucchesi, C., Varga, A., Coindre, J. M.,
Blakemore, S. J., Clawson, A., Suttle, B., McDonald, A. A., Woodruff, M., Ribich, S.,
Hedrick, E., Keilhack, H., Thomson, B., Owa, T., Copeland, R. A., Ho, P. T. C., & Ribrag, V.
(2018, May). Tazemetostat, an EZH2 inhibitor, in relapsed or refractory B-cell non-
Hodgkin lymphoma and advanced solid tumours: a first-in-human, open-label, phase 1
study. *Lancet Oncol, 19*(5), 649-659. https://doi.org/10.1016/s1470-2045(18)30145-1

Jain, P., & Di Croce, L. (2016, May). Mutations and deletions of PRC2 in prostate cancer.
Bioessays, 38(5), 446-454. https://doi.org/10.1002/bies.201500162

Juan, A. H., Wang, S., Ko, K. D., Zare, H., Tsai, P. F., Feng, X., Vivanco, K. O., Ascoli, A. M.,
Gutierrez-Cruz, G., Krebs, J., Sidoli, S., Knight, A. L., Pedersen, R. A., Garcia, B. A.,
Casellas, R., Zou, J., & Sartorelli, V. (2016, Oct 25). Roles of H3K27me2 and H3K27me3
Examined during Fate Specification of Embryonic Stem Cells. *Cell Rep, 17*(5), 1369-1382.
https://doi.org/10.1016/j.celrep.2016.09.087

Karlowee, V., Amatya, V. J., Takayasu, T., Takano, M., Yonezawa, U., Takeshima, Y., Sugiyama,
K., Kurisu, K., & Yamasaki, F. (2019). Immunostaining of Increased Expression of
Enhancer of Zeste Homolog 2 (EZH2) in Diffuse Midline Glioma H3K27M-Mutant
Patients with Poor Survival. *Pathobiology, 86*(2-3), 152-161.
https://doi.org/10.1159/000496691

Kondo, Y. (2014, Nov). Targeting histone methyltransferase EZH2 as cancer treatment. *J
Biochem, 156*(5), 249-257. https://doi.org/10.1093/jb/mvu054

Krill, L., Deng, W., Eskander, R., Mutch, D., Zweizig, S., Hoang, B., Ioffe, O., Randall, L., Lankes,
H., Miller, D. S., & Birrer, M. (2020, Feb). Overexpression of enhance of Zeste homolog 2
(EZH2) in endometrial carcinoma: An NRG Oncology/Gynecologic Oncology Group Study.
Gynecol Oncol, 156(2), 423-429. https://doi.org/10.1016/j.ygyno.2019.12.003

Lavarone, E., Barbieri, C. M., & Pasini, D. (2019, Apr 11). Dissecting the role of H3K27
acetylation and methylation in PRC2 mediated control of cellular identity. *Nat Commun,
10*(1), 1679. https://doi.org/10.1038/s41467-019-09624-w

Lewis, E., & Mislove, R. (1947). New mutants report. *Drosophila Information Service, 21*, 69.

Li, G., Margueron, R., Ku, M., Chambon, P., Bernstein, B. E., & Reinberg, D. (2010, Feb 15). Jarid2
and PRC2, partners in regulating gene expression. *Genes Dev, 24*(4), 368-380.
https://doi.org/10.1101/gad.1886410

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., &
Durbin, R. (2009, Aug 15). The Sequence Alignment/Map format and SAMtools.
Bioinformatics, 25(16), 2078-2079. https://doi.org/10.1093/bioinformatics/btp352

Liao, Y., Smyth, G. K., & Shi, W. (2014, Apr 1). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics, 30*(7), 923-
930. https://doi.org/10.1093/bioinformatics/btt656

Liu, T. P., Lo, H. L., Wei, L. S., Hsiao, H. H., & Yang, P. M. (2015, Feb). S-Adenosyl-L-methionine-
competitive inhibitors of the histone methyltransferase EZH2 induce autophagy and
enhance drug sensitivity in cancer cells. *Anticancer Drugs, 26*(2), 139-147.
https://doi.org/10.1097/cad.0000000000000166

Lue, J. K., & Amengual, J. E. (2018, Oct). Emerging EZH2 Inhibitors and Their Application in
Lymphoma. *Curr Hematol Malig Rep, 13*(5), 369-382. https://doi.org/10.1007/s11899-
018-0466-6
910
911 Margueron, R., Li, G., Sarma, K., Blais, A., Zavadil, J., Woodcock, C. L., Dynlacht, B. D., &
912   Reinberg, D. (2008, Nov 21). Ezh1 and Ezh2 maintain repressive chromatin through
913   different mechanisms. *Mol Cell, 32*(4), 503-518.
914   https://doi.org/10.1016/j.molcel.2008.11.004
915
916 Margueron, R., & Reinberg, D. (2011, Jan 20). The Polycomb complex PRC2 and its mark in life.
917   *Nature, 469*(7330), 343-349. https://doi.org/10.1038/nature09784
918
919 Matsubara, T., Toyokawa, G., Takada, K., Kinoshita, F., Kozuma, Y., Akamine, T., Shimokawa, M.,
920   Haro, A., Osoegawa, A., Tagawa, T., & Mori, M. (2019). The association and prognostic
921   impact of enhancer of zeste homologue 2 expression and epithelial-mesenchymal
922   transition in resected lung adenocarcinoma. *PLoS One, 14*(5), e0215103.
923   https://doi.org/10.1371/journal.pone.0215103
924
925 Mechaal, A., Menif, S., Abbes, S., & Safra, I. (2019, Sep). EZH2, new diagnosis and prognosis
926   marker in acute myeloid leukemia patients. *Adv Med Sci, 64*(2), 395-401.
927   https://doi.org/10.1016/j.advms.2019.07.002
928
929 Nakagawa, M., Koyanagi, M., Tanabe, K., Takahashi, K., Ichisaka, T., Aoi, T., Okita, K., Mochiduki,
930   Y., Takizawa, N., & Yamanaka, S. (2008, Jan). Generation of induced pluripotent stem
931   cells without Myc from mouse and human fibroblasts. *Nat Biotechnol, 26*(1), 101-106.
932   https://doi.org/10.1038/nbt1374
933
934 O'Geen, H., Ren, C., Nicolet, C. M., Perez, A. A., Halmai, J., Le, V. M., Mackay, J. P., Farnham, P.
935   J., & Segal, D. J. (2017, Sep 29). dCas9-based epigenome editing suggests acquisition of
936   histone methylation is not sufficient for target gene repression. *Nucleic Acids Res,
937   45*(17), 9901-9916. https://doi.org/10.1093/nar/gkx578
938
939 Pasini, D., Bracken, A. P., Hansen, J. B., Capillo, M., & Helin, K. (2007, May). The polycomb group
940   protein Suz12 is required for embryonic stem cell differentiation. *Mol Cell Biol, 27*(10),
941   3769-3779. https://doi.org/10.1128/mcb.01432-06
942
943 Pasini, D., Bracken, A. P., Jensen, M. R., Lazzerini Denchi, E., & Helin, K. (2004, Oct 13). Suz12 is
944   essential for mouse development and for EZH2 histone methyltransferase activity. *Embo
945   j, 23*(20), 4061-4071. https://doi.org/10.1038/sj.emboj.7600402
946
Pereira, C. F., Piccolo, F. M., Tsubouchi, T., Sauer, S., Ryan, N. K., Bruno, L., Landeira, D., Santos,
J., Banito, A., Gil, J., Koseki, H., Merkenschlager, M., & Fisher, A. G. (2010, Jun 4). ESCs
require PRC2 to direct the successful reprogramming of differentiated cells toward
pluripotency. *Cell Stem Cell, 6*(6), 547-556. https://doi.org/10.1016/j.stem.2010.04.013

Qi, W., Chan, H., Teng, L., Li, L., Chuai, S., Zhang, R., Zeng, J., Li, M., Fan, H., Lin, Y., Gu, J.,
Ardayfio, O., Zhang, J. H., Yan, X., Fang, J., Mi, Y., Zhang, M., Zhou, T., Feng, G., Chen, Z.,
Li, G., Yang, T., Zhao, K., Liu, X., Yu, Z., Lu, C. X., Atadja, P., & Li, E. (2012, Dec 26).
Selective inhibition of Ezh2 by a small molecule inhibitor blocks tumor cells proliferation.
Proc Natl Acad Sci U S A, 109(52), 21360-21365.
https://doi.org/10.1073/pnas.1210371110

Rai, A. N., Vargas, M. L., Wang, L., Andersen, E. F., Miller, E. L., & Simon, J. A. (2013, Dec).
Elements of the polycomb repressor SU(Z)12 needed for histone H3-K27 methylation,
the interface with E(Z), and in vivo function. *Mol Cell Biol, 33*(24), 4844-4856.
https://doi.org/10.1128/mcb.00307-13

Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A., & Manke, T. (2014, Jul). deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Res, 42*(Web Server issue),
W187-191. https://doi.org/10.1093/nar/gku365

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &
Mesirov, J. P. (2011, Jan). Integrative genomics viewer. *Nat Biotechnol, 29*(1), 24-26.
https://doi.org/10.1038/nbt.1754

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010, Jan 1). edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics, 26*(1),
139-140. https://doi.org/10.1093/bioinformatics/btp616

Schuettengruber, B., Bourbon, H. M., Di Croce, L., & Cavalli, G. (2017, Sep 21). Genome
Regulation by Polycomb and Trithorax: 70 Years and Counting. *Cell, 171*(1), 34-57.
https://doi.org/10.1016/j.cell.2017.08.002

Shan, Y., Liang, Z., Xing, Q., Zhang, T., Wang, B., Tian, S., Huang, W., Zhang, Y., Yao, J., Zhu, Y.,
Huang, K., Liu, Y., Wang, X., Chen, Q., Zhang, J., Shang, B., Li, S., Shi, X., Liao, B., Zhang,
C., Lai, K., Zhong, X., Shu, X., Wang, J., Yao, H., Chen, J., Pei, D., & Pan, G. (2017, Sep 22).
PRC2 specifies ectoderm lineages and maintains pluripotency in primed but not naïve
ESCs. *Nat Commun, 8*(1), 672. https://doi.org/10.1038/s41467-017-00668-4

Shen, X., Kim, W., Fujiwara, Y., Simon, M. D., Liu, Y., Mysliwiec, M. R., Yuan, G. C., Lee, Y., &
Orkin, S. H. (2009, Dec 24). Jumonji modulates polycomb activity and self-renewal
versus differentiation of stem cells. *Cell, 139*(7), 1303-1314.
https://doi.org/10.1016/j.cell.2009.12.003

Shen, X., Liu, Y., Hsu, Y. J., Fujiwara, Y., Kim, J., Mao, X., Yuan, G. C., & Orkin, S. H. (2008, Nov
21). EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2 in
maintaining stem cell identity and executing pluripotency. *Mol Cell, 32*(4), 491-502.
https://doi.org/10.1016/j.molcel.2008.10.016

Shi, B., Behrens, C., Vaghani, V., Riquelme, E. M., Rodriguez-Canales, J., Kadara, H., Lin, H., Lee,
J., Liu, H., Wistuba, I., & Simon, G. (2019, Oct). Oncogenic enhancer of zeste homolog 2
is an actionable target in patients with non-small cell lung cancer. *Cancer Med, 8*(14),
6383-6392. https://doi.org/10.1002/cam4.1855

1000

1001 Skene, P. J., & Henikoff, S. (2017, Jan 16). An efficient targeted nuclease strategy for high-
1002 resolution mapping of DNA binding sites. *Elife, 6*. https://doi.org/10.7554/eLife.21856

1003

1004 Takahashi, K., & Yamanaka, S. (2006, Aug 25). Induction of pluripotent stem cells from mouse
1005 embryonic and adult fibroblast cultures by defined factors. *Cell, 126*(4), 663-676.
1006 https://doi.org/10.1016/j.cell.2006.07.024

1007

1008 Thornton, S. R., Butty, V. L., Levine, S. S., & Boyer, L. A. (2014). Polycomb Repressive Complex 2
1009 regulates lineage fidelity during embryonic stem cell differentiation. *PLoS One, 9*(10),
1010 e110498. https://doi.org/10.1371/journal.pone.0110498

1011

1012 Tian, Z., Li, Z., Zhu, Y., Meng, L., Liu, F., Sang, M., & Wang, G. (2019, Aug). Hypermethylation-
1013 mediated inactivation of miR-124 predicts poor prognosis and promotes tumor growth
1014 at least partially through targeting EZH2/H3K27me3 in ESCC. *Clin Exp Metastasis, 36*(4),
1015 381-391. https://doi.org/10.1007/s10585-019-09974-1

1016

1017 van Mierlo, G., Veenstra, G. J. C., Vermeulen, M., & Marks, H. (2019, Aug). The Complexity of
1018 PRC2 Subcomplexes. *Trends Cell Biol, 29*(8), 660-671.
1019 https://doi.org/10.1016/j.tcb.2019.05.004

1020

1021 Wasenang, W., Puapairoj, A., Settasatian, C., Proungvitaya, S., & Limpaiboon, T. (2019, Jul).
1022 Overexpression of polycomb repressive complex 2 key components EZH2/SUZ12/EED as
1023 an unfavorable prognostic marker in cholangiocarcinoma. *Pathol Res Pract, 215*(7),
1024 152451. https://doi.org/10.1016/j.prp.2019.152451

1025

Wassef, M., Luscan, A., Aflaki, S., Zielinski, D., Jansen, P., Baymaz, H. I., Battistella, A., Kersouani,
C., Servant, N., Wallace, M. R., Romero, P., Kosmider, O., Just, P. A., Hivelin, M., Jacques,
S., Vincent-Salomon, A., Vermeulen, M., Vidaud, M., Pasmant, E., & Margueron, R.
(2019, Mar 26). EZH1/2 function mostly within canonical PRC2 and exhibit proliferation-
dependent redundancy that shapes mutational signatures in cancer. *Proc Natl Acad Sci
U S A, 116*(13), 6075-6080. https://doi.org/10.1073/pnas.1814634116

1032

1033 Wu, X., Scott, H., Carlsson, S. V., Sjoberg, D. D., Cerundolo, L., Lilja, H., Prevo, R., Rieunier, G.,
1034 Macaulay, V., Higgins, G. S., Verrill, C. L., Lamb, A. D., Cunliffe, V. T., Bountra, C., Hamdy,
1035 F. C., & Bryant, R. J. (2019, Jul). Increased EZH2 expression in prostate cancer is
1036 associated with metastatic recurrence following external beam radiotherapy. *Prostate,
1037 79*(10), 1079-1089. https://doi.org/10.1002/pros.23817

1038

1039 Xu, B., On, D. M., Ma, A., Parton, T., Konze, K. D., Pattenden, S. G., Allison, D. F., Cai, L.,
1040   Rockowitz, S., Liu, S., Liu, Y., Li, F., Vedadi, M., Frye, S. V., Garcia, B. A., Zheng, D., Jin, J.,
1041   & Wang, G. G. (2015, Jan 8). Selective inhibition of EZH2 and EZH1 enzymatic activity by
1042   a small molecule suppresses MLL-rearranged leukemia. *Blood, 125*(2), 346-357.
1043   https://doi.org/10.1182/blood-2014-06-581082
1044
1045 Xu, H., Zhang, L., Qian, X., Zhou, X., Yan, Y., Zhou, J., Ge, W., Albahde, M., & Wang, W. (2019,
1046   Oct). GSK343 induces autophagy and downregulates the AKT/mTOR signaling pathway
1047   in pancreatic cancer cells. *Exp Ther Med, 18*(4), 2608-2616.
1048   https://doi.org/10.3892/etm.2019.7845
1049
1050 Yamagishi, M., & Uchimaru, K. (2017, Sep). Targeting EZH2 in cancer therapy. *Curr Opin Oncol,*
1051   *29*(5), 375-381. https://doi.org/10.1097/cco.0000000000000390
1052
1053 Yang, P. M., Hong, Y. H., Hsu, K. C., & Liu, T. P. (2019). p38α/S1P/SREBP2 activation by the SAM-
1054   competitive EZH2 inhibitor GSK343 limits its anticancer activity but creates a druggable
1055   vulnerability in hepatocellular carcinoma. *Am J Cancer Res, 9*(10), 2120-2139.
1056
1057 Yu, W., Zhang, F., Wang, S., Fu, Y., Chen, J., Liang, X., Le, H., Pu, W. T., & Zhang, B. (2017, Apr
1058   13). Depletion of polycomb repressive complex 2 core component EED impairs fetal
1059   hematopoiesis. *Cell Death Dis, 8*(4), e2744. https://doi.org/10.1038/cddis.2017.163
1060
1061 Zhang, M. J., Chen, D. S., Li, H., Liu, W. W., Han, G. Y., & Han, Y. F. (2019). Clinical significance of
1062   USP7 and EZH2 in predicting prognosis of laryngeal squamous cell carcinoma and their
1063   possible functional mechanism. *Int J Clin Exp Pathol, 12*(6), 2184-2194.
1064
1065 Zhang, Q., Han, Q., Zi, J., Ma, J., Song, H., Tian, Y., McGrath, M., Song, C., & Ge, Z. (2019, Sep).
1066   Mutations in EZH2 are associated with poor prognosis for patients with myeloid
1067   neoplasms. *Genes Dis, 6*(3), 276-281. https://doi.org/10.1016/j.gendis.2019.05.001
1068
1069