Genome-wide CRISPRi/a screens in human neurons link lysosomal failure to ferroptosis

Ruilin Tian[1,2,*], Anthony Abarientos[1], Jason Hong[1], Sayed Hadi Hashemi[3], Rui Yan[4], Mike A. Nalls[5,6], Andrew B. Singleton[5], Ke Xu[4], Faraz Faghri[3,5,6], Martin Kampmann[1,7,8,*]

[1] Institute for Neurodegenerative Diseases, Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA, USA

[2] Biophysics Graduate Program, University of California, San Francisco, San Francisco, CA, USA

[3] Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana, IL, USA

[4] Department of Chemistry, University of California, Berkeley, Berkeley, CA, USA

[5] Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA

[6] Data Tecnica International, LLC, Glen Echo, MD, USA

[7] Chan Zuckerberg Biohub, San Francisco, CA, USA

[8] Lead contact

* Correspondence: ruilin.tian@ucsf.edu (R.T.), martin.kampmann@ucsf.edu (M.K.)

SUMMARY
Single-cell transcriptomics provide a systematic map of gene expression in different human cell types. The next challenge is to systematically understand cell-type specific gene function. The integration of CRISPR-based functional genomics and stem cell technology enables the scalable interrogation of gene function in differentiated human cells. Here, we present the first genome-wide CRISPR interference and CRISPR activation screens in human neurons.
We uncover pathways controlling neuronal response to chronic oxidative stress, which is implicated in neurodegenerative diseases. Unexpectedly, knockdown of the lysosomal protein prosaposin strongly sensitizes neurons, but not other cell types, to oxidative stress by triggering the formation of lipofuscin, a hallmark of aging, which traps iron, generating reactive oxygen species and triggering ferroptosis. We also determine transcriptomic changes in neurons following perturbation of genes linked to neurodegenerative diseases. To enable the systematic comparison of gene function across different human cell types, we establish a data commons named CRISPRbrain.

INTRODUCTION

The human body comprises hundreds of different cell types. Even though their genomes are nearly identical, cell types are characterized by vastly different cell biologies, enabling them to fulfill diverse physiological functions. Transcriptomic profiling, fueled by recent advances in single-cell- and single-nucleus-RNA sequencing technologies, has enabled the establishment of a Human Cell Atlas of cell-type specific gene expression signatures ((Regev et al., 2017). In addition to gene expression, gene function can also be cell type-specific, as evidenced by the fact that mutations in broadly expressed or housekeeping genes can lead to cell-type specific defects and disease states. Striking examples are familial mutations causing neurodegenerative diseases, which are often characterized by the selective vulnerability of specific neuronal subtypes, even if the mutated gene is expressed throughout the brain or even throughout the whole body. Cell-type specific gene function is also supported by our recent finding that knockdown of certain genes can have remarkably different impacts on cell survival and gene expression in different isogenic human cell types, including stem cells and neurons (Tian et al., 2019).

Therefore, understanding the function of human genes in different cell types is the next step toward elucidating tissue-specific cell biology and uncovering disease mechanisms. To this end, we recently developed a functional genomics platform, leveraging the strengths of CRISPR interference (CRISPRi) and induced pluripotent stem cell (iPSC) technology, that enables large-scale, multimodal loss-of-function genetic screens in differentiated human cell types, as demonstrated in neurons (Tian et al., 2019).

Here, we present a gain-of-function screening platform in human iPSC-derived neurons based on CRISPR-activation (CRISPRa), which can yield complementary biological insights to CRISPRi screens (Gilbert et al., 2014). We conduct the first genome-wide CRISPRi and CRISPRa screens in human neurons based on a panel of cellular phenotypes, and CROP-seq screens to uncover transcriptional fingerprints of genes associated with neurodegenerative diseases. We apply our functional genomics platforms to systematically identify genetic modifiers of levels of reactive oxygen species and peroxidized lipids, and neuronal survival under oxidative stress, one of the predominant stresses in aging (Finkel and Holbrook, 2000) and neurodegenerative diseases (Barnham et al., 2004). These screens uncovered an unexpected role for the lysosomal protein prosaposin (*PSAP*), knockdown of which caused the formation of lipofuscin, a hallmark of aging, which traps iron, generating reactive oxygen species and triggering ferroptosis. Intriguingly, *PSAP* deficiency caused these dramatic phenotypes only in neurons, but not in iPSCs or HEK293 cells.

These results demonstrate the power of our approach in uncovering novel, cell-type specific human cell biology. We establish a Data Commons, named CRISPRbrain, for systematic

exploration, interactive visualization, and comparison of functional genomics screening results in differentiated human cell types.

RESULTS

Genome-wide CRISPRi and CRISPRa screens reveal genes controlling survival of human neurons

We previously established a CRISPRi platform that enables robust knockdown of endogenous genes and large-scale loss-of-function genetic screens in human iPSC-derived neurons (Tian et al., 2019). Here, we expand our toolbox by establishing a CRISPRa platform for overexpression of endogenous genes and genome-wide gain-of-function screens in human iPSC-derived neurons. We adapted a previously published inducible CRISPRa system, DHFR-dCas9-VPH, whose function had been validated in human iPSCs (Weltner et al., 2018). In this system, the CRISPRa machinery is tagged by a DHFR degron leading to proteasomal degradation of the entire fusion protein. Trimethoprim (TMP) stabilizes the DHFR degron and prevents turnover, thereby inducing CRISPRa activity. As for our CRISPRi platform, an expression cassette for the CRISPRa machinery, CAG promoter-driven DHFR-dCas9-VPH, was stably integrated into the CLYBL safe-harbor locus of an iPSC line with an inducible Neurogenin 2 (*Ngn2*) expression cassette in the AAVS1 safe-harbor locus, which can be efficiently differentiated into homogenous glutamatergic neurons in a highly scalable manner (i^{3}N-iPSC, (Wang et al., 2017) (Figure 1A). A monoclonal line of these CRISPRa-iPSCs was generated and a normal karyotype was confirmed (Figure S1A). We validated the functionality of CRISPRa-iPSCs by confirming robust induction of an endogenous gene, *CXCR4*, in iPSC-derived neurons in a tightly inducible manner (Figure 1B).

We previously conducted a sub-genome scale CRISPRi screen to reveal genes controlling neuronal survival using an sgRNA library targeting 2,325 genes representing the "druggable genome" (Tian et al., 2019). Here, we greatly expanded the screen to target all protein-coding human genes in both loss-of- and gain-of-function screens by CRISPRi and CRISPRa, respectively. To our knowledge, these are the first genome-wide CRISPR screens in human neurons. We lentivirally transduced our CRISPRi and CRISPRa iPSCs with the genome-wide hCRISPRi/a-v2 sgRNA libraries (Horlbeck et al., 2016) and differentiated them into neurons (Figure 1C). For the CRISPRa screen, TMP was added to Day 0 neurons to induce CRISPRa activity. Based on the depletion or enrichment of sgRNAs targeting specific genes at Day 10 compared to Day –3, we identified hit genes for which knockdown or overexpression was toxic or protective for neuronal survival, using our previously published pipeline, MAGeCK-iNC (Tian et al., 2019) (Figure 1D, see Methods).

For CRISPRi, among the top 10 hits with toxic knockdown phenotypes were genes encoding superoxide dismutases *SOD1* and *SOD2*, which protect cells from oxidative stress (Figure 1E). Intriguingly, *SOD1* and *SOD2* are not broadly essential in other human cell types (Figure S1B), including pluripotent stem cells (Ihry et al., 2019; Mair et al., 2019; Yilmaz et al., 2018)) and cancer cells (Hart et al., 2017), suggesting a selective vulnerability to oxidative stress of neurons. Genes encoding subunits of the vacuolar ATPase(V-ATPase) complex, including *ATP6V1H*, *ATP6V1C1,* and *ATP6AP1*, were also among the top hits with toxic knockdown phenotypes. The V-ATPase complex mediates acidification of the endo-lysosomal compartment through ATP hydrolysis-coupled proton transport, and its dysfunction can cause neurodegenerative diseases (Colacurcio and Nixon, 2016). Gene Ontology (GO) analysis revealed additional pathways that were enriched in the top 100 hits with toxic or protective phenotypes in the CRISPRi screen (Figure 1F, left). For example, genes involved in cholesterol biosynthesis were strongly enriched in hits with toxic knockdown phenotypes, suggesting an important role of cholesterol in maintaining neuronal survival, consistent with our previous findings (Tian et al., 2019). Other homeostatic pathways, including iron homeostasis, protein folding, mRNA processing, and autophagy, were also essential for neuronal survival.

For CRISPRa, GO analysis revealed that overexpression of pro- or anti-apoptotic genes showed expected toxic or protective survival phenotypes, respectively, validating our approach (Figure 1F, right). Next, we compared hits from our CRISPRi and CRISPRa screens (Figure 1E). Overall, there was little overlap between CRISPRi and CRISPRa hits, consistent with a previous study comparing parallel CRISPRi and CRISPRa screens in K562 cells (Gilbert et al., 2014). The fact that CRISPRi and CRISPRa screens uncover distinct sets of hit genes can be explained by several factors. First, a gene that is not expressed in neurons will not have a CRISPRi knockdown phenotype, but may have a CRISPRa overexpression phenotype. Indeed, genes expressed at low or undetectable levels were strongly depleted from CRISPRi hits (Figure 1G, left), whereas CRISPRa hits were not restricted by endogenous expression levels (Figure 1G, right). Second, many genes encode proteins that form complexes (such as the V-ATPase complex), for which knockdown of a single component could abrogate the function of the entire complex and result in a phenotype, whereas overexpression of a single subunit by CRISPRa would generally be insufficient to induce an increased function of the complex. Last, knockdown of a single gene may not lead to a phenotype due to redundancy. For example, gene paralogs can compensate for each other's loss-of-function. Paralogs tend to have less deleterious loss-of-function effect in CRISPR-Cas9 screens in different human cell lines (Dandage and Landry, 2019). Consistent with this idea, the percentage of hit genes with paralogs was much lower in our CRISPRi screen compared to that in CRISPRa screen (17.5% (355 / 2032) vs. 37.2% (314 / 845), $P < 0.00001$, Figure 1H). Taken together, CRISPRi and CRISPRa screens can uncover complementary biological insights.

Nevertheless, we identified a number of overlapping hit genes in the two screens (Figure 1E). Many of these genes showed opposing phenotypes on neuronal survival upon CRISPRa induction and CRISPRi repression. Among these genes, we identified known regulators of apoptosis, such as *TP53*, *IKBIP,* and factors specifically of neuronal survival, such as *XIAP* (Kügler et al., 2000; Tsang et al., 2009; Unsain et al., 2013; Xu et al., 1999), *GDPGP1* (Schulz et al., 2020) and *MAP3K12* (Chen et al., 2008; Le Pichon et al., 2017; Tian et al., 2019; Welsbie et al., 2013), as well as genes not previously implicated in neuronal survival, such as *IGF2BP1* and *IGF2BP3*.

For some genes, including several involved in protein homeostasis (e.g. *CUL3*, *FBXO2*, *COMMD1*, and *HSPD1*), perturbations in both directions were detrimental to neuronal survival (i.e. genes in the left lower quadrant of Figure 1E), suggesting their endogenous expression levels are narrowly balanced for optimal survival.

While some CRISPRi and CRISPRa hits were also survival-relevant in other cell types based on previously reported screens, a substantial fraction of hits was neuron-specific (Fig. S1B).

Genome-wide CRISPRi/a screens elucidate pathways controlling neuronal response to oxidative stress

Given the unique vulnerability of neurons to redox imbalance-induced oxidative stress, which is often found in the brain of patients with neurodegenerative diseases (Dalfó et al., 2005; Hensley et al., 1995; Smith et al., 1991; Subbarao et al., 1990; Yao et al., 2003), we sought to apply our functional genomics toolkit to systematically identify factors that are important for redox homeostasis and oxidative stress response in human neurons. We performed screens based on two strategies.

First, we conducted genome-wide CRISPRi and CRISPRa screens to identify modifiers of neuronal survival under oxidative stress conditions (Figure 2A). Standard neuronal culture medium contains a combination of antioxidants, including vitamin E, vitamin E acetate, superoxide dismutase, catalase, and glutathione. To create an environment of chronic low-level oxidative stress, we cultured neurons in medium lacking the above antioxidants (–AO medium). We reasoned that compared to acute harsh treatments to induce ROS, such as adding H_2O_2 or rotenone, –AO medium provided a more physiologically relevant approximation of chronic oxidative stress.

Next, we compared modifiers of neuronal survival in this oxidative stress condition to the modifiers of survival in the standard, unstressed condition (Figure 2B). Interestingly, in the comparison of CRISPRi hits, we identified that *GPX4* (encoding the selenoprotein Glutathione Peroxidase 4) and genes responsible for selenocysteine incorporation into proteins (including

PSTK, *SEPHS2*, and *SEPSECS*) were particularly essential for neurons to survive under oxidative stress (Figure 2B,C). GPX4 utilizes glutathione to reduce peroxidized lipids and thus prevents ferroptosis, which is a non-apoptotic type of cell death caused by iron-dependent lipid peroxidation (Yang et al., 2014). This result suggested that neurons could be susceptible to ferroptosis under oxidative stress conditions. Hits from CRISPRa screens showed high correlation (R = 0.82) between oxidative stress and unstressed conditions, suggesting no strong stress-specific phenotypes for overexpressed genes (Figure S2).

Second, we conducted genome-wide CRISPRi screens for modifiers of levels of ROS and peroxidized lipids in neurons. Specifically, we stained CRISPRi neurons transduced with genome-wide sgRNA libraries with fluorescent indicators of ROS and lipid peroxidation (CellRox and Liperfluo, respectively) and sorted them into high and low fluorescence populations by FACS (Figure 2A). The MAGeCK-iNC pipeline was used to identify hit genes, knockdown of which led to an increase ('high signal'') or decrease ('low signal') in ROS or peroxidized lipids. From these screens, we identified both known and unexpected genetic modifiers of ROS and peroxidized lipid levels. We found that knockdown of components of the electron transport chain increased both ROS and lipid peroxidation levels (Figure 2D). This was expected because ROS are mainly generated by proton leak from the electron transport chain, and knockdown of electron transport chain components could increase proton leakage. Many autophagy-related genes were also common hits in the two screens (Figure 2 D,E), such as *WIPI2*, *ATG9A, ATG13, ATG14*, and *BECN1*, suggesting an important role of autophagy in maintaining redox homeostasis in cells, as reported in previous studies (Bensaad et al., 2009; Taguchi et al., 2012; Wang et al., 2016).

Disruption of genes involved in the mTORC1 pathway, including components of the mTORC1 complex (*MTOR*, *RPTOR*, and *MLST8)* or its activator *RHEB* reduced ROS and/or lipid peroxidation levels, whereas knockdown of *NPRL3*, a mTORC1 inhibitor, induced ROS and lipid peroxidation levels in neurons (Figure 2D). This is consistent with previous observations that increasing mTORC1 signalling induced ROS production (Reho et al., 2019), whereas inhibiting mTORC1 reduced ROS (Nacarelli et al., 2014; Reho et al., 2019; Shin et al., 2011). Moreover, GO term enrichment analysis of lipid peroxidation hits revealed a strong enrichment of peroxisomal genes, knockdown of which increased lipid peroxidation, consistent with the important roles of peroxisomes in redox regulation and degradation of (poly-)unsaturated fatty acids.

FBXO7, a gene associated with Parkinson's disease, whose deficiency was found to cause complex I respiratory impairment and ROS production (Delgado-Camprubi et al., 2017), also increased ROS levels when knocked down in our screen (Figure 2D). Knockdown of other previously characterized ROS regulators, including positive regulators such as *PARP1* (Hocsak et al., 2017), *SAT1* (Ou et al., 2016) and *NOX5* (Bánfi et al., 2001) and negative regulators such

as *PTEN* (Bankoglu et al., 2016) and *FH* (Bardella et al., 2012; Sudarshan et al., 2009) showed the expected effects on ROS and/or peroxidized lipid levels in our screens (Figure 2D).

Interestingly, key regulators of ferroptosis were also hits in the lipid peroxidation screen. *ACSL4,* encoding Acyl-CoA Synthetase Long Chain Family Member 4, which enriches cellular membranes with long PUFAs is required for ferroptosis (Doll et al., 2017; Yuan et al., 2016). ACLS4 inhibition has been shown to prevent ferroptosis (Doll et al., 2017), consistent with a reduction of peroxidized lipids upon knockdown in our screen (Figure 2D). By contrast, knockdown of *CD44*, whose splicing variant CD44v stabilizes the cystine/glutamate antiporter xCT at the plasma membrane and increases cysteine uptake for GSH synthesis, thereby inhibiting ferroptosis (Hasegawa et al., 2016; Ishimoto et al., 2011), increased lipid peroxidation in our screen as expected (Figure 2D).

Surprisingly, we found several strong hit genes with lysosomal functions but not associated with autophagy (Figure 1D). In particular, *PSAP* (encoding the lysosomal protein prosaposin) was one of the strongest hits in both ROS and lipid peroxidation screens (Figure 2D).

To further investigate hit genes from the genome-wide ROS and lipid peroxidation screens in high-throughput, we conducted focused secondary screens with an sgRNA library targeting 730 hit genes (Figure 2A). One secondary screen used lysotracker stain as a readout to uncover redox hits that may also alter lysosomal status. Lysotracker is a fluorescent acidotropic probe for labeling and tracking acidic organelles in live cells, which has been used to detect lysosome localization, quantify the number and sizes of lysosomes (Xu et al., 2014; Zhitomirsky and Assaraf, 2015) and to monitor autophagy (Chikte et al., 2014; DeVorkin and Gorski, 2014). The other secondary screen used the labile Fe^{2+}-detecting probe FeRhoNox-1. Since intracellular labile ferrous iron (Fe^{2+}) can contribute to ROS generation and lipid peroxidation via Fenton reactions (Dixon and Stockwell, 2014), we asked whether some of the redox hits act through changing iron homeostasis.

Our secondary screens uncovered that several of our original ROS/lipid peroxidation hit genes also strongly affected iron and/or lysosome status (Figure 2F). We found that knockdown of many lysosome/autophagy-related genes affected both lysosomal status and iron levels, reflecting the key role of lysosomes in iron homeostasis (Bogdan et al., 2016; Weber et al., 2020). Among these genes, we found *WDR45*, which is involved in autophagosome formation and lysosomal degradation and is also associated with Neurodegeneration with Brain Iron Accumulation (NBIA) in line with previous studies (Haack et al., 2012; Seibler et al., 2018). We also identified other known iron regulators for which knockdown increased iron levels, including *FBXL5,* a negative regulator of iron levels (Muto et al., 2017), the Fe/S cluster biogenesis genes *NFU1* and *NUBPL*, and *MCOLN1*, which encodes an endolysosomal iron release channel (Dong et al., 2008) (Figure 2F). Interestingly, knockdown of genes involved in the electron transport

chain increased labile iron (Figure 2F). Mitochondria are a major site of cellular iron storage and utilization in processes such as heme synthesis and iron-sulfur cluster biogenesis (Ward and Cloonan, 2019). Disruption of the electron transport chain may lead to mitochondrial damage which can cause the subsequent release of labile iron and reduced iron utilization, thus increasing labile iron levels.

In summary, our screens for survival of oxidative stress and levels of ROS and peroxidized lipids uncovered many categories of known redox regulators, validating the sensitivity of our approach and supporting the notion that core mechanisms of redox regulation are conserved across different cell types. A substantial fraction of the hit genes were also modifiers of labile iron levels and/or lysosomal status (Figure 2G).

Loss of prosaposin increases ROS and lipid peroxidation levels in neurons and causes neuronal ferroptosis under oxidative stress

Surprisingly, we identified that knockdown of *PSAP* altered lysosomal status and strongly induced ROS, lipid peroxidation, and iron levels when depleted (Figure 2D,F,G). Given the unexpected link between the known functions of PSAP and its screen phenotypes, we decided to further investigate the underlying mechanisms of *PSAP* in neuronal redox regulation.

PSAP encodes prosaposin, which is a pro-protein that is proteolytically processed by cathepsin D (encoded by *CTSD*) in the lysosome to generate four cleavage products: saposins A, B, C, and D (Hiraiwa et al., 1997). These four saposins, along with the lysosomal protein GM2A (GM2 Ganglioside Activator), function as activators for glycosphingolipid degradation by lysosomal hydrolases (Sandhoff and Harzer, 2013) (Figure 3A). Intriguingly, both *CTSD* and *GM2A* were also hits in at least one of the redox screens, showing similar knockdown phenotypes as *PSAP* (Figure 3B). Together, these findings suggest an important and unexpected role for glycosphingolipid degradation in redox homeostasis.

To validate our screen results using an independent approach, we generated a clonal *PSAP* knockout (KO) iPSC line (Figure 3C, D). Next, we measured ROS and lipid peroxidation levels in WT and *PSAP* KO neurons. We included C11-BODIPY, another indicator for peroxidized lipid, as additional validation for lipid peroxidation levels. Indeed, we observed a substantial increase in ROS and peroxidized lipid levels in *PSAP* KO neurons, confirming our screen results (Figure 3E). Moreover, ROS induction in PSAP KO neurons can be rescued by the overexpression of *PSAP* cDNA, confirming that phenotypes were not driven by off-target genome editing. Interestingly, *PSAP* KO or knockdown in other cell types (iPSCs and HEK293s) did not increase ROS levels (Figure 3F), suggesting a neuron-specific role for *PSAP* in redox regulation.

Next, we asked if increased ROS in *PSAP* KO neurons affects survival. Interestingly, we did not observe a survival defect of *PSAP* KO neurons over two weeks of culture in standard neuronal medium (+AO). Strikingly, however, when we cultured these neurons in the medium lacking antioxidants (–AO), *PSAP* KO caused a dramatic decrease in survival around Day 11, resulting in complete death of all neurons by Day 14 (Figure 3G).

To further investigate the underlying mechanism of cell death, we treated WT and *PSAP* KO neurons with compounds that inhibit different cell death pathways. Intriguingly, the viability of *PSAP* KO neurons under the -AO condition was not rescued by Z-VAD-FMK, a pan-caspase inhibitor that blocks apoptosis, but was fully rescued by ferroptosis inhibitors, including the iron chelator deferoxamine (DFO) and the lipid peroxidation inhibitor ferrostatin-1 (Figure 3H, I), suggesting that mild oxidative stress triggers ferroptosis in *PSAP* KO neurons.

In summary, we validated the unexpected strong hit gene from our unbiased screens, *PSAP,* as a strong redox modifier in human neurons. Loss of *PSAP* increases ROS levels and leads to increased lipid peroxidation, resulting in neuronal ferroptosis in response to mild oxidative stress.

Loss of prosaposin leads to lipofuscin formation and iron accumulation

Given the surprising connection between *PSAP* and ferroptosis, we further investigated the underlying mechanism. Neurons in the following experiments were cultured in standard neuronal medium (+AO) unless otherwise specified. We first asked if the loss of *PSAP* blocked glycosphingolipid degradation, given the canonical function of saposins. Indeed, untargeted lipidomics revealed that almost all glycosphingolipid species accumulated significantly in *PSAP* KO neurons compared to WT neurons (FDR < 0.01, Figure 4A, B, Supplemental Table 1). Ether lipids, which are peroxisome-derived glycerophospholipids, were also enriched in *PSAP* KO neurons. The mechanisms driving ether lipid accumulation and whether this accumulation is protective or maladaptive remain to be investigated. Interestingly, the accumulation of ether lipids was also recently characterized as a feature of hypoxia (Jain et al., 2020).

We also confirmed the accumulation of a specific glycosphingolipid species, GM1 ganglioside, in *PSAP* KO neurons by immunostaining (Figure 4C). Interestingly, we did not observe GM1 accumulation in *PSAP* KO iPSCs (Figure 4D), suggesting a cell-type specific role of *PSAP*. Strikingly, we observed dramatically enlarged lysosomes in *PSAP* KO neurons (Figure 4C), which were also reflected in an increased lysotracker signal by flow cytometry (Figure 4E), as expected based on our screen results (Figure 2F). Again, this phenotype was neuron-specific, as *PSAP* KO did not cause enlarged lysosomes in other cell types, iPSCs or HEK293 cells (Figure 4F).

We observed colocalization of accumulated GM1 and lysosomes by conventional confocal microscopy and by STORM super-resolution microscopy (Rust et al., 2006) (Figure 4C,G), consistent with the notion that the lysosome is the main site of glycosphingolipid degradation.

To further characterize the enlarged lysosome-like structures accumulating in *PSAP* KO neurons, we performed electron microscopy (EM). Remarkably, we observed a large number of electron-dense granules that resembled the structure of lipofuscin (also known as age pigment). Lipofuscin is an insoluble aggregate of peroxidized lipids, proteins and metals in lysosomes of postmitotic cells, such as neurons, which is a hallmark of aging and also associated with several neurodegenerative diseases (Moreno-García et al., 2018). There is currently no consensus on the molecular mechanisms driving lipofuscin formation (Moreno-García et al., 2018). Further supporting the formation of lipofuscin in PSAP KO neurons was the detection of strong autofluorescence (Figure 4I), a hallmark of lipofuscin (Mochizuki et al., 1995). Lipofuscin is known to accumulate iron from lysosomal degradation of iron-rich proteins or organelles (e.g. mitochondria), and accumulated iron in lipofuscin has been proposed to generate ROS through the Fenton reaction (Brunk et al., 1992; Höhn et al., 2010). Indeed, we observed a substantial increase of iron levels in *PSAP* KO neurons using a panel of different iron indicators (Figure 4J, K, S3A, B), This result was consistent with our finding of PSAP as a top hit in our iron level screen (Figure 2F). Iron accumulation occurred in lysosomes and was partially reversed by the iron chelator DFO (Figure 4K, S3A, B). Meanwhile, dramatically increased lipid peroxidation was detected in lysosomes of *PSAP* KO neurons under the -AO condition (Figure 4L), likely due to the accumulated iron, consistent with the strong lipid peroxidation-inducing phenotype of *PSAP* knockdown in our screen (Figure 2D).

Other lysosomal functions were also disrupted in *PSAP* KO neurons, likely due to the formation of lipofuscin. We observed a massive accumulation of autophagosomes in *PSAP* KO neurons as indicated by the increased ratio of PE-conjugated form of LC3B to its unconjugated form (LC3B-II / LC3B-I) by western blot (Figure 4M, N), and increased LC3B puncta by immunostaining (Figure 4O). Treatment with Bafilomycin A1 (BafA1), an inhibitor for degradation of autophagosomes by lysosomes, increased the LC3B-II / LC3B-I ratio in WT, but not *PSAP* KO neurons (Figure 4M), suggesting a blockade of autophagic flux in *PSAP* KO neurons.

Moreover, we performed RNA-seq on WT and *PSAP* KO neurons. Interestingly, a number of genes involved in cholesterol biosynthesis were upregulated in *PSAP* KO neurons (Figure S3C, D). Consistent with the induction of this pathway, we found increased cholesterol levels in lysosomes of *PSAP* KO neurons, as measured by Filipin staining (Figure S3E, F). This finding mirrored the results of previous studies that *PSAP* is a strong genetic modifier of cholesterol

levels (Bartz et al., 2009) and that accumulated glycosphingolipids can lead to the accumulation of cholesterol in lysosomes by inhibiting cholesterol efflux (Glaros et al., 2005; Puri et al., 2003).

In summary, we elucidated that loss of saposins in *PSAP* KO neurons blocks glycosphingolipid degradation in lysosomes, leading to the formation of lipofuscin, which in turn accumulates iron and generates ROS that peroxidized lipids. The accumulation of peroxidized lipids leads to neuronal ferroptosis in the absence of antioxidants (Figure 4P).

Transcriptomic signatures of perturbations of disease-associated genes in human neurons

Over the past decade, genome-wide association studies (GWASs) have uncovered hundreds of genes that are associated with human neurodegenerative diseases (Chang et al., 2017; Kunkle et al., 2019). However, functional characterizations of these risk genes are largely lacking (Gallagher and Chen-Plotkin, 2018). Our CRISPRi and CRISPRa platforms provide a high-throughput approach to systematically interrogate gene function in human neurons. Beyond one-dimensional phenotypes such as survival or fluorescent reporter levels, CRISPR perturbation can be coupled to single-cell RNA sequencing, using CROP-seq or Perturb-Seq strategies (Adamson et al., 2016; Datlinger et al., 2017; Dixit et al., 2016), to provide rich transcriptomic phenotypes.

The hit genes from our unbiased genome-wide CRISPRi and CRISPRa screens included hundreds of genes associated with neurodegenerative diseases (based on DisGeNet annotation (Piñero et al., 2015) and literature research). To better characterize these neurodegenerative disease risk genes, we performed CROP-seq experiments, targeting 184 genes for CRISPRi and 100 genes for CRISPRa with 2 sgRNAs per gene (Figure 5A, see Method for details). Significant on-target gene knockdown or overexpression were detected for most of the targets in the CRISPRi or CRISPRa library, respectively (> 70% at P < 0.05, Figure 5B). Within the population of cells expressing sgRNAs for a specific target gene, the levels of target knockdown or overexpression were heterogeneous in some cases (Figure 5C,D), and we established a machine-learning based strategy to identify cells in which the targeted gene was effectively knocked down or overexpressed (Figure 5C,D, S4A,B, see Methods for details).

We identified differentially expressed genes (DEGs) caused by each perturbation, and determined the similarity of DEGs among all perturbations (see Method for details). This analysis revealed clusters of genes that shared common DEG signatures (Figure 5E). As expected, knockdown of functionally related genes had similar transcriptomic consequences. For example, knockdown several mitochondria-related genes, namely *COX10, NDUFS8, NDUFV1, MRPL10*, and *SOD2,* resulted in similar DEGs, as did knockdown of the anti-apoptotic genes *BNIP1* and *XIAP*. However, we also identified unexpected gene clusters. For example, knockdown of *VPS54*, *PAXIP1*, and *PON2* caused highly correlated transcriptomic changes (Figure 5E, S4C). *VPS54* and *PON2* have been implicated in Amyotrophic Lateral Sclerosis

(ALS) (Schmitt-John et al., 2005; Slowik et al., 2006) whereas *PAXIP1* is associated with Alzheimer's disease (AD) (Chen et al., 2015). Shared transcriptomic changes included: (i) increased expression of *EBF3*, an apoptosis inducer (Zhao et al., 2006) that is also upregulated in the hippocampus of AD model mice (Gu et al., 2018), (ii) decreased expression of components of neurofilaments, namely *NEFL*, *NEFM*, and *NEFH*. Importantly, significantly decreased expression of *NEFL* and *NEFM* was also found in neurons of early-stage AD patients in a recent single-cell transcriptomic study (Mathys et al., 2019) and (iii) decreased expression of other genes important for neuronal function, including *PCDH11X* and *PCDH11Y,* encoding protocadherin proteins, and *SYT2,* encoding synaptotagmin 2.

To identify genes that were co-regulated under different genetic perturbations, we performed weighted gene co-expression network analysis (WGCNA, (Langfelder and Horvath, 2008)). This analysis identified 10 modules for CRISPRi and 8 modules for CRISPRa that were co-regulated across different perturbations (Figure 5F,G). These modules contained genes enriched in various pathways (Figure 5F). Interestingly, we found a cluster of genes including *INSR*, *ATP5F1C*, *SOX5*, *GSX2* and *GBX2* (Figure 5E, CRISPRa) overexpression of which downregulated a gene module related to neurogenesis (M1) and upregulated a gene module related to the cell cycle (M5) (Figure 5G, bottom), suggesting that overexpression of these genes interfered with neuronal differentiation thus kept cells in a proliferating state. This result was expected for *SOX5*, *GSX2* and *GBX2*, which are transcription factors maintaining neural progenitor cell fate and/or their self-renewal (Luu et al., 2011; Martinez-Morales et al., 2010; Méndez-Gómez and Vicario-Abejón, 2012), but unexpected for *INSR* (encoding the insulin receptor) and *ATP5F1C* (encoding a subunit of mitochondrial ATPsynthase), suggesting these genes might have uncharacterized functions in neuronal fate regulation, which require further investigation.

The CROP-seq data also generated hypotheses for disease mechanisms. For example, *NQO1*, encoding the NAD(P)H:Quinone Oxidoreductase 1, is thought to be a cytoprotective factor through its antioxidant functions (Ross and Siegel, 2017). Elevated levels and activity of NQO1 has been found in the brains of patients with different neurodegenerative diseases (Bian et al., 2008; Imaizumi et al., 2012; van Muiswinkel et al., 2004; Raina et al., 1999; SantaCruz et al., 2004; Wang et al., 2000), as well as in patient iPSC-derived neurons (Imaizumi et al., 2012). The upregulation of NQO1 was proposed to be a neuroprotective mechanism against oxidative stress in neurodegenerative diseases. Paradoxically, however, *NQO1* overexpression showed a strong negative impact on neuronal survival in our pooled CRISPRa screen (Figure 1E). This finding suggests the intriguing hypothesis that elevated NQO1 in the context of neurodegenerative diseases is neurotoxic, rather than neuroprotective.

Our CROP-Seq data reveals numerous transcriptomic changes induced by NQO1 overexpression (Figure 5C, Figure 6A-C), providing several potential hypotheses for the mechanisms underlying NQO1 toxicity (Figure 6D).

First, we observed increased expression of many p53 target genes (including p53 itself). A previous study reported that overexpression of *NQO1* stabilizes p53 (Asher et al., 2002). The induced p53 activity may contribute to the strong toxic survival phenotype of NQO1 overexpression in neurons.

Second, *NQO1* overexpression strongly induced the Nrf2 pathway (Figure 6B, C). This finding was surprising because the transcription factor Nrf2 canonically functions upstream of NQO1 by inducing the expression of many antioxidant genes, including *NQO1*, in response to oxidative stress (Ma, 2013). Intriguingly, the Nrf2 pathway is also induced in some neurodegenerative diseases, both in patient brains (Ramsey et al., 2007) and in patient iPSC-derived neurons (Imaizumi et al., 2012). Upregulation of *NQO1* could induce or mimic an oxidative stress condition through unknown mechanisms, in turn activating the Nrf2 pathway, which however is insufficient to cope with the stress. Based on this hypothesis, the elevation of NQO1 levels in patients would actually contribute to disease, rather than being an effective defense mechanism.

Third, the toxicity of *NQO1* overexpression may be caused by other mechanisms, such as downregulation of genes involved in axon development or cholesterol metabolism, which we observed in the CROP-seq data (Figure 6B-D).

Taken together, our CROP-seq results provide a rich resource for investigating the consequences of perturbations of neurodegenerative disease-associated genes in human neurons and for generating testable hypotheses for the neurodegenerative disease mechanisms and potential therapeutic strategies.

CRISPRbrain, an open-access data commons for functional genomics screens in differentiated human cell types

To make the large dataset generated in this study publicly accessible, and to facilitate visualization, interactive exploration, and direct comparison with functional genomics screening data in a broad range of differentiated human cell types generated by many research groups, we developed a data commons, named CRISPRbrain (https://crisprbrain.org) (Figure 7A).

CRISPRbrain currently features screens with "simple" readouts (such as survival and fluorescent reporter levels), as well as screens with complex readouts (such as transcriptomes) Screens can be browsed and searched based on parameters of interest, such as cell type, genotype, mode of CRISPR perturbation, screen method and phenotype, as well as full text search (Figure 7B).

For simple screens, the phenotypes for the perturbed genes can be displayed as volcano plots and rank plots (Figure 7C). Data points can be selected individually or in groups to display

information about the perturbed gene(s). Gene names can be entered to label them in the graph. Data for the entire screen or for selected genes can be exported for offline analysis. Two screens of interest can be compared in a scatter plot (Figure 7D).

For RNA-Seq-based screens, genes of interest can be selected to display the transcriptomic phenotype resulting from perturbation of the gene (Figure 7E). Individual data points can be selected to display information about the transcript, and underlying data can be exported for offline analysis. Alternatively, the results from the entire screen can be explored in a hierarchically clustered heatmap of RNA-Seq phenotypes (Figure 7F).

We invite all research groups to contribute their functional genomics datasets to CRISPRbrain, with the ultimate goal to build a comprehensive atlas of gene function in all human cell types.

DISCUSSION

In this study, we developed a CRISPRa platform in human iPSC-derived neurons to enable robust gene overexpression screens, complementing our previously established CRISPRi platform (Tian et al., 2019). We demonstrated the power of these complementary screening approaches in multiple large-scale screens in human iPSC-derived neurons.

We determine the first comprehensive inventories of genes that, when depleted or activated, modulate the survival of human neurons. Intriguingly, many of these genes only affect survival in neurons, but not stem cells or cancer cells, supporting the notion that neurons have cell-type specific vulnerabilities, which may explain why defects in some generally expressed genes specifically cause neurological diseases.

Neurons, as one of the longest-living cell types in the human body, are challenged by various stresses in aging and disease. Due to their post-mitotic nature, neurons do not have the ability to 'self-renew' by cell division. Therefore, robust stress response mechanisms are required for neurons to maintain long-term health. One of the predominant types of stress in aging and neurodegenerative diseases is oxidative stress (Barnham et al., 2004; Finkel and Holbrook, 2000), which is induced by excessive accumulation of ROS in the cell. The brain is highly susceptible to ROS and ferroptosis, due to its high levels of oxygen consumption, abundant redox-active metals such as iron and copper, limited antioxidants and high levels of PUFAs (Patel, 2016). A large body of evidence has implicated oxidative stress, iron accumulation and ferroptosis in neurodegenerative diseases (Lin and Beal, 2006; Rouault, 2013; Stockwell et al., 2017), yet a comprehensive understanding of how neurons regulate redox homeostasis and maintain survival under oxidative stress is lacking. To this end, we decided to investigate

neuronal redox homeostasis and survival under oxidative stress conditions using our CRISPRi/a screening platform.

Compared to other recent genetic screens focusing on oxidative stress toxicity in human cells (Dubreuil et al., 2020; Reczek et al., 2017), our screens are unique in the following aspects. First, we used post-mitotic neurons instead of cancer cell lines. Second, we induced milder oxidative stress in cells by prolonged culture of cells in an antioxidant-free medium, compared to severe oxidative stress induced by paraquat or H_2O_2 in other studies. Third, we screened not only based on cell survival, but also on direct ROS and lipid peroxidation levels, while other studies focused on survival only. Given these major differences, it is not surprising that our screens identified many unique hits not identified in previous screens.

Numerous novel biological insights have emerged from our rich datasets. *GPX4* and genes related to *GPX4* synthesis are indispensable for neurons to survive oxidative stress. Given the major role of GPX4 in reducing lipid peroxidation and suppressing ferroptosis, this result suggests that lipid peroxidation-induced ferroptosis, rather than other forms of cell death, may be the main cause of neuronal loss under oxidative stress conditions that are commonly found in the brains of patients with neurodegenerative diseases. This is supported by numerous studies reporting high levels of iron and lipid peroxidation in neurodegenerative disease patient brains (Han et al., 2020; Ndayisaba et al., 2019), and by the finding that ferroptosis inhibitors such as iron chelators and the lipid-peroxidation inhibitor Ferrostatin-1 are neuroprotective in animal and cellular models of neurodegenerative diseases (Chen et al., 2013; Kaur et al., 2003; Kupershmidt et al., 2012; Skouta et al., 2014).

We also found that that disruption of lipid metabolism, in particular glycosphingolipid degradation in the lysosome, by loss of prosaposin (*PSAP*), drives the formation of lipofuscin in neurons, which leads to iron accumulation and strongly induces ROS production, oxidizing lipids and leading to neuronal ferroptosis under oxidative stress. While lipofuscin has traditionally been considered a byproduct of aging and as a consequence of defective cellular homeostasis, its physiological and pathological functions have not been well characterized. This is largely due to the lack of robust systems to model lipofuscin given the fact that lipofuscin is normally only formed in aged post-mitotic cells. Our *PSAP* KO neurons provide a reliable genetic system to model and study the biology of lipofuscin in young, live human neurons. Our result suggests a direct pathogenic role of lipofuscin in inducing neuronal ferroptosis. The accumulation of lipofuscin is a pathological hallmark of many degenerative diseases, such as neuronal ceroid lipofuscinosis (NCL) (Anderson et al., 2013) and inherited age-related macular degenerations (Delori et al., 1995; Dorey et al., 1989). It has also been characterized in Alzheimer's disease (Dowson et al., 1992, 1995), Parkinson's disease (Braak et al., 2001), Hungtinton's disease (Braak and Braak, 1992; Goebel et al., 1978) and *GRN*-associated frontotemporal dementia

(FTD) (Ward et al., 2017). Our findings suggest that inhibiting lipofuscin formation or subsequent ferroptosis may serve as new therapeutic strategies for these diseases.

Our results highlight the importance of lipid homeostasis, in particular balanced levels of glycosphingolipids, for neuronal health. Among all tissues in the human body, the brain is one of the richest in lipid content and lipid diversity. Lipids are not only an essential structural component of membranes, but also important signaling molecules in the brain (Bazan, 2005). Therefore, maintaining lipid homeostasis is of vital importance for brain cells, especially neurons with long neurites and dynamic synaptic vesicles release and recycling. Abnormal lipid metabolism has been observed in neurodegenerative diseases (Hallett et al., 2019; Isacson et al., 2019; Yadav and Tiwari, 2014). Accumulation of some glycosphingolipids, especially simple gangliosides, caused by inhibition of lysosome membrane recycling, contributes to neurodegeneration both in cultured neurons and in animal models (Boutry et al., 2018).

Many important questions around neuronal lipid and redox homeostasis and ferroptosis remain to be investigated. For example, since neurons have very long neurites, how do neurons sense and respond to a local lipid peroxidation event on the cell membrane? Are GPX4 or other enzymes recruited to sites of lipid peroxidation, or are peroxidized lipids internalized and delivered to GPX4 or other enzymes for detoxification? What are the sensors and mediators in these processes? How do peroxidized lipids cause death? Is it a passive physical process or a regulated biological program, and are there neuron-specific aspects of ferroptosis? If the latter is true, what are the players mediating cell death downstream of lipid peroxidation? Many of these questions can be readily investigated using our functional genomics platform.

There are several areas for future technology development. First, a robust inducible CRISPRi system that allows temporal control of gene knockdown in mature neurons will help avoiding false-positive screening phenotypes due to interference with the differentiation process. For example, in the CRISPRi survival screen, we unexpectedly identified genes in the N6-methyltransferase writer complex, including *ZC3H13*, *METTL3*, and *METTL4*, as strong protective hits when knocked down (Figure 1E). The N6-methyltransferase writer complex is responsible for m6A RNA modification, which regulates a variety of biological processes including neuronal differentiation (Edens et al., 2019; Li et al., 2018). The inducible CRISPRi system will help to determine whether the strong protective phenotypes of these genes are false positive due to defects in neuronal differentiation. Second, we have previously used our platform for arrayed high-content imaging screens for complex neuronal phenotypes, such as neurite outgrowth (Tian et al., 2019). Such imaging-based screens could be applied to study other neuronal phenotypes, such as electrophysiological signals (by voltage imaging or calcium imaging), synaptic vesicle dynamics and axonal transport, and cell non-autonomous phenotypes such as glia-neuron interaction. Imaging phenotyping can also be coupled with simultaneous transcriptomic profiling, thus allowing the association of gene expression with imaging

phenotypes for a given genetic perturbation. Lastly, by combining our platform with base editor or prime editing technology (Anzalone et al., 2019), we will be able to directly assess the effect of disease-associated mutations in human neurons on various cellular phenotypes in a scalable, massively parallel format. While base editor screens to assess human variants have been recently conducted in cancer cell lines (Hanna et al., 2020), it is of great interest to investigate how variants associated with different human diseases could affect various cell types differently given the fact that a lot of human diseases are tissue-specific.

In the past decade, tremendous efforts have advanced the molecular characterization of various human cell types. For example, through global collaboration in the Human Cell Atlas project, gene expression profiles for >30 human organs in > 4.5M cells have been obtained so far (https://data.humancellatlas.org/). Beyond gene expression, characterizing gene function in different cell types will be the next step towards building complete reference maps for the human body at the cellular level. We anticipate that our iPSC-based functional genomics platforms can be broadly applied to a variety of human differentiated cell types. Our CRISPRbrain Data Commons is complementary to the Human Cell Atlas and serves as an open-access platform for sharing and exploring functional genomics screening data in differentiated human cell types across labs. In contrast to the Cancer Dependency Map (https://depmap.org/), a widely-used database for survival-based functional genomics screens in cancer cells (Tsherniak et al., 2017), CRISPRbrain is the first database focused on screens in non-cancerous, differentiated human cell types for a broad range of phenotypes, including survival, reporter levels, and gene expression changes. Future versions of CRISPRbrain will support other high-dimensional phenotypes, such as imaging and electrophysiological data, and enable cloud computing and machine learning, thus providing researchers with a suite of tools to process, visualize, analyze and contribute functional genomics screening data in different human cell types. Parallel genetic screens across the full gamut of isogenic human cell types will uncover context-specific roles of human genes, leading to a deeper mechanistic understanding of how they control human biology and disease.

AUTHOR CONTRIBUTIONS
R.T. and M.K. conceived this study and wrote the manuscript with input from the other authors. R.T. designed and conducted experiments with help from A.A. and J.H. and guidance from M.K.. R.T. performed data analyses. R.Y. performed STORM imaging with guidance from K.X.. S.H.H., M.A.N. and F.F. developed the CRISPRbrain Data Commons with critical input from R.T. and M.K. and feedback from A.B.S.. All authors reviewed and approved the final manuscript.

ACKNOWLEDGEMENTS
We thank Li Gan, Bruce Conklin for support and advice; Paul Kennedy and Anna Nummy at Cayman Chemical for untargeted lipidomics; Eric Chow (UCSF), Derek Bogdanoff (UCSF), Angela Detweiler (CZI Biohub), Norma Neff (CZI Biohub) and Michelle Tan (CZI Biohub) for

next-generation sequencing; Avi Samelson and Xiaoyan Guo for comments on this manuscript; and Kun Leng, Emmy Li and James Olzmann for discussions. We thank the staff at the University of California Berkeley Electron Microscope Laboratory for advice and assistance in electron microscopy sample preparation and data collection. This research was supported by the Intramural Research Program of the NIH/NINDS, an NIH Director's New Innovator Award (NIH/ NIGMS DP2 GM119139 to M.K.), NIH/NIA grants (R01 AG062359 and R56 AG057528 to M.K.), the NINDS Tau Center Without Walls (NIH/NINDS U54 NS100717 to M.K.), an Allen Distinguished Investigator Award to M.K. (Paul G. Allen Family Foundation, to M.K.), a Chan-Zuckerberg Biohub Investigator Award (to M.K.), and a Tau Consortium Investigator Award (Rainwater Charitable Foundation, to M.K.). K.X. is a Chan Zuckerberg Biohub investigator and acknowledges support from the National Institute Of General Medical Sciences of the National Institutes of Health (DP2GM132681). A. B. S.'s participation was supported in part by the Intramural Research Program of the National Institute on Aging, National Institutes of Health, part of the Department of Health and Human Services (project number ZIA AG000957-16). CRISPRbrain development was supported in part by a collaboration between the Kampmann Lab, UCSF and Data Tecnica International, LLC.

DECLARATION OF INTERESTS

M.K. has filed a patent application related to CRISPRi and CRISPRa screening (PCT/US15/40449) and serves on the Scientific Advisory Board of Engine Biosciences and Casma Therapeutics.

DATA AND CODE AVAILABILITY

All screen datasets are publicly available on the CRISPRbrain website (http://crisprbrain.org/). The accession number for the RNA sequencing datasets reported in this paper is GEO: GSE152988 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE152988).

REFERENCES

Adamson, B., Norman, T.M., Jost, M., Cho, M.Y., Nuñez, J.K., Chen, Y., Villalta, J.E., Gilbert, L.A., Horlbeck, M.A., Hein, M.Y., et al. (2016). A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. Cell *167*, 1867-1882.e21.

Anderson, G.W., Goebel, H.H., and Simonati, A. (2013). Human pathology in NCL. Biochim. Biophys. Acta *1832*, 1807–1826.

Anzalone, A.V., Randolph, P.B., Davis, J.R., Sousa, A.A., Koblan, L.W., Levy, J.M., Chen, P.J., Wilson, C., Newby, G.A., Raguram, A., et al. (2019). Search-and-replace genome editing without double-strand breaks or donor DNA. Nature *576*, 149–157.

Asher, G., Lotem, J., Kama, R., Sachs, L., and Shaul, Y. (2002). NQO1 stabilizes p53 through a

distinct pathway. Proc Natl Acad Sci USA *99*, 3099–3104.

Bánfi, B., Molnár, G., Maturana, A., Steger, K., Hegedûs, B., Demaurex, N., and Krause, K.H. (2001). A Ca(2+)-activated NADPH oxidase in testis, spleen, and lymph nodes. J. Biol. Chem. *276*, 37594–37601.

Bankoglu, E.E., Tschopp, O., Schmitt, J., Burkard, P., Jahn, D., Geier, A., and Stopper, H. (2016). Role of PTEN in Oxidative Stress and DNA Damage in the Liver of Whole-Body Pten Haplodeficient Mice. PLoS ONE *11*, e0166956.

Bardella, C., Olivero, M., Lorenzato, A., Geuna, M., Adam, J., O'Flaherty, L., Rustin, P., Tomlinson, I., Pollard, P.J., and Di Renzo, M.F. (2012). Cells lacking the fumarase tumor suppressor are protected from apoptosis through a hypoxia-inducible factor-independent, AMPK-dependent mechanism. Mol. Cell. Biol. *32*, 3081–3094.

Barnham, K.J., Masters, C.L., and Bush, A.I. (2004). Neurodegenerative diseases and oxidative stress. Nat. Rev. Drug Discov. *3*, 205–214.

Bartz, F., Kern, L., Erz, D., Zhu, M., Gilbert, D., Meinhof, T., Wirkner, U., Erfle, H., Muckenthaler, M., Pepperkok, R., et al. (2009). Identification of cholesterol-regulating genes by targeted RNAi screening. Cell Metab. *10*, 63–75.

Bazan, N.G. (2005). Lipid signaling in neural plasticity, brain repair, and neuroprotection. Mol. Neurobiol. *32*, 89–103.

Bensaad, K., Cheung, E.C., and Vousden, K.H. (2009). Modulation of intracellular ROS levels by TIGAR controls autophagy. EMBO J. *28*, 3015–3026.

Bian, J.-T., Zhao, H.-L., Zhang, Z.-X., Bi, X.-H., and Zhang, J.-W. (2008). Association of NAD(P)H:quinone oxidoreductase 1 polymorphism and Alzheimer's disease in Chinese. J. Mol. Neurosci. *34*, 235–240.

Bogdan, A.R., Miyazawa, M., Hashimoto, K., and Tsuji, Y. (2016). Regulators of iron homeostasis: new players in metabolism, cell death, and disease. Trends Biochem. Sci. *41*, 274–286.

Boutry, M., Branchu, J., Lustremant, C., Pujol, C., Pernelle, J., Matusiak, R., Seyer, A., Poirel, M., Chu-Van, E., Pierga, A., et al. (2018). Inhibition of Lysosome Membrane Recycling Causes Accumulation of Gangliosides that Contribute to Neurodegeneration. Cell Rep. *23*, 3813–3826.

Braak, H., and Braak, E. (1992). Allocortical involvement in Huntington's disease. Neuropathol. Appl. Neurobiol. *18*, 539–547.

Braak, E., Sandmann-Keil, D., Rüb, U., Gai, W.P., de Vos, R.A., Steur, E.N., Arai, K., and Braak, H. (2001). alpha-synuclein immunopositive Parkinson's disease-related inclusion bodies in lower brain stem nuclei. Acta Neuropathol. *101*, 195–201.

Brunk, U.T., Jones, C.B., and Sohal, R.S. (1992). A novel hypothesis of lipofuscinogenesis and cellular aging based on interactions between oxidative stress and autophagocytosis. Mutat. Res.

275, 395–403.

Chang, D., Nalls, M.A., Hallgrímsdóttir, I.B., Hunkapiller, J., van der Brug, M., Cai, F., International Parkinson's Disease Genomics Consortium, 23andMe Research Team, Kerchner, G.A., Ayalon, G., et al. (2017). A meta-analysis of genome-wide association studies identifies 17 new Parkinson's disease risk loci. Nat. Genet. *49*, 1511–1516.

Chen, J.A., Wang, Q., Davis-Turak, J., Li, Y., Karydas, A.M., Hsu, S.C., Sears, R.L., Chatzopoulou, D., Huang, A.Y., Wojta, K.J., et al. (2015). A multiancestral genome-wide exome array study of Alzheimer disease, frontotemporal dementia, and progressive supranuclear palsy. JAMA Neurol. *72*, 414–422.

Chen, J., Marks, E., Lai, B., Zhang, Z., Duce, J.A., Lam, L.Q., Volitakis, I., Bush, A.I., Hersch, S., and Fox, J.H. (2013). Iron accumulates in Huntington's disease neurons: protection by deferoxamine. PLoS ONE *8*, e77023.

Chen, X., Rzhetskaya, M., Kareva, T., Bland, R., During, M.J., Tank, A.W., Kholodilov, N., and Burke, R.E. (2008). Antiapoptotic and trophic effects of dominant-negative forms of dual leucine zipper kinase in dopamine neurons of the substantia nigra in vivo. J. Neurosci. *28*, 672–680.

Chikte, S., Panchal, N., and Warnes, G. (2014). Use of LysoTracker dyes: a flow cytometric study of autophagy. Cytometry A *85*, 169–178.

Colacurcio, D.J., and Nixon, R.A. (2016). Disorders of lysosomal acidification-The emerging role of v-ATPase in aging and neurodegenerative disease. Ageing Res. Rev. *32*, 75–88.

Dalfó, E., Portero-Otín, M., Ayala, V., Martínez, A., Pamplona, R., and Ferrer, I. (2005). Evidence of oxidative stress in the neocortex in incidental Lewy body disease. J. Neuropathol. Exp. Neurol. *64*, 816–830.

Dandage, R., and Landry, C.R. (2019). Paralog dependency indirectly affects the robustness of human cells. Mol. Syst. Biol. *15*, e8871.

Datlinger, P., Rendeiro, A.F., Schmidl, C., Krausgruber, T., Traxler, P., Klughammer, J., Schuster, L.C., Kuchler, A., Alpar, D., and Bock, C. (2017). Pooled CRISPR screening with single-cell transcriptome readout. Nat. Methods *14*, 297–301.

Delgado-Camprubi, M., Esteras, N., Soutar, M.P., Plun-Favreau, H., and Abramov, A.Y. (2017). Deficiency of Parkinson's disease-related gene Fbxo7 is associated with impaired mitochondrial metabolism by PARP activation. Cell Death Differ. *24*, 120–131.

Delori, F.C., Staurenghi, G., Arend, O., Dorey, C.K., Goger, D.G., and Weiter, J.J. (1995). In vivo measurement of lipofuscin in Stargardt's disease--Fundus flavimaculatus. Invest. Ophthalmol. Vis. Sci. *36*, 2327–2331.

DeVorkin, L., and Gorski, S.M. (2014). LysoTracker staining to aid in monitoring autophagy in Drosophila. Cold Spring Harb. Protoc. *2014*, 951–958.

Dixit, A., Parnas, O., Li, B., Chen, J., Fulco, C.P., Jerby-Arnon, L., Marjanovic, N.D., Dionne,

D., Burks, T., Raychowdhury, R., et al. (2016). Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. Cell *167*, 1853-1866.e17.

Dixon, S.J., and Stockwell, B.R. (2014). The role of iron and reactive oxygen species in cell death. Nat. Chem. Biol. *10*, 9–17.

Doll, S., Proneth, B., Tyurina, Y.Y., Panzilius, E., Kobayashi, S., Ingold, I., Irmler, M., Beckers, J., Aichler, M., Walch, A., et al. (2017). ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid composition. Nat. Chem. Biol. *13*, 91–98.

Dong, X.-P., Cheng, X., Mills, E., Delling, M., Wang, F., Kurz, T., and Xu, H. (2008). The type IV mucolipidosis-associated protein TRPML1 is an endolysosomal iron release channel. Nature *455*, 992–996.

Dorey, C.K., Wu, G., Ebenstein, D., Garsd, A., and Weiter, J.J. (1989). Cell loss in the aging retina. Relationship to lipofuscin accumulation and macular degeneration. Invest. Ophthalmol. Vis. Sci. *30*, 1691–1699.

Dowson, J.H., Mountjoy, C.Q., Cairns, M.R., and Wilton-Cox, H. (1992). Changes in intraneuronal lipopigment in Alzheimer's disease. Neurobiol. Aging *13*, 493–500.

Dowson, J.H., Mountjoy, C.Q., Cairns, M.R., and Wilton-Cox, H. (1995). Alzheimer's disease: distribution of changes in intraneuronal lipopigment in the frontal cortex. Dementia *6*, 334–342.

Dubreuil, M.M., Morgens, D.W., Okumoto, K., Honsho, M., Contrepois, K., Lee-McMullen, B., Traber, G.M., Sood, R.S., Dixon, S.J., Snyder, M.P., et al. (2020). Systematic Identification of Regulators of Oxidative Stress Reveals Non-canonical Roles for Peroxisomal Import and the Pentose Phosphate Pathway. Cell Rep. *30*, 1417-1433.e7.

Edens, B.M., Vissers, C., Su, J., Arumugam, S., Xu, Z., Shi, H., Miller, N., Rojas Ringeling, F., Ming, G.-L., He, C., et al. (2019). FMRP Modulates Neural Differentiation through m6A-Dependent mRNA Nuclear Export. Cell Rep. *28*, 845-854.e5.

Finkel, T., and Holbrook, N.J. (2000). Oxidants, oxidative stress and the biology of ageing. Nature *408*, 239–247.

Gallagher, M.D., and Chen-Plotkin, A.S. (2018). The Post-GWAS Era: From Association to Function. Am. J. Hum. Genet. *102*, 717–730.

Gilbert, L.A., Horlbeck, M.A., Adamson, B., Villalta, J.E., Chen, Y., Whitehead, E.H., Guimaraes, C., Panning, B., Ploegh, H.L., Bassik, M.C., et al. (2014). Genome-scale CRISPR-mediated control of gene repression and activation. Cell *159*, 647–661.

Glaros, E.N., Kim, W.S., Quinn, C.M., Wong, J., Gelissen, I., Jessup, W., and Garner, B. (2005). Glycosphingolipid accumulation inhibits cholesterol efflux via the ABCA1/apolipoprotein A-I pathway: 1-phenyl-2-decanoylamino-3-morpholino-1-propanol is a novel cholesterol efflux accelerator. J. Biol. Chem. *280*, 24515–24523.

Goebel, H.H., Heipertz, R., Scholz, W., Iqbal, K., and Tellez-Nagel, I. (1978). Juvenile

Huntington chorea: clinical, ultrastructural, and biochemical studies. Neurology *28*, 23–31.

Gu, C., Chen, C., Wu, R., Dong, T., Hu, X., Yao, Y., and Zhang, Y. (2018). Long Noncoding RNA EBF3-AS Promotes Neuron Apoptosis in Alzheimer's Disease. DNA Cell Biol. *37*, 220–226.

Haack, T.B., Hogarth, P., Kruer, M.C., Gregory, A., Wieland, T., Schwarzmayr, T., Graf, E., Sanford, L., Meyer, E., Kara, E., et al. (2012). Exome sequencing reveals de novo WDR45 mutations causing a phenotypically distinct, X-linked dominant form of NBIA. Am. J. Hum. Genet. *91*, 1144–1149.

Hallett, P.J., Engelender, S., and Isacson, O. (2019). Lipid and immune abnormalities causing age-dependent neurodegeneration and Parkinson's disease. J. Neuroinflammation *16*, 153.

Hanna, R.E., Hegde, M., Fagre, C.R., DeWeirdt, P.C., Sangree, A.K., Szegletes, Z., Griffith, A., Feeley, M.N., Sanson, K.R., Baidi, Y., et al. (2020). Massively parallel assessment of human variants with base editor screens. BioRxiv.

Han, C., Liu, Y., Dai, R., Ismail, N., Su, W., and Li, B. (2020). Ferroptosis and its potential role in human diseases. Front. Pharmacol. *11*, 239.

Hart, T., Tong, A.H.Y., Chan, K., Van Leeuwen, J., Seetharaman, A., Aregger, M., Chandrashekhar, M., Hustedt, N., Seth, S., Noonan, A., et al. (2017). Evaluation and Design of Genome-Wide CRISPR/SpCas9 Knockout Screens. G3 (Bethesda) *7*, 2719–2727.

Hasegawa, M., Takahashi, H., Rajabi, H., Alam, M., Suzuki, Y., Yin, L., Tagde, A., Maeda, T., Hiraki, M., Sukhatme, V.P., et al. (2016). Functional interactions of the cystine/glutamate antiporter, CD44v and MUC1-C oncoprotein in triple-negative breast cancer cells. Oncotarget *7*, 11756–11769.

Hauser, M., Yan, R., Li, W., Repina, N.A., Schaffer, D.V., and Xu, K. (2018). The Spectrin-Actin-Based Periodic Cytoskeleton as a Conserved Nanoscale Scaffold and Ruler of the Neural Stem Cell Lineage. Cell Rep. *24*, 1512–1522.

Hensley, K., Hall, N., Subramaniam, R., Cole, P., Harris, M., Aksenov, M., Aksenova, M., Gabbita, S.P., Wu, J.F., and Carney, J.M. (1995). Brain regional correspondence between Alzheimer's disease histopathology and biomarkers of protein oxidation. J. Neurochem. *65*, 2146–2156.

Hill, A.J., McFaline-Figueroa, J.L., Starita, L.M., Gasperini, M.J., Matreyek, K.A., Packer, J., Jackson, D., Shendure, J., and Trapnell, C. (2018). On the design of CRISPR-based single-cell molecular screens. Nat. Methods *15*, 271–274.

Hiraiwa, M., Martin, B.M., Kishimoto, Y., Conner, G.E., Tsuji, S., and O'Brien, J.S. (1997). Lysosomal proteolysis of prosaposin, the precursor of saposins (sphingolipid activator proteins): its mechanism and inhibition by ganglioside. Arch. Biochem. Biophys. *341*, 17–24.

Hocsak, E., Szabo, V., Kalman, N., Antus, C., Cseh, A., Sumegi, K., Eros, K., Hegedus, Z.,

Gallyas, F., Sumegi, B., et al. (2017). PARP inhibition protects mitochondria and reduces ROS production via PARP-1-ATF4-MKP-1-MAPK retrograde pathway. Free Radic. Biol. Med. *108*, 770–784.

Höhn, A., Jung, T., Grimm, S., and Grune, T. (2010). Lipofuscin-bound iron is a major intracellular source of oxidants: role in senescent cells. Free Radic. Biol. Med. *48*, 1100–1108.

Horlbeck, M.A., Gilbert, L.A., Villalta, J.E., Adamson, B., Pak, R.A., Chen, Y., Fields, A.P., Park, C.Y., Corn, J.E., Kampmann, M., et al. (2016). Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation. Elife *5*.

Ihry, R.J., Salick, M.R., Ho, D.J., Sondey, M., Kommineni, S., Paula, S., Raymond, J., Henry, B., Frias, E., Wang, Q., et al. (2019). Genome-Scale CRISPR Screens Identify Human Pluripotency-Specific Genes. Cell Rep. *27*, 616-630.e6.

Imaizumi, Y., Okada, Y., Akamatsu, W., Koike, M., Kuzumaki, N., Hayakawa, H., Nihira, T., Kobayashi, T., Ohyama, M., Sato, S., et al. (2012). Mitochondrial dysfunction associated with increased oxidative stress and α-synuclein accumulation in PARK2 iPSC-derived neurons and postmortem brain tissue. Mol. Brain *5*, 35.

Isacson, O., Brekk, O.R., and Hallett, P.J. (2019). Novel results and concepts emerging from lipid cell biology relevant to degenerative brain aging and disease. Front. Neurol. *10*, 1053.

Ishimoto, T., Nagano, O., Yae, T., Tamada, M., Motohara, T., Oshima, H., Oshima, M., Ikeda, T., Asaba, R., Yagi, H., et al. (2011). CD44 variant regulates redox status in cancer cells by stabilizing the xCT subunit of system xc(-) and thereby promotes tumor growth. Cancer Cell *19*, 387–400.

Jain, I.H., Calvo, S.E., Markhard, A.L., Skinner, O.S., To, T.-L., Ast, T., and Mootha, V.K. (2020). Genetic screen for cell fitness in high or low oxygen highlights mitochondrial and lipid metabolism. Cell *181*, 716-727.e11.

Kaur, D., Yantiri, F., Rajagopalan, S., Kumar, J., Mo, J.Q., Boonplueang, R., Viswanath, V., Jacobs, R., Yang, L., Beal, M.F., et al. (2003). Genetic or pharmacological iron chelation prevents MPTP-induced neurotoxicity in vivo: a novel therapy for Parkinson's disease. Neuron *37*, 899–909.

Kügler, S., Straten, G., Kreppel, F., Isenmann, S., Liston, P., and Bähr, M. (2000). The X-linked inhibitor of apoptosis (XIAP) prevents cell death in axotomized CNS neurons in vivo. Cell Death Differ. *7*, 815–824.

Kunkle, B.W., Grenier-Boley, B., Sims, R., Bis, J.C., Damotte, V., Naj, A.C., Boland, A., Vronskaya, M., van der Lee, S.J., Amlie-Wolf, A., et al. (2019). Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. Nat. Genet. *51*, 414–430.

Kupershmidt, L., Amit, T., Bar-Am, O., Weinreb, O., and Youdim, M.B.H. (2012). Multi-target, neuroprotective and neurorestorative M30 improves cognitive impairment and reduces

Alzheimer's-like neuropathology and age-related alterations in mice. Mol. Neurobiol. *46*, 217–220.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics *9*, 559.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol. *10*, R25.

Le Pichon, C.E., Meilandt, W.J., Dominguez, S., Solanoy, H., Lin, H., Ngu, H., Gogineni, A., Sengupta Ghosh, A., Jiang, Z., Lee, S.-H., et al. (2017). Loss of dual leucine zipper kinase signaling is protective in animal models of neurodegenerative disease. Sci. Transl. Med. *9*.

Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z., and Zhang, B. (2019). WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. Nucleic Acids Res. *47*, W199–W205.

Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. Nature *443*, 787–795.

Li, M., Zhao, X., Wang, W., Shi, H., Pan, Q., Lu, Z., Perez, S.P., Suganthan, R., He, C., Bjørås, M., et al. (2018). Ythdf2-mediated m6A mRNA clearance modulates neural development in mice. Genome Biol. *19*, 69.

Luu, B., Ellisor, D., and Zervas, M. (2011). The lineage contribution and role of Gbx2 in spinal cord development. PLoS ONE *6*, e20940.

Mair, B., Tomic, J., Masud, S.N., Tonge, P., Weiss, A., Usaj, M., Tong, A.H.Y., Kwan, J.J., Brown, K.R., Titus, E., et al. (2019). Essential gene profiles for human pluripotent stem cells identify uncharacterized genes and substrate dependencies. Cell Rep. *27*, 599-615.e12.

Martinez-Morales, P.L., Quiroga, A.C., Barbas, J.A., and Morales, A.V. (2010). SOX5 controls cell cycle progression in neural progenitors by interfering with the WNT-beta-catenin pathway. EMBO Rep. *11*, 466–472.

Mathys, H., Davila-Velderrain, J., Peng, Z., Gao, F., Mohammadi, S., Young, J.Z., Menon, M., He, L., Abdurrob, F., Jiang, X., et al. (2019). Single-cell transcriptomic analysis of Alzheimer's disease. Nature *570*, 332–337.

Ma, Q. (2013). Role of nrf2 in oxidative stress and toxicity. Annu. Rev. Pharmacol. Toxicol. *53*, 401–426.

Méndez-Gómez, H.R., and Vicario-Abejón, C. (2012). The homeobox gene Gsx2 regulates the self-renewal and differentiation of neural stem cells and the cell fate of postnatal progenitors. PLoS ONE *7*, e29799.

Miyaoka, Y., Chan, A.H., Judge, L.M., Yoo, J., Huang, M., Nguyen, T.D., Lizarraga, P.P., So, P.-L., and Conklin, B.R. (2014). Isolation of single-base genome-edited human iPS cells without antibiotic selection. Nat. Methods *11*, 291–293.

Mochizuki, Y., Park, M.K., Mori, T., and Kawashima, S. (1995). The difference in

autofluorescence features of lipofuscin between brain and adrenal. Zool. Sci. *12*, 283–288.

Moreno-García, A., Kun, A., Calero, O., Medina, M., and Calero, M. (2018). An Overview of the Role of Lipofuscin in Age-Related Neurodegeneration. Front. Neurosci. *12*, 464.

van Muiswinkel, F.L., de Vos, R.A.I., Bol, J.G.J.M., Andringa, G., Jansen Steur, E.N.H., Ross, D., Siegel, D., and Drukarch, B. (2004). Expression of NAD(P)H:quinone oxidoreductase in the normal and Parkinsonian substantia nigra. Neurobiol. Aging *25*, 1253–1262.

Muto, Y., Nishiyama, M., Nita, A., Moroishi, T., and Nakayama, K.I. (2017). Essential role of FBXL5-mediated cellular iron homeostasis in maintenance of hematopoietic stem cells. Nat. Commun. *8*, 16114.

Nacarelli, T., Azar, A., and Sell, C. (2014). Inhibition of mTOR Prevents ROS Production Initiated by Ethidium Bromide-Induced Mitochondrial DNA Depletion. Front Endocrinol (Lausanne) *5*, 122.

Ndayisaba, A., Kaindlstorfer, C., and Wenning, G.K. (2019). Iron in Neurodegeneration - Cause or Consequence? Front. Neurosci. *13*, 180.

Ou, Y., Wang, S.-J., Li, D., Chu, B., and Gu, W. (2016). Activation of SAT1 engages polyamine metabolism with p53-mediated ferroptotic responses. Proc Natl Acad Sci USA *113*, E6806–E6812.

Pang, X.P., Hershman, J.M., Chung, M., and Pekary, A.E. (1989). Characterization of tumor necrosis factor-alpha receptors in human and rat thyroid cells and regulation of the receptors by thyrotropin. Endocrinology *125*, 1783–1788.

Patel, M. (2016). Targeting oxidative stress in central nervous system disorders. Trends Pharmacol. Sci. *37*, 768–778.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. Nat. Methods *14*, 417–419.

Piñero, J., Queralt-Rosinach, N., Bravo, À., Deu-Pons, J., Bauer-Mehren, A., Baron, M., Sanz, F., and Furlong, L.I. (2015). DisGeNET: a discovery platform for the dynamical exploration of human diseases and their genes. Database (Oxford) *2015*, bav028.

Puri, V., Jefferson, J.R., Singh, R.D., Wheatley, C.L., Marks, D.L., and Pagano, R.E. (2003). Sphingolipid storage induces accumulation of intracellular cholesterol by stimulating SREBP-1 cleavage. J. Biol. Chem. *278*, 20961–20970.

Raina, A.K., Templeton, D.J., Deak, J.C., Perry, G., and Smith, M.A. (1999). Quinone reductase (NQO1), a sensitive redox indicator, is increased in Alzheimer's disease. Redox Rep. *4*, 23–27.

Ramsey, C.P., Glass, C.A., Montgomery, M.B., Lindl, K.A., Ritson, G.P., Chia, L.A., Hamilton, R.L., Chu, C.T., and Jordan-Sciutto, K.L. (2007). Expression of Nrf2 in neurodegenerative diseases. J. Neuropathol. Exp. Neurol. *66*, 75–85.

Reczek, C.R., Birsoy, K., Kong, H., Martínez-Reyes, I., Wang, T., Gao, P., Sabatini, D.M., and

Chandel, N.S. (2017). A CRISPR screen identifies a pathway required for paraquat-induced cell death. Nat. Chem. Biol. *13*, 1274–1279.

Regev, A., Teichmann, S.A., Lander, E.S., Amit, I., Benoist, C., Birney, E., Bodenmiller, B., Campbell, P., Carninci, P., Clatworthy, M., et al. (2017). The human cell atlas. Elife *6*.

Reho, J.J., Guo, D.-F., and Rahmouni, K. (2019). Mechanistic target of rapamycin complex 1 signaling modulates vascular endothelial function through reactive oxygen species. J. Am. Heart Assoc. *8*, e010662.

Ross, D., and Siegel, D. (2017). Functions of NQO1 in cellular protection and coq10 metabolism and its potential role as a redox sensitive molecular switch. Front. Physiol. *8*, 595.

Rouault, T.A. (2013). Iron metabolism in the CNS: implications for neurodegenerative diseases. Nat. Rev. Neurosci. *14*, 551–564.

Rust, M.J., Bates, M., and Zhuang, X. (2006). Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). Nat. Methods *3*, 793–795.

Sandhoff, K., and Harzer, K. (2013). Gangliosides and gangliosidoses: principles of molecular and metabolic pathogenesis. J. Neurosci. *33*, 10195–10208.

SantaCruz, K.S., Yazlovitskaya, E., Collins, J., Johnson, J., and DeCarli, C. (2004). Regional NAD(P)H:quinone oxidoreductase activity in Alzheimer's disease. Neurobiol. Aging *25*, 63–69.

Schmitt-John, T., Drepper, C., Mussmann, A., Hahn, P., Kuhlmann, M., Thiel, C., Hafner, M., Lengeling, A., Heimann, P., Jones, J.M., et al. (2005). Mutation of Vps54 causes motor neuron disease and defective spermiogenesis in the wobbler mouse. Nat. Genet. *37*, 1213–1215.

Schulz, A., Sekine, Y., Oyeyemi, M.J., Abrams, A.J., Basavaraju, M., Han, S.M., Groth, M., Morrison, H., Strittmatter, S.M., and Hammarlund, M. (2020). The stress-responsive gene GDPGP1/mcp-1 regulates neuronal glycogen metabolism and survival. J. Cell Biol. *219*.

Seibler, P., Burbulla, L.F., Dulovic, M., Zittel, S., Heine, J., Schmidt, T., Rudolph, F., Westenberger, A., Rakovic, A., Münchau, A., et al. (2018). Iron overload is accompanied by mitochondrial and lysosomal dysfunction in WDR45 mutant cells. Brain *141*, 3052–3064.

Shin, Y.J., Cho, D.Y., Chung, T.Y., Han, S.B., Hyon, J.Y., and Wee, W.R. (2011). Rapamycin reduces reactive oxygen species in cultured human corneal endothelial cells. Curr. Eye Res. *36*, 1116–1122.

Skouta, R., Dixon, S.J., Wang, J., Dunn, D.E., Orman, M., Shimada, K., Rosenberg, P.A., Lo, D.C., Weinberg, J.M., Linkermann, A., et al. (2014). Ferrostatins inhibit oxidative lipid damage and cell death in diverse disease models. J. Am. Chem. Soc. *136*, 4551–4556.

Slowik, A., Tomik, B., Wolkow, P.P., Partyka, D., Turaj, W., Malecki, M.T., Pera, J., Dziedzic, T., Szczudlik, A., and Figlewicz, D.A. (2006). Paraoxonase gene polymorphisms and sporadic ALS. Neurology *67*, 766–770.

Smith, C.D., Carney, J.M., Starke-Reed, P.E., Oliver, C.N., Stadtman, E.R., Floyd, R.A., and

Markesbery, W.R. (1991). Excess brain protein oxidation and enzyme dysfunction in normal aging and in Alzheimer disease. Proc Natl Acad Sci USA *88*, 10540–10543.

Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. [version 2; peer review: 2 approved]. F1000Res. *4*, 1521.

Stockwell, B.R., Friedmann Angeli, J.P., Bayir, H., Bush, A.I., Conrad, M., Dixon, S.J., Fulda, S., Gascón, S., Hatzios, S.K., Kagan, V.E., et al. (2017). Ferroptosis: A regulated cell death nexus linking metabolism, redox biology, and disease. Cell *171*, 273–285.

Subbarao, K.V., Richardson, J.S., and Ang, L.C. (1990). Autopsy samples of Alzheimer's cortex show increased peroxidation in vitro. J. Neurochem. *55*, 342–345.

Sudarshan, S., Sourbier, C., Kong, H.-S., Block, K., Valera Romero, V.A., Yang, Y., Galindo, C., Mollapour, M., Scroggins, B., Goode, N., et al. (2009). Fumarate hydratase deficiency in renal cancer induces glycolytic addiction and hypoxia-inducible transcription factor 1alpha stabilization by glucose-dependent generation of reactive oxygen species. Mol. Cell. Biol. *29*, 4080–4090.

Taguchi, K., Fujikawa, N., Komatsu, M., Ishii, T., Unno, M., Akaike, T., Motohashi, H., and Yamamoto, M. (2012). Keap1 degradation by autophagy for the maintenance of redox homeostasis. Proc Natl Acad Sci USA *109*, 13561–13566.

Tian, R., Gachechiladze, M.A., Ludwig, C.H., Laurie, M.T., Hong, J.Y., Nathaniel, D., Prabhu, A.V., Fernandopulle, M.S., Patel, R., Abshari, M., et al. (2019). CRISPR Interference-Based Platform for Multimodal Genetic Screens in Human iPSC-Derived Neurons. Neuron *104*, 239-255.e12.

Tsang, A.H.K., Lee, Y.-I., Ko, H.S., Savitt, J.M., Pletnikova, O., Troncoso, J.C., Dawson, V.L., Dawson, T.M., and Chung, K.K.K. (2009). S-nitrosylation of XIAP compromises neuronal survival in Parkinson's disease. Proc Natl Acad Sci USA *106*, 4900–4905.

Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S., Gill, S., Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a cancer dependency map. Cell *170*, 564-576.e16.

Unsain, N., Higgins, J.M., Parker, K.N., Johnstone, A.D., and Barker, P.A. (2013). XIAP regulates caspase activity in degenerating axons. Cell Rep. *4*, 751–763.

Wang, C., Chen, S., Yeo, S., Karsli-Uzunbas, G., White, E., Mizushima, N., Virgin, H.W., and Guan, J.-L. (2016). Elevated p62/SQSTM1 determines the fate of autophagy-deficient neural stem cells by increasing superoxide. J. Cell Biol. *212*, 545–560.

Wang, C., Ward, M.E., Chen, R., Liu, K., Tracy, T.E., Chen, X., Xie, M., Sohn, P.D., Ludwig, C., Meyer-Franke, A., et al. (2017). Scalable Production of iPSC-Derived Human Neurons to Identify Tau-Lowering Compounds by High-Content Screening. Stem Cell Reports *9*, 1221–1233.

Wang, Y., Santa-Cruz, K., DeCarli, C., and Johnson, J.A. (2000). NAD(P)H:quinone oxidoreductase activity is increased in hippocampal pyramidal neurons of patients with Aalzheimer's disease. Neurobiol. Aging *21*, 525–531.

Ward, D.M., and Cloonan, S.M. (2019). Mitochondrial iron in human health and disease. Annu. Rev. Physiol. *81*, 453–482.

Ward, M.E., Chen, R., Huang, H.-Y., Ludwig, C., Telpoukhovskaia, M., Taubes, A., Boudin, H., Minami, S.S., Reichert, M., Albrecht, P., et al. (2017). Individuals with progranulin haploinsufficiency exhibit features of neuronal ceroid lipofuscinosis. Sci. Transl. Med. *9*.

Weber, R.A., Yen, F.S., Nicholson, S.P.V., Alwaseem, H., Bayraktar, E.C., Alam, M., Timson, R.C., La, K., Abu-Remaileh, M., Molina, H., et al. (2020). Maintaining iron homeostasis is the key role of lysosomal acidity for cell proliferation. Mol. Cell *77*, 645-655.e7.

Welsbie, D.S., Yang, Z., Ge, Y., Mitchell, K.L., Zhou, X., Martin, S.E., Berlinicke, C.A., Hackler, L., Fuller, J., Fu, J., et al. (2013). Functional genomic screening identifies dual leucine zipper kinase as a key mediator of retinal ganglion cell death. Proc Natl Acad Sci USA *110*, 4045–4050.

Weltner, J., Balboa, D., Katayama, S., Bespalov, M., Krjutškov, K., Jouhilahti, E.-M., Trokovic, R., Kere, J., and Otonkoski, T. (2018). Human pluripotent reprogramming with CRISPR activators. Nat. Commun. *9*, 2643.

Wojcik, M., Hauser, M., Li, W., Moon, S., and Xu, K. (2015). Graphene-enabled electron microscopy and correlated super-resolution microscopy of wet cells. Nat. Commun. *6*, 7384.

Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biol. *19*, 15.

Xu, D., Bureau, Y., McIntyre, D.C., Nicholson, D.W., Liston, P., Zhu, Y., Fong, W.G., Crocker, S.J., Korneluk, R.G., and Robertson, G.S. (1999). Attenuation of ischemia-induced cellular and behavioral deficits by X chromosome-linked inhibitor of apoptosis protein overexpression in the rat hippocampus. J. Neurosci. *19*, 5026–5033.

Xu, M., Liu, K., Swaroop, M., Sun, W., Dehdashti, S.J., McKew, J.C., and Zheng, W. (2014). A phenotypic compound screening assay for lysosomal storage diseases. J. Biomol. Screen. *19*, 168–175.

Yadav, R.S., and Tiwari, N.K. (2014). Lipid integration in neurodegeneration: an overview of Alzheimer's disease. Mol. Neurobiol. *50*, 168–176.

Yang, W.S., SriRamaratnam, R., Welsch, M.E., Shimada, K., Skouta, R., Viswanathan, V.S., Cheah, J.H., Clemons, P.A., Shamji, A.F., Clish, C.B., et al. (2014). Regulation of ferroptotic cancer cell death by GPX4. Cell *156*, 317–331.

Yao, Y., Zhukareva, V., Sung, S., Clark, C.M., Rokach, J., Lee, V.M.Y., Trojanowski, J.Q., and Praticò, D. (2003). Enhanced brain levels of 8,12-iso-iPF2alpha-VI differentiate AD from

frontotemporal dementia. Neurology *61*, 475–478.

Yilmaz, A., Peretz, M., Aharony, A., Sagi, I., and Benvenisty, N. (2018). Defining essential genes for human pluripotent stem cells by CRISPR-Cas9 screening in haploid cells. Nat. Cell Biol. *20*, 610–619.

Yuan, H., Li, X., Zhang, X., Kang, R., and Tang, D. (2016). Identification of ACSL4 as a biomarker and contributor of ferroptosis. Biochem. Biophys. Res. Commun. *478*, 1338–1343.

Zhao, L.Y., Niu, Y., Santiago, A., Liu, J., Albert, S.H., Robertson, K.D., and Liao, D. (2006). An EBF3-mediated transcriptional program that induces cell cycle arrest and apoptosis. Cancer Res. *66*, 9445–9452.

Zhitomirsky, B., and Assaraf, Y.G. (2015). Lysosomal sequestration of hydrophobic weak base chemotherapeutics triggers lysosomal biogenesis and lysosome-dependent cancer multidrug resistance. Oncotarget *6*, 1143–1156.

FIGURES

Figure 1 (legend overleaf)

Figure 1. Genome-wide CRISPRi and CRISPRa screens in human iPSC-derived neurons identify regulators of neuronal survival

(A) Strategy for generating the CRISPRa iPSC line: an inducible CRISPRa construct, CAG promoter-driven DHFR-dCas9-VPH, was stably integrated into the CLYBL safe harbor locus through TALEN-mediated knock-in. dCas9, catalytically dead Cas9. VPH, activator domains containing 4X repeats of VP48, P65, and HSF1.

(B) Functional validation of CRISPRa activity. qPCR quantification of the relative fold change of *CXCR4* mRNA levels in CRISPRa-neurons expressing a *CXCR4* sgRNA as compared to a non-targeting control sgRNA in the presence or absence of trimethoprim (TMP), which stabilizes the DHFR degron (mean +/s sd, n = 3). *CXCR4* levels were normalized to the housekeeping gene *ACTB*.

(C) Strategy for neuronal survival screens. CRISPRi/a iPSCs were transduced with genome-wide sgRNA libraries, containing ~100,000 sgRNAs targeting ~19,000 protein-coding genes and ~1,800 non-targeting control sgRNAs. TMP was added to CRISPRa neurons from Day 0 to induce CRISPRa activity. Frequencies of cells expressing a given sgRNA were determined by next-generation sequencing for Day 10 neurons and Day -3 iPSCs.

(D) Volcano plots summarizing knockdown or overexpression phenotypes and statistical significance (Mann-Whitney U test) for genes targeted in the CRISPRi (left) and CRISPRa (right) screens. Dashed lines: gene score cutoff for hit genes (FDR = 0.05, see Methods)

(E) Comparing hits from CRISPRi and CRISPRa screens. Hit genes with protective or toxic phenotypes in either screen are shown in red or blue, respectively. Genes that are hits in both screens are shown in orange.

(F) Gene Ontology (GO) term enrichment analysis for the top 100 hit genes with protective or toxic phenotypes in CRISPRi (left) and CRISPRa (right) survival screens. Significantly enriched Biological Process terms (FDR < 0.01) are shown.

(G) Expression levels of hit genes and non-hit genes from CRISPRi (left) or CRISPRa (right) screens are shown, binned by order of magnitude.

(H) Percentage of hits with paralogs in CRISPRi and CRISPRa survival screens. A list of human paralog genes was obtained from a previous study (Dandage and Landry, 2019). P value was calculated using Fisher's exact test.

Figure 2 (legend overleaf)

**Figure 2. Genome-wide CRISPRi and CRISPRa screens in human iPSC-derived neurons
identify regulators of oxidative stress survival and redox homeostasis**

(A) Screening strategy. First, survival-based screens were conducted to identify modifiers of
neuronal survival under mild oxidative stress induced by anti-oxidant removal from the neuronal
medium (-AO). Second, FACS-based screens were conducted for modifiers of ROS and lipid

peroxidation levels. Last, secondary screens for lysosomal status and labile iron levels were conducted to further characterize hit genes.

(B) Comparison of gene scores in +AO and -AO conditions for CRISPRi survival screens.

(C) Pathway for selenocysteine incorporation into GPX4. Hit genes are highlighted in orange.

(D) Ranked gene scores from the ROS screen and the lipid peroxidation screen. High-signal hits are shown in red and low signal hits in blue. Genes discussed in the paper are highlighted in orange.

(E) GO term enrichment analysis for the top 100 high-signal and low-signal hits in the ROS screen (left) and the lipid peroxidation screen (right). Significantly enriched Biological Process terms (FDR<0.01) are shown.

(F) Gene scores from the lysosome and iron secondary screens. Genes are color-coded by pathways based on GO annotation.

(G) Heatmap showing gene scores across screens (rows) for genes that are among the top 20 high-signal or low-signal hits in at least one screen (columns). Rows and columns are hierarchically clustered. Genes are color-coded by pathways based on GO annotation.

Figure 3. Loss of prosaposin induces ROS and lipid peroxidation in neurons and causes neuronal ferroptosis in the absence of antioxidants

(A) Prosaposin is processed in the lysosome by cathepsin D (encoded by *CTSD*) into saposin subunits, which function together with GM2A as activators for glycosphingolipid degradation.

(B) Results from the ROS and lipid peroxidation screens, highlighting *PSAP* and the related genes *CTSD* and *GM2A*.

(C) Western blot showing the depletion of prosaposin in the *PSAP* KO iPSC line.

(D) Representative immunofluorescence microscopy images showing the loss of prosaposin in *PSAP* KO neurons. WT and *PSAP* KO neurons were fixed and stained by antibodies against prosaposin (shown in green) and the neuronal marker Tuj1 (shown in purple). Nuclei were counterstained by Hoechst, shown in blue. Scale bars, 20 μm.

(E) ROS levels (as indicated by CellRox) and lipid peroxidation levels (as indicated by Liperfluo and C11-BODIPY) in WT and *PSAP* KO neurons, measured by flow cytometry.

(F) ROS levels in iPSCs and HEK293 cells in WT and *PSAP* KO backgrounds (*PSAP* KO for iPSCs and *PSAP* knockdown by CRISPRi in HEK293s), measured by CellRox via flow cytometry.

(G) Survival curves for WT and *PSAP* KO neurons cultured in normal neuronal medium (+AO) or medium lack of antioxidants (-AO), quantified by imaging using Hoechst stain (all cells) and propidium iodide (PI) (dead cells). Survival fraction is calculated as (total cell count - dead cell count) / total cell count. Data is shown as mean +/- sd, n = 4 culture wells per group. 16 imaging fields were averaged for each well.

(H) Survival fractions of WT and *PSAP* KO neurons treated with different cell death inhibitors under +AO or -AO conditions, quantified by imaging in the same way as for G. Data is shown as mean +/- sd, n = 32 imaging fields per group.

(I) Representative images for the Hoechst (shown in blue) and PI (shown in red) staining in H. Scale bars, 50 μm.

Figure 4 (legend overleaf)

Figure 4. Loss of prosaposin disrupts glycosphingolipid degradation, leads to lipofuscin formation, iron accumulation and impaired autophagy

(A) Untargeted lipidomics comparing abundances of different lipid species in WT and *PSAP* KO neurons. P values were calculated using Student's t-test (n = 3 replicates per group). Dashed line, P value cutoff for FDR < 0.01. Glycosphingolipids are shown in orange and ether lipids in green.

(B) Heatmap showing the abundances of significantly increased or decreased lipids in *PSAP* KO neurons as compared to WT (FDR < 0.01). Enrichment P values for glycosphingolipids and ether lipids were calculated using Fisher's exact test. Lipid abundances were Z score-normalized across samples.

(C) Representative immunofluorescence microscopy images for WT and *PSAP* KO neurons stained with LAMP2 antibodies (shown in green) and GM1 antibodies (shown in red). Nuclei were counterstained by Hoechst, shown in blue. Scale bar, 10 μm.

(D) Representative immunofluorescence microscopy images for WT and *PSAP* KO iPSCs stained with GM1 antibodies (shown in red). Nuclei were counterstained by Hoechst, shown in blue. Scale bar, 20 μm

(E,F) Lysotracker signals measured by flow cytometry in WT and *PSAP* KO neurons (E), WT and *PSAP* KO iPSCs (F, *left*) and WT (no sgRNA) and *PSAP* knockdown (*PSAP* sgRNA) HEK293 cells (F, *right*).

(G) Two-color STORM super-resolution images of *PSAP* KO neurons immunolabeled for LAMP2 (shown in green) and GM1 (shown in magenta). Scale bars, 2 μm.

(H) Electron microscopy images for WT and *PSAP* KO neurons. The red arrow points to a representative lipofuscin structure. Scale bar, 1 μm.

(I) Representative images for autofluorescence in WT and *PSAP* KO neurons. Excitation, UV (405 nm). Emission, FITC (525/20 nm). Scale bar, 10 μm.

(J) Labile iron levels in WT and *PSAP* KO neurons. Neurons were stained with the iron indicators FeRhoNox-1 (*left*) or FerroOrange (*right*) and fluorescence was measured by flow cytometry.

(K) Representative fluorescence microscopy images for WT neurons, *PSAP* KO neurons, and *PSAP* KO neurons treated with 10 μM DFO for 3 days, stained with Lysotracker (shown in

green) and FeRhoNox-1 (shown in red). Nuclei were counterstained by Hoechst, shown in blue. Scale bar, 10 μm.

(L) Representative fluorescence microscopy images for WT and *PSAP* KO neurons under the - AO condition stained with Lysotracker (shown in green) and Liperfluo (shown in red). Nuclei were counterstained by Hoechst, shown in blue. Scale bar, 10 μm.

(M) Western blot showing protein levels of phosphatidylethanolamine (PE)-conjugated LC3B (LC3B-II) and unconjugated LC3B (LC3B-I) in WT and *PSAP* KO neurons in the absence or presence of Bafilomycin A1 (BafA1). β-Actin was used as a loading control. Ratios of LC3B-II to LC3B-I are indicated at the bottom.

(N) Quantification of LC3B-II / LC3B-I ratios for WT and *PSAP* KO neurons (mean +/- sd, n = 6 independent experiments). P value was calculated using Student's t-test.

(O) Representative immunofluorescence microscopy images for WT and *PSAP* KO neurons, stained with LC3B antibodies (shown in red). Nuclei were counterstained by Hoechst, shown in blue. Scale bar, 20 μm.

(P) A model for the mechanism linking prosaposin loss to neuronal ferroptosis. Loss of saposins blocks glycosphingolipid degradation in the lysosome. The build-up of glycosphingolipids leads to lipofuscin formation, which accumulates iron and generates ROS through the Fenton reaction. ROS then peroxidize lipids and cause neuronal ferroptosis in the absence of antioxidants. Other consequences are cholesterol accumulation and impaired autophagy.

Figure 5 (legend overleaf)

Figure 5. CROP-seq reveals transcriptomic responses to perturbations of neurodegenerative disease-associated genes in human iPSC-derived neurons

(A) Hit genes from our screens that are also associated with neurodegenerative diseases were targeted in a CROP-seq screen to detect their knockdown or overexpression effects on gene expression at single-cell resolution in human iPSC-derived neurons.

(B) Summary of on-target knockdown by CRISPRi (*top*) or overexpression by CRISPRa (*bottom*) for all target genes in the CROP-seq libraries. log2FC represents the log2 fold change of the mean expression of a target gene in perturbed cells (i.e. cells expressing sgRNAs targeting that gene) compared to unperturbed cells (i.e. cells expressing non-targeting control sgRNAs). P values were calculated by the Wilcoxon rank-sum test. Target genes are ranked by their expression in unperturbed cells.

(C&D) Examples of CROP-seq results showing on-target knockdown (*TUBB4A* in CRISPRi, C) or overexpression (*NQO1* in CRISPRa, D) and the classification method, shown in a two-dimensional UMAP projection.

(E) Pairwise similarities of differentially expressed genes among perturbations. Similarity scores were determined by the OrderedList package in R (see Methods). Genes in clusters with high similarity are labeled.

(F & G) Eigengene expression of gene modules identified from WGCNA analysis in cells containing different perturbations relative to unperturbed cells (F). Enriched pathways in each module are shown in G.

Figure 6. Overexpression of *NQO1* induces unexpected transcriptome changes in human iPSC-derived neurons that provide hypotheses for its toxicity

(A) Transcriptomic changes induced by *NQO1* overexpression in neurons. Significantly upregulated or downregulated genes (FDR < 0.01) are shown in red or blue, respectively.

(B) Pathway analysis showing enriched pathways in upregulated and downregulated genes in *NQO1*-overexpressing neurons.

(C) String-db association networks of selected pathways enriched in upregulated and downregulated genes. Genes with stronger associations are connected by thicker lines. Colors and sizes of nodes reflect $\log_2$-fold changes ($\log_2$FCs) and significance ($-\log_{10}$P) of differentially expressed genes, respectively.

(D) Hypotheses for potential mechanisms of neurotoxicity of *NQO1* overexpression.

Figure 7. CRISPRbrain, a Data Commons for functional genomics screens in differentiated human cell types.

(A) CRISPRbrain (https://crisprbrain.org) is a Data Commons organizing results from genetic screens for different phenotypes in different human cell types, by different research groups.

(B) Screens can be browsed and searched based on a range of parameters, and full text search.

(C,D) Survival- and FACS-based screens ("simple screens") can be visualized and explored interactively, and compared pairwise.

(E,F) RNA-Seq-based screens can be explored one perturbed gene at a time as MA plots (E), or globally in a hierarchically clustered heatmap.

Figure S1. Karyotyping of the monoclonal CRISPRa-iPSC line, gene paralogs in CRISPRi and CRISPRa hits, and comparison of CRISPRi and CRISPRa screen results for neuronal survival with other published survival screens for different cell types (related to Figure 1)

(A) A normal karyotype was confirmed for the monoclonal CRISPRa-iPSC line.

(B) Venn diagrams comparing CRISPRi and CRISPRa screen results for neuronal survival from this paper with other published survival screens for different human cell types. For CRISPRi, hit genes with toxic phenotypes for the survival of neurons were compared with those for cancer cells ('gold-standard' essential genes, Hart et al., 2017) and pluripotent stem cells (integrated from Ihry et al., 2019; Mair et al., 2019 and Yilmaz et al., 2018); genes that were identified as essential in more than one studies were retained for comparison). Protective hits for the survival of neurons were compared with those for human pluripotent stem cells (integrated from Ihry et al., 2019; Yilmaz et al., 2018. Genes that were identified as essential in both studies were retained for comparison). For CRISPRa, hits were compared with our published survival screen in K562 cells (Horlbeck et al., 2016) reanalyzed using our MAGeCK-iNC pipeline.

Figure S2. Comparing CRISPRa survival screens in +AO and -AO conditions (related to Figure 2)

Each dot represents one gene, and its gene score in the +AO screen was plotted on the x-axis and gene score in the -AO screen on the y-axis. The Pearson correlation coefficient is shown.

Figure S3 (legend overleaf)

Figure S3. Characterization of *PSAP* KO neurons (related to Figure 4)

(A) Labile iron levels in WT and *PSAP* KO neurons with or without DFO treatment. Cells were stained by FeRhoNox-1 and fluorescence was quantified by flow cytometry. Median signal intensities are indicated.

(B) Labile iron levels in WT neurons, *PSAP* KO neurons, and *PSAP* KO neurons with DFO treatment. Cells were stained by Calcein and fluorescence was quantified by flow cytometry. Median signal intensities are indicated.

(C) Gene expression changes in *PSAP* KO neurons as compared to WT. Genes that are significantly upregulated and downregulated in *PSAP* KO neurons are shown in red and blue, respectively (FDR < 0.05). The top 50 up- and down-regulated genes are labeled, and within this set, genes involved in the cholesterol biosynthesis pathway are highlighted in orange.

(D) GO term enrichment analysis for significantly up- and down-regulated genes (FDR < 0.05) in *PSAP* KO neurons. Significantly enriched Biological Process terms are shown (FDR < 0.01).

(E) Cholesterol levels measured by flow cytometry in filipin-stained WT and *PSAP* KO neurons in +AO and -AO conditions.

(F) Representative fluorescence microscopy images of WT and PSAP KO neurons stained with filipin (shown in cyan) and LAMP2 antibodies (shown for *PSAP* KO neurons, in red). Scale bar, 10 μm.

Figure S4. Examples of the CROP-seq classification method, and shared transcriptomic signatures of *VPS54*, *PAXIP1*, and *PON2* knockdown in human iPSC-derived neurons (related to Figure 5)

(A&B) CROP-seq examples showing the application of the outlier detection-based classification method in cases where two sgRNAs targeting the same gene had heterogeneous efficacy (A, *SOX5* in CRISPRa) or the expression level of the target gene was too low to quantify knockdown level (B, *ZNF592* in CRISPRi).

(C)Transcriptomic changes induced by knockdown of VPS54 (*left*), PAXIP1 (*middle*), and PON2 (*right*) in neurons. For each perturbation, the top 200 upregulated and downregulated genes compared to control (i.e. unperturbed cells) are shown in red and blue, respectively. Within this set, shared genes among all three perturbations are highlighted in green.

METHODS

Human iPSCs culture and neuronal differentiation

Human iPSCs (male WTC11 background (Miyaoka et al., 2014)) were cultured in StemFlex Medium (GIBCO/Thermo Fisher Scientific; Cat. No. A3349401) in plates or dishes coated with Growth Factor Reduced, Phenol Red-Free, LDEV-Free Matrigel Basement Membrane Matrix (Corning; Cat. No. 356231) diluted 1:100 in Knockout DMEM (GIBCO/ v; Cat. No. 10829-018). StemFlex Medium was replaced every other day or every day once cells reached 50% confluence. When 80%–90% confluent, cells were dissociated using StemPro Accutase Cell Dissociation Reagent (GIBCO/Thermo Fisher Scientific; Cat. No. A11105-01) at 37°C for 5 min, centrifuged at 200 g for 5 min, resuspended in StemFlex Medium supplemented with 10 nM Y-27632 dihydrochloride ROCK inhibitor (Tocris; Cat. No. 125410) and plated onto Matrigel-coated plates or dishes at desired number. Studies with human iPSCs at UCSF were approved by the Human Gamete, Embryo, and Stem Cell Research (GESCR) Committee.

The CRISPRi- and CRISPRa-iPSC lines used in this study were engineered to express mNGN2 under a doxycycline-inducible system in the AAVS1 safe harbor locus. For their neuronal differentiation, we followed our previously described protocol (Tian et al., 2019). Briefly, iPSCs were pre-differentiated in matrigel-coated plates or dishes in N2 Pre-Differentiation Medium containing the following: Knockout DMEM/F12 (GIBCO/Thermo Fisher Scientific; Cat. No. 12660-012) as the base, 1X MEM Non-Essential Amino Acids (GIBCO/Thermo Fisher Scientific; Cat. No. 11140-050), 1X N2 Supplement (GIBCO/ Thermo Fisher Scientific; Cat. No. 17502-048), 10 ng/mL NT-3 (PeproTech; Cat. No. 450-03), 10ng/mL BDNF (PeproTech; Cat. No. 450-02), 1 μg/mL Mouse Laminin (Thermo Fisher Scientific; Cat. No. 23017-015), 10nM ROCK inhibitor, and 2 μg/mL doxycycline to induce expression of mNGN2. After three days, on the day referred to hereafter as Day 0, pre-differentiated cells were re-plated into BioCoat Poly-D-Lysine-coated plates or dishes (Corning; assorted Cat. No.) in regular Neuronal Medium, which we will refer to as +AO Neuronal Medium, containing the following: half DMEM/F12 (GIBCO/Thermo Fisher Scientific; Cat. No. 11320-033) and half Neurobasal-A (GIBCO/Thermo Fisher Scientific; Cat. No. 10888-022) as the base, 1X MEM Non-Essential Amino Acids, 0.5X GlutaMAX Supplement (GIBCO/Thermo Fisher Scientific; Cat. No. 35050-061), 0.5X N2 Supplement, 0.5X B27 Supplement (GIBCO/Thermo Fisher Scientific; Cat. No. 17504-044), 10ng/mL NT-3, 10ng/mL BDNF and 1 μg/mL Mouse Laminin. For -AO experiments, we used a medium we refer to as -AO Neuronal Medium, in which B-27 Supplement minus antioxidants (GIBCO/Thermo Fisher Scientific; Cat. No. 10889-038) was used instead of regular B27 in the +AO Neuronal Medium. Neuronal Medium was half-replaced every week.

Molecular cloning

The CLYBL-targeting inducible CRISPRa vector pRT43 containing CAG-driven DHFR-dCas9-VPH-T2A-EGFP was generated by sub-cloning DHFR-dCas9-VPH-T2A-EGFP from plasmid PB-CAG-DDdCas9VPH-T2A-GFP-IRES-Neo to the downstream of a CAG promoter in CLYBL-targeting plasmid pUCM-CLYBL-hNIL digested by SalI and EcoRV. (PB-CAG-DDdCas9VPH-T2A-GFP-IRES-Neo was a gift from Timo Otonkoski (Addgene plasmid # 102886 ; http://n2t.net/addgene:102886 ; RRID:Addgene_102886) and pUCM-CLYBL-hNIL

was a gift from Michael Ward (Addgene plasmid # 105841 ; http://n2t.net/addgene:105841 ; RRID:Addgene_105841))

CRISPRa-iPSC cell line generation
WTC11 iPSCs harboring a single-copy of doxycycline-inducible mouse NGN2 at the AAVS1 locus (Ngn2-iPSCs, (Wang et al., 2017)) were used as the parental iPSC line for further genetic engineering. iPSCs were transfected with pRT43 containing DHFR-dCas9-VPH and TALENS targeting the human CLYBL intragenic safe harbor locus (between exons 2 and 3) (pZT-C13-R1 and pZT- C13-L1, gifts from Jizhong Zou (Addgene plasmid # 62196 ; http://n2t.net/addgene:62196 ; RRID:Addgene_62196, and Addgene plasmid # 62197 ; http://n2t.net/addgene:62197 ; RRID:Addgene_62197) using Lipofectamine Stem (Invitrogen/Thermo Fisher Scientific; Cat. No. STEM00003). Monoclonal lines were isolated by limiting dilution and CLYBL integration was confirmed by PCR genotyping. Karyotype testing (Cell Line Genetics) was normal for the clonal line used for further experiments in this study, which we termed CRISPRa iPSCs.

Genome-wide survival-based and FACS-based screens
The genome-wide CRISPRi and CRISPRa libraries hCRISPRi-v2 and hCRISPRa-v2 (Horlbeck et al., 2016), consisting of 7 sublibraries each (H1-H7), were packaged into lentivirus as previously described (Tian et al., 2019). CRISPRi- and CRISPRa-iPSCs were infected by the sgRNA libraries at MOIs of 0.4-0.6 (as measured by the BFP fluorescence from the lentiviral vector) with approximately 1000x coverage per library element. Two days after infection, the cells were selected for lentiviral integration using puromycin (1 µg/mL) for 3 days as the cultures were expanded for the screens. After selection and expansion, a fraction of the cells (Day -3 iPSCs) were harvested and subjected to sample preparation for next-generation sequencing. Another fraction of Day -3 iPSCs, with a cell count corresponding to 1000x coverage per library element, were differentiated into neurons as described in the Human iPSCs Culture and Neuronal Differentiation subsection.

Neurons were cultured in either the +AO or -AO Neuronal Medium (see Human iPSCs Culture and Neuronal Differentiation subsection) for ten days. For the survival screens, Day 10 neurons were harvested and subjected to sample preparation for next-generation sequencing. For the FACS screens, Day 10 CRISPRi neurons cultured in the +AO medium and -AO medium were dissociated using Papain (Worthington; Code: PAP2; Cat. No.LK003178) and stained by CellRox Green (Invitrogen/Thermo Fisher Scientific; Cat. No. C10444) and Liperfluo (Dojindo Molecular Technologies, Inc.; Cat. No. L248-10) respectively (see Cell Staining by Fluorescent Probes) and sorted into high and low signal populations in each screen corresponding to the top 40% and the bottom 40% of the staining signal distribution, followed by sample preparation for next-generation sequencing.

For each screen sample, genomic DNA was isolated using a Macherey-Nagel Blood L kit (Macherey-Nagel; Cat. No. 740954.20). sgRNA-encoding regions were amplified and sequenced on an Illumina HiSeq- 4000 as previously described (Gilbert et al., 2014).

Focused secondary screens

The focused secondary screen library contained 2,190 sgRNAs targeting 730 genes that were hits in at least one of the ROS and lipid peroxidation screens with 3 sgRNAs per gene selected based on their phenotypes in the primary screens, and 100 non-targeting control sgRNAs. A pool of sgRNA-containing oligonucleotides were synthesized by Agilent Technologies and cloned into our optimized sgRNA expression vector as previously described (Gilbert et al., 2014). CRISPRi-iPSCs were transduced with the batch characterization library, puromycin selected, and differentiated into neurons as for the primary screens. Day10 neurons were stained with FeRhoNox-1 (Goryo Chemical; Cat. No. GC901) or Lysotracker-green (Cell Signaling Technology; Cat. No. 8783S) and sorted into high and low signal populations corresponding to the top 40% and bottom 40% of the staining signal distribution. Screen samples were processed and sequenced by next-generation sequencing as described above.

CROP-seq
For the CROP-seq experiments, we included 184 genes for CRISPRi and 100 genes for CRISPRa, which were hits in at least one of our genome-wide pooled screens and were also associated with human neurodegenerative diseases. We curated a list of genes associated with neurodegenerative diseases based on the literature and the DisGeNET database (https://www.disgenet.org/). The CROP-seq libraries included 2 sgRNAs per gene plus 6 non-targeting control sgRNAs, for a total of 374 sgRNAs for CRISPRi and 206 sgRNAs for CRISPRa. Top and bottom strands of sgRNA oligos were synthesized (Integrated DNA Technologies) and annealed in an arrayed format. The annealed sgRNAs were then pooled in equal amounts and ligated into our optimized CROP-seq vector (Tian et al., 2019).

The CROP-seq experiments were carried out similarly as previously described (Tian et al., 2019). Briefly, Day 0 CRISPRi and CRISPRa neurons were infected by the corresponding CROP-seq sgRNA library at a MOI of 0.1-0.2, followed by puromycin selection at 4 µg/ml for 3 days and recovery. On Day 10, neurons were dissociated with Papain and approximately 98,000 CRISPRi neurons and 50,000 CRISPRa neurons were loaded into 10X chips with about 25,000 input cells per lane. Sample preparations were performed using the ChromiumNext GEM Single Cell 3□ Reagent Kits v3.1 (10X Genomics; Cat. No.PN-1000121) according to the manufacturer's protocol. To facilitate sgRNA assignment, sgRNA-containing transcripts were additionally amplified by hemi-nested PCR reactions as described (Tian et al., 2019). The sgRNA-enrichment libraries were separately indexed and sequenced as spike-ins alongside the whole- transcriptome scRNA-seq libraries using a NovaSeq 6000 using the following configuration: Read 1: 28, i7 index: 8, i5 index: 0, Read 2: 91.

Cell Staining by Fluorescent Probes
All stains were performed according to manufacturing protocols. For Filipin staining, cells were washed with PBS three times and fixed with 3% paraformaldehyde for 1 hr at room temperature. Cells were then washed 3 times with PBS and incubated with 0.05 mg/ml Filipin III from *Streptomyces filipinensis* (Sigma; Cat. No. F4767-1MG) in PBS for 2 h at room temperature. Cells were washed with PBS 3 times before analysis. For staining using other live cell probes, cells were washed with PBS and incubated in DMEM containing appropriate concentrations of the probes at 37 °C as detailed below. Cells were washed with PBS before analysis. Concentrations and staining conditions for different probes were as follows: CellRox Green, 2.5 µM for 30 minutes; Lysotracker Green, 50 nM for 5 minutes; FeRhoNox-1, 5 µM for 60

minutes; Liperfluo, 5 µM for 30 minutes; C11-BODIPY, 2.5 µM for 30 minutes; FerroOrange (Dojindo Molecular Technologies, Inc.; Cat. No. F374-10), 1 µM for 30 minutes; Hoechst 33342 (Thermo Fisher Scientific; Cat. No. H3570), 1 µg/ml for 10 minutes; Propidium Iodide (PI)(Thermo Fisher Scientific; Cat. No. P1304MP), 1 µg/ml for 10 minutes.

Immunofluorescence
Cells were washed with PBS and fixed with 4% paraformaldehyde for 15 min at room temperature. After washing with PBS for 3 times, cells were permeabilized with 0.1% Triton X-100 for 10 min and blocked with 5% normal goat serum with 0.01% Triton X-100 in PBS for 1 hr at room temperature. Cells were then incubated with primary antibodies diluted in blocking buffer at 4 °C overnight. After that, cells were washed with PBS for 3 times and incubated with secondary antibodies diluted in blocking buffer for 1 hr at room temperature. Cells were then washed with PBS for 3 times and stained with 10 µg/ml Hoechst 33342 (Thermo Fisher Scientific; Cat. No. H3570) for 10 min. Cells were imaged using a confocal microscope (Leica SP8) or an IN Cell Analyzer 6000 (GE; Cat. No. 28-9938-51). Primary antibodies used for immunofluorescence in this study were as follows: rabbit anti-PSAP antibody (1:50 dilution; Proteintech; Cat. No. 10801-1-AP), mouse anti-LAMP2 antibody (1:100 dilution; abcam; Cat. No. ab25631), rabbit anti-GM1 antibody (1:20 dilution; abcam; Cat. No. ab23943), rabbit anti-LC3B antibody (1:200 dilution; Cell Signaling Technology; Cat. No. 2775S) and chicken anti-TUJ1 antibody (1:500; AVES; Cat. No. TUJ). Secondary antibodies used in this study were as follows: goat anti-rabbit IgG Alexa Fluor 555 (1:500 dilution; abcam; Cat. No. ab150078), goat anti-mouse IgG Alexa Fluor 488 (1:500 dilution; abcam; Cat. No. ab150113) and goat anti-chicken IgG Alexa Fluor 647 (1:500 dilution; abcam; Cat. No. ab150171).

Western blots
Cells were lysed in RIPA buffer and 20-30 µg of total proteins were loaded into NuPAGE 4%–12% Bis-Tris Gels (Invitrogen, Cat# NP0336BOX). Subsequently, the gels were transferred onto nitrocellulose membranes and the membranes were blocked by Odyssey Blocking Buffer (PBS) (LI-COR, Cat#927-50000), followed by overnight incubation with primary antibodies at 4°C. After incubation, the membranes were washed three times with TBST and then incubated with secondary antibodies at room temperature for 1 hr. The membranes were then washed 3 times with TBST and once with TBS and imaged on the Odyssey Fc Imaging system (LI-COR Cat# 2800). Digital images were processed and analyzed using ImageJ.

Primary antibodies used were mouse anti-β-Actin antibody (1:2000 dilution; Cell Signaling Technology; Cat. No. 3700), rabbit anti-PSAP antibody (1:1000 dilution; Proteintech; Cat. No. 10801-1-AP) and rabbit anti-LC3B antibody (1:1000 dilution; Cell Signaling Technology; Cat. No. 2775S). Secondary antibodies were IRDye 680RD goat anti-mouse IgG (1:20,000 dilution; LI-COR; Cat. No. 926-68070) and IRDye 800CW goat anti-rabbit IgG(1:20,000 dilution; LI-COR; Cat. No. 926-32211).

Generating the *PSAP* KO iPSC line
An sgRNA targeting *PSAP* exon 2 (sgRNA sequence: GGACTGAAAGAATGCACCA) was cloned into plasmid px330-mcherry (px330-mcherry was a gift from Jinsong Li (Addgene plasmid # 98750 ; http://n2t.net/addgene:98750 ; RRID:Addgene_98750)). The plasmid was transfected into WT Ngn2-iPSCs using Lipofectamine Stem (Invitrogen/Thermo Fisher

Scientific; Cat. No. STEM00003). Monoclonal lines were isolated in 96-well plates by limiting dilution. One clonal line was selected and frameshift indels were confirmed by Sanger sequencing. Protein level KO was confirmed by western blot and immunofluorescence (Fig. 5C,D). A normal karyotype was confirmed (Cell Line Genetics).

Bulk RNA sequencing

RNA was extracted from cells using the Quick-RNA Miniprep Kit (Zymo; Cat. No. R1054), and 3′-tag RNA-seq was performed by the DNA Technologies and Expression Analysis Core at the UC Davis Genome Center.

Electron Microscopy

Neurons grown on a poly-D-lysine coated 35-mm ibidi μ-Dish (ibidi; Cat. No. 81156) were fixed with 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer (EMS) for at least an hour at room temperature. Samples were further post-fixed with 1% osmium tetroxide and 1.6% potassium ferrocyanide, later dehydrated in graded series of ethanols, and embedded in epon araldite resin. Samples were then trimmed, 70nm sections were cut using Ultra cut E (Leica) and stained with 2% uranyl acetate and Reynold's lead citrate. Images were acquired on a FEI Tecnai 12 120KV TEM (FEI) and data was recorded using UltraScan 1000 Digital Micrograph 3 software (Gatan Inc.)

STORM super-resolution microscopy

Sample preparation and STORM imaging were performed as described previously (Hauser et al., 2018). Samples were fixed in 3% paraformaldehyde and 0.1% glutaraldehyde (Electron Microscopy Sciences) in PBS (Corning) for 30 minutes at room temperature, followed by reduction with 0.1% $NaBH_4$ in PBS for 10 minutes and three times of washing with PBS. Prior to immunostaining, samples were treated by a blocking buffer (BB) of 3% bovine serum albumin (Sigma) and 0.1% Triton-X100 (Sigma) in PBS for 1 hour at room temperature. Primary antibodies were diluted in BB and labeled at 4C overnight. Unbound antibodies were rinsed three times, 10 minutes each time with a washing buffer (WB) prepared by 10X dilution of BB in PBS. Secondary antibodies Dye-labeled secondary antibodies from Invitrogen or Jackson ImmunoResearch Laboratories were diluted in BB and labeled at room temperature for 1 hour, followed by three times of washing with WB.

Before STORM imaging, the sample was mounted in a standard STORM imaging buffer of 5% [w/v] glucose (Sigma), 0.1 M cysteamine (Sigma), 0.8 mg/mL glucose oxidase (Sigma), and 40 mg/mL catalase (Sigma) in 0.1 M Tris-HCl pH 7.5 (Corning). Two-color STORM was carried out on a custom setup modified from a Nikon Eclipse Ti-E inverted fluorescence microscope (Wojcik et al., 2015) by sequentially imaging the Alexa Fluor 647-labeled GM1 and the CF568-labeled LAMP2 using 647 nm and 560 nm excitations, respectively. A strong (~2 kW/cm^2) excitation laser (647 nm or 560 nm) was applied to photoswitch most of the AF647 or CF568 dye molecules into the dark state while also exciting the remaining dye molecules at a low density for single-molecule localization. A weak (0-1 W/cm^2) 405-nm laser was used simultaneously with the 647-nm or 560-nm laser to reactivate dye molecules in the dark state into the emitting state to acquire adequate sampling of the labeled molecules. Images were

collected at 110 frames per second with an electron multiplying charge-coupled device camera (Andor iXon Ultra 897) for ~50,000 frames per image.

Untargeted Lipidomics

The untargeted lipidomics experiment and primary analysis were performed by Cayman Chemical. Briefly, lipids were extracted using a methyl‑tert‑butyl ether (MTBE)‑based liquid‑liquid method. Cell pellets (approximately 100 μL in volume) were thawed on ice and transferred into 8-mL screw-cap tubes before adding 600 μL MeOH, the 600 μL MeOH containing 200 ng each of the internal standards TG(15:0/18:1‑d7/15:0), PC(15:0/18:1‑d7), PE(15:0/18:1‑d7), PG(15:0/18:1‑d7), and PI(15:0/18:1‑d7) (EquiSPLASH, Avanti Polar Lipids), and finally 4 mL MTBE. After vigorous vortexing, the samples were incubated at room temperature on a shaker for 1 h. For phase separation, 1 mL water was added, and samples were vortexed and centrifuged for 10 min at 1000 x g. The upper organic phase of each sample was carefully removed using a Pasteur pipette and transferred into a pre‑weighed empty glass tube. The remaining aqueous phase was re-extracted with 2 mL of clean MTBE/methanol/water 10:3:2.5 (v/v/v). The two upper organic phases were combined and dried under vacuum in a SpeedVac concentrator. The dried lipid extracts were weighed and resuspended in 100 μL isopropanol/acetonitrile 1:1 (v/v) for untargeted lipidomic analysis by LC‑MS/MS. Triplicates of samples for WT and *PSAP* KO neurons were analyzed.

Data analysis

Pooled CRISPR screens

Pooled CRISPR screens were analyzed using the MAGeCK-iNC pipeline as described (Tian et al., 2019). Briefly, raw sequencing reads from next-generation sequencing were cropped and aligned to the reference using Bowtie (Langmead et al., 2009) to determine sgRNA counts in each sample. Counts files for samples subject to comparison were entered into the MAGeCK-iNC software (kampmannlab.ucsf.edu/mageck-inc). Phenotype scores and significance P values were determined for all target genes in the library, as well as for 'negative-control-quasi-genes' that were generated by random sampling from non-targeting control sgRNAs. A gene score was calculated for each gene, which was defined as the product of the phenotype score and $-\log_{10}$(P value). To determine hit genes, a gene score cutoff value was chosen to make sure the false-discovery rate (FDR) is less than 0.05.

CROP-seq

CROP-seq analysis was performed similarly to previously described (Tian et al., 2019). Cellranger (version 3.1.0,10X Genomics) with default parameters was used to align reads and generate digital expression matrices from single-cell sequencing data.

Approximately 58,000 CRISPRi neurons and 38,000 CRISPRa neurons were detected. The mean reads per cell was around 48,000 for CRISPRi and 36,000 for CRISPRa. The median number of genes detected per cell was around 4,341 for iPSCs and 3,100 for CRISPRa.

sgRNA-enrichment libraries were analyzed using methods previously described (Hill et al., 2018) to obtain sgRNA UMI counts for each cell barcode. For a given cell, sgRNA(s) whose UMI counts were greater than 4 standard deviations of the mean UMI counts of all sgRNAs were

assigned to that cell as its identity. Single sgRNAs could be assigned to about 35,000 CRISPRi cells and about 21,000 CRISPRa cells, which were retained for further analysis.

The Scanpy package (version 1.4.6, (Wolf et al., 2018) implemented in Python was used for downstream analysis of the digital expression matrices with mapped sgRNA identities. To ensure data quality, a stringent criterion was applied to filter cells based on the number of genes detected (> 2000 for CRISPRi and > 1500 for CRISPRa) and percentage of mitochondrial transcript counts ($< 0.15\%$). Genes that had less than 0.5 UMIs on average in all perturbation groups were filtered out.

Within the population of cells expressing sgRNAs for a specific target gene, the levels of target knockdown or overexpression were heterogeneous in some cases. This could be due to different efficiencies of the two sgRNAs targeting that gene (see Figure S4A for an example), misassignment of sgRNA identities for some cells or stochastic silencing of the CRISPR machinery in some cells. To select cells in which functional perturbations happened, we leveraged an unsupervised outlier detection method based on the local outlier factor (LOF) using the LocalOutlierFactor function in the Python package scikit-learn (version 0.23.0). A similar strategy was described previously (Adamson et al., 2016). Specifically, for every target gene, we selected two populations of cells, including one population of cells that were mapped with sgRNAs targeting that gene (i.e. perturbed group) and another population of cells that were mapped with non-targeting control sgRNAs (i.e. control group). We then identified differentially expressed genes (DEGs) between the two populations by t-tests at $p < 0.05$. A gene-expression matrix containing only expression of DEGs in cells from the two populations was generated and principal component analysis (PCA) was performed to reduce the matrix to 4 dimensions. Cells in the control group in the 4-dimensional space were used as the training set to fit a LocalOutlierFactor model. Then, the model was used to determine whether a cell in the perturbed group was an 'outlier' based on the extent it deviated from the controls. The 'outliers' were considered as cells in which functional perturbations occurred and were retained for downstream analysis. This classification method was particularly useful for CRISPRi when the basal expression level of the target gene was too low to be detected by single-cell RNA-seq thus did not allow selecting cells based on knockdown level of target gene (see Figure S4B for an example).

DEGs for each perturbation group compared to the control group were then determined by t-tests using the diffxpy package in Python.

A mean gene-expression matrix was generated for different perturbation groups (including the control group) by averaging the normalized expression profile of all cells within that group. This matrix was used for weighted correlation network analysis (WGCNA) using the WGCNA package (version 1.69, (Langfelder and Horvath, 2008)) implemented in R. The blockwiseModules function was used to detect gene modules that were co-regulated in different perturbation groups and to determine eigengene expression of each module in each perturbation

group. Relative eigengene expression values were calculated by subtracting the eigengene expression value of each module in the control group from that in each perturbation group.

RNA-seq

Raw sequencing reads from 3′-tag RNA-seq were mapped to the human reference transcriptome (GRCh38, Ensembl Release 97) using Salmon (v.0.14.139, (Patro et al., 2017)) with the '–noLengthCorrection' option to obtain transcript abundance counts. Gene-level count estimates were obtained using tximport (v.1.8.040, (Soneson et al., 2015)) with default settings. Subsequently, genes with more than 10 counts were retained for differential gene-expression analysis, and adjusted P values (P_{adj}) were calculated using DESeq2 (v.1.20.041, (Pang et al., 1989)).

Pathway enrichment analysis

Gene Ontology (GO) term and Wikipathways enrichment analysis was performed using WebGestalt (WEB-based Gene SeT AnaLysis Toolkit) using the over-representation analysis (ORA) method (Liao et al., 2019).

Lipidomics

Lipostar software (Molecular Discovery) was used for feature detection, noise and artifact reduction, alignment, normalization, and lipid identification. For each lipid, the $\log_2$-fold change and a significance P value by t-test were determined by comparing the abundances of that lipid in WT and *PSAP* KO neurons (samples in triplicates). The Benjamini-Hochberg (BH) method was used to correct for multiple hypothesis testing.

Data commons development

CRISPRbrain (https://crisprbrain.org) was developed as an open-access and cloud-based platform striving to make data and code easily accessible to the scientific community. To enable scientific transparency and replication, each dataset in CRISPRbrain is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0). To handle a large volume of storage and computation demand, CRISPRbrain is deployed on the cloud with elastic resource allocation. We have prioritized a low-latency interactive user experience by pre-computing and caching complex queries. CRISPRbrain is implemented in a manner that is "future-proof", supporting scalability and addition of more complex features.

SUPPLEMENTAL TABLE LEGEND

Untargeted lipidomics data for WT and PSAP KO neurons. P values were calculated using Student's t-test and were corrected for multiple testing using the Benjamini-Hochberg procedure.