Now you see me, now you don't: Modulation of neural responses to targets

moving in three-dimensional optic flow

Sarah Nicholas[1] and Karin Nordström[1,2]*

[1]Centre for Neuroscience, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia

[2]Department of Neuroscience, Uppsala University, Box 593, 751 24 Uppsala, Sweden

*email: karin.nordstrom@flinders.edu.au

ORCIDs

Sarah Nicholas: 0000-0002-5555-9421

Karin Nordström: 0000-0002-6020-6348

12
13

Abstract

For the human observer, it is difficult to follow the motion of small objects, especially when these are displayed against background motion. However, insects efficiently do this, as evidenced by their ability to capture prey, pursue conspecifics or defend territories, often in foliage-rich areas. This ability has been attributed to insect optic lobe neurons sharply tuned to target motion, as these respond robustly to targets even against moving backgrounds. However, their presumed post-synaptic counterparts, the target selective descending neurons (TSDNs), do not, suggesting inhibition by widefield motion. Importantly, though, previous experiments used panoramic background images, which not only lacked 3D cues, but also high-contrast features, experienced in natural flight. To mitigate this, we here use widefield motion consisting of many targets moving coherently to simulate the type of 3D optic flow that would be generated by the animal's own motion through the world. We show that TSDNs are inhibited by syn-directional optic flow, consistent with previous findings. However, and more surprisingly, we show that TSDNs are facilitated by optic flow in the opposite direction to target motion. Our results thereby emphasize that the responses of descending neurons cannot simply be inferred from their presumed pre-synaptic counterparts.

Introduction

The survival of many animals often depends on their ability to visually detect small moving objects or targets, as these could represent predators, prey or territorial intruders. Efficient target detection is a computationally challenging task which becomes even more difficult when done against visual clutter. Despite this, many insects successfully detect small moving targets and engage in highly acrobatic pursuits, often in visually complex environments. For example, male *Eristalis tenax* hoverflies establish territories in foliage rich areas within which they act as sentry, on alert for intruders or female conspecifics, and ready to engage in pursuit as soon as a target is detected[1].

Initial target detection can be facilitated by behavioral mechanisms that make the surrounding background stationary, thereby making the target stand out as the only thing that moves. For example, many insects and vertebrates alike visualize prey or conspecifics against the bright sky[2], or from a stationary stance, such as perching[3-5] or hovering[6,7]. However, as soon as the pursuer moves its own head or body, this creates self-generated optic flow[8]. To achieve reliable and continuous identification of a target, the independent motion of the target needs to be discriminated from such self-generated optic flow. That insects do this successfully is a remarkable feat considering their small brains and low-resolution compound eyes[9]. This suggests a high level of optimization, which makes insects excellent for understanding the underlying neural mechanisms.

Predatory dragonflies and territorial hoverflies, which both pursue targets, have sharply size-tuned small target motion detector (STMD) neurons in their optic lobes[10,11]. Impressively, some STMD neurons respond robustly to targets in clutter, even without

relative velocity differences[11,12], suggesting that they could support behavioral pursuit

against self-generated optic flow. However, the target selective descending neurons

(TSDNs), which are thought to be post-synaptic to STMDs[13], do not respond to targets

moving across syn-directional background motion[14]. This is peculiar as descending neurons

are more likely to directly drive behavioral output[15,16].

However, the STMDs that respond most robustly to targets against background motion, in

both hoverflies and dragonflies[11,12], do not project to pre-motor areas, but contralaterally

through the protocerebrum, with outputs in the heterolateral lobula[11,17]. This suggests a

role in modulating the responses of other optic lobe neurons, rather than providing direct

input to descending neurons[13]. In contrast, the STMDs that are more likely to provide direct

input to the TSDNs, based on their small receptive fields, and outputs in pre-motor areas of

the lateral mid- brain[18], where at least in the dragonfly, TSDNs have input dendrites[19,20], are

inhibited by background motion. Indeed, the response to a target moving against syn-

directional background motion is smaller than the response to a target traversing a white

background[18].

The TSDN response to targets against background motion is inversely correlated with the

response of optic flow sensitive descending neurons to the same background motion[14]. This

suggests that the target detection pathway is inhibited by input from the widefield system,

possibly post-synaptic to the STMDs. Importantly, however, the backgrounds used in

previous STMD[11,12,18] and TSDN[14] experiments consisted of panoramic images that were

displaced across the visual monitor. Most often, these were moved horizontally, with a

black, high-contrast target also moving horizontally. Even if the backgrounds had spatial

image statistics resembling nature[21] they lacked the high-contrast, hard edges of the target.

Where high-contrast features are present in naturalistic backgrounds, these will often

generate STMD responses[12]. Furthermore, the backgrounds lacked the depth cues inherent

in self-generated optic flow. This is crucial as self-generated optic flow not only reflects the

speed and direction of the animal's own movements, but also the distance to objects within

the environment[8]. During translational movements, for example, nearby objects move

faster across the retina than distant objects do. By contrast, in rotational optic flow all

surrounding objects move across the retina at a constant velocity, irrespective of their

distance from the viewer. Many insects use this information to estimate object size and

distance. For example, walking *Drosophila* use this to estimate relative distance prior to

approaching the nearest object[22], bees discern the height of objects prior to landing[23] and

locusts and praying mantises assess distance before attacking prey[24-26]. Hoverflies can move

in nearly every direction, including backwards[27,28], and in doing so generate different types

of translational and rotational optic flow. However, a flat, artificial image sliding across a

visual monitor[11,12,14,18] provides none of these relative motion cues, and none or very few

high-contrast cues, which would be experienced during such natural behaviors.

To redress this deficiency, we quantified the responses of *Eristalis* TSDNs to targets moving

against 6 types of optic flow, three translations and three rotations. By calibrating for the

hoverfly's position relative to the visual monitor, the simulated optic flow[29] mimics the self-

generated optic flow seen during motion through a 3D world. For this purpose, we used an

optic flow stimulus consisting of many targets moving coherently. The individual features

that make up the optic flow thus contain the same high-contrast edges as the target itself,

against which the target's motion can only be determined by its independent trajectory. We

found that this optic flow alone did not generate any TSDN response, despite it consisting of

similar "targets", even when moving in the neuron's preferred direction. Consistent with

previous results[14] we found that optic flow in the same direction as the target inhibits the

TSDN response. Importantly, this inhibition occurs even if the optic flow only precedes the

target motion and does not appear concurrently. By quantifying responses to different

target and background combinations, we show that the inhibition is likely inherited from

pre-synaptic STMDs, instead of occurring post-synaptic to the STMDs as we previously

suggested[14]. More surprisingly, though, we show for the first time that optic flow in the

opposite direction to target motion increases the TSDN response. Together, our results

emphasize that the responses of descending neurons cannot simply be inferred from their

presumed pre-synaptic counterparts in the optic lobes, and if we want to understand the

neural correlate of behavior we need to study the neurons closer to motor output.

Results

TSDN responses to small targets are slightly reduced by stationary background clutter

We recorded extracellularly from target selective descending neurons (TSDNs) in male

Eristalis hoverflies. TSDNs respond selectively to the motion of small targets[14,29], as shown in

the example where a round, black target traverses a white background (Extended Data

Movie 1, 2, Fig. 1a, b, blue line indicates target motion). As shown previously[14,29], TSDNs

typically have no spontaneous activity (Fig. 1b).

We used a 3D optic flow stimulus built of many targets that move coherently[29] to simulate

the types of retinal flow fields experienced during self-generated optic flow. Against this

optic flow, the target can only be identified by its independent trajectory. When the optic

flow and the target are both stationary, there are no spatial differences which would permit

the identification of the target (Fig. 1a, right). The stationary optic flow, which appeared

0.5 s prior to target motion, sometimes evokes 1-2 spikes (beginning of red line, Fig. 1b,

right, Extended Data Fig. 1, "Stationary"). Nevertheless, when the target moves across this

stationary optic flow (Extended Data Movie 3, 4, Fig. 1a, right), the TSDN response remains

robust compared with targets moving across the white background (Fig. 1b, c, compare left

and right data).

The responses of an individual TSDN are stable over long recording times, as illustrated in

the spike raster plot from 18 trials spanning over 22 minutes' recording in the same neuron

(Fig. 1d). The resulting spike histograms highlight how similar the responses are whether the

target traverses a white background or the stationary optic flow (Fig. 1e). For quantification

across neurons, we calculated the mean spike frequency for the duration of target motion,

after excluding the first and last 40 ms (analysis window indicated by dotted box, Fig. 1c-e).

Using this, we find that there is a small response reduction to targets moving across

stationary optic flow (grey, Fig. 1f) compared with a white background (black, Fig. 1f).

We find that the response across neurons is highly variable (coefficient of variation 49% and

51%, respectively, Fig. 1f). This could reflect the internal state of the individual animals or

correspond to a real difference between TSDNs. In dragonfly TSDNs, spike rates vary not

only across neuron types, but also within neuron types[19,30]. To take this variability into

account, subsequent data from each neuron are normalized to its own mean response to a

target moving across a white background (black, Fig. 1f).

The TSDN response to target motion is strongly affected by sideslip and yaw optic flow

We next investigated how TSDNs respond to targets moving across either sideslip or yaw

optic flow. Importantly, both sideslip and yaw result in horizontal motion in the frontal

visual field. The control data from one TSDN show the response to a target moving over

stationary optic flow (grey, Fig. 2a). For each neuron, we normalized this to its mean

response to a target moving over a white background (grey, Fig. 2b). When the target moves

over sideslip optic flow in the same direction as the target, the TSDN response is strongly

inhibited, as seen in the example neuron (Fig. 2a) and across neurons (Fig. 2b, "Sideslip

+50"). We found that yaw optic flow moving in the same direction as the target, also

strongly inhibits the TSDN response, as seen in the example neuron (Fig. 2a) as well as

across neurons (Fig. 2b, "Yaw +50"). This complete inhibition is striking considering that the

target moved along an independent trajectory from the remaining optic flow (Extended

Data Movie 5, 6), but consistent with previous work using sliding background images[14].

When the sideslip optic flow moves in the opposite direction to the target the TSDN

response is strongly enhanced (Extended Data Movie 7, 8, Fig. 2a, "Sideslip -50"). This

response increase to target motion against sideslip optic flow in the opposite direction is

substantial (mean 71.1%) and consistent across neurons (Fig. 2b). Similarly, yaw optic flow

in the opposite direction to the target also facilitates the TSDN response by 84.9%, as seen

in the example data (Fig. 2a) and across neurons (Fig. 2b, "Yaw -50"). Such response

facilitation was not seen when displaying targets moving over panoramic background

images, in either TSDNs[14] or target tuned neurons in the fly optic lobes[11,18,31].

The background optic flow itself, contained ca. 1200 "targets" moving coherently in front of

the fly. When the sideslip moves in the TSDN's preferred direction (Fig. 2, "Sideslip +50") we

expect tens of these "background targets" to move through the receptive field of the TSDN

(Extended Data Movie 5, 6). Despite this, across 222 repetitions in 12 TSDNs, preferred

direction sideslip optic flow generated a response in only 18% of the repetitions (Extended

Data Fig. 1a, "Sideslip +50"). In the instances where preferred direction sideslip optic flow

generated a response, the amplitude of the response was less than one-fifth of the response

to a target traversing a white background (Extended Data Fig. 1b, "Sideslip +50"). In

addition, when the background sideslips in the anti-preferred direction, we see small

breakthrough responses in 9% of repetitions (213 repetitions, "Sideslip -50", Extended Data

Fig. 1). Taken together, this suggests that the coherent motion of the background is treated

differently from the independent motion of a target.

The response to targets moving against other types of 3D optic flow is inhibited

We next investigated the effect different types of 3D optic flow has on TSDN responses to

target motion. The example data from one TSDN show that the response to targets is more

strongly inhibited by lift optic flow moving downwards in the visual field (Fig. 3a, "Lift +50")

compared to lift moving upwards (Fig. 3a, "Lift -50"). Across neurons, however, both

directions of lift optic flow suppress the TSDN response (by 44.0%, "Lift +50"; and 34.2%,

"Lift -50", Fig. 3b). When the target is instead displayed again pitch optic flow, which

provides similar vertical, local motion in the frontal visual field, the response is suppressed

by similar levels (57.1%, "Pitch +50", and 54.9%, "Pitch -50", Fig. 3b). In addition, the

response to target motion is inhibited to roughly similar levels when displayed against

thrust or roll optic flow in either direction (Fig. 3).

In summary, when displaying a target moving horizontally in the TSDN's preferred direction,

optic flow in the opposite direction facilitates the response, syn-directional optic flow

completely inhibits the response, and optic flow in any other direction inhibits the response

by ca. 50% (Fig. 2, 3). The relative lack of responses to the optic flow itself can be seen in

Extended Data Fig. 1.

TSDN responses to target motion is inhibited by preceding optic flow

In dragonflies, STMD responses to target motion can be primed by preceding target motion,

thus facilitating responses to longer target trajectories[32,33]. Above, we showed that TSDN

responses to target motion are affected by sideslip optic flow (Fig. 2). To determine if this

effect can be primed in a similar manner, we moved preceding optic flow for 1 s (green, Fig.

4a), but stopped it to be stationary (red, Fig. 4a) concurrent with target motion (blue, Fig.

4a).

In the control experiment the optic flow was stationary preceding and concurrent with

target motion (grey, Fig. 4b). We found that when sideslip optic flow preceded target

motion in the same direction, this was enough to significantly inhibit the TSDN response,

even if the optic flow concurrent with target motion was stationary (black, Fig. 4b). This

suggests that the inhibition of TSDN responses can be primed by preceding optic flow. We

did the inverse experiment where the sideslip optic flow moves concurrently with target

motion and we varied the preceding optic flow. We found that whether the optic flow was

in motion or stationary prior to the target had no significant effect on the TSDN inhibition

(Fig. 4c).

We next investigated whether facilitation by sideslip optic flow in the opposite direction to

target motion can be primed. We found that preceding sideslip optic flow in the opposite

direction to target motion did not increase the TSDN response (black, Fig. 4d) compared to

control (grey, Fig. 4d). When the sideslip optic flow moves in the opposite direction

concurrent with target motion, the TSDN response is facilitated (Fig. 4e). However, whether

the sideslip was in motion or was stationary prior to the target had no significant effect on

this TSDN facilitation (Fig. 4e).

TSDN inhibition by syn-directional optic flow is likely inherited from pre-synaptic STMDs

The data above show that TSDN responses to horizontal target motion are almost

completely inhibited if the optic flow moves in the same direction as the target (Fig. 2), and

by 34% if displayed against lift, which moves perpendicular to the target (grey crosses, Fig.

5c, d, re-plotted from Fig. 2b, 3b). This suggests that the target detection pathway receives

inhibitory input from widefield motion. We assume that TSDNs receive input from

STMDs[11,18] or other target tuned neurons[31,34-36], potentially directly (dashed lines,

pictograms, Fig. 5c, d). Inhibition from widefield motion could take place post-synaptic to

the STMDs, but pre-synaptic to TSDNs (pictogram, Fig. 5c). If so, we would not expect the

normalized TSDN responses to change if the target changes direction, but the optic flow

remains the same (black lines, Fig. 5d). In this scenario, the TSDN inhibition would be

correlated with the optic flow, irrespective of target direction. Alternatively, inhibition from

the widefield motion could take place prior to the pre-synaptic STMDs, so that the STMDs

themselves are unable to respond to targets against syn-directional background motion

(pictogram, Fig. 5d). This could e.g. be caused by direct inhibition from widefield motion, or

from the underlying model generating target selectivity itself. If this is true, the TSDN

response would be most strongly inhibited when target and optic flow move in the same

direction, whereas targets moving perpendicular to the optic flow would be more

detectable (black lines, Fig. 5d).

To investigate these two hypotheses we recorded from TSDNs that respond robustly not

only to horizontal target motion (grey, Fig. 5a, b), but also to vertical target motion (black,

Fig. 5a, b). As before, we found that when the target moves horizontally against sideslip

optic flow in the same direction as the target, the response is strongly inhibited (left, Fig.

5e). When the target instead moves vertically but the optic flow remains horizontal, so that

the target now moves perpendicular to the optic flow, the TSDN response is significantly

larger (right, Fig. 5e). In the inverse experiment, when the target moves against lift optic

flow, the TSDN response is more strongly inhibited when the target moves in the same

direction (right, Fig. 5f), compared with when the target moves horizontally (left, Fig. 5f).

These results therefore support our second hypothesis (Fig. 5d), that the inhibition by

widefield motion is inherited from the pre-synaptic STMDs, or other target tuned neurons.

TSDN response facilitation by background motion is linked to target direction

Above, we showed that inhibitory input from the widefield motion system is likely inherited

from the pre-synaptic STMDs (Fig. 5). To determine if facilitation (Fig. 2, 3) could also be

caused by input from the widefield system, we tested the analogous two hypotheses in the

same neurons as Figure 5. If facilitation takes place post-synaptic to the STMDs (pictogram,

Fig. 6a), we would expect the TSDN facilitation to be correlated with the optic flow,

irrespective of target direction (black lines, Fig. 6a). Alternatively, if facilitation takes place

prior to the pre-synaptic STMDs (pictogram, Fig. 6b), the TSDN response would be stronger

when target and background move in the opposite direction compared to targets moving

perpendicular to the optic flow (black lines, Fig. 5d).

We found that when the target moves horizontally with the background side-slipping in the

opposite direction to the target, the response is significantly larger compared to the

response to a vertical target on the same optic flow (Fig. 6c). In the inverse experiment,

when we display targets against lift optic flow, there was no difference depending on the

direction of the target (Fig. 6d). There is thus no significant facilitation when the target

moves vertically against lift optic flow in the opposite direction (right, Fig. 6d). Taken

together, our data (Fig 6c, d) do not support either hypothesis (Fig. 6a, b) making it unclear

where the facilitation by optic flow comes from (Fig. 2, 3, 6).

Discussion

We generated optic flow using the coherent motion of thousands of "targets" within a

simulated 3D space surrounding the hoverfly, and projecting the can 1200 targets anterior

of the fly onto a 2D visual monitor[29]. We showed that this 3D optic flow stimulus modulates

the TSDN response to target motion. In particular, we found that optic flow in the opposite

direction to target motion strongly enhances the TSDN response, that optic flow in the same

direction virtually abolishes the TSDN response, and that other types of optic flow decrease

the TSDN response to a lesser degree (Fig. 2, 3). We found that preceding syn-directional

optic flow was enough to inhibit the TSDN response, even if the optic flow was stationary

during target motion (Fig. 4). Using different directions of target motion and optic flow, we

showed that the inhibition by is likely inherited from the STMDs (Fig. 5), but that the source

of the facilitation remains unclear (Fig. 6).

TSDN input

The pre-synaptic input of TSDNs has not been verified, but is likely to come from target-

tuned neurons in the optic lobes[13]. Our experiments changing the direction of target motion

and optic flow (Fig. 5) suggests that the lack of TSDN responses to targets against syn-

directional optic flow is inherited from the pre-synaptic neurons. Indeed, early target

motion models relied on inhibitory feedback from the widefield system[37]. This is

conceptually an appealing model, as by simply subtracting self-generated background

motion, the independent motion of a target will stand out[38-40]. This model explains

physiological responses of blowfly lobula plate Figure Detecting (FD) cells, where the

response to small object motion is inhibited by background motion in the same and in the

opposite direction[39]. However, as FD cells are not as sharply tuned to small targets as TSDNs

are[37], and FD cells often respond to features within the background[41], as opposed to TSDNs

(Extended Data Fig. 1) they are unlikely to provide the TSDN input.

As some STMDs respond robustly to targets moving in visual clutter, even without relative

motion cues, they cannot rely on the FD model, which prompted the development of an

elementary STMD model[42]. The elementary STMD model uses the unique spatiotemporal

profile created by the motion of a dark target where its leading edge creates a dark contrast

change (OFF), quickly followed by the bright contrast change (ON) of the target's trailing

edge. By separating and half-wave rectifying the ON and OFF contrast signals from the same

point in space, and adding lateral inhibition from neighboring channels, a selective response

to small, dark, moving targets is generated[42]. The elementary STMD model provides robust

predictions of physiological responses of dragonfly STMDs[12,43]. However, as we see strong

inhibition of the TSDN target response when targets move over syn-directional optic flow

(Fig. 2, 3), and this inhibition takes place pre-synaptic to the TSDNs (Fig. 5), this argues

against TSDN input from an elementary STMD type model.

Importantly, as well, the elementary STMD model predicts[42] that background features with

the correct spatio-temporal profile will generate a neural response. Indeed, this prediction

is confirmed in dragonfly STMD recordings[12]. In our TSDN experiments the background

consisted of hundreds of targets moving coherently (Fig. 1a)[29]. Therefore, we expect tens of

these to move through the TSDN receptive field at any one time. These targets have varying

size, from the smallest we can render on our screen (1 pixel, below the resolution of the

hoverfly eye[44]) to 37° diameter, larger than what we expect TSDNs to respond to[14].

Nevertheless, when the optic flow moves in the neuron's preferred direction we would

expect some of these targets to have the right size to trigger a TSDN response, if the STMD

model underlies TSDN tuning. However, the lack of strong, and even consistent responses to

"targets" in preferred direction optic flow ("Sideslip +50", "Yaw +50", Extended Data Fig. 1),

also argue against an elementary STMD model type input to the TSDNs.

STMDs and FD cells are not the only object tuned neurons in the fly optic lobes. In

Drosophila, lobula columnar (LC) neurons have recently received more attention as possible

substrates of figure-ground discrimination behaviors[45]. Whereas some of these respond

better to bars than to small objects, LC11 and LC26 appear to respond selectively to small

targets[31,46]. Indeed, as LC11 neurons are clearly size-tuned, they would physiologically

classify as STMDs[47], and might be the *Drosophila* correlates of the columnar small-field

STMDs described in *Eristalis*[18]. LC11 neurons are especially interesting as they are clearly

involved in behavioral responses to small objects[36]. A model based on LC11 physiology

generates small object selectivity using a combination of center-surround antagonism and

rapid adaptation[36]. Indeed, LC11 neurons are strongly inhibited by background motion, and

this inhibition is believed to take place upstream of LC11[35], i.e. consistent with our

conceptual model (Fig. 5b). However, arguing against LC11 providing input to TSDNs is the

finding that they are equally inhibited by background motion in the same as well as in the

opposite direction to object motion[34], inconsistent with TSDNs (Fig. 2). In addition, LC11

neurons are not directional[35,46], so if they provide the input to the TSDNs, directionality has

to be added at some stage.

STMDs are a very heterogeneous group[11]. Some STMDs' responses to target motion are

slightly inhibited by syn-directional background motion[18], and other STMDs even respond

weakly to features in the background[11], but a complete lack of response to targets in clutter

are unpublished. More work on the pre-synaptic input to the TSDNs is clearly needed,

linking any variability in ability to see targets in clutter with neuro-anatomy, to determine

which neurons provide direct input to the TSDNs that we recorded from here.

Behavioral relevance

The facilitation of TSDN responses to target motion displayed on optic flow in the opposite

direction (Fig. 2b) is a novel observation, and difficult to reconcile with the existing models

of small target selectivity discussed above[36,39,42]. Indeed, whereas the inhibition is clearly

inherited from the pre-synaptic STMDs (Fig. 5), it is unclear where the facilitation is

generated (Fig. 6). Interestingly, we only saw facilitation to horizontal target motion (Fig. 2,

6), and not to vertical target motion (Fig. 6d), even if the neurons clearly respond to these

targets (Fig. 5a, b). Furthermore, whereas inhibition could be primed by preceding motion

(Fig. 4b), facilitation could not (Fig. 5c, e). All of these experiments (Fig. 4-6) point to

facilitation being generated somewhere else than inhibition.

Our own previous work, using panoramic background images, showed that TSDN responses

to small targets are significantly reduced when the target and the background move in

opposite direction[14]. This suggests that there is something fundamentally different about

the image background used previously[14] and the 3D optic flow used here. Indeed, the 3D

optic flow used here consisted of "targets" moving coherently (Extended Data Movie 5-8),

so that the target itself is impossible to identify in a stationary screen shot (Fig. 1a).

Furthermore, 3D depth cues are important for many insects[22,23,25], but not available when

viewing flat images. These differences could underlie the facilitation of TSDN responses.

Nevertheless, the facilitation could make behavioral sense. Prior to initiating target pursuit,

male *Eristalis* hoverflies predict the flight course required to successfully intercept the

target. This prediction is predominantly based on the angular velocity of the target[48]. To

successfully execute an interception style flight course, the hoverfly turns in the direction

that the target is moving[48]. In doing so, the hoverfly creates self-generated optic flow in the

opposite direction to the target's motion. If the TSDNs that we have recorded from here are

involved in such planning of pursuit, they would be facilitated when the target moves in the

opposite direction to the self-generated optic flow (Fig. 2, 3), which might be beneficial.

Once an interception style pursuit has started, if it is perfect the target's position remains

fixed on the pursuer's eye. If the retinal position of the target slips, male *Syritta* hoverflies

correct this by turning towards the target at an angular velocity proportional to the target's

angular position on the eye[28]. Such turns towards the target could generate optic flow in

different directions depending on the target's angular position. If this generates optic flow

in the same direction as the target is moving, the TSDNs would be inhibited (Fig. 2), likely

making them less suitable during flight. However, reconstructing retinal flow fields as

experienced during actual pursuits[49] will help resolve how often the target and the optic

flow actually move in the same direction. Furthermore, replaying such reconstructed

pursuits to TSDNs may determine whether they would respond during actual pursuits.

Importantly, as opposed to the case of a pursuing hoverfly, we recorded from immobilized

animals, with no neuronal feedback from the flight motor circuits[50,51]. It is possible that the

TSDN responses that we recorded are affected by the animals being immobilized. Indeed, in

two descending neurons types that control landing behaviors in *Drosophila,* visual responses

are suppressed in non-flight states[52]. Octopamine, which has been shown to mimic flight

behavior, restores responses in one of these neurons, while the other may receive direct

feedback from flight motor circuits[52]. This prevents inappropriate landing behaviors from

occurring during non-flight periods. Similarly, in *Drosophila*, the responses of bar tuned

lobula columnar LC12 and LC15 neurons are suppressed by simultaneous background

motion, with the effect reversed by chlordimeform, an octopamine agonist[31]. However, the

inability of LC11 neurons to respond to targets against backgrounds is not restored with

chlordimeform[31]. While technically difficult to record from TSDNs during actual flight, it

should be possible to mimic flight be treating the animal with chlordimeform or

octopamine. Another possibility is that TSDNs are not involved in active target pursuit and

that another class of neurons is recruited during flight. Indeed, there are other descending

neurons in the insect ventral nerve chord, including those that also respond to looming

stimuli, that respond to small targets[19,29].

Methods

Electrophysiology

Eristalis tenax hoverflies were reared and maintained as previously described[53]. Prior to electrophysiology, the male hoverfly was immobilized ventral side up using a beeswax and resin mixture. A small hole was cut at the anterior end of the thorax to expose the cervical connective, which was then raised slightly and supported using a small wire hook. A sharp polyimide-insulated tungsten microelectrode (2 MOhm, Microprobes, Gaithersburg, USA) was inserted into the cervical connective. The animal was grounded via a silver wire inserted into the ventral part of the hole. This also served as a recording reference.

Extracellular signals were amplified at 100x gain and filtered through a 10- to 3000-Hz bandwidth filter on a DAM50 differential amplifier (World Precision Instruments, Sarasota, USA), with 50 Hz noise removed with a HumBug (Quest Scientific, North Vancouver, Canada). The data were digitized via Powerlab 4/30 (ADInstruments, Sydney, Australia) and acquired at 40 kHz with LabChart 7 Pro software (ADInstruments). Single units were discriminated by amplitude and half-width using Spike Histogram software (ADInstruments).

Visual stimuli

Eristalis males were placed ventral side up, centered and perpendicular to an Asus LCD screen (Asus, Taipai, Taiwan) at a distance of 6.5 cm. The screen had a refresh rate of 165 Hz, a linearized contrast with a mean illuminance of 200 Lux and a spatial resolution of 2560 x 1440 pixels, giving a projected screen size of 155° x 138°. Visual stimuli were

displayed using custom written software based on the Psychophysics toolbox[54,55] in Matlab

(Mathworks, Natick, USA).

TSDNs were identified as described previously[14,29]. In short, TSDNs have small receptive

fields located in the dorso-frontal visual field, and sharp size tuning with a peak response to

targets subtending 3°- 6°, with no response to larger bars, to looming or to widefield

stimuli[29].

All targets were black and round with a diameter of 15 pixels, moving at a velocity of

900 pixels/s for 0.48 s. When converted to angular values and taking the small frontal

receptive fields of TSDNs into account, this corresponds to a diameter of 3° and a velocity of

130°/s[14]. Unless otherwise stated, each target trajectory travelled in each neuron's

preferred direction and across the center of its receptive field. Between repetitions, we

varied the target elevation slightly, to minimize habituation[14]. A minimum of 4 s was left

between stimulus presentations. Stimulus order was randomized.

3D optic flow was generated as previously described[29]. Briefly, the optic flow consisted of a

simulated 3D space, a cube with 4 m sides, filled with 2 cm diameter spheres at a density of

100 per m^3, with the hoverfly placed in the center. The coherent motion of these ca 6400

spheres around the hoverfly was used to simulate self-generated optic flow. The ca 1200

spheres anterior to the hoverfly were projected onto the screen, with the size of the

resultant circles indicating the distance from the hoverfly. Circles closer than 6 cm were not

displayed. Six types of optic flow were simulated: three translations at 50 cm/s (sideslip, lift

and thrust) and three rotations at 50°/s (yaw, pitch and roll). Unless otherwise stated optic

flow was displayed for 0.48 s prior to the target. Both target motion and optic flow ended

simultaneously.

Data analysis and statistics

We recorded from 24 TSDNs in 24 male hoverflies. We kept data from all TSDNs that

showed a robust response to a target moving over a white background (Extended Data

Movie 1, 2). We repeated this control throughout the recording, and only kept data from

neurons that showed a consistent response to this control. We only kept data from

experiments with a minimum 9 repetitions. The data from repetitions within a neuron were

averaged, and shown as spike histograms (mean ± sem) with 1 ms resolution, after

smoothing with a 20 ms square-wave filter. For quantification across neurons, we calculated

the mean spike rate for each neuron from the spike histogram for the duration of target

motion, after excluding the first and last 40 ms (dotted boxes, Fig. 1b-e). Due to the

response variation across animals (Fig. 1f), the mean spike rate for each TSDN was

normalized to its own mean response to a target moving over a white background.

Data analysis was performed in Matlab and statistical analysis in Prism 7.0c for Mac OS X

(GraphPad Software, San Diego, USA). Throughout the paper n refers to the number of

repetitions within one neurons, and N to the number of neurons. The sample size, type of

test and P value is indicated in each figure legend. We performed either two-sided t-tests,

one-way RM ANOVAs, followed by Dunnett's post hoc test for multiple comparisons or two-

way RM ANOVAs, followed by either Bonferroni's or Tukey's post hoc test for multiple

comparisons. All data have been deposited to DataDryad.

Acknowledgements

451 We thank the Botanic Gardens of Adelaide for their ongoing support. This research was

452 funded by the US Air Force Office of Scientific Research (AFOSR, FA9550-19-1-0294), the

453 Australian Research Council (ARC, DP170100008, DP180100144 and FT180100289), and the

454 Flinders Foundation.

Author Contributions

456 K.N. conceptualized the study and developed the research plan. Both authors designed the

457 study. S.N. performed all experiments and data analyses and prepared the figures. Both

458 authors wrote the manuscript and contributed to the discussion. K.N. supervised all the

459 work.

Competing Interests Statement

461 The authors declare no competing interests.

References

Wellington, W. & Fitzpatrick, S. Territoriality in the drone fly, *Eristalis tenax* (Diptera, Syrphidae). *Can Entomol* **113**, 695-704 (1981).

Zeil, J. Sexual dimorphism in the visual system of flies: The free flight behavior of male Bibionidae (Diptera). *J Comp Physiol A* **150**, 395-412 (1983).

Fabian, S. T., Sumner, M. E., Wardill, T. J., Rossoni, S. & Gonzalez-Bellido, P. T. Interception by two predatory fly species is explained by a proportional navigation feedback controller. *Journal of The Royal Society Interface* **15** (2018).

Schröder, R., Linkem, C. N., Rivera, J. A. & Butler, M. A. Should I stay or should I go? Perching damselfly use simple colour and size cues to trigger flight. *Animal Behaviour* **145**, 29-37, doi:https://doi.org/10.1016/j.anbehav.2018.08.015 (2018).

O'Rourke, C. T., Pitlik, T., Hoover, M. & Fernández-Juricic, E. Hawk eyes II: diurnal raptors differ in head movement strategies when scanning from perches. *PLoS One* **5**, e12169, doi:10.1371/journal.pone.0012169 (2010).

Fitzpatrick, S. M. & Wellington, W. G. Contrasts in the territorial behavior of three species of hover flies (Diptera: Syrphidae). *Can Entomol* **115**, 559-566 (1983).

Corvidae, E. L., Bierregaard, R. O. & Peters, S. E. Comparison of wing morphology in three birds of prey: correlations with differences in flight behavior. *J Morphol* **267**, 612-622, doi:10.1002/jmor.10425 (2006).

Koenderink, J. J. Optic flow. *Vision Res* **26**, 161-179, doi:10.1016/0042-6989(86)90078-7 (1986).

Land, M. F. Visual acuity in insects. *Annu Rev Entomol* **42**, 147-177, doi:DOI 10.1146/annurev.ento.42.1.147 (1997).

10   O'Carroll, D. Feature-detecting neurons in dragonflies. *Nature* **362**, 541-543 (1993).

11   Nordström, K., Barnett, P. D. & O'Carroll, D. C. Insect detection of small targets

moving in visual clutter. *PLoS Biol* **4**, 378-386 (2006).

12   Wiederman, S. D. & O'Carroll, D. C. Discrimination of features in natural scenes by a

dragonfly neuron. *J Neurosci* **31**, 7141-7144, doi:10.1523/JNEUROSCI.0970-11.2011

(2011).

13   Nordström, K. Neural specializations for small target detection in insects. *Curr Opin*

Neurobiol **22**, 272-278, doi:S0959-4388(11)00231-5 [pii]

10.1016/j.conb.2011.12.013 [doi] (2012).

14   Nicholas, S., Supple, J., Leibbrandt, R., Gonzalez-Bellido, P. T. & Nordström, K.

Integration of small- and wide-field visual features in Target-Selective Descending

Neurons of both predatory and non-predatory dipterans. *J Neurosci* **38**, 10725-

10733, doi:10.1523/jneurosci.1695-18.2018 (2018).

15   Namiki, S., Dickinson, M. H., Wong, A. M., Korff, W. & Card, G. M. The functional

organization of descending sensory-motor pathways in *Drosophila*. *eLife* **7**, e34272,

doi:10.7554/eLife.34272 (2018).

16   Chen, C. L. *et al.* Imaging neural activity in the ventral nerve cord of behaving adult

Drosophila. *Nat Commun* **9**, 4390, doi:10.1038/s41467-018-06857-z (2018).

17   Geurten, B. R. H., Nordström, K., Sprayberry, J. D. H., Bolzon, D. M. & O'Carroll, D. C.

Neural mechanisms underlying target detection in a dragonfly centrifugal neuron. *J*

Exp Biol **210**, 3277-3284, doi:doi: 10.1242/jeb.008425 (2007).

18   Barnett, P. D., Nordström, K. & O'Carroll, D. C. Retinotopic organization of small-

field-target-detecting neurons in the insect visual system. *Curr Biol* **17**, 569-578

(2007).

19 Gonzalez-Bellido, P. T., Peng, H., Yang, J., Georgopoulos, A. P. & Olberg, R. M. Eight

pairs of descending neurons in the dragonfly give wing motor centers accurate

population vector of prey direction. *Proc Natl Acad Sci USA* **110**, 676-701,

doi:www.pnas.org/cgi/doi/10.1073/pnas.1210489109 (2013).

20 Olberg, R. M. Object- and self-movement detectors in the ventral nerve cord of the

dragonfly. *J Comp Physiol A* **141**, 327-334 (1981).

21 Dyakova, O., Lee, Y.-J., Longden, K. D., Kiselev, V. G. & Nordström, K. A higher order

visual neuron tuned to the spatial amplitude spectra of natural scenes. *Nat Commun*

6, 8522, doi:10.1038/ncomms9522 (2015).

22 Schuster, S., Strauss, R. & Gotz, K. G. Virtual-reality techniques resolve the visual

cues used by fruit flies to evaluate object distances. *Curr Biol* **12**, 1591-1594,

doi:10.1016/s0960-9822(02)01141-7 (2002).

23 Lehrer, M., Srinivasan, M. V., Zhang, S. W. & Horridge, G. A. Motion cues provide the

bee's visual world with a third dimension. *Nature* **332**, 356-357,

doi:10.1038/332356a0 (1988).

24 Collett, T. S. Peering - Locust Behavior Pattern for Obtaining Motion Parallax

Information. *Journal of Experimental Biology* **76**, 237-241 (1978).

25 Sobel, E. C. The locust's use of motion parallax to measure distance. *Journal of*

comparative physiology. A, Sensory, neural, and behavioral physiology **167**, 579-588,

doi:10.1007/bf00192653 (1990).

26 Nityananda, V. *et al.* A Novel Form of Stereo Vision in the Praying Mantis. *Curr Biol*

28, 588-593.e584, doi:10.1016/j.cub.2018.01.012 (2018).

27 Geurten, B. R., Kern, R., Braun, E. & Egelhaaf, M. A syntax of hoverfly flight

prototypes. *J Exp Biol* **213**, 2461-2475, doi:213/14/2461 [pii]

10.1242/jeb.036079 [doi] (2010).

28   Collett, T. S. & Land, M. F. Visual control of flight behaviour in the hoverfly, *Syritta*

pipiens L. *J Comp Physiol A* **99**, 1-66 (1975).

29   Nicholas, S., Leibbrandt, R. & Nordström, K. Visual motion sensitivity in descending

neurons in the hoverfly. *J Comp Physiol A* **206**, 149-163, doi:10.1007/s00359-020-

01402-0 (2020).

30   Supple, J. A. *et al.* Binocular Encoding in the Damselfly Pre-motor Target Tracking

System. *Curr Biol* **30**, 645-656.e644, doi:10.1016/j.cub.2019.12.031 (2020).

31   Städele, C., Keleş, M. F., Mongeau, J. M. & Frye, M. A. Non-canonical receptive field

properties and neuromodulation of Feature-Detecting neurons in flies. *Curr Biol*,

doi:10.1016/j.cub.2020.04.069 (2020).

32   Wiederman, S. D., Fabian, J. M., Dunbier, J. R. & O'Carroll, D. C. A predictive focus of

gain modulation encodes target trajectories in insect vision. *eLife* **6**,

doi:10.7554/eLife.26478 (2017).

33   Nordström, K., Bolzon, D. M. & O'Carroll, D. C. Spatial facilitation by a high-

performance dragonfly target-detecting neuron. *Biol Lett* **7**, 588-592,

doi:rsbl.2010.1152 [pii]

10.1098/rsbl.2010.1152 [doi] (2011).

34   Keleş, M. F., Hardcastle, B. J., Städele, C., Xiao, Q. & Frye, M. A. Inhibitory

interactions and columnar inputs to an object motion detector in *Drosophila*. *Cell*

Rep **30**, 2115-2124.e2115, doi:10.1016/j.celrep.2020.01.061 (2020).

35   Keleş, M. F. & Frye, M. A. Object-detecting neurons in *Drosophila*. *Curr Biol* **27**, 680-

687, doi:10.1016/j.cub.2017.01.012 (2017).

36 Tanaka, R. & Clark, D. A. Object-displacement-sensitive visual neurons drive freezing

in *Drosophila*. *Curr Biol*, doi:10.1016/j.cub.2020.04.068 (2020).

37 Egelhaaf, M. On the neuronal basis of figure-ground discrimination by relative

motion in the visual system of the fly. III. Possible input circuitries and behavioral

significance of the FD-cells. *Biol Cybern* **52**, 267-280 (1985).

38 Warzecha, A. K., Egelhaaf, M. & Borst, A. Neural circuit tuning fly visual interneurons

to motion of small objects. I. Dissection of the circuit by pharmacological and

photoinactivation techniques. *J Neurophysiol* **69**, 329-339,

doi:10.1152/jn.1993.69.2.329 (1993).

39 Egelhaaf, M. On the neuronal basis of figure-ground discrimination by relative

motion in the visual system of the fly. II. Figure-detection cells, a new class of visual

interneurones. *Biol Cybern* **52**, 195-209 (1985).

40 Reichardt, W. & Poggio, T. Figure-ground discrimination by relative movement in the

visual-system of the fly. Part I: Experimental results. *Biol Cybern* **35**, 81-100 (1979).

41 Liang, P., Heitwerth, J., Kern, R., Kurtz, R. & Egelhaaf, M. Object representation and

distance encoding in three-dimensional environments by a neural circuit in the visual

system of the blowfly. *J Neurophysiol* **107**, 3446-3457, doi:jn.00530.2011 [pii]

10.1152/jn.00530.2011 [doi] (2012).

42 Wiederman, S. D., Shoemaker, P. A. & O'Carroll, D. C. A model for the detection of

moving targets in visual clutter inspired by insect physiology. *PLoS ONE* **3**, e2784,

doi:10.1371/journal.pone.0002784 (2008).

43 Wiederman, S. D., Shoemaker, P. A. & O'Carroll, D. C. Correlation between OFF and

ON Channels Underlies Dark Target Selectivity in an Insect Visual System. *J Neurosci*

33, 13225-13232, doi:33/32/13225 [pii]

10.1523/JNEUROSCI.1277-13.2013 [doi] (2013).

44   Straw, A. D., Warrant, E. J. & O'Carroll, D. C. A 'bright zone' in male hoverfly (*Eristalis*

582   *tenax*) eyes and associated faster motion detection and increased contrast

583   sensitivity. *J Exp Biol* **209**, 4339-4354, doi:10.1242/jeb.02517 (2006).

584   45   Aptekar, J. W., Keles, M. F., Lu, P. M., Zolotova, N. M. & Frye, M. A. Neurons forming

585   optic glomeruli compute figure-ground discriminations in *Drosophila*. *J Neurosci* **35**,

586   7587-7599, doi:10.1523/jneurosci.0652-15.2015 (2015).

587   46   Wu, M. *et al.* Visual projection neurons in the *Drosophila* lobula link feature

588   detection to distinct behavioral programs. *eLife* **5**, e21022, doi:10.7554/eLife.21022

589   (2016).

590   47   Nordström, K. & O'Carroll, D. C. Feature detection and the hypercomplex property in

591   insects. *Trends Neurosci* **32**, 383-391, doi:S0166-2236(09)00091-5 [pii]

592   10.1016/j.tins.2009.03.004 [doi] (2009).

593   48   Collett, T. S. & Land, M. F. How hoverflies compute interception courses. *J Comp*

594   *Physiol A* **125**, 191-204 (1978).

595   49   Wardill, T. *et al.* A novel interception strategy in a miniature robber fly with extreme

596   visual acuity. *Curr Biol* **27**, 854-859 (2017).

597   50   Fujiwara, T., Cruz, T. L., Bohnslav, J. P. & Chiappe, M. E. A faithful internal

598   representation of walking movements in the Drosophila visual system. *Nat Neurosci*

599   **20**, 72-81, doi:10.1038/nn.4435 (2017).

600   51   Kim, A. J., Fitzgerald, J. K. & Maimon, G. Cellular evidence for efference copy in

601   *Drosophila* visuomotor processing. *Nat Neurosci* **18**, 1247-1255,

602   doi:10.1038/nn.4083 (2015).

52   Ache, J. M., Namiki, S., Lee, A., Branson, K. & Card, G. M. State-dependent

decoupling of sensory and motor circuits underlies behavioral flexibility in

Drosophila. *Nat Neurosci* **22**, 1132-1139, doi:10.1038/s41593-019-0413-4 (2019).

53   Nicholas, S., Thyselius, M., Holden, M. & Nordström, K. Rearing and long-term

maintenance of *Eristalis tenax* hoverflies for research studies. *JoVE*, e57711,

doi:doi:10.3791/57711 (2018).

54   Brainard, D. H. The Psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

55   Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision* **10**, 437-442 (1997).

Figure Legends

Figure 1. The TSDN response to target motion is robust against stationary optic flow

a, Pictograms of the round, black target, with a diameter of 3°, traversing a white

background (left) or stationary optic flow (right) at 130°/s. **b,** Raw data trace from an

extracellular TSDN recording. Timing of stimulus presentation indicated by colored bars

(blue, target, and red, optic flow). **c,** Magnification of the raw data traces shown in panel **b.**

d, Spike raster of the same neuron in response to 18 repeated trials of a target on a white

background (left) or on stationary optic flow (right). **e,** Response shown as spike histograms

of the mean ± sem shown with 1 ms resolution, after smoothing with a 20 ms square-wave

filter. **f,** The mean spiking response of different TSDNs was significantly reduced when the

target moved across stationary optic flow compared with a white background (N = 24;

P = 0.0019, two-sided paired t-test). In **b-e** the dotted box shows the analysis window used

for quantification. **P < 0.01.

Figure 2. The TSDN response to target motion is affected by horizontal optic flow

a, The response of a single TSDN to a target traversing stationary optic flow (left, grey) or

sideslip or yaw in either the same direction or opposite direction to the target (see red

pictograms). All histograms from a single TSDN (mean ± sem, n = 18) shown with 1 ms

resolution after smoothing with a 20 ms square-wave filter. **b**, The mean response across

TSDNs (N = 12) to a target traversing stationary optic flow (left, grey), or sideslip or yaw in

either the same direction ("+50") or opposite direction ("-50") to the target after

normalizing the data to the response to a target traversing a white background (one-way

RM ANOVA followed by Post hoc Dunnett's test: sideslip +50, P < 0.0001; sideslip -50,

P < 0.0001, yaw +50, P < 0.0001; yaw -50, P < 0.0001). Sideslip optic flow was shown at

50 cm/s and yaw optic flow at 50°/s. The line shows the mean and the filled data points are

from the neuron shown in panel **a**. ****P < 0.0001.

Figure 3. The TSDN response to target motion is affected by 3D optic flow

a, The response of a single TSDN to a target traversing stationary optic flow (left, grey,

replotted from Fig. 2a), or different types of optic flow (see red pictograms). Translations

were shown at 50 cm/s and rotations at 50°/s. All histograms shown as mean ± sem (n = 18)

at 1 ms resolution, after smoothing with a 20 ms square-wave filter. **b**, The mean response

across TSDNs (N = 12) to a target traversing stationary optic flow (left, grey, replotted from

Fig. 2b), or different types of optic flow (one-way RM ANOVA followed by Post hoc

Dunnett's test: lift +50, P < 0.0001; lift -50, P = 0.0006; pitch +50, P < 0.0001; pitch -50,

P < 0.0001; thrust +50, P < 0.0001; thrust -50, P < 0.0001; roll +50, P = 0.0020; roll -50,

P < 0.0001). The lines show the mean and the filled data points are from the neuron shown

in panel **a**. **P < 0.01, ***P < 0.001 and ****P < 0.0001.

Figure 4. The TSDN inhibition by syn-directional sideslip can be primed

a, Typical raw data trace from a single TSDN to a target presented on sideslip optic flow,

which moved in the same direction as the target prior to target presentation, then remained

stationary during target motion. Timing is color coded: green, preceding optic flow; red,

concurrent optic flow; blue, target motion. **b**, TSDN target responses are significantly

inhibited by preceding sideslip in the same direction as the target, compared with preceding

stationary optic flow (N = 7, two-way RM ANOVA followed by Post hoc Bonferroni's test:

P = 0.0005). **c**, There was no significant difference between TSDN responses to targets

moving across sideslip optic flow in the same direction, whether this moved or not

preceding target motion (N = 7, two-way RM ANOVA: Post hoc Bonferroni's test:

P = 0.4704). **d**, There was no significant difference between TSDN responses to targets

moving across stationary optic flow, whether this moved or not preceding target motion

(N = 7, two-way RM ANOVA: Post hoc Bonferroni's test: P > 0.9999). The grey data is the

same as in panel **b**. **e**, There was no significant difference between TSDN responses to

targets displayed on sideslip optic flow in the opposite direction to target motion, whether

the optic flow moved or not preceding target motion (N = 7, two-way RM ANOVA: Post hoc

Bonferroni's test: P > 0.9999). **b-e**, The lines show the mean. NS P > 0.05 and ***P < 0.001.

Figure 5. The inhibition of TSDN target responses by syn-directional sideslip comes from

the pre-synaptic input

a, The response of a single TSDN to a target traversing the white background horizontally

(left, grey) or vertically (right, black). All histograms shown as mean ± sem (n = 18) at 1 ms

resolution, after smoothing with a 20 ms square-wave filter. **b**, The mean spiking response

of the same 8 TSDNs to targets moving horizontally (grey, left) or vertically (black, right)

over a white background (N = 8). **c**, TSDNs are likely to receive input from STMDs (dashed

arrow). If the inhibition from widefield motion is post-synaptic to the STMDs, the inhibition

will depend on the direction of optic flow. For example, since the TSDN response is strongly

inhibited by sideslip optic flow (grey data, replotted from Fig. 2), changing the direction of

target motion should not affect this inhibition. Similarly, since the TSDN response is less

inhibited by lift optic flow (grey data, replotted from Fig. 3), changing the direction of target

motion should not affect this. **d**, If the inhibition from widefield motion is pre-synaptic to

the STMDs, the inhibition will depend on the direction of target motion. For example,

changing the target direction from horizontal (grey data, replotted from Fig. 2), to vertical

when displayed against sideslip optic flow, should increase the response. Similarly, changing

the target direction to syn-directional when displayed against lift optic flow, should increase

the inhibition, and decrease the response. **e**, TSDN responses to targets moving across

sideslip optic flow are significantly increased when the target direction changes from syn-

directional to vertical (N = 8, two-way RM ANOVA followed by Post hoc Tukey's test:

P = 0.0413). **f**, TSDN responses to targets moving across lift optic flow are significantly

decreased when the target direction changes from perpendicular to syn-directional (N = 8,

two-way RM ANOVA: Post hoc Tukey's test: P = 0.0014). **e-f**, The lines show the mean.

*P < 0.05 and **P < 0.01.

Figure 6. There is no facilitation to targets moving across lift optic flow

a, If the facilitation of TSDNs is due to input from the widefield system post-synaptic to the

STMDs, the facilitation will depend on the direction of optic flow. For example, since the

TSDN response is strongly facilitated by sideslip optic flow (grey data, replotted from Fig. 2),

changing the direction of target motion should not affect this facilitation. Similarly, since the

TSDN response is not facilitated by lift optic flow (grey data, replotted from Fig. 3), changing

the direction of target motion should not result in any facilitation. **b**, If the facilitation is

inherited from the STMDs, the facilitation will depend on the direction of target motion. For

example, changing the target direction from horizontal (grey data, replotted from Fig. 2), to

vertical when displayed against sideslip optic flow, should increase the response. Similarly,

changing the target direction to vertical when displayed against the opposing direction of

lift optic flow, should increase the response. **c**, TSDN responses to targets moving across

sideslip optic flow are significantly decreased when the target direction changes from

moving in the opposite direction to perpendicular to the optic flow (N = 8, two-way RM

ANOVA: Post hoc Tukey's test: P = 0.0017). **d**, TSDN responses to targets moving across lift

optic flow are not significantly affected when the target direction changes from

perpendicular to moving in the opposite direction to the optic flow (N = 8, two-way RM

ANOVA: Post hoc Tukey's test: P = 0.5394) **c-d**, The lines show the mean. NS P > 0.05 and

**P < 0.01

Extended Data

Extended Data Figure 1. TSDN response to 3D optic flow

a, To determine the TSDN response to the optic flow we first quantified each neuron's spontaneous rate, i.e. the response to a white background. We used this as a threshold. For each neuron we then quantified the number of trials in which the response to the optic flow was above this spontaneous rate. This was converted to a percentage of responses. The data here show this percentage of responses across 12 TSDNs as mean ± sem. **b**, In each neuron, we quantified the mean response for those trials that were classified as responders (in panel **a**). This mean response was normalized to each neuron's mean response to a target traversing a white background. Data are shown as mean ± sem (N = 12).

Extended Data Movie 1. TSDN response to a target over a white background

The movie depicts the stimulus as displayed on the screen, with the hoverfly positioned centered and perpendicular to the screen. The hoverfly was ventral side up, but we have rotated the movie for display purposes for the dorsal side to be up. The colored lines show the outline of the receptive field, mapped as described previously[29]. The red data at the bottom of the movie show the response of an example TSDN.

Extended Data Movie 2. TSDN response to a target over a white background

Same as Movie 1, but slowed down 10 times.

Extended Data Movie 3. TSDN response to a target over stationary optic flow

The movie depicts the stimulus as displayed on the screen, with the colored lines showing

the outline of the receptive field. The red data at the bottom of the movie show the

response of an example TSDN. In this case the optic flow was stationary, and appeared 0.5 s

before the target.

Extended Data Movie 4. TSDN response to a target over stationary optic flow

Same as Movie 3, but slowed down 10 times.

Extended Data Movie 5. TSDN response to a target over syn-directional sideslip optic flow

The movie depicts the response of the same example TSDN to the same target motion as in

previous movies, but now traversing sideslip optic flow appearing 0.5 s before the target.

The hoverfly was positioned at a distance of 6.5 cm. At this viewing distance, the resulting

optic flow simulates the type of optic flow that would be generated by the hoverfly

sideslipping through the world at 50 cm/s. The simulated optic flow consisted of ca 6400

spheres, of which ca 1200 are projected onto the screen at any one time.

Extended Data Movie 6. TSDN response to a target over syn-directional sideslip optic flow

Same as Movie 5, but slowed down 10 times. Note the many optic flow "targets" of the

correct size that move through the receptive field (colored contour lines).

Extended Data Movie 7. TSDN response to a target over opposite direction sideslip optic

flow

The movie depicts the response of the same example TSDN to the same target motion as in

previous movies, but now traversing sideslip optic flow in the opposite direction to the

target.

Extended Data Movie 8. TSDN response to a target over opposite direction sideslip optic

flow

Same as Movie 7, but slowed down 10 times.

a

White background

Stationary

b

200 μV

200 ms

200 μV

200 ms

Target
Optic flow

c

200 μV

100 ms

200 μV

100 ms

d

Trial number

100 ms

Trial number

100 ms

e

White
background

Stationary

100 spikes/s

100 ms

f

Mean response (spikes/s)

**

White
background

Stationary

a

Target

100 spikes/s

100 ms

b

Normalized response

Optic flow

Stationary

Sideslip

Yaw

+ 50   − 50   + 50   − 50

****   ****   ****   ****

a

Target

100 spikes/s

100 ms

b

Normalized response

Optic flow

Stationary

Lift
+ 50   − 50

Pitch
+ 50   − 50

Thrust
+ 50   − 50

Roll
+ 50   − 50

**** *** **** **** **** **** ** ****