Facilitation of neural responses to targets moving in three-dimensional optic

flow

Sarah Nicholas[1] and Karin Nordström[1,2]*

[1]Flinders Health and Medical Research Institute, Flinders University, GPO Box 2100, Adelaide,

SA, 5001, Australia

[2]Department of Neuroscience, Uppsala University, Box 593, 751 24 Uppsala, Sweden

*email: karin.nordstrom@flinders.edu.au

ORCIDs

Sarah Nicholas: 0000-0002-5555-9421

Karin Nordström: 0000-0002-6020-6348

Classification

Biological sciences: Neuroscience

Keywords

Target motion, insect vision, motion vision, background motion, hoverfly, TSDN, wide-field

motion, inhibition, facilitation

Abstract

For the human observer, it can be difficult to follow the motion of small objects, especially

when they move against background clutter. However, insects efficiently do this, as

evidenced by their ability to capture prey, pursue conspecifics, or defend territories, even in

highly textured surrounds. This behavior has been attributed to optic lobe neurons that are

sharply tuned to the motion of small targets, as these neurons respond robustly even to a

target moving against background motion. However, the target selective descending

neurons (TSDNs), that more directly control behavioral output, do not. Importantly, though,

the backgrounds used previously not only lacked 3D motion cues, but also high-contrast

features, both of which would be encountered during natural behaviors. To redress this

deficiency, we here use backgrounds consisting of many targets moving coherently to

simulate the type of 3D optic flow that would be generated by an insect's own motion

through the world. We show that hoverfly TSDNs do not respond to this type of optic flow,

even though it contains features with spatio-temporal profiles similar to optimal targets.

However, TSDN responses are inhibited when this optic flow is shown together with a

target. More surprisingly, TSDNs are facilitated by horizontal, frontal optic flow in the

opposite direction to target motion. We show that these interactions are likely inherited

from the pre-synaptic neurons, and argue that the facilitation could benefit the initiation of

target pursuit.

Significance statement

Target detection in visual clutter is a difficult computational task that insects, with their poor resolution compound eyes and small brains, do successfully and with extremely short behavioral delays. We here show that target neurons do not respond to widefield motion consisting of a multitude of "targets", suggesting that the hoverfly visual system interprets coherent widefield motion differently from the motion of individual targets. In addition, we show that widefield motion in the opposite direction to target motion increases the neural response. This is an incredibly non-intuitive finding, and difficult to reconcile with current models for target selectivity, but has behavioral relevance.

/body

Introduction

The survival of many animals often depends on their ability to visually detect small moving

objects or targets, as these could represent predators, prey, mates or territorial intruders

(1). Efficient target detection is a computationally challenging task, which becomes even

more difficult when done against visual clutter. Despite this, many insects successfully

detect targets, followed by highly acrobatic pursuits, often in visually complex

environments. For example, male *Eristalis tenax* hoverflies establish territories in foliage rich

areas, on alert for intruders and ready to engage in high-speed pursuit (2).

Initial target detection can be facilitated by behaviors that render the background

stationary, thus making the target the only thing that moves. Many insects and vertebrates

thus visualize targets against the bright sky (3), or from a stationary stance, such as perching

(4-6) or hovering (7, 8). However, as soon as the pursuer moves, this creates self-generated

optic flow (9), against which the independent motion of the target needs to be

discriminated. That insects do this successfully is remarkable considering their small brains

and low-resolution compound eyes (10), suggesting a high level of optimization.

Predatory dragonflies and territorial hoverflies, which both pursue targets, have sharply

tuned small target motion detector (STMD) neurons in their optic lobes (11, 12).

Impressively, some STMDs respond robustly to targets in clutter, even without relative

velocity differences (12, 13), suggesting that they could support behavioral pursuit against

self-generated optic flow. However, the target selective descending neurons (TSDNs), which

are thought to be post-synaptic to STMDs (14), do not respond to targets moving across syn-

directional background motion (15). This is peculiar as descending neurons more directly

drive behavioral output (16, 17). Importantly, however, the backgrounds used in previous

STMD (12, 13, 18) and TSDN (15) experiments consisted of panoramic images that were

displaced across the visual monitor, lacking the 3D cues associated with real pursuits (19,

20). In addition, even if some backgrounds had spatial statistics resembling natural images

(21) they lacked the high-contrast edges often found in natural scenes. Where naturalistic

backgrounds contained small, high-contrast features, these would often generate STMD

responses (13). Indeed, it is highly likely that STMDs generate their ability to detect targets

in clutter by being sharply tuned to the target's unique spatio-temporal profile (22).

To investigate TSDN responses to targets in background motion, under more dynamic

conditions, we used an optic flow stimulus consisting of many "targets" simulated to move

coherently around the fly's point of view (23). We use this to simulate the type of optic flow

generated during translations and rotations through the world. Importantly, the individual

components that make up the optic flow contain the same high-contrast edges as the target

itself, against which target motion can only be determined by its independent trajectory.

We quantified the responses of *Eristalis* TSDNs to targets moving against 6 types of optic

flow (23), three translations and three rotations. We found that optic flow on its own did

not generate any TSDN response, even when moving in the neuron's preferred direction,

and despite it consisting of many "targets". Consistent with previous results (15), we found

that optic flow in the same direction as the target inhibits the TSDN response. Importantly,

inhibition occurred even if the optic flow only preceded target motion and did not appear

concurrently. Most strikingly we showed that optic flow in the opposite direction to target

motion increased the TSDN response. This facilitation required opposing horizontal optic

flow in the frontal visual field. Such neural facilitation is a novel observation.

Results

3D optic flow strongly affects the TSDN response to target motion

We recorded extracellularly from target selective descending neurons (TSDNs) in male

Eristalis tenax hoverflies. TSDNs respond strongly to the motion of a small, dark target (15,

23), traversing a white background (Fig. 1A, B, Movie 1, 2, Fig. S1A-C). We used a 3D optic

flow stimulus simulating the coherent motion of many "targets" (23) around the hoverfly's

position. Against this optic flow, representing retinal flow fields experienced during flight

through the world, the target can only be identified by its independent trajectory. Indeed,

when both are stationary, there are no features identifying the target (Fig. 1A). Despite this,

TSDNs respond strongly to targets moving across stationary optic flow (Fig. 1B; Movie 3, 4;

Fig. S1A-C).

For quantification, we calculated the mean spike frequency for the duration of target

motion, after excluding the first and last 40 ms (dotted box, Fig. S1A-C). We found a small,

significant response reduction to targets moving across stationary optic flow compared with

a white background (Fig. 1B). As the response across neurons was variable (N=34,

coefficient of variation 60% and 63%, respectively, Fig. 1B), we normalized the response

from each neuron to its own mean response to a target moving across a white background.

We found that when the target moved horizontally across sideslip optic flow moving in the

same direction, the TSDN response was strongly inhibited (Fig. 1C, D, "Sideslip +50"),

compared with control where the target moved over stationary optic flow (grey, Fig. 1C, D).

Yaw optic flow in the same direction as the target also strongly inhibited the TSDN response

(Fig. 1C, D, "Yaw +50"). This complete inhibition is striking (Movie 5, 6), but consistent with

previous TSDN work using sliding background images (15).

In contrast, when the sideslip moved in the opposite direction to the target the TSDN

response was strongly enhanced (Movie 7, 8, Fig. 1C, D, "Sideslip -50", mean increase

71.1%). Similarly, yaw optic flow in the opposite direction to the target also facilitated the

TSDN response, by 84.9% (Fig. 1C, D, "Yaw -50"). Such response facilitation was not seen

when displaying targets moving over panoramic background images, in either TSDNs (15) or

other target tuned neurons in the fly optic lobes (12, 18, 24).

At any one time the optic flow itself contained ca. 1200 "targets" moving coherently on the

display in front of the fly (23). When the sideslip moved in the TSDN's preferred direction

tens of these "optic flow targets" moved through the TSDN receptive field (Movie 5, 6).

Despite this, preferred direction sideslip without target motion only generated a response in

18% of 222 repetitions across 12 TSDNs (Fig. S1D, "Sideslip +50"). In addition, when

preferred direction sideslip did generate a TSDN response, this was less than one-fifth of the

response to a target traversing a white background (Fig. S1E, "Sideslip +50"). This suggests

that the coherent motion of the "targets" in the optic flow is treated differently from the

motion of a single target.

We next investigated the effect other types of 3D optic flow had on TSDN responses to

target motion. We found that the target response was inhibited by lift (mean 44.0%

response for downwards lift, "Lift +50"; mean 34.3% for upwards lift, "Lift -50", Fig. S2).

When the target was displayed against pitch, which provides similar vertical motion in the

frontal visual field, the response was suppressed to similar levels (57.1%, "Pitch +50";

54.9%, "Pitch -50", Fig. S2). In addition, the target response was inhibited to roughly similar

levels when displayed against thrust or roll optic flow in either direction (Fig. S2). No optic

flow generated a substantial TSDN response when displayed on its own (Fig. S1D, E).

Inherited optic flow interactions

Our data show that optic flow significantly affects the TSDN response to target motion (Fig.

1, S2). Where does the optic flow pathway interact with the target motion pathway? If

widefield sensitive neurons interact directly with the TSDNs, as previously suggested (15),

the optic flow should act independently. That is, if the target changes direction but the optic

flow remains the same, the influence the optic flow has on the TSDN response should

remain similar. In contrast, if the interaction is inherited from pre-synaptic neurons, we

expect the TSDN response to show significant interaction between target direction and optic

flow direction.

We investigated this by recording from TSDNs that respond robustly both to horizontal

(grey, Fig. S3) and vertical target motion (black, Fig. S3), displayed against either sideslip or

lift (Fig. 2). As above (Fig. 1, S2), the TSDN response was inhibited when a horizontally

moving target was displayed against sideslip or lift (Fig. 2A). If the target instead moved

vertically against sideslip or lift, a two-way ANOVA showed that neither the target direction

nor the optic flow direction had a significant effect on the TSDN response (Fig. 2A).

However, there was a significant interaction between target direction and optic flow

direction (P=0.0002, two-way ANOVA), consistent with the hypothesis that the interaction is

inherited from the pre-synaptic neurons.

As above (Fig. 1, S2), the TSDN response was facilitated when a target moved horizontally

against sideslip in the opposite direction, but inhibited against lift (Fig. 2B). When comparing

this to vertical target motion, a two-way ANOVA showed that both the target direction and

the optic flow direction significantly affected the TSDN response (Fig. 2B, P=0.0092 and

P=0.0228, respectively). In addition, there was a significant interaction between target

direction and optic flow direction (P=0.0010, two-way ANOVA, Fig. 2B), consistent with the

hypothesis that the interaction is inherited from pre-synaptic neurons.

Presynaptic neurons

From which pre-synaptic neurons do the TSDNs inherit these interactions? TSDNs have been

proposed to get their input from STMDs (25), but this has never been shown conclusively,

and in addition, there are many other target tuned neurons in the fly optic lobes (e.g. 24,

26-29). We can investigate this by looking at the underlying target tuning mechanisms,

which can be distilled down into three fundamentally different concepts. For example,

visual neurons can become target tuned by receiving inhibitory feedback from the widefield

system (29-31), or by using center-surround antagonism together with rapid adaptation (28,

32). Alternatively, they can use an elementary STMD model, which is tuned to the unique

spatio-temporal profile of a moving target, with a dark contrast change (OFF) from the

leading edge followed by a bright contrast change (ON) by the trailing edge. Importantly,

while the first two mechanisms rely on comparisons from neighboring points in space, the

elementary STMD compares input from one point on space (22, 28, 33). Therefore, the first

two models will respond equally well to the motion of a target, to the motion of a leading

OFF edge, and the motion of a trailing ON edge (black, Fig. 3, redrawn from (28, 34)). In

contrast, the elementary STMD model only responds strongly to the target (grey, Fig. 3,

redrawn from (34)).

Our results show that TSDNs do not respond well to a leading OFF edge, or to a trailing ON

edge (white, Fig. 3). However, a complete target, where the leading edge is rapidly followed

by a trailing edge, gives a robust TSDN response (white, Fig. 3). Indeed, the physiological

responses (white, Fig. 3) match the elementary STMD model output (grey, Fig. 3). Since

STMD physiology also matches the elementary STMD model output (13, 34), this suggests

that TSDNs receive input from STMDs.

TSDN responses are inhibited by preceding optic flow

In dragonflies, STMD target responses can be primed by preceding target motion, thereby

facilitating responses to longer target trajectories (35, 36). To investigate if the optic flow

can similarly prime TSDN responses, we displayed preceding optic flow for 1 s (green, Fig.

4A), followed by optic flow concurrent (red, Fig. 4A) with target motion (blue, Fig. 4A). We

found that when the sideslip moved in the same direction as the target, both the preceding

and concurrent optic flow had a significant effect on the TSDN response (2-way ANOVA,

$P=0.0032$ for preceding optic flow and $P=0.0003$ for concurrent optic flow). A Bonferroni

multiple comparison's test showed that the preceding optic flow had a significant effect

when the optic flow was stationary concurrent with target motion (Fig. 4B), but not when it

was moving (Fig. 4C).

In the inverse experiment, when the sideslip moved in the opposite direction to the target,

only the concurrent optic flow had a significant effect on the TSDN response (2-way ANOVA,

P=0.58 for preceding, and P=0.0056 for concurrent optic flow). Thus, preceding sideslip in

the opposite direction to target motion did not facilitate the TSDN response (Fig. 4D).

Neither did the facilitation depend on whether the optic flow was in motion or stationary

prior to target motion (Fig. 4E).

Frontal optic flow is essential

Our results show that yaw and sideslip optic flow have similar effects on TSDN target

responses (Fig. 1). As both sideslip and yaw contain substantial motion in the frontal visual

field (37), we next asked if frontal optic flow is required. We investigated this by limiting the

spatial extent of sideslip to either cover the ipsilateral, dorsal, ventral or contralateral

position on the screen (Fig. 5A). Note that only the dorsal position covers the TSDN

receptive field (Fig. 5A). In the other three positions, the sideslip was spatially separated

from the receptive field (Fig. 5A).

We found that when sideslip moved in the same direction as the target, the TSDN response

was inhibited if the optic flow covered the full, dorsal or ventral position on the screen (Fig.

5B), compared with the stationary control (grey, Fig. 5B). However, when optic flow only

covered the ipsilateral or contralateral positions, there was no inhibition (Fig. 5B),

suggesting that frontal optic flow is required. Our results are consistent with inhibition

driven by panoramic background images, which do not have to spatially overlap the target

trajectory, or have a large spatial extent (15).

We found that when sideslip moved in the opposite direction to the target, the TSDN

response was facilitated if the optic flow covered the full, dorsal or ventral positions on the

screen (Fig. 5C). When the optic flow was limited to the ipsilateral or contralateral positions,

there was no facilitation (Fig. 5C). This suggests that the optic flow does not have to spatially

overlap with the TSDN receptive field. However, there has to be frontal opposite direction

sideslip for facilitation to take place.

Discussion

We generated optic flow using the coherent motion of thousands of "targets" within a

simulated 3D space (23) and recorded from TSDNs, which respond to the motion of small

targets (Fig. 1). We found that even when the "targets" in the optic flow moved in the

neuron's preferred direction, there was no response (Fig. S1D, E). However, when the optic

flow was displayed together with a target, this modulated the TSDN's target response (Fig.

1, S2). In particular, optic flow in the opposite direction to target motion strongly enhanced

the TSDN response, whereas optic flow in the same direction inhibited the response (Fig. 1).

The effect optic flow has on the TSDN response to target motion (Fig. 1, S2) could be

inherited from the neurons that are pre-synaptic to the TSDNs, or alternatively, TSDNs could

receive input from widefield motion sensitive neurons, as previously suggested (15).

However, we found e.g. that while the response to a target moving horizontally across syn-

directional sideslip was strongly inhibited, if the target moved vertically against the same

sideslip, the TSDN response was much larger (Fig. 2A). This argues against direct input from

the widefield motion vision pathway, as we would expect the same optic flow to have the

same effect on the TSDN response, irrespective of target direction. Instead, we found

significant interactions between target direction and optic flow direction (Fig. 2), making it

likely this is inherited from the pre-synaptic neurons.

What are the pre-synaptic candidates? Even if TSDNs have been suggested to get input from

STMDs (25), there are other optic lobe neurons tuned to the motion of targets, also referred

to as objects, or figures (24, 26-29). For example, blowfly lobula plate figure detection (FD)

cells receive inhibitory input from widefield motion sensitive neurons (38), thereby making

them respond to the independent motion of a figure (29-31). However, FD cells are not as

sharply tuned to small targets (33) as TSDNs (15), and they are inhibited by background

motion in the same and in the opposite direction (29), unlike our findings (Fig. 1C, D),

making it unlikely that they provide the TSDN input.

Another option are the anatomically defined *Drosophila* lobula columnar (LC) neurons (39).

Whereas some LC neurons respond better to bars or widefield motion, LC11 and LC26

neurons respond strongly to small targets (24, 40). Besides being sharply size-tuned, LC11

neurons are clearly involved in behavioral responses to small objects (27, 28). LC11 small

object selectivity can be explained using a combination of center-surround antagonism and

rapid adaptation (28), making them inhibited by background motion in either direction (26),

inconsistent with our TSDN facilitation (Fig. 1C, D). Potentially, facilitation could be achieved

by making the influence of the surround directional (32). However, as the optic flow did not

have to cover the TSDN receptive field to have an effect ("ventral", Fig. 5), such center-

surround mechanisms are unlikely.

Importantly, as both FD cells and LC11 neurons rely on an input from at least two points in

space (28, 33), they respond to high-contrast changes associated with sweeping edges (28,

34), as well as to complete targets (black, Fig. 3). In contrast, TSDNs respond much better to

a complete target, than to either OFF or ON contrast changes (white, Fig. 5), i.e. consistent

with input from elementary STMDs (22). Since the elementary STMD model provides robust

predictions of physiological STMD responses in dragonflies (13, 34), we conclude that it is

likely that TSDNs get their input from STMDs.

At a first glance, this seems counterintuitive, as some STMDs respond robustly to targets in

background motion, even without relative motion cues (12), whereas we saw inhibition to

targets displayed against syn-directional optic flow (Fig. 1C, D). In addition, the elementary

STMD model predicts that target-like features in the background should generate a

response (22). In our TSDN experiments, despite the optic flow consisting of "targets" with

optimal spatio-temporal profiles (Movie 5, 6), there were no consistent responses to

preferred direction optic flow (e.g. "Sideslip +50", "Yaw +50", Fig. S1D, E, see also raw data,

Fig. 4A).

However, STMDs are a heterogeneous group (12), and in some STMDs responses to target

motion are decreased by syn-directional background motion (18). Indeed, the centrifugal

STMDs that respond most robustly to targets against background motion (12, 13), and that

respond to background features with target-like profiles (13), do not project to pre-motor

areas, but contralaterally through the protocerebrum, with outputs in the heterolateral

lobula (12, 41). This suggests a role in modulating the responses of other optic lobe neurons,

rather than providing direct input to descending neurons (14). In contrast, the STMDs that

have small receptive fields, and outputs in pre-motor areas of the lateral mid-brain, and that

are therefore more likely to be pre-synaptic to the TSDNs, are often inhibited by background

motion (18).

Our data suggest that inhibition and facilitation are not necessarily generated by the same

mechanism. For example, most types of optic flow generated inhibition (Fig. S2), but only

opposite direction yaw and sideslip led to facilitation (Fig. 1C, D). In addition, unlike

facilitation, inhibition could be primed (Fig. 4). Importantly, the facilitation of TSDN

responses to target motion displayed on optic flow in the opposite direction (Fig. 1C, D) is a

novel observation. In previous work using panoramic background images, TSDN responses

were significantly reduced when the background moved in the opposite direction to the

target (15). This suggests that facilitation is not generated by a widefield motion sensitive

neuron, as such a neuron would respond to panoramic background images (15) as well as to

optic flow (23).

The elementary STMD model predicts a response to the "targets" in the optic flow (22), as

these have correct spatio-temporal profiles. While the TSDN response matches the

elementary STMD model (Fig. 3), TSDNs do not respond to optic flow when displayed alone

(Fig. S1D, E), therefore suggesting pre-synaptic inhibition from the widefield motion

pathway (Fig. 2). However, STMDs with input from elementary STMDs, but without

inhibition from widefield motion, should respond to the "targets" in optic flow, as

evidenced by dragonfly centrifugal STMDs (13). In contrast, background images lacking high-

contrast features do not generate a response in centrifugal STMDs (12, 13). Therefore, such

STMDs may drive the TSDN facilitation. Given facilitation requires frontal, horizontal motion

(Fig. 5), these neurons would likely have frontal receptive fields. The centrifugal STMDs have

large receptive fields (12, 41), suggesting that ipsilateral or contralateral optic flow should

also drive facilitation, which it did not (Fig. 5). Importantly, however, there are at least 20

different STMD types in the hoverfly lobula (12), suggesting that future investigation of

STMD responses to targets in clutter are required to identify the neurons driving facilitation.

Nevertheless, the facilitation could make behavioral sense. Prior to initiating target pursuit,

male *Eristalis* hoverflies have been described to predict the flight course required to

successfully intercept the target, based predominantly on the target's angular velocity (42).

To successfully execute an interception style flight course, the hoverfly turns in the direction

that the target is moving (42). In doing so, the hoverfly creates self-generated optic flow in

the opposite direction to the target's motion. If the TSDNs that we have recorded from here

are involved in such planning of pursuit, they would be facilitated when the target moves in

the opposite direction to the self-generated optic flow (Fig. 1C, D), which might be beneficial

for controlling behavioral output. Reconstructing retinal flow fields as experienced during

actual pursuits (19) could help address this question.

Materials and Methods

Electrophysiology

Eristalis tenax hoverflies were reared and maintained as previously described (43). For

electrophysiology, a male hoverfly was immobilized ventral side up using a beeswax and

resin mixture. A small hole was cut at the anterior end of the thorax to expose the cervical

connective, which was then raised slightly and supported using a small wire hook, for

insertion of a sharp polyimide-insulated tungsten microelectrode (2 MOhm, Microprobes,

Gaithersburg, USA). The animal was grounded via a silver wire inserted into the ventral part

of the hole.

Extracellular signals were amplified at 100x gain and filtered through a 10- to 3000-Hz

bandwidth filter on a DAM50 differential amplifier (World Precision Instruments, Sarasota,

USA), with 50 Hz noise removed with a HumBug (Quest Scientific, North Vancouver,

Canada). The data were digitized via Powerlab 4/30 (ADInstruments, Sydney, Australia) and

acquired at 40 kHz with LabChart 7 Pro software (ADInstruments). Single units were

discriminated by amplitude and half-width using Spike Histogram software (ADInstruments).

Visual stimuli

Eristalis males were placed ventral side up, centered and perpendicular to an Asus LCD

screen (Asus, Taipai, Taiwan) at 6.5 cm distance. The screen had a refresh rate of 165 Hz, a

linearized contrast with a mean illuminance of 200 Lux, and a spatial resolution of 2560 x

1440 pixels, giving a projected screen size of 155° x 138°. Visual stimuli were displayed using

custom written software based on the Psychophysics toolbox (44, 45) in Matlab

(Mathworks, Natick, USA).

TSDNs were identified as described (15, 23). In short, we mapped the receptive field of each

neuron by scanning a target horizontally and vertically at 20 evenly spaced elevations and

azimuths (23), to calculate the local motion sensitivity and local preferred direction. We

then scanned targets of varying height through the small, dorso-frontal receptive fields (Fig.

4A) to confirm that each neuron was sharply size tuned with a peak response to targets

subtending 3°- 6°, with no response to larger bars, to looming or to widefield stimuli (15,

23).

Unless otherwise mentioned, targets were black and round with a diameter of 15 pixels,

moving at a velocity of 900 pixels/s for 0.48 s. When converted to angular values and taking

the small frontal receptive fields of TSDNs into account, this corresponds to an average

diameter of 3° and a velocity of 130°/s (15). Unless otherwise stated, each target travelled in

each neuron's preferred horizontal direction (i.e. left or right) and across the center of its

receptive field. Between repetitions, we varied the target elevation slightly, to minimize

habituation (15). There was a minimum 4 s between stimulus presentations. Stimulus order

was randomized.

For input mechanism experiment (Fig. 3), targets were black and square with a side of 15

pixels, moving at a velocity of 900 pixels/s. OFF and ON edges had a height of 15 pixels. All

stimuli started at the far edge of the screen moving in each neuron's preferred direction,

across the entire width of the screen.

3D optic flow was generated as previously described (23). Briefly, the optic flow consisted of

a simulated cube with 4 m sides, filled with 2 cm diameter spheres at a density of

100 per m^3, with the hoverfly placed in the center. The coherent motion of these ca 6400

spheres around the hoverfly was used to simulate self-generated optic flow. The ca 1200

spheres anterior to the hoverfly were projected onto the screen, with their size indicating

the distance from the hoverfly. Circles closer than 6 cm were not displayed. Six types of

optic flow were simulated: three translations at 50 cm/s (sideslip, lift and thrust) and three

rotations at 50°/s (yaw, pitch and roll). Unless otherwise stated optic flow was displayed for

0.48 s prior to the target. Both target motion and optic flow disappeared simultaneously.

In most experiments the optic flow covered the entire visual display. In some experiments,

we limited the spatial extent of the optic flow, into 4 spatial positions (Fig. 5A). TSDN

receptive fields tend to be located slightly offset from the visual midline, with preferred

direction of motion away from the midline (Fig. 5A). We defined the lateral parts of the

display as either ipsilateral or contralateral based on the preferred direction of each TSDN.

Data analysis and statistics

We recorded from 34 TSDNs in 34 male hoverflies. We kept data from all TSDNs that

showed a robust response to a target moving over a white background (Fig. 1B, Movie 1, 2).

We repeated this control throughout the recording, and only kept data from neurons that

responded consistently. We only kept data from experiments with a minimum 9 repetitions.

The data from repetitions within a neuron were averaged, and shown as spike histograms

(mean ± sem) with 1 ms resolution, after smoothing with a 20 ms square-wave filter. For

quantification across neurons in most cases, we calculated the mean spike rate for each

neuron from the spike histogram for the duration of target motion, after excluding the first

and last 40 ms (dotted boxes, Fig. S1B-E). We normalized the responses to each neuron's

own mean response to a target moving over a white background.

For the model experiment (Fig. 3), we calculated the mean spike rate across 0.48 s from the

spike histogram when the target traversed each neuron's receptive field. We normalized

the data from each neuron to the sum of the responses to the three stimuli (ON, OFF,

target).

Data analysis was performed in Matlab and statistical analysis in Prism 7.0c for Mac OS X

(GraphPad Software, San Diego, USA). Throughout the paper n refers to the number of

repetitions within one neuron, and N to the number of neurons. The sample size, type of

test and P value is indicated in each figure legend. We performed paired t-tests, one-way

ANOVAs, followed by Dunnett's post hoc test for multiple comparisons, or two-way RM

ANOVAs, followed by Bonferroni's post hoc test for multiple comparisons. All data have

been deposited to DataDryad.

Acknowledgements

We thank lab members for constructive feedback, and the Botanic Gardens of Adelaide for

their ongoing support. This research was funded by the US Air Force Office of Scientific

Research (AFOSR, FA9550-19-1-0294), the Australian Research Council (ARC, DP170100008,

DP180100144 and FT180100289), and the Flinders Foundation.

References

1. P. T. Gonzalez-Bellido, S. T. Fabian, K. Nordström, Target detection in insects: optical, neural and behavioral optimizations. *Curr Opin Neurobiol* **41**, 122-128 (2016).

2. W. Wellington, S. Fitzpatrick, Territoriality in the drone fly, *Eristalis tenax* (Diptera, Syrphidae). *Can Entomol* **113**, 695-704 (1981).

3. J. Zeil, Sexual dimorphism in the visual system of flies: The free flight behavior of male Bibionidae (Diptera). *J Comp Physiol A* **150**, 395-412 (1983).

4. S. T. Fabian, M. E. Sumner, T. J. Wardill, S. Rossoni, P. T. Gonzalez-Bellido, Interception by two predatory fly species is explained by a proportional navigation feedback controller. *Journal of The Royal Society Interface* **15** (2018).

5. R. Schröder, C. N. Linkem, J. A. Rivera, M. A. Butler, Should I stay or should I go? Perching damselfly use simple colour and size cues to trigger flight. *Animal Behaviour* **145**, 29-37 (2018).

6. C. T. O'Rourke, T. Pitlik, M. Hoover, E. Fernández-Juricic, Hawk eyes II: diurnal raptors differ in head movement strategies when scanning from perches. *PLoS One* **5**, e12169 (2010).

7. S. M. Fitzpatrick, W. G. Wellington, Contrasts in the territorial behavior of three species of hover flies (Diptera: Syrphidae). *Can Entomol* **115**, 559-566 (1983).

8. E. L. Corvidae, R. O. Bierregaard, S. E. Peters, Comparison of wing morphology in three birds of prey: correlations with differences in flight behavior. *J Morphol* **267**, 612-622 (2006).

9. J. J. Koenderink, Optic flow. *Vision Res* **26**, 161-179 (1986).

10. M. F. Land, Visual acuity in insects. *Annu Rev Entomol* **42**, 147-177 (1997).

11.   D. O'Carroll, Feature-detecting neurons in dragonflies. *Nature* **362**, 541-543 (1993).

12.   K. Nordström, P. D. Barnett, D. C. O'Carroll, Insect detection of small targets moving

428   in visual clutter. *PLoS Biol* **4**, 378-386 (2006).

429   13.   S. D. Wiederman, D. C. O'Carroll, Discrimination of features in natural scenes by a

430   dragonfly neuron. *J Neurosci* **31**, 7141-7144 (2011).

431   14.   K. Nordström, Neural specializations for small target detection in insects. *Curr Opin*

432   *Neurobiol* **22**, 272-278 (2012).

433   15.   S. Nicholas, J. Supple, R. Leibbrandt, P. T. Gonzalez-Bellido, K. Nordström, Integration

434   of small- and wide-field visual features in Target-Selective Descending Neurons of

435   both predatory and non-predatory dipterans. *J Neurosci* **38**, 10725-10733 (2018).

436   16.   S. Namiki, M. H. Dickinson, A. M. Wong, W. Korff, G. M. Card, The functional

437   organization of descending sensory-motor pathways in *Drosophila*. *eLife* **7**, e34272

438   (2018).

439   17.   C. L. Chen *et al.*, Imaging neural activity in the ventral nerve cord of behaving adult

440   *Drosophila*. *Nat Commun* **9**, 4390 (2018).

441   18.   P. D. Barnett, K. Nordström, D. C. O'Carroll, Retinotopic organization of small-field-

442   target-detecting neurons in the insect visual system. *Curr Biol* **17**, 569-578 (2007).

443   19.   T. Wardill *et al.*, A novel interception strategy in a miniature robber fly with extreme

444   visual acuity. *Curr Biol* **27**, 854-859 (2017).

445   20.   R. M. Olberg, A. H. Worthington, J. L. Fox, C. E. Bessette, M. P. Loosemore, Prey size

446   selection and distance estimation in foraging adult dragonflies. *J Comp Physiol A* **191**,

447   791-797 (2005).

448 21. O. Dyakova, Y.-J. Lee, K. D. Longden, V. G. Kiselev, K. Nordström, A higher order

449   visual neuron tuned to the spatial amplitude spectra of natural scenes. *Nat Commun*

450   **6**, 8522 (2015).

451 22. S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, A model for the detection of

452   moving targets in visual clutter inspired by insect physiology. *PLoS ONE* **3**, e2784

453   (2008).

454 23. S. Nicholas, R. Leibbrandt, K. Nordström, Visual motion sensitivity in descending

455   neurons in the hoverfly. *J Comp Physiol A* **206**, 149-163 (2020).

456 24. C. Städele, M. F. Keleş, J. M. Mongeau, M. A. Frye, Non-canonical receptive field

457   properties and neuromodulation of Feature-Detecting neurons in flies. *Curr Biol*

458   10.1016/j.cub.2020.04.069 (2020).

459 25. K. Nordström, D. C. O'Carroll, Feature detection and the hypercomplex property in

460   insects. *Trends Neurosci* **32**, 383-391 (2009).

461 26. M. F. Keleş, B. J. Hardcastle, C. Städele, Q. Xiao, M. A. Frye, Inhibitory interactions

462   and columnar inputs to an object motion detector in *Drosophila*. *Cell Rep* **30**, 2115-

463   2124.e2115 (2020).

464 27. M. F. Keleş, M. A. Frye, Object-detecting neurons in *Drosophila*. *Curr Biol* **27**, 680-687

465   (2017).

466 28. R. Tanaka, D. A. Clark, Object-displacement-sensitive visual neurons drive freezing in

467   *Drosophila*. *Curr Biol* 10.1016/j.cub.2020.04.068 (2020).

468 29. M. Egelhaaf, On the neuronal basis of figure-ground discrimination by relative

469   motion in the visual system of the fly. II. Figure-detection cells, a new class of visual

470   interneurones. *Biol Cybern* **52**, 195-209 (1985).

30.   A. K. Warzecha, M. Egelhaaf, A. Borst, Neural circuit tuning fly visual interneurons to

motion of small objects. I. Dissection of the circuit by pharmacological and

photoinactivation techniques. *J Neurophysiol* **69**, 329-339 (1993).

31.   W. Reichardt, T. Poggio, Figure-ground discrimination by relative movement in the

visual-system of the fly. Part I: Experimental results. *Biol Cybern* **35**, 81-100 (1979).

32.   T. Collett, Visual neurones in the anterior optic tract of the privet hawk moth. *J Comp*

Physiol **78**, 396-433 (1972).

33.   M. Egelhaaf, On the neuronal basis of figure-ground discrimination by relative

motion in the visual system of the fly. III. Possible input circuitries and behavioral

significance of the FD-cells. *Biol Cybern* **52**, 267-280 (1985).

34.   S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, Correlation between OFF and ON

Channels Underlies Dark Target Selectivity in an Insect Visual System. *J Neurosci* **33**,

13225-13232 (2013).

35.   S. D. Wiederman, J. M. Fabian, J. R. Dunbier, D. C. O'Carroll, A predictive focus of gain

modulation encodes target trajectories in insect vision. *eLife* **6** (2017).

36.   K. Nordström, D. M. Bolzon, D. C. O'Carroll, Spatial facilitation by a high-performance

dragonfly target-detecting neuron. *Biol Lett* **7**, 588-592 (2011).

37.   M. O. Franz, H. G. Krapp, Wide-field, motion-sensitive neurons and matched filters

for optic flow fields. *Biol Cybern* **83**, 185-197 (2000).

38.   A. K. Warzecha, A. Borst, M. Egelhaaf, Photo-ablation of single neurons in the fly

visual system reveals neural circuit for the detection of small moving objects.

Neurosci Lett **141**, 119-122 (1992).

39.   J. W. Aptekar, M. F. Keles, P. M. Lu, N. M. Zolotova, M. A. Frye, Neurons forming

optic glomeruli compute figure-ground discriminations in *Drosophila*. *J Neurosci* **35**,

7587-7599 (2015).

40.   M. Wu *et al.*, Visual projection neurons in the *Drosophila* lobula link feature

detection to distinct behavioral programs. *eLife* **5**, e21022 (2016).

41.   B. R. H. Geurten, K. Nordström, J. D. H. Sprayberry, D. M. Bolzon, D. C. O'Carroll,

Neural mechanisms underlying target detection in a dragonfly centrifugal neuron. *J*

Exp Biol **210**, 3277-3284 (2007).

42.   T. S. Collett, M. F. Land, How hoverflies compute interception courses. *J Comp*

Physiol A **125**, 191-204 (1978).

43.   S. Nicholas, M. Thyselius, M. Holden, K. Nordström, Rearing and long-term

maintenance of *Eristalis tenax* hoverflies for research studies. *JoVE*

doi:10.3791/57711, e57711 (2018).

44.   D. H. Brainard, The Psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

45.   D. G. Pelli, The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision* **10**, 437-442 (1997).

Figure Legends

Figure 1. The TSDN response to target motion is affected by horizontal optic flow

A) Pictograms of the round, black target, with a diameter of 3°, traversing a white

background (left), or stationary optic flow (right), at 130°/s. **B)** The mean spiking response of

different TSDNs was significantly reduced when the target moved across stationary optic

flow compared with a white background (N=34; ***P < 0.001, two-tailed paired t-test). **C)**

The TSDN response to a target traversing stationary optic flow (left, grey), sideslip or yaw in

either the same or the opposite direction to the target (red pictograms). All histograms from

a single TSDN (mean ± sem, n=18) shown with 1 ms resolution after smoothing with a 20 ms

square-wave filter. **D)** The mean response across TSDNs (N=12) to a target traversing

stationary optic flow (left, grey), sideslip or yaw, in the same direction ("+50") or opposite

direction ("-50") to the target, after normalizing the data to each neuron's own response to

a target traversing a white background. Sideslip was simulated at 50 cm/s and yaw at 50°/s.

The line shows the mean and the filled data points correspond to the neuron in panel C. In

panel D significance is shown with **** for P < 0.0001 following a one-way ANOVA followed

by Dunnett's multiple comparison's test done together with the data shown in Fig. S2B.

Figure 2. Interactions between optic flow and target direction

A) The data show TSDN responses to targets moving either horizontally or vertically across

sideslip or lift, in the same direction as the target, or perpendicular to the target, as

indicated by the pictograms. N=8. Two-way ANOVA showed P=0.2959 for target direction,

P=0.7639 for optic flow direction and P=0.0002 for the interaction between target and optic

flow. **B)** TSDN responses to targets moving horizontally or vertically across sideslip or lift, in

the opposite direction to the target, or perpendicular to the target, as indicated by the

pictograms. Two-way ANOVA showed P=0.0092 for target direction, P=0.0228 for optic flow

direction and P=0.0010 for the interaction between target and optic flow. Same N=8 in

panels A and B.

Figure 3. Elementary STMD input to TSDNs

Responses to a leading OFF edge, trailing ON edge, or a complete black target, with a side of

3°, traversing a white background at 130°/s. The black data show the predicted output from

a motion detector that compares luminance changes from at least 2 points in space. Data

replotted from (34), after normalizing to its own sum. The grey data show the predicted

output from an elementary STMD (ESTMD), which compares luminance changes from one

point in space. Data replotted from (34), after normalizing to its own sum. The white data

show the TSDN response to the same three stimuli (N=6) after normalizing the data from

each neuron to its own sum.

Figure 4. The TSDN inhibition by syn-directional sideslip can be primed

A) Raw data trace from one TSDN to a target moving across sideslip moving in the same

direction as the target preceding target motion, then being stationary concurrent with

target motion. Timing is color coded: green, preceding optic flow; red, concurrent optic

flow; blue, target motion. **B)** TSDN target responses are significantly inhibited by preceding

sideslip in the same direction as the target, compared with preceding stationary optic flow.

C) There was no significant difference between TSDN responses to targets moving across

syn-directional sideslip, when the sideslip moved preceding target motion compared with

preceding stationary optic flow (two-way ANOVA for the data in panels B and C, followed by

Post hoc Bonferroni's test: ***P=0.0004. **D)** There was no significant difference between

TSDN responses to targets moving across stationary optic flow, whether this moved in the

opposite direction or not preceding target motion. The grey data is the same as in panel B.

E) There was no significant difference between TSDN responses to targets on sideslip in the

opposite direction to target motion, whether the optic flow moved or not preceding target

motion (two-way ANOVA for data in panels D and E, ns). Data from the same N=7 in panels

B-E, with the lines showing the mean.

Figure 5. Frontal optic flow is essential

A) A pictogram of the separation of the optic flow into four distinct positions: ipsilateral (IL),

dorsal (D), ventral (V) and contralateral (CL). The color coding shows the receptive field of

an example TSDN, and the arrows the local motion sensitivity. **B)** TSDN responses to targets

moving across syn-directional sideslip are significantly inhibited compared to stationary

control (grey) if the sideslip covers the full, dorsal or ventral screen. **C)** TSDN responses to

targets moving across opposite direction sideslip are significantly facilitated compared to

stationary control (grey, same data as in panel B) if the sideslip covers the full, dorsal or

ventral visual field. Panels B and C show data from the same neurons (N=10, one-way

ANOVA followed by Dunnett's multiple comparisons test, with *P<0.05, **P<0.01).

Supplementary Information

Figure S1. The TSDN response to target motion and optic flow

A) Raw data trace from an extracellular TSDN recording. Timing of stimulus presentation

indicated by colored bars (blue, target, and red, optic flow). **B)** Magnification of the raw data

traces shown in panel A. **C)** Spike raster of the same neuron in response to 18 repeated

trials of a target on a white background (left) or on stationary optic flow (right). **D)** To

determine the TSDN response to the optic flow we first quantified each neuron's

spontaneous rate, i.e. the spikes generated when viewing a white background, and used this

as a threshold. For each neuron we then quantified the number of trials in which the

response to the optic flow was above this threshold. This was converted to a percentage of

responses. The data here show this percentage of responses across 12 TSDNs as mean ±

sem. **E)** In each neuron, we quantified the mean response for those trials that were

classified as responders (in panel D). This mean response was normalized to each neuron's

mean response to a target traversing a white background, as throughout the rest of the

paper. Data are shown as mean ± sem (N=12).

Figure S2. The TSDN response to target motion is affected by 3D optic flow

A) The response from one TSDN to a target traversing stationary optic flow (left, grey,

replotted from Fig. 1C), or different types of optic flow (see red pictograms). All histograms

shown as mean ± sem (n=18) at 1 ms resolution, after smoothing with a 20 ms square-wave

filter. **B)** The mean response across TSDNs (N=12, same neurons as Fig. 1D) to a target

traversing stationary optic flow (left, grey, replotted from Fig. 1D), or different types of optic

flow. Stars indicate significance, one-way ANOVA followed by Dunnett's multiple

comparisons test (**P<0.01, ***P<0.001 and ****P<0.0001). Note that the one-way ANOVA

was done together with the data shown in Fig. 1D. Translations were simulated at 50 cm/s

and rotations at 50°/s. The lines show the mean and the filled data points correspond to the

neuron in panel A.

Figure S3. The TSDN response to vertical target motion

A) The TSDN response to a target traversing a white background horizontally (left, grey) or

vertically (right, black). Histograms shown as mean ± sem (n=18) at 1 ms resolution, after

smoothing with a 20 ms square-wave filter. **B)** The mean spiking response of 8 TSDNs to

targets moving horizontally (grey, left) or vertically (black, right) over a white background.

Movie 1. TSDN response to a target over a white background

The movie depicts the stimulus as displayed on the screen, with the hoverfly positioned

centered and perpendicular to the screen. The hoverfly was positioned ventral side up, but

we have rotated the movie for display purposes so the dorsal side is up. The colored lines

show the outline of the receptive field, mapped as described previously (23). The red data at

the bottom of the movie show the response of an example TSDN.

Movie 2. TSDN response to a target over a white background

Same as Movie 1, but slowed down 10 times.

Movie 3. TSDN response to a target over stationary optic flow

The movie depicts the stimulus as displayed on the screen, with the colored lines showing

the outline of the receptive field. The red data at the bottom of the movie show the

response of an example TSDN. In this case the optic flow was stationary, and appeared 0.5 s

before the target.

Movie 4. TSDN response to a target over stationary optic flow

Same as Movie 3, but slowed down 10 times.

Movie 5. TSDN response to a target over syn-directional sideslip optic flow

The movie depicts the response of the same example TSDN to the same target motion as in

previous movies, but now traversing sideslip optic flow appearing 0.5 s before the target.

The hoverfly was positioned at a distance of 6.5 cm. At this viewing distance, the resulting

optic flow simulates the type of optic flow that would be generated by the hoverfly side-

slipping through the world at 50 cm/s. The simulated optic flow consisted of ca 6400

spheres, of which roughly 1200 are projected onto the screen at any one time.

Movie 6. TSDN response to a target over syn-directional sideslip optic flow

Same as Movie 5, but slowed down 10 times. Note that many of the optic flow "targets" of

the correct size move through the receptive field (colored contour lines).

Movie 7. TSDN response to a target over opposite direction sideslip optic flow

The movie depicts the response of the same example TSDN to the same target motion as in

previous movies, but now traversing sideslip optic flow in the opposite direction to the

target.

Movie 8. TSDN response to a target over opposite direction sideslip optic flow

Same as Movie 7, but slowed down 10 times.

A

White background | Stationary

B

Mean response (spikes/s)

White Background | Stationary

Target

***

C

100 spikes/s

100 ms

D

Normalized response

Target

Optic flow

Stationary | Sideslip | Yaw

+50 | -50 | +50 | -50

****

A

Target

100 spikes/s

100 ms

B

Normalized response

Optic flow

Stationary

Lift   + 50   − 50

Pitch   + 50   − 50

Thrust   + 50   − 50

Roll   + 50   − 50

**** *** **** **** **** **** ** ****

A

100 spikes/s

100 ms

B

Mean response (spikes/s)

Target