1

Facilitation of neural responses to targets moving against optic flow

Sarah Nicholas[1] and Karin Nordström[1,2]*

[1]Flinders Health and Medical Research Institute, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia

[2]Department of Neuroscience, Uppsala University, Box 593, 751 24 Uppsala, Sweden

*email: karin.nordstrom@flinders.edu.au

ORCIDs

Sarah Nicholas: 0000-0002-5555-9421

Karin Nordström: 0000-0002-6020-6348

Classification

Biological sciences: Neuroscience

Keywords

Target motion, insect vision, motion vision, background motion, hoverfly, TSDN, widefield motion, response suppression, neural facilitation, sensori-motor

Abstract

For the human observer, it can be difficult to follow the motion of small objects, especially

when they move against background clutter. In contrast, insects efficiently do this, as

evidenced by their ability to capture prey, pursue conspecifics, or defend territories, even in

highly textured surrounds. We here recorded from target selective descending neurons

(TSDNs) which likely subserve these impressive behaviors. To simulate the type of

background optic flow that would be generated by the pursuer's own movements through

the world, we used the coherent motion of a perspective distorted sparse dot field. We

show that hoverfly TSDN responses to target motion are suppressed when such background

optic flow moves in the same direction as the target. Indeed, the neural responses are

strongly attenuated against both translational sideslip as well as rotational yaw. More

strikingly, we show that TSDNs are facilitated by background optic flow in the opposite

direction to the target, if the target moves horizontally. Furthermore, we show that a small,

frontal spatial window of background optic flow is enough to fully facilitate or attenuate

TSDN responses to target motion. We argue that the TSDN response facilitation could be

beneficial in modulating corrective turns during target pursuit.

Significance statement

Target detection in visual clutter is a difficult computational task that insects, with their poor resolution compound eyes and small brains, do successfully and with extremely short behavioral delays. We here show that the responses of target selective descending neurons are attenuated by background motion in the same direction as target motion, but facilitated by opposite direction background motion. This finding is important for understanding conspecific pursuit behavior, since these descending neurons likely control behavioral output. The facilitation that we describe would come into effect if the hoverfly is subjected to background motion in one direction, but the target it is pursuing moves in the opposite direction, and could therefore be used to modulate gaze stabilizing corrective turns.

/body

Introduction

The survival of many animals often depends on their ability to visually detect small moving

objects or targets, as these could represent predators, prey, mates or territorial intruders

(1). Efficient target detection is a computationally challenging task, which becomes even

more difficult when done against visual clutter. Despite this, many insects successfully

detect targets, followed by highly acrobatic pursuits, often in visually complex

environments. For example, male *Eristalis tenax* hoverflies establish territories in foliage rich

areas, on alert for intruders or potential mates and ready to engage in high-speed pursuit

(2).

Initial target detection can be facilitated by behaviors that render the background

stationary, thus making the target the only thing that moves. Many insects and vertebrates

indeed visualize targets against the bright sky (3), or from a stationary stance, such as

perching (4-6) or hovering (7, 8). However, as soon as the pursuer moves, its own

movement creates self-generated widefield motion across the retina, often referred to as

optic flow (9), or background motion (10). In addition to self-generated background optic

flow, when a pursuer is subjected to involuntary deviations away from their flight path, for

example by a gust of wind, this also generates background optic flow. Quickly correcting

such unplanned course deviations is essential for successfully navigating through the world.

For example, a flying insect being pushed sideways to the right by a gust of wind, will

experience optic flow in the opposite direction. To correct for this, in response to leftward

optic flow, the insect will use its wings to perform a corrective optomotor response to the

left (11), and/or stabilize its gaze by moving its head (12). Recent evidence, suggests that

efference copies suppress the visual neurons sensitive to optic flow, as the optomotor

response could otherwise counteract voluntary turns (13).

During pursuit, the pursuer is thus subjected to visual motion that could originate from the

independent motion of objects within the environment, from its own intentional

movements, or from movement imposed by external forces, such as a gust of wind.

Importantly, how the insect moves also affects the type of optic flow it will experience. For

example, during translations distant features move slower than closer ones, whereas during

rotations, all features move at the same angular velocity irrespective of distance from the

observer (14). Many insects use the information such relative motion provides to estimate

the distance to features in the surround. *Eristalis tenax* hoverflies, for example, make both

small translational head movements when perched and larger side-to-side movements

during flight (15, 16). Moreover, many flying insects, including *Eristalis,* show behavioral

segregation between rotational and translational movements, through a saccadic flight and

gaze strategy (16-18). How this may influence target detection is currently not known.

The ability of insects to successfully pursue targets in clutter (4, 19) is thus remarkable, and

suggests a high level of optimization, making the underlying neural mechanisms interesting

to study. Indeed, insects that efficiently pursue targets, including predatory dragonflies and

robberflies, as well as territorial hoverflies, have higher-order neurons that are sharply

tuned to the motion of small, dark targets, identified in the optic lobes (20, 21) and the

descending nerve cord (22, 23). Target tuned neurons often have receptive fields (24-26) in

the part of the compound eye that has the best optics (27, 28). Target selective descending

neurons (TSDNs) project to the thoracic ganglia (25, 29) where wing and head movements

are controlled (30), and electrically stimulating dragonfly TSDNs leads to wing movements

(31). Taken together, this suggests that TSDNs subserve target pursuit. However, how TSDNs

respond to targets moving against translational and rotational optic flow is unknown.

We quantified the responses of *Eristalis* TSDNs to targets moving against 6 types of optic

flow, three translations and three rotations. The background optic flow consisted of the

coherent motion of thousands of "targets", so that the target could only be distinguished by

its relative motion. We found that optic flow in the same direction as the target attenuated

the TSDN response to the target, whether the optic flow moved in a rotational or

translational manner. We found that perpendicular optic flow attenuated the TSDN target

response, but to a lesser degree than syn-directional background motion. This suggests that

the vector divergence between the target and the background is important. Most strikingly,

we found that optic flow in the opposite direction to target motion increased the TSDN

response, if the target moved horizontally. Such neural facilitation is a novel observation.

We found that a small spatial extent of frontal motion was sufficient to elicit both TSDN

attenuation and facilitation, whether spatially overlapping with target motion or not. As

descending neurons control behavioral output (30, 31), the response attenuation and

facilitation could play a role in modulating optomotor or gaze stabilizing corrective turns as

needed during target pursuit.

Results

TSDNs respond to target motion but not to background optic flow

We simulated background optic flow using the coherent motion of a sparse dot field (24). By

projecting the individual features in the sparse dot field onto a screen anterior of the

hoverfly, their spatial location, and simulated z-depth, over time, provided the type of optic

flow that would be generated by self-motion (24). In optic flow sensitive descending

neurons, this stimulus elicits strong direction-selective responses, similar to the responses

to widefield sinusoidal gratings (24), or large moving images (22) with naturalistic statistics

(32). We confirmed that our optic flow stimulus did not generate a response in target

selective descending neurons (TSDNs) in male *Eristalis tenax* hoverflies (Fig. S1). For

example, translational sideslip to the left only generated a response in 18% of 222

repetitions across 12 TSDNs (Fig. S1A). Furthermore, when leftward sideslip did generate a

TSDN response, this was much less than the response to a target traversing a white

background (Fig. S1B). This lack of TSDN response confirms that the optic flow stimulus is

qualitatively similar to other types of widefield background motion.

In contrast to the lack of response to the optic flow, TSDNs responded strongly to the

motion of a small, dark target (22, 24) traversing a white background (compare spiking

response during stimulation with the lack of activity before stimulation, Fig. 1, left, Movie 1,

2), as well as when the target traversed the stationary sparse dot field (Fig. 1, right, Movie 3,

4). Note, that since the sparse dot field consisted of hundreds of "targets", there were no

spatial characteristics separating the individual target from the optic flow when both were

stationary (Fig. 1A, right). For quantification across neurons, we calculated the mean spike

frequency for the duration of target motion, except for the first and last 40 ms of each 500

ms target trajectory (dotted box, Fig. 1B, C). We found a small, but significant response

reduction to targets moving across a stationary dot field compared with a white background

(Fig. 1D). As the response across neurons was variable (N=30, coefficient of variation 63%

and 67%, respectively, Fig. 1D), we normalized the response from each neuron to its own

mean response to a target moving across a white background.

Background optic flow modulates the TSDN target response

To investigate the effect the background optic flow had on TSDN responses to target motion

we simulated different types of translations (Fig. 2A) at 50 cm/s and rotations (Fig. 2B) at 50

deg/s. During translations distant features move slower than closer ones, whereas during

rotations, all features move at the same angular velocity irrespective of distance from the

observer (14). We first simulated translational sideslip and found that when the target

moved horizontally across syn-directional sideslip, the TSDN response was strongly

attenuated (Fig. 2C, D, "Left Sideslip", Movie 5, 6), compared with control where the target

moved over a stationary dot field (grey, Fig. 2C, D, Movie 3, 4). Similar effects of syn-

directional background motion were previously seen when using panoramic images with

naturalistic statistics (22).

To investigate the effect the relative motion of individual features within the optic flow had

on TSDN responses to target motion, we next simulated rotational yaw at 50 deg/s. We

found that yaw optic flow in the same direction as the target also strongly attenuated the

TSDN response (Fig. 2C, D, "Left Yaw"). This suggests that rotational and translational optic

flow have the same effect on TSDN responses to target motion, and that the relative motion

of features within the optic flow is less important.

We next investigated the effect opposite-direction background optic flow had on TSDN

responses to target motion. We found that when the target moved against counter-

directional sideslip the TSDN response was strongly enhanced (Movie 7, 8, Fig. 2C, D, "Right

Sideslip", mean increase 71.1%). Similarly, yaw optic flow in the opposite direction to the

target also facilitated the TSDN response, by 84.9% (Fig. 2C, D, "Right Yaw"). Such response

facilitation was not seen when displaying targets over counter-directional background

motion using panoramic images, in either TSDNs (22) or other target tuned neurons in the

fly optic lobes (21, 26, 33).

Frontal optic flow is required and sufficient

Our data show that yaw and sideslip have similar effects on TSDN responses to target

motion (Fig. 2). As both sideslip and yaw contain substantial local motion in the frontal

visual field (34), we next asked if frontal optic flow is required. We investigated this by

limiting the spatial extent of the sideslip to either cover the ipsilateral, dorsal, ventral or

contralateral position on the screen (Fig. 3A). Note that only the dorsal position covers the

TSDN receptive field (Fig. 3A). In the other three positions, the sideslip optic flow was

spatially separated from the receptive field (Fig. 3A), and thus the target trajectory.

We found that when sideslip moved in the same direction as the target, the TSDN response

was attenuated if the optic flow covered the full, dorsal or ventral position on the screen

(Fig. 3B), compared with the stationary control (grey, Fig. 3B). However, when the optic flow

only covered the ipsilateral or contralateral positions, there was no difference compared

with control (Fig. 3B), suggesting that frontal optic flow is both required and sufficient. Our

results are consistent with previous work using moving images with naturalistic statistics,

which did not have to spatially overlap with the target trajectory, nor have a large spatial

extent, to attenuate TSDN responses (22).

We next found that when sideslip moved in the opposite direction to the target, the TSDN

response to target motion was facilitated if the optic flow covered the full, dorsal or ventral

positions of the screen (Fig. 3C). When the sideslip was limited to the ipsilateral or

contralateral positions, there was no TSDN response facilitation (Fig. 3C). This suggests that

the optic flow does not have to spatially overlap with the target trajectory. However, there

has to be frontal, opposite direction optic flow for facilitation to take place.

In summary, our results show that a small spatial window of optic flow in either the dorsal

or ventral visual field is enough to strongly attenuate (Fig. 3B) or facilitate (Fig. 3C) the TSDN

response to target motion.

Vector divergence between target and background optic flow affects TSDNs response

Our data above show that syn-directional optic flow strongly attenuates TSDN responses to

target motion, whereas counter-directional optic flow facilitates the response (Fig. 2, 3).

This suggests that the level of vector divergence between the target and the background

influences the TSDN responses, so that maximum attenuation is generated at minimum

vector divergence, whereas maximum facilitation is generated at maximum divergence. To

explore this in further detail, we quantified the effect other types of optic flow had on TSDN

responses to target motion.

We found that the TSDN target response was suppressed by lift, which moves perpendicular

to the target (mean 44.0% response for downwards lift, "Lift Down"; mean 34.3% for

upwards lift, "Lift Up", Fig. 4). When the target was displayed against pitch, which also

provides perpendicular motion in the frontal visual field, the response was suppressed to

similar levels (57.1%, "Pitch Down"; 54.9%, "Pitch Up", Fig. 4). The TSDN response to target

motion was also attenuated when the target was displayed against thrust, which provides

perpendicular motion along the animal's anterio-posterior axis (44.3%, "Thrust Towards";

58.4%, "Thrust Away", Fig. 4). Thus, perpendicular background optic flow attenuates the

TSDN response (Fig. 4), but not as much as syn-directional motion (Fig. 2).

We next looked at roll optic flow, which creates no local motion straight ahead of the fly,

but opposite direction local motion in the dorsal and ventral visual fields. For example,

during counter-clockwise roll ("Roll CCW", Fig. 4) the target and the background optic flow

move in the same direction through the dorsal receptive field, but in opposite directions in

the ventral visual field. Therefore, the TSDN would receive a combination of maximum and

minimum vector divergence signals from the "dorsal" and "ventral" parts (as shown in Fig.

3). We found that the response attenuation was stronger against counter-clockwise roll

(75.3%, "Roll CCW", Fig. 4), than against clockwise roll (31.9%, "Roll CW", Fig. 4).

Thus, our data (Fig. 2, 4) support the notion that the level of vector divergence influences

the TSDN response. To explore this in further detail, we recorded from TSDNs that respond

robustly both to horizontal (grey, Fig. 5) and vertical target motion (black, Fig. 5). We found

that when the target moved horizontally, the TSDN response was strongly attenuated

against syn-directional sideslip (2nd column, Fig. 5C), less attenuated against lift in either

direction (last two columns, Fig. 5C), and strongly facilitated when displayed against

counter-directional sideslip (3rd column, Fig. 5C), consistent with previous results (Fig. 2, 4).

In the same TSDN neurons, we next moved the target vertically. We found that the TSDN

response was completely attenuated against syn-directional lift (last column, Fig. 5D), and

less attenuated against sideslip in either direction (2nd and 3rd column, Fig. 5D), which moves

perpendicular to the target. This supports the suggestion that maximum TSDN response

attenuation is generated at minimum vector divergence between the target and

background. However, when the vertical target was displayed against counter-directional

lift (4th column, Fig. 5D), there was no response facilitation. This suggests that maximum

vector divergence on its own is not enough to explain the TSDN response facilitation to

target motion.

Responses to ON-OFF edges consistent with 1-point correlation

TSDNs have been proposed to get their input from STMDs (35). As opposed to our TSDN

data, hoverfly STMD responses are rarely suppressed by syn-directional background motion,

and not facilitated by counter-directional background motion (21). As there are many other

target tuned neurons in the fly optic lobes (e.g. 33, 36-39) it is possible that TSDNs do not

get their input from STMDs. We can investigate the potential input using the underlying

target tuning mechanisms, which can be distilled down into three fundamentally different

concepts. For example, visual neurons can become target tuned by receiving inhibitory

feedback from the widefield system (39-41), or by using center-surround antagonism

together with rapid adaptation (38, 42). Alternatively, they can use an elementary STMD

model, which is tuned to the unique spatio-temporal profile of a moving target, with a dark

contrast change (OFF) from the leading edge followed by a bright contrast change (ON) by

the trailing edge. Importantly, while the first two mechanisms rely on comparisons from

neighboring points in space, the elementary STMD compares input from one point on space

(38, 43, 44). Therefore, the first two models will respond similarly to the motion of a target,

to the motion of a leading OFF edge, and the motion of a trailing ON edge (black, Fig. 6,

redrawn from (38, 45)). In contrast, the elementary STMD model only responds strongly to

the target (grey, Fig. 6, redrawn from (45)).

Our results show that TSDNs do not respond well to a leading OFF edge, or to a trailing ON

edge (white, Fig. 6). However, a complete target, where the leading edge is rapidly followed

by a trailing edge, gives a robust TSDN response (white, Fig. 6). Indeed, the physiological

responses (white, Fig. 6) match the elementary STMD model output (grey, Fig. 6). Since

STMD physiology also matches the elementary STMD model output (45, 46), this suggests

that TSDNs receive input from STMDs.

Discussion

We found that optic flow moving in the same direction as a target almost completely

attenuated the TSDN target response (Fig. 2), even when the optic flow only covered a small

part of the frontal visual field (Fig. 3). More strikingly, we found that optic flow in the

opposite direction to target motion increased the TSDN response (Fig. 2), if the target

moved horizontally (Fig. 5). We also found that perpendicular optic flow attenuated the

TSDN target response (Fig. 3, 4), but less than syn-directional motion, suggesting that the

vector divergence between the target and the background is important for response

suppression. However, we found that vector divergence is not enough to explain the

response facilitation (Fig. 3C).

We show for the first time, that TSDN responses to target motion are facilitated by counter-

directional optic flow (Fig. 2). Such neural facilitation has not been seen in previous work in

TSDNs (22) or STMDs (21, 26, 46), or other target tuned neurons in the optic lobes (33, 36,

39). Instead, in our previous work TSDN responses were significantly reduced when a

background image with naturalistic statistics moved in the opposite direction to the target

(22). Since the background used previously drive optic flow sensitive descending neurons as

well as sinusoidal gratings do (22, 47), and as well as the sparse dot field used here (24), this

suggests that facilitation is not generated by widefield motion in general. Indeed, this

suggests that there is something fundamentally different about the optic flow used here,

and the backgrounds in previous work. One difference is that we here use perspective

distorted optic flow, where the individual features move faster the closer they are during

translations, whereas during rotations they all move at the same angular velocity

irrespective of distance from the viewer. However, we found that the facilitation was strong

whether the target was displayed over sideslip or yaw (Fig. 2), suggesting that the relative

motion of the features within the background optic flow is not important.

Another difference is that the background used here consisted of the coherent motion of

thousands of "targets", whereas previous TSDN work used a background image with

naturalistic statistics (22). We currently do not know how STMDs, which are likely pre-

synaptic to the TSDNs (Fig. 6, and see Ref. 35), respond to this stimulus. However, we do

know that STMDs generate their ability to detect targets in clutter by being sharply tuned to

the target's unique spatio-temporal profile (43), with a dark-contrast OFF edge quickly

followed by a bright-contrast ON edge (Fig. 6). Indeed, if naturalistic backgrounds contain

small, high-contrast features, these often generate STMD responses (46). However, despite

our optic flow consisting of "targets", we saw no consistent TSDN responses even to

preferred direction motion (e.g. "Sideslip Left", "Yaw Left", Fig. S1, Movie 5, 6). This

suggests that TSDNs do not filter contrasting targets embedded within a background, like

some STMDs do (21, 46).

It is thus difficult to determine whether the facilitation is driven by active facilitation onto

the TSDN, or if it is inherited from upstream processes. If it is generated by neural input, and

since the facilitation required frontal optic flow, but this did not have to spatially overlap

with the TSDN receptive field or the target trajectory (Fig. 3A, C), the facilitation would

come from a direction-selective neuron type with a frontal visual field. Since counter-

directional lift did not facilitate the TSDN response if the target moved vertically rather than

horizontally (Fig. 5C), this suggests that such a neuron would have to be most sensitive to

horizontal motion, and less to vertical motion. Since some STMDs might respond to features

within the optic flow used here (46), and furthermore, since there are some STMDs that

respond both to target motion and to features within background motion (21), these could

potentially play a role in this. Future investigation of STMD and TSDN responses are clearly

required to explore the underlying mechanisms.

In contrast to the facilitation, we found that the sparse dot field stimulus suppressed the

TSDN response to target motion when these moved in the same direction (Fig. 2). This

response suppression could be driven by active inhibition from widefield motion detectors,

as previously suggested (22), or inherited from upstream processes. We previously showed

that TSDN responses to target motion are suppressed by background motion consisting of

an image with naturalistic statistics, and that background stimuli that drive optic flow

sensitive neurons suppress TSDN responses to target motion, whether spatially overlapping

with the target trajectory or not (22). This suggests that TSDNs are suppressed by widefield

motion in general. However, note that since we found maximum TSDN response

suppression at minimum vector divergence, whereas perpendicular motion suppressed the

response less (Fig. 4, 5; and see ref. 22), the response suppression has to contain some type

of vector divergence comparison.

Nevertheless, our findings make behavioral sense. Prior to initiating target pursuit, male

Eristalis hoverflies predict the flight course required to successfully intercept the target,

based predominantly on the target's angular velocity (48). To successfully execute an

interception style flight course, the hoverfly turns in the direction that the target is moving

(48). In doing so, the hoverfly creates self-generated optic flow in the opposite direction to

the target's motion. In this case the TSDNs would be facilitated (Fig. 2), which could be

beneficial.

Once pursuit is initiated, if the pursuer drifts rightward due to e.g. a gust of wind, this

induces leftward optic flow across the frontal retina, which would evoke a leftward

optomotor response and/or gaze stabilizing turn. If the target is also moving leftward, in the

direction of the corrective maneuver, this maneuver would also pursue the target and no

TSDN signal would be required. Indeed, we found that the TSDNs were quiet under such

conditions ("Left", Fig. 2). However, if during the same pursuit the target instead moves

rightward, opposite to the direction of the pursuer's corrective maneuver, then the TSDNs

would be strongly facilitated ("Right", Fig. 2). Considering that TSDNs project to motor

command centers in the thoracic ganglia (25, 31), this TSDN facilitation could then

presumably override corrective turns. The TSDN response modulation by optic flow shown

here could thus be beneficial for controlling behavioral output.

We only saw a response facilitation if the target moved horizontally, and not vertically (Fig.

5). This facilitation difference could be affected by the fact that the TSDNs that we recorded

from responded better to horizontal motion than to vertical target motion (Fig. 5A). In

addition, since we do not know where the TSDNs project to we do not know if the TSDNs

control head movements, wing movements, or maybe both. Indeed, recent work suggests

that gaze stabilizing turns by the head, and the wing optomotor response, are controlled

independently, and by different visual components (12, 49, 50). Furthermore, the difference

could be explained by different behavioral requirements to vertical and horizontal target

deviations. For example, whereas horizontal deviations require the left and right wings to

move differently, to induce a turn, vertical deviations would likely require the left and right

wings to perform similar corrective behaviors. Finally, it is important to note that even if the

responses to vertical target motion were not facilitated, neither were they suppressed (4th

column, Fig. 5D). There would thus still be a strong neural signal projected to the thoracic

ganglia.

Materials and Methods

Electrophysiology

Eristalis tenax hoverflies were reared and maintained as previously described (51). For

electrophysiology, a male hoverfly was immobilized ventral side up using a beeswax and

resin mixture. A small hole was cut at the anterior end of the thorax to expose the cervical

connective, which was then raised slightly and supported using a small wire hook, for

insertion of a sharp polyimide-insulated tungsten microelectrode (2 MOhm, Microprobes,

Gaithersburg, USA). The animal was grounded via a silver wire inserted into the ventral part

of the hole.

Extracellular signals were amplified at 100x gain and filtered through a 10- to 3000-Hz

bandwidth filter on a DAM50 differential amplifier (World Precision Instruments, Sarasota,

USA), with 50 Hz noise removed with a HumBug (Quest Scientific, North Vancouver,

Canada). The data were digitized via Powerlab 4/30 (ADInstruments, Sydney, Australia) and

acquired at 40 kHz with LabChart 7 Pro software (ADInstruments). Single units were

discriminated by amplitude and half-width using Spike Histogram software (ADInstruments).

Visual stimuli

Eristalis males were placed ventral side up, centered and perpendicular to an Asus LCD

screen (Asus, Taipai, Taiwan) at 6.5 cm distance. The screen had a refresh rate of 165 Hz, a

linearized contrast with a mean illuminance of 200 Lux, and a spatial resolution of 2560 x

1440 pixels, giving a projected screen size of 155° x 138°. Visual stimuli were displayed using

custom written software based on the Psychophysics toolbox (52, 53) in Matlab

(Mathworks, Natick, USA).

TSDNs were identified as described (22, 24). In short, we mapped the receptive field of each neuron by scanning a target horizontally and vertically at 20 evenly spaced elevations and azimuths (24), to calculate the local motion sensitivity and local preferred direction. We then scanned targets of varying height through the small, dorso-frontal receptive fields (Fig. 3A) to confirm that each neuron was sharply size tuned with a peak response to targets subtending 3°- 6°, with no response to larger bars, to looming or to widefield stimuli (22, 24).

Unless otherwise mentioned, targets were black and round with a diameter of 15 pixels, moving at a velocity of 900 pixels/s for 0.48 s. When converted to angular values and taking the small frontal receptive fields of TSDNs into account, this corresponds to an average diameter of 3° and a velocity of 130°/s (22). Unless otherwise stated, each target travelled in each neuron's preferred horizontal direction (i.e. left or right) and across the center of its receptive field. Between repetitions, we varied the target elevation slightly, to minimize habituation (22). There was a minimum 4 s between stimulus presentations. Stimulus order was randomized.

Optic flow was generated as previously described (24). Briefly, the optic flow consisted of a simulated cube with 4 m sides, filled with 2 cm diameter spheres at a density of 100 per m^3, with the hoverfly placed in the center. The coherent motion of these ca 6400 spheres around the hoverfly was used to simulate self-generated optic flow. The ca 1200 spheres anterior to the hoverfly were projected onto the screen, with their size indicating the distance from the hoverfly. Circles closer than 6 cm were not displayed. Six types of optic flow were simulated: three translations at 50 cm/s (sideslip, lift and thrust) and three rotations at 50°/s (yaw, pitch and roll). Unless otherwise stated optic flow was displayed for 0.48 s prior to the target. Both target motion and optic flow disappeared simultaneously.

In most experiments the optic flow covered the entire visual display. In some experiments, we limited the spatial extent of the optic flow, into 4 spatial positions (Fig. 3A). TSDN receptive fields tend to be located slightly offset from the visual midline, with preferred direction of motion away from the midline (Fig. 3A). We defined the lateral parts of the display as either ipsilateral or contralateral based on the preferred direction of each TSDN.

 Data analysis and statistics

We recorded from 30 TSDNs in 30 male hoverflies. We kept data from all TSDNs that

showed a robust response to a target moving over a white background (Fig. 1, Movie 1, 2).

We repeated this control throughout the recording, and only kept data from neurons that

responded consistently. We only kept data from experiments with a minimum 9 repetitions.

The data from repetitions within a neuron were averaged, and shown as spike histograms

(mean ± sem) with 1 ms resolution, after smoothing with a 20 ms square-wave filter. For

quantification across neurons, we calculated the mean spike rate for each neuron from the

spike histogram for the duration of target motion, after excluding the first and last 40 ms of

each 0.48 s target trajectory (dotted boxes, Fig. 1B, C), unless otherwise indicated. We

normalized the responses to each neuron's own mean response to a target moving over a

white background.

Data analysis was performed in Matlab and statistical analysis in Prism 7.0c for Mac OS X

(GraphPad Software, San Diego, USA). Throughout the paper n refers to the number of

repetitions within one neuron, and N to the number of neurons. The sample size, type of

test (paired t-tests or one-way ANOVAs, followed by Dunnett's post hoc test for multiple

comparisons) and P value is indicated in each figure legend. All data have been deposited to

DataDryad.

Acknowledgements
416   We thank current and past lab members for constructive feedback, and the Botanic Gardens

417   of Adelaide for their ongoing support. Malin Thyselius provided the hoverfly pictogram in

418   Fig. 2. Our research was funded by the US Air Force Office of Scientific Research (AFOSR,

419   FA9550-19-1-0294), the Australian Research Council (ARC, DP170100008, DP180100144 and

420   FT180100289), and the Flinders Foundation.

References

1.   P. T. Gonzalez-Bellido, S. T. Fabian, K. Nordström, Target detection in insects: optical, neural and behavioral optimizations. *Curr Opin Neurobiol* **41**, 122-128 (2016).

2.   W. Wellington, S. Fitzpatrick, Territoriality in the drone fly, *Eristalis tenax* (Diptera, Syrphidae). *Can Entomol* **113**, 695-704 (1981).

3.   J. Zeil, Sexual dimorphism in the visual system of flies: The free flight behavior of male Bibionidae (Diptera). *J Comp Physiol A* **150**, 395-412 (1983).

4.   S. T. Fabian, M. E. Sumner, T. J. Wardill, S. Rossoni, P. T. Gonzalez-Bellido, Interception by two predatory fly species is explained by a proportional navigation feedback controller. *Journal of The Royal Society Interface* **15** (2018).

5.   R. Schröder, C. N. Linkem, J. A. Rivera, M. A. Butler, Should I stay or should I go? Perching damselfly use simple colour and size cues to trigger flight. *Animal Behaviour* **145**, 29-37 (2018).

6.   C. T. O'Rourke, T. Pitlik, M. Hoover, E. Fernández-Juricic, Hawk eyes II: diurnal raptors differ in head movement strategies when scanning from perches. *PLoS One* **5**, e12169 (2010).

7.   S. M. Fitzpatrick, W. G. Wellington, Contrasts in the territorial behavior of three species of hover flies (Diptera: Syrphidae). *Can Entomol* **115**, 559-566 (1983).

8.   E. L. Corvidae, R. O. Bierregaard, S. E. Peters, Comparison of wing morphology in three birds of prey: correlations with differences in flight behavior. *J Morphol* **267**, 612-622 (2006).

9.   J. J. Koenderink, Optic flow. *Vision Res* **26**, 161-179 (1986).

10.   W. Reichardt, M. Egelhaaf, A. Guo, Processing of figure and background motion in

the visual system of the fly. *Biol Cybern* **61**, 327-345 (1989).

11.   W. Reichardt, T. Poggio, Visual control of orientation behaviour in the fly. Part I. A

quantitative analysis. *Q Rev Biophys* **9**, 311-375 (1976).

12.   J. L. Fox, M. A. Frye, Figure-ground discrimination behavior in *Drosophila*. II. Visual

influences on head movement behavior. *J Exp Biol* **217**, 570-579 (2014).

13.   A. J. Kim, J. K. Fitzgerald, G. Maimon, Cellular evidence for efference copy in

Drosophila visuomotor processing. *Nat Neurosci* **18**, 1247-1255 (2015).

14.   M. Egelhaaf, R. Kern, J. P. Lindemann, Motion as a source of environmental

information: a fresh view on biological motion computation by insect brains. *Front*

Neural Circuits **8** (2014).

15.   W. G. Wellington, S. M. Fitzpatrick, Territoriality in the Drone Fly, Eristalis-Tenax

(Diptera, Syrphidae). *Can Entomol* **113**, 695-704 (1981).

16.   B. R. Geurten, R. Kern, E. Braun, M. Egelhaaf, A syntax of hoverfly flight prototypes. *J*

Exp Biol **213**, 2461-2475 (2010).

17.   C. Schilstra, J. H. Hateren, Blowfly flight and optic flow. I. Thorax kinematics and

flight dynamics. *The Journal of Experimental Biology* **202**, 1481 (1999).

18.   N. Boeddeker, L. Dittmar, W. Stürzl, M. Egelhaaf, The fine structure of honeybee

head and body yaw movements in a homing task. *Proceedings of the Royal Society B:*

Biological Sciences **277**, 1899-1906 (2010).

19.   T. S. Collett, Angular tracking and the optomotor response. An analysis of visual

reflex interaction in a hoverfly. *J Comp Physiol A* **140**, 145-158 (1980).

20.   D. O'Carroll, Feature-detecting neurons in dragonflies. *Nature* **362**, 541-543 (1993).

21.   K. Nordström, P. D. Barnett, D. C. O'Carroll, Insect detection of small targets moving

in visual clutter. *PLoS Biol* **4**, 378-386 (2006).

22.   S. Nicholas, J. Supple, R. Leibbrandt, P. T. Gonzalez-Bellido, K. Nordström, Integration

of small- and wide-field visual features in Target-Selective Descending Neurons of

both predatory and non-predatory dipterans. *J Neurosci* **38**, 10725-10733 (2018).

23.   M. A. Frye, R. M. Olberg, Visual receptive field properties of feature detecting

neurons in the dragonfly. *J Comp Physiol A* **177**, 569-576 (1995).

24.   S. Nicholas, R. Leibbrandt, K. Nordström, Visual motion sensitivity in descending

neurons in the hoverfly. *J Comp Physiol A* **206**, 149-163 (2020).

25.   P. T. Gonzalez-Bellido, H. Peng, J. Yang, A. P. Georgopoulos, R. M. Olberg, Eight pairs

of descending neurons in the dragonfly give wing motor centers accurate population

vector of prey direction. *Proc Natl Acad Sci USA* **110**, 676-701 (2013).

26.   P. D. Barnett, K. Nordström, D. C. O'Carroll, Retinotopic organization of small-field-

target-detecting neurons in the insect visual system. *Curr Biol* **17**, 569-578 (2007).

27.   A. D. Straw, E. J. Warrant, D. C. O'Carroll, A 'bright zone' in male hoverfly (*Eristalis*

tenax) eyes and associated faster motion detection and increased contrast

sensitivity. *J Exp Biol* **209**, 4339-4354 (2006).

28.   G. A. Horridge, The separation of visual axes in apposition compound eyes. *Philos*

Trans R Soc Lond B Biol Sci **285**, 1-59 (1978).

29.   R. M. Olberg, Identified target-selective visual interneurons descending from the

dragonfly brain. *J Comp Physiol A* **159**, 827-840 (1986).

30.   C. T. Hsu, V. Bhandawat, Organization of descending neurons in *Drosophila*

melanogaster. *Sci Rep* **6**, 20259 (2016).

31. R. M. Olberg (1978) Visual and multimodal interneurons in dragonflies. (University

of Washington, Seattle).

32. O. Dyakova, M. Mueller, M. Egelhaaf, K. Nordström, Image statistics of the

environment surrounding freely behaving hoverflies. *J Comp Physiol A* **205**, 373-385

(2019).

33. C. Städele, M. F. Keleş, J. M. Mongeau, M. A. Frye, Non-canonical receptive field

properties and neuromodulation of Feature-Detecting neurons in flies. *Curr Biol*

10.1016/j.cub.2020.04.069 (2020).

34. M. O. Franz, H. G. Krapp, Wide-field, motion-sensitive neurons and matched filters

for optic flow fields. *Biol Cybern* **83**, 185-197 (2000).

35. K. Nordström, D. C. O'Carroll, Feature detection and the hypercomplex property in

insects. *Trends Neurosci* **32**, 383-391 (2009).

36. M. F. Keleş, B. J. Hardcastle, C. Städele, Q. Xiao, M. A. Frye, Inhibitory interactions

and columnar inputs to an object motion detector in *Drosophila*. *Cell Rep* **30**, 2115-

2124.e2115 (2020).

37. M. F. Keleş, M. A. Frye, Object-detecting neurons in *Drosophila*. *Curr Biol* **27**, 680-687

(2017).

38. R. Tanaka, D. A. Clark, Object-displacement-sensitive visual neurons drive freezing in

Drosophila. *Curr Biol* 10.1016/j.cub.2020.04.068 (2020).

39. M. Egelhaaf, On the neuronal basis of figure-ground discrimination by relative

motion in the visual system of the fly. II. Figure-detection cells, a new class of visual

interneurones. *Biol Cybern* **52**, 195-209 (1985).

40.   A. K. Warzecha, M. Egelhaaf, A. Borst, Neural circuit tuning fly visual interneurons to

513   motion of small objects. I. Dissection of the circuit by pharmacological and

514   photoinactivation techniques. *J Neurophysiol* **69**, 329-339 (1993).

515   41.   W. Reichardt, T. Poggio, Figure-ground discrimination by relative movement in the

516   visual-system of the fly. Part I: Experimental results. *Biol Cybern* **35**, 81-100 (1979).

517   42.   T. Collett, Visual neurones in the anterior optic tract of the privet hawk moth. *J Comp*

518   *Physiol* **78**, 396-433 (1972).

519   43.   S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, A model for the detection of

520   moving targets in visual clutter inspired by insect physiology. *PLoS ONE* **3**, e2784

521   (2008).

522   44.   M. Egelhaaf, On the neuronal basis of figure-ground discrimination by relative

523   motion in the visual system of the fly. III. Possible input circuitries and behavioral

524   significance of the FD-cells. *Biol Cybern* **52**, 267-280 (1985).

525   45.   S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, Correlation between OFF and ON

526   Channels Underlies Dark Target Selectivity in an Insect Visual System. *J Neurosci* **33**,

527   13225-13232 (2013).

528   46.   S. D. Wiederman, D. C. O'Carroll, Discrimination of features in natural scenes by a

529   dragonfly neuron. *J Neurosci* **31**, 7141-7144 (2011).

530   47.   S. Nicholas, K. Nordström, Persistent firing and adaptation in optic flow sensitive

531   descending neurons. *Curr Biol* **30**, 2739-2748.e2732 (2020).

532   48.   T. S. Collett, M. F. Land, How hoverflies compute interception courses. *J Comp*

533   *Physiol A* **125**, 191-204 (1978).

49.   J. L. Fox, J. W. Aptekar, N. M. Zolotova, P. A. Shoemaker, M. A. Frye, Figure-ground

discrimination behavior in *Drosophila*. I. Spatial organization of wing-steering

responses. *J Exp Biol* **217**, 558-569 (2014).

50.   B. Cellini, J. M. Mongeau, Active vision shapes and coordinates flight motor

responses in flies. *Proc Natl Acad Sci U S A* **117**, 23085-23095 (2020).

51.   S. Nicholas, M. Thyselius, M. Holden, K. Nordström, Rearing and long-term

maintenance of *Eristalis tenax* hoverflies for research studies. *JoVE*

doi:10.3791/57711, e57711 (2018).

52.   D. H. Brainard, The Psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

53.   D. G. Pelli, The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision* **10**, 437-442 (1997).

Figure Legends

Figure 1. TSDN response to target motion

A) Pictograms of the round, black target, with a diameter of 3°, traversing a white background (left), or stationary optic flow (right), at 130°/s. **B)** Raw data trace from an extracellular TSDN recording. Timing of stimulus presentation indicated by colored bars (blue, target, and red, sparse dot field). **C)** Magnification of the raw data traces shown in panel B. **D)** The mean spiking response of different TSDNs was significantly reduced when the target moved across a stationary sparse dot field compared with a white background (N=30; ***P = 0.0005, two-tailed paired t-test).

Figure 2. TSDN response to target motion is affected by horizontal background optic flow

A) A pictogram indicating the three different types of translations, as seen by the hoverfly. Sideslip was displayed to the left or the right, lift up or down, and thrust towards or away from the hoverfly, all at 50 cm/s. **B)** The three different types of rotations used. As these were simulated to rotate around the hoverfly's position (24), the optic flow displayed on the screen can be described as yaw to the left or the right, pitch up or down, and clockwise or counter-clockwise roll, all at 50 deg/s. **C)** The TSDN response to a target traversing a stationary sparse dot field (left, grey), sideslip or yaw in either the same or the opposite direction to the target (red pictograms). All histograms from a single TSDN (mean ± sem, n=18) shown with 1 ms resolution after smoothing with a 20 ms square-wave filter. **D)** The mean response across TSDNs (N=12) to a target traversing stationary optic flow (left, grey), sideslip or yaw, in the same direction (Left") or opposite direction ("Right") to the target, after normalizing the data to each neuron's own response to a target traversing a white background. The line shows the mean and the filled data points correspond to the neuron in panel C. In panel D significance is shown with **** for $P < 0.0001$ following a one-way ANOVA followed by Dunnett's multiple comparison's test done together with the data shown in Fig. 4.

Figure 3. Frontal optic flow is required and sufficient

A) A pictogram of the separation of the background optic flow into four distinct positions:

ipsilateral (IL), dorsal (D), ventral (V) and contralateral (CL). The color coding shows the

receptive field of an example TSDN, and the arrows its local motion sensitivity. **B)** TSDN

responses to targets moving across syn-directional sideslip are significantly inhibited

compared to stationary control (grey) if the sideslip covers the full, dorsal or ventral screen.

C) TSDN responses to targets moving across opposite direction sideslip are significantly

facilitated compared to stationary control (grey, same data as in panel B) if the sideslip

covers the full, dorsal or ventral visual field. Panels B and C show data from the same

neurons (N=10, one-way ANOVA followed by Dunnett's multiple comparisons test, with

*P<0.05, **P<0.01).

Figure 4. The TSDN response to target motion is attenuated by perpendicular optic flow

A) The response from one TSDN to a target traversing a stationary sparse dot field (left,

grey, replotted from Fig. 2C), or different types of optic flow (red pictograms). All histograms

shown as mean ± sem (n=18) at 1 ms resolution, after smoothing with a 20 ms square-wave

filter. **B)** The mean response across TSDNs (N=12, same neurons as Fig. 2D) to a target

traversing a stationary sparse dot field (left, grey, replotted from Fig. 2D), or different types

of optic flow. Stars indicate significance, one-way ANOVA followed by Dunnett's multiple

comparisons test (**P<0.01, ***P<0.001 and ****P<0.0001), done together with the data

shown in Fig. 2D. Translations were simulated at 50 cm/s and rotations at 50°/s. The lines

show the mean and the filled data points correspond to the neuron in panel A.

Figure 5. Vector divergence between optic flow and target

A) The TSDN response to a target traversing a white background horizontally (left, grey) or

vertically (right, black). Histograms shown as mean ± sem (n=18) at 1 ms resolution, after

smoothing with a 20 ms square-wave filter. **B)** The mean spiking response of 8 TSDNs to

targets moving horizontally (grey, left) or vertically (black, right) over a white background.

The filled data points correspond to the neuron in panel A. **C)** TSDN responses to targets

moving horizontally across sideslip or lift, in the same direction as the target, or

perpendicular to the target, as indicated by the pictograms. Significance shown using one-

way ANOVA followed by Dunnett's multiple comparisons test, with *P=0.0232, **P=0.006,

***P=0.0003, ****P<0.0001. **D)** TSDN responses to targets moving vertically across sideslip

or lift, in the opposite direction to the target, or perpendicular to the target, as indicated by

the pictograms. Significance shown using one-way ANOVA followed by Dunnett's multiple

comparisons test, with *P=0.0198, **P=0.0021, ****P<0.0001. Same N=8 in panels B-D.

Figure 6. Elementary STMD input to TSDNs

Responses to a leading OFF edge, trailing ON edge, or a complete black target, with a side of

3°, traversing a white background at 130°/s. The black data show the predicted output from

a motion detector that compares luminance changes from at least 2 points in space. Data

replotted from (45), after normalizing to its own sum. The grey data show the predicted

output from an elementary STMD (ESTMD), which compares luminance changes from one

point in space. Data replotted from (45), after normalizing to its own sum. The white data

show the TSDN response to the same three stimuli (N=6) after normalizing the data from

each neuron to its own sum.

Supplementary Information

Figure S1. TSDNs do not respond to optic flow

A) To determine the TSDN response to the optic flow we first quantified each neuron's spontaneous rate, i.e. the spikes generated when viewing a white background, and used this as a threshold. For each neuron we then quantified the number of trials in which the response to the optic flow was above this threshold. This was converted to a percentage of responses. The data here show this percentage of responses across 12 TSDNs as mean ± sem. **B)** In each neuron, we quantified the mean response for those trials that were classified as responders (in panel A). This mean response was normalized to each neuron's mean response to a target traversing a white background, as throughout the rest of the paper. Data are shown as mean ± sem (N=12).

Movie 1. TSDN response to a target over a white background

The movie depicts the stimulus as displayed on the screen, with the hoverfly positioned centered and perpendicular to the screen. The hoverfly was positioned ventral side up, but we have rotated the movie for display purposes so the dorsal side is up. The colored lines show the outline of the receptive field, mapped as described previously (24). The red data at the bottom of the movie show the response of an example TSDN.

Movie 2. TSDN response to a target over a white background

Same as Movie 1, but slowed down 10 times.

Movie 3. TSDN response to a target over stationary optic flow

The movie depicts the stimulus as displayed on the screen, with the colored lines showing the outline of the receptive field. The red data at the bottom of the movie show the response of an example TSDN. In this case the optic flow was stationary, and appeared 0.5 s before the target.

Movie 4. TSDN response to a target over stationary optic flow

Same as Movie 3, but slowed down 10 times.

Movie 5. TSDN response to a target over syn-directional sideslip

The movie depicts the response of the same example TSDN to the same target motion as in
previous movies, but now traversing sideslip appearing 0.5 s before the target. The hoverfly
was positioned at a distance of 6.5 cm. At this viewing distance, the resulting optic flow
simulates the type of optic flow that would be generated by the hoverfly side-slipping
through the world at 50 cm/s. The simulated optic flow consisted of ca 6400 spheres, of
which roughly 1200 are projected onto the screen at any one time.

Movie 6. TSDN response to a target over syn-directional sideslip

Same as Movie 5, but slowed down 10 times. Note that many of the optic flow "targets" of
the correct size move through the receptive field (colored contour lines).

Movie 7. TSDN response to a target over opposite direction sideslip

The movie depicts the response of the same example TSDN to the same target motion as in
previous movies, but now traversing sideslip in the opposite direction to the target.

Movie 8. TSDN response to a target over opposite direction sideslip

Same as Movie 7, but slowed down 10 times.

658
659

A White background | Stationary

B 200 μV | 200 ms

Target
Optic flow

C 200 μV | 100 ms

D Mean response (spikes/s)

***

Target

White Background | Stationary

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6