Fine-mapping the consequences of site-specific glycan installation by shotgun scanning glycomutagenesis

Mingji Li[1†], Xiaolu Zheng[1†], Sudhanshu Shanker[2], Thapakorn Jaroentomeechai[1], Ilkay Koçer[1], Josef Byrne[1], Emily C. Cox[3], Qin Fu[4], Sheng Zhang[4], Jason W. Labonte[2,5], Jeffrey J. Gray[2]* and Matthew P. DeLisa[1,3,4]*

[1]Robert F. Smith School of Chemical and Biomolecular Engineering, Cornell University, Ithaca, New York 14853 USA

[2]Department of Chemical and Biomolecular Engineering, Johns Hopkins University, Baltimore, Maryland 21218 USA

[3]Biomedical and Biological Sciences, College of Veterinary Medicine, Cornell University, Ithaca, New York 14853 USA

[4]Cornell Institute of Biotechnology, Cornell University, Ithaca, New York 14853 USA

[5]Department of Chemistry, Franklin & Marshall College, Lancaster, Pennsylvania 17604 USA

*Address correspondence to: (1) Matthew P. DeLisa, Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University, Ithaca, NY 14853 USA. Tel: 607-254-8560; Email: md255@cornell.edu; and (2) Jeffrey J. Gray, Department of Chemical and Biomolecular Engineering, Johns Hopkins University, Baltimore, Maryland 21218 USA. Tel: 410-516-5313; Email: jgray@jhu.edu.

[†]These authors contributed equally to this work.

Abstract

N-linked glycosylation serves to diversify the proteome and is crucial for the folding and activity of numerous cellular proteins. Consequently, there is great interest in uncovering the rules that govern how glycosylation modulates protein properties so that the effects of site-specific glycosylation can be rationally exploited and eventually even predicted. Towards this goal, we describe a combinatorial strategy termed shotgun scanning glycomutagenesis (SSGM) that enables systematic investigation of the structural and functional consequences of glycan installation along a protein backbone. The utility of this approach was demonstrated with three different acceptor proteins, namely bacterial immunity protein Im7, bovine pancreatic ribonuclease A, and a human anti-HER2 single-chain Fv antibody, all of which were found to tolerate *N*-glycan attachment at a large number of positions and with relatively high efficiency. The stability and activity of many glycovariants was measurably altered by the *N*-linked glycan in a manner that critically depended on the precise location of the modification. Comparison of the results with calculations of simple geometrics and Rosetta energies suggested that glycosylation effects on protein activity may be predictable. By enabling a workflow for mapping glycan-mediated effects on acceptor proteins, glycomutagenesis opens up possibilities for accessing unexplored regions of glycoprotein structural space and engineering protein variants with designer biophysical and biological properties.

Introduction

Glycosylation of asparagine residues is one of the most abundant and structurally complex protein post-translational modifications [1, 2] and occurs in all domains of life [3]. Owing to their relatively large size and hydrophilicity or simply their presence at definite locations, asparagine-linked (*N*-linked) glycans can significantly alter protein properties including biological activity, chemical solubility, folding and stability, immunogenicity, and serum half-life.[4, 5] Hence, glycosylation effectively increases the diversity of the proteome by enriching the repertoire of protein characteristics beyond that dictated by the twenty canonical amino acids. For example, accumulating evidence indicates that the immune system diversifies the repertoire of antigen specificities by exclusively targeting the antigen-binding sites of immunoglobulins (IgGs) with post-translational modifications, in particular *N*-linked glycosylation [6]. Moreover, the profound effect of glycans on proteins has prompted widespread glycoengineering efforts to rationally manipulate key glycosylation parameters (*e.g.*, glycan size and structural composition, glycosite location and occupancy) as a means to optimize therapeutic proteins, vaccine formulations, and industrial enzymes [7-10]. Despite some notable successes, the routine use of glycosylation as a strategy for engineering proteins with advantageous properties is currently limited by our inability to generalize and predict how glycosylation affects protein structure and function.

Indeed, deciphering the "glycosylation code" that defines the interplay between a glycan and its underlying acceptor protein represents a grand challenge that requires access to large collections of chemically defined glycoproteins in sufficient quantities for characterization [11]. Such products are difficult to obtain because of the intrinsic variability with respect to the site of glycan attachment (macroheterogeneity) and glycan structure (microheterogeneity). This variability stems from the untemplated nature of glycosylation, which, unlike protein synthesis that involves a coding template, is defined by the relative activities of a number of glycosyltransferase enzymes. One way to circumvent this issue is through computational approaches that enable *in silico* exploration of glycosylation-induced effects on protein folding and stability [12, 13]; however, these involve a trade-off between molecular detail and glycoprotein size, with full-atomistic molecular dynamics simulations typically limited to only short glycopeptides (~10 residues) [12]. Another option

is chemical synthesis, which can furnish defined glycopeptides for investigating the local

effects of *N*-linked glycans on peptide conformation [14]. Although this approach is also not

amenable to full-length proteins, advances in expressed protein ligation (EPL) have

opened the door to convergent assembly of chemically synthesized glycopeptides with

recombinantly expressed protein domains to form larger glycoproteins bearing complex

N-glycans installed at discrete sites [15]. Using this technology, Imperiali and colleagues

created a panel of seven site-specifically glycosylated variants of the bacterial immunity

protein Im7 modified with the disaccharide *N*,*N'*-diacetylchitobiose (GlcNAc$_2$) and

assessed the kinetic and thermodynamic consequences of glycan installation at defined

locations [16]. Unfortunately, EPL is a technically demanding procedure, requiring manual

construction of each individual glycoprotein and thus limiting the number of testable

glycosite designs to just a small handful. Hence, there remains an unmet need for

scalable techniques that can rapidly supply large collections of discretely glycosylated

proteins for investigating the influence of *N*-glycans on the properties of conformationally

complex proteins.

Herein, we describe s̲hotgun s̲canning g̲lyco m̲utagenesis (SSGM), a reliable and

generalizable workflow for fine-mapping the effects of site-specific glycan installation at

every possible location of a protein backbone (**Fig. 1**). SSGM is inspired by protein

engineering methods based on alanine-scanning mutagenesis that have proven useful

for probing the contributions of individual amino acid sidechains to the properties of

proteins [17]. In combinatorial variations of this method, rather than requiring many mutant

proteins to be manually produced and separately characterized, combinatorial protein

libraries are used in conjunction with high-throughput screens or selections thereby

enabling a single library to simultaneously provide information about many different

residues [18, 19]. Along conceptually similar lines, SSGM is a glycoprotein-focused method

that involves design and construction of combinatorial acceptor protein libraries whereby:

(i) each member of the library carries a single *N*-glycosite "mutation" introduced at a

defined position along the protein backbone; and (ii) the complete ensemble of glycan

acceptor sites (sequons) in the library effectively covers every possible position in the

target protein. The resulting SSGM libraries are expressed using *N*-glycosylation-

competent bacteria in the context of glycoSNAP (g̲lyco̲sylation of s̲ecreted *N̲*-linked

Figure 1. Fine-mapping the effects of glycan installation. Schematic of shotgun scanning
glycomutagenesis (SSGM), a glycoprotein engineering method based on combinatorial protein libraries in
which glycosylation "sequon walking" is used to introduce an acceptor site at every possible position along
the protein backbone. Note that the multi-residue nature of a sequon (*e.g.*, N-X-S/T or D/E-X_1-N-X_2-S/T
where X, X_1, $X_2 \neq$ P) necessitates insertion or replacement of up to four additional amino acid substitutions
at each position. The resulting library is expressed in glycoengineered bacteria, providing an opportunity
for each library member to be expressed and glycosylated in a manner that is compatible with high-
throughput screening via glycoSNAP to interrogate the glycosylation phenotype of individual variants. By
integrating bacterial SSGM libraries with multiplexable assays, the biochemical and biophysical properties
of each glycovariant can be individually interrogated.

a̲c̲c̲e̲p̲t̲o̲r̲ p̲r̲o̲t̲e̲i̲n̲s̲) [20], a versatile high-throughput screen based on glycosylation and

extracellular secretion of the small (10 kDa in its mature form) *Escherichia coli* protein

YebF [21].

Using this new glycoprotein engineering tool, we constructed and screened three

different SSGM libraries corresponding to bacterial immunity protein Im7, bovine

pancreatic ribonuclease A (RNase A), and a human single-chain Fv antibody specific for

HER2 (scFv-HER2). Our results revealed that installation of *N*-glycans was tolerated at a

large number of positions and within all types of secondary structure, and the observed

N-glycosylation efficiency was quite high in the majority of cases. For many of these

glycovariants, the presence of *N*-glycans significantly impacted protein stability and/or

activity in a manner that critically depended on the precise location of the modification.

We computed simple geometric measures and Rosetta energetic estimates [22, 23] and

assessed which factors correlated with the sequon substitution or glycosylation effects on

stability and activity. Taken together, these findings demonstrate the utility of SSGM for

rapidly cataloging the biological and biophysical effects of *N*-glycan installation at single

amino acid resolution and in a protein-agnostic manner, thereby opening the door to

largely unexplored regions of glycoprotein structural space and providing a new tool for

engineering protein variants with advantageous properties.

Results

Reliable detection of acceptor protein glycosylation by glycoSNAP screening. To

enable screening of SSGM libraries, we hypothesized that target proteins of interest could

be fused to YebF and subsequently interrogated for N-glycosylation in a high-throughput

fashion by glycoSNAP screening (**Fig. 2a**). Briefly, YebF modified with an artificial

glycosite (*e.g.*, N-X-S/T or D/E-X_1-N-X_2-S/T where X, X_1, $X_2 \neq$ P) is expressed in the

presence of heterologous N-glycosylation machinery in *E. coli* cells that are bound to a

nitrocellulose filter membrane. Following secretion out of filter-bound colonies, putatively

glycosylated YebF is captured on a second nitrocellulose membrane, which is probed

with antibodies or lectins to detect N-linked glycans. In this way, glycoSNAP creates a

convenient genotype–glycophenotype linkage for facile scoring (glycosylated versus

aglycosylated) of YebF proteins secreted from individual bacterial colonies. We initially

focused on *E. coli* Im7 for several reasons: (i) it is a small, globular 87-residue protein

that lacks disulfide bonds and is well expressed in the periplasm where bacterial N-

glycosylation occurs [24]; (ii) although not a native glycoprotein, Im7 modified at its C-

terminus with a DQNAT glycosylation tag has been glycosylated by the *Campylobacter*

jejuni N-glycosylation machinery in *E. coli* [24]; (iii) crystal structures are available for wild-

type (wt) Im7 [25] and for Im7 in complex with its cognate toxin colicin E7 (ColE7) [26]; and

(iv) a limited set of seven Im7 variants was previously generated to determine the effects

of $GlcNAc_2$ attachment on folding and stability [16], providing some useful reference points

for comparison.

To determine whether Im7 was compatible with the glycoSNAP procedure, *E. coli*

strain CLM24 was co-transformed with a plasmid encoding YebF-Im7 that was modified

at its C-terminus with a DQNAT glycosylation tag [27] along with two additional plasmids,

one encoding glycosyltransferase (GT) enzymes for the biosynthesis of the N-glycan and

the other encoding the oligosaccharyltransferase (OST) for transfer of the resulting N-

glycan to acceptor proteins. To minimize microheterogeneity so that modified acceptor

Figure 2. Construction and interrogation of SSGM libraries. (a) Schematic of SSGM library construction using multiplex inverse PCR approach. The resulting plasmid library, encoding acceptor protein variants with glycosite substitutions at every possible position, was used to co-transform *E. coli* strain CLM24 along with two additional plasmids encoding the requisite *N*-glycosylation machinery from *C. jejuni*. The resulting bacterial library was plated on solid agar, after which colonies and their secreted glycoproteins were replica plated on nitrocellulose membranes as described in the text. (b) Immunoblot analysis of acceptor proteins in colony secretions (left) and extracellular supernatant fractions (right) derived from *E. coli* CLM24 carrying a plasmid encoding either YebF-Im7 or YebF-Im7^{DQNAT} along with plasmids encoding *N*-glycosylation machinery with either wild-type *Cj*PglB (wt) or an inactive mutant (mut). (c) Same as in (b) but with YebF-RNase A and YebF-RNase A^{N34} in colony secretions (left) and periplasmic fractions (right). Blots were probed with anti-polyhistidine antibody (α-His) to detect acceptor protein and SBA or hR6 serum to detect the glycan. Bottom color panels in (b) and (c) depict overlay of α-His and SBA blots (merge). Arrows denote aglycosylated (g0) and singly glycosylated (g1) forms of YebF-Im7^{DQNAT} or YebF-RNase A^{N34}. Molecular weight (M_W) markers are indicated at left. Results are representative of at least three biological replicates.

proteins all carried identical glycans, we created a system for producing homogeneous

N-glycans with the structure GalNAc$_5$(Glc)GlcNAc, which is one of several structurally

related glycan donors that can be efficiently transferred to target proteins in *E. coli* by the

C. jejuni OST PglB (*Cj*PglB) [28, 29]. This is significant because it enables differences in

glycosylation efficiency to be attributed to accessibility of a given acceptor site rather than

poor compatibility between the lipid-linked oligosaccharide (LLO) donor and the OST.

When plated on solid agar and subjected to the colony-blotting method, cells

expressing YebF-Im7^{DQNAT}, or a control YebF-Im7 construct that lacked the glycosylation

tag, were able to secrete the fusion into the extracellular medium as evidenced by cross-

reaction of an anti-His antibody with the membranes (**Fig. 2b**). However, only the strain

expressing YebF-Im7^{DQNAT} in the presence of wt *Cj*PglB, but not a *Cj*PglB variant rendered inactive by two active-site mutations (D54N and E316Q) [20], gave rise to colonies that reacted with soybean aggluntinin (SBA) (**Fig. 2b**), a lectin that binds terminal GalNAc residues in the minimal *C. jejuni N*-glycan [28]. The colony blotting results were corroborated by immunoblot analysis of culture supernatants, which revealed that YebF-Im7 and YebF-Im7^{DQNAT} were both secreted into the extracellular medium but only the latter was glycosylated as evidenced by the appearance of a higher molecular weight band in the blot probed with glycan-specific antiserum (**Fig. 2b**). As expected, no glycan-specific signal was detected in colony blots or immunoblots corresponding to cells carrying the mutant *Cj*PglB enzyme (**Fig. 2b**). Importantly, the predominant glycan attached to YebF-Im7^{DQNAT} corresponded to GalNAc$_5$(Glc)GlcNAc, which represented >98% of all detected glycoforms as confirmed by mass spectrometry (**Supplementary Fig. 1**). Collectively, these results confirmed the compatibility of bacterial Im7 with our glycosylation workflow, yielding homogenously modified acceptor proteins that were readily detected by glycoSNAP screening.

Rapid identification of acceptor site permissiveness using SSGM. Next, the plasmid encoding YebF-Im7 was mutagenized to create a library of Im7 gene sequences, each carrying an individual sequon substitution and cumulatively covering all positions in the Im7 protein. Mutagenesis was performed using multiplex inverse PCR [30] with a set of divergent abutting primers that were designed to amplify the entire plasmid and introduce an acceptor asparagine residue at every position in the Im7 gene (with the two upstream and two downstream residues being changed to DQ and AT, respectively), thereby yielding a highly focused plasmid library enriched with in-frame clones each bearing a single DQNAT acceptor motif at a defined position (**Fig. 2a**). The resulting plasmid library was introduced into strain CLM24 carrying the requisite *N*-glycosylation machinery, after which the library-transformed cells were plated on solid agar and subjected to glycoSNAP screening. From one membrane, we detected a total of ~200 glycosylation-positive colonies, of which 20 were randomly chosen for further analysis. Sequencing confirmed that a single in-frame DQNAT motif was present in each isolated hit, with the Im7^{N37} and Im7^{N58} variants (where the superscript denotes the location of the asparagine residue) occurring three and two times, respectively (**Fig. 3a**). The hits were fairly evenly

Figure 3. Exhaustive *N*-glycosylation of bacterial Im7 and its consequences. (a) Primary sequence and predicted secondary structure for *E. coli* Im7 immunity protein. Asterisks denote location and frequency of glycosite hits isolated using SSGM. Predicted structures adapted from PDB ID 1AYI. (b) Immunoblot analysis of supernatant fractions from CLM24 cells carrying plasmids encoding YebF-Im7 fusions with sequon mutations at indicated position and requisite *N*-glycosylation machinery. Blots were probed with anti-polyhistidine antibody (α-His) to detect acceptor protein (top panel) and hR6 serum against the glycan (bottom panel). Markers for aglycosylated (g0) and singly glycosylated (g1) forms of acceptor proteins are indicated at right. Molecular weight (*M*$_W$) markers are indicated at left. Asterisk indicates construct with mutation that introduced stop codon just before 6xHis tag, preventing α-His detection. Results are representative of at least three biological replicates. (c) Mapping of *in vivo* glycosylation efficiency onto three-dimensional structure of Im7 in complex with ColE7 (left). Heatmap analysis of the glycosylation efficiency was determined based on densitometric quantification of the percent glycosylated (defined as g1/[g0+g1] ratio) for each acceptor protein in the anti-His immunoblot. Detailed interactions between ColE7 and Im7, highlighting sidechains of Im7 in the regions of α1-loop12-α2 (residues 19-39; middle) and loop23-α3-loop34 (residues 46-63; right). Heatmap analysis of change in binding activity was determined by normalizing activity measured for glycosylated sequon variant by aglycosylated counterpart. (d) Binding activity of glycosylated (gray bars) and aglycosylated (white bars) YebF-Im7 variants recovered from supernatants was measured by ELISA with ColE7 as immobilized antigen. All data were normalized to binding activity measured for aglycosylated YebF-Im7 lacking a sequon (wt), such that values greater than 1 (denoted by dashed red line) indicate enhanced binding activity relative to wt Im7. Dashed boxes correspond to two regions (Region 1: residues 23-33; Region 2: residues 58-69) that have many variants with increased activity. Data are average of three biological replicates and error bars represent standard deviation of the mean. (e) DSF analysis of 15 most active YebF-Im7 variants with and without glycosylation. *T*$_m$ calculated as midpoint of thermal transition between native and unfolded states. Dashed line indicates *T*$_m$ for wt YebF-Im7 (38.6 ± 1.0 °C). Black bars are average of three independent replicates with error bars reported as standard error of the mean.

distributed throughout the entire Im7 sequence and situated in every type of secondary structure including bends, turns, and α-helices, consistent with X-ray crystallographic data showing that occupied glycosylation sites can occur on all secondary structural elements [31]. Immunoblot analysis confirmed that each of the selected clones was efficiently glycosylated (**Fig. 3b**).

To exhaustively explore glycosylation sequence space, we manually constructed all possible individual Im7 sequon variants (80 in total) using the multiplex PCR primer pairs to introduce DQNAT sequons at every position of the protein. A strikingly large number (79 out of 80) of these variants were found to be glycosylated, many with an efficiency that was at or near 100% as estimated from densitometry of the anti-His blot (**Fig. 3c** and **Supplementary Fig. 2a**). Because glycosylation by *Cj*PglB can occur both before and after protein folding is completed (**Supplementary Fig. 2b**) [32, 33], the secondary and tertiary structure around a glycosylation site is likely to have a direct effect on the extent to which a given site is occupied, if at all. Indeed, it has been observed that sequons located in structurally defined regions of folded acceptor proteins are poorly glycosylated and that partial unfolding is required to increase glycosylation efficiency at these sites [33, 34]. To determine if the structural context for any of the Im7 sequon variants was a determinant for the timing and efficiency of glycosylation, we performed *in vitro* glycosylation reactions in which already folded but yet-to-be glycosylated YebF-Im7 proteins derived from culture supernatants were incubated with purified *Cj*PglB and glycan donor. Remarkably, there was near perfect agreement between the *in vitro* and *in vivo* glycosylation results, with nearly all of the purified Im7 variants undergoing highly efficient glycosylation that was at or near 100% with few exceptions (**Supplementary Fig. 2a**). The observation that so many Im7 variants were efficiently glycosylated by the *Cj*PglB enzyme *in vitro* (*i.e.*, after folding had been completed) indicates that each sequon was located in either a structurally compliant position (*e.g.*, flexible and surface-exposed loops) within the folded protein or in a region of the protein that became partially unfolded during the *in vitro* glycosylation reaction. While broad accessibility is certainly plausible given the small size and simple topology of Im7, we cannot rule out the contribution of conformational destabilizing effects caused by replacement of five-residue stretches of native amino acids in the protein. Regardless of the exact reason, these results indicate

that Im7 was extremely tolerant to the installation of *N*-glycans over its entire structure both *in vivo* and *in vitro*.

Structural and functional consequences of Im7 glycosylation. To exhaustively determine the effect of glycan attachment on acceptor protein properties, we first quantified binding activity of all 80 Im7 sequon variants with and without glycosylation by subjecting each to multiwell enzyme-linked immunosorbent assay (ELISA) using purified ColE7 as immobilized antigen. Native Im7 interacts with ColE7, a 60-kDa bacterial toxin that is cytotoxic in the absence of the cognate Im7 inhibitor [35]. With an eye towards multiplexibility, we chose to assay YebF-Im7 fusions directly because: (i) it obviated the need for molecular reformatting of the expression constructs; (ii) the fusions could be isolated as relatively pure species from cell-free supernatants, bypassing the need for extensive purification; and (iii) the introduction of the small YebF domain had no measurable effect on ColE7-binding activity (**Supplementary Fig. 3a**). Whereas nearly two thirds of the YebF-Im7 fusions were either unaffected by glycosylation or rendered inactive by introduction of the DQNAT motif alone, particularly in a contiguous stretch between residues 50-57 of Im7, the remaining one third exhibited significantly altered binding activity that was attributable to the presence of the *N*-glycan (**Fig. 3d**). These glycosylation-induced effects were clearly dependent on the precise location of the modification. Indeed, some of the most striking increases in binding activity for glycosylated variants over their aglycosylated counterparts were observed to occur at the transition between different types of secondary structure (*e.g.*, variants Im7^{N33}, Im7^{N58} and Im7^{N65}). These results were particularly noteworthy in light of the elevated probability of finding naturally occurring sequons in locations where secondary structure changes [31].

Among the Im7 glycovariants whose activity was most significantly affected both positively and negatively by *N*-glycosylation, the majority were located in two distinct regions covering residues 23–33 and 58–69 (**Fig. 3d**). These regions occurred within the two arms of Im7 (one located in α1–loop12–α2 from residue 19 to 39 and the other in loop23–α3–loop34 from residue 46 to 63) that interact extensively with a continuous region in ColE7 in the crystal structure (**Fig. 3c**) [26]. The two interfaces are charge-complementary, and charge interactions are largely responsible for the tight and specific binding between the two proteins; hence, it was not surprising that binding activity was

sensitive to *N*-glycan attachment in the vicinity of these interfaces. It should be pointed out that the presence of an *N*-glycan in some of these positions was uniquely modulatory, as substitution of DQNAT alone in these same locations generally had little effect on activity, as evidenced by the comparable ColE7 binding measured for aglycosylated Im7 variants versus wt Im7 (**Supplementary Fig. 3b**).

To determine whether any of the glycosylation-induced increases in binding activity were related to stabilization of the native fold, the most active Im7 glycovariants were subjected to differential scanning fluorimetry (DSF) with SYPRO Orange dye in a real-time PCR instrument. Previous studies showed that melting temperature (T_m) values obtained by DSF correlated well with those determined by circular dichroism (CD) thermal denaturation [36]. Here too, we observed excellent agreement between these two methods, which both yielded T_m values for wt Im7 (~39 ^{o}C, **Supplementary Fig. 3c and d**) that agreed with a previously reported value [35]. Importantly, the presence of the small YebF domain did not significantly alter the T_m value for Im7 (**Supplementary Fig. 3d**), consistent with its lack of effect on ColE7-binding activity. We also confirmed that DSF results obtained using YebF-Im7 derived directly from cell-free supernatants were indistinguishable from those obtained with more extensively purified YebF-Im7 (**Supplementary Fig. 3d**). Using DSF, the average T_m values for glycosylated and aglycosylated versions of each Im7 variant were measured, and the change in unfolding temperature, ΔT_m, was calculated such that a positive ΔT_m signified an increase in structural order and a reduced conformational flexibility due to appending a glycan. Several of the variants exhibited positive ΔT_m values, with the largest increases corresponding to glycan installation at N33, N59, N60, N65 and N80 (**Fig. 3e**). Conversely, glycans at N10, N58, and N64 caused the largest decreases in T_m, indicative of glycan-induced protein structural changes that destabilized the protein.

To test whether protein-structure analyses could explain the observed effects of sequon substitution and glycosylation, we modeled the sequon-substituted variants and generated ensembles of glycan conformations for Im7 alone and in the context of Im7 bound to E7. We calculated simple geometric measures (secondary structure, burial, distance to the binding site, and surface area) and Rosetta energy estimates (stability and interface score) (**Supplementary Figs. 4** and **5**; **Supplementary Results**).

Surprisingly, none of these factors correlated with either stability or activity of the

aglycosylated or glycosylated Im7 constructs. Only the attractive van der Waals energy

had a weak correlation with aglycosylated binding activity (**Supplementary Fig. 6**),

possibly because of Im7's marginal stability and the known flexibility of helix α4 [37]. The

ensembles of glycan conformations for several glycovariants in the context of Im7 bound

to E7 (**Supplementary Fig. 7**) indicated that (i) the glycan and the bound protein often

interact to change the binding activity positively or negatively (**Supplementary Results**)

and (ii) enhanced binding appears to be mediated by multiple low-energy glycan

conformations making favorable interactions with E7.

Extension of SSGM to a protein with more complex topology. We next turned our

attention to bovine RNase A. Like Im7, RNase A has been intensely studied from a

structure–function standpoint and has been pivotal to understanding many aspects of

enzymology, biological chemistry, and protein folding and stability. We chose RNase A

because (i) it is a relatively small, basic protein, containing 124 residues but with a more

complex topology than Im7, with all major types of secondary structure, namely α-helices,

β-sheets, and turns, represented; (ii) the natively glycosylated form of RNase A, namely

RNase B, contains a single *N*-linked oligosaccharide at N34 and a crystal structure is

available [38]; (iii) glycosylation at N34 has no apparent effect on the secondary or tertiary

structure [38] but does appear to alter the thermal stability [39] although this is controversial

[40]; and (iv) RNase A modified with an optimal bacterial sequon at the native N34

glycosylation site (RNase A^{N34}) can undergo *Cj*PglB-dependent glycosylation *in vivo* and

in vitro [32, 33]. For these reasons, RNase A represented an ideal target for SSGM.

Extracellular secretion of glycosylated YebF-RNase A^{N34} was observed in colony

blots and immunoblots (**Fig. 2c**), confirming the compatibility of RNase A with glycoSNAP

screening. An SSGM library was created by subjecting YebF-RNase A plasmid DNA to

the multiplex inverse PCR method, after which glycoSNAP screening was performed

using CLM24 cells carrying plasmids encoding the library and the *C. jejuni* glycosylation

machinery. A total of ~100 glycosylation-positive colonies were randomly selected from

two membranes and subjected to sequencing analysis. Of the 97 sequence-confirmed

hits, only 50 were unique, as many of the sequences were isolated multiple times (*e.g.*,

seven times each for RNase A^{N41} and RNase A^{N122}; **Fig. 4a**). The sequons of these hits

Figure 4. Glycomutagenesis of bovine pancreatic RNase A. (a) Primary sequence and predicted secondary structure for bovine pancreatic RNase A. Asterisks denote location and frequency of glycosite hits isolated using SSGM. Predicted structures adapted from PDB ID 1RBX. (b) Mapping of *in vivo* (left) and *in vitro* (right) glycosylation efficiency onto three-dimensional structure of RNase A. Heatmap analysis of glycosylation efficiency was determined based on densitometric quantification of percent glycosylated (defined as g1/[g0+g1] ratio) for each acceptor protein in anti-His immunoblot. (c) Enzymatic activity of glycosylated (gray bars) and aglycosylated (white bars) RNase A variants recovered from culture supernatants. All data were normalized to binding activity measured for aglycosylated YebF-RNase A lacking a sequon (wt). Data are average of three biological replicates and error bars represent standard deviation of the mean. (d) DSF analysis of YebF-RNase A variants with and without glycosylation. T_m was calculated as midpoint of thermal transition between native and unfolded states. Dashed line indicates T_m for wt YebF-RNase A (59.0 $\pm$ 0.1 °C). Black bars are average of three independent replicates with error bars reported as standard error of the mean.

were uniformly distributed throughout the primary sequence and found in every type of secondary structural element, akin to the results with Im7. Immunoblot analysis confirmed that all selected clones were glycosylated, and the efficiency for most was at or near 100% as estimated by densitometry analysis of the anti-His blots (**Fig. 4b** and **Supplementary Fig. 8**). To investigate whether the structural context of the sequon impacted the possible timing of PglB-mediated glycan installation, we performed *in vitro* glycosylation of folded RNase A variants. While some of the variants were glycosylated

as efficiently *in vitro* as they were *in vivo* (*e.g.*, RNase A[N46] and RNase A[N64]), an unexpectedly large number of variants showed significantly lower levels of glycosylation under *in vitro* conditions (**Fig. 4b** and **Supplementary Fig. 8**). Most notably among these were variants N34, N35, N36, N43, N51, N61, N69, N72, N80, N89, and N104, which were all efficiently glycosylated *in vivo* but underwent little or no detectable glycosylation *in vitro*. These sequons occur at locations that were likely to be accessible to the OST during translation/translocation when the proteins are unfolded but became inaccessible after the protein completed folding. Indeed, the native *N*-glycosylation site at N34 is located in a structured domain, suggesting that the poor *in vitro* glycosylation at this specific location (and perhaps also at the nearby N36 and N43 sites) was due to sequon inaccessibility in the folded state. Such folding-dependent recognition of this site has been observed previously [32, 33] and, together with the results presented here, supports a model whereby *in vivo* glycosylation of these particular sequons involves glycan installation prior to folding, either co- or post-translocationally (**Supplementary Fig. 2b**).

To determine the consequences of glycosylation at the 50 unique sites, the ability of glycosylated and aglycosylated versions of each sequon variant to catalyze the hydrolysis of the phosphodiester bonds in RNA was evaluated. While the addition of YebF had little to no effect on RNase A activity (**Supplementary Fig. 9a**), more than half of the RNase A variants were inactivated by substitution of the DQNAT sequon (**Fig. 4c**). To determine if this might be due to the substitution of five residues in the target protein, a requirement for optimal recognition by *Cj*PglB [41], we mutated RNase A more conservatively at a select number of sites. Specifically, we generated minimal sequons (D-X-N-X-T/S or X-X-N-X-T/S, where X represents the native amino acid), which in most cases required only 1 or 2 amino acid changes. Each of these mutants was completely inactive except for RNase A[N55] with a DVNAT sequon, which showed some activity but was still significantly less active than the wt enzyme (**Supplementary Fig. 9b**). Hence, even relatively minor sequence perturbations at these positions, in addition to the less subtle substitution with DQNAT, were all capable of inactivating RNase A. More careful inspection revealed that the majority of variants with little to no activity corresponded to the substitution of sequons in locations that would be predicted to disrupt catalytically important residues or disulfide bonds (**Fig. 4c** and **Supplementary Results**).

Among the variants that retained function, only eight (sequons at N34, N35, N36,

N51, N53, N61, N89, and N104) showed activity that was on par (>50%) with wt RNase

A but none were more active than their aglycosylated counterpart (**Fig. 4c**). In the case

of RNase A^{N119}, introduction of the DQNAT sequence completely abrogated catalytic

activity, consistent with previous findings that the relative activity of an H119N mutant was

reduced to less than 1% of wt RNase A, with k_{cat}/K_M values reduced by 100- to 1000-fold

depending on the substrate used [42]. Despite the importance of this residue for catalysis,

glycosylation at this position partially restored enzymatic activity, indicating an *N*-glycan-

dependent gain-of-function.

To determine whether glycosylation impacted stability, we again used DSF to

analyze the most active RNase A glycovariants along with RNase A^{N93}, which was

randomly chosen as a representative inactive variant. The measured T_m values for wt

YebF-RNase A and its unfused counterpart were both ~59 $^{\circ}$C (**Supplementary Fig. 9c**),

in close agreement with previous findings [40], while the T_m values for all the YebF-RNase

A variants spanned a range from 58–63 $^{\circ}$C (**Fig. 4d**). Most exhibited positive ΔT_m values

compared to their aglycosylated counterpart, including the RNase A^{N119} variant,

suggesting that the restoration of activity caused by glycan attachment at N119 also

served to stabilize the protein. In contrast, RNase A^{N89} and RNase A^{N93} exhibited large

negative ΔT_m values that coincided with slightly weakened activity due to glycan

attachment in the case of N89 and complete inactivation in the case of N93.

Next, we measured geometries and calculated Rosetta metrics to determine the

effect of sequon substitution and glycosylation on RNase A activity. For the activity of

aglycosylated and glycosylated RNase A variants in the context of all seven metrics, the

only weakly correlated metric was the total Rosetta energy score (R^2 = 0.14 for

aglycosylated activity and R^2 = 0.12 for glycosylated activity; **Supplementary Figs. 4, 5**).

These metrics may be less useful for RNase A because the activities are primarily

explained by the disruption of the active site and the disulfide bonds, which are not

captured in these metrics.

Investigation of IgG variable domain glycosylation using SSGM. We next

investigated antibody variable domain glycosylation, a phenomenon that is observed for

~15% of serum IgGs and contributes to diversification of the B-cell antibody repertoire [6].

Although glycan installation within the variable domains of Fab arms has been long known, the rules governing the selection of *N*-glycosylation sites in Fab domains that emerge during somatic hypermutation and the functional consequences of the attached glycans remain poorly understood. To systematically investigate this phenomenon using SSGM, the two variable domains, V_H and V_L, from the human anti-HER2 monoclonal antibody were joined by a flexible linker to form scFv-HER2 that was subsequently modified at its N-terminus with YebF and at its C-terminus with a DQNAT motif. Extracellular secretion of glycosylated YebF-scFv-HER2^{DQNAT} was observed in colony blots and immunoblots (**Fig. 5a**), confirming the compatibility of scFv-HER2 with glycoSNAP screening. Because variable domain glycosylation was subject to selection mechanisms that depend on the nature of the antigen [6], we modified the SSGM strategy to enable dual screening of glycosylation and antigen-binding activity by labeling colonies with SBA lectin and the extracellular domain (residues 1-652) of human HER2 (HER2-ED), which was avidly bound by scFv-HER2^{DQNAT} fused to YebF (**Supplementary Fig. 10a**). In this way, two-color screening could be used to identify colonies that were positive both for glycosylation and for antigen binding, as demonstrated with the YebF-scFv-HER2^{DQNAT} construct (**Fig. 5a**). Next, we constructed and screened an SSGM library, after which two-color glycoSNAP screening was performed with CLM24 cells carrying plasmids encoding the library and the *C. jejuni* glycosylation machinery. A total of ~60 dual-positive hits were isolated from membranes, of which 21 were determined to be non-redundant (*e.g.*, N58 in V_L and N42 in V_H were each isolated 12 times) (**Fig. 5b**) and subsequently confirmed for glycosylation by immunoblot analysis (**Supplementary Fig. 10b**). The sequons of these hits were sparsely distributed throughout the primary sequence, with a large proportion clustering just after the second and third complementarity-determining regions (CDRs) of the V_L domain and also in the flexible linker, suggesting a strong selection bias for specific sites that both tolerated glycosylation and retained function. Interestingly, a few of the identified sequons occurred in CDR2 of the V_L domain and CDR1 and CDR2 of the V_H domain, consistent with naturally occurring IgG repertoires in which *N*-glycosites were observed preferentially in the CDRs [6].

In terms of function, all 21 scFv-HER2 glycovariants exhibited HER2-ED binding activity above background (**Fig. 5c**), which was expected given that the screening

Figure 5. Probing scFv antibody glycosylation using glycomutagenesis. (a) Immunoblot analysis of acceptor proteins in colony secretions (left and middle) and periplasmic fractions (right) derived from *E. coli* CLM24 carrying plasmids encoding scFv-HER2^{DQNAT} and requisite *N*-glycosylation machinery with either wild-type *Cj*PglB (wt) or an inactive mutant (mut). Blots were probed with anti-polyhistidine antibody (α-His) to detect acceptor protein, SBA or hR6 serum to detect the glycan, and HER2-ED to detect antibody binding. Bottom color panels depict overlay of α-His and SBA blots or SBA and HER2 blots (merge). Arrows denote aglycosylated (g0) and singly glycosylated (g1) forms of scFv-HER2^{DQNAT}. Molecular weight (M_W) markers are indicated at left. Results are representative of at least three biological replicates. (b) Frequency and position of *N*-glycosylation sites in scFv-HER2^{DQNAT} glycovariants isolated using SSGM. (c) Binding activity of glycosylated (gray bars) and aglycosylated (white bars) scFv-HER2^{DQNAT} variants as measured by ELISA with HER2-ED as immobilized antigen. All data were normalized to binding activity measured for aglycosylated scFv-HER2 lacking a sequon (wt), such that values greater than 1 (denoted by dashed line) indicate enhanced binding activity relative to wt scFv-HER2. Data are average of three biological replicates and error bars represent standard deviation of the mean. (d) DSF analysis of YebF-scFv-HER2 variants with and without glycosylation. T_m was calculated as midpoint of thermal transition between native and unfolded states. Dashed line indicates T_m for wt YebF-scFv-HER2 (68.2 ± 0.1 °C). Black bars are average of three independent replicates with error bars reported as standard error of the mean.

process was adapted to include antigen binding. Importantly, nine of these glycovariants (N58, N64, and N109 in V_L; N3, N4, N9, N10 in linker; N42 and N113 in V_H) exhibited increased binding compared to their aglycosylated counterpart, and most of these were also more active than the parental scFv-HER2. For the five clones exhibiting the greatest increase in activity due to glycosylation, we measured T_m values and found that glycan attachment in general did not affect stability (**Fig. 5d**). However, the one exception was N64 V_L which experienced a 2.6 °C increase in T_m due to the addition of the *N*-glycan. Overall, these results are in agreement with several previous studies showing that variable region glycans contribute to antibody binding characteristics and stability in a manner that depends on the precise location of the glycan [6, 43] and suggest that glycosylation in this region may be a useful strategy for fine-tuning the performance of IgG antibodies and their derivatives.

Next, we compared the experimental binding activity for scFv-HER2 with multiple geometric and Rosetta metrics. Unlike Im7 or RNase A, scFv-HER2 activity correlated with many of our metrics. First, sequon burial reduces the binding affinity of scFv-HER2 for its antigen both in the aglycosylated ($R^2 = 0.21$) and glycosylated states ($R^2 = 0.43$) (**Supplementary Fig. 4b** and **Fig. 6a**). Similarly, the closer the sequon was to the paratope, the greater the likelihood of reduced activity ($R^2 = 0.20$ and 0.23 for the aglycosylated and glycosylated variants, respectively; **Supplementary Fig. 4c** and **Fig. 6b**). The buried surface area correlated with the glycosylated state activity ($R^2 = 0.19$, **Supplementary Fig. 5e**). The strongest predictors, however, were the Rosetta scores. The aglycosylated antibody–antigen complex total score correlated with experimental binding activity ($R^2 = 0.49$, **Supplementary Fig. 4f**). For the glycosylated state, the activity correlated with both the total Rosetta score ($R^2 = 0.49$, **Fig. 6c**) and the interface score ($R^2 = 0.63$, **Supplementary Fig. 5g**). These Rosetta scores are primarily driven by the van der Waals complementarity and to a lesser extent electrostatics (**Supplementary Figs. 6** and **11**).

For the aglycosylated activities, we selected three variants for deeper analysis: two variants that had low binding activity and a poor Rosetta score (N36 V_L, N113 V_H; black circles in **Supplementary Fig. 6a**) and one variant with high activity and a favorable Rosetta score (N108 V_L; red circle in **Supplementary Fig. 6a**). Both N36 V_L and N113

Figure 6: Quantitative analysis of scFv-HER2 variants. The structure of scFv-HER2 V_L (red) and V_H (blue) domains in complex with HER2 protein (gray) is shown at bottom left. Regression analyses of log activity ratio (glycosylated / wild-type) versus (a) burial of sequon substitution site, (b) distance of closest HER2 residue from the sequon substitution site, and (c) total Rosetta score. In all three panels, the dark red lines are the respective regression lines. Colors of dots in (c) show the respective secondary structure of the sequon substitution site. Orange, green, and blue correspond to α-helix, β-strand, and loop regions, respectively. N58 V_L (red circle) has the highest glycosylated binding activity increase and is discussed in the text. (d) Wild-type representation of sites used for analysis of sequon substitution (36 V_L, 108 V_L, and 113 V_H) and glycosylation (58 V_L). Side-chain colors reflect their respective secondary structures. (e) Glycan arrangement (orange sticks) from eight low energy conformations of glycosylated N58 V_L variant of scFv-HER2, revealing possible glycan-HER2 interaction responsible for binding activity improvement.

V_H sites are situated on β-strands in compact regions of the anti-HER2 antibody on the side opposite the antigen-binding site (**Fig. 6d**, green sticks). The reduced stability arises from the steric clash of substituting a sequon inside (or near) a close-packed region of the protein (Rosetta terms for steric clashes (vdW_rep) of 90.2 and 79.8 Rosetta energy units (REU) for the N36 V_L and N113 V_H, respectively). When glycosylated, the clashes worsen in the Rosetta models, corresponding to low activity (black circles in **Supplementary Fig. 6a**). On the other hand, site N108 V_L is located at the C-terminal end of V_H (**Fig. 6d**, blue sticks). Sequon substitution had a relatively small effect on the electrostatic interactions (−6.2 REU) and a greater effect on the repulsive van der Waals

terms (−28.0 REU), indicating that new side chains are acceptable in less compact

regions. A similar outcome was reported following substitution mutation of a human

monoclonal antibody [44].

To understand how *N*-glycosylation was able to improve binding activity, we

selected mutant N58 V_L because the aglycosylated variant was 25.6% more active than

the wt scFv-HER2 and glycan addition improves the binding an additional 1.83-fold.

Residue N58 V_L resides in the turn between strands 1 and 2 (**Fig. 6d**, blue backbone).

From Rosetta-generated glycosylated structures, the low-energy states showed

interfacial contacts between the glycan and the surface residues of HER2 (**Fig. 6e**),

improving both the total Rosetta score and the interface score (red circle in **Fig. 6c** and

Supplementary Fig. 5g) and explaining the binding activity improvement as resulting

from favorable glycan-antigen contacts.

Discussion

In this study, we developed a new protein engineering workflow called SSGM that enables

comprehensive mapping of the structural and functional consequences of site-specific

glycan installation for any protein of interest. By combining synthetic glycosite libraries

with a high-throughput screen for protein glycosylation [20], we showed that a single SSGM

experiment allowed the effects of many different site-directed glycan "mutations" to be

probed for their contribution to the properties of a protein. In this way, SSGM is

conceptually analogous to combinatorial alanine-scanning mutagenesis, which allows

systematic determination of the importance of individual amino acids to protein structure

and function [17-19]. Here, we demonstrated the utility and generality of the SSGM technique

using three different acceptor proteins namely bacterial Im7, bovine RNase A, and human

scFv-HER2. Consistent with the known modulatory effects of *N*-glycans [4, 5], many

glycovariants of these proteins exhibited detectably altered stability and activity that

resulted from covalent attachment of *N*-glycans at precise locations in the protein

backbone. Fine-mapping of these effects was made possible by systematic "sequon

walking" experiments whereby a glycosylation site was introduced at every possible

position of a protein, as we demonstrated for Im7. We imagine that sequon walking on a

large, even proteome-wide, scale could provide access to datasets that might allow the

effects of glycosylation to be further generalized and predicted. Even though an unprecedentedly large number of intact glycoprotein variants (151 in total) were evaluated here, larger scale studies might require higher throughput techniques for resolving glycosylation efficiency such as recently described workflows that leverage mass spectrometry [45, 46]. Nonetheless, the fact that *N*-glycan attachment significantly increased the binding activity of several Im7 and scFv-HER2 variants suggests that glycomutagenesis may become a useful tool for custom engineering of proteins with advantageous biophysical and biological properties.

A unique aspect of the SSGM method is the ability to generate combinatorial glycosite libraries and comprehensively assess which positions in a protein can tolerate glycan installation. The three proteins studied here were found to be efficiently *N*-glycosylated at an unexpectedly large number of positions and in all types of secondary structure although loops tended to be more receptive to glycosylation, consistent with the observation that naturally occurring *N*-glycans also exist on all forms of secondary structure [31]. Closer inspection of the data revealed some additional trends. For instance, installing *N*-glycans in the center of α-helices negatively affected activity (*e.g.*, positions 19, 42, 72 in Im7) whereas those installed at the transition between different types of secondary structure and at turns between motifs promoted enhanced activity and, in some cases, stability (*e.g.*, positions 33, 49, 58, 59, 60, 61, 65, 67, 68, 69, 78, 80 in Im7). These findings generally agreed with the folding and stability effects contributed by $GlcNAc_2$ disaccharide attachment to discrete locations in Im7 [16] and also provide clues for why natural *N*-glycosylation sites occur with elevated frequency in turns and bends and especially at points of change in secondary structure and with low frequency within ordered helices [31]. Despite the overall agreement with previous studies, a few notable differences emerged. For example, in our hands, Im7 glycosylated at position 27 with the $GalNAc_5(Glc)GlcNAc$ heptasaccharide was more active but equally stable as its aglycosylated counterpart, whereas an EPL-derived Im7 modified with chitobiose at residue 27 was significantly more stable than unmodified Im7 (and activity data was not reported) [16]. Likewise, RNase A^{N34} glycosylated with $GalNAc_5(Glc)GlcNAc$ exhibited activity that was nearly identical to that of aglycosylated RNase A^{N34} (and wt RNase A), whereas the attachment of oligomannose glycans at N34 was previously observed to

reduce activity by more than threefold [47]. These discrepancies would suggest that distinct

glycan structures (*i.e.,* $GalNAc_5(Glc)GlcNAc$ versus $Man_{5-9}GlcNAc_2$ versus $GlcNAc_2$)

attached to the same site in RNase A have distinct effects. This is not entirely surprising

in light of studies that have measured different biological and/or biophysical effects

depending on the structure of the attached glycan [48]. Thus, in the future it will of interest

to adapt SSGM for use with alternative glycan structures, including for example

$Man_3GlcNAc_2$ or other human *N-* and *O-*linked glycans that have been engineered in *E.*

coli [32, 49, 50], so that the consequences of varying glycan structures at discrete locations

can be systematically investigated.

The studies performed here also provided insight on the possible timing and impact

of glycosylation with respect to the folding process. For instance, Im7 tolerated a glycan

at almost every position, even when the target asparagine side chain pointed inward and

was considered buried (*e.g.*, positions N7, N68, and N76). Because these buried

positions physically cannot be glycosylated by PglB when the target protein is in the folded

state, they must either be glycosylated co-translationally or during a fluctuation to a

partially unfolded state that provides access to that site. Then, after glycosylation,

because Im7 presumably cannot fold back into the native structure, it must adopt a

different conformation to accommodate the newly added glycan, which would be feasible

in light of the fact that Im7 is very flexible [37]. In the case of RNase A, several sites were

identified (*e.g.*, N34, N36) that could be efficiently glycosylated *in vivo* but underwent little

to no glycosylation *in vitro* (in the already folded state), providing clear evidence for glycan

installation prior to folding and in a manner that may resemble the co-translocational

process in mammalian cells [51]. The overall less efficient glycosylation seen for many

RNase A variants is also consistent with the protein adopting a more stable folded

structure compared to Im7 and providing less accessibility to buried sites.

Using the results of the SSGM studies, we also investigated explanatory factors

that could be used to predict *a priori* the functional effect of glycosylation from the

sequence. Here, we tested four simple structural metrics (secondary structure, distance

to the active site, burial of the glycosylation site in the native fold, and change in buried

surface area upon binding) and the change in energy of the protein alone and in complex

with its substrate upon addition of the glycan (estimated with Rosetta). Whereas RNase

A activity was more sensitive to sequon substitution than glycosylation, the effects of glycosylation on the function of Im7 and scFv-HER2 were quite interesting. Collectively, the results indicated that interactions between the glycan and the bound protein can alter binding activity (positively or negatively) and that enhanced binding likely arises from low-energy glycan conformations making favorable interactions with the binding partner. In the case of scFv-HER2, mutant N58 V_L exhibited significantly higher binding activity compared to wt scFv-HER2. While part of the increase was from the sequon substitution alone, perhaps from the additional contacts of the long Q57 side chain or from a stabilizing effect of the sequon on the CDR L2 loop (residues 51-57 in V_L), most of the effect was from the *N*-glycan itself. At residue N58 V_L, the heptameric glycan creates new contacts between scFv-HER2 and HER2-ED and buries more surface area upon binding. Glycans attached near (but not within) the binding site have previously been reported to increase antigen-binding affinity [52]. In fact, the Im7 variant that underwent the largest increase in binding activity upon glycosylation (at residue N58 in Im7) also acquired new contacts with its binding partner, E7, through the glycan, which strengthened binding activity by 3.5-fold. Thus, it is an intriguing possibility that binding affinity of a protein might be purposefully increased by rational introduction of *N*-glycans at predetermined locations, with SSGM serving as the discovery engine for precisely pinpointing the sites within a protein that lead to glycan-mediated affinity enhancement.

Materials and Methods

Strains and culture conditions. *E. coli* strain DH5α was used for all molecular biology, including plasmid construction, site-directed mutagenesis, and SSGM library construction. BL21(DE3) was used to purify ColE7 that was used to measure Im7 binding activity in ELISA format. All glycosylation studies were performed using *E. coli* strain CLM24 [53], which was initially grown at 37 °C in Luria–Bertani (LB) medium containing appropriate antibiotics at the following concentrations: 20 μg/mL chloramphenicol (Cm), 100 μg/mL trimethoprim (Tmp), and 50 μg/mL spectinomycin (Spec). When cells reached mid-log phase, protein expression was induced by adding 0.1 mM isopropyl-β-D-thiogalactoside (IPTG) and 0.2% (v/v) L-arabinose, after which cells were grown at 30 °C for 16–20 h.

Plasmid construction. For expression of the glycosylation machinery, plasmid pMAF10 encoding *Cj*PglB [53] along with either plasmid pMW07-pglΔB [20] or pMW07-pglΔBCDEF were used. The latter plasmid was constructed by deleting the *pglCDEF* genes from plasmid pMW07-pglΔB [20], resulting in a "minimal" *C. jejuni* glycan biosynthesis pathway that excluded the *pglB* gene encoding the OST and also the *pglCDEF* genes encoding enzymes for synthesis and transfer of bacillosamine as described previously [28]. This deletion can be complemented by *E. coli* WecA, a sugar-phosphate transferase that transfers GlcNAc phosphate to undecaprenol phosphate, therefore initiating LLO biosynthesis. It should be noted that while pMW07-pglΔB encodes the bacillosaome-related *pglCDEF* genes, we did not detect the presence of bacillosamine in any of the glycoforms produced by cells carrying this plasmid. A derivative of pMAF10 that encoded a catalytically inactive version of *Cj*PglB carrying two active-site mutations (D54N and E316Q) [20] was used as a negative control. The plasmids pTrc99S-YebF-Im7 and pTrc99S-YebF-Im7^{DQNAT} were constructed by inserting cloning cassettes YebF^{N24L}-XbaI-Im7-SalI-FLAG-6xHis and YebF^{N24L}-XbaI-Im7-BamHI-DQNAT-SalI-FLAG-6xHis, respectively, into the SacI and HindIII sites of pTrc99S [54]. The genes encoding RNase A and scFv-HER2 were PCR amplified from plasmids pTrc-ssDsbA-RNaseA [20] and pMAZ360–cIgG–Herceptin [55], respectively, and cloned into the cassette between XbaI and SalI sites in pTrc99S-YebF-Im7 or XbaI and BamHI sites in pTrc99S-YebF-Im7^{DQNAT}, replacing Im7 and Im7^{DQNAT}, respectively. The pTrc-spDsbA-POI plasmids (where POI corresponds to each of the proteins of interest, namely Im7, RNase A, and scFv-HER2) were cloned by one-step PCR integration of primers encoding the *E. coli* DsbA signal peptide (spDsbA) into each pTrc99S-YebF-POI plasmid as templates followed by Gibson assembly. PCR products were subjected to DpnI digestion to remove parental plasmid. The resulting PCR products were assembled by Gibson assembly and used to transform *E. coli* cells to obtain the desired plasmids. Plasmid pET28-ColE7 (H569A) was constructed by inserting DNA encoding the ColE7 H569A variant [56] bearing a C-terminal 6×His tag (Integrated DNA Technologies) into the NcoI and SalI sites of pET28a. All plasmids were confirmed by DNA sequencing at the Biotechnology Resource Center of the Cornell Institute of Biotechnology.

SSGM library construction. SSGM mutagenesis libraries were constructed by multiplex inverse PCR [30] followed by T4 ligation. Each of the pTrc99S-YebF-POI plasmids was used as template for PCR amplification using primer sets specifically designed such that the DNA sequence 5′-GAT CAG AAT GCG ACC-3′ was included in the 5′ end of every forward primer to enable substitution of the adjacent five amino acids with DQNAT. Prior to PCR, the forward primers were phosphorylated using T4 polynucleotide kinase (New England Biolabs) to facilitate T4 ligation later. PCR reactions were performed using Phusion polymerase (New England Biolabs), and the PCR products were gel-purified from the product mixtures with a ratio of 1:1 to get rid of non-specific PCR products. The resulting PCR products were self-assembled using T4 ligase (New England Biolabs) to obtain the desired SSGM plasmid libraries, which were subsequently used to transform highly competent DH5α cells and then isolated using a QIAprep Spin Miniprep Kit (Qiagen) according to manufacturer's instructions.

GlycoSNAP assay. Screening of SSGM libraries was performed using glycoSNAP as described previously [20]. Briefly, *E. coli* strain CLM24 carrying pMW07-pglΔB and pMAF10 was transformed with corresponding SSGM library plasmids, and the resulting transformants were grown on 150-mm LB-agar plates containing 20 μg/mL Cm, 100 μg/mL Tmp, and 50 μg/mL Spec overnight at 37 °C. The second day, nitrocellulose transfer membranes were cut to fit 150-mm plates and pre-wet with sterile phosphate-buffered saline (PBS) before placement onto LB-agar plates containing 20 μg/mL Cm, 100 μg/mL Tmp, 50 μg/mL Spec, 0.1 mM IPTG, and 0.2% (w/v) ʟ-arabinose. Library transformants were replicated onto 142-mm cellulose nitrate membrane filters (Whatman, 0.45 μm), which were then placed colony-side-up on transfer membranes and incubated at 30 °C for 16 h. The nitrocellulose transfer membranes were washed in Tris-buffered saline (TBS) for 10 min, blocked in 5% bovine serum albumin for 30 min and probed for 1 h with fluorescein-labeled SBA (Vector Laboratories, catalog # FL-1011) and Alexa Fluor 647® (AF647)-conjugated anti-His antibody (R&D Systems, catalog # IC0501R) or HER2-ED (R&D Systems, catalog # 10126-ER) that was conjugated with Alexa Fluor 647™ (AF647) (Thermo Fisher Scientific, catalog # A37573) following the manufacturer's instructions. All positive hits were re-streaked onto fresh LB-agar plates containing 20 μg/mL Cm, 100 μg/mL Tmp, 50 μg/mL Spec, and grown overnight at 37 °C. Individual

colonies were grown in liquid culture and subjected to DNA sequencing to confirm the

location of glycosites and to protein glycosylation analysis as described below.

Protein isolation. For Western blot analysis and protein activity assays, cell-free culture

supernatants were generated by subjecting 1.5 mL of cells that had been induced for 16

h to centrifugation at 13,4000 × g at 4 °C for 2 min. Periplasmic fractions were generated

by subjecting 3 mL of 16-h-induced cultures to centrifugation at 13,400 × g for 2 min. The

resulting pellets were resuspended in 300 μL of 0.4 M arginine and incubated at 4 °C for

1 h with gentle shaking. After centrifugation at 13,400 × g for 2 min, the supernatant

containing periplasmic extracts was collected. For stability assays, YebF-Im7, YebF-

RNase A, and YebF-scFv-HER2 variants were purified from supernatant fractions and

soluble lysate fractions. To prepare the latter, cells expressing YebF-RNase A variants

were harvested by centrifugation at 6000 × g at 4 °C for 20 min and the pellets were

resuspended in PBS buffer supplemented with 10-mM imidazole followed by cell lysis

using a Emulsiflex-C5 Homogenizer (Avestin) at 16,000–18,000 psi. The resulting lysate

was clarified by centrifugation at 15,000 × g for 30 min at 4 °C to collect the soluble

fraction. All soluble fractions, or supernatant fractions supplemented with 10-mM

imidazole, were then applied twice to a gravity flow column loaded with Ni-NTA resin at

room temperature and washed with PBS containing 20-mM imidazole until the

concentration was lower than 0.1 mg/mL. Proteins were eluted in 2.5 mL of PBS with 250

mM imidazole. The eluted proteins were desalted using PD10 Desalting Columns (GE

Healthcare) and stored at 4 °C.

To produce ColE7 for ELISA experiments, an overnight culture BL21(DE3) cells

carrying plasmid pET28a-ColE7 (H569A) was used to inoculate 1 L of LB supplemented

with 50 μg/mL kanamycin. Cells were grown at 37 °C until mid-log phase and then were

induced with 0.1 mM IPTG for 16 h at 16 °C before being harvested. Following

centrifugation at 10,000× g, pellets were resuspended in PBS buffer supplemented with

10-mM imidazole and lysed at 16,000–18,000 psi using an Emulsiflex-C5 homogenizer

(Avestin). The lysate was clarified by centrifugation at 15,000 × g for 30 min at 4 °C and

the collected soluble fraction was mixed with Ni-NTA resin for 2 h at 4 °C. The mixture

was then applied to a gravity flow column and washed with 5 column volumes of PBS

containing 20 mM imidazole. Proteins were eluted in 4 column volumes of PBS with 250-

mM imidazole. The eluted protein was desalted and concentrated to 5 mg/mL in PBS

buffer using Ultra Centrifugal Filters with 10-kDa molecular weight cut-off (Amicon®) and

stored at 4 °C.

Western blotting analysis. Supernatant or periplasmic fractions were diluted 3:1 in 4×

Laemmli sample buffer (Bio-Rad) and were boiled at 100 °C for 10 min. The treated

samples were subjected to SDS-polyacrylamide gel electrophoresis on 10% Mini-

PROTEAN® TGX™ Precast Protein Gels (Bio-Rad). The separated protein samples

were then transferred to nitrocellulose membranes. Following transfer, the membranes

were blocked with 5% milk (w/v) in TBST (TBS, 0.1% Tween 20) and were probed with

horseradish peroxidase (HRP) conjugated anti-His antibody (Abcam, catalog # ab1187)

or the *C. jejuni* heptasaccharide glycan-specific antiserum hR6 for 1 h. For the latter, goat

anti-rabbit IgG (HRP) (Abcam, catalog # ab205718) was used as the secondary antibody

to detect hR6 antiserum. After washing three times with TBST for 10 min, the membranes

were visualized using a ChemiDoc™ MP Imaging System (Bio-Rad).

***In vitro* glycosylation.** Methods for purification of *C. jejuni* PglB and isolation of LLOs

from glycoengineered *E. coli* were described previously [57]. *In vitro* glycosylation was

carried out in 30-µL reactions containing either 20 µL of supernatant fraction containing

aglycosylated YebF-Im7 or 20 µL of periplasmic fraction containing YebF-RNase A, 2 µg

of purified *Cj*PglB, and 5 µg extracted LLOs in *in vitro* glycosylation buffer (10-mM

HEPES, pH 7.5, 10-mM $MnCl_2$, and 0.1% (w/v) *n*-dodecyl-β-D-maltoside (DDM)).

Reaction mixtures were incubated at 30 °C for 16 h and stopped by adding 10 µL of 4×

Laemmli sample buffer containing 5% β-mercaptoethanol followed by boiling at 100 °C

for 15 min, after which they were subjected to Western blot analysis.

ELISA. Binding activity for Im7 and scFv-HER2 was determined by standard ELISA.

Briefly, Costar 96-well ELISA plates (Corning) were coated overnight at 4 °C with 50 µL

of 5 µg/mL purified ColE7 in 0.05-M sodium carbonate buffer (pH 9.6) for Im7 variants

and 50 µL of 0.2 µg/mL HER2-ED (Sino Biological, catalog # 10004-HCCH) in PBS buffer

for scFv-HER2 variants. After blocking with 5% (w/v) non-fat milk in PBS for 1 h at room

temperature, the plates were washed three times with PBST (PBS, 0.05% (v/v) Tween-

20) and incubated with serially diluted aglycosylated and glycosylated YebF-Im7 and

YebF-scFv-HER2 glycovariants for 1 h at room temperature. After washing three times

with PBST, 50 µL of 1:2,500-diluted HRP-conjugated anti-DDDK tag antibody (Abcam,

catalog # ab49763) for Im7 variants or 50 µL of 1:5,000-diluted HRP-conjugated anti-

6×His tag antibody (Abcam, catalog # ab1187) for scFv-HER2 variants, both in 1% PBST,

was added to each well for 1 h. Plates were washed three times and then developed using

50 µL 1-Step Ultra TMB-ELISA substrate solution (ThermoFisher).

RNase A activity assay. The enzymatic activity of RNase A variants was assayed using

RNaseAlert[®]-1 Kit (Integrated DNA Technologies) according to the manufacturer's

protocol. Each of the 80-times-diluted supernatant samples were normalized to have an

OD_{600} equivalent to the positive control strain expressing wt RNase A. Samples were then

mixed with 20 pmol of RNase A substrate and 10 µL of 10× RNaseAlert Buffer and

incubated in RNase-free black 96 well microplates (Fisher) at 37 °C for 30 min.

Fluorescence values were measured at 490 nm/520 nm excitation/emission wavelengths.

Thermal stability analysis. Far-UV CD spectroscopy of purified Im7 (50-mM sodium

phosphate, 400-mM sodium sulfate, pH 7.4) as a function of temperature was carried out

in a 0.1-cm cuvette on a spectropolarimeter. Far-UV CD spectra were acquired between

200 nm and 260 nm with a step resolution of 1 nm. Melting temperatures of purified

glycovariants was determined using high-throughput DSF as previously described [58].

Briefly, 5–10 µg of proteins were mixed with Protein Thermal Shift[TM] Buffer and Protein

Thermal Shift[TM] Dye purchased as Protein Thermal Shift Dye Kit[TM] (Thermo Fischer

Scientific) according to manufacturer's instructions. A melting curve was generated by

monitoring fluorescence at 465 nm/610 nm excitation/emission wavelengths while

increasing temperature from 10 °C to 90 °C at a rate of 0.06 °C/s on an Applied Biosystem

ViiA 7 instrument (Life Technologies). To calculate T_m values, the collected data were

analyzed by nonlinear regression analysis using the Boltzmann equation in Prism 8.4.2

(GraphPad).

Mass spectrometry analysis of protein glycosylation. Proteins samples (~2 µg) were

separated by SDS-PAGE gel and bands corresponding to glycosylated YebF-Im7[DQNAT]

were excised and subjected to in-gel digestion by trypsin followed by extraction of tryptic

peptides essentially as described [50]. Gel slices were washed and then destained by

treatment with a 1:1 mixture of acetonitrile (Fisher Chemical) and 50 mM aqueous

NH_4HCO_3 followed by treatment with 100% acetonitrile. After destaining, gel pieces were

reduced and alkylated with 10 mM dithiothreitol (Roche) and 50 mM iodoacetamide (Acros Organics). The glycoproteins were directly digested by adding trypsin (w/w = 1:10) in 50 mM NH_4HCO_3 to the gel pieces and incubating overnight at 37 °C. The tryptic peptides were extracted by 50% acetonitrile with 5% formic acid (Fisher Chemical) and 75% acetonitrile with 5% formic acid. Extracted peptides were pooled and dried by SpeedVac. The tryptic peptides were suspended in 24 μL of 0.5% formic acid and 10 μL was injected into an UltiMate3000 RSLCnano (Dionex) coupled to an Orbitrap Fusion mass spectrometer (Thermo-Fisher Scientific) as described [59] with slight modifications. The peptides were injected onto a PepMap C18 RP nano trapping column (5 μm, 100 μm i.d x 20 mm) at 20 μL/min flow rate for rapid sample loading, and separated on an Acclaim PepMap C18 nano column (3 μm, 75 μm x 25cm, Thermo Fisher Scientific). The tryptic peptides were eluted in a 90 min gradient of 5% to 23% to 35% solvent B (95% acetonitrile, 0.1% formic acid) corresponding to 3 to 73 to 93 min, respectively, at 300 nL/min. The 90-min gradient was followed by a 9-min ramping to 90% B, a 9-min hold at 90% B and quick switch to 5% B and 95% solvent A (2% acetonitrile, 0.1% formic acid) for 1 min. The column was re-equilibrated with 95% A for 25 min prior to the next run. The Orbitrap Fusion was operated in positive ion mode with nanospray voltage set at 1.7 kV and source temperature at 275 °C. The MS survey scan was acquired at a resolving power of 120,000 (FWHM at m/z 200) across m/z 350-1800, which was followed by a "top speed" data-dependent electron-transfer dissociation (ETD) MS/MS scan (cycle time of 4 s) supplemented with higher-energy collision dissociation (EThcD) fragmentation workflow for precursor peptides with 3–6 charges. EThcD fragmentation was acquired using calibrated charge-dependent ETD parameters supplemented with 15% collisional energy in ion trap detector with Automatic Gin Control (AGC) of 3e4 and maximum injection time of 118 s.

Data analysis was performed by Byonic v3.6 (Protein Metrics) searching software against an *E. coli* database containing the YebF-Im7 protein, and an in-house generated *N*-linked glycan database with additional diBacNAc-containing glycans. The peptide search parameters were as follows: two missed cleavage for full trypsin digestion with fixed carbamidomethyl modification of cysteine and Q to Pyro-E on N-terminal Q, variable modifications of methionine oxidation and deamidation on asparagine/glutamine

residues. The peptide mass tolerance was 10 ppm and fragment mass tolerance values

for EThcD spectra was 0.6 Da. Both the maximum number of common and rare

modifications were set at two. Identified peptides and glycopeptides were filtered for a

[log Prob] value >3 and glycopeptide mass error <5 ppm. The search results were

exported to excel files for final analysis and graphical presentation.

Protein structure preparation. Initial coordinates of structures used for analysis were

obtained from the Protein Data Bank at RCSB.org as follows: the Im7–E7 complex (PDB:

2JBG), RNAse A bound to four-nucleotide-long DNA (PDB: 1RCN, DNA: ATAA) and the

scFv-Her2–ErbB2 complex (PDB: 3WSQ). For RNase A, we modeled an eight-

nucleotide-long RNA at the same binding site of the DNA. We used X3DNA [60] to identify

nucleotide arrangement parameters from DNA, as a template, to model RNA. We

replaced the thymine residue of bound DNA with its RNA counterpart uracil. Since an

initial analysis showed that four-nucleotide-long RNA is small for an interface-interaction

calculation, we added two extra nucleotides to both ends of the modeled RNA to make it

eight nucleotides long. Finally, to remove any unwanted clashes coming from the base

sugar (deoxyribose to ribose) and the nucleotide change (thymine to uracil), we used the

Rosetta relax protocol with a restraint on protein backbone atoms to generate a pool of

100 RNA–protein structures ("decoys"). By manual inspection, we removed decoys

showing RNAs outside of the binding site of RNase A. From the pool of remaining

structures, we selected a candidate with the best Rosetta score for our study.

Geometric calculations. Secondary structure was identified using the DSSP protocol in

PyRosetta [61]. Burial of a glycosylated residue was represented by a count of C_β atoms

within 8.5 Å from the C_β atom of the selected amino acid residue (C_α for alanine). The

distance of a glycosylated residue from its binding partner was estimated by the C_α–C_α

distance from the glycosylated residue to the closest residue of the binding partner. For

RNase A, the distance of glycosylated residue from the active site residue His-119 was

used instead, as RNase A does not bind to a protein. This distance was used for both the

"activity ratio" and the "activity improvement probability" analyses. Interface solvent-

accessible surface area (SASA) was measured using the Rosetta interface analyzer [62].

Rosetta measures. Rosetta's REF15 score function [22] was used to estimate the stability

of a protein–protein complex. An upper-bound cutoff of 100 REU is used to limit the effect

of bad conformers on the correlation, while for Im7, this value was 60 REU. To represent the binding energy of a complex, the interface score was calculated with the Rosetta Interface Analyzer [62] by subtracting the score of the separated monomers from the complex.

Rosetta protocols for sequon substitution and glycomutagenesis. We used the RosettaCarbohydrate framework [23] and a new glycomutagenesis protocol to input a wt PDB file and generate all possible sequon-substituted variants and glycosylated variants. Coordinate files after sequon substitution and side chain optimization steps were used as aglycosylated variants for analysis. Calculations were carried out in Rosetta release version 2020.06 and PyRosetta4 release 247 (www.rosettacommons.org). The command line for the glycomutagenesis calculation was:

```
glycomutagenesis.linuxgccrelease -in:file:s <PDB> -include_sugars
-nstruct <length of sequence selected for glycomutagenesis>
-n_cycles 100 -out:path:pdb ./output -out:path:score ./output
```

The command line for the relaxation runs was:

```
relax.linuxgccrelease -s <PDB> -nstruct 100
-relax:default_repeats 5
-relax:bb_move false
-out:path:pdb ./output_relax -out:path:score ./output_relax
```

The command for analyzing aglycosylated structures was:

```
InterfaceAnalyzer.default.linuxgccrelease -s <PDB>
-interface A_B #(For chain A and B)
-out:path:pdb ./output -out:path:score ./output
```

and the command for glycosylated structures was:

```
InterfaceAnalyzer.default.linuxgccrelease -s <pdb file>
-interface AB_C #(For [protein A + Glycan B] and Protein C)
-fixedchains A B #(For Protein A + Glycan B)
-out:path:pdb ./output -out:path:score ./output
```

Acknowledgements. We thank Markus Aebi for providing strain CLM24 and hR6 serum used in this work. The authors also thank Mike Jewett, Milan Mrksich, Eric Sundberg, Sophia Hulbert, José-Marc Techner, Weston Kightlinger, Liang Lin, Jessica Stark, and Sai Pooja Mahajan for helpful discussions of the manuscript. This work was supported by the Defense Threat Reduction Agency (HDTRA1-15-10052 and HDTRA1-20-10004 to

M.P.D.), National Science Foundation (CBET-1159581, CBET-1264701, CBET-1936823 to M.P.D.), and National Institutes of Health (1R01GM137314 to M.P.D., 1R01GM127578 to M.P.D. and J.J.G., and 1S10 OD017992-01 to S.Z.). The work was also supported by seed project funding (to M.P.D.) through the National Institutes of Health-funded Cornell Center on the Physics of Cancer Metabolism (supporting grant 1U54CA210184). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health. T.J. was supported by a Royal Thai Government Fellowship and also a Cornell Fleming Graduate Scholarship.

Author Contributions. M.L. and X.Z. designed and performed all research, analyzed all data, and wrote the paper. S.S., T.J., I.K., J.B., E.C.C., J.W.L., and J.J.G. designed and performed research, and analyzed data. M.P.D. directed research, analyzed data, and wrote the paper.

Data availability. All data generated or analyzed during this study are included in this article (and its supplementary information) or are available from the corresponding authors on reasonable request.

Competing Interests. M.P.D. has a financial interest in Ajuta Therapeutics, Inc., Glycobia, Inc., SwiftScale Biologics, Inc., and Versatope, Inc. M.P.D.'s interests are reviewed and managed by Cornell University in accordance with their conflict of interest policies. J.J.G. is an unpaid board member of the Rosetta Commons. Under institutional participation agreements between the University of Washington, acting on behalf of the Rosetta Commons, Johns Hopkins University may be entitled to a portion of revenue received on licensing Rosetta software including some methods developed in this article. As a member of the Scientific Advisory Board, J.J.G. has a financial interest in Cyrus Biotechnology. Cyrus Biotechnology distributes the Rosetta software, which may include methods mentioned in this article. J.J.G.'s arrangements have been reviewed and approved by the Johns Hopkins University in accordance with its conflict of interest policies. All other authors declare no competing interests.

References

1.   Khoury, G.A., Baliban, R.C. & Floudas, C.A. Proteome-wide post-translational
modification statistics: frequency analysis and curation of the swiss-prot
database. *Sci Rep* **1** (2011).
2.   Walsh, C.T., Garneau-Tsodikova, S. & Gatto, G.J., Jr. Protein posttranslational
modifications: the chemistry of proteome diversifications. *Angew Chem Int Ed*
Engl **44**, 7342-7372 (2005).
3.   Abu-Qarn, M., Eichler, J. & Sharon, N. Not just for Eukarya anymore: protein
glycosylation in Bacteria and Archaea. *Curr Opin Struct Biol* **18**, 544-550 (2008).
4.   Varki, A. Biological roles of glycans. *Glycobiology* **27**, 3-49 (2017).
5.   Mitra, N., Sinha, S., Ramya, T.N. & Surolia, A. N-linked oligosaccharides as
outfitters for glycoprotein folding, form and function. *Trends Biochem Sci* **31**, 156-
163 (2006).
6.   van de Bovenkamp, F.S. et al. Adaptive antibody diversification through N-linked
glycosylation of the immunoglobulin variable region. *Proc Natl Acad Sci U S A*
115, 1901-1906 (2018).
7.   Beckham, G.T. et al. Harnessing glycosylation to improve cellulase activity. *Curr*
Opin Biotechnol **23**, 338-345 (2012).
8.   Wang, L.X., Tong, X., Li, C., Giddens, J.P. & Li, T. Glycoengineering of antibodies
for modulating functions. *Annu Rev Biochem* **88**, 433-459 (2019).
9.   Berti, F. & Adamo, R. Antimicrobial glycoconjugate vaccines: an overview of classic
and modern approaches for protein modification. *Chem Soc Rev* **47**, 9015-9025
(2018).
10. Van Landuyt, L., Lonigro, C., Meuris, L. & Callewaert, N. Customized protein
glycosylation to improve biopharmaceutical function and targeting. *Curr Opin*
Biotechnol **60**, 17-28 (2019).
11. Rich, J.R. & Withers, S.G. Emerging methods for the production of homogeneous
human glycoproteins. *Nat Chem Biol* **5**, 206-215 (2009).
12. Bosques, C.J., Tschampel, S.M., Woods, R.J. & Imperiali, B. Effects of
glycosylation on peptide conformation: a synergistic experimental and
computational study. *J Am Chem Soc* **126**, 8421-8425 (2004).
13. Shental-Bechor, D. & Levy, Y. Effect of glycosylation on protein folding: a close look
at thermodynamic stabilization. *Proc Natl Acad Sci U S A* **105**, 8256-8261 (2008).
14. Buskas, T., Ingale, S. & Boons, G.J. Glycopeptides as versatile tools for
glycobiology. *Glycobiology* **16**, 113R-136R (2006).
15. Payne, R.J. & Wong, C.H. Advances in chemical ligation strategies for the
synthesis of glycopeptides and glycoproteins. *Chem Commun (Camb)* **46**, 21-43
(2010).
16. Chen, M.M. et al. Perturbing the folding energy landscape of the bacterial immunity
protein Im7 by site-specific N-linked glycosylation. *Proc Natl Acad Sci U S A* **107**,
22528-22533 (2010).
17. Morrison, K.L. & Weiss, G.A. Combinatorial alanine-scanning. *Curr Opin Chem Biol*
5, 302-307 (2001).
18. Weiss, G.A., Watanabe, C.K., Zhong, A., Goddard, A. & Sidhu, S.S. Rapid mapping
of protein functional epitopes by combinatorial alanine scanning. *Proc Natl Acad*
Sci U S A **97**, 8950-8954 (2000).

19. Gregoret, L.M. & Sauer, R.T. Additivity of mutant effects assessed by binomial mutagenesis. *Proc Natl Acad Sci U S A* **90**, 4246-4250 (1993).

20. Ollis, A.A., Zhang, S., Fisher, A.C. & DeLisa, M.P. Engineered oligosaccharyltransferases with greatly relaxed acceptor-site specificity. *Nat Chem Biol* **10**, 816-822 (2014).

21. Zhang, G., Brokx, S. & Weiner, J.H. Extracellular accumulation of recombinant proteins fused to the carrier protein YebF in Escherichia coli. *Nat Biotechnol* **24**, 100-104 (2006).

22. Alford, R.F. et al. The Rosetta All-Atom Energy Function for Macromolecular Modeling and Design. *J Chem Theory Comput* **13**, 3031-3048 (2017).

23. Labonte, J.W., Adolf-Bryfogle, J., Schief, W.R. & Gray, J.J. Residue-centric modeling and design of saccharide and glycoconjugate structures. *J Comput Chem* **38**, 276-287 (2017).

24. Mansell, T.J., Guarino, C. & DeLisa, M.P. Engineered genetic selection links in vivo protein folding and stability with asparagine-linked glycosylation. *Biotechnol J* **8**, 1445-1451 (2013).

25. Dennis, C.A. et al. A structural comparison of the colicin immunity proteins Im7 and Im9 gives new insights into the molecular determinants of immunity-protein specificity. *Biochem J* **333 (Pt 1)**, 183-191 (1998).

26. Ko, T.P., Liao, C.C., Ku, W.Y., Chak, K.F. & Yuan, H.S. The crystal structure of the DNase domain of colicin E7 in complex with its inhibitor Im7 protein. *Structure* **7**, 91-102 (1999).

27. Fisher, A.C. et al. Production of secretory and extracellular N-linked glycoproteins in Escherichia coli. *Appl Environ Microbiol* **77**, 871-881 (2011).

28. Yates, L.E. et al. Glyco-recoded Escherichia coli: Recombineering-based genome editing of native polysaccharide biosynthesis gene clusters. *Metab Eng* **53**, 59-68 (2019).

29. Wacker, M. et al. N-linked glycosylation in Campylobacter jejuni and its functional transfer into E. coli. *Science* **298**, 1790-1793 (2002).

30. Kanwar, M., Wright, R.C., Date, A., Tullman, J. & Ostermeier, M. Protein switch engineering by domain insertion. *Methods Enzymol* **523**, 369-388 (2013).

31. Petrescu, A.J., Milac, A.L., Petrescu, S.M., Dwek, R.A. & Wormald, M.R. Statistical analysis of the protein environment of N-glycosylation sites: implications for occupancy, structure, and folding. *Glycobiology* **14**, 103-114 (2004).

32. Valderrama-Rincon, J.D. et al. An engineered eukaryotic protein glycosylation pathway in Escherichia coli. *Nat Chem Biol* **8**, 434-436 (2012).

33. Kowarik, M. et al. N-linked glycosylation of folded proteins by the bacterial oligosaccharyltransferase. *Science* **314**, 1148-1150 (2006).

34. Silverman, J.M. & Imperiali, B. Bacterial N-glycosylation efficiency Is dependent on the structural context of target sequons. *J Biol Chem* **291**, 22001-22010 (2016).

35. Juraja, S.M. et al. Engineering of the Escherichia coli Im7 immunity protein as a loop display scaffold. *Protein Eng Des Sel* **19**, 231-244 (2006).

36. Lavinder, J.J., Hari, S.B., Sullivan, B.J. & Magliery, T.J. High-throughput thermal scanning: a general, rapid dye-binding thermal shift screen for protein engineering. *J Am Chem Soc* **131**, 3794-3795 (2009).

37. Whittaker, S.B., Spence, G.R., Gunter Grossmann, J., Radford, S.E. & Moore, G.R. NMR analysis of the conformational properties of the trapped on-pathway folding intermediate of the bacterial immunity protein Im7. *J Mol Biol* **366**, 1001-1015 (2007).

38. Williams, R.L., Greene, S.M. & McPherson, A. The crystal structure of ribonuclease B at 2.5-A resolution. *J Biol Chem* **262**, 16020-16031 (1987).

39. Arnold, U. & Ulbrich-Hofmann, R. Kinetic and thermodynamic thermal stabilities of ribonuclease A and ribonuclease B. *Biochemistry* **36**, 2166-2172 (1997).

40. Grafl, R., Lang, K., Vogl, H. & Schmid, F.X. The mechanism of folding of pancreatic ribonucleases is independent of the presence of covalently linked carbohydrate. *J Biol Chem* **262**, 10624-10629 (1987).

41. Chen, M.M., Glover, K.J. & Imperiali, B. From peptide to protein: comparative analysis of the substrate specificity of N-linked glycosylation in C. jejuni. *Biochemistry* **46**, 5579-5585 (2007).

42. Panov, K.I. et al. Ribonuclease A mutant His119 Asn: the role of histidine in catalysis. *FEBS Lett* **398**, 57-60 (1996).

43. van de Bovenkamp, F.S. et al. Variable domain N-linked glycans acquired during antigen-specific immune responses can contribute to immunoglobulin G antibody stability. *Front Immunol* **9**, 740 (2018).

44. Krause, J.C. et al. An insertion mutation that distorts antibody binding site architecture enhances function of a human antibody. *mBio* **2**, e00345-00310 (2011).

45. Kightlinger, W. et al. Design of glycosylation sites by rapid synthesis and analysis of glycosyltransferases. *Nat Chem Biol* **14**, 627-635 (2018).

46. Techner, J.M. et al. High-throughput synthesis and analysis of intact glycoproteins using SAMDI-MS. *Anal Chem* **92**, 1963-1971 (2020).

47. Rudd, P.M. et al. Glycoforms modify the dynamic stability and functional activity of an enzyme. *Biochemistry* **33**, 17-22 (1994).

48. Hanson, S.R. et al. The core trisaccharide of an N-linked glycoprotein intrinsically accelerates folding and enhances stability. *Proc Natl Acad Sci U S A* **106**, 3131-3136 (2009).

49. Kightlinger, W. et al. A cell-free biosynthesis platform for modular construction of protein glycosylation pathways. *Nat Commun* **10**, 5404 (2019).

50. Natarajan, A. et al. Engineering orthogonal human O-linked glycoprotein biosynthesis in bacteria. *Nat Chem Biol* **16**, 1062-1070 (2020).

51. Ruiz-Canada, C., Kelleher, D.J. & Gilmore, R. Cotranslational and posttranslational N-glycosylation of polypeptides by distinct mammalian OST isoforms. *Cell* **136**, 272-283 (2009).

52. van de Bovenkamp, F.S., Hafkenscheid, L., Rispens, T. & Rombouts, Y. The emerging importance of IgG Fab glycosylation in immunity. *J Immunol* **196**, 1435-1441 (2016).

53. Feldman, M.F. et al. Engineering N-linked protein glycosylation with diverse O antigen lipopolysaccharide structures in Escherichia coli. *Proc Natl Acad Sci U S A* **102**, 3016-3021 (2005).

54. Ollis, A.A. et al. Substitute sweeteners: diverse bacterial oligosaccharyltransferases with unique N-glycosylation site preferences. *Sci Rep* **5**, 15237 (2015).

55. Robinson, M.P. et al. Efficient expression of full-length antibodies in the cytoplasm of engineered bacteria. *Nat Commun* **6**, 8072 (2015).

56. Kortemme, T. et al. Computational redesign of protein-protein interaction specificity. *Nat Struct Mol Biol* **11**, 371-379 (2004).

57. Jaroentomeechai, T. et al. Single-pot glycoprotein biosynthesis using a cell-free transcription-translation system enriched with glycosylation machinery. *Nat Commun* **9**, 2686 (2018).

58. Ericsson, U.B., Hallberg, B.M., Detitta, G.T., Dekker, N. & Nordlund, P. Thermofluor-based high-throughput stability optimization of proteins for structural studies. *Anal Biochem* **357**, 289-298 (2006).

59. Song, M. et al. IRE1alpha-XBP1 controls T cell function in ovarian cancer by regulating mitochondrial activity. *Nature* **562**, 423-428 (2018).

60. Lu, X.J. & Olson, W.K. 3DNA: a software package for the analysis, rebuilding and visualization of three-dimensional nucleic acid structures. *Nucleic Acids Res* **31**, 5108-5121 (2003).

61. Chaudhury, S., Lyskov, S. & Gray, J.J. PyRosetta: a script-based interface for implementing molecular modeling algorithms using Rosetta. *Bioinformatics* **26**, 689-691 (2010).

62. Stranges, P.B. & Kuhlman, B. A comparison of successful and failed protein interface designs highlights the challenges of designing buried hydrogen bonds. *Protein Sci* **22**, 74-82 (2013).