Interconnected assembly factors regulate the biogenesis of mitoribosomal large subunit in trypanosomes

Victor Tobiasson[1†], Ondřej Gahura[2†], Shintaro Aibara[1,3†], Rozbeh Baradaran[1], Alena Zíková[2*], Alexey Amunts[1*]

[1] Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, 17165 Solna, Sweden.

[2] Institute of Parasitology, Biology Centre, Czech Academy of Sciences, Ceske Budejovice, Czech Republic, Faculty of Science, University of South Bohemia, Ceske Budejovice, Czech Republic.

[3] Current address: Department of Molecular Biology, Max-Planck-Institute for Biophysical Chemistry, Göttingen, Germany.

† The authors contributed equally to this work.

* correspondence to: azikova@paru.cas.cz, amunts@scilifelab.se

Abstract

Mitoribosomes synthesize essential proteins encoded in the mitochondrial genome. They consist of the ribosomal RNA and protein components, coordinated assembly of which is critical for function. Divergent mitoribosomes with reduced RNA and increased protein mass were described in *Trypanosoma brucei*, a mammalian parasite causing African Trypanosomiases. Here, we report that the large subunit of *T. brucei* mitoribosome requires at least 15 biogenesis factors, which we characterized in complex with an assembly intermediate using cryo-EM. These assembly factors form a protein network that spans over 180 Å and connects between functional regions, allowing to coordinate their maturation. The conserved proteins mt-RbgA and mt-EngA are stably bound together at the subunit interface and mitochondrial acyl-carrier proteins play a structural role in constraining the L1 stalk. The characterized assembly intermediate forms an extensive network connecting between functionally important regions of the mitoribosome.

Impact statement

Biogenesis of *Trypanosoma brucei* mitoribosomal large subunit requires a protein network of assembly factors that connects between its functional regions.

Introduction

Mitoribosomes differ from bacterial and cytosolic ribosomes in the ribosomal RNA (rRNA) and protein content, overall size, and structure [1, 2]. In mitochondria of *Trypanosoma brucei*, the mitoribosome consists of over a hundred components, and the ratio of protein to rRNA is unusually high [3, 4]. Since the rRNA forms a compact core of the mitoribosome, and proteins are mostly peripherally associated, an architecture based on the reduced rRNA and supernumerary mitoribosomal proteins suggests a specialized assembly pathway. Indeed, structural characterization of the *T. brucei* mitoribosomal small subunit (mtSSU) provided insights into its assembly pathway that have not been previously known [5]. Compared to the mtSSU, no information is available on the mitoribosomal large subunit (mtLSU) assembly in *T. brucei*.

The mtLSU accommodates the peptidyl transferase center (PTC) that forms peptide bonds between amino acids, tRNA binding sites, the L7/L12 stalk that is responsible for the recruitment of translation factors, the L1 stalk, the central protuberance (CP) that facilitates communication between various functional sites, and exit tunnel for egress of synthesized protein. In bacteria, our understanding of LSU assembly comes from characterization of its final maturation stages [6-8], studies on incomplete LSU particles as a result of protein depletion [9], and *in vitro* reconstitution studies with purified ribosomal RNA and protein components [10]. These studies identified different LSU precursors with assembly factors that play key roles by binding to rRNA components [11]. In mitochondria of *T. brucei*, the mtLSU is lacking many of the rRNA components involved in the canonical pathways. Whether dedicated assembly factors have evolved is not known.

Here, we determine the cryo-EM structure of a native *T. brucei* mtLSU assembly intermediate (pre-mtLSU) with 15 assembly factors. It suggests a biogenesis relying on an extensive protein network formed by assembly factors that connects the maturation of the PTC, L1 and L7/L12 stalks, as well as the CP. Comparison with two bacterial assembly intermediates [12, 13] provides insights into the conserved GTPases EngA and RbgA bound at the subunit interface.

Results and Discussion

Structural determination and composition of the native pre-mtLSU complex

We used a procyclic strain grown in low-glucose medium that maintains translationally active mitochondria. Mitoribosomal complexes were purified directly from *T. brucei* mitochondria. During image processing, in addition to intact monosomes, we detected a pool of free subunits. We focused the analysis on this population and isolated pre-mtLSUs through 3D classification.

896,263 particles were picked using Warp [14], and further processed using RELION [15]. We performed reference-based 3D classification with references generated from a preliminary classification of a screening data set. This resulted in 207,788 particles corresponding to the mtLSU shape but distinct from that of a mature mtLSU of which we found 152,816 particles.

Refinement of angular assignments and subsequent classification using fine-angular searches
with a solvent mask identified 32,339 pre-mtLSU particles (Figure 1-figure supplement 1). To
improve the angles further, the particles were subjected to masked auto-refinement. Following
the CTF refinement, we obtained a reconstruction of a pre-mtLSU that likely reflects a stable
premature complex. This is evidenced by the presence of densities corresponding to conserved
ribosomal assembly factors.

The cryo-EM reconstruction was refined to 3.50 Å resolution (Figure 1-figure supplement 1).
This allowed to build a ~2.2 MDa complex and assign 15 assembly factors, as well as additional
mitoribosomal proteins, directly from the density (Figure 1, Figure 1-figure supplement 2). Three
distinct features define the overall premature complex: 1) the rRNA domain V is well resolved
and the subunit interface consists almost entirely of newly identified assembly factors associated
with two conserved GTPases; 2) the L7/L12 stalk proteins are missing, and its rRNA platform is
not folded, instead assembly factors occupy similar positions; 3) the CP is absent, and the L1
stalk is shifted inward ~30 Å. Due to these features, compositional and conformational changes
are required for the maturation of the pre-mtLSU. In terms of the mitoribosomal proteins, 18
previously identified proteins are missing from the structure of the pre-mtLSU. Seven of these
have bacterial homologs (uL10m, uL12m, uL16m, bL27m, bL31m, bL33m and bL36m) and the
rest are mitochondria specific (Figure 1, Figure 1-figure supplement 3). Additionally, we
assigned sequences to two previously unidentified mtLSU proteins, which we named mL109 and
mL110, and observed the presence of mitoribosomal protein uL14m which was not modeled in
the mtLSU (Figure 1-figure supplement 2).

Following the previously identified mitoribosomal small subunit assembly factors [5], we adopt
a similar nomenclature for the mitochondria specific large subunit factors. Therefore, we
reference them as *T. brucei* Large subunit Assembly Factor(s) (TbLAF1-10), while proteins with
bacterial homologs are referred to as their bacterial names with the prefix "mt-".

The identified assembly factors of the mitoribosome include two homologs of bacterial GTPase
assembly factors mt-EngA and mt-RbgA, a homolog of the ribosome silencing factor mt-RsfS, a
DEAD-box helicase (TbLAF10), two pseudouridinases (TbLAF4 and TbLAF6), a
methyltransferase (TbLAF5), two copies of the mitochondrial acyl-carrier protein (mt-ACP), a
LYR-motif-containing protein L0R8F8 (TbLAF1), and other lineage-specific proteins with
previously unassigned functions (TbLAF2, 3, 7, 8 and 9). In the model, we included only the
parts for which clear secondary structure is apparent, and other regions with only partial density
visible were modeled as UNK1-13 (Supplementary file 2).

Figure 1. Structure of *T. brucei* pre-mtLSU with 15 assembly factors. Left, the overall
modeled structure of the pre-mtLSU (white, rRNA blue) with models of assembly factors
(shades of purple) covering the subunit interface, CP, L7/L12 stalk and connecting to the L1
stalk. Right, structure of the mature mtLSU (PDB ID 6HIX) with 18 additional mitoribosomal
proteins (shades of orange) absent from pre-mtLSU.

GTPase mt-RbgA is structurally linked to the mitoribosomal core via specific assembly linkers

We started the structural analysis by searching for similar assembly intermediate architectures in
bacterial counterparts. Particles with an absent CP were reported previously in RbgA-depleted
Bacillus subtilis cells. RbgA was then added *in vitro* and shown to bind to the complex, which
identified its role as an assembly factor [13]. RbgA is from the Ras GTPase family containing a
typically low intrinsic GTPase activity, which is increased in the presence of mature LSU
subunits [16]. It has N-terminal GTPase domain and C-terminal helical domain, which forms a
five-helix bundle [17]. In the pre-mtLSU structure, we found a conserved mt-RbgA homolog.

Studies in yeasts reported that deletion of mt-RbgA (yeast Mtg1) results in respiration deficiency
[18]. In *B. subtilis*, where this assembly factor is essential, the LSU:RbgA complex showed that
the N-terminal domain overlaps with rRNA H69 and H71, and the C-terminal helical domain
interacts with H62 and H64 [13]. In this position, RbgA displaces the P-site and interacts with
the surrounding rRNA, including H92 and H93.Therefore, binding of RbgA requires contacts
with rRNA.
In our map of *T. brucei* pre-mtLSU, the corresponding regions forming the rRNA binding site
for mt-RbgA are not observed. The nucleotide binding site in mt-RbgA is empty, and fitting a
ligand results in clashes with the protein (Figure 2b). However, the comparison of our structure
with the *B. subtilis* LSU:RbgA complex (PDB ID 6PPK) shows nearly identical conformation of
the factor on the pre-mtLSU complex (Figure 2-figure supplement 1). Even the weak interaction
between the mt-RbgA C-terminal domain and the mitoribosomal protein uL14m is conserved
(Figure 2).
We found that the conserved position of mt-RbgA in *T. brucei* is maintained through two
specialized assembly linkers (Figure 2). The first linker is established between the C-terminal
domain and the TbLAF5 N-terminal helix. The latter adopts a crescent shape around the C-
terminal domain of mt-RbgA, forming a series of contacts with four out of its five helices
(Figure 2). The second linker is provided by TbLAF4 approaching from the mitoribosomal core.
It interacts with the mt-RbgA C-terminal domain and contributes a β-strand to a shared β-sheet
(Figure 2-figure supplement 2). Together, RbgA is anchored to the flexible core via two
dedicated factors that compensate for the lack of rRNA contacts.

TbLAF4 belongs to the family of the site-specific RluD pseudouridine synthases that is involved
in the bacterial LSU assembly via formation of pseudouridines at positions 1911, 1915 and 1917
(E. coli numbering) in the H69 end-loop [19, 20]. In pre-mtLSU structure, TbLAF4 encircles the
immature rRNA nucleotides U1017-U1021 and G1073-U1087 (Figure 3-figure supplement 2),
and its active site is occupied by uridine 1017 likley present in H89-93 of the mature rRNA
(Figure 2-figure supplement 2B). The N-terminal domain is positioned at the distance of ~80 Å
towards the L7/L12 stalk, and this connection plays a key role in coordinating the factors
between the different functional sites (Figure 3).

TbLAF5 belongs to the family of SpoU RNA methyltransferases, but appears to have a closed
active site that does not allow the binding of its typical S-adenosyl methionine cofactor (Figure
2-figure supplement 2A). It is located peripherally, and bound to the mitoribosome via a C-
terminal 24-residue helix interacting with rRNA H41/42, and via TbLAF8 (Figure 2-figure
supplement 2). Therefore, TbLAF4 and TbLAF5/TbLAF8 perform a structural scaffolding role
for binding mt-RbgA.

Figure 2. Binding of the mt-RgbA and mt-EngA to the subunit interface. (**A**) mt-RgbA is bound to TbLAF4 and TbLAF5, which are connected to the L7/L12 stalk; mt-EngA is associated with TbLAF9 and TbLAF10, which are connected to the CP. (**B**) A short helix of mt-RgbA (yellow) interacts with a flipped A894 nucleotide from H75 (white). Two GTPs in the GTPase domains on mt-EngA are shown as sticks. The residues forming electrostatic and hydrophobic interactions between mt-EngA and mt-RgbA are shown in the top right inset. Clashes between superimposed GTP in its putative binding site on mt-RgbA are shown in the bottom right inset.

168 **Figure 3. Network of interactions between the assembly factors in pre-mtLSU.** (**A**)
169 Assembly factors shown on the background of the pre-mtLSU density map, featuring the
170 interconnection. (**B**) Schematic of protein-protein network. The node size represents the relative
171 molecular mass of the protein. All the assembly factors are interconnected with each other.

GTPase mt-EngA is stabilized via protein extension

In the subunit interface, we identified another stably bound conserved GTPase homolog, mt-EngA. It contains two tandemly arranged GTPase domains and a C-terminal K homology (KH) domain, which is pointed towards the PTC. We could model two GTPs in the GTPase domains (Figure 2B). The positioning is identical to bacteria, suggesting a functional conservation. The assembly factor occupies the space between the PTC and the E-site (Figure 2-figure supplement 1), and a role in chaperoning rRNA has been proposed [12]. The comparison with *E. coli* LSU:EngA complex reveals conformational differences that further highlight the unconventional nature of the *T. brucei* protein-rich system, and its role in the stabilization of the conserved assembly factors.

Firstly, the N-terminal GTPase domain is extended by 60 residues, with residues 101-108 stabilizing a helix-turn-helix motif (275-305), which remained unresolved in the bacterial complex (Figure 2-figure supplement 1B). This motif is important for the stabilization of EngA, because one helix is stacked against a helix of TbLAF10, whereas the other forms a helical bundle with an unidentified protein (Figure 2-figure supplement 3). TbLAF10 is one of the largest mitoribosomal assembly factors, which spans 110 Å through the rRNA core to the CP (Figure 2 and 3, Figure 3-figure supplement 1). It has the contact area with rRNA of ~8400 Å and keeps the nucleotides U904-A917 as well as U953-A965 (putative H80-83) in a premature conformation (Figure 3).

Secondly, the N-terminal residues 72-75 stabilize a short helix (residues 367-374), which is buried within rRNA groove via Arg367 and Arg369 (Figure 2b). It disrupts the local structure of

H75 and stabilizes the flipped nucleotide A894 (Figure 2b). This loop is also highly charged in
the corresponding *E.coli* structure, but does not adopt the helical conformation. Finally, the N-
terminus forms additional contacts with five mitoribosomal proteins (bL28m, bL35m, bL19m,
mL64, mL74), a stabilizing protein mass that compensates for the missing rRNA in this region.
Overall, while the N-terminal GTPase domain aligns well with the bacterial EngA, its interacting
partners in our structure are proteinaceous and specific to *T. brucei*.

The conserved globular domains of mt-EngA are associated with the pre-mtLSU core via
TbLAF9. Its three helices from the N-terminus encloses the N-terminal GTPase domain helix
230-242 (Figure 2- figure supplement 3). Here, TbLAF9 replaces the missing rRNA H82, H88
and protein L1, which binds the EngA N-terminal GTPase domain in bacteria. TbLAF9 spans
over 100 Å to the top of the CP, where it also stabilizes unwound rRNA (Figure 1-figure
supplement 3, Figure 3-figure supplement 1). Thus, mt-EngA is stably bound via protein
extension and also associated with the protein-based scaffold of assembly factors, including the
high molecular weight TbLAF9 and TbLAF10, which are connected to the CP (Figure 3).

The module mt-RbgA:mt-EngA coordinates maturation of interfacial rRNA

The process of the LSU assembly is dynamic with cooperative action of different assembly
factors [9, 11]. Although RbgA and EngA have previously been visualized separately on the
bacterial LSU though deletion and reconstitution experiments, our cryo-EM structure shows both
factors simultaneously associated with the mt-LSU and with each other. This is attributed to the
stabilization by the protein linkers and extensions at the subunit interface. The presence of both
factors rationalizes why rRNA domain V is unexpectedly well resolved compared to the mature
mtLSU [4]. We were able to model 33% more nucleotides relative to the mature LSU (Figure 4),
thereby showing that the H89-93 region does not occupy the expected final position and
highlighting a need for prominent remodeling during maturation (Figure 4, Figure 4- figure
supplement 1).

The contacts between mt-RbgA and mt-EngA are formed between the N-terminal domain and
KH domains, respectively (Figure 2b). The shared surface area is ~540 Å, and each of the
domains is also associated with conserved rRNA. The formed contacts between mt-RbgA and
mt-EngA include electrostatic interactions, as well as hydrophobic residues (Figure 2b). Since
the structures and positions of both factors are conserved, these results indicate that a similar
overall mode of interaction could take place in other translational systems.

Figure 4. Tertiary structure of rRNA in pre-mtLSU (A) and mature mtLSU (B). Shown from the subunit interface (left) and sideview (right). (A) Two views of rRNA related by 90° are shown with each domain in a different color. Domain V is more structured in pre-mtLSU, and H89-93 adopt a different conformation. Domain II that is responsible for L7/L12 stalk is largely disordered.

Progressive maturation of the L7/L12 stalk

The L7/L12 stalk is a universal mobile element that associates translational protein factors with the A-site. It typically consists of the rRNA platform that connects to the flexible factor-recruiting series of bL12m copies. The ubiquitous ribosomal proteins uL10m, uL11m and mitochondria-specific mL43 link the different components together. In our structure, most of the protein constituents of the stalk are missing and others adopted conformational changes (Figure 5).

In the region of the platform, at least three proteins (uL16m, bL36m, mL88) associated with the rRNA in the mature mtLSU are absent. Consistently, the rRNA platform is not folded, as the folding relies on the missing mitoribosomal proteins. Instead, the N-terminal domain of TbLAF4 extends from the subunit interface to occupy part of the space left by uL16m absence (Figure 5A and B). It binds two specific assembly TbLAF2 and TbLAF7 (Figure 5A). TbLAF2 mediates further contacts with the core of the pre-mtLSU. TbLAF7 consists of 7-stranded beta-barrel, 12 α-helices, and 63-residue tail inserted into the mitoribosomal core. The structure suggests that both TbLAF2 and TbLAF7 need to dissociate for the missing mitoribosomal proteins to assemble (Figure 5A).

In the factor-recruiting region, instead of the terminal bL12m copies, TbLAF3 and density corresponding to UNK6 form a protrusion (Figure 5A). They comprise a protein continuum of at least 13 helices associated with each other, and attached to the platform region through 30-residue C-terminal tail of TbLAF3, forming a helical bundle with mL75 (Figure 5A and C). Overall, this protein module extends ~70 Å from the core in a similar fashion to the L7/L12 stalk, and both appear to be mutually exclusive.

The position of the uL10m N-terminus, which links the stalk to the body, is occupied by TbLAF3 C-terminus. It interacts with mL43, resulting in its helix being bent by 90° (Figure 5B). This conformational change and the lack of uL10m is correlated with ~15 Å shift of uL11m to occupy the formed void (Figure 5B). Nevertheless, TbLAF3 is only peripherally associated with mL43, and therefore in principle the protrusion can be replaced by the mature L7/L12 stalk independently.

Together, our structure proposes a working model for the L7/L12 stalk maturation through four steps: 1) TbLAF4, which is extended from the subunit interface anchoring mt-RgbA, has to be removed. 2) TbLAF2:TbLAF7 is evicted from the ribosomal core; 3) rRNA platform is folded; 4) uL16m, bL36m, and mL88 are recruited to the rRNA platform; 5) TbLAF3 is removed, and uL11m, mL43 adopt their mature conformations; 6) bL10m and bL12m stalk proteins are associated.

Figure 5. Assembly of the L7/L12 stalk. (**A**) In pre-mtLSU, TbLAF4 extends from the subunit
interface to occupy the position of uL16m in the mature mtLSU. TbLAF2 and TbLAF7 are
bound at the stalk base to the unfolded rRNA H41/42. TbLAF3 and additional protein form a
protrusion similar to bL10m:bL12m. Other mitoribosomal protein removed for clarity. (**B**)
Conformational changes from pre-mtLSU (green) to mature mtLSU (white) include mL43 and
uL11m. (**C**) TbLAF3, mL75, and UNK protein form continuum of at least 13 helices that is
peripherally associated.

Below the L7/L12 stalk we identified three additional proteins: an ortholog of the bacterial
ribosome silencing factor mt-Rsf, a LYR (leucine-tyrosine-arginine) motif containing protein
L0R8F8 (TbLAF1), as well as an associated mt-ACP (Figure 1 and 3). The human and fungi
mitochondrial counterparts of these three proteins have been shown to be involved in late
assembly stages of the mitoribosome [21,22]. In our structure, the module is further stabilized by
mL85, however unlike in human and fungi, it does not appear to obstruct the binding of the
mature mtSSU, likely because of the reduced rRNA and different overall shape. This suggests
that in order for this protein module to act according to the same mechanism as found in the

286 mitoribosomes from other species, a complementary structural domain from pre-mtSSU would
287 be required for steric hindrance.

Assembly of the CP is linked to the subunit interface and L1 maturation via mt-ACP

The most prominent architectural feature of the pre-mtLSU complex is the absence of the CP,
which is a universal ribosomal element that defines the shape of the LSU and forms bridges with
the SSU head. In mitoribosomes, the CP is particularly protein-rich [23-28]. The acquisition of
those mitochondria-specific proteins took place in the early evolution of the mitoribosome [29].

In the *T. brucei* pre-mtLSU, all the CP mitoribosomal proteins are missing and the high
molecular weight assembly factors TbLAF6 (69 kDa) and TbLAF9 (67 kDa) are present (Figure
1 and Figure 6A). TbLAF6 binds on the solvent side of the CP covering the otherwise exposed
rRNA, which only engages in limited base pairing. This assembly factor is annotated as a
putative TruD family pseudouridine synthase. However, in our structure, it displays a two-strand
antiparallel β-sheet near the catalytic site protruding into the ribosomal core interacting with four
mitoribosomal proteins (uL15m, bL28m, bL35m, mL74) (Figure 6B). TbLAF9 is an exclusively
helical protein, comprised of at least 29 helices. It binds on top of the rRNA, providing an
additional protective protein cap (Figure 6A).
Two of the TbLAF9 helices interface with a four-helix bundle, which we identified as mt-ACP
(mt-ACP2) in the density with a local resolution of 3.5 Å (Figure 1-supplement figure 2). Since
mt-ACP proteins are known to form interactions with Leu-Tyr-Arg (LYR)-motif proteins, we
compared the mt-ACP1-TbLAF1 module from the subunit interface with the CP mt-ACP2
region (Figure 6C). The helices of the LYR-motif protein TbLAF1 align well with an unassigned
density corresponding to three helices associated with the mt-ACP2. The interactions in both
cases are mediated by the 4′-phosphopantetheine (4′-PP) modification of mt-ACP, resembling
the canonical interactions between mt-ACP and the LYR-motif proteins. The 4′-PP appears to be
acylated as indicated by the density (Figure 6C).
The importance of the mt-ACP2 protein module in our structure is of twofold. First, it directly
binds the L1 stalk and locks it in an inward facing nonfunctional conformation (Figure 1).
Second, it is involved in mt-EngA stabilization by forming a U-shaped continuum from TbLAF9
on the solvent side of the CP to the subunit interface (Figure 3). It therefore contributes to the
protein network connecting between the various functional sites. Thus, mt-ACP2 is a principal
partner in the mitoribosomal assembly pathway. In the pre-mtSSU, mt-ACP was also identified
as one of the assembly factors [5]. In addition, ACPs in mitochondria act as subunits of the
electron transport chain [29] and Fe-S cluster assembly complexes [30]. This further supports the
proposed concept that mt-ACPs could act as signaling molecules in an intramitochondrial
metabolic state sensing circuit [31].

Figure 6. The CP assembly intermediate. (**A**) TbLAF6 and TbLAF9 form the CP in the pre-mtLSU. (**B**) TbLAF6 and mL64 elements are inserted through rRNA loop corresponding to H83. Conformational change of mL64 from pre-mtLSU to mature mtLSU (white) is indicated. (**C**) Comparison between the mt-ACP1:TbLAF1 (left) and the CP mt-ACP2 region (right). The density (white) for acylated 4′-PP is indicated. Bottom panel, comparison with mt-ACP and associated LYR-motif proteins from complex I (PDB ID 5LNK) and human mitoribosome (PDB ID 5OOM) shows the canonical interactions.

At the CP, the assembly factors cooperatively bind unwound rRNA nucleotides U934-U953
(H83) (Figure 6A). Remarkably, the rRNA forms a loop 25 Å in diameter that contains the
TbLAF6 β-sheet and mL64 C-terminal helix, both inserted from the opposite directions (Figure
B). The conserved helix of mL64 is shifted in our pre-mtLSU structure ~30 Å from the final
location on the mature LSU, where it aligns the E-site (Figure 6B). Interestingly, this is also one
of the most conserved mitoribosomal proteins across species [32], and its role in tRNA
translocation has been established [33]. To switch to the conserved and mature position, the
extended C-tail of mL64 has to liberate from the rRNA loop and then undergo a large
conformational shift towards the L1 stalk. Subsequently, the C-tail is inserted to its mature
position, where it contacts CP components absent from the assembly intermediate. Since the L1
stalk is shifted, the maturation towards a mature LSU is likely to occur in a concerted manner.
343

Conclusions
Together, our structure of the *T. brucei* pre-mtLSU in complex with at least 15 assembly factors
reveals that mt-RbgA and mt-EngA are held by protein linkers and interact with each other. The
analysis of the different sections shows that high molecular weight TbLAF4, TbLAF9 and
TbLAF10 proteins form a network that spans over 180 Å and connects the subunit interface with
the progressive maturation of the L7/L12 stalk, L1 stalk, and the assembly of the CP. This design
would allow biogenesis steps towards the mature mtLSU to be coordinated. In this regard, the
tight binding of the mt-ACP2 with partner proteins at the CP emphasizes its structural role in
forming the connections. For the L7/L12 stalk, the structure further suggests a sequential
mechanism of assembly, and protein replacement that can be a final control. Overall, the protein-
based communication allows the formation of the here reported stable assembly intermediate that
is ready to progress through the biogenesis pathway.

356
357 **Figure 1-figure supplement 1.** Cryo-EM data processing. (**A**) Representative micrograph. (**B**)
358 Processing workflow. (**C**) Fourier shell correlation (FSC) curves. Resolution is estimated based
359 on the 0.143 FSC cut-off criterion (red line).
360
361

362
Figure 1-figure supplement 2. (**A**) final map colored by local resolution. (**B**) Models for
individual assembly factors and newly identified proteins colored by local resolution of the
density in the corresponding regions.

366
367 **Figure 1-figure supplement 3.** Comparison of the cryo-EM maps between pre-mtLSU and
368 mature mtLSU in two 90°-related views. Assembly factors (shades of purple) are indicated.

Figure 2-figure supplement 1. The binding of mt-RbgA and mt-EngA at the mtLSU interface.
(**A**) Comparison between pre-mtLSU and bacterial counterparts *E. coli* 50S:EngA (PDB ID
3J8G) and *B. subtilis* 45S:RbgA (PDB ID 6PPK) shows nearly identical positions of the factors
on their ribosomal complexes. (**B**) Comparison between mt-RbgA:mt-EngA module from the
pre-mtLSU and superimposed bacterial counterparts combined from the two structures from (A)
shows nearly identical conformations of the factors. The N-terminal extension of mt-EngA
(orange) is buried in the mitoribosomal core and stabilizes the binding, as well as 275-305 region
(orange). The comparison suggests that a similar mode of interaction would exist in bacterial
ribosomal assembly intermediates, where both factors present.

394
395 **Figure 2-figure supplement 2.** (**A**) The active site of TbLAF4 (yellow) is occupied by uridine
396 1048. (**B**) The factor TbLAF4 (yellow) binds mt-RbgA (pale-yellow) via a shared β-sheet
397 (circled). The methyltransferase site of TbLAF5 does not allow for the binding of S-adenosyl
398 methionine cofactor (white sticks and surface) due to clashes with the protein residues (red).
399
400
401
402

403
404 **Figure 2-figure supplement 3.** N-terminal extension (yellow) of mt-EngA stabilizes helix-turn-
405 helix (275-305), which forms interaction with TbLAF10 on the other side (bottom right panel),
406 and a helical bundle with an unidentified protein UNK that is in contact with TbLAF9.
407
408

Figure 3-figure supplement 1. Binding of assembly factors to rRNA. For each panel rRNA is shown with an individual protein characterized in the structure for the first time. Bottom right panel illustrates the total RNA that is involved in the interactions (yellow) with the assembly factors.

414
415   **Figure 4-figure supplement 1.** Secondary structure rRNA diagram derived from the model and
416   colored by domain. Unbuilt sections that appear in the mature mtLSU are shown in gray.
417   Domains in Roman numerals.

Supplementary File 1. Cryo-EM data collection, refinement and validation statistics

	Consensus map
Data collection and processing	
Magnification	130000x
Voltage (kV)	300
Electron exposure (e–/Å^2)	35
Defocus range (μm)	-0.8 ~ -3.5
Pixel size (Å)	1.05
Symmetry imposed	C1
Initial particle images (no.)	896,263
Final particle images (no.)	32,339
Map resolution (Å) FSC threshold 0.143	3.50
Map resolution range (Å)	3.0~10
Refinement	
Map sharpening B factor (Å^2)	-70
Model composition	
Non-hydrogen atoms	146831
Protein residues	17415
Ligands	10
B factors min/max/avg (Å^2)	
Protein	17/172/68
Nucleotide	22/281/57
Ligand	35/194/61
R.m.s. deviations	
Bond lengths (Å)	0.002
Bond angles (°)	0.46
Validation	
MolProbity score	1.65
Clashscore	6.7
Poor rotamers (%)	0.35
Ramachandran plot	
Favored (%)	95.6
Allowed (%)	4.10
Disallowed (%)	0.03

 Supplementary File 2. Summary of pre-mtLSU components

Alias	Chain ID	TriTrypDB Gene ID (Lister strain 427)	TriTrypDB Gene ID (reference strain TREU927)	Uniprot ID (reference strain TREU927)	Full size	Modeled residues	Comment
12S rRNA	AA	N/A	N/A	N/A	1176	1-205, 254-264, 270-356, 369-380, 404-413, 445-450, 456-534, 541-582, 591-594, 796-883, 887-967, 980-999, 1004-1008, 1012-1021, 1071-1090, 1095-1176	
uL3m	AE	Tb427.03.5610	Tb927.3.5610	Q580R4	473	38-265, 272-404	
uL4m	AF	Tb427tmp.02.3810	Tb927.11.6000	Q385G8	351	18-459	
bL9m	AI	Tb427.05.3410	Tb927.5.3410	Q57UC5	263	9-220	
uL11m	AK	Tb427.02.4740	Tb927.2.4740	N/A	342	26-200, 207-235, 239-306	239-306 built as UNK
uL13m	AN	Tb427.04.1070	Tb927.4.1070	Q580D5	202	10-180	
uL14m	XG	Tb427.04.930	Tb927.4.930	Q580C1	217	20-107, 114-189	
uL15m	AP	Tb427.05.3980	Tb927.5.3980	Q57U68	374	10-136, 150-322, 354-363	
bL17m	AR	Tb427.08.5860	Tb927.8.5860	Q57YI7	301	11-266	
bL19m	AT	Tb427.01.1210	Tb927.1.1210	Q4GZ98	144	2-139	
bL20m	AU	Tb427tmp.01.1930	Tb927.11.10170	Q383R2	213	10-140, 162-205	
bL21m	AV	Tb427.07.4140	Tb927.7.4140	Q57UP4	188	6-185	
uL22m	AW	Tb427.07.2760	Tb927.7.2760	Q57Y86	278	2-278	
uL23m	AX	Tb427tmp.03.0260	Tb927.11.870	Q387G3	246	64-228	
uL24m	AY	Tb427.03.1710	Tb927.3.1710	Q57ZE0	378	1-311, 318-340	
bL28m	A1	Tb427.06.4040	Tb927.6.4040	Q586A2	241	10-226	
uL29m	A2	Tb427tmp.160.5240	Tb927.9.7170	Q38EM7	471	9-233, 248-471	
uL30m	A3	Tb427tmp.211.0230	Tb927.9.8290	Q38ED8	218	51-200	
bL32m	A5	Tb427.04.2330	Tb927.4.2330	Q584F4	80	26-80	2Fe-2S cluster binding

bL35m	A8	Tb427.10.1870	Tb927.10.1870	Q38C55	181	40-181	
mL41	Ae	Tb427tmp.01.1600	Tb927.11.9830	Q383U6	197	47-161	
mL42	Af	Tb427tmp.01.1840	Tb927.11.10080	Q383S1	189	41-173	
mL43	Ag	Tb427.04.4600	Tb927.4.4600	Q583E5	260	2-186	
mL49	Al	Tb427.05.3110	Tb927.5.3110	Q57Z82	218	37-101, 114-218	
mL52	Ao	Tb427tmp.02.2250	Tb927.11.4650	Q385V2	152	19-151	
mL53	Ap	Tb427.07.2990	Tb927.7.2990	N/A	309	16-303	
mL63	At	Tb427.07.7010	Tb927.7.7010	Q57XS1	154	10-154	
mL64	Av	Tb427tmp.01.3500	Tb927.11.11630	Q383B7	242	27-222	NAD binding
mL67	BA	Tb427tmp.55.0016	Tb927.11.1630	Q386Z1	831	27-83, 130-328, 335-542, 562-824	
mL68	BB	Tb427.10.600	Tb927.10.600	Q38CI0	541	62-258, 264-294, 304-341, 346-450	264-294, 304-341 built as UNK
mL70	BD	Tb427.06.4200	Tb927.6.4200	Q586Y7	547	105-521	
mL71	BE	Tb427.07.3460	Tb927.7.3460	Q57WG1	449	11-190, 228-448	
mL72	BF	Tb427.06.3930	Tb927.6.3930	Q585Z1	426	26–62, 118-421	
mL74	BH	Tb427.10.7380	Tb927.10.7380	Q38AM5	349	89-314	
mL75	BI	Tb427.10.380	Tb927.10.380	Q38CK0	342	20-342	
mL76	BJ	Tb427tmp.01.2340	Tb927.11.10570	Q383M2	333	173-333	
mL77	BK	Tb427.06.2480	Tb927.6.2480	Q584Q8	386	84-156, 188-233, 254-269, 280-386	254-259 built as UNK
mL78	BL	Tb427.10.11050	Tb927.10.11050	Q389N4	312	31-130, 141-197, 216-265, 281-306	
mL80	BN	Tb427.06.1440	Tb927.6.1440	Q585A3	302	53-266	
mL81	BO	Tb427tmp.02.3230	Tb927.11.5530	Q385L5	262	36-193, 210-262	

mL83	BQ	Tb427.07.3430	Tb927.7.3430	Q57WF8	231	16-200
mL84	BR	Tb427.06.4080	Tb927.6.4080	Q586A6	205	11-205
mL85	BS	Tb427tmp.160.2250	Tb927.9.3640	Q38FG8	198	20-163

456 **Supplementary File 3. Contacts of assembly factors with rRNA.** Regions and nucleotides of
457 respective rRNA domains are color-coded as in Figure 4. The underscored regions are in mature
458 conformation in the pre-mtLSU.

Assembly factor	Contacts with rRNA	
	Region	Nucleotides
mt-EngA	H74, H75, H88, H90-91	868-874, 891-898, 958-967, 994-999, 1004
mt-RbgA	H90-91	1021, 1071
mt-RsfS	H95	1126-1132, 1138-1141
TbLAF2	H39, H43, H72, H74, H88, H93	132-135 347-351, 404, 821-829, 832-835, 860-862, 904, 965-677, 985-992, 1079-1082, 1119
TbLAF3	H43	411-412, 446
TbLAF4	H39, H90-93	351-353, 857-859, 992-994, 1008, 1012-1021, 1071-1087, 1122-1124
TbLAF5	H43, H90, H95	409-413, 1020-1021, 1136-1137
TbLAF6	H13, H28, H37, H80-81, H88	65-66, 164-166, 313, 328, 908, 917-919, 935-939, 944-953
TbLAF7	H41, H43	407, 448-450
TbLAF8	H95	1131-1140
TbLAF9	H80-81, H88	908-909, 915-916, 919-924, 930-940, 952-957, 961
TbLAF10	H26, H32, H33, H35a, H37, H39, H51, H72, H73-75, H80-81, H93	127-139, 199-200, 270-276, 289-293, 306-316, 344-355, 497, 550, 555, 826-828, 847-848, 850-855, 860-862, 870, 903-922, 930-933, 947-980, 955-965, 983 -992, 1085-1089

Materials and Methods

Strains and growth conditions

T. brucei procyclic Lister strain 427 was grown in SDM-80 medium supplemented with 10%
fetal bovine serum. Mitochondria were isolated as described earlier [34]. 1.5×10^{11} cells were
harvested, washed in 20 mM sodium phosphate buffer pH 7.9 with 150 mM NaCl and 20 mM
glucose, resuspended in 1 mM Tris-HCl pH 8.0, 1 mM EDTA, and disrupted by 10 stroke in 40
ml Dounce homogenizer. The hypotonic lysis was stopped by immediate addition of 1/6 volume
of 1.75 M sucrose. Crude mitochondria were pelleted (15 min at 16000 xg, 4°C), resuspended in
20 mM Tris-HCl pH 8.0, 250 mM sucrose, 5 mM $MgCl_2$, 0.3 mM CaCl2 and treated with 5
μg/ml DNase I for 60 min on ice. DNase I treatment was stopped by addition of one volume of
the STE buffer (20 mM Tris-HCl pH 8.0, 250 mM sucrose, 2 mM EDTA) followed by
centrifugation (15 min at 16000 xg, 4°C). The pellet was resuspended in 60% Percoll in STE and
loaded on the bottom of six 10-35% Percoll gradient in STE in polycarbonate tubes for SW28
rotor (Beckman). Gradients were centrifuged for 1 hour at 24000 rpm, 4°C. The middle diffused

phase containing mitochondrial vesicles (15-20 ml per tube) was collected, washed twice in the
STE buffer, snap-frozen in liquid nitrogen and stored at -80°C.

Purification of mitoribosomes

Mitochondria were further purified further using a stepped sucrose gradient (60 %, 32 %, 23 %,
15%) in a low ionic strength buffer (50 mM HEPES/KOH pH 7.5, 5 mM MgOAc, 2 mM
EDTA). A thick pellet at the 60-32% interface was collected and lysed by mixing with 5
volumes of detergent containing lysis buffer (25 mM HEPES/KOH pH 7.5, 100 mM KCl, 15
mM MgOAc, 1.7 % Triton X-100, 2 mM DTT, Complete-EDTA Free Protease Inhibitor). The
lysate was centrifuged at 30,000 xg twice, saving the supernatant after each spin. The
supernatant was then subjected to differential PEG precipitation; PEG 10,000 was added to reach
a concentration of 1.5 % (w/v) and incubated on ice for 10 mins, followed by a spin at 30,000 xg.
The supernatant was transferred to a fresh tube, and PEG 10,000 was added to reach a
concentration of 8 % (w/v) then incubated on ice for 10 mins, followed by a spin at 30,000 xg.

The pellet was then resuspended in 800 µl of lysis buffer and then layered onto a 34% sucrose
cushion (25 mM HEPES/KOH pH 7.5, 100 mM KCl, 15 mM MgOAc, 1.0 % Triton X-100, 2
mM DTT, Complete-EDTA Free Protease Inhibitor) in a TLA120.2 centrifuge tube (0.4 ml of
cushion per tube). Mitoribosomes were pelleted through the cushion by centrifugation at 231,550
xg for 45 min. Pelleted mitoribosomes were resuspended using a total of 100 µl of resuspension
buffer (25 mM HEPES/KOH pH 7.5, 100 mM KCl, 15 mM MgOAc, 0.01 % β-DDM, 2 mM
DTT). The resuspended mitoribosomes were then layered onto a continuous 15-30 % sucrose
gradient and centrifuged in a TLS55 rotor for 120 min at 213,626 xg. The gradient was
fractionated manually, and fractions containing mitoribosome as judged by the 260 nm
absorbance were pooled and buffer exchanged in a centrifugal concentrator.

Cryo-EM and model building

For cryo-EM analysis, 3 µL of the sample at a concentration of OD260 3.5, was applied onto a
glow-discharged (20 mA for 30 seconds) holey-carbon grid (Quantifoil R2/2, copper, mesh 300)
coated with continuous carbon (of ~3 nm thickness) and incubated for 30 seconds in a controlled
environment of 100% humidity and 4 ºC temperature. The grids were blotted for 3 seconds,
followed by plunge-freezing in liquid ethane, using a Vitrobot MKIV (FEI/Thermofischer). The
data was collected on a FEI Titan Krios (FEI/Thermofischer; Scilifelab, Stockholm, Sweden, and
ESRF, Grenoble, Frane) transmission electron microscope operated at 300 keV, using C2
aperture of 70 µm; slit width of 20 eV on a GIF quantum energy filter (Gatan). A K2 Summit
detector (Gatan) was used to collect images at a pixel size of 1.05 Å (magnification of 130,000X)
with a dose of ~35 electrons/Å2 fractionated over 20 frames. A defocus range of 0.8 to 3.5 µm
was applied.

19,158 Micrographs (after bad images were removed based on real and reciprocal space features)
were collected across 5 non-consecutive data acquisition sessions and processed together using
RELION. 896,263 particles were picked using Warp and coordinates were imported into
RELION for particle extraction at an initial binning factor of two. The particles were subjected to

supervised 3D classification using references generated previously in a screening dataset, which was started based on the Trypanosoma cytosolic ribosome as an initial model. This crude separation classified the 207,788 particles as mtLSU-like, and the remaining as mature mtLSU-like, SSU-like or monosomes. This subset was subjected to auto-refinement separately to improve the angular assignments and then classified further using fine-angular searches with a solvent mask applied. From the mtLSU-like particles, 32,339 particles were retained as pre-mtLSU of good quality and the rest were discarded as non-particles. The retained pre-mtLSUs were then subjected to auto-refinement once more to improve the angles further, this time applying a solvent mask during the refinement procedure, and then the 3D reconstructions obtained were used as a reference for CTF refinement to improve the reconstruction. The final map was then estimated for local resolution using RELION and sharpened with a B-factor appropriate for the reconstruction as estimated automatically using the postprocessing procedure.

Model building was done using *Coot* 0.9 [35]. First the model of the mature mtLSU (PDB ID:6HIX) was fitted to the density. Chains present in the pre-mtLSU were then individually fitted and locally refined. Additional chains were modeled de-novo and sequences assigned by direct comparison with the density. All models were refined iteratively using PHENIX [36] realspace refinement and validated using MolProbity [37]. The data collection, model refinement and validation statistics are presented in Supplementary data file1. All figures were prepared either in Chimera [38] or ChimeraX [39] with additional graphical elements created using Inkscape.

Data availability

The electron density map has been deposited in EMDB under accession code EMD-XXX. The model has been deposited in PDB under accession code XXXX. All data is available in the paper or Supplementary Information.

Acknowledgements

The authors thank the SciLifeLab cryo-EM and mass spectrometry facilities. This work was supported by the Swedish Foundation for Strategic Research (FFL15:0325), Ragnar Söderberg Foundation (M44/16), Swedish Research Council (NT_2015-04107), Cancerfonden (2017/1041), European Research Council (ERC-2018-StG-805230), Knut and Alice Wallenberg Foundation (2018.0080), EMBO Young Investigator Program to A.A., and by and by Czech Science Foundation (18-17529S) and ERD fund (CZ.02.1.01/0.0/0.0/16_019/0000759) to A.Z., and by Czech Science Foundation (20-04150Y) to O.G. The cryo-EM facility is funded by the Knut and Alice Wallenberg, Family Erling Persson, and Kempe foundations.

Author contributions

Victor Tobiasson: Formal analysis, Investigation, Validation, Visualization, Writing - original draft, Writing - review and editing

Ondřej Gahura: Conceptualization, Data curation, Formal analysis, Investigation, Validation, Visualization, Writing - original draft, Writing - review and editing

Shintaro Aibara: Conceptualization, Data curation, Formal analysis, Writing - review and editing

Rozbeh Baradaran: Investigation, Validation, Visualization, Writing - review and editing

Alena Zíková: Conceptualization, Investigation, Funding acquisition, Project administration, Supervision, Writing - original draft, Writing - review and editing

Alexey Amunts: Conceptualization, Investigation, Funding acquisition, Project administration, Resources, Supervision, Validation, Writing - original draft, Writing - review and editing

References

1. Ott, M., Amunts, A. & Brown, A. (2016) Organization and Regulation of Mitochondrial Protein Synthesis, *Annu Rev Biochem*.

2. Greber, B. J. & Ban, N. (2016) Structure and Function of the Mitochondrial Ribosome, *Annu Rev Biochem.* **85**, 103-32.

3. Zikova, A., Panigrahi, A. K., Dalley, R. A., Acestor, N., Anupama, A., Ogata, Y., Myler, P. J. & Stuart, K. (2008) Trypanosoma brucei mitochondrial ribosomes: affinity purification and component identification by mass spectrometry, *Mol Cell Proteomics.* **7**, 1286-96.

4. Ramrath, D. J. F., Niemann, M., Leibundgut, M., Bieri, P., Prange, C., Horn, E. K., Leitner, A., Boehringer, D., Schneider, A. & Ban, N. (2018) Evolutionary shift toward protein-based architecture in trypanosomal mitochondrial ribosomes, *Science.* **362**.

5. Saurer, M., Ramrath, D. J. F., Niemann, M., Calderaro, S., Prange, C., Mattei, S., Scaiola, A., Leitner, A., Bieri, P., Horn, E. K., Leibundgut, M., Boehringer, D., Schneider, A. & Ban, N. (2019) Mitoribosomal small subunit biogenesis in trypanosomes involves an extensive assembly machinery, *Science.* **365**, 1144-1149.

6. Li, N., Chen, Y., Guo, Q., Zhang, Y., Yuan, Y., Ma, C., Deng, H., Lei, J. & Gao, N. (2013) Cryo-EM structures of the late-stage assembly intermediates of the bacterial 50S ribosomal subunit, *Nucleic Acids Res.* **41**, 7073-83.

7. Ni, X., Davis, J. H., Jain, N., Razi, A., Benlekbir, S., McArthur, A. G., Rubinstein, J. L., Britton, R. A., Williamson, J. R. & Ortega, J. (2016) YphC and YsxC GTPases assist the maturation of the central protuberance, GTPase associated region and functional core of the 50S ribosomal subunit, *Nucleic Acids Res.* **44**, 8442-55.

8. Jomaa, A., Jain, N., Davis, J. H., Williamson, J. R., Britton, R. A. & Ortega, J. (2014) Functional domains of the 50S subunit mature late in the assembly process, *Nucleic Acids Res.* **42**, 3419-35.

9. Davis, J. H., Tan, Y. Z., Carragher, B., Potter, C. S., Lyumkis, D. & Williamson, J. R. (2016) Modular Assembly of the Bacterial Large Ribosomal Subunit, *Cell.* **167**, 1610-1622 e15.

10. Nikolay, R., Hilal, T., Qin, B., Mielke, T., Burger, J., Loerke, J., Textoris-Taube, K., Nierhaus, K. H. & Spahn, C. M. T. (2018) Structural Visualization of the Formation and

Activation of the 50S Ribosomal Subunit during In Vitro Reconstitution, *Mol Cell.* **70**, 881-893 e3.

11. Davis, J. H. & Williamson, J. R. (2017) Structure and dynamics of bacterial ribosome biogenesis, *Philos Trans R Soc Lond B Biol Sci.* **372**.

12. Zhang, X., Yan, K., Zhang, Y., Li, N., Ma, C., Li, Z., Zhang, Y., Feng, B., Liu, J., Sun, Y., Xu, Y., Lei, J. & Gao, N. (2014) Structural insights into the function of a unique tandem GTPase EngA in bacterial ribosome assembly, *Nucleic Acids Res.* **42**, 13430-9.

13. Seffouh, A., Jain, N., Jahagirdar, D., Basu, K., Razi, A., Ni, X., Guarne, A., Britton, R. A. & Ortega, J. (2019) Structural consequences of the interaction of RbgA with a 50S ribosomal subunit assembly intermediate, *Nucleic Acids Res.* **47**, 10414-10425.

14. Tegunov, D. & Cramer, P. (2019) Real-time cryo-electron microscopy data preprocessing with Warp, *Nat Methods.* **16**, 1146-1152.

15. Kimanius, D., Forsberg, B. O., Scheres, S. H. & Lindahl, E. (2016) Accelerated cryo-EM structure determination with parallelisation using GPUs in RELION-2, *Elife.* **5**.

16. Achila, D., Gulati, M., Jain, N. & Britton, R. A. (2012) Biochemical characterization of ribosome assembly GTPase RbgA in Bacillus subtilis, *J Biol Chem.* **287**, 8417-23.

17. Pausch, P., Steinchen, W., Wieland, M., Klaus, T., Freibert, S. A., Altegoer, F., Wilson, D. N. & Bange, G. (2018) Structural basis for (p)ppGpp-mediated inhibition of the GTPase RbgA, *J Biol Chem.* **293**, 19699-19709.

18. Barrientos, A., Korr, D., Barwell, K. J., Sjulsen, C., Gajewski, C. D., Manfredi, G., Ackerman, S. & Tzagoloff, A. (2003) MTG1 codes for a conserved protein required for mitochondrial translation, *Mol Biol Cell.* **14**, 2292-302.

19. Gutgsell, N. S., Deutscher, M. P. & Ofengand, J. (2005) The pseudouridine synthase RluD is required for normal ribosome assembly and function in Escherichia coli, *RNA.* **11**, 1141-52.

20. Gutgsell, N. S., Del Campo, M., Raychaudhuri, S. & Ofengand, J. (2001) A second function for pseudouridine synthases: A point mutant of RluD unable to form pseudouridines 1911, 1915, and 1917 in Escherichia coli 23S ribosomal RNA restores normal growth to an RluD-minus strain, *RNA.* **7**, 990-8.

21. Brown, A., Rathore, S., Kimanius, D., Aibara, S., Bai, X. C., Rorbach, J., Amunts, A. & Ramakrishnan, V. (2017) Structures of the human mitochondrial ribosome in native states of assembly, *Nat Struct Mol Biol.* **24**, 866-9.

22. Itoh, Y., Naschberger, A., Mortezaei, N., Herrmann, J. & Amunts, A. (2020) Analysis of translating mitoribosome reveals functional characteristics of protein synthesis in mitochondria of fungi, *bioRxiv* 2020.01.31.929331

23. Amunts, A., Brown, A., Bai, X. C., Llacer, J. L., Hussain, T., Emsley, P., Long, F., Murshudov, G., Scheres, S. H. & Ramakrishnan, V. (2014) Structure of the yeast mitochondrial large ribosomal subunit, *Science.* **343**, 1485-9.

24. Amunts, A., Brown, A., Toots, J., Scheres, S. H. W. & Ramakrishnan, V. (2015) Ribosome. The structure of the human mitochondrial ribosome, *Science.* **348**, 95-98.

25. Greber, B. J., Bieri, P., Leibundgut, M., Leitner, A., Aebersold, R., Boehringer, D. & Ban, N. (2015) Ribosome. The complete structure of the 55S mammalian mitochondrial ribosome, *Science.* **348**, 303-8.

26. Greber, B. J., Boehringer, D., Leibundgut, M., Bieri, P., Leitner, A., Schmitz, N., Aebersold, R. & Ban, N. (2014) The complete structure of the large subunit of the mammalian mitochondrial ribosome, *Nature.* **515**, 283-6.

27. Waltz, F., Soufari, H., Bochler, A., Giege, P. & Hashem, Y. (2020) Cryo-EM structure of the
RNA-rich plant mitochondrial ribosome, *Nat Plants.* **6**, 377-383.
28. Tobiasson, V. & Amunts, A. (2020) Ciliate mitoribosome illuminates evolutionary steps of
mitochondrial translation, *Elife.* **9**, e59264.
29. Zhu, J., King, M.S., Yu, M., Klipcan, L., Leslie, A.G. and Hirst, J. (2015) Structure of
subcomplex Iβ of mammalian respiratory complex I leads to new supernumerary subunit
assignments. *Proceedings of the National Academy of Sciences*, **112**, 12087-12092.
30. Van Vranken, J.G., Jeong, M.Y., Wei, P., Chen, Y.C., Gygi, S.P., Winge, D.R. and Rutter, J.
(2016). The mitochondrial acyl carrier protein (ACP) coordinates mitochondrial fatty acid
synthesis with iron sulfur cluster biogenesis. *Elife*, **5**, e17828.
31. Masud, A.J., Kastaniotis, A.J., Rahman, M.T., Autio, K.J. and Hiltunen, J.K. (2019).
Mitochondrial acyl carrier protein (ACP) at the interface of metabolic state sensing and
mitochondrial function. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, **1866**,
118540.
32. Petrov, A. S., Wood, E. C., Bernier, C. R., Norris, A. M., Brown, A. & Amunts, A. (2019)
Structural Patching Fosters Divergence of Mitochondrial Ribosomes, *Mol Biol Evol.* **36**, 207-
219.
33. Aibara, S., Modelska, A., Singh, V. & Amunts, A. (2020) Structural basis of mitochondrial
translation, *Elife.* in press.
34. Schneider, A., Charriere, F., Pusnik, M. & Horn, E. K. (2007) Isolation of mitochondria from
procyclic Trypanosoma brucei, *Methods Mol Biol.* **372**, 67-80.
35. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. (2010) Features and development of
Coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501.
36. Liebschner, D., Afonine, P.V., Baker, M.L., Bunkóczi, G., Chen, V.B., Croll, T.I., Hintze, B.,
Hung, L.W., Jain, S., McCoy, A.J. and Moriarty, N.W. (2019) Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta
Crystallogr D Struct Biol* **75**, 861–877.

37. Williams, JS. et al. (2018) MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Science* **27**, 293-315.
38. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C. and
Ferrin, T.E. (2004) UCSF Chimera – a visualization system for exploratory research and
analysis. *Journal of Computational Chemistry* **25**, 1605–1612.
39. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H. and
Ferrin, T.E. (2018) UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Science **27,** 14–25.