1

Size-dependent protein segregation creates a spatial switch for Notch and APP signaling

Minsuk Kwak[1,2,3,4,5,6,*], Kaden M. Southard[1,2,*], Nam Hyeong Kim[1,2,3,6], Ramu Gopalappa[4,8], Woon Ryoung Kim[1,2,3], Minji An[4,5], Hyun Jung Lee[1,2,3], Justin Farlow[2], Anastasios Georgakopoulos[9], Nikolaos K. Robakis[9], Daeha Seo[10], Hyeong Bum Kim[4,5,8,11,12], Yong Ho Kim[6,7], Jinwoo Cheon[4,5,13], Zev J. Gartner[2,14,†], Young-wook Jun[1,2,3,4,5†]

[1]Department of Otolaryngology, University of California, San Francisco, CA, USA

[2]Department of Pharmaceutical Chemistry, University of California, San Francisco, CA, USA

[3]Helen Diller Family Cancer Comprehensive Center (HDFCCC), University of California, San Francisco, CA, USA

[4]Center for Nanomedicine, Institute for Basic Science (IBS), Seoul, Republic of Korea

[5]Graduate Program of Nano Biomedical Engineering (Nano BME), Advanced Science Institute, Yonsei University, Seoul, Republic of Korea

[6]SKKU Advanced Institute of Nanotechnology (SAINT), Sungkyunkwan University, Suwon, Republic of Korea

[7]Department of Biomedical Engineering, Sungkyunkwan University, Suwon, Republic of Korea

[8]Department of Pharmacology, Yonsei University College of Medicine, Seoul, Republic of Korea

[9]Department of Psychiatry and Neuroscience, Icahn School of Medicine at Mount Sinai, New York, USA

[10]Department of Emerging Materials Science, DGIST, Daegu, Republic of Korea

[11]Brain Korea 21 Plus Project, Yonsei University College of Medicine, Seoul, Republic of Korea

[12]Severance Biomedical Science Institute, Yonsei University College of Medicine, Seoul, Republic of Korea

[13]Department of Chemistry, Yonsei University, Seoul, Republic of Korea

[14]Chan Zuckerberg Biohub, University of California San Francisco, CA, USA

*These authors contributed equally.

†Correspondence should be addressed to Z.J.G. (zev.gartner@ucsf.edu) or Y.J. (young-wook.jun@ucsf.edu).

Aberrant cleavage of Notch and amyloid precursor protein (APP) by γ-secretase is implicated in developmental disorders, cancer, and dementia. How cleavage is regulated in space and time is unclear. Here, we report that size-dependent protein segregation acts as a spatial switch regulating the dynamic distribution of these receptors relative to their enzymatic partner, γ-secretase. Because the size of the extracellular domain (ECD) greatly exceeds the width of the junctional cleft, Notch is excluded from cadherin-based adherens junctions (cadAJs) where γ-secretase localizes, prohibiting the enzyme-substrate interaction. ECD elimination by ADAM 10/17 drives a spatial rearrangement of Notch into cadAJs where γ-secretase cleavage occurs. Spatial mutation of Notch by decreasing (or increasing) the ECD size promotes (or inhibits) downstream signaling, respectively, regardless of ligand-presentation or S2 cleavage. Elimination of cadAJs abolishes Notch activation, confirming the indispensable role of cadAJs in regulating Notch-γ-secretase interactions. We further show that this spatial switch regulates APP cleavage and amyloid beta production. Therefore, size-dependent protein segregation creates membrane domains with unique proteolytic activities, choreographing the proteolytic sequence while suppressing nonspecific signal activation.

Notch is a highly conserved mediator for contact-dependent cell-cell communication, which orchestrates

diverse functions in metazoans[1-5]. Tight control of Notch signal activation is essential for many

developmental processes, while dysregulation of Notch activation can cause severe disease including

developmental, neurological, and immunological disorders and cancer[1-8]. Accordingly, to enable precise

signal regulation, receptor activation occurs through multiple steps, independently gated by chemical

(e.g., ligand-receptor interactions, posttranslational modifications) and mechanical cues[4,5,9-27]. However,

many signaling processes involving physical contact between two cells – so called juxtacrine signaling

processes – are also regulated by spatial cues[28-31]. As exemplified by the kinetic segregation model in

immune cells[28,32-43], spatial rearrangements of these signaling molecules modify their physical and

biochemical environments to facilitate receptor activation. Notch, which also mediates signal exchange

by physical contact, is subjected to similar structural and spatial rearrangements[44,45]. To test whether

Notch signaling is regulated by these spatial rearrangements, we investigated the spatial dynamics of

Notch and its signaling partners at cellular interfaces. Specifically, we focused on four key signaling

molecules – Notch, delta-like ligand 1 (Dll1), ADAM 10, and γ-secretase, relative to cadherin-based

adherens junctions (cadAJs); the cell-cell adhesions that initiate many structural and spatial changes by

bringing and holding the membranes together[46-49] (**Fig. 1a**).

CadAJs segregate Notch from γ-secretase, preventing interactions

To map the distribution of Notch signaling components, we generated a series of U2OS cells expressing

recombinant Notch1, Dll1, and/or epithelial cadherin (E-cadherin, Ecad) proteins. To facilitate the imaging

of these proteins, we fused them with self-labeling tags (SNAP or Halo-tags) and/or fluorescent proteins

(EGFP or mCherry) at the N- or C-termini, respectively (**Extended Data Table 1**). The endogenous Notch

processing enzymes (*i.e.* ADAM 10, and γ-secretase) were immunostained and then imaged by confocal

microscopy. With the exception of ADAM 10 which exhibited no preferential distribution relative to cadAJs

(**Extended Data Fig. 1a**), all other proteins exhibited cadAJ-dependent localization (**Fig. 1b, c**). γ-

secretase (visualized by staining with an anti-presenilin-1 antibody) was strongly enriched at the cadAJs

with negligible non-junctional membrane signal (**Fig. 1b, c, Extended Data Fig. 1b, c**)[50]. In contrast, both

Notch and Dll1 were excluded from cadAJs (**Fig. 1b, c**) and consequentially γ-secretase (**Extended Data**

Fig. 2a, b). Notch exclusion from cadAJs was observed in multiple contexts, including different cadherin

types, cell types, and cell polarization states (**Extended Data Fig. 3**). Quantitative analysis using the

Manders' overlap coefficient to calculate fractional overlap with E-cadherin also confirmed the enrichment

of γ-secretase (0.85 ± 0.21) at cadAJs and the exclusion of Notch (0.24 ± 0.19) and Dll1 (0.26 ± 0.18)

from cadAJs (**Extended Data Fig. 2c, d**). These observations suggest two mechanisms by which cadAJs

might influence Notch signaling: first, cadAJs recruit γ-secretase; second, cadAJs segregate Notch

ligands and receptors from γ-secretase to prevent their interactions.

To interrogate how cadAJs drive the spatial segregation of the enzyme (*i.e.* γ-secretase) and

substrate (*i.e.* Notch) pair, we employed our recently developed single-cell perturbation tool,

mechanogenetics, where magnetofluorescent nanoparticles (MFNs) control the location and mechanical

loading of targeted receptors hence cell signaling[26,27,51]. Specifically, we clustered E-cadherin using

mechanogenetics to generate artificial cadAJs that recapitulate the functional and signaling roles of native

cadAJs[26,27,51] (**Fig. 1d, e**; See methods for more details). We then monitored the consequence of cadAJ

formation on the spatial distributions of γ-secretase, Notch, and associated proteins (e.g. Flotillin-1, Flot1)

relative to the artificial cadAJ (**Fig. 1f, g**). Consistent with the native cell-cell cadAJs, γ-secretase was

localized at the artificial cadAJ (**Fig. 1f and Extended Data Fig. 4a, b**). We also observed colocalization

of Flot1, a protein enriched in spatially discrete and ordered membrane microdomains at the artificial

cadAJs[52-56] (**Fig. 1f and Extended Data Fig. 4a, b**). This observation, along with analysis of native cell-

cell junctions (**Extended Data Fig. 4, c-g**), molecular dynamics (MD) simulation (**Extended Data Fig. 4,**

h-j), and cholesterol depletion experiments (**Extended data Figs. 4k**), suggests that the cadAJs recruit

and stabilize γ-secretase through a common spatially discrete and ordered membrane microdomains (for

more discussion, see **Supplementary Discussion**).

Surprisingly, we observed an intense Notch signal at the artificial cadAJ (**Fig. 1g and Extended**

Data Fig. 5), in contrast to the depleted Notch signal at native cell-cell cadAJs (**Fig. 1, a-c**). The

extraordinarily large size (extended length = 136 nm[12], compacted height by electron microscopy > 20

nm[57]) of the Notch extracellular domain (NECD) suggested a potential explanation for these contradicting

observations. Specifically, Notch could be excluded from cadAJs due to the NECD size greatly exceeding

the narrow intermembrane cleft created by native cadAJs (20 nm)[47,58]. On the other hand, artificial cadAJs

generated by MFNs are free of membrane juxtaposition and hence allow for Notch diffusion and

accumulation, presumably through its association with γ-secretase or other components of the membrane

microdomains[59]. These observations fit a model wherein the size-based physical constraint caused by

cell-cell cadAJs segregates Notch from γ-secretase, preventing interactions that would otherwise serve

to colocalize and concentrate the enzyme-substrate pair.

Size-dependent segregation choreographs the Notch proteolytic sequence

Upon activation by juxtacrine ligand-receptor interactions, Notch is processed by ADAM 10/17 (S2

cleavage) and then by γ-secretase (S3 cleavage), sequentially[1-5]. Since each of proteolytic events leads

to a dramatic reduction in the size of NECD, we investigated how the size changes during activation

correlate with its spatial distribution. We plated U2OS cells co-expressing SNAP-N^{FL}-mCherry and Ecad-

GFP on a Dll4-coated substrate to trigger Notch activation. Cells were also treated with TAPI2 (an ADAM

10/17 inhibitor which stabilizes full-length Notch) or DAPT (a γ-secretase inhibitor that stabilizes the S2-

cleaved Notch having a minimal extracellular domain) to capture Notch intermediates (**Fig. 2a**). To

quantify the spatial redistribution of each Notch intermediate relative to the cadAJ, we measured the

average mCherry fluorescence signal inside (I_{IN}) and outside (I_{OUT}) of the cadAJ and estimated an

enrichment ratio (I_{IN}/I_{OUT}, See Methods). An intensity ratio of 1 represents no cadAJ-dependent

distribution as validated with DiI dyes (I_{IN}/I_{OUT} = 1.07, **Fig. 2c and Extended Data 6a**), and values less

than or greater than 1 represent exclusion from or enrichment within cadAJs, respectively. With TAPI2, we observed minimal mCherry fluorescence signal from cadAJs (I_{IN}/I_{OUT} = 0.50, **Fig. 2b**(ii)**, c**(ii)), indicating that the ligand-receptor interaction did not alter the spatial distribution of Notch before S2 cleavage. In contrast, when TAPI2 was removed to activate S2 cleavage and DAPT was added to inhibit S3 cleavage, we observed strong enrichment of the mCherry signal at cadAJs (I_{IN}/I_{OUT} = 2.14, **Fig. 2b**(iii)**, c**(iii)), indicative of relocalization and concentration of the Notch extracellular domain truncation (NEXT), the product of S2 cleavage, into cadAJs. When γ-secretase activity was rescued by washing out DAPT, the initially intense mCherry signal at cadAJs gradually disappeared (**Fig. 2b**(iv)**, c**(iv)**, and Extended Data Fig. 6b, c**), presumably corresponding to release of Notch intracellular domain (NICD) from the membrane. These results suggest a role for the size-dependent protein segregation as a spatial switch that regulates the distribution of Notch intermediates, thereby choreographing the sequential steps in Notch proteolysis. According to this model, the mass of NECD poses a physical constraint preventing entry of Notch to the narrow space between membranes in the cadAJ cleft where γ-secretase is localized. Hence, γ-secretase cannot process the full-length Notch before S2 cleavage. Following S2 cleavage, removal of NECD relieves the physical constraint, allowing Notch to enter into the cadAJ cleft. This facilitates a productive Notch-γ-secretase interaction[59], S3 cleavage, and then downstream signaling.

To explore how size-dependent segregation controls Notch signaling, we first generated a series of U2OS cell lines stably expressing Notch variants with different truncation lengths: a partial deletion of the EGF repeats (NΔEGF$_{1-25}$, approximate height: 48 nm), complete deletion of the EGF repeats but retention of negative regulatory region (NΔEGF, approximate height: 10 nm), and a complete removal of NECD (NEXT, approximate height: 4 nm) (**Fig. 2d**). We fused these Notch variants with SNAP- and mCherry-tags at their N- and C-termini, to differentially image the extracellular and intracellular domains. All cells were treated with TAPI2 and DAPT to prevent any potential proteolysis of the variants. Consistent with predictions based on size-dependent protein segregation, NΔEGF$_{1-25}$, the Notch variant with an ECD taller than the height of the intermembrane cadAJ cleft, was excluded from cadAJs (I_{IN}/I_{OUT} = 0.57) (**Fig. 2e, f**). NEXT with an ECD smaller than the junctional height was enriched at cadAJs (I_{IN}/I_{OUT} = 2.39) (**Fig. 2e, f**). Interestingly, we observed binary localizations of NΔEGF (intermediate height) relative to cadAJs, with a mean I_{IN}/I_{OUT} value of 1.32 (**Fig. 2e, f**). Some cadAJs displayed NΔEGF enrichment (**Fig. 2e**(bottom left), **f**(right)), consistent with the size-based prediction. Meanwhile, other cadAJs excluded NΔEGF (**Fig. 2e**(bottom right)**, f**(right)). Considering the fact that cadherin cluster densities within cadAJs vary with the size, type, and degree of junction maturation[60,61] and that the glycosylated negative regulatory region domain[62] is susceptible to steric crowding, this unanticipated exclusion might result from a lateral crowding effect in high-density cadAJs.

We then investigated the functional consequences of NΔEGF relocalization to cadAJs by measuring the impact on downstream signaling process. Focusing on the subset of cadAJs showing strong enrichment for Notch signal, we inferred γ-secretase activity by measuring the changes in extracellular SNAP (labeled with SNAP surface dye) and intracellular mCherry fluorescence signals of

NΔEGF after DAPT removal while maintaining TAPI2 concentration. This construct has different
fluorescence markers on the extracellular and intracellular domains of the protein, allowing us to
differentially map the S2 and S3 cleavage events of the construct. While SNAP fluorescence signal
remained strong, the mCherry signal at the cadAJ was negligible (**Fig. 2g**). Similarly, colocalization
analysis of multiple cadAJs using Pearson correlation coefficients showed more significant signal
reduction in the mCherry channel than in SNAP, indicating selective release of NICD from cadAJs (**Fig.**
2h). We measured the kinetics of S3 processing by tracking mCherry intensity at the cadAJ in single cells.
Initially intense mCherry signals (average I_{IN}/I_{OUT} = 2.89 ± 1.15) rapidly dissipated within the first 2 hours
following DAPT removal (average I_{IN}/I_{OUT} = 0.81 ± 0.42), reaching a plateau at 4 hours (I_{IN}/I_{OUT} = 0.65 ±
0.14) (**Fig. 2i, j, and Extended Data Fig. 6d, e**). Removing DAPT did not elicit a significant reduction in
mCherry signal intensity from the non-cadAJ membrane, indicating that the S3 cleavage activity was
strongly localized at the cadAJ (**Extended Data Fig. 6f**). To confirm that the observed decrease in
mCherry signal at the cadAJ corresponds to the S3 cleavage, we performed western blot analysis of the
cells expressing the Notch variants. We cultured cells with TAPI2 to decouple γ-secretase processing
from S2 cleavage, and measured cleaved NICD levels by immunoblotting with Notch anti-V1744
antibodies. Cells expressing N^{FL} or NΔEGF$_{1-25}$ resulted in no or minimal NICD, respectively (**Fig. 2k, l**).
Whereas, cells expressing NΔEGF produced a significant amount of NICD (**Fig. 2k, l**). Cells expressing
NEXT exhibited the highest NICD production, about a 4-fold increase compared with that of NΔEGF (**Fig.**
2k, l). The observed NICD production levels were proportional to the enrichment ratio (I_{IN}/I_{OUT}) of the
Notch variants at cadAJs, suggesting the essential role of size-dependent protein segregation as a spatial
switch that controls Notch activation. Additionally, the substantial NICD production from the cells
expressing NΔEGF indicates that, when localized together, γ-secretase can process Notch, bypassing
S2 cleavage. Ligand-independent activation of Notch receptors with an intact S2 site was observed
previously in Notch truncation variants or synNotch constructs[63-65], but the mechanism of this activation
has been unclear so far. Our observations support the notion that colocalization of Notch with γ-secretase
is sufficient to trigger S3 proteolysis and signaling.

Spatial mutation alters Notch signaling
Prevailing models of Notch proteolysis by γ-secretase are based on the notion that S2-cleavage of Notch
serves to potentiate the cleavage by modifying the molecular interface at the enzyme-substrate pair[66,67].
For example, it has been suggested that γ-secretase selectively recognizes S2-cleaved Notch (*i.e.,* NEXT)
through hydrogen bonding between a glutamate residue in nicastrin and the new N-terminus of NEXT[66].
Another model proposed that S2 cleavage serves to reduce steric repulsion between nicastrin and NECD,
strengthening their interaction[67]. However, another key feature of S2 cleavage is that it generates a
smaller molecular intermediate that can uniquely access cadAJs, thereby colocalizing Notch with γ-
secretase and significantly increasing its concentration near the enzyme active site. This suggests a third

model, wherein γ-secretase activity on full length Notch and its intermediates is blocked by maintaining
Notch concentrations below the K_M for γ-secretase through spatial segregation.

195   To explicitly test the effect of Notch spatial localization relative to cadAJs and γ-secretase on the
signaling, we designed two experiments that induce spatial mutation of Notch. First, we employed a DNA-
mediated crosslinking strategy to enhance NΔEGF – a Notch variant that exhibited binary localizations
relative to cadAJs and relatively low Notch activation – enrichment at the cadAJ. We generated cells co-
expressing SNAP-NΔEGF-mCherry and Halo-Ecad-GFP, and treated the cells with complementary BG-
and chloroalkane-modified oligonucleotides in the presence of TAPI2 and DAPT (**Fig. 3a**). Notch-E-
cadherin heterodimers were formed efficiently as evidenced by the appearance of a higher molecular
weight band corresponding to the DNA-linked complex on western blots (**Extended Data Fig. 7a**).
Compared to untreated cells (I_{IN}/I_{OUT} = 1.32 ± 1.06), we observed further enrichment of NΔEGF at cadAJs
in the presence of DNA crosslinking (I_{IN}/I_{OUT} = 1.89 ± 0.91) (**Fig. 3b**). We then maintained cells in TAPI2
but removed DAPT to allow S3 cleavage. We observed decreases in mCherry signal at cadAJs after
DAPT removal, indicating efficient S3 cleavage without S2 cleavage (**Extended Data Fig. 7b-d**).
Interestingly, in western blots, we observed increased V1744-terminated NICD levels from the cells
treated with DNA crosslinkers, compared with the untreated control (**Fig. 3c**). There results suggest that
new molecular interfaces produced by mechanical activation and S2 cleavage are not necessary when
γ-secretase is concentrated with its substrate. Considering that DNA crosslinking (molecular weight =
21.4 kD) increases the ECD size of NΔEGF, the observed increase in NICD production cannot be
explained by the nicastrin-induced steric repulsion model. Hence, we concluded the result that increased
concentration of the NΔEGF at cadAJs facilitated its interaction with γ-secretase and promoted S3
cleavage.

215   To further test the importance of size-dependent spatial segregation, we induced spatial mutation
of NEXT (*i.e.* the Notch variant that showed the strongest enrichment at the cadAJ and activation) by
chemically conjugating it with macromolecules. Specifically, we conjugated BG-modified polymers and
proteins, to the extracellular SNAP tag (4.0 nm) of the variant via BG-SNAP chemistry (**Fig. 3d**). Grafting
of these macromolecular pendants onto NEXT increases the size of the Notch construct but does not
modify the N-terminal amine accessibility for hydrogen bonding with nicastrin. To interrogate the size-
dependent spatial mutation of NEXT systematically, we used a series of pendants with different
hydrodynamic sizes, that includes polyethylene glycol with an average molecular weight of 3.4 kD
(PEG3.4k, 2.5 nm), branched PEG20k (*b*PEG20k, 4.0 nm), linear PEG20k (ℓPEG20k, 8.0 nm), DNA-
streptavidin conjugates (DNA-stv, 9.5 nm), and human immunoglobulin G (hIgG, 12 nm) (**Fig. 3d and**
Extended Data Fig. 8a). In the presence of DAPT, we observed a size-dependent distribution of NEXT
at the cadAJ, where the larger pendant resulted in a greater decrease in mCherry signal at the cadAJ.
With pendants smaller than 5 nm (*i.e.* PEG3.4k and *b*PEG20*k*), NEXT remained enriched at the cadAJ
with I_{IN}/I_{OUT} of 2.21 and 2.01, respectively (**Fig. 3d, e**). When ℓPEG20k, DNA-stv, or hIgG were added,
we observed binary localizations of NEXT that were enriched at or excluded from the cadAJs, with mean

I_{IN}/I_{OUT} values of 1.06, 0.82, 0.98, respectively (**Fig. 3d, e**). These observations were similar to the spatial

behaviors of NΔEGF having a comparable ECD size, where only less dense cadAJs allowed Notch

colocalization. We then examined the signaling consequences of each size-dependent spatial mutation

of NEXT. Following S3 cleavage, NICD traffics to the nucleus, allowing us to measure nuclear mCherry

signal as a proxy for Notch pathway activation. The PEG3.4k or bPEG20k addition did not significantly

alter nuclear mCherry signal of NEXT, compared with cells with no pendant addition (**Fig. 3f, g and**

Extended Fig. 9b). Conjugation of ℓPEG20k, DNA-stv (9.5 nm) resulted in substantial decrease in

nuclear mCherry signal to 0.39 and 0.37, respectively (**Fig. 3f, g and Extended Fig. 8b**). hIgG addition

slightly further suppressed nuclear mCherry signal to 0.27 (**Fig. 3f, g and Extended Fig. 8b**). Finally, we

summarized the S3-cleaved NICD production of all Notch variants as a function of the Notch enrichment

factor, I_{IN}/I_{OUT}, in **Fig. 3h**, clearly visualizing the spatial dependence of S3 cleavage.

The cadAJ-mediated spatial switch is indispensable for Notch signaling

Given the significant role of cadAJs in coordinating the spatial dynamics of Notch and γ-secretase, we

next interrogated Notch signal activation in cells lacking cadAJs. To minimize physical contact between

cells and hence cadAJ formation, we sparsely plated cells expressing SNAP-Notch-Gal4 (SNAP-N^{FL}-

Gal4) receptors on a Dll4-coated substrate. We used a UAS-Gal4 reporter system that detects Notch

activation with nuclear mCherry signal[15,23,26]. After 16 hours from cell seeding, we measured the mCherry

fluorescence in cells having no prior contact with other cells. For comparison, we also analyzed the

mCherry signal in cells with robust cadAJs within high-density culture. While the cells with physical

contacts with other cells exhibited robust increase in nuclear mCherry fluorescence signal, those without

cell-cell contact elicited no increase in signal (**Fig. 4a, b and Supplementary Video 1**). Reestablishing

the cadAJs by plating cells on a substrate coated with Ecad-Fc and Dll4-Fc rescued the Notch signaling

of the solitary cells (**Fig. 4a, c, and Supplementary Video 2**). These results support that cadAJs are

required for Notch processing at the cell surface and downstream signaling. Critically, E-cadherin seems

to function in this capacity in a manner that is independent of its role in mediating cell-cell contact. To

further test this notion, we knocked out the gene encoding E-cadherin (CDH1) in the reporter cell line via

CRISPR-Cas9, then plated the cells at high density on Dll4-Fc coated plates (**Extended Data Fig 9**, **a-**

f). Strikingly, E-cadherin knockout (Ecad-KO) resulted in abrogation of Notch activation even with robust

cell-cell contact (**Fig. 4d**, **e, and Extended Data Fig. 9g**). Reintroduction of plasmids encoding E-

cadherin or N-cadherin into Ecad-KO cells recovered Notch activation with substantial nuclear mCherry

signal, confirming the indispensable role of cadAJs in Notch signaling (**Fig. 4d, e and Extended Data**

Fig. 9g).

Size-dependent spatial dynamics and proteolysis of amyloid precursor proteins

The proteolysis processes of amyloid precursor protein (APP) plays a central role in amyloid beta (Aβ)

pathology, which can cause failures in many organs such as brain, heart, kidney, and vasculatures[68-71].

Interestingly, APP has a strikingly similar topology and proteolytic cleavage sequence to that of Notch.

APP possesses a large ECD (> 20 nm) and is processed by α- or β-secretase and then γ-secretase[68-71].
These characteristics motivated us to investigate the role of the size-dependent protein segregation in
APP processing. We generated U2OS cells co-expressing APP-GFP and SNAP-N-cadherin (SNAP-
Ncad) and monitored the cell surface spatial dynamics of APP intermediates relative to N-cadherin based
AJs (NcadAJs) using proteolytic enzyme inhibitors. Similar to Notch, we found that APP was initially
excluded from NcadAJs, diffused into the NcadAJs after the ECD shedding by α- or β-secretase, and
then was processed by γ-secretase within it (**Fig. 4f**).

APP can be processed along two pathways by γ-secretase – one pathway occurs at the cell
surface to generate more soluble isoforms including p3 or $A\beta_{40}$, the other pathway occurs internally to
produce less soluble and pathogenic $A\beta_{42}$[68-73]. Given our previous observation that loss of cadAJs leads
to a decrease in cell surface γ-secretase, we hypothesized that APP processing would be biased under
these conditions towards $A\beta_{42}$. We tested this hypothesis by constructing U2OS cell lines lacking both E-
and N-cadherins (cad-KO cells) using CRISPR-Cas9 (**Extended Data Fig. 9h, i**). We then transfected
plain U2OS cells or cad-KO cells with a plasmid encoding APP sequence and measured Aβ production
by ELISA. Consistent with the hypothesis, cad-KO cells produced higher $A\beta_{42}$, the isoform prone to
severe fibril aggregation, compared to cells with endogenous cadherin expression (**Fig. 4g**).

Discussion
Notch displays several unique features that are distinct from other juxtacrine receptors. For example,
unlike classical receptor signaling, the Notch ligand-receptor interaction (chemical switch) is converted
into intracellular signaling through a sequence of additional signaling switches acting by mechanical,
enzymatic, and spatial mechanisms. These include unfolding of the negative regulatory region domain
(mechanical switch), S2- and S3- cleavages (enzymatic switch), and finally translocation of the NICD
from the cell membrane to the nucleus (spatial switch)[4,5]. Our study reveals that Notch additionally
integrates a spatial switch at the cell surface to tightly choreograph the enzymatic cleavage sequence
prior to NICD release. Previously, it was thought that this enzymatic sequence was regulated by
modification of the molecular interface between Notch and nicastrin after S2-cleavage[66,67]. Our model
does not rule out the contribution of the nicastrin-Notch chemical interface during this process, but our
study strongly suggests that the spatial switch is the major regulator of Notch-γ-secretase interaction and
signaling, functioning by increasing the concentration of the γ-secretase substrate to the point that it is
efficiently processed by the enzyme. Particularly, our mechanogenetic and the NΔEGF cleavage
experiment shown in Figure 1(d-g) and 2(e-l), respectively, support the notion that Notch with an intact
S2 site is effectively engaged and then processed by γ-secretase when the spatial constraints are
removed.

The operating principle of the spatial switch is closely related to another unique feature of Notch
receptor: the unusually tall NECD. The functional residues responsible for ligand binding are located near
the N-terminus, which protrudes above the crowded cell surface, where they are poised to engage ligands

on neighboring cells. Surprisingly, however, it has been also shown that replacing the EGF-like domain

repeats with a relatively smaller ligand binding domain (e.g. synNotch) maintains the receptor function[74,75].

Why then does Notch receptor bear such a massive ECD? Our study provides insight into this question,

where the large ECD is crucial for its spatial segregation from γ-secretase thereby minimizing nonspecific

ligand-independent activation. Low level NICD production was observed even for the Notch variants with

a partial EGF truncation (NΔEGF$_{1-25}$), and levels gradually increased upon successive truncations to the

NECD size. Our model also explains previous observations where synNotch with a relatively small ECD

exhibited significant ligand-independent activation (10-50% of ligand-induced activation)[64].

We also show that the size-dependent spatial segregation regulates APP cleavage and Aβ

production. It has been previously shown that γ-secretase presenting in different subcellular

compartments cleaves APP into diverse Aβ isoforms[70,73,76-78]. Our study shows that, after the ECD

cleavage, cadAJ potentiates cell surface processing of APPs within the junction, yielding Aβ_{40}

predominantly, while removal of cadAJ produces more Aβ_{42}. To establish the relevance of this

observation to APP processing will require further investigation in a neuronal system, but our results in

model cell lines are consistent with the predominant secretion of Aβ from synapse, where N-cadherin

junctions localize[79,80].

The spatial switch is analogous to the kinetic segregation model of T cell activation, where large

CD45 phosphatase is excluded from T cell receptor (TCR) immunological synaptic clefts[28,32-42]. However,

there are several distinct features of the Notch spatial switch compared to the kinetic segregation model.

First, unlike the immunological synapse where TCR and CD45 remain constant in size throughout

activation, Notch undergoes a dramatic size change during the course of cell surface activation, enabling

its dynamic spatial redistribution and sequential proteolysis. Second, the role of cadAJs in Notch signaling

is not limited to creating a physical barrier, but also plays the critical role of recruiting γ-secretase to

facilitate processing of S2-cleaved Notch at the cell surface. Third and finally, the consequences of size-

dependent segregation on signaling are reversed compared to the immunological synapse. While spatial

segregation of CD45 enables sustained TCR phosphorylation and downstream signaling, Notch

segregation from cadAJs inhibits signal activation. Our result extends the relevance of size-dependent

spatial segregation models beyond immune cells, [28,32-43], supporting the notion that the size-dependent

protein segregation can serve as a general mechanism for regulating a broad range of receptor signaling

at the cell-cell interface, including Notch and APPs.

Overall, a spatial switching mechanism based on size-dependent protein segregation not only

sheds light on the mechanism underlying the sequential proteolysis of Notch and APPs, but also may

extend to other receptors processed by γ-secretase. Finally, we anticipate further implications of our work

in other areas of research such as providing new design principles for synthetic receptors like synNotch,

as well as new therapeutic approaches that target Notch and APP signaling by spatial mutation in cancer

and neurodegenerative diseases.

1.   Artavanis-Tsakonas, S., Rand, M. D. & Lake, R. J. Notch Signaling: Cell Fate Control and Signal
Integration in Development. *Science* **284,** 770–776 (1999).
2.   Fortini, M. E. Notch signaling: the core pathway and its posttranslational regulation. *Dev. Cell* **16,**
633–647 (2009).
3.   Louvi, A. & Artavanis-Tsakonas, S. Notch signalling in vertebrate neural development. *Nat. Rev.*
Neurosci. **7,** 93–102 (2006).
4.   Bray, S. J. Notch signalling in context. *Nat. Rev. Mol. Cell Biol.* **17,** 722–735 (2016).
5.   Kovall, R. A., Gebelein, B., Sprinzak, D. & Kopan, R. The Canonical Notch Signaling Pathway:
Structural and Biochemical Insights into Shape, Sugar, and Force. *Dev. Cell* **41,** 228–241 (2017).
6.   Koch, U. & Radtke, F. Notch and cancer: a double-edged sword. *Cell. Mol. Life Sci.* **64,** 2746–
2762 (2007).
7.   South, A. P., Cho, R. J. & Aster, J. C. The double-edged sword of Notch signaling in cancer.
Semin. Cell Dev. Biol. **23,** 458–464 (2012).
8.   Lai, E. C. Notch signaling: control of cell communication and cell fate. *Development* **131,** 965–
973 (2004).
9.   Luca, V. C. *et al.* Notch-Jagged complex structure implicates a catch bond in tuning ligand
sensitivity. *Science* **355,** 1320–1324 (2017).
10.   Haines, N. & Irvine, K. D. Glycosylation regulates Notch signalling. *Nat. Rev. Mol. Cell Biol.* **4,**
786–797 (2003).
11.   D'Souza, B., Miyamoto, A. & Weinmaster, G. The many facets of Notch ligands. *Oncogene* **27,**
5148–5167 (2008).
12.   Luca, V. C. *et al.* Structural biology. Structural basis for Notch1 engagement of Delta-like 4.
Science **347,** 847–853 (2015).
13.   Moloney, D. J. *et al.* Fringe is a glycosyltransferase that modifies Notch. *Nature* **406,** 369–375
(2000).
14.   Brückner, K., Perez, L., Clausen, H. & Cohen, S. Glycosyltransferase activity of Fringe
modulates Notch-Delta interactions. *Nature* **406,** 411–415 (2000).
15.   Sprinzak, D. *et al.* Cis-interactions between Notch and Delta generate mutually exclusive
signalling states. *Nature* **465,** 86–90 (2010).
16.   Nandagopal, N. *et al.* Dynamic Ligand Discrimination in the Notch Signaling Pathway. *Cell* **172,**
869–880.e19 (2018).
17.   Lebon, L., Lee, T. V., Sprinzak, D., Jafar-Nejad, H. & Elowitz, M. B. Fringe proteins modulate
Notch-ligand cis and trans interactions to specify signaling states. *Elife* **3,** e02950 (2014).
18.   Shaya, O. *et al.* Cell-Cell Contact Area Affects Notch Signaling and Notch-Dependent Patterning.
Dev. Cell **40,** 505–511.e6 (2017).
19.   Weinmaster, G. & Fischer, J. A. Notch Ligand Ubiquitylation: What Is It Good For? *Dev. Cell* **21,**
134–144 (2011).

20.   Baron, M. *et al.* Multiple levels of Notch signal regulation. *Mol. Membr. Biol.* **19,** 27–38 (2002).

21.   Yamamoto, S. *et al.* A mutation in EGF repeat-8 of Notch discriminates between Serrate/Jagged
and Delta family ligands. *Science* **338,** 1229–1232 (2012).

22.   Wang, X. & Ha, T. Defining single molecular forces required to activate integrin and notch
signaling. *Science* **340,** 991–994 (2013).

23.   Gordon, W. R. *et al.* Mechanical Allostery: Evidence for a Force Requirement in the Proteolytic
Activation of Notch. *Dev. Cell* **33,** 729–736 (2015).

24.   Meloty-Kapella, L., Shergill, B., Kuon, J., Botvinick, E. & Weinmaster, G. Notch ligand
endocytosis generates mechanical pulling force dependent on dynamin, epsins, and actin. *Dev.*
Cell **22,** 1299–1312 (2012).

25.   Shergill, B., Meloty-Kapella, L., Musse, A. A., Weinmaster, G. & Botvinick, E. Optical tweezers
studies on Notch: single-molecule interaction strength is independent of ligand endocytosis. *Dev.*
Cell **22,** 1313–1320 (2012).

26.   Seo, D. *et al.* A Mechanogenetic Toolkit for Interrogating Cell Signaling in Space and Time. *Cell*
165, 1507–1518 (2016).

27.   Kim, J.-W. *et al.* Single-cell mechanogenetics using monovalent magnetoplasmonic
nanoparticles. *Nat. Protoc.* **12,** 1871–1889 (2017).

28.   Bell, G. I. Models for the specific adhesion of cells to cells. *Science* **200,** 618–627 (1978).

29.   Manz, B. N. & Groves, J. T. Spatial organization and signal transduction at intercellular junctions.
Nat. Rev. Mol. Cell Biol. **11,** 342–352 (2010).

30.   Schmid, E. M. *et al.* Size-dependent protein segregation at membrane interfaces. *Nat. Phys.* **12,**
704–711 (2016).

31.   Salaita, K. *et al.* Restriction of receptor movement alters cellular response: physical force
sensing by EphA2. *Science* **327,** 1380–1385 (2010).

32.   Dustin, M. L. & Shaw, A. S. Costimulation: Building an Immunological Synapse. *Science* **283,**
649–650 (1999).

33.   Aricescu, A. R. *et al.* Structure of a tyrosine phosphatase adhesive interaction reveals a spacer-
clamp mechanism. *Science* **317,** 1217–1220 (2007).

34.   Monks, C. R., Freiberg, B. A., Kupfer, H., Sciaky, N. & Kupfer, A. Three-dimensional segregation
of supramolecular activation clusters in T cells. *Nature* **395,** 82–86 (1998).

35.   Dustin, M. L. Making a little affinity go a long way: a topological view of LFA-1 regulation. *Cell*
Adhes. Commun. **6,** 255–262 (1998).

36.   Grakoui, A. *et al.* The immunological synapse: a molecular machine controlling T cell activation.
Science **285,** 221–227 (1999).

37.   James, J. R. & Vale, R. D. Biophysical mechanism of T-cell receptor triggering in a reconstituted
system. *Nature* **487,** 64–69 (2012).

38. Cordoba, S.-P. *et al.* The large ectodomains of CD45 and CD148 regulate their segregation from
and inhibition of ligated T-cell receptor. *Blood* **121,** 4295–4302 (2013).
39. Dustin, M. L. & Groves, J. T. Receptor signaling clusters in the immune synapse. *Annu. Rev.*
Biophys. **41,** 543–556 (2012).
40. Wülfing, C. & Davis, M. M. A receptor/cytoskeletal movement triggered by costimulation during T
cell activation. *Science* **282,** 2266–2269 (1998).
41. Choudhuri, K., Wiseman, D., Brown, M. H., Gould, K. & van der Merwe, P. A. T-cell receptor
triggering is critically dependent on the dimensions of its peptide-MHC ligand. *Nature* **436,** 578–
582 (2005).
42. Groves, J. T. & Kuriyan, J. Molecular mechanisms in signal transduction at the membrane. *Nat.*
Struct. Mol. Biol. **17,** 659–665 (2010).
43. Bakalar, M. H. *et al.* Size-Dependent Segregation Controls Macrophage Phagocytosis of
Antibody-Opsonized Targets. *Cell* **174,** 131–142.e13 (2018).
44. Luty, W. H., Rodeberg, D., Parness, J. & Vyas, Y. M. Antiparallel segregation of notch
components in the immunological synapse directs reciprocal signaling in allogeneic Th: DC
conjugates. *The Journal of Immunology* **179,** 819–829 (2007).
45. Narui, Y. & Salaita, K. Membrane tethered delta activates notch and reveals a role for spatio-
mechanical regulation of the signaling pathway. *Biophys. J.* **105,** 2655–2665 (2013).
46. Yap, A. S. & Kovacs, E. M. Direct cadherin-activated cell signaling: a view from the plasma
membrane. *J. Cell Biol.* **160,** 11–16 (2003).
47. Harrison, O. J. *et al.* The extracellular architecture of adherens junctions revealed by crystal
structures of type I cadherins. *Structure* **19,** 244–256 (2011).
48. Adams, C. L., Chen, Y. T., Smith, S. J. & Nelson, W. J. Mechanisms of epithelial cell-cell
adhesion and cell compaction revealed by high-resolution tracking of E-cadherin-green
fluorescent protein. *J. Cell Biol.* **142,** 1105–1119 (1998).
49. Chen, C. S., Tan, J. & Tien, J. Mechanotransduction at cell-matrix and cell-cell contacts. *Annu.*
Rev. Biomed. Eng. **6,** 275–302 (2004).
50. Kouchi, Z. *et al.* p120 catenin recruits cadherins to gamma-secretase and inhibits production of
Abeta peptide. *J. Biol. Chem.* **284,** 1954–1961 (2009).
51. Kwak, M. *et al.* Small, Clickable, and Monovalent Magnetofluorescent Nanoparticles Enable
Mechanogenetic Regulation of Receptors in a Crowded Live-Cell Microenvironment. *Nano Lett.*
19, 3761–3769 (2019).
52. Causeret, M., Taulet, N., Comunale, F., Favard, C. & Gauthier-Rouvière, C. N-cadherin
association with lipid rafts regulates its dynamic assembly at cell-cell junctions in C2C12
myoblasts. *Mol. Biol. Cell* **16,** 2168–2180 (2005).

53.   Taulet, N. *et al.* N-cadherin/p120 catenin association at cell-cell contacts occurs in cholesterol-
447   rich membrane domains and is required for RhoA activation and myogenesis. *J. Biol. Chem.*
448   **284,** 23137–23145 (2009).

449   54.   Yap, A. S., Gomez, G. A. & Parton, R. G. Adherens Junctions Revisualized: Organizing
450   Cadherins as Nanoassemblies. *Dev. Cell* **35,** 12–20 (2015).

451   55.   Guillaume, E. *et al.* Flotillin microdomains stabilize cadherins at cell-cell junctions. *J. Cell. Sci.*
452   **126,** 5293–5304 (2013).

453   56.   Vetrivel, K. S. *et al.* Spatial segregation of gamma-secretase and substrates in distinct
454   membrane domains. *J. Biol. Chem.* **280,** 25892–25900 (2005).

455   57.   Kelly, D. F. *et al.* Molecular structure and dimeric organization of the Notch extracellular domain
456   as revealed by electron microscopy. *PLoS ONE* **5,** e10532 (2010).

457   58.   Brasch, J., Harrison, O. J., Honig, B. & Shapiro, L. Thinking outside the cell: how cadherins drive
458   adhesion. *Trends Cell Biol.* **22,** 299–310 (2012).

459   59.   Polacheck, W. J. *et al.* A non-canonical Notch complex regulates adherens junctions and
460   vascular barrier function. *Nature* **552,** 258–262 (2017).

461   60.   Indra, I. *et al.* Spatial and temporal organization of cadherin in punctate adherens junctions.
462   *Proc. Natl. Acad. Sci. U.S.A.* **115,** E4406–E4415 (2018).

463   61.   Takeichi, M. Dynamic contacts: rearranging adherens junctions to drive epithelial remodelling.
464   *Nat. Rev. Mol. Cell Biol.* **15,** 397–410 (2014).

465   62.   Takeuchi, H. & Haltiwanger, R. S. Role of glycosylation of Notch in development. *Semin. Cell*
466   *Dev. Biol.* **21,** 638–645 (2010).

467   63.   Palmer, W. H. & Deng, W.-M. Ligand-Independent Mechanisms of Notch Activity. *Trends Cell*
468   *Biol.* **25,** 697–707 (2015).

469   64.   Yang, Z.-J., Yu, Z.-Y., Cai, Y.-M., Du, R.-R. & Cai, L. Engineering of an enhanced synthetic
470   Notch receptor by reducing ligand-independent activation. *Commun. Biol.* **3,** 116 (2020).

471   65.   Mumm, J. S. *et al.* A ligand-induced extracellular cleavage regulates gamma-secretase-like
472   proteolytic activation of Notch1. *Mol. Cell* **5,** 197–206 (2000).

473   66.   Shah, S. *et al.* Nicastrin functions as a gamma-secretase-substrate receptor. *Cell* **122,** 435–447
474   (2005).

475   67.   Bolduc, D. M., Montagna, D. R., Gu, Y., Selkoe, D. J. & Wolfe, M. S. Nicastrin functions to
476   sterically hinder γ-secretase-substrate interactions driven by substrate transmembrane domain.
477   *Proc. Natl. Acad. Sci. U.S.A.* **113,** E509–18 (2016).

478   68.   Sala Frigerio, C. & De Strooper, B. Alzheimer's Disease Mechanisms and Emerging Roads to
479   Novel Therapeutics. *Annu. Rev. Neurosci.* **39,** 57–79 (2016).

480   69.   O'Brien, R. J. & Wong, P. C. Amyloid precursor protein processing and Alzheimer's disease.
481   *Annu. Rev. Neurosci.* **34,** 185–204 (2011).

70.   Selkoe, D. J. & Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol.*
Med. **8,** 595–608 (2016).

71.   Pepys, M. B. Amyloidosis. *Annu. Rev. Med.* **57,** 223–241 (2006).

72.   Ehehalt, R., Keller, P., Haass, C., Thiele, C. & Simons, K. Amyloidogenic processing of the
Alzheimer beta-amyloid precursor protein depends on lipid rafts. *J. Cell Biol.* **160,** 113–123
(2003).

73.   De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164,** 603–615
(2016).

74.   Morsut, L. *et al.* Engineering Customized Cell Sensing and Response Behaviors Using Synthetic
Notch Receptors. *Cell* **164,** 780–791 (2016).

75.   Engineering T Cells with Customized Therapeutic Response Programs Using Synthetic Notch
Receptors. *Cell* **167,** 419–432.e16 (2016).

76.   Chow, V. W., Mattson, M. P., Wong, P. C. & Gleichmann, M. An overview of APP processing
enzymes and products. *Neuromolecular Med.* **12,** 1–12 (2010).

77.   Yang, G. *et al.* Structural basis of Notch recognition by human γ-secretase. *Nature* **565,** 192–197
(2019).

78.   Bolduc, D. M., Montagna, D. R., Seghers, M. C., Wolfe, M. S. & Selkoe, D. J. The amyloid-beta
forming tripeptide cleavage mechanism of γ-secretase. *Elife* **5,** 08928 (2016).

79.   Yuksel, M. & Tacal, O. Trafficking and proteolytic processing of amyloid precursor protein and
secretases in Alzheimer's disease development: An up-to-date review. *Eur. J. Pharmacol.* **856,**
172415 (2019).

80.   Arikkath, J. N-cadherin: stabilizing synapses. *J. Cell Biol.* **189,** 397–398 (2010).

Acknowledgement

The authors thank Drs. Stephen Blacklow (Harvard U.) and C. Miller (King's College London) for the kind gifts of Notch and APP plasmids, respectively. We also thank Dr. Daniel Fletcher (UC Berkeley), Mr. Ari Joffe (UC Berkeley), and Dr. Duaa Al-Rawi (Stanford U.) for helpful discussion. For reagents, technical supports, and discussions we thank the Kim, Cheon, Gartner, and Jun laboratories, as well as the Nikon Imaging Center and Wynton at UCSF. M.K. was supported by a Life Science Research Foundation fellowship as the Shurl and Kay Curci Foundation fellow, and by Burroughs Wellcome Travel Fund. This work was supported by 2020R1A2C4002533 (D.S.) and NRF-2017R1A2B3004198 (H.K.), HI17C0676 from Korean Ministry of Health and Welfare (H.K.), 5R01AG008200 from National Institute on Aging (NIA) and the National Institute of Health (NIH) (N.K.R.), IBS-R026-D1 from IBS (H.K. and J.C.), NRF-2019R1A2C1085712 (Y.H.K.), the UCSF Center for Cellular Construction (an NSF Science and Technology Center, no. DBI-1548297) (Z.J.G.), U01CA244109 from the National Cancer Institute (Z.J.G), 1R01GM112081, 1R01GM126542-01, and R35GM134948 from the National Institute of General Medical Science (NIGMS) and the NIH (Y.J.), and the UCSF Program for Breakthrough Biomedical Research (PBBR) funded in part by the Sandler Foundation (Y.J.). Z.J.G. is a Chan Zuckerberg BioHub Investigator.

Author contributions

M.K., K.M.S., Z.J.G., and Y.J. conceived the ideas and designed research; M.K. and K.M.S. constructed plasmids, generated cell lines, and performed confocal microscopy. M.K. performed mechanogenetics, truncation study, spatial mutation, immunoblot analysis, reporter cell assay, and APP experiment. K.M.S. performed Notch exclusion and activation experiments, designed truncation study, wrote custom python image analysis scripts. N.K. performed coarse-grained MD simulation. R.G. generated cadherin-KO cells. W.R.K. performed flotillin staining and mechanogenetic experiment. M.A. and H.L. synthesized magnetic nanoparticles and helped confocal imaging. J.F. and D.S. performed initial proof-of-concept experiment. A.G. and N.K.R. provided anti-PS1 antibodies. H.K., Y.H.K., and J.C. oversaw and supervised CRISPR-Cas9 KO experiment, MD simulation, and magnetic nanoparticle synthesis, respectively. Z.J.G. oversaw and supervised all spatial mapping and mutation experiments. Y.J. oversaw and supervised all aspects of the experiment. M.K. and K.M.S. analyzed data. M.K., K.M.S., Z.J.G., and Y.J. wrote the manuscript.

Competing interests

The authors declare no competing interest.

Additional information

Supplementary Information is available for this paper.

Figure 1. Spatial segregation of Notch receptors and ligands from cadAJs prevents their
interactions with γ-secretase. a, A schematic showing spatial distribution of Notch signaling
components at the juxtaposed cell-cell interface. **b,** Representative confocal fluorescence images
showing presenilin-1 (PS1), Notch1, and Dll1 distributions relative to E-cadherin-based cadherin junction
(cadAJ). (top) A maximum projection image. Scale bar, 5 μm. (bottom) z-resliced images. Scale bar, 3
μm. PS1 was strongly enriched at the cadAJ, whereas Notch1 and Dll1 were segregated from the cadAJ.
c, Line profiles quantifying fluorescence from E-cadherin (green), PS1 (cyan), Notch (red), and Dll1
(magenta) along a representative section of the cadAJ (a white dashed line in panel **b**, z-resliced images).
d, A schematic showing mechanogenetic interrogation of γ-secretase and Notch distribution relative to
the artificial cadAJs. Artificial cadAJs were formed by clustering Ecad-GFP labeled with
magnetofluorescent nanoparticles (MFNs) by application of an external micromagnetic tweezer (μMT).
DAPT was used to inhibit γ-secretase activity during mechanogenetic interrogation of Notch/γ-secretase
interaction in absence of the membrane juxtaposition. **e,** Epifluorescence images showing the formation
of an artificial cadAJ by mechanogenetics. After stimulation by μMT, vivid MFN and E-cadherin signals
at the magnetic focus were seen, indicating formation of cadAJs. Scale bar, 5 μm. **f,** Confocal
fluorescence images of E-cadherin, PS1, and Flotillin-1 (Flot1) at the artificial cadAJ. Scale bar, 2 μm.
Line profiles of E-cadherin, PS1 and Flot1 signals along a white dashed line. $\Delta I/I_0$ represents a fold
change relative to nonjunctional membrane signals. **g,** Confocal fluorescence images showing E-
cadherin and Notch distributions at the artificial cadAJs after μMT application. Strong accumulation of
Notch signals at the artificial cadAJ was clearly seen. Line profiles of MFN, E-cadherin, and Notch signals
along a white dashed line. $\Delta I/I_0$ represents a fold change relative to fluorescence intensity before
stimulation. Scale bar, 2 μm.
561
562

Figure 2. Size-dependent segregation of Notch at the cadAJs controls the proteolytic sequence.
a, A schematic to capture the spatial distribution of Notch intermediates during the cell-surface activation pathway. **b,** Confocal z-resliced images showing Notch distribution (red) relative to cadAJ (green) from the cells without Dll4 activation (i), treated with Dll4 and TAPI2 (ii), treated with Dll4 and DAPT (iii), and washed out to remove DAPT inhibition (iv). Scale bar, 3 μm. **c,** Quantification of Notch signal enrichment at the cadAJs during the activation. Notch enrichment (I_{IN}/I_{OUT}) is calculated as the ratio of average Notch fluorescence intensity within cadAJs (I_{IN}) and outside cadAJ (I_{OUT}). The enrichment factor of Dil is present as a control showing cadAJ-independent distribution. ****$P < 0.0001$, ns: non-significant, one-way ordinary ANOVA followed by Tukey's multiple comparison testing. **d,** Schematics of Notch variants with different truncation lengths, in comparison with the cadAJ intermembrane cleft. **e,** Confocal fluorescence images showing spatial distribution of the Notch variants (red) relative to the cadAJs (green). To prevent any ligand-independent activation, cells were incubated with TAPI-2 and DAPT. (left) Maximum

projections of confocal z-stacks are shown. Scale bar, 5 µm. (right) Confocal z resliced images along the white dashed lines in the maximum projection images. Scale bar, 2 µm. **f,** Quantification of the enrichment factor (I_{IN}/I_{OUT}) of Notch variants relative to the cadAJs. A box-plot showing binary localization of NΔEGF which are defined as either excluded (yellow) or enriched (orange) is shown on the right. **P < 0.01, ***P < 0.001, ns: non-significant, one-way ordinary ANOVA followed by Tukey's multiple comparison testing. **g,** Confocal *z*-resliced images showing the distribution of extracellular SNAP (purple) and intracellular mCherry (red) tags of NΔEGF relative to Ecad-AJs (green) after DAPT removal. Scale bar, 3 µm. **h,** A box-whisker plot showing Pearson correlation coefficients of extracellular-SNAP (purple) and intracellular-mCherry (red) domains relative to the cadAJs before and after DAPT washout (***P < 0.001; ns, not significant; n = 9 biological replicates; two-tailed unpaired Student's t-test). **i,** Time series of confocal z-resliced images showing the enrichment of NΔEGF (red) at the cadAJ (green) under DAPT treatment (t = 0), and the dissipation during DAPT washout (t ≥ 2). Scale bar, 3 µm. **j,** Single-cell traces showing the time-course of the decline of NΔEGF enrichment factor at the cadAJs during DAPT washout (mean ± s.e.m.; n = 4 independent single-cell experiments; see Extended Data Figure 6, d-f). **k, l,** Western blot analyses of cleaved NICD levels in the cells stably expressing N^{FL}, $NΔEGF_{1-25}$, NΔEGF, and NEXT. All cells were transfected with Ecad-GFP and incubated with TAPI2 for 24 hr. β-actin levels represent the loading control. A representative image of immunoblotting **(k)**, and quantification **(l)** of cleaved NICD levels. The average intensity of each NICD band relative to respective β-actin band was quantified and then normalized to that of NEXT (mean ± s.d.; n = 4 biological replicates; *P < 0.05, ****P < 0.0001, one-way ANOVA followed by Tukey's multiple comparison test). (**c, f, h**) Boxes and whiskers indicate the interquartile and the full ranges, respectively. Colored lines and (+) marks indicate median and mean, respectively.

Figure 3. Spatial mutation directs Notch activation, regardless of ligand presentation or S2 cleavage. a, A schematic describing DNA-mediated crosslinking strategy to enhance NΔEGF localization at the cadAJ. **b,** Confocal z-resliced images showing intense NΔEGF fluorescence (red) enriched at the cadAJ (green) after the DNA crosslinking. Scale bar, 3 μm. Quantification of enrichment (I_{IN}/I_{OUT}) without (n = 33) with DNA crosslinker treatment (n = 29), indicating an increase in enrichment after the crosslinking (*P < 0.05; two-tailed Mann-Whitney-Wilcoxon test). **c,** Western blot analyses showing increased S3-cleaved NICD levels in the NΔEGF cells treated with the DNA crosslinker. (top) A representative gel image showing immunoblotting for NICD and β-actin. (bottom) Quantification of cleaved NICD levels. The average intensity of NICD band was normalized to that of β-actin band in each sample. (mean ± s.d.; ** P < 0.01; n = 5 biological replicates; ordinary one-way ANOVA). **d,** Spatial mutation of NEXT via chemical ligation of macromolecular pendants (denoted as P). 10 μM of BG-modified polymers or proteins were conjugated to the extracellular SNAP tag of NEXT. Cartoons depicting shape and hydrodynamic size of different pendants are shown. Confocal images showing size-dependent spatial mutation of NEXT (red) at the cadAJs (green). The top row shows maximum projection images of the cells treated with the indicated pendants. Scale bar, 5 μm. The middle row shows confocal z resliced images along the white dashed lines in the maximum projection images. Yellow and green arrowheads indicate the cadAJs enriches with and those that excludes Notch, respectively. Scale bar, 3

μm. The bottom row shows line profiles quantifying fluorescence signals from NEXT (red) and E-cadherin (green) along the white lines in the z-resliced images. Images and line profiles are representative of n ≥ 15 biological replicates. **e,** Quantification in I_{IN}/I_{OUT} of NEXT with macromolecular pendants (n ≥ 15). **f,** Confocal fluorescence images of nuclear mCherry signals resulting from nuclear translocation of cleaved NICD (red) in NEXT cells treated with macromolecular pendants. DAPI signals (blue) indicate cell nucleus. Scale bar, 5 μm. **g,** Quantification of nuclear mCherry fluorescence for NEXT cells treated with the pendants. Cells expressing N^{FL} were used as a negative control. Data shown are from single cells (n ≥ 180) identified in large-area confocal images. **h,** A plot representing the NICD level of various Notch variants as a function of the enrichment factor (I_{IN}/I_{OUT}, mean ± s.e.m.; n ≥ 15 biological replicates). All Notch variants with different truncation length, DNA crosslinking, or pendant addition used in Figures 2 and 3 were included. The Notch variants showing exclusion, mixed distribution, and enrichment were indicated with red, yellow, and green background colors, respectively (median ± s.e.m.; n ≥ 4 for western blot for NICD levels; n ≥ 180 for nuclear mCherry fluorescence). (**e, g**) Boxes and whiskers indicate the interquartile and the full ranges, respectively. Black lines and (+) marks indicate median and mean, respectively. **P < 0.01; ****P < 0.001; ns, not significant; ordinary one-way ANOVA followed by Tukey's multiple comparison.

Figure 4. The cadAJ-mediated spatial switch regulates Notch and APP signaling. a, Representative
fluorescence images showing Notch activation in U2OS SNAP-N^{FL}-Gal4 reporter cell lines in different
cellular environments: Group of cells (left), solitary cells with no prior contact (middle), and solitary cells
plated on an Ecad-Fc coated substrate (right). All cells were plated on Dll4-Fc coated substrates. Scale
bars, 20 µm (see supplementary video 1, 2). **b,** Quantification of Notch activation by measuring H2B-
mCherry fluorescence changes (intensity at t = 16 hr minus intensity at t = 0 hr) in cells within a group (n
= 152), solitary cells (n = 50). **c,** Quantification of Notch activation in solitary cells cultured on an Ecad-
Fc-coated substrate (n = 27) and those cultured without ligands (n = 27). **d,** Representative confocal
images of H2B-mCherry fluorescence in U2OS SNAP-N^{FL}-Gal4 reporter cells (wt), E-cadherin knockout
cells (Ecad-KO), Ecad-KO cells with recombinant E-cadherin transfection (Ecad-KO + Ecad), and Ecad-
KO cells with N-cadherin transfection (Ecad-KO + Ncad). Cytosol labeled with CMFDA dye was shown
for wt and Ecad-KO cells. E-cadherin and N-cadherin were shown for Ecad-KO + Ecad and Ecad-KO +
Ncad cells. Scale bar, 100 µm. **e,** Quantification of Notch activation in the wt, Ecad-KO, Ecad-KO + Ecad,
and Ecad-KO + Ncad cells. Data shown are from n ≥ 50 single cells analyzed from two independent
experiments. **f,** (left) Confocal fluorescence maximum projection (left) and z-resliced images of U2OS
cells co-expressing Ncad-mCherry (green) and APP-EGFP (red). To capture the APP intermediates, cells
were cultured with a combination of α-, β-, and γ-secretase inhibitors. Scale bars, 10 µm (maximum
projection) & 3 µm (inset). **g,** Aβ$_{42}$/Aβ$_{40}$ ratio measured by ELISA in wild-type cells or CDH1/2 KO cells
(mean ± SEM, n = 3, *P< 0.05, two-tailed paired Student's t test). (**b, c**, and **e**) Boxes and whiskers
indicate the interquartile and full ranges, respectively. Black lines and (+) marks indicate median and
mean, respectively. **P < 0.01; ****P < 0.001; ns, not significant; unpaired two-tailed t test in (**b**) and (**c**);
ordinary one-way ANOVA followed by Tukey's multiple comparison in (**e**).

658
659
Extended Data Figure 1. Distribution of ADAM10 and γ-secretase at the cell-cell interface. a,
Representative confocal fluorescence images showing ADAM10 distribution relative to E-cadherin-based
adherens junction (cadAJ). U2OS cells were transfected with Ecad-GFP (green), and stained with
ADAM10 antibody (red). Z-resliced images were reconstructed along the white dashed lines. Scale bars,
5 µm and 3 µm for maximum intensity projection and z-resliced images, respectively. **b,** Additional
immunofluorescence confocal images showing endogenous presenilin-1 (PS1, magenta) distribution
relative to cadAJs (green). Paxillin was also imaged as a negative control (blue) Scale bars, 10 µm, 3
µm, and 3 µm for maximum intensity projection, zoomed-in, z-resliced images, respectively. **c,** Cross-
correlation analysis of PS1, ADAM10, and Notch distributions over E-cadherin. PS1 fluorescence
intensities exhibited strong positive correlation with cadAJs, while Notch showed nearly zero correlation.
ADAM10 showed a weakly positive correlation over cadAJs. The cadAJs were set as x = 0 in the analyses.
The shading along the curves represent error bars (s.e.m.; n ≥ 9).

Extended Data Figure 2. Notch exclusion from cadAJs is consistent for several cadherin family proteins, host cell types, and cell polarization states. A series of cell lines including U2OS, MDCK, and HUVEC were imaged. **a, b,** U2OS (**a**) and MDCK (**b**) cells were co-transfected with plasmids encoding Ecad-GFP (green) as well as SNAP-N^{FL}-mCherry. Notch receptors were labeled with BG-Alexafluor647 (magenta). Line profiles show fluorescence signal intensities along the white lines. Scale bars 10 µm, 5 µm, and 2 µm for maximum intensity projection, zoomed-in, and z-resliced images, respectively. **c,** HUVECs were transfected with SNAP-N^{F}-mCherry and labeled with SNAP surface 647. VE-cadherin was immunostained with an anti-vascular endothelial cadherin (VE-cad) antibody. Line profiles show fluorescence signal intensities along the white lines. Scale bars 5 µm and 2 µm for maximum intensity projection and z-resliced images, respectively. **d,** Polarized MDCK cells grown on a transwell filter. Scale marked every 5 µm for 3D construction and Scale bar is 2 µm for z-resliced images. **e**, A schematic representation of Notch localization in a polarized epithelial cell.

686
687
Extended Data Figure 3. Quantitative analysis of PS1, Notch, Dll distribution relative to cadAJs
and additional confocal images representing their spatial distribution. a, Confocal images of U2OS
cells co-expressing Ecad-GFP (green) and SNAP-N^{FL}-mCherry (red). Cells were also immunostained
with anti-PS1 antibodies (cyan). Line profiles show fluorescence signal intensities along the white lines.
Scale bars, 10 µm, 2 µm, and 2 µm for maximum intensity projection, zoomed-in, and z-resliced images,
respectively. **b**, Paired analysis of Manders' overlap coefficients of E-cadherin and Notch signals over
PS1 in multiple cells (n = 9). ***P < 0.001. **c,** Box-whisker plots showing Manders' overlap coefficients
(MOCs) of PS1 (cyan), Notch (red), and Dll1 (purple) relative to cadAJs. Each dot represents the MOC
of a selected cadAJ. Boxes and whiskers denote the inner-quartile and full ranges. Colored lines and (+)
marks indicate median and mean, respectively (****P < 0.0001; ns, not significant; n ≥ 11; ordinary one-
way ANOVA followed by Tukey's multiple comparison test). **d**, Confocal fluorescence images of U2OS
cells co-expressing Ecad-GFP (green) and Halo-Dll1. Halo-Dll1 was labeled with Halo-660 dyes
(magenta). Scale bars, 5 µm, 1 µm, and 1 µm for maximum intensity projection, zoomed-in, and z-resliced
images, respectively.

Extended Data Figure 4. Interrogation of the mechanism underlying γ-secretase recruitment into cadAJs through mechanogenetics, cell-cell cadAJ spatial mapping, molecular dynamics simulation, and cholesterol depletion experiments. a, b, Additional mechanogenetic experiments showing PS1 and Flot1 localizations at cadAJs. E-cadherin and Flot1 were labeled with fluorescent tags. Endogenous PS1 was immunostained after fixation. Scale bar, 20 μm. **c, d,** Representative confocal fluorescence images showing the PS1 and Flot1 distribution relative to native cell-cell cadAJs. (left) A

maximum projection image of merged channels. Scale bar, 10 μm. (center) Magnified images showing greater details of the boxed region. Scale bar, 2 μm. (right) Z-resliced images showing the sections of the cadAJs. Scale bar, 2 μm. Line profiles of fluorescence signals from E-cadherin, PS1, and Flot1 along the white dashed lines in the z-resliced images. **e,** Cross-correlation analysis of E-cadherin and PS1 over Flot1. Both Flot1 and PS1 fluorescence intensities exhibited strong positive correlation with the cadAJ. The shades along the curves represent error bars (s.e.m.; n = 7). **f,** Manders' overlap coefficients of E-cadherin and Flot-1 signals over PS1. Two-tailed paired Student's t test, ns: not significant. **g,** Quantitative assessment of Flot1 and Notch colocalization with E-cadherin. Pearson correlation coefficient is shown. Each circle is the score of a single cadAJ. Overlaid are box and whisker plots. **** P<0.0001. Two-tailed unpaired Student's t test. **h,** Snapshot images of coarse-grained MD simulation of a lipid bilayer comprising of 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DIPC; gray), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC; yellow), N-palmitoyl-O-phosphocholineserine (PPCS; green), 1,2-dipalmitoyl-sn-glycero-3-phosphoserine (DPPS; blue), immobilized DPPS (pink), and cholesterol (CHOL; grey). Left and right panels represent the simulation results without and with a partial (30%) DPPS immobilization, respectively. **i, j,** Diffusion coefficients (**i**) and binding energy (**j**) of individual lipid components in a lipid bilayer with or without a partial (30%) DPPS immobilization. Both decreases in lipid diffusion and increased binding energy indicate lipid microdomain stabilization due to DPPS immobilization. **k,** Confocal fluorescence images showing PS1 distribution at an artificial cadAJ after cholesterol depletion with MβCD treatment. No PS1 recruitment was seen at the cadAJ, suggesting that γ-secretase recruitment to the cadAJ requires lipid microdomain formation at the cadAJ.

732
733
Extended Data Figure 5. Additional confocal images representing recruitment of Notch receptors
to the artificial cadAJ. a, Pre-stimulation images of a subcellular locus of the cell shown in Fig. 1g. Initial
dim MFN (magenta), E-cadherin (green), and Notch (red) signals were evenly distributed. Scale bar, 2
μm. **b, c,** Additional artificial cadAJs showing Notch recruitment. Time-lapse images (c) were acquired
before micromagnetic tweezer (μMT) stimulation and then at 10, 20, 30 minutes of the μMT application.
Gradual MFN and E-cadherin clustering were clearly seen, followed by Notch accumulation to the cadAJ.
Scale bar, 2 μm. **d.** Kinetics of signal enrichments at the artificial cadAJ shown in the panel c.
741

Extended Data Figure 6. Confocal images showing the gradual release of NICD from S2-cleaved Notch concentrated within cadAJs. a, Representative confocal images and enrichment factors (I_{IN}/I_{OUT}) of DiI membrane staining dye distribution relative to cadAJs. **b,** Representative time-course z-resliced images showing S2-cleaved Notch at cadAJs as a function of time after DAPT removal. The NICD signal (red) at the cadAJ gradually decreases, indicating NICD release. Images shown here are not from identical cells, but represent a general trend of NICD signal at cadAJs for each time point. **c,** Quantification I_{IN}/I_{OUT} ratio as a function of time after DAPT washout. **d-f,** Time traces of NICD release from U2OS cells expressing NΔEGF (red) and Ecad-GFP (green). **d,** Cells showing strong NΔEGF enrichment at cadAJs under DAPT treatment were chosen. DAPT was removed while keeping TAPI2 concentration constant and time series confocal images were acquired at 0, 2, 4, and 6 hrs from DAPT removal. The white arrows indicate the locations of two representative cadAJs shown and analyzed in Fig. 2i, j. **e,** Single-cell traces (n = 4) of enrichment factor. **f,** Quantification of changes in NICD signal from these four cells at the cadAJs and non-cadAJ membrane, at t = 0 (green, before washout) and t = 6 hr (red, after DAPT wash-out). CadAJs and non-cadAJ membrane were detected based on thresholding and automatic segmentation using the custom-built script. Average mCherry fluorescence signals from cadAJs and non-cadAJ membranes were also measured using the same script. Intracellular-mCherry signals significantly decreased at the cadAJs, but not at non-cadAJ membranes (* P < 0.05, ns: non-significant, paired two-tailed Student's t-test, n = 4). Scale bars, 5 μm.

762
763
Extended Data Figure 7. Spatial mutation of NΔEGF using DNA-mediated crosslinking. a. A
representative western blot of lysate from cells after 2 hr incubation with or without DNA crosslinkers.
The blot was labelled with anti-SNAP and anti-β-actin antibodies. The expected mass of NΔEGF and its
complex with the Notch construct and E-cadherin forming a heterodimer is 90 kD and 230 kD, respectively.
β-actin detection was used to assess protein loading. Predicted bands representing Notch-E-cadherin
heterodimers (heterodimer) and SNAP-NΔEGF-mCherry monomers are indicated with solid lines.
Molecular weights are indicated on the left. **b.** Representative maximum intensity projection confocal
images showing the distribution of NΔEGF relative to the cadAJs after crosslinking. Cells were treated
with or without DAPT. Scale bar, 10 μm. **c, d,** Removal of DAPT induced S3 cleavage of NΔEGF at the
cadAJs. **c,** Single-cell confocal images showing intracellular-mCherry signals at the cadAJ under DNA
and DAPT treatment (left) and after washing out DAPT (right). Removing DAPT elicited a significant
reduction in mCherry signal intensity from the cadAJ. Scale bar, 5 μm. **d.** Paired analysis of multiple cells
expressing NΔEGF (n = 6) in enrichment factor (I_{IN}/I_{OUT}) after DAPT washout. Each dot represents I_{IN}/I_{OUT}
value before and after DAPT washout from a single cell. Each line corresponds to the I_{IN}/I_{OUT} changes
before and after DAPT washout in a same single cell (**$P<0.01$, paired two-tailed Student's t test, n = 6).
779

Extended Data Figure 8. Spatial mutation of NEXT using chemical ligation of macromolecular
pendants. a, Dynamic light scattering spectra of BG-modified macromolecules used in the experiment
to induce spatial mutation of NEXT in Fig. 3d-g. **b,** Larger area (1 x 1 mm) confocal fluorescence images
shown in **Fig. 3f**. Scale bar, 200 μm.

Extended Data Figure 9. Generation of U2OS SNAP-N^{FL}-Gal4 fluorescence reporter cell lines
lacking E- and N-cadherins via CRISPR/Cas9. a. Schematic representation of human CDH1 gene
structure and targeted segmental deletion sites. The sixteen exons are shown in orange boxes (E1-E16).
Red arrowheads indicate the sgRNA-binding sites (E1LT1, E2RT1, E13LT1, E13LT2, E14RT1 and
E14RT2). The targeted segmental deletion of 940bp and 4.6kb is shown with a black line with red arrow
tips, respectively. Purple arrows represent PCR primers used for the T7E1 assay and detection of alleles
with targeted deletions, respectively. **b, c,** T7E1 assays to evaluate the efficiency of each sgRNAs. U2OS
SNAP-N^{FL}-Gal4 reporter cells were analyzed after transfection with plasmids encoding Cas9 and each
sgRNA for CDH1 (**b**) and CDH2 (**c**). Mutation frequencies (indel [%]) were calculated from the band
intensities. Untransfected cells were used as controls (CONT). The red arrows and asterisk indicate the
uncleaved and expected cleaved position of DNA bands generated by cleavage by T7E1, respectively.
M: Marker lane. **d,** Identification of clones containing the entire segmental deletion in both target alleles
for selected clones. The wild-type and deleted alleles were detected by PCR using different primer pairs
at exon12 and 13 target regions, respectively. The black arrows indicate the expected positions of the
amplicons with wild type allele or allele with deletions. The red arrows indicate the candidate clones

containing the entire segmental deletion in both target alleles. **e,** Sanger sequencing of the CDH1 wild-
type (wt) allele and alleles containing deletions of the segmental CDH1 gene. Cas9 recognition sites are
shown in red and the protospacer adjacent motif (PAM) sequences are shown in blue bold characters.
Dashes indicate deleted bases. The number of occurrences is shown in the first set of parentheses (e.g.,
X6, X3 and X6 indicate how many times each sequence was observed). The cleavage sites are indicated
by red arrows. The sequence length of the consequent large deletion is shown in the second set of
parentheses. **f,** CDH1 mRNA expression levels in CDH1 KO clone #3 and #4 were determined using
qRT-PCR. CDH1 expression levels in the selected clones containing a segmental deletion were
quantified relative to CDH1 mRNA levels of the wild-type U2OS SNAP-N[FL]-Gal4 cells. Clone #3 was used
for subsequent experiments. **g,** Western blot analyses of cleaved NICD levels in the U2OS SNAP-Notch-
Gal4 reporter cells (wt), Ecad-KO, and Ecad-KO cells with recombinant E-cadherin transfection (Ecad-
KO + Ecad). (top) A representative image of immunoblotting. (bottom) Quantification of cleaved NICD
levels. The average intensity of NICD bands relative to β-actin bands was quantified and then normalized
to that of Ecad-KO cells (mean ± s.d.; ***P < 0.001; n = 5 biological replicates; one-way ANOVA followed
by Tukey's multiple comparison test). **h,** PCR-based evaluation of targeted segmental deletions in double
KO (CDH1 and CDH2) single-cell derived clones. U2OS SNAP-N[FL]-Gal4 cells were transfected with
plasmids encoding Cas9 and two sets of sgRNA pairs, each targeting CDH1 and CDH2, respectively.
The red arrow indicates the expected position of the amplicon from the region containing the segmental
targeted deletion M: Marker lane. **i,** To obtain the bi-allelic KO clones, we validated bi-allelic deletion in
both CDH1 and CDH2 genes from the indicated clones. The red arrows indicate the expected positions
of the amplicons with wild type allele or allele with deletions. Clone #11 (red color), showing bi-allelic
deletion of both CDH1 and CHD2, was used for further subsequent experiments. M: Marker lane**.**
824

Extended Data Table 1. Cell lines used in this study.

Cell line	Stable expression*,**	CRISPR-KO	Transient expression 1*,**	Transient expression 2*,**	Figure
Plain U2OS			Halo-Dll1	Ecad-GFP	1(b-d) EDF[#] 3a
			Ecad-GFP	Flot1-Halo	1(e-g) EDF 3(d-f)
			SNAP-Ecad-GFP	Flot1-Halo	1j EDF 5(a-d)
			Ecad-GFP		EDF 1(a-c)
			APP-GFP	SNAP-Ncad	4f
T-rex U2OS (doxycycline inducible)	SNAP-N^{FL}-mCherry (N^{FL})		Ecad-GFP		1(b-d) 2(b, c) 2(k, l) 3h EDF 2a EDF 3(b,c) EDF 7(a,b)
			Halo-Ecad-GFP		1(k, l) EDF 6(a,b) EDF 8(a-d)
	SNAP-$N\Delta EGF_{1-25}$-mCherry ($N\Delta EGF_{1-25}$)		Ecad-GFP		2(e, f) 2(k, l) 3h
	SNAP-$N\Delta EGF$-mCherry ($N\Delta EGF$)		Ecad-GFP		2(e-l) 3h EDF 7(c-e)
			Halo-Ecad-GFP		3(a-c) 3h
	SNAP-NEXT-mCherry (NEXT)		Ecad-GFP		2(e, f) 2(k, l) 3(d-h) EDF 9b
	SNAP-N^{FL}-Gal4 & UAS-H2B-mCherry				4(a-e)
		Ecad	Ecad-GFP		4(d, e)
			Ncad-GFP		4(d, e)
	SNAP-N^{FL}-Gal4 & UAS-H2B-mCherry (uninduced)		APP-GFP	SNAP-Ncad	4g
		Ecad & Ncad	APP-GFP		4g
MDCK			SNAP-N^{FL}-mC	Ecad-GFP	EDF 2(b,d)
HUVEC			SNAP-N^{FL}-mC		EDF 2c

*X-Y represents a fusion protein of X with Y.

**X-Y-Z represents a Y protein fused with X- and Z- tag proteins at its N- and C-termini, respectively.

[#]Extended Data Figure

Abbreviations:

    <u>Notch receptors and ligands</u>

    N^{FL}: full length Notch

    $N\Delta EGF_{1-25}$: Notch with EGF domain truncation from 1-25.

    $N\Delta EGF$: Notch with complete EGF repeat domain truncation.

    NEXT: Notch with extracellular domain truncation

    Dll1: Delta like ligand 1

839   <u>Cadherins</u>
840   Ecad: epithelial (E) cadherin
841   Ncad: neural (N) cadhrin
842   VEcad: vascular endothelial cadherin
843
844   <u>Amyloid precursor proteins</u>
845   APP: amyloid precursor proteins with deletion of YENPTY motifs
846
847   <u>Self-labeling and fluorescent protein tags</u>
848   SNAP: SNAP tag
849   Halo: Halo tag
850   GFP: green fluorescent protein
851
852   <u>Others</u>
853   Flot1: flotillin 1
854   H2B: histone 2B
855   UAS: upstream activation sequence

METHODS

Cloning, cell culture, transfection, and cell line generation

Cloning

Plasmid constructs used in this study are listed in **Extended Data Table 1**. All constructs used in this
paper were assembled using standard restriction enzyme-based cloning, in fusion cloning, and/or Gibson
isothermal assembly. The maps, sequences, and construction details of all plasmids are available upon
request. All constructs were sequenced to confirm mutation. Complete details of all cloning procedures
are available upon request.

Flag-human Notch1 (N^{FL})-Gal4 and pGF1-UAS-H2B-mCherry were gifts from S. Blacklow (Harvard
University). Flag-human N^{FL}-Gal4 was provided in a Tet-ON Flp-IN vector (pcDNA5). SNAP-N^{FL}-mCherry
and SNAP-N^{FL}-Gal4 were constructed as previously reported[1]. All Notch1 variants with partial or full
extracellular domain truncation were constructed by linearizing and amplifying SNAP-hN1-mCherry
vector via inverse PCR while omitting the sequence corresponding the ECD truncation. Notch
ectodomain sequences of amino acid 23-981, 23-1426, 23-1709 were deleted for SNAP-ΔEGF_{1-25}-
mCherry. SNAP-ΔEGF-mCherry. and SNAP-NEXT-mCherry. respectively.

Ecad-GFP was purchased from Addgene (Addgene plasmid # 28009;
http://n2t.net/addgene:28009). SNAP-Ecad-GFP and Halo-Ecad-GFP were constructed first by
linearizing and amplifying Ecad-EGFP vector via inverse PCR. SNAP- and Halo-tags were then inserted
in frame with E-cadherin, downstream of the E-cadherin pro-peptide (amino acid #155) and upstream of
EGFP, using Gibson assembly (NEB). pCMV6-Flotillin-1-Halo (Flot1-Halo) was constructed by replacing
the myc-tag within the pCMV6-Flotillin1-myc (purchased from Origene (MR206823)) with the Halo-Tag
sequence in frame with Flot1 using In-Fusion cloning (Clontech). Human amyloid precursor protein1-
EGFP (APP-EGFP) was created by cloning human APP695 (a gift from C. Miller of King's College London)
into a pEGFP-N1 plasmid (Clontech). To facilitate membrane distribution mapping, all APP constructs
used in confocal imaging lack YENPTY motifs. APP$^{\Delta YENPTY}$-EGFP was constructed by first linearizing the
vector via inverse PCR, and deleting the sequence of final 15 amino acids upstream of C-terminus of
APP (amino acid 681-695), which includes YENPTY motif (amino acid 682-687) using Gibson assembly.

Tissue culture

Human U2OS cells were cultured in McCoy's 5A media supplemented with 10% heat-inactivated FBS
(Life Technologies) and 1% Pen-Strep (Life Technologies) and passaged with 0.05% trypsin-EDTA.
MDCK cells were grown in Eagle's Minimum Essential Medium (UCSF cell culture facility) supplemented
with 10% heat-inactivated FBS and 1% Pen-Strep. MDCKs Cells were lifted by treating for 10 minutes
with PBS Ca^{2+} and Mg^{2+} free with 0.05% EDTA followed by trypsinization. HUVECs were purchased from
ATCC (ATCC® CRL-1730™) and grown in Endothelial Cell Growth Medium-2 (EGM-2, Lonza CC-3162)
for up to 4 passages (0.05% trypsin-EDTA) after thawing. All cell lines were maintained at 37°C in a
humidified incubator with 5% CO_2 and passaged every 2-3 days, depending on confluency, using 0.05%
Trypsin-EDTA (UCSF cell culture facility). For polarized MDCK culture, **c**ells were for use on a transwell
filter (0.4 µm, collagen coated) at high density.

Transfection and cell line generation

All cell lines expressing recombinant proteins used in this study are listed in **Extended Data Table 1**.
U2OS stable cell lines were constructed from parental U2OS T-rex cell lines (Flp-IN, Tet-ON engineered
cell line, gift from S. Blacklow). Constructs were inserted into the engineered Flp-IN site by co-transfection
with a plasmid containing the Flp-recombinase (pOG44) via electroporation with the Neon Transfection
System (ThermoFisher) according to manufacturer's protocol (Shock Conditions: 1230V, 10ms, 4 pulses,
number of cells 5 x 10^6). The amount of total DNA used was 10 µg/well: 1 µg of DNA containing the
desired construct and 9 µg pOG44. Cells transfected with desired plasmids were incubated in a selection
medium containing 400 µg/mL hygromycin (Invitrogen) for at least 10 days. All cells with Notch truncation
and reporter were further sorted for inducible expression of Notch variants via fluorescence-activated cell
sorting (FACS) on a FacsAria2 (BD) by staining for the appropriate tag (SNAP or Halo) with fluorescently
tagged antibody. For single-cell monoclonal population establishment, fluorescently-positive bulk-sorted
populations were plated into 96 well plates at 0.2 cells/well by serial dilution and grown in selection

medium. Each clonal cell population was tested and selected based on the levels of Notch reporter activity or Notch membrane expression. U2OS cells expressing recombinant proteins transiently were generated by transfecting plasmids encoding desired proteins using Neon-based electroporation. Cells were allowed to settle in a 6-well cell culture dish post electroporation for 6-8 hr. To remove dead cells, cells were lifted and re-plated on a fibronectin-coated glass bottom dish with 1×10^5 cells per well density. MDCK cells were plated at 70% density then transfected with N^{FL}-mCherry utilizing Lipofectamine 3000 (ThermoFisher) or Neon electroporation according to the manufacturer's protocol (Shock Conditions: 1,650V, 20ms, 1 pulse, number of cells 5×10^6). HUVEC cells were transfected via electroporation with SNAP-N^{FL} via the BioRad Gene Pulser system (250 V, 20 ms square wave, 1×10^6 cells/mL Gene Pulser Electroporation Buffer, 5 µg/mL SNAP-N^{FL}-mC). All cells transiently expressing recombinant proteins were incubated for 24-48 hr from the transfection, and then used for further analyses.

Live cell mechanogenetics experiment

Mechanogenetics experiments were performed as previously described in Kwak et al., 2019 with some modifications[2,3]. Monovalent magnetofluorescent nanoparticles (MFNs) were synthesized as previously described[2].

Micro-magnetic tweezers (µMT) set up. The µMT was set up and aligned on the inverted microscope with point-scanning confocal and total internal reflection fluorescence (TIRF) imaging capabilities (Nikon) as previously described[2,3]. The needle probe – NdFeB magnet assembly was attached to the z-translation stage (Sutter Instrument, MP-325) and its location was carefully aligned with the microscopic objective lens while observing the dummy substrate filled with DPBS. The µMT tip was positioned at the center of the objective oculus with bright-field illumination using the X-Y translation stage linked to PIMikroMove (Physik Instrumente) and µManager (UCSF). Using the z translation stage, the µMT was carefully lowered to set the height of the tip to 10 µm above the focal plane while recording the X-Y coordinates and the z-position of the needle probe.

Preparation of cells expressing recombinant Flotilin-1 for mechanogenetics experiments. U2OS cells were co-transfected with SNAP-Ecad-GFP (5 µg) and Flot1-Halo (5 µg) plasmids using Neon electroporation. 24 hr later, cells were re-plated on a #1.5 glass-bottomed dish (MatTek, d = 10 mm) coated with collagen at a density of 1×10^5 cells per dish. To fluorescently label Flot1-Halo, cells were treated in a complete McCoy medium containing 3.5 µM cell membrane permeable Halo-ligand 660 dye (Promega) for 30 minutes at 37°C. Cells were washed three times with DPBS, incubated with a phenol red-free complete medium, and then mechanogenetically stimulated (see below). For the cholesterol depletion experiment, we also treated cells 10 mM of methyl-β-cyclodextrine (MβCD) (Sigma-Aldrich) in serum-free McCoy's 5A medium for 30 min at 37°C and washed with a complete medium three times. To label SNAP-Ecad-GFP with MFNs, cells were first treated with 5 µM of an oligonucleotide bearing benzylguanine (BG-$T_{60}ACTG_{10}$) for 45 minutes at 37°C, washed two times with 10 ml of serum free medium, and then incubated with serum-free medium containing 10 nM monovalent MFNs bearing complementary sequence ($T_{60}CAGT_{10}$) and 0.5% alkali casein for 10 min at 37°C, 5% CO_2. Cells were washed with 10 ml of complete medium two times, and then incubated with phenol red-free medium for mechanogenetic experiments on a confocal or wide-field epifluorescence microscope.

Preparation of cells expressing human Notch1 receptor for the mechanogenetic experiment. Inducible U2OS cells stably integrated with SNAP-N^{FL}-mCherry were transfected with Halo-Ecad-GFP (10 µg) using Neon electroporation. 24 h later, cells were re-plated on a collagen (or fibronectin)-coated glass-bottomed dish as previously described. To induce surface expression of SNAP-N^{FL}-mCherry. cells were incubated with complete medium containing doxycycline (Sigma, 2 µg/mL) for 18 h. To inhibit γ-secretase activity, cells were treated with DAPT (5 µM) and further incubated for 6 h. Cells were treated with 5 µM of an oligonucleotide bearing chloroalkane (Cl-$T_{60}ACTG_{10}$) for 45 minutes at 37°C and labeled with MFNs via the procedure described above.

Mechanogenetic regulation of artificial E-cadherin junctions. To induce MFN and hence cadherin clustering, the µMT was carefully directed towards a targeted subcellular location until the tip-to-membrane distance (*d*) reached 10 µm. As the tip approached the target membrane, the formation of an artificial E-cadherin junctions (cadAJs) was monitored every 5 minutes. After 30 min of mechanogenetic

966 stimulation, the spatial distribution of MFNs, artificial cadAJs, and Flot1 (or Notch1) was monitored using
967 time-lapse confocal fluorescence imaging. Cells were immediately fixed with 4% paraformaldehyde (Life
968 Technologies) in DPBS for 15 minutes and washed with DPBS 3 times for 5 minutes before
969 immunostaining.

Fluorescence labeling and Immunostaining

Fluorescence labeling of cells expressing SNAP- or Halo-tag proteins

Cells expressing SNAP- and/or Halo-tagged fusion proteins were labeled with BG- and/or chloroalkane
functionalized fluorescence dyes, respectively. Dox-inducible cell lines grown on a collagen I-coated
substrate were treated with doxycycline (2 µg/ml) 24 hr before labeling. Cells with transient receptor
expression were labeled with dyes 48 hr post-transfection. Dye-labeling was performed by treating the
cells with 5 µM fluorescence dye in serum containing media for 30 min. Cells were then washed 3 times
with complete media. For live cell imaging, cells were incubated with phenol-red free complete media.
For imaging of fixed cells, cells were washed with PBS, fixed with 4% PFA in PBS for 10 min, and then
washed thoroughly with PBS.

Immunofluorescence staining

Fixed cells were permeabilized with 0.5% tween-20 diluted in PBS for 15 minutes, then blocked by
incubation with blocking buffer (5% normal goat serum and 1% BSA in 1xPBS) for 1 hr in room
temperature. For immunostaining of surface ADAM10 and ADAM17 expression, the sample were directly
proceeded to blocking without a permeabilization step. Cells were incubated overnight at 4°C or 2 hr at
25°C with the primary antibodies: mouse monoclonal anti-ADAM10 (1:100; Santa Cruz Biotechnology),
rabbit polyclonal anti-PS1[4,5] (1:50), mouse polyclonal anti-Paxillin (1:500; BD Bioscience), rabbit
polyclonal anti-myc tag (1:200; abcam), mouse monoclonal anti VE-cadherin (1:400, BD Biosciences),
and Alexa Fluor 488 Phalloidin (1:400 dillution of 200 units/mL stock ; life technologies). All antibodies
were diluted in the 0.5x blocking buffer (2.5% normal goat serum and 0.5% BSA in 1xPBS). Following
primary antibody incubation, cells were washed in PBS for 5 min four times and incubated with Alexa
Fluor 405-conjugated goat anti-mouse (1:400, PS-1), Alexa Fluor 594-conjugated goat anti-mouse (1:400;
Paxillin), Alexa Fluor 488-conjugated goat anti-mouse (1:400; VE-cadherin), and Alexa Fluor 647-
conjugated goat anti-mouse (1:500, ADAM10), as appropriate. Nucleus staining was performed using
Hoechst 33342 (ThermoFisher) diluted in 1xPBS (1 µg/mL) for 15 min and then washed once. Cells were
incubated at room temperature in the dark for 1 hr, washed with PBS for 5 min three times, and imaged
by epifluorescence, confocal, or TIRF microscopy. Epifluorescence and TIRF microscopy was performed
with 60x Apo TIRF, 1.40 NA or 100x Apo TIRF, 1.49 NA oil objectives (Nikon) on a Nikon Ti Eclipse
microscope equipped with OBIS solid-state lasers (488, 552, and 647 nm, Coherent Inc.), a 300W Xenon
lamp (Sutter Instrument, Lambda LS), a motorized stage (ASI, MS-2000), and a temperature- and CO_2-
controlled stage top incubator (Okolab, Bold Line). Unless otherwise noted, confocal microscopy was
performed using Plan-Apo 60x, 1.4 NA or Plan-Apo 100x, 1.4 NA oil objectives (Nikon) on a Nikon A1R
laser scanning confocal microscope. Images were acquired using Galvano scanning mode and confocal
zoom of 3-4x magnification.

1005

Monitoring dynamic spatial localization of Notch intermediates

To activate Notch, we plated cells expressing SNAP-N[FL]-mCherry on a substrate coated with Dll4 fused
with a Fc fragment (Dll4-Fc)[6]. Briefly, a glass bottom dish (Lab-Tek II Chambered Coverglass,
ThermoFisher, or 7-mm glass-bottomed dish, MatTek) was coated with fibronectin (Hamster, 5 µg/ml)
and Dll4-Fc (2.5 µg/ml) for 1 hr at 37°C, and washed thoroughly with 10 ml of PBS. A negative control
dish was also prepared by coating it with fibronectin only. U2OS cells co-expressing SNAP-N[FL]-mCherry
and Ecad-GFP were plated and incubated with doxycycline (2 µg/mL), TAPI-2 (100 µM), and/or DAPT (5
µM). Different combinations of inhibitors were used to capture the respective intermediates (See **Fig. 2b**).
After 48 hr, cells were labeled with SNAP-647 (NEB, 5 µM) and then fixed as detailed above. Inhibitor
concentrations were maintained during wash and fixation steps. For DAPT washout experiments cells
were plated and activated via Dll4-Fc ligand as described above. DAPT inhibitor was removed by washing

in media at each time point (0, 0.5, 1.5, 3, 6, and 12 hrs). At each time point, cells were washed three
times with large volumes of PBS and fixed 4% with PFA. The spatial distribution of Notch intermediates
and cadAJs was monitored using spinning disk confocal fluorescence microscopy (Zeiss Cell Observer
Z1), equipped with Yokagawa spinning disk and Evolve 512 EMCCD Camera (Photometrics). Images
were obtained with Plan-Apo 63x, 1.4 NA or Plan-Apo 100x 1.46 NA oil objectives (Zeiss) with solid-state
lasers of 405, 488, 561 nm, and 647 nm. The microscope was controlled with Zeiss Zen software (Zeiss).

1023

Plasma membrane staining using DiI dyes

U2OS cells expressing Ecad-GFP were plated on a fibronectin-coated glass bottom dish (MatTek, D =
7.0 mm) at a density of 1×10^4 cells per dish. After 48 hr, the dish was filled with a complete McCoy's 5A
medium containing 5 µM of DiI plasma membrane labeling dyes (Invitrogen) and incubated for 10 min at
37°C, 5% CO_2. Cells were then washed 3 times with complete medium. The spatial distribution of cadAJs
and DiI membrane staining and Ecad-GFP was monitored using spinning disk confocal microscopy. For
live cell imaging, cells were incubated with phenol-red free media.

1031

Single-cell cleavage kinetics study of SNAP-NΔEGF-mC

A 6-channel µ-slide flow chamber (Ibidi, Vi 0.4) was coated with fibronectin (2.5 µg/mL) for 1 hr at 37°C
and washed with PBS four times. U2OS cells co-expressing SNAP-NΔEGF-mCherry and Ecad-GFP
were plated on the µ-slide flow chamber by applying 60 µL of single cell suspension at a density of 3 x
10^5 cells/mL. After 3 hr, the channel was filled with a complete McCoy's 5A medium containing
doxycycline (2 µg/mL), TAPI-2 (100 µM), and DAPT (5 µM). Cells were grown 48 hr in normal growth
medium to reach 70-80% confluency and form cadherin adherens junctions. Cells were labeled with BG-
Alexa Fluor 647 (NEB) for 30 min to stain cell surface NΔEGF. Multiple cells with stable cadAJs were
identified using large-area epi-fluorescence scanning (500 µm x 500 µm), and the spatial distribution of
SNAP-NΔEGF-mCherry at cadAJs under TAPI-2 and DAPT inhibition was imaged by confocal z-stack
(step size = 0.2 µm, total range of z stacks = 10 µm) scanning from basal to apical membranes. Then,
DAPT was removed and replaced from the cells by flowing complete medium containing doxycycline and
TAPI-2 at a flow rate of 50 µl/min for 10 minutes using a syringe pump. Localization of extracellular
(NECD) and intracellular (NICD) domain at the cadAJs before and during DAPT washout was monitored
every 30 minutes in multiple color channels (NICD, mCherry; NECD, AF647; cadAJ, GFP) by time-lapse
confocal z-stack microscopy for 12 hr. Time-lapse live cell confocal imaging was performed using a 60x
Plan-Apo oil objective (NA 1.4) on a Nikon A1 laser scanning confocal microscope equipped with an
environmental chamber maintaining cells at 37°C, 5% CO_2.

1050

Western Blot analysis

U2OS cells co-expressing Notch variants and Ecad-GFP (or Halo-Ecad-GFP) were incubated with culture
media containing doxycycline (2 µg/mL) and TAPI-2 (100 µM) in a 6-well plate at a density of 1×10^6 cells
per well. After 24 hr, cells were washed with ice-cold DPBS twice and lysed in RIPA (Invitrogen) or 1%
NP-40 (Invitrogen) supplemented with complete protease and phosphatase inhibitor cocktail (100x; Cell
Signaling Technology) at 4°C while gently shaking for 30 minutes. Insoluble fractions were removed by
centrifugation of the cell lysates at 13,000 r.p.m. for 10 minutes. Total protein concentrations in lysates
were determined by a BCA assay (Bio-Rad). 20 µg of whole cell lysates were then mixed with 4x Laemmli
sample buffer (Bio-Rad) with 10% β-mercaptoethanol (BME) and heated to 95°C for 5 minutes. For
western blot analysis of DNA-crosslinked heterodimers, the cell lysates were mixed with 4x Laemmli
sample buffer (Bio-Rad) but no BME and were used without boiling. Samples were then loaded into a 4-
15% Mini-Protein TGX precast gel (Bio-Rad) and were run at 70 V for 30 minutes and then 120 V for 45
minutes. Separated proteins were transferred to a PVDF membrane using the Trans Turbo Blot system
(Bio-Rad). Membranes were blocked for 1hr at room temperature in blocking solution (5% w/v nonfat dry
milk in 1x TBST). The membranes were probed with anti-V1744 NICD antibody (1:1000; Cell Signaling
Technology #4147), anti-SNAP (1:1000; NEB), anti-Notch1 (1:1000, Cell Signaling Technology #3447 or
#4380), and anti-β-actin (1:5000; Cell Signaling Technology #4970) antibodies overnight at 4°C with

gentle rocking. The membranes were washed in TBST three times for 5 minutes and incubated with an
anti-rabbit HRP conjugated antibody (Cell Signaling Technology, # at 1:2000 for NICD, mCherry, SNAP
detection and at 1:10000 for β-actin detection). The target proteins were visualized by
chemiluminescence using an ECL detection kit and a ChemiDoc MP imaging system (Bio-Rad).
Quantification of band intensities by densitometry was carried out using the Image Lab software (Bio-
Rad).

1074

Spatial mutation of SNAP-NΔEGF-mCherry via DNA crosslinking

DNA-mediated crosslinking of SNAP-NΔEGF-mCherry with Halo-Ecad-GFP. DNA crosslinkers including benzylguanine (BG)- and chloroalkane (Cl) modified oligonucleotides were synthesized as previously described[2,7]. To prepare 10x crosslinking DNA stock solution, complementary BG- and Cl-modified oligonucleotides were hybridized in situ. BG-$T_{10}(ACTG)_5$ and Cl-$T_{10}(CAGT)_5$ were mixed at equimolar concentration (20 µM) in PBS, incubated at 95°C on a dry heat block for 5 min, and slowly cooled down to room temperature for 2 hr. U2OS cells co-expressing SNAP-NΔEGF-mCherry and Halo-Ecad-GFP were cultured in a 6-well plate for western blot analysis at a density of 1×10^6 cells per mL or in a channel of an Ibidi µ-slide for confocal imaging analysis at a density of 3×10^5 cells per mL. Cells were grown to 70-80% confluency for typically 24 hr, followed by overnight incubation with complete medium containing doxycycline (2 µg/mL), TAPI-2 (100 µM) and DAPT (5 µM). Cells were then serum starved with 2 ml of serum-free medium with doxycycline, DAPT, and TAPI-2 for 6 hrs. Before adding DNA crosslinkers, cells were washed and placed in 450 µl of serum-free media. 50 µl of prewarmed 10x DNA crosslinker stock solution were added to each well and incubated at 37°C. Western blot analysis to validate receptor crosslinking were performed after 30-minute incubation of the DNA crosslinkers as detailed above.

Live cell confocal time-lapse imaging. After overnight incubation with the DNA crosslinkers, imaging was performed on an inverted laser scanning confocal microscope (Nikon A1) equipped with an environmental chamber at 37°C and 5% CO_2. Images were obtained with a Plan-Apochromat 60x, 1.4 NA oil objective (Nikon) with solid-state lasers of 405, 488, 561 nm, and 647 nm. Additionally, the microscope was equipped with Ti-E Perfect Focus System (Nikon). To examine the effect of DNA-mediated crosslinking on spatial distribution of SNAP-NΔEGF-mCherry at cadAJs, multiple cadAJs were imaged in entirety from basal to apical sides for Halo-Ecad-GFP and SNAP-NΔEGF-mCherry using a 488 nm and 561 nm laser respectively, for a 12 µm range at a z-step size of 0.25 µm. To monitor dissipation of SNAP-NΔEGF-mCherry at cadAJs upon removal of DAPT inhibition, fresh phenol red-free McCoy's 5A medium containing doxycycline and TAPI-2 was introduced into the channel using a syringe pump for 10 min, and confocal z-stack images of the previously selected cadAJs were acquired every 30 minutes for 6 hr. Images were acquired using NIS-element software (Nikon), and image post-processing and analyses were done using Fiji/ImageJ and custom-built codes.

Spatial mutation of SNAP-NEXT-mCherry via molecular pendant addition

Synthesis of BG-modified polyethylene glycol (PEG). Amine-functionalized PEGs with different molecular weights and structures were purchased from Creative PEGWorks (NH$_2$-PEG3.4k), Sigma (NH$_2$-*b*PEG20k), and NanoCS (NH$_2$-*ℓ*PEG20k) and used without further purification. BG-functionalization of PEGs was performed by amine-NHS (N-hydroxysuccinimide; NEB) coupling reaction. Briefly, NH$_2$-PEG (0.5 µmol), BG-GLH-NHS (2.4 mg, 5 µmol), and N,N-dimethylaminopyridine (0.73 mg, 6 µmol; Sigma) were dissolved in anhydrous dimethylsulfoxide (DMSO). The mixture was reacted overnight with constant shaking. Crude products were recovered by evaporating DMSO using a Speed-Vac concentrator (Vacufuge, Eppendorf), reconstituted in 500 µl deionized water, and removed insoluble precipitates by centrifugation at 14,000 r.p.m. for 10 minutes. The BG-modified PEG was then dissolved in 200 µl in deionized water and purified by reverse-phase high performance liquid chromatography with an Agilent Eclipse XDB C-18, 5 µm, 4.6 x 250 mm column using an elution gradient of 5-75% acetonitrile in 0.02% trifluoroacetic acid.

Synthesis of BG-modified DNA-streptavidin conjugates. DNA oligonucleotides bearing biotin- and BG-functional groups were synthesized by reacting biotin-$(ACTG)_5$-NH$_2$ (IDT DNA) with BG-GLH-NHS as described above. Equimolar amounts of streptavidin (10 nmol) and BG-DNA-biotin (10 nmol) were

dissolved in PBS (0.5 ml) for 2 hr, forming streptavidin-BG complex. The solution was concentrated to
approximately 50 µl using an Amincon centrifugal filter (MWCO: 30k) and then diluted again with 0.45 ml
of PBS. This concentration and reconstitution step was repeated three times to remove unconjugated
DNA.

Synthesis of BG-modified human IgG. hIgG (10 mg) and BG-GLA-NHS (0.82 mg) were dissolved in 850
µl of PBS and 150 µl of anhydrous DMSO, respectively. Two solutions were mixed and reacted for 2 hr
at room temperature with gentle shaking. The solution was desalted with NAP-10 and then with NAP25
pre-equilibrated with PBS. The proteins were further concentrated until the final volume is 300-500 µl
using Amicon centrifugal filter (MWCO: 30k). The IgG concentration was determined by measuring the
absorbance at 280 nm.

Spatial mutation of SNAP-NEXT-mCherry using the BG-modified macromolecules. U2OS cells co-
expressing SNAP-NEXT-mCherry and Ecad-GFP were incubated with a complete McCoy's 5A medium
containing doxycycline (2 µg/ml), TAPI-2 (100 µM), DAPT (5 µM), and respective BG-modified
macromolecules (10 µM). After 24 h, cells were fixed and imaged by confocal microscopy to determine
the enrichment factor of SNAP-NEXT-mCherry at cadAJs. Images were taken with a 100x objective and
3x confocal zoom. 20 stage positions per each treatment were manually selected and their coordinates
were stored in the computer. In each position, confocal z stacks of DAPI, Ecad-GFP and Notch-mCherry
were acquired for a 12 µm range at a Z step-size of 0.25 µm to monitor the cadAJs in their entirety from
basal to apical sides. To assess the levels of Notch activation, a set of identical experiment but without
DAPT was performed. After 24 h, cells were fixed, stained with DAPI, and imaged by confocal microscopy
to determine nuclear mCherry signals. Images were taken with a 60x objective and 1x confocal zoom. 5
stage positions per each condition were selected manually. For each position, a confocal large-area scan
of DAPI, Ecad-GFP, and SNAP-NEXT-mCherry was acquired for a 1 mm x 1 mm area.

1143

Plate-bound Dll4 Notch activation in high-density grouped versus solitary cells

To activate Notch, we plated SNAP-N^{FL}-Gal4 reporter cells on a substrate coated with Dll4-Fc as
detailed above. Two different cell seeding densities were used: We plated cells with a density of 1 x 10^3
cells per 10 mm glass-bottomed dish (MatTek, No. 1.5 glass), predominantly yielding solitary cells. We
also plated cells with a density of 1 x 10^4 cells per dish, predominantly yielding high-density grouped
cells.

Plate-bound E-cadherin Notch activation experiment. Glass-bottomed dishes (MatTek, #1.5, D = 10 mm)
were coated with recombinant human E-cadherin-Fc (50 µg/ml, R&D systems), recombinant human Dll4-
Fc (2.5 µg/ml, Sino Biological), and fibronectin (5 µg/ml, Sino Biological) diluted in PBS for 1 hr at 37°C,
and rinsed with 10 ml PBS with calcium and magnesium (UCSF cell culture facility). The U2OS SNAP-
N^{FL}-Gal4 reporter cells were transfected with Ecad-GFP (10 µg) via electroporation, incubated overnight,
and re-plated onto a fibronectin, E-cadherin-Fc, and Dll4-Fc coated glass-bottomed dish at a density of
0.3 x 10^5 cells/ml, same as the solitary cell assay. Negative control experiment was performed with the
cells plated on E-cadherin-Fc and fibronectin coated glass-bottomed dishes without Dll4-Fc coating.

Time-lapse epifluorescence imaging. All cells were treated with 2 µg/ml doxycycline (sigma-aldrich) at
the time of plating. 2 hr post-plating, cells were imaged using time-lapse microscopy. For a high-density
cell seeding assay, several groups of cells having cell-cell contacts were manually identified and their
coordinates were stored. For a solitary cell assay, a number of solitary cells without any prior cell-cell
contact were manually identified and their coordinates were stored. While maintaining live cells on a
microscope stage with a top stage incubator, time-lapse fluorescence images were acquired in GFP and
mCherry channels. In each position, the microscope (Nikon) first found focuses using the Perfect Focus
System (Nikon), took a DIC image, and two fluorescent images (GFP, mCherry). Epifluorescence images
were obtained with an inverted microscope (Nikon, Ti Eclipse) equipped with 300W Xenon lamp (Sutter
Instrument, Lambda LS), a motorized stage (ASI, MS-2000), and a temperature- and CO$_2$-controlled
stage top incubator (Okolab, Bold Line). Images were taken with 40x (CFI Plan fluor, N.A. 1.3, Nikon)
objective lens. The microscopy setup was controlled using µ-manager software.

1170

CRISPR editing to generate E-cadherin and N-cadherin knockout mutants

CRISPR/Cas9 was used to knock out E-cadherin and N-cadherin expression from U2OS SNAP-N^{FL}-Gal4 reporter cells. The genes coding for the E-cadherin (CDH1) and N-cadherin (CDH2) protein from homo sapiens (gene ID: ENSG00000039068 and ENSG00000170558) were truncated by a CRISPR/Cas9 paired sgRNAs excision strategy[8,9] For the fragment deletion of genomic DNA, we used a pair of gRNAs against the target locus of CDH1 (Exon 1 & 2 (940bp deletion) or 13 & 14 (~4712bp deletion) (**Extended Data Fig. 9b**) and CDH2 (Exon 1 & 2 (~29,255bp deletion) (**Extended Data Fig. 9c**) genes.

sgRNA design and expression vector cloning. Cas9 and sgRNAs were expressed using the CMV promoter-driven Cas9-2A-mRFP-2A-Puro plasmid (hereafter, Cas9-puro vector) and the hU6 promoter-driven sgRNA plasmid (Toolgen), respectively. To design sgRNAs for fragmental deletion of target loci of genes, all candidate sgRNA target sites with a protospacer adjacent motif (PAM; 5'-NGG-3') within the coding sequence of CDH1 and CDH2 were initially identified. For efficient deletion, selected sgRNAs for the candidate target sites were evaluated with DeepSpCas9 sgRNA prediction tool (http://deepcrispr.info/DeepSpCas9/)[10]. sgRNAs with high DeepSpCas9 score were selected and sgRNA oligonucleotides annealed and cloned into the vector as previously described[11]. Sequences of the vectors and sgRNAs listed here are available upon request.

Generation of single-cell derived knock-out clones. For CDH1 knockout, SNAP-N^{FL}-Gal4 reporter cells were transfected with plasmid mixtures containing Cas9-puro, U6-sgRNA encoding individual sgRNAs at a weight ratio of 1:2 using the Neon system. For CDH1 & CDH2 knockout, cells were transfected with plasmid mixtures containing Cas9-puro, U6-sgRNA targeting CDH1 loci, and U6-sgRNA targeting CDH2 loci at a weight ratio of 1:1:1 using the Neon system. One day after transfection, puromycin was added to the culture media at a final concentration of 2.5 µg ml^{-1}. Three days after transfection, the pooled cells were analyzed for the indel efficiency of sgRNA pairs using T7E1 assay. To obtain single cell-derived clones containing the fragment deletion, we plated the cells after puromycin selection into 96-well plates at an average density of 0.25 cells/well. 14 days after plating, individual clones were isolated and analyzed using PCR and gel electrophoresis of genomic DNA to check the deletion and wild-type alleles as previously described[8]. We next sequenced the genomic DNA of the clones containing targeted deletions to check if the two cleavage sites were joined by the generation of indels. Sequencing of genomic regions including the target sequence was performed as previously described[12]. Briefly, PCR amplicons that included the junction regions of the deleted lncRNA target sites were cloned into the T-Blunt vector (Promega) and sequenced using universal M13FP or RP primers.

T7E1 assay. The T7E1 assay was performed as previously described[8,13]. Briefly, genomic DNA was isolated using the Wizard Genomic DNA purification Kit (Promega) according to the manufacturer's instructions. The region including the target site was nested PCR-amplified using appropriate primers. The amplicons were denatured by heating and annealed to allow the formation of heteroduplex DNA, which was treated with 5 units of T7 endonuclease 1 (NEB) for 20 min at 37°C followed by analysis using 2% agarose gel electrophoresis. Mutation frequencies were calculated as previously described based on the band intensities using Image J software and the following equation[14]: mutation frequency (%) = 100 × (1 − (1 − fraction cleaved)$^{1/2}$), where the fraction cleaved is the total relative density of the cleavage bands divided by the sum of the relative density of the cleavage bands and uncut bands.

RT-PCR. Total RNA was extracted from wild-type SNAP-N^{FL}-Gal4 reporter (WT) cells and knockout clonal cells using TRIzol (Ambion) or an RNeasy Kit (QIAGEN), after which complementary DNA (cDNA) synthesis was performed using a DiaStarTM RT Kit (SolGent Co., Ltd.). The synthesized cDNA was subjected to quantitative PCR (qPCR) in triplicate using an Applied Biosystems StepOnePlus Real Time PCR System with PowerSYBR Green PCR Master Mix (Applied Biosystems). Gene expression was normalized to that of the CDH1 gene in WT cells. Error bars represent the standard deviation (s.d.) of the mean of triplicate reactions. Primer sequences for qPCR are available upon request.

Notch activation assay. WT cells, CRISPR *CDH1* knock-out cells (CDH1$^{-/-}$), *CDH1*$^{-/-}$ transfected with E-cadherin-GFP (CDH1$^{-/-}$ + E-cad), and *CDH1*$^{-/-}$ transfected with N-cadherin (CDH1$^{-/-}$ + N-cad) were plated on 8 well Nunc Lab-Tek II Chambered Coverglass pre-coated with recombinant human Dll4-Fc (2.5 µg/ml) and fibronectin (5.0 µg/ml) as previously described. All cells were plated at a density of 30,000 cells per well. After 24 hr incubation with doxycycline (2 µg/ml), cell cytoplasm and nucleus were stained with

CellTracker CMFDA dye (Invitrogen) and Hoechst 33342 (ThermoFisher), respectively. The cells were then fixed with 4% PFA for 15 minutes at room temperature and proceeded to epifluorescence and confocal imaging.

Aβ40 and Aβ42 ELISA

1×10^6 of wild-type or CDH1/CDH2 knockout cells were transfected with APP$^{\Delta YENPTY}$-EGFP (10 µg) and plated on 6-well tissue culture plate. After 48 hr incubation, cells were lysed using 100 µl of RIPA buffer supplemented with 1x protease/phosphatase inhibitor cocktail (ThermoFisher) for 10 minutes on ice. The cells were scraped off the plates using a cell scraper, and the lysates were transferred to 1.5 ml microcentrifuge tubes and incubated for 20 minutes at 4°C with gentle agitation. Lysates were centrifuged at 14,000xg 10 minutes at 4°C to remove insoluble portions, and the supernatant were transferred to a new tube and stored at -80C. Cell lysates were analyzed for Aβ40 and Aβ42 contents using Aβ ELISA kits from Invitrogen. All ELISAs were performed according to the manufacturer's protocols. Briefly, 50 µl/well of samples, followed by 50 µl/well of detection antibodies, were applied to Aβ40 or Aβ42 coated 96-well plate, and incubated for 3 hr at room temp with shaking at 300 rpm. After washing step, 100 µl of HRP-IgG solution was applied and incubated for 30 min at room temp with shaking. After washing step, 100 µl of stabilized substrate solution was applied and incubated for 30 min at room temp with shaking. Finally, 100 µl of stop solution was applied. Absorbance at 450 nm was read and analyzed using a plate reader (Biotek Synergy 2). Wells were washed with wash buffer 4 times between each incubation step. Standard curves were generated using recombinant Aβ40 and Aβ42 provided by the manufacturer (Invitrogen).

Image Processing and Analysis

Cadherin junction colocalization analysis. Colocalization analysis was carried out in ImageJ, using thresholding to identify cadAJs and then applying the JACOP plugin to quantify colocalization using Pearson coefficient, Mander's overlap coefficients, and cross-correlation analysis.

Confocal 3D z-stack image processing. Custom python code was used for automatic segmentation and junction intensity ratio analysis for Notch activation and truncation studies. Code is available at (**https://github.com/kmsouthard/JunctionAnalysis**) In brief, resliced z-stacks of cell-cell interfaces were thresholded to identify the cadAJs and membrane Notch signal. Notch membrane intensity was measured inside the junction and outside the junction for each cell forming the junction. The ratio of junctional intensity was calculated as ratio = $I_{junc}/(I_{cell1}+ I_{cell2})$ as deviations from the expected intensity at the junction is a function of the sum of each cell's expression level.

Intracellular mCherry nuclear translocation analysis. mCherry nuclear translocation analysis was carried out in ImageJ. GFP images were used for automated identification of cell edges and segmentation of single cells. DAPI images were used for automated identification of nucleus by implementing the Otsu thresholding method. Nuclear mCherry fluorescence data was extracted from nuclear segments by calculating the integrated fluorescence within the nucleus and subtracting a background scattering signals. In Fig 3h, nuclear mCherry fluorescence intensities for NEXT cells treated with the macromolecular pendants were rescaled to make the intensity of N^{FL} and NEXT to 0.002 and 1.0, respectively, which are identical to the normalized band intensities of N^{FL} and NEXT measured by western blot.

Quantification of single-cell fluorescence. Single-cell tracking and nuclear mCherry fluorescence signal analysis of UAS-Gal4 reporter cells were performed with ImageJ, as previously described[1,15,16].

Western blot quantification. Quantification of band intensities by densitometry was carried out using the Image Lab software (Bio-Rad). Band intensities of NICD in each lane were normalized by band intensities of loading control, β-actin in the corresponding lane.

Estimation of Protein heights

Protein heights including extended Notch height was estimated by measuring the structural size of each domain (i.e. EGF, NRR, SNAP) in Pymol (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.) and then creating an additive estimate based on the number of domains in the full-length Notch construct and each Notch truncation.

Molecular dynamics simulations

All MD simulations were conducted using the GROMACS package[17]. Polarized MARTINI 2.2 parameters were used for the simulations[18,19]. The system was composed of 54 PPCS, 54 DPPC, 108 DPPS, 288 DIPC, 216 CHOL, and 10108 water molecules with a molar composition of lipid was CHOL:PPCS:DPPC:DIPC = 3.0 : 1.5 : 1.5 : 4.0 in upper layer and CHOL:DPPS:DIPC = 3.0 : 3.0 : 4.0 in lower layer. To create immobilized lipids, we increased the mass of the phosphorus atom within DPPS by a factor of 1,000, keeping everything else the same in the parameter file. The pressure was set at 1.0 bar with a semi-isotropic parrinello-rahman coupling with compressibility 4.5×10^{-5} $bar^{-1}$and the temperature was set to 295 K using nose-hoover coupling. Each system was neutralized and brought to a concentration of 0.15 M with randomly placed sodium and chloride ions. We employed the LINCS algorithm to constrain to bond lengths[20]. A time step of 20 fs was used with an update of the neighbor list every 10 steps, which are typical values employed in MARTINI simulations. Each simulation was run afterwards for 12 µs, the last 3 µs of which was used for analysis. The MD simulations were analyzed using the in-built GROMACS tools. MDAnalysis libraries[21,22] were used for calculating diffusion constant of lipid component and *g_energy* was used for calculating interlayer interaction.

Statistical analysis. Statistical analysis was performed in GraphPad Prism 8.0 (GraphPad) or Microsoft Excel. Figure legends indicate all statistical tests used in the figure. Unless otherwise noted in the figure legends, statistical differences were determined using Student's t-test (two-tailed unpaired or paired t-test, depending on the experiment) when only two groups were compared or by ordinary one-way ANOVA followed by Tukey posthoc test when multiple groups were analyzed. The number of samples ('n') used for each experimental analysis is indicated in the figure legends. Sample sizes of sufficient power were chosen on the basis of general standards accepted by the field. In all cases, statistical significance was assumed for probability (P) < 0.05. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001; ns indicates when no significant difference was detected.

Data availability. All raw images, immunoblot gel images, and analyzed data are available on request. Additional data that support the findings of this study are available from the corresponding author upon reasonable request.

References for Methods

1. Seo, D. *et al.* A Mechanogenetic Toolkit for Interrogating Cell Signaling in Space and Time. *Cell* **165,** 1507–1518 (2016).

2. Kwak, M. *et al.* Small, Clickable, and Monovalent Magnetofluorescent Nanoparticles Enable Mechanogenetic Regulation of Receptors in a Crowded Live-Cell Microenvironment. *Nano Lett.* **19,** 3761–3769 (2019).

3. Kim, J.-W. *et al.* Single-cell mechanogenetics using monovalent magnetoplasmonic nanoparticles. *Nat Protoc* **12,** 1871–1889 (2017).

4. Georgakopoulos, A. *et al.* Presenilin-1 forms complexes with the cadherin/catenin cell-cell adhesion system and is recruited to intercellular and synaptic contacts. *Molecular Cell* **4,** 893–902 (1999).

5. Marambaud, P. A presenilin-1/gamma-secretase cleavage releases the E-cadherin intracellular domain and regulates disassembly of adherens junctions. *The EMBO Journal* **21,** 1948–1956 (2002).

6. Andrawes, M. B. *et al.* Intrinsic Selectivity of Notch 1 for Delta-like 4 Over Delta-like 1. *J. Biol. Chem.* **288,** 25477–25489 (2013).

7. Liang, S. I. *et al.* Phosphorylated EGFR Dimers Are Not Sufficient to Activate Ras. *Cell Rep* **22,** 2593–2600 (2018).

8. Lee, H. J. *et al.* En bloc and segmental deletions of human XIST reveal X chromosome inactivation-involving RNA elements. *Nucleic Acids Res.* **47,** 3875–3887 (2019).

9. Pulido-Quetglas, C. *et al.* Scalable Design of Paired CRISPR Guide RNAs for Genomic Deletion. *PLoS Comput Biol* **13,** e1005341 (2017).

10. Kim, H. K. *et al.* SpCas9 activity prediction by DeepSpCas9, a deep learning–based model with high generalization performance. *Sci Adv* **5,** eaax9249 (2019).

11. Ramakrishna, S. *et al.* Gene disruption by cell-penetrating peptide-mediated delivery of Cas9 protein and guide RNA. *Genome Research* **24,** 1020–1027 (2014).

12. Ramakrishna, S. *et al.* Surrogate reporter-based enrichment of cells containing RNA-guided Cas9 nuclease-induced mutations. *Nat Commun* **5,** 167 (2014).

13. Gopalappa, R., Suresh, B., Ramakrishna, S. & Kim, H. H. Paired D10A Cas9 nickases are sometimes more efficient than individual nucleases for gene disruption. *Nucleic Acids Res.* **46,** e71–e71 (2018).

14. Guschin, D. Y. *et al.* A rapid and general assay for monitoring endogenous gene modification. *Methods Mol. Biol.* **649,** 247–256 (2010).

15. Nandagopal, N. *et al.* Dynamic Ligand Discrimination in the Notch Signaling Pathway. *Cell* **172,** 1–32 (2018).

16. Sprinzak, D. *et al.* Cis-interactions between Notch and Delta generate mutually exclusive signalling states. *Nature Publishing Group* **465,** 86–90 (2010).

17. Van Der Spoel, D. *et al.* GROMACS: fast, flexible, and free. *J Comput Chem* **26,** 1701–1718 (2005).

18. de Jong, D. H. *et al.* Improved Parameters for the Martini Coarse-Grained Protein Force Field. *J Chem Theory Comput* **9,** 687–697 (2013).

19. Marrink, S. J., Risselada, H. J., Yefimov, S., Tieleman, D. P. & de Vries, A. H. The MARTINI force field: coarse grained model for biomolecular simulations. *J Phys Chem B* **111,** 7812–7824 (2007).

20. Hess, B., Bekker, H., Berendsen, H. & Fraaije, J. LINCS: A linear constraint solver for molecular simulations. *J Comput Chem* **18,** 1463–1472 (1997).

21. Michaud-Agrawal, N., Denning, E. J., Woolf, T. B. & Beckstein, O. MDAnalysis: a toolkit for the analysis of molecular dynamics simulations. *J Comput Chem* **32,** 2319–2327 (2011).

22. Chavent, M. *et al.* Methodologies for the analysis of instantaneous lipid diffusion in md simulations of large membrane systems. *Faraday Discuss.* **169,** 455–475 (2014).