Initial spindle positioning at the oocyte center protects against incorrect

kinetochore-microtubule attachment and aneuploidy in mice

Jessica N. Kincade and Ahmed Z. Balboula*

Animal Sciences Research Center, University of Missouri, Columbia, MO 65211, USA

Running title: Central spindle positioning protects against aneuploidy

Keywords: Aneuploidy, Nucleus Positioning, Spindle Positioning, Oocyte, Meiosis

*Corresponding author: Ahmed Z. Balboula, Animal Sciences Research Center, University of Missouri, Columbia, MO 65211, USA, abalboula@missouri.edu

ABSTRACT

Spindle formation and positioning are two critical events that must be regulated tightly to avoid erroneous chromosome segregation. Following nuclear envelope breakdown (NEBD), the spindle is predominantly assembled centrally before migrating towards the cortex to allow the first asymmetric division. The biological significance of primary central positioning of the spindle has not yet been studied. Because the spindle forms where NEBD occurs, nucleus position greatly influences the site of spindle assembly. We find that nucleus positioning is a dynamic process that depends on maternal age. Moreover, we show that preferential initial spindle positioning at the oocyte center is an insurance mechanism to avoid premature exposure to cortical Cdc42 signaling that hinders proper kinetochore-microtubule attachments, thereby protecting against the development of aneuploid gametes. These findings will help in understanding why female gametes are notoriously associated with high rates of aneuploidy, the leading genetic cause of miscarriage and infertility in women.

INTRODUCTION

Infertility is a globally prevalent issue affecting approximately 122 million women worldwide [1]. Errors in chromosome segregation during meiosis result in aneuploidy, the leading genetic cause of infertility, spontaneous abortion, and congenital birth defects such as down syndrome[2,3]. Aneuploidy occurs in females of all ages, but women of advanced age experience elevated risk of aneuploidy [4]. Given that chromosome segregation errors occur most frequently in female gametes (as opposed to male gametes), particularly within the first meiotic division (Meiosis I, MI)[5,6], it is crucial to understand the molecular mechanisms regulating female MI.

Oogenesis is a complex process that begins during fetal development in female mammalian species, shortly after conception. During late pregnancy, oocytes enter MI and duplicate their chromosomes [7]. Shortly after birth, all germ cells undergo a prolonged arrest that is maintained until puberty. Hormonal cues prior to ovulation stimulate a subset of oocytes to resume meiosis [7]. In fully-grown prophase I-arrested oocytes, the nucleus (germinal vesicle, GV) localizes mainly at the center [8]. *In vitro*, peripherally localized nuclei migrate towards the center of the oocyte, regulated by a meshwork of cytoplasmic actin. This meshwork of cytoplasmic actin creates a gradient of velocity (higher near the cortex, lower near the center) induced by Myosin Vb activity and relocates the nucleus centrally through gentle movement governed by active diffusion, rather than exerting a force on the nucleus [9-11]. However, the biological significance of central nucleus positioning is largely unknown.

Upon meiotic resumption, prophase I-arrested oocytes undergo nuclear envelope breakdown (NEBD). Throughout the NEBD process, the microtubule organizing centers (MTOCs) undergo

a three-step fragmentation before clustering and sorting into two dominant spindle poles that are

integral in forming a bipolar spindle at the center of metaphase I (Met I) oocytes [12-15]. The

spindle dictates the plane of cell division [16], and, therefore, must migrate to the oocyte cortex to

ensure the highly asymmetric meiotic division [16,17]. Extruding a small polar body through a

highly asymmetrical division is critical as it allows the egg to retain a majority of the

cytoplasmic store of maternal RNAs and proteins synthesized during the growth phase which are

essential for fertilization and early embryonic development [18,19]. However, the necessity of

central spindle assembly prior to cortical spindle migration is unknown.

Here we show that nucleus positioning is a dynamic process that is affected by maternal age. A

majority of oocytes collected from sexually mature mice had a centrally positioned nucleus,

whereas a majority of oocytes from prepubertal mice had a peripherally positioned nucleus. We

find that peripheral nucleus positioning is associated with chromosome misalignment, abnormal

kinetochore-microtubule (K-MT) attachments and aneuploidy. Importantly, we find that central

spindle positioning post-NEBD is critical to avoid deleterious cortical Cdc42 signals that hinder

the fidelity of forming kinetochore-microtubule (K-MT) attachments, thereby protecting against

aneuploidy in mammalian eggs.

RESULTS

Nucleus Positioning – A dynamic process

Because the spindle forms where NEBD occurs whether at the center (Supplementary Fig. 1A)

or at the cortex (Supplementary Fig. 1B), we sorted full-grown prophase I-arrested oocytes

according to the positioning of the nucleus into three groups: central, intermediate, and

peripheral; herein after, these groups are referred to as central GV oocytes, intermediate GV

oocytes, or peripheral GV oocytes. Peripheral GV oocytes had a distance of 5 µm or less from

the nucleus to the nearest cortex, intermediate GV oocytes remained at a distance of 5 to 15 µm

from the nucleus to the nearest cortex, and central GV oocytes maintained more than 15 µm

distance from the nucleus to the nearest cortex (Fig. 1A). First, we observed the proportions of

these groups of oocytes *in vitro* after collection. Oocytes with a centrally located nucleus were

the most prevalent, contributing nearly 50.96% of all oocytes collected. Oocytes with an

intermediately positioned nucleus accounted for nearly 24.22% of all collected oocytes, while

peripheral oocytes represented only 24.81% of collected oocytes (Fig. 1B).

To determine whether nucleus positioning proportions *in vitro* are similar to those *in vivo*,

ovaries were collected at four ages (2 weeks, 3 weeks, 5 weeks, and 8 weeks) in order to display

nucleus positioning throughout reproductive development. All oocytes and their corresponding

nucleus positioning were examined from histological sections of ovaries stained with

Hematoxylin and Eosin (Fig. 1C,D). Similar to our *in vitro* results, ovaries from mice aged 8

weeks showed a majority of oocytes with a centrally positioned nucleus (63.52%). Interestingly,

younger mice (2 & 3 weeks of age) showed a majority of peripherally located nucleus (50.46%

and 55.63%, respectively), while mice aged 5 weeks showed very similar proportions of oocytes

with both a centrally and peripherally located nucleus (42.86% and 43.25%, respectively, Fig.

1D,E), suggesting that, before sexual maturation, oocytes have an eccentrically positioned

nucleus (likely incompetent) prior to central relocation after sexual maturation.

Oocytes with a peripherally located nucleus associate with higher rates of chromosome

misalignment and aneuploidy

Upon meiotic resumption, NEBD is shortly followed by the formation of a bipolar spindle and

the alignment of chromosomes at the metaphase plate. The occurrence of peripheral nucleus

positioning is correlated with a decreased maturation rate [8]. To assess the meiotic progression,

oocytes were collected from mice aged 6-8 weeks and matured for 16 h (Met II). The majority

(~92%) of centrally located GV oocytes were able to undergo NEBD. In contrast, peripherally

located GV oocytes showed a significant decrease (~61%) in the percentage of NEBD (Figure

2A). However, using time-lapse microscopy, we found no significant difference in the timing of

NEBD between centrally located GV oocytes and peripherally located GV oocytes (Figure 2B).

Similarly, a majority (~81%) of central GV oocytes were able to extrude the first polar body

(PBE), while peripheral GV oocytes were less successful in extruding the first polar body

(~59%, Fig. 2C). Once again, there was no significant difference in the timing of PBE when both

centrally located GV oocytes and peripherally located GV oocytes were compared (Fig. 2D).

To assess the chromosome alignment and spindle morphology among groups, oocytes were

matured *in vitro* for either 7 h (Met I) or 16 h (Met II), fixed, and immunostained with α-tubulin

to label the spindle and stained with DAPI to label DNA. Chromosome alignment was assessed

according to the parameters previously described [20]. We found a significant increase in the

percentage of chromosome misalignment at both Met I or Met II in oocytes with a more

peripherally positioned nucleus when compared to oocytes with a centrally located nucleus.

Intermediate GV oocytes also showed significantly higher rates of chromosome misalignment

when compared to central GV oocytes at Met I, but not at Met II (Fig. 2E,F,H,I). Spindle

morphology was assessed using the criterion that a normal meiotic spindle is barrel shaped with

two clearly defined spindle poles. Spindle morphology did not vary significantly among groups

either at Met I or Met II (Fig. 2E,G,H,J). Because chromosome misalignment is highly

associated with aneuploidy, we then assessed aneuploidy in Met II eggs using *in situ*

chromosome counting [21,22]. Strikingly, segregation errors occurred in oocytes with a peripherally

located nucleus (~50.15%) approximately 5 times more frequently than in oocytes with a

centrally located nucleus (~10.37%, Fig. 2K,L).

Nuclear envelope breakdown occurs during central relocation of peripherally located nucleus

To understand the behavior of the nucleus and its positioning during MI, we employed time-

lapse microscopy to image live oocytes. When peripheral GV oocytes were arrested at prophase I

(using milrinone, a phosphodiesterase inhibitor) to prevent NEBD, only ~15% of the oocytes

remained peripheral within 3 h of incubation, while the majority (~85%) relocated to the center

of the oocyte (Supplementary Fig. 2 and Movie 1), confirming that the nucleus favors central

movement, as previously reported [9,23,24]. When the oocytes were allowed to mature in milrinone-

free medium, nuclei initially located at the periphery of the cell underwent 10 ± 1.256 μm on

average of central movement before experiencing NEBD. Intermediately located nucleus and

centrally located nucleus underwent central movement and traveled a distance of 6 ± 0.6024 μm

and 4 ± 0.6783 μm, respectively, before NEBD (Supplementary Fig. 3A). However,

chromosome alignment in oocytes with a peripherally located nucleus remained relatively closer

to the oocyte cortex in comparison to chromosome alignment in oocytes with a central nucleus

(Supplementary Fig. 3B,C). It is of note that the timing of chromosome alignment did not vary

significantly among groups (Supplementary Fig. 3B,D). Why oocytes favor central nucleus

positioning is not fully understood.

Incorrect kinetochore-microtubule attachments are increased in oocytes with a peripherally located nucleus

While aneuploidy can occur as a result of many different errors, one important cause of aneuploidy is a weakened or non-functional spindle assembly checkpoint (SAC). Typically, the SAC is inactivated when MTs are attached correctly to all kinetochores, but a weakened or non-functional SAC may allow the progression of the cell to anaphase without proper K-MT attachments, a phenomenon that is highly associated with aneuploidy. To assess the SAC function in oocytes with a central, intermediate, or peripheral nucleus, oocytes were treated with nocodazole, a MT depolymerizing agent that is known to maintain the SAC active and thereby prevent PBE by inducing Met I arrest [22,25]. Expectedly, positive control oocytes treated by ZM447439, an Aurora Kinase B/C inhibitor that is known to disrupt the SAC function at a high dose (10µM), extruded a PB even in the presence of nocodazole [26]. In contrast to ZM447439 treated oocytes, all other oocytes (central, intermediate, and peripheral) failed to extrude a PB in the presence of nocodazole, suggesting that the SAC is intact, functional, and therefore not contributing to aneuploidy in peripheral GV oocytes (Supplementary Fig. 4).

Improper K-MT attachment is another major contributor of aneuploidy. Because cortically facing centromeres have relatively high rates of unstable K-MT attachments when compared to those away from the oocyte cortex [27] and peripheral nucleus positioning correlates with peripheral spindle assembly (near the cortex), we hypothesized that K-MT attachments are perturbed in peripheral GV oocytes. In meiotic oocytes, even if the SAC is satisfied by present K-MT attachment, it is not necessary that all attachments are correct and homologous chromosomes are bioriented [20,28]. To assess K-MT attachments, Met I oocytes (~7 h) were exposed to a brief cold shock (6 minutes) in order to depolymerize labile unattached MTs and maintain stable MT fibers

that are attached to kinetochores. The oocytes were immunostained with α-tubulin (to label

MTs), CREST (to label kinetochores), stained with DAPI (to label DNA), and imaged using

confocal microscopy. K-MT attachments were scored as normal, unattached, or abnormally

attached (merotelic or syntelic) (Fig. 3A). Nucleus positioning had no effect on the percentage of

unattached kinetochores among all groups (Fig. 3A,B). Importantly, oocytes with peripherally

located nucleus displayed ~ four-fold increase in the percentage of incorrect K-MT attachments,

compared to oocytes with a centrally located nucleus (Fig. 3A,C). Moreover, the percentages of

peripheral (37.93 ± 9.17%) and intermediate (39.13 ± 10.41%) oocytes having two or more

abnormally attached kinetochores were significantly higher than those in the central group (7.14

± 4.02%) (Fig. 3D). These results suggest that improper K-MT attachments are the likely cause,

at least in part, of increased aneuploidy in oocytes with a peripherally located nucleus. These

results led us to hypothesize that initial spindle formation and its positioning at the oocyte center

serves as mechanism to ensure avoidance of cortical influence on the establishment of K-MT

attachments, prior to timely spindle migration to the cortex.

Incidence of aneuploidy directly associated with the proximity of early spindle positioning to

the cortex

To confirm that the cortical influence during spindle formation is the cause, at least partially, of

the high rate of aneuploidy in peripheral GV oocytes, we tracked the movement of the nucleus

using live time-lapse microscopy. Oocytes with a peripherally located nucleus demonstrated one

of three types of movement: oocytes that were able to achieve complete relocation of the nucleus

to the center prior to NEBD were termed 'corrected', oocytes that demonstrated central

movement prior to NEBD but did not achieve full relocation were termed 'migratory', and

oocytes that failed to make a relocation attempt and remained at the cortex through NEBD were

termed 'peripheral' (Fig. 4A,B and supplementary movies 2-4). Strikingly, oocytes that were

able to successfully relocate their nucleus to the center showed relatively lower rates of

aneuploidy (comparable to that of oocytes beginning with a central nucleus) when compared to

oocytes that failed nucleus relocation and underwent NEBD at the periphery of the oocyte (Fig.

4C).

It is technically feasible to shift the position of the spindle within the oocyte without noticeable

adverse effects [29,30]. Therefore, to further confirm the relationship between aneuploidy and the

proximity of early spindle positioning to the cortex, we relocated the spindle prematurely to the

cortex through mild centrifugation, a technique proven successful in murine oocytes (Fig. 4D) [31].

Central GV oocytes were matured *in vitro* for 4 h, until prometaphase (the time prior to

establishing correct K-MT attachments [32]). These prometaphase oocytes were centrifuged,

examined live under the microscope and resorted into two groups according to their spindle

positioning: those maintaining central positioning (control central) and those with induced

peripheral positioning (induced peripheral) prior to *in vitro* maturation for an additional 13 h

(Met II). Notably, oocytes with induced peripheral positioning showed significantly higher rates

of aneuploidy (58.33%) when compared to oocytes maintaining central spindle positioning after

centrifugation (16.67%, Fig. 4E); thus suggesting that peripheral GV oocytes are not inherently

erroneous, but instead are subjected to increased levels of cortical influence that is detrimental to

the fidelity of chromosome segregation.

Inhibition of the Cdc42 cortical gradient permits normal formation of K-MT attachment in

oocytes with peripheral spindle formation

Recent study showed that Cdc42 signals coming from the oocyte cortex can influence K-MT

attachments through tyrosination of MTs close to the cortex [27]. In mitotic cells, tyrosinated MTs

are associated with abnormal K-MT attachments [33]. From these findings, we hypothesized that

premature exposure to increased cortical Cdc42 signals in oocytes with a peripherally positioned

spindle during prometaphase could hinder the oocyte's ability to establish and/or stabilize correct

K-MT attachments. To this end, oocytes were matured *in vitro* for 5 h. Based on spindle position

using live imaging, oocytes were sorted into two groups: central and peripheral oocytes. At this

point, the treatment groups were supplemented with a Cdc42 inhibitor (ML141), whereas control

oocytes were supplemented with dimethyl sulfoxide (DMSO) for an additional 2 h (Fig. 4F). Met

I oocytes were exposed to a 6 minutes cold shock followed by the assessment of K-MT

attachments using immunocytochemistry and confocal microscopy. K-MT attachments were

scored as normal, unattached, or abnormally attached (merotelic or syntelic). The number of

unattached kinetochores did not differ significantly among groups (Fig. 4G). Expectedly, oocytes

with peripherally positioned spindles (during prometaphase I) exhibited a significant increase in

the percentage of abnormal K-MT attachments, compared to those with a centrally positioned

spindle. Importantly, inhibition of cortical Cdc42 signals using ML141 rescued, at least in part,

the percentage of abnormal K-MT attachments in non-treated oocytes with a peripherally

positioned spindle (Fig. 4H). These results suggest that avoidance of the cortical cdc42 signals

during prometaphase I is critical in protecting the oocyte's ability to establish correct K-MT

attachments and further undergo faithful chromosome segregation (Fig. 5).

DISCUSSION

Most errors contributing to aneuploid embryos occur in female gametes, specifically within the

first meiotic division [5,6]; therefore, it is crucial to understand the tightly regulated mechanisms

that ensure faithful chromosome segregation. Our findings suggest that central nucleus

positioning serves as a preemptive insurance mechanism to ensure that the spindle forms at the

center of the oocyte and avoids deleterious cortical Cdc42 signals. These cortical signals are

capable of affecting the fidelity of proper K-MT attachment formation. Indeed, when the spindle

is prematurely exposed to cortical signals, due to peripheral nucleus positioning or premature

induction of cortical positioning of the spindle, the oocyte experiences significantly increased

levels of abnormal K-MT attachment and aneuploidy.

Abnormal K-MT attachments during Met I is a major cause of aneuploidy in mammalian oocytes

[22,34,35]. Our results showed that peripheral GV oocytes that displayed a correction mechanism by

relocating their nuclei to the center (before spindle assembly) displayed a significant decrease in

the percentage of abnormal K-MT attachments when compared to uncorrected oocytes,

suggesting that the cortical signals may have an influence on the ability of the oocyte to establish

correct K-MT attachments. In mitotic cells, tyrosinated MTs form early K-MT attachments, but

correctly established attachments become detyrosinated and stabilize as spindle tension

increases. Incorrect attachments remain tyrosinated (unstable) and are thus more likely to

depolymerize [33].In addition, recent evidence suggests that cortical Cdc42 signals have the

capability to influence K-MT attachments and centromeric selection in meiotic oocytes through

spindle asymmetry and MT tyrosination. Consistently, we found that inhibition of Cdc42 largely

rescued increased abnormal K-MT attachments in peripherally assembled and positioned spindle.

Taken together, our results suggest that primary spindle formation and positioning at the center

of the oocyte may serve as an essential protection mechanism against cortical signals capable of

influencing improper K-MT attachments.

Nucleus positioning in meiotic oocytes is often studied only *in vitro* or at a single time point *in*

vivo and is thus not well understood. There has been a long-standing debate regarding the

position of the nucleus in mammalian oocytes. For example, a previous study found that the

majority of prophase I-arrested oocytes have a peripherally located nucleus, and this cortical

positioning of the nucleus is likely influenced by gap junctions and increased cumulus cell

contact [36]. On the other hand, other studies demonstrated that the majority of prophase I-arrested

oocytes have a centrally positioned nucleus [8,37,38]. Interestingly, in the first study [36], the oocytes

were collected from mice aged 21 days, whereas in the other studies [8,37,38], the oocytes were

collected from mice aged more than 6-8-week old. Our results– that the majority of oocytes

collected from 2-3-week old mice exhibited a peripherally located nucleus, whereas the majority

of oocytes from 6-8-week old mice exhibited a centrally located nucleus– clearly explain these

previous findings and advance our understanding of an important phenomenon in mammalian

oocytes.

Although the nucleus shifts from a peripheral position in prepubertal oocytes towards a central

position in sexually mature mouse oocytes, this is likely not the only shift during the

reproductive life. Indeed, oocytes from aged mice have a majority of peripherally positioned

nuclei [8]. Why and how nucleus positioning is a dynamic process during oocyte development

remains unknown. It is unlikely that age-dependent nucleus positioning is due to the difference

in the response to gonadotropins because FSHβ knockout mice showed similar proportions of

central *vs*. peripheral GV oocytes to those of WT mice (Kincade, J., Almamun, Marshall, K., M.,

Rivera, R. and Balboula, A., unpublished data). During oocyte meiosis, prophase I-arrested

oocytes have a cytoplasmic mesh of F-actin-coated vesicles that experience increased movement

near the cortex and decreased movement near the center. Accordingly, the gradient of F-actin

creates a gentle force that encourages the nucleus to position at the center of the cell through

active diffusion [9,10]. Analyzing the cytoskeletal changes within the oocyte during its development

and growth may reveal more insight into the underlying mechanisms behind nucleus positioning

throughout oogenesis.

Current *in vitro* fertilization (IVF) technologies maintain much room for improvement, as only

12.7% of naturally ovulated IVF cycles in humans produce a pregnancy and only 8.8% of those

pregnancies result in live births [39]. Although technologies and protocols have advanced since the

first IVF birth in 1978, the success rate in producing a live birth in women under the age of 35

through IVF remains less than 50% [40]. To date, there are no definitive morphological markers of

predictive oocyte quality and successful IVF that are particularly non-invasive [41]. Gametes from

human females are notoriously associated with high rates of aneuploidy [5,42]. In addition, human

oocytes with a peripherally located nucleus are associated with poor maturation rates compared

to those with a centrally located nucleus [37]. Therefore, understanding the association between

peripherally located nucleus and the incidence of aneuploidy may allow for the selection of

oocytes predisposed to correct segregation of chromosomes. Additionally, arresting the cell and

enabling the nucleus to relocate to central positioning may be a promising approach to increase

the overall efficacy of IVF practices, allowing for increases in blastocyst rate, pregnancy rate,

and live birth rate in all species.

MATERIAL AND METHODS

Ethics

All laboratory animals were managed, and experiments conducted, in compliance with the

University of Missouri (Animal Care Quality Assurance Reference Number, 9695).

Mouse strains

Unless otherwise specified, female CF-1 WT mice were used. *Cep192-eGfp* reporter mice were

generated by integrating a construct harboring the EGFP reporter gene into CF-1 mouse genome

via CRISPR/Cas9-mediated homology-directed repair. The EGFP reporter was fused at the C-

terminus of the endogenous mouse Cep192. Homozygous breeding pairs of *Cep192-eGfp*

reporter mice were used to maintain the colony.

Oocyte collection and in vitro maturation (IVM)

Full-grown GV oocytes were collected from CF-1 female mice (not primed by pregnant mare

serum gonadotropin) aged 6-8 weeks. Oocyte collection and culture were carried out as

previously described [22,43]. Cumulus oocyte complexes (COCs) were collected and cultured in

bicarbonate-free minimal essential medium (MEM) supplemented with 3mg/ml

polyvinylpyrrolidone (PVP) and 25 mM Hepes (pH 7.3) under mineral oil (MilliporeSigma, St.

Louis, MO, USA # M4659, # P2307, # H3784, # M8410). Oocytes were manually denuded and

sorted into 3 groups: Central GV, Intermediate GV, Peripheral GV oocytes. Careful rolling of the

oocyte ensured correct classification. Oocytes were then transferred to Chatot, Ziomek, and

Bavister (CZB) medium [44] supplemented with 1µM glutamine (MilliporeSigma, # G8549) under

mineral oil and matured at 37˚C with humidified 5% CO_2 in an incubator for either 7, 14, or 16 h.

Nocodazole (Sigma #M1404), monastrol (Sigma #M8515), ML141 (Sigma #SML0407)and

ZM447439 (Tocris #2458) were dissolved in DMSO and added to CZB culture medium at a final

concentration of 5 µm, 100 µM, 500 nM and 10 µm, respectively. SiR-tubulin and SiR-DNA

were added to the maturation medium (at a final concentration of 100 nM) during live imaging to

label MTs and DNA, respectively.

Immunocytochemistry

After maturation, oocytes were fixed in a freshly prepared solution of 3.7% paraformaldehyde

(MilliporeSigma, # P6148) dissolved in phosphate buffer saline (PBS). Oocytes were incubated

in permeabilization solution (0.1% Triton X-100 in PBS) for 20 minutes and blocking solution

(0.3% BSA and 0.01% Tween-20 in PBS) for 20 minutes before staining. Oocytes were

incubated at room temperature for one h with the primary antibody before undergoing 3

consecutive washes in blocking solution for 9 minutes each. Oocytes were then incubated at

room temperature for one h in the secondary antibody solution. Following another 3 consecutive

washes in blocking solution at 9 minutes each, oocytes were mounted on glass slides using

Vectashield containing 4',6-Diamidino-2-Phenylindole, Dihydrochloride (DAPI; Vector

Laboratories, Burlingame, CA, USA) in order to stain the DNA. Fluorescence was observed

using a 40X oil objective using Leica TCS SP8 confocal microscope or Leica DMI8 fluorescence

microscope and oocytes were imaged using 3 µm Z-intervals. Oocytes were analyzed using NIH

ImageJ software (National Institute of Health, Bethesda, MD USA).

The following primary antibodies were used in immunofluorescence: conjugated α-tubulin-AlexaFluor 488 (Life Technologies # 322 588; 1:75), CREST autoimmune serum (Antibodies Incorporated # 15-234; 1:25).

In situ chromosome counting

Oocytes matured *in vitro* for 14 h in CZB medium supplemented with L-glutamine under mineral oil were transferred to CZB containing 100 µm monastrol (MilliporeSigma, # M8515) and incubated an extra 2 h; monastrol is an Eg5-kinesin inhibitor which induces monopolar spindle formation and thus results in a rosette of chromosome distribution [21,22]. After treatment with monastrol, oocytes at Met II stage were fixed in a freshly prepared solution of 2.5% paraformaldehyde (MilliporeSigma, # P6148) and stained with CREST autoimmune serum (Antibodies Incorporated # 15-234; 1:25) to label the kinetochores. DAPI was used to label the DNA. Fluorescence was observed using a 40X oil objective using a Leica DMI8 microscope and oocytes were imaged using 0.5 µm Z-intervals in order to observe all kinetochores. Oocytes were analyzed individually and were scored either as euploid (containing 40 kinetochores) or as aneuploid (containing ± 40 kinetochores).

Assessment of kinetochore-microtubule attachment

Cells were matured *in vitro* for 7 h in CZB medium supplemented with L-glutamine under mineral oil. The oocytes were placed on ice for 6 minutes in a 96 well dish containing chilled MEM. Exposing the oocytes to cold shock depolymerizes unattached MTs; however, when MTs have established end-on attachment to a kinetochore, it becomes cold stable [45]. Oocytes were fixed using a freshly prepared 2.5% paraformaldehyde solution and immunostained with anti-

human CREST (to label kinetochores) and α-tubulin (to label MTs) antibodies. DAPI was used

to label DNA. Cells were imaged using a Leica TCS SP8 confocal microscope and a x63 oil

objective, taking images at 0.5 µm Z-intervals. Kinetochores were analyzed for unattached,

normal, or abnormal attachments (syntelic - both kinetochores attached to one side of the spindle

or merotelic - a kinetochore maintaining attachment to both sides of the spindle). Oocytes were

analyzed using NIH ImageJ Software.

Time-lapse microscopy

Full-grown oocytes were imaged using a 40X oil objective on Leica TCS SP8 confocal

microscope or Leica DMI8 microscope within a microenvironmental chamber maintaining an

atmosphere of 5% CO_2 and temperature of 37˚C in humidified air. Imaging acquisition began

before NEBD. Images were captured at 5 µm Z-intervals. Image sequences were analyzed using

NIH ImageJ Software.

Ovary collection and histological evaluation

Ovaries from CF-1 mice of varying ages (2 weeks, 3 weeks, 5 weeks, and 8 weeks) were

collected, submerged, and stored in buffered zinc formalin overnight. Tissues were then moved

to 70% ethanol for storage until undergoing a 10-h protocol using a Leica ASP300S enclosed

tissue processor. After processing, tissues were embedded in paraffin wax and cooled fully using

the Leica EG1150 C/H pair. Embedded tissues were sectioned at 8 µm thickness using a Leica

Histocore AutoCut microtome. Representative samples were taken for each age by taking 3

consecutive sections periodically (every 4, 6, 8, or 10 sections respectively). Tissues were

stained with Hematoxylin and Eosin (Abcam, # 245880) before being mounted with permount

resin (Thermo Fisher Scientific, Inc. # SP15). Oocytes and their corresponding nucleus

positioning were examined using 10X or 40X objective on a Leica DMI8 microscope.

Statistical analysis

Tests used to evaluate the statistical significance of the findings reported include One-way

ANOVA and Student t-test using GraphPad Prism. The Tukey post hoc test was utilized to

determine statistical significance between groups. P values of < 0.05 were considered significant.

All data are displayed as means $\pm$ SEM.

Declaration of interest

All authors declare that there are no conflicts of interest that may prejudice or bias the research

reported.

Funding

This research was supported by laboratory start-up funding from the University of Missouri to

AZB.

Acknowledgements

The authors would like to thank all members of the Balboula lab at the University of Missouri,

Columbia for dedicated help and encouragement. The authors would like to thank Rocio Rivera,

Heide Schatten for critical reading of the manuscript and valuable discussion, and Yuksel Agca,

Pramod Dhakal, Thomas Spencer, the Animal Modeling Core and the Molecular Cytology Core

for equipment usage and valuable discussion.

Author Contributions

AZB conceived the project. AZB and JK designed and analyzed experiments. JK performed experiments. AZB and JK wrote the manuscript.

Figure Legends

Figure 1: Nucleus positioning is a dynamic process throughout the development

(A) Full-grown GV oocytes were collected, imaged and classified into one of three groups: central GV, intermediate GV, or peripheral GV according to the distance measured from the nucleus to the nearest cortex. The scale bar represents 50 µm. Quantification of the average distance from the nucleus to the nearest cortex within groups (right graph). One-way ANOVA and Tukey's post hoc test were performed to analyze the data. (B) Representative images of oocytes with a central, intermediate, and peripherally located nucleus. The scale bars represent 100 µm. Quantification of the contribution of each group to all oocytes collected (right graph). One-way ANOVA and Tukey's post hoc test were performed to analyze the data. (C) Histological analysis of nucleus positioning *in vivo* at different ages. Ovaries from 2, 3, 5, and 8 weeks old mice were collected, fixed, and stained with Hematoxylin and Eosin. Sections were observed through 10X and 40X objective magnification and classified into one of the three positioning groups: central, intermediate, peripheral. The scale bar represents 100 µm. (D) Representative images of nucleus positioning in 3 and 8 weeks old mice. The white box emphasizes the location of the magnified image within the ovary. The scale bar represents 100 µm. (E) Quantification of the proportion of oocytes sorted according to their nucleus positioning (central, intermediate, peripheral) in 2, 3, 5, and 8 week old mice. Two-way ANOVA and Sidak's multiple comparison test were performed to determine if the groups differed

significantly. Data are displayed as mean ± SEM. Values with asterisks vary significantly, * P

<0.05, ** P <0.01, *** P <0.001, **** P <0.0001. The total number of analyzed oocytes (from 3

independent replicates) is specified above each graph.

Figure 2: Oocytes with a peripherally located nucleus have higher rates of chromosome

misalignment and aneuploidy.

Full-grown prophase I-arrested oocytes were collected and sorted into central, intermediate and

peripheral groups based on nucleus positioning and imaged live using time-lapse microscopy

during *in vitro* maturation. (A) Quantification of the percentage of oocytes that underwent

NEBD. (B) Quantification of the average time of NEBD. (C) Quantification of the percentage of

first polar body extrusion (1st PBE). (D) Quantification of the average time of PBE calculated

from oocytes successfully extruded the PB. (E-J) Full-grown prophase I oocytes were collected

and sorted into central, intermediate and peripheral groups based on nucleus positioning and *in*

vitro matured for 7 h (Met I) or 16 h (Met II. Oocytes were fixed and immunostained with α-

tubulin to label the microtubules (the spindle). DNA was labeled by DAPI. (E) Representative

confocal images of Met I oocytes. (F) Quantification of the percentage of chromosome

misalignment in Met I oocytes. (G) Quantification of the percentage of abnormal spindle

morphology in Met I oocytes. (H) Representative confocal images of Met II oocytes. (I)

Quantification of the percentage of chromosome misalignment in Met II oocytes. (J)

Quantification of the percentage of abnormal spindle morphology in Met II oocytes. (K) Full-

grown prophase I oocytes were collected and sorted into central, intermediate and peripheral

groups based on nucleus positioning and *in vitro* matured for 14 h (Met II). Oocytes were treated

with monastrol for 2 h, fixed and immunostained with CREST antibody to label kinetochores.

DNA was labeled by DAPI. Oocytes were analyzed individually and scored either as euploid

(containing 40 kinetochores) or as aneuploid (containing ± 40 kinetochores). Shown are

representative images. (L) Quantification of the percentage of aneuploid eggs. The scale bar

represents 10 μm. One-way ANOVA and Tukey's post hoc tests were performed to determine

statistical significance among groups. Data are displayed as mean ± SEM. Values with asterisks

vary significantly, * P <0.05, *** P <0.001, **** P <0.0001. The total number of analyzed

oocytes (from at least 3 independent replicates) is specified above each graph.

Figure 3: Oocytes with peripherally located nucleus experience higher levels of incorrect

kinetochore-microtubule attachments

(A) Representative images of kinetochore-microtubule (K-MT) attachments in oocytes with a

central, intermediate, or peripherally located nucleus. Panels on the right (from top to bottom)

display normal K-MT attachment, unattached kinetochores and abnormal K-MT attachment. (B)

Quantification of the percentage of unattached kinetochores and (C) abnormal K-MT

attachments. (D) Quantification of the oocytes with 2 or more incorrect K-MT attachments. One-

way ANOVA and Tukey's post hoc test were performed to analyze the data. Data are displayed

as mean ± SEM. Values with asterisks vary significantly, * P <0.05, ** P <0.01, **** P <0.0001.

The total number of analyzed oocytes (from 4 independent replicates) is specified above each

graph.

Figure 4: Incidence of incorrect K-MT attachments and aneuploidy associate with spindle

proximity to the cortex

(A) Schematic depicting the determination of corrected, migratory, and peripheral oocytes. (B) Representative images of the nucleus movement prior to NEBD. Images were taken every 30 minutes (Z-interval of 5 μm). The black circle denotes the primary nucleus positioning, while the green circle denotes the final nucleus position prior to NEBD. (C) Quantification of the incidence of aneuploidy in metaphase II oocytes from the oocytes categorized in B. One-way ANOVA and Tukey's post hoc test were performed to determine statistical significance. (D) Schematic depicting the induction of peripheral spindle positioning by mild centrifugation in oocytes collected with a centrally positioned nucleus. (E) Quantification of the incidence of aneuploidy in oocytes with induced peripherally positioned nucleus and centrally positioned nucleus after centrifugation. Student t-test was performed to determine statistical significance. Data are displayed as mean ± SEM. Values with asterisks vary significantly, * P <0.05. (F) Schematic depicting the treatment of collected oocytes with ML141, a Cdc42 inhibitor, in order to inhibit the cortical influence upon kinetochore-microtubule (K-MT) attachment formation. (G,H) Quantification of the average incidence of unattached kinetochores and abnormal K-MT attachment, respectively. One-way ANOVA and Tukey's post hoc test were performed to determine statistical significance. Data are displayed as mean ± SEM. Values with asterisks vary significantly, * P <0.05, ** P <0.01. The total number of analyzed oocytes (from 3 independent replicates) is specified above each graph.

Figure 5: A summary schematic depicting the deleterious influence of cortical Cdc42 upon proper K-MT attachments in oocytes with a peripherally positioned spindle that is at least partially contributory to the development of aneuploid gametes. The lower section of the

schematic demonstrates the ability of the oocyte to establish correct K-MT attachments in

oocytes with a centrally positioned spindle and thus avoid chromosome mis-segregation.

Supplementary Figure 1: Full-grown prophase I-arrested oocytes were collected from *Cep192-*

eGfp reporter mice, sorted into central (A) and peripheral (B) groups based on the positioning of

their nucleus and *in vitro* matured in CZB medium containing SiR-tubulin (to label microtubules,

magenta). Spindle poles were labeled by CEP192-EGFP (MT organizing centers, green).

Oocytes were imaged live using time-lapse confocal microscopy. Shown are representative

fluorescence and bright-field (lower panels) image sequence depicting the location of NEBD and

consequent spindle formation. Images were taken every 1 h (Z-projection of 13 sections every 5

μm). The scale bar represents 50 μm.

Supplementary Figure 2: (A) Representative bright-field images of time-lapse microscopy of

the movement of a peripherally positioned nucleus during prophase I arrest in oocytes incubated

with milrinone. Images were taken every 10 minutes using a Z-interval of 5 μm. The black circle

identifies the initial positioning of the nucleus, while the green circle denotes the final

positioning of the nucleus immediately prior to NEBD. (B) Quantification of the percentage of

peripheral and intermediate GV oocytes achieving full relocation of the nucleus to a central

position within 3 h of incubation with milrinone.

Supplementary Figure 3: Full-grown prophase I-arrested oocytes were collected and sorted into

central, intermediate, and peripheral groups based on the positioning of their nucleus and imaged

live using time-lapse microscopy during *in vitro* maturation. (A) Quantification of the change in

the distance of the nucleus to the nearest cortex from the initial position of the nucleus at the

prophase I-arrested stage to the final destination of the nucleus immediately prior to NEBD. (B)

Representative fluorescence and bright-filed (lower panels) image sequence depicting

chromosome alignment (labeled by SiR-DNA, red) in oocytes with a central (upper) and

peripheral (lower) nucleus. Images were taken every 45 minutes using a Z-interval of 7 μm.

Scale bars represent 50 μm. (C) Quantification of the distance of chromosomes at the time of

alignment to the nearest cortex. (D) Quantification of the time of chromosome alignment. One-

way ANOVA and Tukey's post hoc test were performed to determine statistical significance.

Data are displayed as mean ± SEM. Values with asterisks vary significantly, ** $P < 0.01$, *** P

<0.001. The total number of analyzed oocytes (from 3 independent replicates) is specified above

each graph.

Supplementary Figure 4: (A) Full-grown prophase I-arrested oocytes were collected, sorted

(into central, intermediate, and peripheral groups based on the positioning of their nucleus) and

in vitro matured in the presence of nocodazole. Randomly selected oocytes were *in vitro* matured

in the presence of nocodazole and ZM447439 (positive control group). The percentage of first

polar body extrusion (PBE) was assessed at 16 h (Met II). One-way ANOVA and Tukey's post

hoc test were performed to determine statistical significance. Data are displayed as mean ± SEM.

Values with asterisks vary significantly, **** $P < 0.0001$. The total number of analyzed oocytes

(from 4 independent replicates) is specified above the graph.

Supplementary Movie 1: Time-lapse microscopy of germinal vesicle (GV) oocytes cultured in

milrinone-containing medium to maintain prophase I arrest. DIC images were captured every 10

minutes at Z-intervals of 5 µm. The scale bar represents 50 µm.

Supplementary Movie 2: Time-lapse microscopy of peripheral germinal vesicle (GV) oocyte

showing corrected nucleus relocation during *in vitro* maturation (Same oocyte used in the top

panels of Fig. 4B). DIC images were captured every 10 minutes at Z-intervals of 5 µm. The scale

bar represents 50 µm.

Supplementary Movie 3: Time-lapse microscopy of peripheral germinal vesicle (GV) oocyte

showing migratory nucleus relocation during *in vitro* maturation (Same oocyte used in the

middle panels of Fig. 4B). DIC images were captured every 10 minutes at Z-intervals of 5 µm.

The scale bar represents 50 µm.

Supplementary Movie 4: Time-lapse microscopy of peripheral germinal vesicle (GV) oocyte

that failed to relocate its nucleus during *in vitro* maturation (Same oocyte used in the lower

panels of Fig. 4B). DIC images were captured every 10 minutes at Z-intervals of 5 µm. The scale

bar represents 50 µm.

REFERENCES

Sun, H. *et al.* Global, regional, and national prevalence and disability-adjusted life-years for infertility in 195 countries and territories, 1990-2017: results from a global burden of disease study, 2017. *Aging (Albany NY)* **11**, 10952-10991, doi:10.18632/aging.102497 (2019).

Jacobs, P. A., Baikie, A. G., Court Brown, W. M. & Strong, J. A. The somatic chromosomes in mongolism. *Lancet* **1**, 710, doi:10.1016/s0140-6736(59)91892-6 (1959).

Lejeune, J., Turpin, R. & Gautier, M. [Mongolism; a chromosomal disease (trisomy)]. *Bull Acad Natl Med* **143**, 256-265 (1959).

Gruhn, J. R. *et al.* Chromosome errors in human eggs shape natural fertility over reproductive life span. *Science* **365**, 1466-1469, doi:10.1126/science.aav7321 (2019).

Hassold, T., Hall, H. & Hunt, P. The origin of human aneuploidy: where we have been, where we are going. *Hum Mol Genet* **16 Spec No. 2**, R203-208, doi:10.1093/hmg/ddm243 (2007).

Hall, H. E. *et al.* The origin of trisomy 22: evidence for acrocentric chromosome-specific patterns of nondisjunction. *Am J Med Genet A* **143A**, 2249-2255, doi:10.1002/ajmg.a.31918 (2007).

Gilbert, S. in *Developmental Biology* Vol. 6th edition   (Sinauer Associates, Inc, 2000).

Brunet, S. & Maro, B. Germinal vesicle position and meiotic maturation in mouse oocyte. *Reproduction* **133**, 1069-1072, doi:10.1530/REP-07-0036 (2007).

Almonacid, M. *et al.* Active diffusion positions the nucleus in mouse oocytes. *Nat Cell Biol* **17**, 470-479, doi:10.1038/ncb3131 (2015).

Almonacid, M., Terret, M. E. & Verlhac, M. H. Control of nucleus positioning in mouse oocytes. *Semin Cell Dev Biol* **82**, 34-40, doi:10.1016/j.semcdb.2017.08.010 (2018).

Colin, A. *et al.* Active diffusion in oocytes nonspecifically centers large objects during prophase I and meiosis I. *J Cell Biol* **219**, doi:10.1083/jcb.201908195 (2020).

Schuh, M. & Ellenberg, J. Self-organization of MTOCs replaces centrosome function during acentrosomal spindle assembly in live mouse oocytes. *Cell* **130**, 484-498, doi:10.1016/j.cell.2007.06.025 (2007).

Clift, D. & Schuh, M. A three-step MTOC fragmentation mechanism facilitates bipolar spindle assembly in mouse oocytes. *Nat Commun* **6**, 7217, doi:10.1038/ncomms8217 (2015).

Balboula, A. Z. *et al.* Haspin kinase regulates microtubule-organizing center clustering and stability through Aurora kinase C in mouse oocytes. *J Cell Sci* **129**, 3648-3660, doi:10.1242/jcs.189340 (2016).

Luksza, M., Queguigner, I., Verlhac, M. H. & Brunet, S. Rebuilding MTOCs upon centriole loss during mouse oogenesis. *Dev Biol* **382**, 48-56, doi:10.1016/j.ydbio.2013.07.029 (2013).

McCarthy, E. K. & Goldstein, B. Asymmetric spindle positioning. *Curr Opin Cell Biol* **18**, 79-85, doi:10.1016/j.ceb.2005.12.006 (2006).

Longo, F. J. & Chen, D. Y. Development of cortical polarity in mouse eggs: involvement of the meiotic apparatus. *Dev Biol* **107**, 382-394 (1985).

Ma, J., Zeng, F., Schultz, R. M. & Tseng, H. Basonuclin: a novel mammalian maternal-effect gene. *Development* **133**, 2053-2062, doi:10.1242/dev.02371 (2006).

19   Menezo, Y. J. Paternal and maternal factors in preimplantation embryogenesis:
interaction with the biochemical environment. *Reprod Biomed Online* **12**, 616-621,
doi:10.1016/s1472-6483(10)61188-1 (2006).
20   Lane, S. I., Yun, Y. & Jones, K. T. Timing of anaphase-promoting complex activation in
mouse oocytes is predicted by microtubule-kinetochore attachment but not by bivalent
alignment or tension. *Development* **139**, 1947-1955, doi:10.1242/dev.077040 (2012).
21   Duncan, F. E., Chiang, T., Schultz, R. M. & Lampson, M. A. Evidence that a defective
spindle assembly checkpoint is not the primary cause of maternal age-associated
aneuploidy in mouse eggs. *Biol Reprod* **81**, 768-776, doi:10.1095/biolreprod.109.077909
(2009).
22   Balboula, A. Z. & Schindler, K. Selective disruption of aurora C kinase reveals distinct
functions from aurora B kinase during meiosis in mouse oocytes. *PLoS Genet* **10**,
e1004194, doi:10.1371/journal.pgen.1004194 (2014).
23   Sanfins, A., Lee, G. Y., Plancha, C. E., Overstrom, E. W. & Albertini, D. F. Distinctions
in meiotic spindle structure and assembly during in vitro and in vivo maturation of mouse
oocytes. *Biol Reprod* **69**, 2059-2067, doi:10.1095/biolreprod.103.020537 (2003).
24   Sanfins, A., Plancha, C. E., Overstrom, E. W. & Albertini, D. F. Meiotic spindle
morphogenesis in in vivo and in vitro matured mouse oocytes: insights into the
relationship between nuclear and cytoplasmic quality. *Hum Reprod* **19**, 2889-2899,
doi:10.1093/humrep/deh528 (2004).
25   Balboula, A. Z., Schindler, K., Kotani, T., Kawahara, M. & Takahashi, M. Vitrification-
induced activation of lysosomal cathepsin B perturbs spindle assembly checkpoint
function in mouse oocytes. *Mol Hum Reprod* **26**, 689-701, doi:10.1093/molehr/gaaa051
(2020).
26   Lane, S. I., Chang, H. Y., Jennings, P. C. & Jones, K. T. The Aurora kinase inhibitor
ZM447439 accelerates first meiosis in mouse oocytes by overriding the spindle assembly
checkpoint. *Reproduction* **140**, 521-530, doi:10.1530/REP-10-0223 (2010).
27   Akera, T. *et al.* Spindle asymmetry drives non-Mendelian chromosome segregation.
Science **358**, 668-672, doi:10.1126/science.aan0092 (2017).
28   Sebestova, J., Danylevska, A., Novakova, L., Kubelka, M. & Anger, M. Lack of response
to unaligned chromosomes in mammalian female gametes. *Cell Cycle* **11**, 3011-3018,
doi:10.4161/cc.21398 (2012).
29   Schuh, M. & Ellenberg, J. A new model for asymmetric spindle positioning in mouse
oocytes. *Curr Biol* **18**, 1986-1992, doi:10.1016/j.cub.2008.11.022 (2008).
30   Tatham, B. G. *et al.* Centrifugation of bovine oocytes for nuclear micromanipulation and
sperm microinjection. *Hum Reprod* **11**, 1499-1503,
doi:10.1093/oxfordjournals.humrep.a019425 (1996).
31   Metchat, A. *et al.* An actin-dependent spindle position checkpoint ensures the asymmetric
division in mouse oocytes. *Nat Commun* **6**, 7784 (retracted), doi:10.1038/ncomms8784
(2015).
32   Kitajima, T. S., Ohsugi, M. & Ellenberg, J. Complete kinetochore tracking reveals error-
prone homologous chromosome biorientation in mammalian oocytes. *Cell* **146**, 568-581,
doi:10.1016/j.cell.2011.07.031 (2011).
33   Ferreira, L. T. *et al.* alpha-Tubulin detyrosination impairs mitotic error correction by
suppressing MCAK centromeric activity. *J Cell Biol* **219**, doi:10.1083/jcb.201910064
(2020).

34 Orr, B., Godek, K. M. & Compton, D. Aneuploidy. *Curr Biol* **25**, R538-542,
doi:10.1016/j.cub.2015.05.010 (2015).
35 Lane, M., Bavister, B. D., Lyons, E. A. & Forest, K. T. Containerless vitrification of
mammalian oocytes and embryos. *Nat Biotechnol* **17**, 1234-1236, doi:10.1038/70795
(1999).
36 Barrett, S. L. & Albertini, D. F. Cumulus cell contact during oocyte maturation in mice
regulates meiotic spindle positioning and enhances developmental competence. *J Assist*
Reprod Genet **27**, 29-39, doi:10.1007/s10815-009-9376-9 (2010).
37 Levi, M., Ghetler, Y., Shulman, A. & Shalgi, R. Morphological and molecular markers
are correlated with maturation-competence of human oocytes. *Hum Reprod* **28**, 2482-
2489, doi:10.1093/humrep/det261 (2013).
38 Almonacid, M. *et al.* Active Fluctuations of the Nuclear Envelope Shape the
Transcriptional Dynamics in Oocytes. *Dev Cell* **51**, 145-157 e110,
doi:10.1016/j.devcel.2019.09.010 (2019).
39 Nargund, G. *et al.* Cumulative conception and live birth rates in natural (unstimulated)
IVF cycles. *Hum Reprod* **16**, 259-262, doi:10.1093/humrep/16.2.259 (2001).
40 Technology, S. f. A. R. (2019).
41 Rienzi, L., Vajta, G. & Ubaldi, F. Predictive value of oocyte morphology in human IVF:
a systematic review of the literature. *Hum Reprod Update* **17**, 34-45,
doi:10.1093/humupd/dmq029 (2011).
42 Hassold, T. & Hunt, P. To err (meiotically) is human: the genesis of human aneuploidy.
Nat Rev Genet **2**, 280-291, doi:10.1038/35066065 (2001).
43 Stein, P. & Schindler, K. Mouse oocyte microinjection, maturation and ploidy
assessment. *J Vis Exp*, doi:10.3791/2851 (2011).
44 Chatot, C. L., Ziomek, C. A., Bavister, B. D., Lewis, J. L. & Torres, I. An improved
culture medium supports development of random-bred 1-cell mouse embryos in vitro. *J*
Reprod Fertil **86**, 679-688, doi:10.1530/jrf.0.0860679 (1989).
45 Rieder, C. L. The structure of the cold-stable kinetochore fiber in metaphase PtK1 cells.
Chromosoma **84**, 145-158, doi:10.1007/BF00293368 (1981).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Pro-Met I   Cdc42   Met II   Aneuploid

Pro-Met I   Cdc42   Met II   Euploid

A

Cep192/SiR-tubulin/Bright-field

B

Cep192/SiR-tubulin/Bright-field

Supplementary Figure 1

A

B

n=90

% Oocytes with nucleus relocation

Intermediate   Peripheral

A

n=61

Peripheral
Intermediate
Central

***

Change in distance to cortex (μm)
from GV to NEBD

B

Central GV oocyte

-00:45   00:00   01:30   03:00   04:30   06:00   07:30   09:00

Peripheral GV oocyte

-00:45   00:00   01:30   03:00   04:30   06:00   07:30   09:00

SiR-DNA/Bright-field

C

n=47

Peripheral
Intermediate
Central

**
***

Distance of chromosomes to nearest
cortex at chromosome alignment (μm)

D

n=47

% oocytes with aligned chromosomes

Central
Intermediate
Peripheral

Time of chromosome alignment (min)

Supplementary Figure 3