Iron-meditated fungal starvation by Lupine rhizosphere-associated and extremotolerant

Streptomyces sp. S29 desferrioxamine production

Scott A. Jarmusch[a,c]*, Diego Lagos-Susaeta[b†], Emtinan Diab[a†], Oriana Salazar[b], Juan A.

Asenjo[b], Rainer Ebel[a], Marcel Jaspars[a]*

[a] Marine Biodiscovery Centre, Department of Chemistry, University of Aberdeen, Old

Aberdeen AB24 3UE, Scotland, U.K.

[b] Centre for Biotechnology and Bioengineering, University of Chile, Beauchef 850,

Santiago, Chile.

[c] Department of Medicinal Chemistry – Pharmacognosy, Uppsala University,

Biomedicinskt Centrum, Uppsala, Sweden 751 23

[†]Authors contributed equally to this work

*Corresponding authors: scott.jarmusch@ilk.uu.se; m.jaspars@abdn.ac.uk

ABSTRACT

Siderophores are iron-chelating compounds that aid iron uptake, one of the key strategies

for microorganisms to carve out ecological niches in microbially diverse environments.

Desferrioxamines are the principal siderophores produced by *Streptomyces* spp. Their

biosynthesis has been well studied and as a consequence, the chemical potential of the

pathway continues to expand. With all of this in mind, our study aimed to explore

extremotolerant and Lupine rhizosphere-derived *Streptomyces* sp. S29 for its potential

antifungal capabilities. Cocultivation of isolate S29 was carried out with *Aspergillus niger*

and *Botrytis cinerea*, both costly fungal phytopathogens in the wine industry. The results

indicate that not only is *Streptomyces* sp. S29 extraordinary at producing hydroxamate

siderophores but uses siderophore production as a means to 'starve' the fungi of iron.

High resolution LC-MS/MS followed by GNPS molecular networking was used to observe

the datasets for desferrioxamines and guided structure elucidation of new

desferrioxamine analogues. Comparing the new chemistry, using tools like molecular

networking and MS2LDA, with the known biosynthesis, we show that the chemical

potential of the desferrioxamine pathway has further room for expansion.

INTRODUCTION

The application of *Streptomyces* secondary metabolites goes beyond drug discovery, as they are one of the most important bacteria for crop protection. Their biotechnological application as rhizosphere or endophytic bacteria is proven to confer protection against soilborne pathogens like fungi, nematodes, and other bacteria.[1,2] They serve as an important factor in 'disease suppressive soils', with several mechanisms of action including synthesis of plant growth regulators[3], antibiotic production[4], secretion of volatile compounds[5], and siderophore production.[6] The last of these is not only a way for bacteria to uptake vital metals from the surrounding soil but also to sequester metals in order to gain a competitive edge against other soilborne organisms.

Siderophores are iron-chelating compounds that aid metal uptake in iron deficient environments.[7] These secondary metabolites are produced by all microbes and four types of siderophore-mediated social interactions have been observed: uptake among clonal cells (sharing), cheating (which can only happen amongst microbes with the same uptake receptor), locking away, and metabolite competition (whichever microbe makes the most cost efficient and effective chelator).[8] Desferrioxamines (DFO) are the most widely observed group of siderophores in *Streptomyces* spp., used principally for Fe^{3+} scavenging[7,9], as well as some other heavy metals.[10–12] Since the vast majority of organisms require Fe^{3+} in order to maintain proper cell wall function[7], as one scavenges for itself, it simultaneously starves other competing organisms, rendering them potentially unviable.

The biosynthetic machinery of desferrioxamines has been well studied[13–15] and as a consequence, the chemical potential of the biosynthetic pathway continues to expand.

The substrate specificity of DesC-like proteins, similar to acyl CoA-dependent

transferases, seems to be the key factor in producing diverse desferrioxamine

scaffolds.[13,15] It has been shown that DesC can accommodate for acetyl, succinyl, and

myristol-CoA, with the latter providing insight into the production of acyl

desferrioxamines.[13] Additionally, aryl desferrioxamines, first isolated from *Micrococcus*

luteus KLE1011[16] and later from *Streptomyces sp.* MA37[17], showed further substrate

flexibility, where the latter communication described a new biosynthetic gene cluster

(BGC) which contained a DesC homolog (LgoC). This enzyme promiscuity fits ideally with

the Screening Hypothesis first proposed by Firn and Jones, where secondary metabolism

stems from metabolic pathways that 'maximize diversity and minimize cost'.[18]

The precursors required to drive siderophore biosynthesis have been

hypothesized to originate from metabolite pools (Figure 1). *Shewanella putrefaciens* has

been used for many of these studies, showing when precursors like putrescine are absent

to drive putrebactin production, lysine-based desferrioxamines are then produced.[19,20]

The last step in the desferrioxamine 'life-cycle' (Figure 1) is the export/import of the DFOs

and ferrioxamines (FOs), respectively. Import of ferrioxamines is mainly determined by

the protein DesE. Its role in the binding and uptake of FOs has been shown in multiple

studies[13,21], but most importantly, this and all other siderophore uptake mechanisms are

specific to their producing organisms, leading to the competition mechanisms described

previously.

Due to the modular nature of desferrioxamines and the presence of peptide bonds

throughout their structure, mass spectrometry-based studies have been utilized

extensively to not only study their biological impact but also for elucidation of new

structures. Previous studies have established standard fragmentation patterns of these

siderophores, even as far as elucidating new structures solely based on mass

spectrometry.[22–26] This standardized fragmentation makes the discovery of siderophores

much easier thanks to metabolomics-based tools like molecular networking. Global

Natural Products Social Molecular Networking (GNPS) is the most widely used platform,

allowing for correlations to be drawn amongst large data sets using MS/MS.[27]

With all of this in mind, our study aimed to explore extremotolerant, rhizosphere-

derived *Streptomyces* sp. S29, for its potential plant protective capabilities. Cocultivation

of S29 with *Aspergillus niger* and *Botrytis cinerea*, both costly fungal phytopathogens in

the wine industry, were carried out in a fashion to allow S29 to dominate the culture, thus

eliciting a chemical response from the addition of each fungus. High resolution LC-MS/MS

followed by GNPS molecular networking was used to analyse the data sets for

desferrioxamines and guided structure elucidation of new desferrioxamines. The results

of the present study indicate that not only is *Streptomyces* sp. S29 extraordinary at

producing hydroxamate siderophores, totalling 16 new analogues and 9 previously

identified ones confirmed through MS/MS structure elucidation, but uses siderophore

production as a means to 'starve' the fungi of iron. When grown in coculture with the fungi,

9 new analogues and 16 previously described ones were identified by MS/MS. Using

MS2LDA to further explore desferrioxamine chemical space, we observed that it is

potentially three times larger than previously anticipated. These results point to

Streptomyces sp. S29 being an important biocontrol agent in the plant rhizosphere.

RESULTS AND DISCUSSION

***Streptomyces* sp. S29 cocultivation and LC-MS of desferrioxamine-rich fractions.**

Preliminary fungal bioassay screening of the lupine rhizosphere *Streptomyces* culture collection showed *Streptomyces* sp. S29 inhibited the growth of *Botrytis cinerea* when grown in coculture on solid medium (Figure S32). In order to induce production of potential antifungal metabolites, the bacteria and fungi were cocultivated in liquid culture in a fashion to allow for *Streptomyces* sp. to be the dominating microbe. Bioassay guided fractionation lead to no antifungal leads, yet, post-fermentation streaks showed no fungal growth (data not shown). Upon LC-MS analysis of the extracts, an abundance of DFOs was observed, leading to the hypothesis that this *Streptomyces* sp. might use a different mechanism than producing antimycotics to fight off the fungal co-cultivate, as seen previously in human bacterial infections.[28] Therefore, *Streptomyces* sp. S29 was also grown in monoculture on iron deficient medium to induce production of DFOs in an attempt to compare siderophore production to the coculture experiments. *Aspergillus niger* and *Botrytis cinerea* were chosen as the fungal phytopathogens due to their established pathology of wine producing grapes [29,30] and the importance of wine as Chile's main agricultural export, a ~ $2 billion (USD) industry.[31]

Extracts and fractions were evaluated for the presence of two DFOs (B and D1) during the fractionation steps, in order to trace the fractions that contained the highest abundance of siderophores. Presence of DFOs was further evaluated at the final stages of separation based on the molecular families generated from molecular networking, utilizing the known DFOs in the GNPS library as anchor points, which further aided structure elucidation. Based on the LC-MS results, the butanol partitions contained the

largest abundance of DFOs, and therefore this data was used to generate a molecular

network using GNPS. The standard settings of the molecular networking workflow were

reduced from 6 to 3 matched fragments in order to identify molecular families which

include closely related known DFOs. A large number of analogues found in this study did

not cluster into molecular families due to the varying nature of CID-MS/MS but were

identified manually using retention time and molecular formula comparisons.

Molecular networking and MS/MS based structure elucidation. The molecular

network generated contained 1208 molecular features of which 165 originated from the

media and instrument blanks included in the workflow. The hydroxamate siderophores

were distributed amongst five subnetworks (Figure 2). Closer examination of each

molecular family showed new analogues linking to known DFOs deposited in the GNPS

library. Subnetwork 1 (a1) contained a number of new acyl DFO D-like analogues that

are first described in this study (Figure 3). Subnetwork 2 (a2) contained various Fe^{3+} and

Al^{3+} complexed FOs (Figure S1). Within this network a new analogue was identified, C_{11}

monounsaturated acyl DFO-B, while its unbound form was not observed in culture (Figure

S24). Subnetworks 3 and 4 (a3 and a4) each contained one known siderophore,

bisucaberin-02 and legonoxamine B, respectively (Figure S2). Subnetwork 5 (a5)

contains one known DFO, legonoxamine A, and one additional new desferrioxamine,

legonoxamine E (*m/z* 537.2924, Figure S9). For the remaining nodes in a5, it was not

possible to assign definite structures elucidated solely based on mass spectrometry

(Figure S3).

MS/MS based structure elucidation was carried out in a similar fashion to studies

conducted previously (Figure 4).[22–25] Early observations of desferrioxamine B and D1

fragmentation showed an initial loss of the N-terminus, corresponding to the loss of either 118.1 or 160.1. Using this as a foundation, structures were either started with a *N*-hydroxycadaverine or a *N*-hydroxy-*N*-acetylcadaverine, respectively. The remainder of structure elucidation was completed stepwise: (1) predicting the molecular formula from the accurate mass and collection of MS/MS data, (2) filling out the structures with the typical desferrioxamine moieties of *N*-hydroxycadaverine and succinate, and (3) taking the remainder of the molecular formula and placing an appropriate acid at the C-terminus that either has precedent from previous studies or could be a potential binding partner with an acyltransferase like coenzyme A. Overall, 25 new DFO analogues were identified in the molecular network and through manual data processing (Table S1). These range from short acyl chain DFO, acyl desferrioxamine D-like (DFO-D) analogues which have not previously been described, aryl DFOs, and long chain dicarboxylate DFOs (Figure 5 and Table S1). Additionally, 25 known hydroxamate siderophores were detected, including known acyl and aryl DFOs, giving further credence to the new structure assignments. Their identification and observed presence in monoculture and/or coculture are present in Table S1.

Acyl DFO production by *Streptomyces* sp. S29. *Streptomyces* sp. S29 was able to produce seven new acyl DFOs similar to those previously described.[23,24] Of these seven, five were short acyl chain derivatives (C_3, C_4, hydroxy-C_4, C_5, and C_6) and two were medium monounsaturated acyl chained derivatives (C_{10} and C_{11}). The placement of the alkene at the position α to the carbonyl is based off previous studies also reporting DFOs containing unsaturated acyl chains.[24,32] We propose henceforth that the previously known 'acyl DFOs' should be renamed 'acyl DFO-B' due to the observation in this study

of analogues that contain a *N*-acetyl-*N*-hydroxycadaverine terminus, similar to

desferrioxamine D. The new acyl DFO-D analogues discovered here also range from

short, medium and long acyl chained derivatives (Table S1). It is likely these acyl DFOs

are branched analogues as previous studies noted that it was unlikely the chains were

fully linear since *Streptomyces* spp. utilise branched starter units in fatty acid

synthesis.[23,24]

Streptomyces coelicolor has shown broad substrate specificity in regard to medium

and long acyl chain incorporation into desferrioxamine structures[13], but not short acyl

chained groups, where *Streptomyces* sp. S29 can facilitate production of short, medium,

and long acyl chained DFO-B analogues. Traxler et al. showed the ability of other

actinobacteria to react to the presence of *S. coelicolor*. through the production of acyl

DFO-B analogues.[23] Interestingly, we see this same phenomenon for the coculture of

Streptomyces sp. S29 and the fungi. Medium and long chained acyl DFO-Bs were almost

exclusively produced in coculture whereas medium and long chained acyl DFO-Ds were

produced under both mono- and coculture conditions. It was hypothesized these long-

chained derivatives might be a way for the uptake of these analogues to bypass the DesE

uptake mechanism, instead they 'hand-off' the Fe^{3+} via membrane bound

ferrioxamines.[13,33]

Aryl DFO production by *Streptomyces* sp. S29. The production of seven new

aryl DFOs related to legonoxamine A and B, which were also observed in this study (Table

S1), shows the biosynthetic flexibility of *Streptomyces* sp. S29. These structures range

from being smaller, potential oligomers like legonoxamine C (*m/z* 351.1912, Figure S6)

to the larger legonoxamine H (*m/z* 665.3893, Figure S20). Two interesting legonoxamine

derivatives include the putrescine-containing legonoxamine G (Figures 5B and S16) and

legonoxamine A glycoside (Figures 5C and S31).

Legonoxamine G contains a *N*-hydroxyputrescine moiety due to the initial loss of

104 Da instead of the typical 118 Da associated to *N*-hydroxycadaverine. Putrebactin is

the archetypal putrescine containing siderophore[34], and as its biosynthetic gene cluster

has been shown to lack genes coding for production of putrescine, it is assumed that

instead, this precursor is sequestered from metabolite pools.[19] Interestingly, researchers

have also found that under putrescine-depleted conditions, *Shewanella putrefaciens*

shifts to producing desferrioxamine B when cadaverine precursors are available.[35] In

desferrioxamine biosynthesis, incorporation of the putrescine precursor has only been

shown once before in desferrioxamine A[36] and now in this study with the discovery of

legonoxamine G.

Legonoxamine A glycoside was observed using the predicted molecular formula

($C_{37}H_{62}N_6O_{13}$) and the loss of a sugar (-162 Da). An additional putative glycosylated DFO

(*m/z* 899.4827), with a predicted molecular formula of $C_{38}H_{70}N_6O_{18}$, was observed in the

mass spectrum but upon selected ion monitoring (SIM), produced insufficient fragment

ions for structural characterisation. Based on molecular formula alone, the structure of

this glycosylated desferrioxamine could be a C_2 acyl DFO-B diglycoside. The production

of glycosylated DFOs has only been reported once before: nocardamin glucuronide,

isolated from *Streptomyces* sp. 80H647.[37] There are only a few reports of microbial

enzymes that produce glucuronides[37], yet, the addition of glucose or other hexoses are

more commonplace in bacterial natural products.[38] We propose legonoxamine A

glycoside undergoes a similar glycosylation mechanism to nocardamin glucuronide,

occurring on a *N*-hydroxyl.

Dicarboxylic acid desferrioxamine analogues. Two new desferrioxamines,

observed only in coculture extracts, with dicarboxylic acid C-termini were also elucidated

(Figure 6A-B). Dodecanedioic DFO-B (Figure 6A and S26) has an *m/z* of 745.5086,

corresponding to the molecular formula of $C_{36}H_{68}N_6O_{10}$. Tridecanedioic DFO-B (Figure

6B and S28) has an *m/z* of 759.5221, corresponding to the molecular formula of

$C_{37}H_{70}N_6O_{10}$. This is the first reported instance of dicarboxylic acids larger than succinate

being incorporated into DFO structures yet there are examples of larger dicarboxylic acids

being bound to CoA, like pimelic acid, therefore we cannot rule the possibility out. Based

on formula predictions and the lack of further fragmentation, the proposed structures are

based on the biosynthesis and the mass spectrometric data obtained. Further

investigation into these dicarboxylic acid products is currently underway.

Ferrioxamines and DFO presence in pure and coculture experiments. A

number of metal complexed ferrioxamines were observed in subnetwork 2 (Figure S1).

Fe^{3+} and Al^{3+} are the only metals complexed with the variety of desferrioxamines present,

similar to studies before (Figure S1).[24,26] The binding efficiencies for Fe^{3+} and Al^{3+} are

similar, therefore there seems to be no overwhelming preference for either.[39] The

importance of iron has already been addressed in regard to bacteria and cell maintenance

but Al^{3+} is toxic and has no direct role in metabolism and seems to be an unintended

biological activity of these metabolites.

The molecular network was also used to evaluate the production of the DFOs

between the two cocultivations with *A. niger* and *B. cinerea* and the *Streptomyces* sp.

S29 monoculture. The results showed that 25 of the 50 were present in the monoculture,

whereas all of the remaining were found in coculture extracts only (Table S1). There was

no significant difference in the siderophores produced under cocultivation with *A. niger* or

B. cinerea, indicating the potential broad response to a fungal microbe in the environment.

This result might point to DFO and siderophore production as a common chemical

response to other microbes, whether it be bacterial or fungal, in close environmental

proximity to the primary organism.

MS2LDA analysis. In order to further evaluate the overall chemical space that

DFOs inhabit in the molecular network, MS2LDA was performed. MS2LDA is a technique

which identifies molecular 'fingerprints' in fragmentation patterns and assigns these into

motifs (13 Mass2Motifs were manually curated). By merging the two networks (GNPS

molecular network and MS2LDA) after motifs were annotated, it is clear that further DFO

chemical space is there to be discovered. Figure S4 represents the overall results of this

merged network where 4A highlights the previously identified nodes (55) and edges (573)

from the original molecular networking job and 4B highlights all of the nodes (147) and

edges (1158) that contain DFO motifs identified through MS2LDA. This tripling of nodes

by inclusion of the MS2LDA workflow shows a clear expansion of DFO chemical space

that other studies up to this date have yet been able to show.

In order to evaluate the accuracy of this analysis, a characterized DOF in this study

(legonoxamine E, Figure 5A) and an unknown DFO observed using MS2LDA (*m/z*

650.37, $[M+2H]^{2+}$) were more closely analysed (Figure S5). The unknown

desferrioxamine is a singleton (Figure S5B) in both the original network and the merged

network, yet, MS2LDA identifies it contains DFO motifs and upon closer inspection and

prediction of its molecular formula ($C_{59}H_{103}N_{12}O_{20}$), it seems to be a novel dodecadentate

DFO, which would point to it being able to bind two moles of Fe^{3+}. Elucidation of its full

structure was not able to be completed with MS/MS data alone, therefore, further studies

are required to characterise its structure.

Streptomyces sp. S29 siderophore biosynthetic gene clusters. DFOs are

biosynthesized by nonribosomal peptide synthetase independent siderophore (NIS)

pathways and the presence of at least one gene encoding a homolog to IucA/IucC has

become recognized as a marker of NIS BGCs.[40,41] DFO biosynthesis is encoded by the

highly conserved _des_ BGC (_desEFABCD_ genes) found in many _Streptomyces_ species [42].

Using FeGenie[43], the identification of iron-related genes revealed 27 siderophore

biosynthesis and 11 transport related genes in the _Streptomyces_ sp. S29 genome. These

siderophore biosynthesis related genes are grouped in five biosynthetic gene clusters

and antiSMASH [44] predictions reveal that three of the five contain at least one gene

homolog to IucA/IucC and the other two correspond to NRPS dependent pathways. In _S._

coelicolor A3(2), transcription of _des_ BGC is repressed by the divalent metal-dependent

regulatory (DmdR) protein DmdR1 and derepressed by iron limitation.[45] BLASTN/P

analysis revealed a putative DmdR repressor in _Streptomyces_ sp. S29 (s29_002782) with

89% nucleotide and 93% amino acid identity to DmdR1 (AJ271797.1 / CAC28070.1).

BLASTN analyses identified eight putative DmdR-binding sites (iron boxes) in

Streptomyces sp. S29 by query of putative iron boxes found upstream in different ORFs

in _S. coelicolor_ A3(2) genome.[13]

The three NIS BGCs (Figure S31A-C) were all independently evaluated against

the two known desferrioxamine BGCs. All three ORFs are present in numerous

Streptomyces spp. yet based on the biosynthetic genes, only those found in regions 7

and 9 of the antiSMASH analysis are potentially related to *des*. Region 7 (Figure S31A)

contains a siderophore BGC in *Streptomyces* sp. S29 with high homology to *desA-D*

genes and to the ferric-siderophore uptake and utilization genes (*desE-F*). This *des*

cluster follows the same structure as many *Streptomyces* spp., including the position of

iron boxes upstream putative *desE* (s29_002291) and *desA* (s29_002293) genes (Fig

S31). DesC of the *des* BGC from *Streptomyces* sp. S29 (s29_2295) exhibits high identity

to DesC from S. coelicolor (72%) and to LgoC from *Streptomyces* sp. MA37 (69%).

Regions 9 and 14 are each siderophore BGCs but the gene clusters and its products are

uncharacterized and unknown. Region 9 (Figure S31B) comprises the essential genes

required to be a NIS BGC related to *des*, like an iron transporter, a IucA/IucC family

siderophore biosynthesis protein and an acyltransferase (s29_4785) with moderate

homology to DesC (30%) and LgoC (32%). Region 14 (Figure S31C) lacks an

acyltransferase which points to this gene cluster not being related to *des*.

Acyltransferases involved in hydroxamate biosynthesis normally exhibit narrow

substrate tolerance (e.g. IucB, which shows a high degree of specificity towards acetyl-

CoA).[13] The array of DFOs produced by *S. coelicolor* (B, E, G1 and amphiphilic) and aryl

DFOs (legonoxamine A-B) produced by *Streptomyces* sp. MA37, has been attributed to

substrate tolerant acyltransferases DesC and LgoC. [13,46] Senges et al. acknowledged in

their evaluation of the *Streptomyces chartreusis* secretome that in iron depleted minimal

medium they also observed additional desferrioxamines produced and that the overall

biosynthetic potential may be much broader than what has been predicted.[25] While the

biosynthesis of the *des* gene cluster in *Streptomyces* sp. S29 needs further studying, the

chemical potential shown in this study highlights its ability for producing a wide range of

DFOs.

In conclusion, the chemistry observed in the *Streptomyces* sp. S29 secretome

shows it potential as an agent to prevent fungal infection, evidenced in the overproduction

of desferrioxamines observed via mass spectrometry. New analogues were observed in

both monoculture and cocultivation studies with fungal phytopathogens, *A. niger* and *B.*

cinerea. *Streptomyces* sp. S29 seems to produce an arsenal of desferrioxamine

analogues using any available precursor materials available whether that be putrescine,

cadaverine, aryl, acyl, or long chained dicarboxylates. The role of bacterial communities

in plant and crop protection is becoming more important due to the growing world

population, with a strong emphasis on improving strategies at preventing phytopathogenic

infection. Overall, the production of 25 new and 25 previously reported siderophores

indicates this *Streptomyces* sp. should be studied further for its plant protective properties

as a biocontrol agent.

EXPERIMENTAL SECTION

Strain collection

Streptomyces sp. S29 was isolated from a soil sample collected from Lupine plant

(*Lupinus oreophilus*) roots originating from a polyextreme enviroment at Quebrada

Nacimiento, Socaire, San Pedro de Atacama, Atacama desert (Northern Chile)

[23°37'06''S;67°50'56''W - 3,646 m.a.s.l]. One aliquot was heated at 100 [°C] for 5 [min]

and directly sprinkled onto Gause #1 and Starch-Casein plates, both complemented with

Nystatin, Cycloheximide and Nalidixic acid. Another aliquot was resuspended in 1⁄4

Ringer solution, heated at 60 [°C] for 10 [min] and dilutions of $10^{-1/2}$ and 10^{-1} were spread

onto agar plates. Plates were incubated at 30 [°C] for ~1 week and colonies were

subcultured on ISP2 plates. Based on morphological characteristics of the colony and

antifungal activity experiments against *B. cinerea* and *A. niger*, 16S rRNA sequence of

S29 isolate was used to identify and confirm its taxonomy using EzBioCloud database (3)

(data not shown).

Phytopathogens

Aspergillus niger and *Botrytis cinerea*, biochemically characterized field samples, were

provided by Laboratorio de microbiología, Facultad de Ciencias naturales, matemática y

medio ambiente, Universidad Tecnológica Metropolitana (UTEM), Santiago, Chile.

Molecular identification of these strains was carried out by sequencing of internal

transcribed spacer (ITS) region amplified by PCR using universal primers ITS1 and ITS4

(supplementary data).

Streptomyces sp. S29 genome sequencing and assembly

Streptomyces sp. S29 total DNA was extracted from liquid culture using DNeasy

UltraClean Microbial Kit (Qiagen). Whole genome sequencing (WGS) of the strain was

performed by an Illumina MiSeq (2 x 300 PE). Nanopore sequencing was performed using

a Rapid Sequencing Kit (SQK-RAD004) and MinIon Flow Cell (R9.4.1). The sequence

was assembled using Canu, polished using Racon, Illumina dataset was mapped onto

the contigs using BWA mem and then corrected using Pilon. The genome was annotated

using NCBI pgap software (6). Raw reads from short-read (Illumina) and long-read

(Nanopore) sequencing were deposited in the NCBI Sequence Read Archive (SRA): SRA

accession number XXXXX. Complete genome sequences were deposited in GenBank

via the NCBI's submission portal: Genome accession number XXXXX.

Cocultivation and extraction

A set of experiments were carried out on these three organisms in order to determine the

effect of the fungal culture being added to the *Streptomyces* sp. (a) *Streptomyces* sp. S29

alone: a single colony was aseptically transferred to 100 mL of iron deficient medium

(ISP2) in order to induce production of siderophores and was fermented for 7 days. (b)

Streptomyces sp. S29 + *Aspergillus niger*: 100 mL of bacterial culture was grown for 5

days. Simultaneously, an independent axenic culture of *A. niger* was grown for 3 days

(10 mL of PDA liquid medium). On the fifth day, the fungal culture was transferred

aseptically to the bacterial culture and allowed to grow an addition 2 days. (c)

Streptomyces sp. S29 + *Botrytis cinerea*: the same procedure as described in experiment

b was followed.

Each liquid fermentation included Diaion HP-20 resin (3 g/50 mL) that was

sterilised with culture medium. After 7 days, the resin was vacuum filtered, subsequently

washed five times with milli-q water and finally macerated exhaustively in methanol (3×).

The methanolic extract was then dried on a rotary evaporator and subsequently

underwent Kupchan partitioning.[47] Finally, the *sec*-butanol fraction underwent C18 solid-

phase extraction (Phenomenex), with each separation yielding 5 fractions of increasing

methanol and these samples were subsequently prepared for LC-MS studies.

The 100% methanol fractions from the *Botrytis* and *Aspergillus* co-culture were

subjected to further separation via reversed phase-HPLC with a Waters Sunfire C18

semi-preparative column (5 µM, 100Å, 250 × 10 mm) wand a solvent system of A–95/5

MeOH/H_2O and B-MeOH. The Agilent 1200 HPLC utilized a solvent gradient of 0% B to

100% B at 2.0 mL/min over 50 min.

Liquid chromatography – mass spectrometry

All runs were analysed using an Agilent 1290 UHPLC coupled to Bruker MAXIS II Q-ToF

mass spectrometer. LC utilizes a Phenomenex Kinetex XB-C_{18} (2.6 mM, 100 × 2.1 mm)

column with a mobile phase of 5% MeCN+0.1% formic acid to 100% MeCN+0.1% formic

acid in 11 min. Bruker MAXIS II has a mass range of *m/z* 100-2000, capillary voltage 4.5

kV, nebulizer gas 5.0 bar, dry gas 12.0 l/min, and dry temperature of 220 °C. MS/MS

experiments were conducted under Auto MS/MS scan mode with a step collision energy

from 80-200%. Samples were prepared at a concentration of 0.3 mg/mL. Raw data files

were analysed using Bruker Daltonics 3.6.

GNPS Molecular Networking and MS2LDA

A molecular network was created using the online workflow (https://ccms-

ucsd.github.io/GNPSDocumentation/) on the GNPS website (http://gnps.ucsd.edu).[27]

The data was filtered by removing all MS/MS fragment ions within ± 17 Da of the precursor

m/z. MS/MS spectra were window filtered by choosing only the top 6 fragment ions in the

± 50 Da window throughout the spectrum. The precursor ion mass tolerance was set to

2.0 Da and a MS/MS fragment ion tolerance of 0.5 Da. A network was then created where

edges were filtered to have a cosine score above 0.7 and more than 3 matched peaks.

Further, edges between two nodes were kept in the network if and only if each of the

nodes appeared in each other's respective top 10 most similar nodes. Finally, the

maximum size of a molecular family was set to 0, and the lowest scoring edges were

removed from molecular families until the molecular family size was below this threshold.

The spectra in the network were then searched against GNPS' spectral libraries. The

library spectra were filtered in the same manner as the input data. All matches kept

between network spectra and library spectra were required to have a score above 0.7

and at least 6 matched peaks.

For documentation on running MS2LDA, the user is referred to the original publication

and guidelines, which are also present at ms2lda.org.[48] All spectral images in the

supplementary information were created using Metabolomics Spectrum Resolver

(https://metabolomics-usi.ucsd.edu/).[49]

All siderophores dereplicated and elucidated in this study can be found in Table

S1. The structure elucidation of each new derivative and its corresponding MS/MS

spectra were generated using Metabolomics Spectrum Resolver are displayed in the

supplementary information (Table S2 and Figures S6-S31). All MS/MS spectra were also

annotated and deposited in the GNPS library, as well as the original datasets (Table S2).

ASSOCIATED CONTENT

Supplementary Information

Tables of known and new desferrioxamine derivatives (Table S1) and their corresponding

CCMS Library IDs and Metabolomics Resolver Images (Table S2), Desferrioxamine

containing subnetworks, new desferrioxamine analogues with annotated structures and

MS^2 spectra, preliminary fungal bioassay image, ITS regions for fungal phytopathogens.

Author Contributions

422 DLS[†] and ED[†] contributed equally to this work.

Acknowledgements

SAJ would like to thank the University of Aberdeen for funding doctoral studies through

and Elphinstone Scholarship. DL would like to thank the Agencia Nacional de

Investigación y Desarrollo [ANID] for funding doctoral studies through 'Beca nacional de

doctorado' Scholarship. ED would like to thank the Ministry of Higher Education and

Scientific Research - Sudan, together with the University of Khartoum, for joint funding of

masters studies. We would like to thank to Valeria Razmilic and Jean Franco Castro for

their valuable advice and work in the setup of Lupine *Streptomyces* culture collection. We

would also like to thank the support team at GNPS and Justin J.J. van der Hooft at

MS2LDA for help with data deposition and for help at any stage of running the relevant

workflows.

References

R. G. Expósito, I. de Bruijn, J. Postma and J. M. Raaijmakers, *Frontiers in Microbiology*, 2017, 8.

O. S. Olanrewaju and O. O. Babalola, *Applied Microbiology and Biotechnology*, 2019, 103, 1179–1188.

Y. Goudjal, O. Toumatia, N. Sabaou, M. Barakate, F. Mathieu and A. Zitouni, *World Journal of Microbiology and Biotechnology*, 2013, **29**, 1821–1829.

O. S. Olanrewaju and O. O. Babalola, *Applied Microbiology and Biotechnology*, 2019, 103, 1179–1188.

S. E. Jones and M. A. Elliot, *Trends in Microbiology*, 2017, 25, 522–531.

V. C. Verma, S. K. Singh and S. Prakash, *Journal of Basic Microbiology*, 2011, **51**, 550–556.

M. Sandy and A. Butler, *Chemical Reviews*, 2009, **109**, 4580–4595.

J. Kramer, Ö. Özkaya and R. Kümmerli, *Nature Reviews Microbiology*, 2020, 18, 152–163.

R. Saha, N. Saha, R. S. Donofrio and L. L. Bestervelt, *Journal of Basic Microbiology*, 2013, **53**, 303–317.

B. Casentini and M. Pettine, *Applied Geochemistry*, 2010, **25**, 1688–1698.

K. E. Jones, K. L. Batchler, C. Zalouk and A. M. Valentine, *Inorganic Chemistry*, 2017, **56**, 1264–1272.

U. Neubauer, G. Furrer and R. Schulin, *European Journal of Soil Science*, 2002, **53**, 45–55.

J. L. Ronan, N. Kadi, S. A. McMahon, J. H. Naismith, L. M. Alkhalaf and G. L. Challis, *Philosophical Transactions of the Royal Society B: Biological Sciences*, , DOI:10.1098/rstb.2017.0068.

T. Schupp, U. Waldmeier and M. Divers, *FEMS Microbiology Letters*, 1987, **42**, 135–139.

F. Barona-Gómez, U. Wong, A. E. Giannakopulos, P. J. Derrick and G. L. Challis, *Journal of the American Chemical Society*, 2004, **126**, 16282–16283.

A. D'Onofrio, J. M. Crawford, E. J. Stewart, K. Witt, E. Gavrish, S. Epstein, J. Clardy and K. Lewis, *Chemistry and Biology*, 2010, **17**, 254–264.

F. Maglangit, M. H. Tong, M. Jaspars, K. Kyeremeh and H. Deng, *Tetrahedron Letters*, 2019, **60**, 75–79.

R. D. Firn and C. G. Jones, *Natural Product Reports*, 2003, 20, 382–391.

S. Rütschlin, S. Gunesch and T. Böttcher, *Cell Chemical Biology*, 2017, **24**, 598-604.e10.

C. Z. Soe, A. A. H. Pakchung and R. Codd, *Chemistry & Biodiversity*, 2012, **9**, 1880–1890.

V. H. Tierrafría, H. E. Ramos-Aboites, G. Gosset and F. Barona-Gómez, *Microbial Biotechnology*, 2011, **4**, 275–285.

T. Pluháček, K. Lemr, D. Ghosh, D. Milde, J. Novák and V. Havlíček, *Mass Spectrometry Reviews*, 2016, **35**, 35–47.

M. F. Traxler, J. D. Watrous, T. Alexandrov, P. C. Dorrestein and R. Kolter, *mBio*, , DOI:10.1128/mBio.00459-13.

24   A. M. Sidebottom, A. R. Johnson, J. A. Karty, D. J. Trader and E. E. Carlson,
ACS Chemical Biology, 2013, **8**, 2009–2016.
25   C. H. R. Senges, A. Al-Dilaimi, D. H. Marchbank, D. Wibberg, A. Winkler, B.
Haltli, M. Nowrousian, J. Kalinowski, R. G. Kerr and J. E. Bandow, *Proceedings of*
the National Academy of Sciences of the United States of America, 2018, **115**,
2490–2495.
26   W. J. Moree, V. v. Phelan, C. H. Wu, N. Bandeira, D. S. Cornett, B. M. Duggan
and P. C. Dorrestein, *Proceedings of the National Academy of Sciences of the*
United States of America, 2012, **109**, 13811–13816.
27   M. Wang, J. J. Carver, V. v. Phelan, L. M. Sanchez, N. Garg, Y. Peng, D. D.
Nguyen, J. Watrous, C. A. Kapono, T. Luzzatto-Knaan, C. Porto, A. Bouslimani,
A. v. Melnik, M. J. Meehan, W. T. Liu, M. Crüsemann, P. D. Boudreau, E.
Esquenazi, M. Sandoval-Calderón, R. D. Kersten, L. A. Pace, R. A. Quinn, K. R.
Duncan, C. C. Hsu, D. J. Floros, R. G. Gavilan, K. Kleigrewe, T. Northen, R. J.
Dutton, D. Parrot, E. E. Carlson, B. Aigle, C. F. Michelsen, L. Jelsbak, C.
Sohlenkamp, P. Pevzner, A. Edlund, J. McLean, J. Piel, B. T. Murphy, L. Gerwick,
C. C. Liaw, Y. L. Yang, H. U. Humpf, M. Maansson, R. A. Keyzers, A. C. Sims, A.
R. Johnson, A. M. Sidebottom, B. E. Sedio, A. Klitgaard, C. B. Larson, C. A. P.
Boya, D. Torres-Mendoza, D. J. Gonzalez, D. B. Silva, L. M. Marques, D. P.
Demarque, E. Pociute, E. C. O'Neill, E. Briand, E. J. N. Helfrich, E. A.
Granatosky, E. Glukhov, F. Ryffel, H. Houson, H. Mohimani, J. J. Kharbush, Y.
Zeng, J. A. Vorholt, K. L. Kurita, P. Charusanti, K. L. McPhail, K. F. Nielsen, L.
Vuong, M. Elfeki, M. F. Traxler, N. Engene, N. Koyama, O. B. Vining, R. Baric, R.
R. Silva, S. J. Mascuch, S. Tomasi, S. Jenkins, V. Macherla, T. Hoffman, V.
Agarwal, P. G. Williams, J. Dai, R. Neupane, J. Gurr, A. M. C. Rodríguez, A.
Lamsa, C. Zhang, K. Dorrestein, B. M. Duggan, J. Almaliti, P. M. Allard, P.
Phapale, L. F. Nothias, T. Alexandrov, M. Litaudon, J. L. Wolfender, J. E. Kyle, T.
O. Metz, T. Peryea, D. T. Nguyen, D. VanLeer, P. Shinn, A. Jadhav, R. Müller, K.
M. Waters, W. Shi, X. Liu, L. Zhang, R. Knight, P. R. Jensen, B. Palsson, K.
Pogliano, R. G. Linington, M. Gutiérrez, N. P. Lopes, W. H. Gerwick, B. S. Moore,
P. C. Dorrestein and N. Bandeira, *Nature Biotechnology*, 2016, 34, 828–837.
28   B. D. Corbin, E. H. Seeley, A. Raab, J. Feldmann, M. R. Miller, V. J. Torres, K.
L. Anderson, B. M. Dattilo, P. M. Dunman, R. Gerads, R. M. Caprioli, W. Nacken,
W. J. Chazin and E. P. Skaar, *Science*, 2008, **319**, 962–965.
29   P. A. G. Elmer and T. Reglinski, *Plant Pathology*, 2006, 55, 155–177.
30   G. Perrone, A. Susca, G. Cozzi, K. Ehrlich, J. Varga, J. C. Frisvad, M. Meijer, P.
Noonim, W. Mahakamchanakul and R. A. Samson, in *Studies in Mycology*,
Centraalbureau voor Schimmelculturen, 2007, vol. 59, pp. 53–66.
31   ODEPA - Oficina de Estudios y Políticas Agrarias, Vinos,
https://www.odepa.gob.cl/rubros/vinos-y-alcoholes, (accessed May 26, 2020).
32   M. Luo, E. A. Fadeev and J. T. Groves, *Journal of the American Chemical*
Society, 2005, **127**, 1726–1736.
33   J. S. Martinez, G. P. Zhang, P. D. Holt, H. T. Jung, C. J. Carrano, M. G.
Haygood and A. Butler, *Science*, 2000, **287**, 1245–1247.

34   K. M. Ledyard, A. Butler, K. M. Ledyard and A. Butler, *Structure of putrebactin,*
524   *a new dihydroxamate siderophore produced by Shewanella putrefaciens*, 1997,
525   vol. 2.
526   35   C. Z. Soe, A. A. H. Pakchung and R. Codd, *Directing the Biosynthesis of*
527   *Putrebactin or Desferrioxamine B in Shewanella putrefaciens through the*
528   *Upstream Inhibition of Ornithine Decarboxylase*, .
529   36   W. Keller-Schierlein, P. Mertens, V. Prelog and A. Walser, *Helvetica Chimica*
530   *Acta*, 1965, **48**, 710–723.
531   37   J. A. v. Lopez, T. Nogawa, Y. Futamura, T. Shimizu and H. Osada, *Journal of*
532   *Antibiotics*, 2019, **72**, 991–995.
533   38   S. I. Elshahawi, K. A. Shaaban, M. K. Kharel and J. S. Thorson, *Chemical*
534   *Society Reviews*, 2015, 44, 7591–7697.
535   39   T. C. Johnstone and E. M. Nolan, *Dalton Transactions*, 2015, 44, 6320–6339.
536   40   G. L. Challis, *ChemBioChem*, 2005, 6, 601–611.
537   41   N. Kadi and G. L. Challis, *Methods in Enzymology*, 2009, 458, 431–457.
538   42   F. Barona-Gómez, S. Lautru, F. Francois-Xavier, P. Leblond, J. L. Pernodet and
539   G. L. Challis, *Microbiology*, 2006, **152**, 3355–3366.
540   43   A. I. Garber, K. H. Nealson, A. Okamoto, S. M. McAllister, C. S. Chan, R. A.
541   Barco and N. Merino, *Frontiers in Microbiology*, 2020, **11**, 37.
542   44   K. Blin, S. Shaw, K. Steinke, R. Villebro, N. Ziemert, S. Y. Lee, M. H. Medema
543   and T. Weber, *Nucleic acids research*, 2019, **47**, W81–W87.
544   45   F. J. Flores and J. F. Martín, *Biochemical Journal*, 2004, **380**, 497–503.
545   46   F. M. Maglangit, S. Alrashdi, J. Renault, L. Trembleau, C. Victoria, M. H. Tong,
546   S. Wang, K. Kyeremeh and H. Deng, *Organic & Biomolecular Chemistry*, ,
547   DOI:10.1039/D0OB00320D.
548   47   W. Houssen and M. Jaspars, in *Natural Products Isolation*, ed. A. Sarker, S.;
549   Latfi, Z.; Gray, Humana Press, Second., 2006, p. 380.
550   48   J. J. J. van der Hooft, J. Wandy, M. P. Barrett, K. E. V. Burgess and S. Rogers,
551   *Proceedings of the National Academy of Sciences of the United States of*
552   *America*, 2016, **113**, 13738–13743.
553   49   M. Wang, S. Rogers, W. Bittremieux, C. Chen, P. C. Dorrestein, E. L.
554   Schymanski, T. Schulze, S. Neumann and R. Meier, *bioRxiv*, 2020,
555   2020.05.09.086066.
556

Figure 1. Desferrioxamine 'life-cycle'. (A) metabolite pools are generated in the bacterium, allowing for resources to be feed into the biosynthetic machinery. (B) desferrioxamine biosynthesis is dictated by the presence or absence of the proper precursors in the metabolite pools. The substrate flexibility of DesC allows for precursor organic acids to be integrated into a wide variety of structures. (C) the mature desferrioxamines are then transported into the extracellular environment to sequester Fe3+ and finally, (D), the iron-bound ferrioxamine

metabolites are then imported via specified uptake mechanisms (desE or other membrane bound mechanisms). Created with BioRender.com.

Figure 2. Molecular network containing all sec-butanol SPE fractions from monocultures, cocultures, and instrument and medium blanks. Node size represents *m/z* of each molecular feature. Nodes are connected if the cosine score meets the threshold set in the molecular networking workflow (≥0.7) and if the minimum amount of matched fragment peaks is met (minimum of 3). Circled molecular families represent those containing desferrioxamines.

Figure 3. Desferrioxamine subnetwork 1. Desferrioxamine B (561.322), desf-05 (575.375), and C$_{11}$ acyl DFO (701.513) were identified (represented as hexagons). Desferrioxamine D1 (603.367) and acyl ferrioxamine 1/legonoxamine A (679.396) were also identified through fragmentation comparison. All features with * represent new desferrioxamines identified through MS/MS fragmentation pattern analysis.

Figure 4. Molecular network and MS/MS based structure determination. (A) molecular networks generated using GNPS were dereplicated and potentially unknown ions were found in the Raw data. (B) Predicted molecular formula were generated and the most likely candidate formula was chosen. MS/MS spectra were analysed for the initial loss of -118 or -160 to determine the N-terminal moiety of *N*-hydroxycadaverine or *N*-acetylhydroxycadaverine, respectively. (C) Based on known structure of desferrioxamines, substructures were proposed for each new metabolite and lastly, (D) putative structures were confirmed using the MS/MS data.

Figure 5. Representation of desferrioxamine structural diversity of *Streptomyces* sp. S29. (A-B) New legonoxamine derivatives were some of the most abundant

desferrioxamines present in the culture, represented here by legonoxamine G (contains N-terminal *N*-hydroxyputrescine) and legonoxamine A glycoside, respectively. (C-D) New acyl DFOs were also observed, including acyl DFO-Ds and monounsaturated acyl DFO-Bs.

Figure 6. Dicarboxylic acid analogues produced under coculture conditions, (A) Dodecandioic DFO-B and (B) Tridecandioic DFO-B. MS/MS spectra can be found in supplementary data (Figures S26 and S28, respectively).

565