2   1

1   Increases in parieto-occipital alpha-band power reflect involuntary spatial

2   attention due to a task-distracting deviant sound

Annekathrin Weise[1], Thomas Hartmann[1], Fabrice Parmentier[2], Philipp Ruhnau[3,4], Nathan Weisz[1]

[1]CCNS and Division of Physiological Psychology, Paris Lodron University of Salzburg, Salzburg,

Austria

[2] Department of Psychology, Institut Universitaris de Recerca, University of the Balearic Islands,

Palma, Spain

[3]Department of Neurology, Otto-von-Guericke University, Magdeburg, Germany

[4] Center for Behavioral Brain Sciences, Otto-von-Guericke-University, Magdeburg, Germany

14   Address correspondence to:

15   Annekathrin Weise,

16   Paris-Lodron Universität Salzburg, Division of Physiological Psychology, Hellbrunnerstraße 34/II,

17   5020 Salzburg, Austria,

18   e-mail: annekathrinweise@gmail.com

Abstract

Imagine you are focusing on the traffic on a busy city street to ride a bike safely. Suddenly, the siren of an ambulance rings out. This unpredictable sound can involuntarily capture your attention and compromise performance. Related to this, the present study addresses two questions: 1) Does a shift of *spatial* attention contribute to this type of distraction?; and 2) Does oscillatory alpha activity reflect involuntary spatial attention? We harnessed a crossmodal paradigm in which participants responded as fast and as accurately as possible to the location of a visual target (left or right of fixation). Each target was preceded by a task-irrelevant sound, usually the same (i.e. standard) animal sound. Rarely, it was a novel (i.e., deviant) environmental sound. 50% of the deviants occurred on the same side as the target, and 50% occurred on the opposite side of the target. As expected, responses were slower to targets that followed a task-distracting deviant compared to a standard. Crucially, responses were faster when targets followed deviants on the same versus different side, suggesting a *spatial* shift of attention. Magnetoencephalographic data provides corroborating support for this effect: left-hemispheric visual alpha power increased in response to the left deviant, indicating a disengagement of visual areas when visual information is outside the locus of involuntary attention. Upon the occurrence of the right deviant, right-hemispheric alpha power increased in regions functionally linked to auditory processing and attentional reorienting. Overall, our findings strengthen the view that alpha power is modulated with an involuntary shift of spatial attention. They further suggest that involuntary attention has a similar impact on sensory processing as voluntary attention, thus challenging currently held claims to the contrary.

Keywords: alpha oscillations, deviant distraction, involuntary attention, magnetoencephalography, spatial attention

Highlights

- Lateralized, task-distracting deviant sounds capture spatial attention.

- Responding is faster when a target follows a deviant on the same side.

- Deviant sounds alter visual oscillatory alpha power in a spatially selective manner.

- Those changes are driven by alpha power increases for the ipsilateral deviant.

- Thus, visual areas that process deviant-unrelated information are disengaged.

Introduction

Efficient cognitive functioning requires not only the ability to filter out task-irrelevant events to better

engage in the task at hand, but also to let changes in the soundscape that are potentially relevant for

behavior break through attention filters to adequately adapt or react to them. While such a trade-off

can be advantageous, it may also come at a cost. To exemplify this, imagine the following scenario:

you are riding a bicycle on a busy city street. While filtering out street noises helps you focus on the

traffic to ride safely, some of the task-irrelevant events may still involuntarily catch your attention.

Those may be behaviorally relevant and require further action, such as the siren of an ambulance, or

they may be completely unimportant, such as someone nearby yelling at another person across the

street. In any case, the task-irrelevant events may distract you from your current task (Parmentier,

2014).

Task-distracting events occur unexpectedly in the soundscape. In an experimental setting, these so-

called deviants usually violate predictions by deviating from an otherwise (more or less) regular

sound sequence consisting of so-called standards. Distraction due to auditory deviants has been

extensively studied via event-related potentials and behavioral measures (e.g. Bendixen et al., 2007;

Schröger and Wolff, 1998; Wetzel et al., 2012). It is well known that deviant distraction is

accompanied by distinct event-related markers (not focused on here; for reviews, see Escera et al.,

2000; Parmentier, 2014) and a behavioral effect (referred to as behavioral distraction hereafter). In a

task in which participants attend and respond to a target, behavioral distraction is defined as the

slowing of responses following the presentation of a task-distracting event. This is caused by an

involuntary (stimulus driven, bottom-up, or exogenous) shift of attention from the task towards the

deviant, which actually reflects a time penalty due to attention shifting to the deviant and then back to

the task (Parmentier et al., 2008; Schröger, 1996).

The present study has two goals for investigating the cognitive and neural underpinnings of deviant

distraction in a crossmodal scenario, such as the one described above. The first goal is to address the

nature of the shift of attention. Especially in scenarios in which the auditory deviant and the visual

target occur at different location,s there are several types of attention shifts - not necessarily mutually

exclusive - that may contribute to behavioral distraction (Parmentier et al., 2008; Parmentier, 2014).

The attention shift of interest in the current study is the spatial shift from the location of the target to

the location of the deviating sound. The second goal is to probe whether neural oscillatory alpha

activity reflects a shift of involuntary spatial attention due to a deviant, which has been widely

overlooked, to date. Before we outline how we will address these goals in detail, we will briefly

review the state-of-the-art on those topics.

1) Does a shift of *spatial* attention contribute to behavioral distraction?

In the field of deviant distraction, the nature of the involuntary shift of attention in scenarios in which

the deviant and target occur in close temporal succession, but at different locations, is still relatively

poorly understood (for a review, see Parmentier, 2014). One might assume that a deviant sound can

capture spatial attention originally allocated to the upcoming task-relevant event, and that this process

is reversed as soon as the target appears. Given that such spatial shifts require time to complete (Cheal

and Gregory, 1997; Luck et al., 1996; Shiu and Pashler, 1994), they might actually contribute to

behavioral distraction (Parmentier, 2014).

Still, strong evidence for the hypothesis of a *spatial shift of involuntary attention* is lacking. There are

only a few reports on this topic that were not able to exclude alternative (though not mutually

exclusive) interpretations (Parmentier et al., 2008), or that found ambiguous results (Corral and

Escera, 2008). However, support for a *spatial shift of involuntary attention* comes from behavioral

data outside of the deviant distraction research field. Those studies applied an exogenous spatial

cueing paradigm, in which a task-irrelevant sound, such as a noise burst, appeared lateralized (left or

right) briefly before the presentation of a visual target either on the same or opposite side.

Importantly, the location of the sound neither predicted the location, or the type, of the upcoming

visual target. Results showed that if the visual target appeared on the same side as the cue, perceptual

processing (Spence and Driver, 1997) and response time improved (Dufour, 1999; Feng et al., 2014;

McDonald et al., 2000), compared to a situation in which these events occurred on opposite sides.

This pattern of results indicated that the cues were effective in generating an involuntary shift of

spatial attention.

Further support comes from event-related potentials (e.g. Feng et al., 2014; Matusz et al., 2016;

McDonald et al. 2013) and neural oscillatory activity (Feng et al., 2017; Störmer et al., 2016)

collected during the exogenous spatial cueing paradigm. These measures have shown that an auditory

cue can bias neural activity localized to the visual cortex in the hemisphere contralateral to the

location of the sound. The magnitude of these lateralized activations over parieto-occipital areas were

informative about the perception of a subsequent visual target (Feng et al., 2014; Feng et al., 2017;

Störmer et al., 2019).

2) Does oscillatory alpha activity reflect an involuntary shift of spatial attention?

The question concerning whether oscillatory alpha activity reflects an involuntary shift of spatial

attention has been motivated by studies of voluntary (goal driven, top-down, or endogenous) spatial

attention in which alpha power has already been well-characterized and used as a neural marker (e.g.

Banerjee et al., 2011; Deng et al., 2019; Rihs et al., 2009; Sauseng et al., 2005; Thut et al., 2006;

Worden et al., 2000; Wöstmann et al., 2016). This research made use of endogenous spatial cueing

tasks, allowing attention to be modulated in a goal-driven manner via cues that guided participants'

attention to certain target locations (left or right). Irrespective of whether spatial attention was directed

to a visual or auditory target, sustained oscillatory alpha power appeared to be lateralized, especially

over parieto-occipital areas. It was also higher in the ipsilateral hemisphere, compared to the

contralateral hemisphere, relative to the attended target location. This lateralization is often driven by

an ipsilateral increase and/or a contralateral decrease in alpha power relative to the attended side,

thought to reflect the inhibition of brain regions that process task-irrelevant information at unattended

locations, or facilitate processing of task-relevant information at attended locations, respectively (for

reviews, see Hanslmayr, 2011; Klimesch, 2012; for a controversial view see Palva and Palva, 2007).

Evidence is slowly accumulating that alpha power is not only an indicator of voluntary, but also

involuntary, (spatial) attention (Feng et al., 2017; Störmer et al., 2016; Weise et al., 2016). Still, to

date, research on this topic is sparse, and there is only one study so far that has particularly focused on

deviant distraction (Weise et al., 2016). That magnetoencephalography (MEG) study used a

crossmodal distraction paradigm wherein, on each trial, a visual target followed a task-irrelevant

sound. Across trials, sounds were presented in an oddball paradigm. The standard was an animal

noise, and the deviant was a novel environmental sound. All sounds were presented binaurally. The

deviant induced low alpha power in the pretarget time window compared to standards with sources in

the occipital, parietal, and supratemporal cortices, which correlated with response speed for the

deviants only. While this study provides the first evidence that alpha power can reflect involuntary

attention shifts, it is still unknown whether alpha power can also reflect shifts in involuntary *spatial*

attention.

Initial support for the latter comes from two electroencephalographic (EEG) studies employing an

exogenous spatial cueing task. These studies found a bilateral decrease in occipital alpha power, while

the alpha power decrease contralateral to the cue was more pronounced. The alpha power modulation

built up rapidly (starting around 200 to 300 milliseconds (ms) after cue onset), and was rather short-

lived, thus resembling the properties of involuntary attention (Feng et al., 2017; Störmer et al., 2016).

Also, the authors proposed that involuntary attention to the sound has a "purely facilitatory influence

on processing targets on the cued side, with no inhibitory influence on targets on the opposite side.

This lack of an early inhibitory influence may represent a fundamental difference between involuntary

and voluntary attention" (Feng et al., 2017). No doubt, such a strong assumption based on two very

similar tasks clearly asks for more research exploiting variants of exogenous spatial cueing

paradigms, preferably using different conceptual frameworks, to either strengthen or reject this view.

Moreover, the observed changes in alpha power in these studies were accompanied by evoked activity

- the so-called auditory-evoked contralateral occipital positivity - that is known for its link to

involuntary spatial attention (McDonald et al., 2013; Feng et al., 214; Störmer et al., 2019). Critically,

both alpha power and evoked activity shared similarities in the time course, scalp topography, and

source estimation, making it difficult to interpret whether they reflect the same cognitive mechanism

or two dissociable mechanisms (Feng et al., 2017; Störmer et al., 2016).

Current approach

To tackle the two questions within the framework of deviant distraction, we employed a unique

crossmodal paradigm that combines a distraction task (Escera et al., 1998; Ruhnau et al., 2013) with

an exogenous spatial cueing task (e.g. Feng et al., 2017; Störmer et al., 2016). Participants were

instructed to respond as fast and accurately as possible, as to whether a visual target appeared on the

left or right. Each target followed a task-irrelevant sound. Similar to the distraction task, 80% of the

sounds were standards, while 20% of the sounds were deviants. Similar to the exogenous spatial

cueing task, the deviant was non-informative about the spatial position of the upcoming target. That

is, 50% of the deviants occurred on the same side as the target (Congruent condition) and the other

50% of the deviants occurred on the opposite side of the target (Incongruent condition).

At the behavioral level, we intended to reproduce a distraction effect, as indicated by longer reaction

times (RTs) to a visual target that follows a deviant versus a standard (Escera et al., 1998; Parmentier

et al., 2008; Ruhnau et al., 2013). This behavioral distraction effect can be taken as a sign of an

involuntary shift of attention from the task towards the deviant sound (Parmentier, 2014). Crucially,

we hypothesized a *spatial* shift of attention, which should be indicated by shorter RTs in the

Congruent versus Incongruent condition (i.e., when the visual target follows the auditory deviant on

the same versus different side).

In addition to collecting behavioral data, we recorded MEG to shed light on neural oscillatory activity,

with an emphasis on alpha power and its link to involuntary spatial attention. We expected the deviant

to induce a lateralized, short-lived alpha power modulation over parieto-occipital sensors that builds

up rapidly after stimulus onset (Feng et al., 2017; Störmer et al., 2016; Weise et al., 2016), thereby

matching the characteristics of involuntary spatial attention (for a review, see Corbetta and Shulman,

2002). We predicted that neural sources of this alpha power modulation would be located in 1)

parieto-occipital areas, known for their role in spatial attention, even in purely auditory tasks (Deng et

al., 2019; Deng et al., 2020; Feng et al., 2017; Störmer et al., 2016); 2) auditory areas, due to the

processing of the attention-capturing sound (ElShafei et al., 2018; Frey et al., 2014; Weisz et al.,

2014; Wöstmann et al., 2016); and 3) areas linked to the ventro-frontoparietal attention system,

comprising the temporoparietal junction and the ventral frontal cortex, which have been functionally

linked to attention reorienting (for reviews, see Corbetta et al., 2008; Vossel et al., 2014).

Method

The study was approved by the local ethics committee of the Paris Lodron Universität Salzburg and

adhered to the code of ethics of the World Medical Association (Declaration of Helsinki).

Participants

30 healthy young adults participated in the MEG experiment (12 males, mean age ± STD: 27 ± 6 yrs,

4 left-handed, 1 ambidextrous). Volunteers gave written, informed consent prior to their participation.

All participants reported normal hearing and normal or corrected-to-normal vision, and none reported

a history of neurological or psychiatric diseases. Volunteers received 10€/hour (h) or 1 credit point/h

as compensation for their participation.

Stimuli and Procedure

The experiment was programmed in MATLAB 9.1 (The MathWorks, Natick, Massachusetts, U.S.A)

using the open source Psychophysics Toolbox (Brainard 1997; Kleiner et al. 2007) and o_ptb, an

additional class-based abstraction layer on top of the Psychophysics Toolbox (https://gitlab.com/thht/

o_ptb). Stimuli were presented with precise timing using the VPixx System (DATAPixx2 display

driver, PROPixx DLP LED Projector, TOUCHPixx response box by VPixx Technologies, Canada).

The Blackbox2 Toolkit (The Black Box ToolKit Ltd, Sheffield, UK) was used to measure and correct

for timing inaccuracies between triggers and the visual and auditory stimulation. Note that the tubes

of the MEG-proof sound system caused a time-delay between sound and trigger of 16.5 ms that was

corrected within the stimulation protocol.

We employed a cross-modal paradigm combining a distraction task with an exogenous spatial cueing

task. Participants performed a visual-spatial two-alternative forced choice task (see Figure 1).

Throughout the stimulation, a frog was presented at the center of the screen that participants were

asked to fixate at all times. Every trial, a visual target in the shape of a fly was presented on the screen

next to the frog. On 50% of trials, the target appeared on the left side of the frog. On the other 50% of

trials, it appeared on the right side. Participants held their left thumb over a left-side button, and their

right thumb over a right-side button. The participants' task was to indicate on which side the fly had

been presented, by pressing the left or right button. The mapping between the response buttons and

the target location was counterbalanced across participants. The target appeared for maximally 1

second (s); otherwise, a response terminated the trial.

(A) Trial structure and timing

(B) Sound sequence

Legend

D Deviant (10% left, 10% right)
S   Standard (80%, binaural)

(C) Locations of deviant and target in different conditions

	Congruent Location		Incongruent Location	
Left	**Deviant Target**		**Deviant**	**Target**
Right		**Deviant Target**	**Target**	**Deviant**

Figure 1: Cross-modal paradigm combining a distraction task with an exogenous spatial cueing task.
(A) Trial structure and timing. During each trial, the to-be-ignored sound (illustrated as an orange bar)
preceded the task-relevant visual stimulus (illustrated as a blue bar). Participants had to respond
whether the target occurred to the left or right of the fixation frog. (B) Exemplary sound sequence.
The task-irrelevant sounds were presented in an oddball paradigm: standards were presented binaural

on 80% of trials, deviants were presented left (10%) or right (10%). (C) A visual representation of the two conditions (Congruent location, Incongruent location) and the corresponding locations (left, right) of deviants and targets.

Additionally, on each trial, a sound of 0.5 s duration (with, additionally, 10 ms rise and fall times) was presented that had to be ignored. This sound always appeared after a 0.8 s silent interval and preceded the visual target by 0.6 s. That is, a silent interval of 0.1 s separated both the auditory and visual event. The sound was either a binaurally presented standard (buzzing mosquito, p = 0.8) or a monaurally presented deviant. Deviants were chosen from a pool of 56 different environmental sounds (e.g., speech, animal voices, tool noise, etc.; p = 0.2) selected from a commercial CD (1,111 Geräusche, Döbeler Cooperations, Hamburg, Germany). All sounds were equalized for overall intensity (RMS). The trial order within the whole stimulation was pseudorandomized such that a deviant was preceded by at least two standards. One auditory-visual trial lasted maximally 2.6 s, but was usually shorter (due to the relatively fast RT).

Within one experimental block, we presented two different conditions as a function of where the deviant sound and the visual target appeared. In the Congruent condition, both the deviant and target occurred on the same side (i.e., both appeared on the left or right). In the Incongruent condition, the deviant and target occurred on opposite sides. For example, if the deviant occurred on the left, the visual target occurred on the right. Note, each deviant could occur only once within a block, and only once every three blocks, for a total of four times in the whole experiment -- twice on the left (once in the Incongruent and once in the Congruent condition) and twice on the right (once in the Incongruent and once in the Congruent condition).

The experiment consisted of seven blocks in total, each lasting ∼5 minutes (min) and consisting of 160 trials (128 standards, 16 deviants presented on the left, 16 deviants presented on the right). Thus, the stimulation (without any breaks) lasted maximally ~35 min and resulted in a total of 112 left and 112 right deviants, half Congruent and half Incongruent. The auditory stimuli were presented at 50 dB

above sensation threshold level. The auditory stimulation was delivered via MEG-compatible

pneumatic in-ear headphones (SOUNDPixx, VPixx technologies, Canada). The visual stimuli were

presented inside of the magnetically shielded room using a projector (PROPixx DLP LED Projector),

via a back projection screen and a mirror system.

Before the start of the experiment, individual sensation threshold was determined using a 200-ms

1000 Hertz (Hz) tone presented via a staircased threshold procedure, measured separately for the right

and left ear. Thereafter, participants practiced the task with fewer trials to familiarize themselves with

the experiment. Deviants were chosen randomly from the pool of environmental sounds (N=56).

These were played backwards so that participants did not become familiar with those sounds.

MEG recording and preprocessing

Before MEG recording, five head position indicator (HPI) coils were placed on the left and right

mastoid, on the upper right and lower right forehead, and on the left middle forehead. Anatomical

landmarks (nasion and left/right preauricular points), the HPI locations, and at least 300 headshape

points were sampled using a Polhemus FASTTRAK digitizer. Six pre-gelled and self-adhesive

Ag/AgCl electrodes (Ambu Neuroline 720) were attached to record the vertical and horizontal

electrooculogram (EOG) and the electrocardiogram (ECG). One additional electrode - placed at the

right shoulder - served as ground.

Participants were in a seated position during data recording. A chin rest was used to avoid head

movements during and across different blocks of recording. The eye, heart, and magnetic signals of

the brain were recorded at 1000 Hz (hardware filters: 0.03 - 330 Hz) in a standard passive

magnetically shielded room (AK3b, Vacuumschmelze, Germany) using a whole-head MEG (Elekta

Neuromag Triux, Elekta Oy, Finland). Signals were captured by 102 magnetometers and 204

orthogonally placed planar gradiometers at 102 different positions.

MEG data were preprocessed and analyzed offline using Fieldtrip (Oostenveld et al., 2011), an open

source toolbox for Matlab (www.mathworks.com), and custom-written functions. Maxfilter (version

2.2.15, Elekta Neuromag, Finland) was applied to the continuous MEG raw data using a signal space

separation algorithm (Taulu and Kajola, 2005; Taulu and Simola, 2006) provided by the MEG

manufacturer. This allowed us to 1) automatically remove and interpolate the data of bad channels; 2)

remove artifacts from the MEG signal (50 Hz line-noise and its harmonics, 16.7 Hz train noise, and

muscle activity with origins beyond the head); and 3) compensate, offline, for different head positions

across blocks by realigning the data to a common standard head position (-trans default Maxfilter

parameter) based on the head position measured at the beginning of each block. Thereafter,

continuous data were visually inspected and sample points of extensive artifacts were stored

separately for each block and participant.

Two different pipelines served for further preprocessing of the MEG data. First, preprocessing of the

data was optimized for independent component analysis (ICA). Therefore, MEG data were filtered

offline using a 100 Hz lowpass filter (sinc FIR, kaiser window, order 3624, cutoff (-6 dB) 100 Hz)

and a 1-Hz highpass filter (sinc FIR, kaiser window, order 908, cutoff (-6 dB) 1 Hz). After rejecting

sample points covering extensive artifacts, runica ICA was applied in order to identify components

originating from eye movements and heartbeat. The second preprocessing pipeline was optimized for

further data analysis. Specifically, cleaned MEG data were filtered offline using a 35 Hz lowpass filter

(sinc FIR, kaiser window, order 3624, cutoff (-6 dB) 35 Hz). Data were downsampled to 256 Hz.

Thereafter, a 0.1 Hz highpass filter was applied (sinc FIR, kaiser window, order 9056, cutoff (-6 dB)

0.1 Hz).

Filtered continuous data (originating from the second preprocessing pipeline) were cleaned from the

ICs (on average two per participant). Data were segmented into 3-s epochs (-1.5 s to 1.5 s time-locked

to the onset of the sound) and cleaned from artifacts that were identified via visual inspection (see

above). For further analysis, we used deviant trials only. Trials in which deviants occurred on the left

or right are, hereafter, referred to as Deviant Left or Deviant Right, respectively. Only those deviant

trials for which participants responded once, correctly, and within the predefined response window

(100 to 800 ms following the target), entered the analysis. This resulted in 106.53/112 (SD = 3.38)

Deviant Left trials and 106.93/112 (SD = 2.89) Deviant Right trials, on average.

Data analyses

Behavioral Analysis

The following trials were excluded from the analysis of behavioral data: trials without responses,

trials including responses faster than 100 ms or slower than 800 ms after target onset, and trials that

included more than one button press. Individual RTs and hit rates were calculated.

Statistical analyses. We used jamovi (version 1.2.22.0, https://www.jamovi.org/) for the statistical

analyses of behavioral data. To determine the behavioral distraction effect on RTs and hit rates, we

applied two-way repeated-measures analyses of variance (ANOVAs) using the factors sound type

(deviant, standard) and deviant location (left, right). Note that we used standards before the

corresponding deviants for the analyses. To determine the behavioral effect of the spatial shift of

attention on RTs and hit rates, we applied two-way repeated-measures ANOVAs using the factors

congruence type (congruent, incongruent) and deviant side (left, right). Note that for statistical

purposes, hit rates were transformed to rationalized arcsine units (rau) to account for non-normality of

proportion data (Studebaker, 1985). We reported the generalized eta-squared (η^2_G; Bakeman, 2005) as

a measure of effect size. Where appropriate, we reported mean values (M) and the standard error of

the mean (SEM) in the format M ± SEM.

MEG Data Analyses

For the analyses of MEG data in sensor and source space, we solely used the signal recorded at the

gradiometers. Data were detrended to remove slow frequency shifts.

Sensor space analyses.

Time-frequency analysis. Time-frequency analysis was performed on the preprocessed single trial

data between 1 and 30 Hz (1 Hz steps) using Morlet wavelets with a width of 5 cycles, for each

participant and condition, separately. The analysis time window was shifted in steps of 50 ms between

-1.5 to 1.5 s relative to deviant onset. Note that power was first computed for each individual

gradiometer. In a second step, the power for each gradiometer pair was summed in order to yield an

orientation independent estimate. Thereafter, single trial spectral power was log transformed with a

base of 10. For each participant, trials were averaged for Deviant Left and Deviant Right, separately.

In order to statistically examine alpha power modulations in dependence of the deviant side (Deviant

Left, Deviant Right) in both hemispheres, we performed non-parametric, cluster-based dependent-

samples t-tests, with Monte-Carlo randomization (Maris and Oostenveld, 2007). The tests were

restricted to the alpha frequency range (8 – 14 Hz) and the time window ranging from 0.2 to 0.6 s

following the deviant sound and preceding the target (cf. Feng et al., 2017; Störmer et al., 2016;

Weise et al., 2016). Since we had a priori hypotheses concerning the direction of the lateralized alpha

power modulations expected in each hemisphere, we applied paired-sample, one-tailed Student's t-

tests for the left-hemispheric channels and the right-hemispheric channels, separately. The Student's t-

tests were applied for each sensor-time-frequency bin separately for each hemisphere by testing the

alpha power values of Deviant Left against the alpha power values of Deviant Right. Based on this,

clusters were formed among neighboring sensors of the respective hemisphere and corresponding

time-frequency bins that exceeded an a priori chosen threshold (p = 0.05). The sum of the t-values

within the respective cluster was used as the cluster-level statistic. The cluster with the maximum sum

was subsequently used as the test statistic. By permuting the data between the two conditions and

recalculating the test statistic 10,000 times, we obtained a reference distribution of maximum cluster

t-values to evaluate the significance of the to-be-tested contrast. The power distributions between the

two conditions were considered to be significantly different from each other if no more than 5% of the

sum-t-values of all of the 10,000-permuted data were larger than the sum-t-value of the to-be-tested

cluster.

In addition, we ran the time frequency analysis a second time using the same parameter as described

above, but with a preceding step that phase-locked activity; that is, the evoked signal averaged across

trials of the respective condition was subtracted from every single trial, to increase sensitivity to non-

phase locked signals, and to reduce contamination by phase-locked signals. The results are

comparable to those obtained via the time frequency analysis of phase-locked activity, and can be

found in the supplementary material.

We further investigated the cause of the alpha power modulation observed in the data (Figure 3).

Therefore, we analyzed the time courses of ipsi- and contralateral alpha power for the right centro-

temporal and the left parieto-occipital region of interest (ROI), respectively, that had been identified

via the preceding cluster-based permutation analysis (see above). Each ROI consisted of three

channels that showed prominent deviant-related effects in the deviant–target interval. Data were

pooled within each ROI and across frequencies (right centro-temporal ROI: 8 – 14 Hz, left parieto-

occipital ROI: 9-13 Hz; see results). Alpha power was submitted to non-parametric, cluster-based,

dependent-samples t-tests with Monte-Carlo randomization (n = 10,000; Maris and Oostenveld,

2007), contrasting the data at each sample point in the deviant-target interval (0.2 to 0.6 s following

the deviant) with the mean alpha power of the baseline period (-0.4 to -0.2 s; Feng et al., 2017). Since

we had no a priori hypotheses on the direction of the alpha power modulations expected in the

respective ROI, we applied paired-sample, two-tailed Student's t-tests.

Source space analyses.

Source projection of time series data. For each participant, realistically shaped, single-shell head

models (Nolte, 2003) were computed by co-registering the participants' head shapes either with their

structural MRI (18 participants) or—when no individual MRI was available (12 participants)—with a

standard brain from the Montreal Neurological Institute (MNI, Montreal, Canada), warped to the

individual head shape. A grid with 3 millimeter resolution based on an MNI template brain was

morphed to fit the brain volume of each participant. The leadfield was calculated for each grid point

and the forward model was computed for gradiometers only.

To project the preprocessed single-trial sensor data into source space (i.e., to the points of the grid),

we applied the linearly constrained minimum variance (LCMV) spatial filter (Van Veen et al., 1997).

We followed a procedure described for single virtual sensors (http://www.fieldtriptoolbox.org/tutorial/

virtual_sensors/) and extended it to 146,400 points covering the gray matter of the brain. The

covariance matrix across all trials (including Deviant Left and Deviant Right conditions) was

calculated and used to obtain a LCMV beamformer spatial filter for each of the grid points (for a

similar approach, see Neuling et al., 2015; Ruhnau et al., 2016). The covariance window for the

calculation of the beamformer filter was based on a 2-s time window centered at deviant onset. To

optimize the analysis in source space (i.e., increase spatial resolution using a high-definition grid and

at the same time compensate for computation time), we divided the individual brain into 333 parcels

(Gordon et al., 2016; http://www.nil.wustl.edu/labs/petersen/Resources.html) based on the

'Anatomical Automatic Labeling' [AAL] atlas (Tzourio-Mazoyer et al., 2002; provided by the

fieldtrip toolbox). That is, for each anatomical area, the spatial filters of the corresponding grid points

were averaged to obtain one spatial filter per parcel. The resulting 'averaged filter' was then

multiplied with the sensor level data to obtain time series data for each of the 333 parcels. A

regularization parameter (lambda = 5%) was applied.

Time-frequency analysis. For each participant, time-frequency analysis was performed on the single

trial data in source space between 1 and 30 Hz (1 Hz steps) using Morlet wavelets with a width of 5

cycles. The analysis time window was shifted in steps of 50 ms between -1.5 to 1.5 s relative to

deviant onset. For each participant, trials were averaged separately for Deviant Left and Deviant

Right.

To localize the deviant induced sensor level effects in source space, data were restricted to the time-

frequency ranges covering the alpha power effect (right centro-temporal ROI: 8 – 14 Hz and 0.3 - 0.5

s, left posterior ROI: 9 - 13 Hz and 0.2 - 0.6 s). Alpha power was averaged over frequency and time. T

values were calculated via a paired-sample, one-tailed Student's t-test for each of the parcels and the

corresponding time-frequency bin, contrasting the alpha power values between Deviant Left and

Deviant Right. The respective labels of localized brain regions were identified with an anatomical

brain atlas (AAL atlas; Tzourio-Mazoyer et al., 2002) and a network parcellation atlas (Gordon et al.,

2016).

Results

Behavioral data

In terms of accuracy, task performance was generally very good, as indicated by the high hit rate

(>0.97) for targets, regardless of whether they were preceded by deviants or standards (Figure 2A).

The two-way repeated measures ANOVA on hit rates shows a significant main effect for factor

stimulus type ($F(1, 29) = 18.51$, $p < .001$, $\eta^2_G = .087$), with higher hit rates for deviant trials (0.99 ±

0.003) than standard trials (0.97 ± 0.004). There was neither a significant main effect for factor

deviant side, or a significant interaction. This result may be linked to an increase in nonspecific

arousal upon the occurrence of a deviant (Näätänen, 1992), which nevertheless facilitates task

performance. Hit rates did not differ when deviant and target occurred on the congruent vs.

incongruent side (Figure 2B), as indicated by the two-way repeated-measures ANOVA yielding

neither significant main effects for the factors congruence type and deviant side, or a significant

interaction.

As depicted in Figure 2A, RTs were prolonged when targets followed a deviant versus a standard.

This is substantiated by the two-way repeated measures ANOVA on RTs, yielding a significant main

effect for stimulus type ($F(1, 29) = 39.68$, $p < .001$ $\eta^2_G = .037$; deviants: 296 ± 9 ms; standards: 278 ±

8 ms). There was neither a significant main effect for factor deviant side, or a significant interaction.

Overall, these results point to a behavioral distraction effect due to an attention shift towards the task-

distracting deviant. As depicted in Figure 2B, RTs were shorter when deviant and target occurred on

the congruent versus incongruent side. This is substantiated by the two-way repeated measures

ANOVA on RTs, yielding a significant main effect for congruence type ($F(1, 29) = 4.59$, $p < .041$ η^2_G

$= .002$; congruent: 293 ± 9 ms; incongruent: 298 ± 10 ms). There was neither a significant main effect

for factor deviant side, or a significant interaction. Overall, these results point to a *spatial* shift of

attention towards the task-distracting deviant sound.

Figure 2: Bar plots of RTs and hit rates in response to visual targets to determine a distraction effect

(A) or a shift of spatial attention (B) caused by the deviant. (A) Prolonged RTs and higher hit rates for

targets following deviants presented on the left or right side versus standard before deviant. (B)

Shortened RTs for targets following deviants on the same versus different side in the Incongruent and

Congruent conditions. Data were analysed with respect to the deviant side (left, right). Hit rates did

not differ between conditions. Error bars represent SEM.

Alpha power analysis

We analyzed alpha-band (8 - 14 Hz) power to probe whether attentional reorienting towards the

location of the deviant sound is also reflected in neural oscillatory activity, as indexed via the current

RT data. We followed a methodological approach used to measure voluntary attention (Banerjee et

al., 2011; Deng et al., 2020; Haegens et al., 2011; Wöstmann et al., 2016) and contrasted alpha power

for the cued / attended target positions (attend left versus attend right). Accordingly, we contrasted

alpha power for the left and right attention-capturing, task-distracting deviants (Deviant Left condition

vs. Deviant Right condition). Since we had a priori hypotheses concerning the direction of the

lateralized alpha power modulations expected in each hemisphere, we applied the corresponding, non-

parametric, cluster-based permutation statistics (using paired-sample, one-tailed Student's t-tests) on

the left-hemispheric and the right-hemispheric channels, separately. Overall, the results clearly show

that alpha power to lateralized deviants is modulated in a spatially selective manner (Figure 3).

Specifically, the results for the left hemispheric channels show that alpha power is significantly higher

ipsilateral to the deviant location in the time window from about 0.2 s to 0.6 s following the auditory

deviant and preceding the visual target (see Fig. 3 A). It mainly covers the frequencies from 9 - 13 Hz.

This effect is most pronounced over left parieto-occipital sensors (p = .023), as highlighted via the

scalp distribution of the alpha power modulation for the frequency range 9 - 13 Hz and the time range

0.2 to 0.6 s following the deviant (see Fig. 3 C). The main generators of this sensor level effect are

estimated in occipital regions, including primary and secondary visual cortex (see Fig. 3 E).

The results for the right hemispheric channels show that alpha power, including the frequencies 8 - 14

Hz, is significantly lower contralateral to the deviant location in the time window from about 0.3 s to

0.5 s following the auditory deviant and preceding the visual target (see Fig. 3 B). This effect is most

pronounced over right centro-temporal sensors (p = .012), as highlighted via the scalp distribution of

the alpha power modulation for the frequency range 8 - 14 Hz and the time range 0.3 to 0.5 s

following the deviant (see Fig. 3 D). The main generators of the sensor level effect are estimated in

parieto-occipital, auditory, and somatosensory regions, as well as in regions belonging to the ventral

attention system (see Fig. 3 F).

Figure 3: Log10-transformed alpha power modulation to lateralized deviants based on group-

averaged data for Deviant Left minus Deviant Right, with a focus on the left hemispheric parieto-

occipital effect (left column) and the right hemispheric centro-temporal effect (right column). (A and

B) Time frequency representations highlighting (A) high alpha power ipsilateral to the deviant

location at left-hemispheric posterior gradiometers (average across three channels showing a

prominent effect, marked in C with a star), and (B) low alpha power contralateral to the deviant

location at right-hemispheric centro-temporal gradiometer (average across three channels showing a

prominent effect, marked in D with a star). The black box marks the time-frequency range (time: 0.2 -

0.6 s, frequency: 8 - 14 Hz) used for the statistical cluster analysis. (C and D) Topoplot of the alpha

power distribution (C: 9 -13 Hz, D: 8-14 Hz) in the time window showing a significant alpha power

modulation (C: 0.2 - 0.6 s, D: 0.3 - 0.5 s). Stars indicate channels on which there was a prominent

statistical effect. (E and F) Source estimation of the (E) left hemispheric parieto-occipital alpha power

effect and (F) the right hemispheric centro-temporal alpha power effect.

To test whether the observed alpha lateralization (see Fig. 3) was driven by a decrease or increase in

alpha-band power, we analyzed the time courses of alpha power for the right hemispheric centro-

temporal ROI and the left hemispheric parieto-occipital ROI, separately, and the corresponding ipsi-

and contralateral deviant. We contrasted the alpha power following the deviant with the mean alpha

power of the baseline period, via a non-parametric cluster-based permutation statistic. Since we had

no a priori hypotheses on the direction of the alpha power modulations, we applied paired-sample,

two-tailed Student's t-tests. The results show that the right/ipsilateral deviant caused a significant

increase in alpha power in the right-hemispheric centro-temporal ROI, in the time range 0.2 to 0.4 s

after sound onset ($p = .017$, see Fig. 4 A). Furthermore, for the left/ipsilateral deviant, we found a

significant increase in alpha power in the left-hemispheric parieto-occipital ROI, in the time range 0.2

to 0.3 s after sound onset ($p = .044$, see Fig. 4 B). Additionally, the right/contralateral deviant caused

a significant decrease in alpha power in the left-hemispheric parieto-occipital ROI, in the time range

0.5 to 0.6 s after sound onset ($p = .044$, see Fig. 4 B).

Figure 4: What drives the alpha power modulations? The ipsilateral deviant conditions caused a

significant increase in alpha power compared to baseline. Deviant-locked time courses of contra- and

ipsilateral log10-transformed and group-averaged alpha power for (A) the right centro-temporal ROI

(8 - 14 Hz) and (B) the left parieto-occipital ROI (9 -13 Hz). Red traces indicate deviant ipsilateral

and blue traces indicate deviant contralateral. Channels belonging to the ROI are marked with stars in

the corresponding channel layout. Significant time samples are illustrated by the horizontal red and

blue traces (*p < .05). Upward directed arrows indicate an alpha power increase, and the downward

directed arrow indicates an alpha power decrease. Gray areas indicate the baseline interval (-0.4 to -

0.2 s) and the test interval (0.2 to 0.6 s). The orange and green bar indicate the presentation time of

deviant and target, respectively.

Discussion

Our MEG study aimed to answer two questions concerning the cognitive and neural underpinnings of

behavioral distraction: 1) Does a shift of *spatial* attention, due to a lateralized deviant sound,

contribute to behavioral distraction? 2) Does neural oscillatory alpha activity reflect this involuntary

spatial attention shift? Our RT and neural oscillatory MEG data provide a positive answer to each of

these questions, which we will discuss in more detail in the following paragraphs.

1) The shift of involuntary *spatial* attention contributes to behavioral distraction

As expected, our behavioral distraction effect - that is, the longer RTs to targets that followed a

deviant versus a standard - shows that the deviant captured attention involuntarily. This strengthens

previous findings that have been obtained via a roughly comparable behavioral distraction task

numerous times before (Bendixen et al., 2007; Parmentier et al., 2008; Schröger and Wolff, 1998;

Wetzel et al., 2012).

Importantly, we observed an advantage in response speed when deviants and targets occurred on the

same versus opposite side. This clearly speaks in favor of a *spatial* shift of attention, which has been

proposed, though not directly tested, earlier (Parmentier et al., 2008; Weise et al., 2016). That is,

when the deviant and target occur on the same side, the time penalty due to the spatial shift is lower

compared to when both events occur on opposite sides. This finding is in line with behavioral data

obtained via the exogenous cueing paradigm (McDonald et al., 2000), and receives further support

from our MEG data, as will be discussed below.

Note that we do not claim that the spatial component of the attention shift fully accounts for the

overall behavioral distraction effect; in fact, several cognitive determinants may contribute to it. A

second potential determinant that could be at play in a crossmodal situation, is the involuntary shift of

attention across sensory modalities (i.e., from vision to audition). In this case, the time penalty

accumulated by a shift of attention from the visual to the auditory channel (i.e., upon the onset of the

deviant), and its re-orientation towards the visual channel (i.e., upon the onset of the target), would

contribute to behavioral distraction (Parmentier, 2014). This idea receives support from behavioral

studies showing that RTs are slower for a target in an unexpected modality compared to a target in an

expected modality (Boulter, 1977; Spence et al., 2001).

The RT data seem to be at odds with the hit rates, at first glance. While the RTs indicate a decrease in

performance for targets following a deviant versus a standard, the hit rates rather reflect an increase.

However, this apparent contradiction can be reconciled by linking the RTs to the costs of the attention

shift towards the deviant (Schröger and Wolff, 1998; Parmentier et al., 2008; for a review, see

Parmentier 2014), and by linking the hit rates to an increase in nonspecific arousal that improves

cognitive functioning (Näätänen, 1992). The current findings are well in line with previous research

on deviant distraction, showing both costs and benefits (SanMiguel et al., 2010; Wetzel et al., 2012,

Wetzel et al, 2019).

2) Oscillatory alpha power reflects an involuntary shift of spatial attention

Our study is the first to show that a lateralized deviant sound modulates alpha power in a spatially

selective way, which cannot be attributed to stimulus-driven evoked activity (Figure 3). Similarly to

recent EEG data collected during an exogenous spatial cueing paradigm (Feng et al., 2017; Störmer et

al., 2016), we observed a lateralized alpha power modulation over parieto-occipital sensors that

occurred relatively early and was rather short-lived, thereby closely matching the characteristics of

involuntary attention both in timing and in the build-up phase (Corbetta and Shulman, 2002). Our

whole-brain analyses estimated that the main sources of this modulation occurred in the alpha band in

the visual cortex, thus confirming and extending earlier EEG data (Feng et al., 2017; Störmer et al.,

2016) using an MEG approach.

Of utmost importance, we found a left-hemispheric alpha power modulation over parieto-occipital

sensors that was higher ipsilateral versus contralateral to the deviant location (Figure 3 A, C). This

effect was driven by an alpha power increase (relative to baseline) due to the left deviant (Figure 4 B).

The current finding is in line with that obtained in endogenous spatial cueing tasks, in which attention

was voluntarily guided to the location of either a visual (Kelly et al., 2006; Rihs et al., 2009; Worden

et al., 2000) or auditory target (Banerjee et al., 2011; ElShafei et al., 2018; Frey et al., 2014; Müller

and Weisz, 2012). In those tasks, lateralized alpha power was larger over the ipsilateral than the

contralateral hemisphere, relative to the attended target location. This alpha power modulation was

driven by 1) an alpha power decrease in the contralateral hemisphere and/or an alpha power increase

ipsilateral to the attended side (e.g. Haegens et al., 2011); 2) a bilateral alpha power decrease that was

maximal over the hemisphere contralateral to the attended position (e.g. Thut et al., 2006); or 3) a

bilateral alpha power increase that was maximal over the hemisphere ipsilateral to the attended side

(e.g. Banerjee et al., 2011; Worden et al., 2000). The pattern that is actually observed, reflecting

mechanisms of suppression or enhancement at play in attention biasing, probably relies on divergent

task demands of the respective study (Kelly et al., 2006; Thut et al., 2006). In any case, based on those

studies, it has been widely acknowledged that decreased alpha power reflects the processing of task-

relevant information in the voluntarily attended space, and increased alpha power indicates the

inhibition of processes associated with task irrelevant information in the unattended space (for a

review, see Klimesch, 2012). In a similar vein, our left-hemispheric parieto-occipital alpha power

increase due to the left deviant can be linked to the suppression of irrelevant information in the

unattended space, in that deviant-induced involuntary spatial attention is *not* captured. This active

inhibition of visual brain areas may free up resources for brain areas contralateral to the deviant site

that process relevant information (Jensen and Mazaheri, 2010; Meeuwissen et al., 2011), which are

also consequently involved in attention reorienting towards the deviant location. This interpretation

receives support from the current behavioral data, showing that deviants capture spatial attention

(Figure 2B). Furthermore, it is in line with findings suggesting that a sound can engage visual

processing at the same location (Feng et al., 2014; Feng et al., 2017; McDonald et al., 2013; Störmer

et al., 2016).

Importantly, the current pattern of results contradicts the prevailing view on alpha power in the

context of involuntary attention. Earlier studies used exogenous spatial cueing paradigms to

demonstrate a bilateral alpha power decrease that was more prominent in the hemisphere contralateral

to the task-irrelevant sound. Based thereon, the authors suggested that a shift in involuntary attention

exclusively facilitates target processing on the cued side, whereas target processing on the opposite

side is not negatively impacted (Feng et al., 2017; Störmer et al., 2016). This conclusion was

substantiated by the characteristic pattern of alpha power modulation preceding correct and incorrect

discriminations of valid and invalid targets (Feng et al., 2017). Our results - demonstrating an increase

of parieto-occipital alpha power - challenge that conclusion, as well as the prevalent notion that the

"lack of an early inhibitory influence may represent a fundamental difference between involuntary

orienting and voluntarily directed attention" (Feng et al., 2017, page 326). The difference between the

current and previous findings is that the current study did not only apply an exogenous spatial cueing

task, but also a distraction task, thus resulting in different task demands (Kelly et al., 2006; Thut et al.,

2006).

3) Oscillatory alpha power indicates deviant-related processes beyond involuntary spatial

attention

In addition to the current evidence pointing to a spatial shift of attention, the current   right-

hemispheric alpha power modulation over centro-temporal sites (Figure 3 B, D) gives more insights

into the processing of deviant-related information in a crossmodal scenario. Similar to the left

hemispheric visual alpha power modulation, the right-hemispheric effect was driven by an increase in

alpha power, relative to baseline, due to the right/ipsilateral deviant (Figure 4 A). This reflects

inhibition of the underlying brain regions, namely the parieto-occipital, auditory, and somatosensory

areas, as well as the ventral attention network (Figure 3 F). This active inhibition leads to an increased

allocation of resources to contralateral brain regions that are responsible for processing relevant

deviant-related information (Jensen and Mazaheri, 2010; Meeuwissen et al., 2011). Specifically, the

left auditory cortex should be engaged in processing the right deviant, and somatosensory cortex

should be involved in processing the anticipated behavioral relevance of the right deviant. Based

thereon, we can draw further conclusions about the nature of attention reorienting in crossmodal

scenarios. The current modulation in auditory alpha power fits the notion that, in crossmodal

scenarios, attention is not only shifted spatially, but also across modalities (i.e., from vision to

audition; Parmentier, 2014; Weise et al., 2016). This interpretation supports the involvement of a

resource-limited supramodal attention mechanism that operates independent of stimulus modality

(Banerjee et al., 2011; Deng et al., 2020; Wöstmann et al., 2016). Therefore,, when attention shifts

from vision to audition, auditory processing is facilitated at the cost of diminished visual processing.

This explanation accounts for our behavioral result showing longer RTs to visual targets that follow a

deviant versus a standard (cf. Ruhnau et al., 2013; Weise et al., 2016), which cannot be attributed

exclusively to the spatial shift of attention (see above).

The pattern of auditory alpha power modulation observed in the current study contributes to the

growing number of findings highlighting distinct alpha generators in the auditory cortex. So far, this

has mainly been shown using endogenous (spatial) cueing paradigms (Billig et al., 2019; Frey et al.,

2014; ElShafei et al., 2018; Müller and Weisz, 2012; Weisz et al., 2014; Wöstmann et al., 2016). Our

data importantly extend those results to the field of involuntary spatial attention.

The right-hemispheric alpha power modulation in regions related to somatosensory processing may

point to the allocation of resources in contralateral brain regions in preparation for action. Though

auditory deviants were not relevant to the task, the brain seems to classify those sounds as

behaviorally relevant. This receives support from the engagement of the ventral attention network,

which is activated when behaviorally relevant events are detected (Corbetta et al., 2000). Note that the

observed somatosensory alpha power modulation cannot be explained by preparatory processes

related to the task for two reasons: 1) Participants responded to the left and right target with both

hands, and the button-response assignment was counterbalanced across participants; 2) The alpha

power modulation occured in the pre-target interval, whereas the responding hand was only

determined once the target stimulus appeared.

The observed right hemispheric alpha power modulation in areas linked to the ventral attention

network fits to the right hemispheric bias of that network known mainly from fMRI studies (e.g.

Downar et al., 2000; Corbetta et al., 2000; Vossel et al., 2012). Given that this fronto-parietal network

has been functionally linked to stimulus-driven attention reorienting (for reviews, see Corbetta and

Shulman, 2002; Corbetta et al., 2008), our MEG data establish a strong link between neural

oscillatory activity in the alpha range and involuntary attention. One reason why the current MEG

data showed sources in the fronto-parietal network, whereas previous EEG data did not, (Feng et al.,

2017; Störmer et al., 2016) might be due to the different stimulus material that was used. Earlier

studies used salient noise bursts, whereas we used environmental sounds that have real-world

behavioral relevance (e.g., baby crying, telephone ringing, etc.). In fact, previous fMRI studies

suggest that behaviorally relevant events activate the ventral attention network, whereas salient, but

noninformative, events do not (Downar et al., 2001). Furthermore, the observed alpha power

modulation in the ventral attention network fits the view that this neural network is particularly

engaged whenever environmental events elicit a change in the current task, and especially when these

events are rare, as in the current study (for a review, see Corbetta and Shulman, 2008). Even though

only the visual targets were task-relevant, the auditory deviants produced a clear behavioral

distraction effect, which can be regarded as a temporary change in the task.

4) Conclusion

Here, we show that lateralized deviant sounds cause a shift of involuntary spatial attention and

contribute to behavioral distraction. This is evident in our reaction time data, which demonstrates that

responses are faster when the target follows an auditory deviant on the same side compared to the

opposite sides. Additionally, our MEG data clearly show that oscillatory alpha power is shaped in a

spatially selective manner. Most importantly, attention-capturing, task-distracting deviant sounds

induced high visual alpha power in the left hemisphere. These attention-related changes were driven

by alpha power increases due to the left deviant, suggesting disengagement of visual areas that

process task-irrelevant information outside the locus of involuntary attention. Together with the

behavioral data, our oscillatory data strengthen the view that alpha power reflects a shift of

involuntary spatial attention. Critically, the current results challenge the prevailing view that

involuntary attention capture only affects facilitatory mechanisms.

Acknowledgments

We thank Jonas Heilig for his help in collecting and preprocessing the data. We also thank Fred

Seifter for his technical support in the lab.

Funding: This work was supported by the Deutsche Forschungsgemeinschaft (DFG, WE 6085/1-1).

Data and code availability

The preprocessed MEG and behavioral data used to generate the figures and calculate the statistics are

available at the OSF public repository (https://osf.io/qt95u/). The corresponding data analysis code is

available at the corresponding author's GitLab repository (https://gitlab.com/aweise/crossfrog-final-

analysis). The original raw data are available, upon reasonable request, from the corresponding

author.

710
711
712

References

Bakeman, R. (2005). Recommended effect size statistics for repeated measures designs.

Behavior Research Methods, *37*(3), 379–384. https://doi.org/10.3758/bf03192707

Banerjee, S., Snyder, A. C., Molholm, S., & Foxe, J. J. (2011). Oscillatory alpha-band

mechanisms and the deployment of spatial attention to anticipated auditory and visual target

locations: Supramodal or sensory-specific control mechanisms? *The Journal of*

Neuroscience, *31*(27), 9923–9932. https://doi.org/10.1523/JNEUROSCI.4660-10.2011

Bendixen, A., Roeber, U., & Schröger, E. (2007). Regularity extraction and application in

dynamic auditory stimulus sequences. *Journal of Cognitive Neuroscience*, *19*(10), 1664–

1677. https://doi.org/10.1162/jocn.2007.19.10.1664

Billig, A. J., Herrmann, B., Rhone, A. E., Gander, P. E., Nourski, K. V., Snoad, B. F., …

Johnsrude, I. S. (2019). A sound-sensitive source of alpha oscillations in human non-

primary auditory cortex. *Journal of Neuroscience*, *39*(44), 8679–8689.

https://doi.org/10.1523/JNEUROSCI.0696-19.2019

Boulter, L. R. (1977). Attention and reaction times to signals of uncertain modality. *Journal of*

Experimental Psychology. Human Perception and Performance, *3*(3), 379–388.

https://doi.org/10.1037//0096-1523.3.3.379

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Cheal, M. L., & Gregory, M. (1997). Evidence of limited capacity and noise reduction with

single-element displays in the location-cuing paradigm. *Journal of Experimental*

Psychology. Human Perception and Performance, *23*(1), 51–71.

https://doi.org/10.1037//0096-1523.23.1.51

Corbetta, M., Kincade, J. M., Ollinger, J. M., McAvoy, M. P., & Shulman, G. L. (2000).

Voluntary orienting is dissociated from target detection in human posterior parietal cortex.

Nature Neuroscience, *3*(3), 292–297. https://doi.org/10.1038/73009

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in

the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215.

Corbetta, Maurizio, Patel, G., & Shulman, G. L. (2008). The reorienting system of the human

brain: From environment to theory of mind. *Neuron*, *58*(3), 306–324.

https://doi.org/10.1016/j.neuron.2008.04.017

Corral, M.-J., & Escera, C. (2008). Effects of sound location on visual task performance and

electrophysiological measures of distraction. *Neuroreport*, *19*(15), 1535–1539.

https://doi.org/10.1097/WNR.0b013e3283110416

Deng, Y., Choi, I., & Shinn-Cunningham, B. (2020). Topographic specificity of alpha power

during auditory spatial attention. *NeuroImage*, *207*, 116360.

https://doi.org/10.1016/j.neuroimage.2019.116360

Deng, Y., Reinhart, R. M., Choi, I., & Shinn-Cunningham, B. G. (2019). Causal links between

parietal alpha activity and spatial auditory attention. *ELife*, *8*.

https://doi.org/10.7554/eLife.51184

Downar, J., Crawley, A. P., Mikulis, D. J., & Davis, K. D. (2000). A multimodal cortical

network for the detection of changes in the sensory environment. *Nature Neuroscience*,

3(3), 277–283. https://doi.org/10.1038/72991

Downar, J., Crawley, A. P., Mikulis, D. J., & Davis, K. D. (2001). The effect of task relevance

on the cortical response to changes in visual and auditory stimuli: An event-related fMRI

study. *NeuroImage*, *14*(6), 1256–1267. https://doi.org/10.1006/nimg.2001.0946

Dufour, A. (1999). Importance of attentional mechanisms in audiovisual links. *Experimental*

Brain Research, *126*(2), 215–222. https://doi.org/10.1007/s002210050731

ElShafei, H. A., Bouet, R., Bertrand, O., & Bidet-Caulet, A. (2018). Two sides of the same coin:

Distinct sub-bands in the alpha rhythm reflect facilitation and suppression mechanisms

during auditory anticipatory attention. *ENeuro*, ENEURO.0141-18.2018.

https://doi.org/10.1523/ENEURO.0141-18.2018

Escera, C., Alho, K., Schröger, E., & Winkler, I. (2000). Involuntary attention and distractibility

as evaluated with event-related brain potentials. *Audiology and Neuro-Otology*, *5*(3–4),

151–166. http://dx.doi.org/10.1159/000013877

Escera, C., Alho, K., Winkler, I., & Näätänen, R. (1998). Neural mechanisms of involuntary

attention to acoustic novelty and change. *Journal of Cognitive Neuroscience*, *10*(5), 590–

604. http://dx.doi.org/10.1162/089892998562997

Feng, W., Störmer, V. S., Martinez, A., McDonald, J. J., & Hillyard, S. A. (2014). Sounds activate visual cortex and improve visual discrimination. *The Journal of Neuroscience*, *34*(29), 9817–9824. https://doi.org/10.1523/JNEUROSCI.4869-13.2014

Feng, W., Störmer, V. S., Martinez, A., McDonald, J. J., & Hillyard, S. A. (2017). Involuntary orienting of attention to a sound desynchronizes the occipital alpha rhythm and improves visual perception. *NeuroImage*, *150*, 318–328. https://doi.org/10.1016/j.neuroimage.2017.02.033

Frey, J. N., Mainy, N., Lachaux, J.-P., Müller, N., Bertrand, O., & Weisz, N. (2014). Selective modulation of auditory cortical alpha activity in an audiovisual spatial attention task. *The Journal of Neuroscience*, *34*(19), 6634–6639. https://doi.org/10.1523/JNEUROSCI.4813-13.2014

Gordon, E. M., Laumann, T. O., Adeyemo, B., Huckins, J. F., Kelley, W. M., & Petersen, S. E. (2016). Generation and evaluation of a cortical area parcellation from resting-state correlations. *Cerebral Cortex*, *26*(1), 288–303. https://doi.org/10.1093/cercor/bhu239

Haegens, S., Händel, B. F., & Jensen, O. (2011). Top-down controlled alpha band activity in somatosensory areas determines behavioral performance in a discrimination task. *Journal of Neuroscience*, *31*(14), 5197–5204. https://doi.org/10.1523/JNEUROSCI.5199-10.2011

Hanslmayr, S., Gross, J., Klimesch, W., & Shapiro, K. L. (2011). The role of α oscillations in temporal attention. *Brain Research Reviews*, *67*(1–2), 331–343. https://doi.org/10.1016/j.brainresrev.2011.04.002

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10.3389/fnhum.2010.00186

Kelly, S. P., Lalor, E. C., Reilly, R. B., & Foxe, J. J. (2006). Increases in alpha oscillatory power reflect an active retinotopic mechanism for distracter suppression during sustained visuospatial attention. *Journal of Neurophysiology*, *95*(6), 3844–3851.

https://doi.org/10.1152/jn.01234.2005

Kleiner, M., Brainard, D., Pelli, D. (2007). "What's new in Psychtoolbox-3?" *Perception*, 36,

ECVP Abstract Supplement.

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored

information. *Trends in Cognitive Sciences*, *16*(12), 606–617.

https://doi.org/10.1016/j.tics.2012.10.007

Luck, S. J., Hillyard, S. A., Mouloua, M., & Hawkins, H. L. (1996). Mechanisms of visual-

spatial attention: Resource allocation or uncertainty reduction? *Journal of Experimental*

Psychology. Human Perception and Performance, *22*(3), 725–737.

https://doi.org/10.1037//0096-1523.22.3.725

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data.

Journal of Neuroscience Methods, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Matusz, P. J., Retsa, C., & Murray, M. M. (2016). The context-contingent nature of cross-modal

activations of the visual cortex. *NeuroImage*, *125*, 996–1004.

https://doi.org/10.1016/j.neuroimage.2015.11.016

McDonald, J. J., Teder-Sälejärvi, W. A., & Hillyard, S. A. (2000). Involuntary orienting to sound

improves visual perception. *Nature*, *407*(6806), 906–908. https://doi.org/10.1038/35038085

McDonald, John J., Störmer, V. S., Martinez, A., Feng, W., & Hillyard, S. A. (2013). Salient

sounds activate human visual cortex automatically. *The Journal of Neuroscience*, *33*(21),

9194–9201. https://doi.org/10.1523/JNEUROSCI.5902-12.2013

Meeuwissen, E. B., Takashima, A., Fernández, G., & Jensen, O. (2011). Increase in posterior

alpha activity during rehearsal predicts successful long-term memory formation of word

sequences. *Human Brain Mapping*, *32*(12), 2045–2053. https://doi.org/10.1002/hbm.21167

Müller, N., & Weisz, N. (2012). Lateralized auditory cortical alpha band activity and

interregional connectivity pattern reflect anticipation of target sounds. *Cerebral Cortex*,

22(7), 1604–1613. https://doi.org/10.1093/cercor/bhr232

Näätänen, R. (1992). *Attention and brain function*. Hillsdale, NJ: Erlbaum.

Neuling, T., Ruhnau, P., Fuscà, M., Demarchi, G., Herrmann, C. S., & Weisz, N. (2015).

Friends, not foes: Magnetoencephalography as a tool to uncover brain dynamics during

transcranial alternating current stimulation. *NeuroImage*, *118*, 406–413.

https://doi.org/10.1016/j.neuroimage.2015.06.026

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. *Physics in*

Medicine and Biology, *48*(22), 3637. https://doi.org/10.1088/0031-9155/48/22/002

Palva, S., & Palva, J. M. (2007). New vistas for alpha-frequency band oscillations. *Trends in*

Neurosciences, *30*(4), 150–158. https://doi.org/10.1016/j.tins.2007.02.001

Parmentier, F. B. R. (2014). The cognitive determinants of behavioral distraction by deviant

auditory stimuli: A review. *Psychological Research*, *78*(3), 321–338.

https://doi.org/10.1007/s00426-013-0534-4

Parmentier, F. B. R., Elford, G., Escera, C., Andrés, P., & San Miguel, I. (2008). The cognitive

locus of distraction by acoustic novelty in the cross-modal oddball task. *Cognition*, *106*(1),

408–432. https://doi.org/10.1016/j.cognition.2007.03.008

Rihs, T. A., Michel, C. M., & Thut, G. (2009). A bias for posterior alpha-band power

suppression versus enhancement during shifting versus maintenance of spatial attention.

NeuroImage, *44*(1), 190–199. https://doi.org/10.1016/j.neuroimage.2008.08.022

Ruhnau, P., Herrmann, B., Maess, B., Brauer, J., Friederici, A. D., & Schröger, E. (2013).

Processing of complex distracting sounds in school-aged children and adults: Evidence from

EEG and MEG data. *Frontiers in Psychology*, *4*. https://doi.org/10.3389/fpsyg.2013.00717

Ruhnau, P., Neuling, T., Fuscá, M., Herrmann, C. S., Demarchi, G., & Weisz, N. (2016). Eyes

wide shut: Transcranial alternating current stimulation drives alpha rhythm in a state

dependent manner. *Scientific Reports*, *6*, 27138. https://doi.org/10.1038/srep27138

SanMiguel, I., Linden, D., & Escera, C. (2010). Attention capture by novel sounds: Distraction

versus facilitation. *European Journal of Cognitive Psychology*, *22*(4), 481–515.

https://doi.org/10.1080/09541440902930994

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., …

Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with

human EEG alpha activity. *European Journal of Neuroscience*, *22*(11), 2917–2926. https://

doi.org/10.1111/j.1460-9568.2005.04482.x

Schröger, E. (1996). A neural mechanism for involuntary attention shifts to changes in auditory

stimulation. *Journal of Cognitive Neuroscience*, *8*, 527–539.

http://dx.doi.org/10.1162/jocn.1996.8.6.527

Schröger, E., & Wolff, C. (1998). Behavioral and electrophysiological effects of task-irrelevant

sound change: A new distraction paradigm. *Brain Research. Cognitive Brain Research*,

7(1), 71–87.

Shiu, L., & Pashler, H. (1994). Negligible effect of spatial precuing on identification of single

digits. *Journal of Experimental Psychology: Human Perception and Performance*, *20*(5),

1037–1054. https://doi.org/10.1037/0096-1523.20.5.1037

Spence, C., & Driver, J. (1997). Audiovisual links in exogenous covert spatial orienting.

Perception & Psychophysics, *59*(1), 1–22. https://doi.org/10.3758/BF03206843

Spence, C., Nicholls, M. E. R., & Driver, J. (2001). The cost of expecting events in the wrong

sensory modality. *Perception & Psychophysics*, *63*(2), 330–336.

https://doi.org/10.3758/BF03194473

Störmer, V., Feng, W., Martinez, A., McDonald, J., & Hillyard, S. (2016). Salient, irrelevant

sounds reflexively induce alpha rhythm desynchronization in parallel with slow potential

shifts in visual cortex. *Journal of Cognitive Neuroscience*, *28*(3), 433–445.

https://doi.org/10.1162/jocn_a_00915

Störmer, V. S. (2019). Orienting spatial attention to sounds enhances visual processing. *Current

Opinion in Psychology*. https://doi.org/10.1016/j.copsyc.2019.03.010

Studebaker, G. A. (1985). A „rationalized" arcsine transform. *Journal of Speech and Hearing

Research*, *28*(3), 455–462. https://doi.org/10.1044/jshr.2803.455

Taulu, S., & Simola, J. (2006). Spatiotemporal signal space separation method for rejecting

nearby interference in MEG measurements. *Physics in Medicine and Biology*, *51*(7), 1759–

1768. https://doi.org/10.1088/0031-9155/51/7/008

Taulu, Samu, & Kajola, M. (2005). Presentation of electromagnetic multichannel data: The

signal space separation method. *Journal of Applied Physics*, *97*(12), 124905. https://doi.org/

10.1063/1.1935742

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). Alpha-band

electroencephalographic activity over occipital cortex indexes visuospatial attention bias

and predicts visual target detection. *Journal of Neuroscience*, *26*(37), 9494–9502.

https://doi.org/10.1523/JNEUROSCI.0875-06.2006

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., …

Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a

macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*,

15(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Van Veen, B. D., van Drongelen, W., Yuchtman, M., & Suzuki, A. (1997). Localization of brain

electrical activity via linearly constrained minimum variance spatial filtering. *IEEE*

Transactions on Bio-Medical Engineering, *44*(9), 867–880.

https://doi.org/10.1109/10.623056

Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and ventral attention systems: Distinct

neural circuits but collaborative roles. *The Neuroscientist*, *20*(2), 150–159.

https://doi.org/10.1177/1073858413494269

Vossel, S., Weidner, R., Driver, J., Friston, K. J., & Fink, G. R. (2012). Deconstructing the

architecture of dorsal and ventral attention systems with dynamic causal modeling. *Journal*

of Neuroscience, *32*(31), 10637–10648. https://doi.org/10.1523/JNEUROSCI.0414-12.2012

Weise, A., Hartmann, T., Schröger, E., Weisz, N., & Ruhnau, P. (2016). Cross-modal distractors

modulate oscillatory alpha power: The neural basis of impaired task performance.

Psychophysiology, *53*(11), 1651–1659. https://doi.org/10.1111/psyp.12733

Weisz, N., Müller, N., Jatzev, S., & Bertrand, O. (2014). Oscillatory alpha modulations in right

auditory regions reflect the validity of acoustic cues in an auditory spatial attention task.

Cerebral Cortex, *24*(10), 2579–2590. https://doi.org/10.1093/cercor/bht113

Wetzel, N., Scharf, F., & Widmann, A. (2019). Can't Ignore—Distraction by task-irrelevant

sounds in early and middle childhood. *Child Development*, *90*(6), e819–e830.

https://doi.org/10.1111/cdev.13109

Wetzel, N., Widmann, A., & Schröger, E. (2012). Distraction and facilitation—Two faces of the

same coin? *Journal of Experimental Psychology. Human Perception and Performance*,

38(3), 664–674. https://doi.org/10.1037/a0025856

Worden, M. S., Foxe, J. J., Wang, N., & Simpson, G. V. (2000). Anticipatory biasing of

visuospatial attention indexed by retinotopically specific alpha-band

electroencephalography increases over occipital cortex. *The Journal of Neuroscience*, *20*(6),

RC63.

Wöstmann, M., Herrmann, B., Maess, B., & Obleser, J. (2016). Spatiotemporal dynamics of

auditory attention synchronize with speech. *Proceedings of the National Academy of

Sciences*, *113*(14), 3873–3878. https://doi.org/10.1073/pnas.1523357113