Increases in parieto-occipital alpha-band power reflect involuntary spatial

attention due to a task-distracting deviant sound

Annekathrin Weise[1], Thomas Hartmann[1], Fabrice Parmentier[2], Philipp Ruhnau[3,4], Nathan Weisz[1]

[1]CCNS and Division of Physiological Psychology, Paris Lodron University of Salzburg, Salzburg,

Austria

[2] Department of Psychology, Institut Universitaris de Recerca, University of the Balearic Islands,

Palma, Spain

[3]Department of Neurology, Otto-von-Guericke University, Magdeburg, Germany

[4] Center for Behavioral Brain Sciences, Otto-von-Guericke-University, Magdeburg, Germany

Address correspondence to:

Annekathrin Weise,

Paris-Lodron Universität Salzburg, Division of Physiological Psychology, Hellbrunnerstraße 34/II,

5020 Salzburg, Austria,

e-mail: annekathrinweise@gmail.com

Abstract

Imagine you are focusing on the traffic of a busy city street to ride the bike safely. Suddenly, the siren of an ambulance rings out. This unpredictable sound can involuntarily capture your attention and compromise performance. Related to that the present study addresses two questions: 1) Does a shift of *spatial* attention contribute to this type of distraction? And 2) does oscillatory alpha activity reflect involuntary spatial attention? We harnessed a crossmodal paradigm where participants responded as fast and as correct as possible whether a visual target appeared on the left or right. Each target was preceded by a task-irrelevant sound, usually the same (i.e. standard) animal sound. Rarely, it was a novel (i.e. deviant) environmental sound. Crucially, 50% of the deviants occurred at the same and the other 50% at the opposite side as the target. As expected, responses were slower to targets that followed a task-distracting deviant vs. standard. Crucially, responses were faster when targets followed deviants at the same vs. different side, suggesting a *spatial* shift of attention. Corroborating support comes from our magnetoencephalographic data. Left-hemispheric visual alpha power increased in response to the left deviant, indicating that visual areas processing information outside the locus of involuntary attention are disengaged. Upon the occurrence of the right deviant right-hemispheric alpha power increased in regions functionally linked to auditory processing and attentional reorienting. Overall, our findings strengthen the view that alpha power indicates an involuntary shift of spatial attention. They further suggest that involuntary attention has a similar impact on target processing as voluntary attention, thus, challenging previous reports claiming otherwise.

Keywords: alpha oscillations, deviant distraction, involuntary attention, magnetoencephalography, spatial attention

Highlights

-    Lateralized, task-distracting deviant sounds capture involuntary spatial attention.

-    Responding is faster when a target follows a deviant at the same side.

-    Deviant sounds alter visual oscillatory alpha power in a spatially selective manner.

-    Those changes are driven by alpha power increases for the ipsilateral deviant.

-    Thus, visual areas that process deviant-unrelated information are disengaged.

Introduction

Efficient cognitive functioning requires not only the ability to filter out task-irrelevant events to better

engage in the task at hand but also to let changes in the soundscape that are potentially relevant for

behaviour break through attention filters in order to adequately adapt or react to them. While such a

trade-off can be advantageous it may also come at a cost. To exemplify this, imagine the following

scenario: You are riding a bike on a busy city street. While filtering out street noises helps focusing on

the traffic to bike safely, some of the task-irrelevant events may still involuntarily catch your

attention. Those may be behaviorally relevant and require further action such as the siren of an

ambulance or may be completely unimportant such as someone nearby yelling at another person

across the street. In any case, the task-irrelevant events may distract you from your current task

(Parmentier, 2014).

Task-distracting events occur unexpectedly in the soundscape. In an experimental setting these so-

called deviants usually violate predictions by deviating from an otherwise (more or less) regular

sound sequence consisting of so-called standards. The distraction due to auditory deviants has been

extensively studied via event-related potentials and behavioral measures (e.g. Bendixen et al., 2007;

Schröger and Wolff, 1998; Wetzel et al., 2012). It is well known that deviant distraction is

accompanied by distinct event-related markers (not focused on here; for reviews see Escera et al.,

2000; Parmentier, 2014) and a behavioral effect (referred to as behavioral distraction hereafter). In a

task where participants attend and respond to a target, behavioral distraction is defined as the slowing

down of responses following the presentation of the task-distracting event. This is caused by an

involuntary (stimulus driven, bottom-up or exogenous) shift of attention from the task towards the

deviant, actually reflecting the time penalties due to the shift of attention to the deviant and back to

the task (Parmentier et al., 2008; Schröger, 1996).

The present study aims to tackle two questions concerning the cognitive and neural underpinnings of

deviant distraction in a crossmodal scenario such as the one described above: 1) The first question

addresses the nature of the shift of attention. Especially in scenarios where the auditory deviant and

the visual target occur at different locations there are several types of attention shifts - not necessarily

mutually exclusive - that may contribute to behavioral distraction (Parmentier et al., 2008; Parmentier,

2014). The attention shift of interest in the current study is the spatial shift from the location of the

target to the location of the deviating sound. The second goal is to probe whether neural oscillatory

alpha activity reflects a shift of involuntary spatial attention due to a deviant which has been widely

overlooked to date. Before we will outline how we will address these questions in detail, we will

briefly review the state-of-the-art on those topics.

1) Does a shift of *spatial* attention contribute to behavioral distraction?

In the field of deviant distraction the nature of the involuntary shift of attention in scenarios where

deviant and target occur in close temporal succession but at different locations is still relatively poorly

understood (for a review, see Parmentier, 2014). One might assume that a deviant sound can capture

spatial attention originally allocated to the upcoming task-relevant event and that this process is

reversed as soon as the target appears. Given that such spatial shifts require time to complete (Cheal

and Gregory, 1997; Luck et al., 1996; Shiu and Pashler, 1994) they might actually contribute to

behavioral distraction (Parmentier, 2014).

Still, strong evidence for the hypothesis of a *spatial shift of involuntary attention* is lacking. There are

only few reports on that topic that could either not exclude alternative (though not mutually exclusive)

interpretations (Parmentier et al., 2008) or that found ambiguous results (Corral and Escera, 2008).

However, support for a *spatial shift of involuntary attention* comes from behavioral data outside the

deviant distraction research field. Those studies applied an exogenous spatial cueing paradigm, in

which a task-irrelevant sound, such as a noise burst, appeared lateralized (left or right) briefly before a

visual target either at the same or opposite side. Importantly, the location of the sound did neither

predict the location nor the type of the upcoming visual target. Results showed that if the visual target

appeared at the same side as the cue, perceptual processing (Spence and Driver, 1997) and response

time improved (Dufour, 1999; Feng et al., 2014; McDonald et al., 2000) compared to a situation when

these events occurred at opposite sides. This pattern of results indicated that the cues were effective in

generating an involuntary shift of spatial attention.

Further support comes from event-related potentials (e.g. Feng et al., 2014; Matusz et al., 2016;

McDonald et al. 2013) and neural oscillatory activity (Feng et al., 2017; Störmer et al., 2016)

collected during the exogenous spatial cueing paradigm. They showed that an auditory cue can bias

neural activity localized to the visual cortex in the hemisphere contralateral to the location of the

sound. The magnitude of these lateralized activations over parieto-occipital areas were informative

about the perception of a subsequent visual target (Feng et al., 2014; Feng et al., 2017; Störmer et al.,

2019).

2) Does oscillatory alpha activity reflect an involuntary shift of spatial attention?

The question whether oscillatory alpha activity reflects an involuntary shift of spatial attention has

been motivated by studies on voluntary (goal driven, top-down or endogenous) spatial attention where

alpha power has already been well characterized and used as a neural marker (e.g. Banerjee et al.,

2011; Deng et al., 2019; Rihs et al., 2009; Sauseng et al., 2005; Thut et al., 2006; Worden et al., 2000;

Wöstmann et al., 2016). That research made use of endogenous spatial cueing tasks allowing attention

to be modulated in a goal-driven manner via cues that guided participants' attention to certain target

locations (left or right). Irrespective of whether spatial attention was directed to a visual or an auditory

target, sustained oscillatory alpha power appeared to be lateralized especially over parieto-occipital

areas. It was higher in the ipsilateral than in the contralateral hemisphere relative to the attended target

location. Often, the lateralization has been driven by an ipsilateral increase and/or a contralateral

decrease in alpha power relative to the attended side, reflecting the inhibition of brain regions

processing task-irrelevant information at unattended locations or facilitated processing of task-

relevant information at attended locations, respectively (for reviews, see Hanslmayr, 2011; Klimesch,

2012; for a controversial view see Palva and Palva, 2007).

Evidence is slowly accumulating that alpha power is not only an indicator of voluntary but also

involuntary (spatial) attention (Feng et al., 2017; Störmer et al., 2016; Weise et al., 2016). Still, to

date research on that topic is sparse and there is only one study so far focusing especially on deviant

distraction (Weise et al., 2016). That magnetoencephalography (MEG) study used a crossmodal

distraction paradigm wherein on each trial a visual target followed a task-irrelevant sound. Across

trials sounds were presented in an oddball paradigm. The standard was an animal noise, the deviant

was a novel environmental sound. All sounds were presented binaurally. The deviant induced low

alpha power in the pretarget time window as compared to standards with sources in the occipital,

parietal and supratemporal cortices that correlated with response speed for the deviants only. While

this study provides first indices that alpha power can reflect involuntary attention shifts it still needs to

be tested whether it can do so for involuntary *spatial* attention.

Initial support for the latter comes from two electroencephalographic (EEG) studies employing an

exogenous spatial cueing task. They found a bilateral decrease in occipital alpha power, while the

alpha power decrease contralateral to the cue was more pronounced. The alpha power modulation

built up rapidly (starting ~around 200 to 300 ms after cue onset) and was rather short-lived, thus,

resembling the properties of involuntary attention (Feng et al., 2017; Störmer et al., 2016). Also, the

authors proposed that involuntary attention to the sound has a "purely facilitatory influence on

processing targets on the cued side, with no inhibitory influence on targets on the opposite side. This

lack of an early inhibitory influence may represent a fundamental difference between involuntary and

voluntary attention" (Feng et al., 2017). No doubt, such a strong assumption based on two very

similar tasks clearly asks for more research exploiting variants of exogenous spatial cueing paradigms

preferrably in different conceptual frameworks to either strengthen or reject this view. Moreover, the

changes in alpha power were accompanied by evoked activity - the so-called auditory-evoked

contralateral occipital positivity - that is known for its link to involuntary spatial attention (McDonald

et al., 2013; Feng et al., 214; Störmer et al., 2019). Critically, both alpha power and evoked activity

shared similarities in the time course, scalp topography and source estimation making it difficult to

interpret whether they reflect the same or two dissociable cognitive mechanisms (Feng et al., 2017;

Störmer et al., 2016).

Current approach

To tackle the two questions within the framework of deviant distraction we employ a unique

crossmodal paradigm that combines a distraction task (Escera et al., 1998; Ruhnau et al., 2013) with

an exogenous spatial cueing task (e.g. Feng et al., 2017; Störmer et al., 2016). Participants were

instructed to respond as fast and as correctly as possible whether a visual target appeared on the left or

right. Each target followed a task-irrelevant sound. Similar to the distraction task, 80% of the sounds

were standards, while 20% of the sounds were deviants. Similar to the exogenous spatial cueing task,

the deviant was non-informative about the spatial position of the upcoming target. That is, 50% of the

deviants occurred at the same side as the target. That condition is referred to as the Congruent

condition. The other 50% of the deviants occurred at the opposite side as the target. That condition is

called the Incongruent condition.

On the behavioral level we intend to reproduce a distraction effect, indicated by longer reaction times

to the visual target that follows a deviant vs. a standard (Escera et al., 1998; Parmentier et al., 2008;

Ruhnau et al., 2013). This behavioral distraction effect can be taken as a sign of an involuntary shift

of attention from the task towards the deviant sound (Parmentier, 2014). Crucially, we hypothesize a

spatial shift of attention, that should be indicated by shorter reaction times in the Congruent vs.

Incongruent condition, i.e. when the visual target follows the auditory deviant at the same vs. different

side.

Besides collecting behavioral data we recorded MEG to shed light on neural oscillatory activity with

an emphasis on alpha power and its link to involuntary spatial attention. We expect the deviant to

induce a lateralized, short-lived alpha power modulation over parieto-occipital sensors that builds up

rapidly after stimulus onset (Feng et al., 2017; Störmer et al., 2016; Weise et al., 2016), thereby

matching the characteristics of involuntary spatial attention (for a review see Corbetta and Shulman,

2002). We predict neural sources of this alpha power modulation to be located in 1) parieto-occipital

areas, known for their role in spatial attention even in purely auditory tasks (Deng et al., 2019; Deng

et al., 2020; Feng et al., 2017; Störmer et al., 2016), 2) auditory areas due to the processing of the

attention-capturing sound (ElShafei et al., 2018; Frey et al., 2014; Weisz et al., 2014; Wöstmann et

al., 2016), as well as 3) areas linked to the ventro-frontoparietal attention system, comprising the

temporoparietal junction and the ventral frontal cortex, that have functionally been linked to

attentional reorienting (for reviews see Corbetta et al., 2008; Vossel et al., 2014).

Method

The study was approved by the local ethics committee of the Paris Lodron Universität Salzburg and

was conducted following the code of Ethics of the World Medical Association (Declaration of

Helsinki).

Participants

30 healthy young adults participated in the MEG experiment (12 males, mean age ± STD: 27 ± 6 yrs,

4 left-handed, 1 ambidexter). Volunteers gave written informed consent prior to their participation.

All participants reported normal hearing, normal or corrected-to-normal vision, and none reported a

history of neurological or psychiatric diseases. Volunteers received 10€/h or 1 credit point/h as

compensation for their participation.

Stimuli and Procedure

The experiment was programmed in MATLAB 9.1 (The MathWorks, Natick, Massachusetts, U.S.A)

using the open source Psychophysics Toolbox (Brainard 1997; Kleiner et al. 2007) and o_ptb, an

additional class-based abstraction layer on top of the Psychophysics Toolbox (https://gitlab.com/thht/

o_ptb). Stimulus presentation and exact timing was ensured by using the VPixx System (DATAPixx2

display driver, PROPixx DLP LED Projector, TOUCHPixx response box by VPixx Technologies,

Canada). The Blackbox2 Toolkit (The Black Box ToolKit Ltd, Sheffield, UK) was used to measure

and correct for timing inaccuracies between triggers and the visual and auditory stimulation. Note that

the tubes of the MEG-proof sound system caused a time-delay between sound and trigger of 16.5 ms

that was corrected within the stimulation protocol.

We employed a cross-modal paradigm combining a distraction task with an exogenous spatial cueing

task. Participants performed a visual-spatial two-alternative forced choice task (see Figure 1).

Throughout the stimulation a frog was presented at the center of the screen that participants were

asked to fixate on at all times. Every trial a visual target in the shape of a fly was presented on the

screen next to the frog. In 50% of the trials the target appeared on the left side of the frog. In the other

50% of the trials it appeared on the right side. The participants' task was to respond via their left or

right thumb and distinct buttons where the fly had been presented. The mapping between the response

buttons and the target location was counterbalanced across participants. Both, the target presentation

and the response window, was maximal 1 s in duration following the onset of the visual target or

ended as soon as the participant responded.

(A) Trial structure and timing

(B) Sound sequence

Left 🔊	S S **D** S S S **D** S S S	**Legend**	
	... ...	**D** Deviant (10% left, 10% right)	
Right 🔊	S S S S S S S **D** S	S Standard (80%, binaural)	

(C) Locations of deviant and target in different conditions

	Congruent Location		Incongruent Location	
Left	**Deviant Target**		**Deviant**	**Target**
Right		**Deviant Target**	**Target**	**Deviant**

Figure 1: Cross-modal paradigm combining a distraction task with an exogenous spatial cueing task.
(A) Trial structure and timing. During each trial the to-be-ignored sound (indicated as an orange bar)
preceded the task-relevant visual stimulus (indicated as a blue bar). Participants had to respond
whether the target occurred at the left or right next to the fixation frog. (B) Exemplary sound
sequence. The task-irrelevant sounds were presented in an oddball paradigm: Standards were
presented binaural on 80% of the trials, deviants were presented left (10%) or right (10%). (C) Listing
of the two different conditions (Congruent Location, Incongruent location) and the corresponding
locations (left, right) of deviant and corresponding target presentation.

Additionally, on each trial a sound of 0.5 s duration (10 ms rise and fall times) was presented that had

to be ignored. This sound always appeared after a 0.8 s silent interval and preceded the visual target

by 0.6 s. That is, a silent interval of 0.1 s separated both the auditory and visual event. The sound was

either a binaurally presented standard (buzzing mosquito, $p = 0.8$) or a monaurally presented deviant.

Deviants were chosen out of a pool of 56 different environmental sounds (e.g., speech, animal voices,

tool noise, etc.; $p = 0.2$) selected from a commercial CD (1,111 Geräusche, Döbeler Cooperations,

Hamburg, Germany). All sounds were equalized for overall intensity (RMS). The trial order within

the whole stimulation was pseudorandomized such that a deviant was preceded by at least two

standards. One auditory-visual trial lasted maximally 2.6 s but was usually shorter (due to the

relatively fast reaction time).

Within one experimental block we presented two different conditions as a function of where the

deviant sound and the visual target appeared. In the Congruent condition both the deviant and target

occurred at the same side, i.e. both appeared on the left or right. In the Incongruent condition, the

deviant and target occurred at opposite sides. That is, while the deviant occurred e.g. on the left the

visual target occurred on the right or vice versa. Noteworthy, each deviant could occur only once

within a block and once every three blocks while it could appear four times in total (twice on the left

[once in the Incongruent and once in the Congruent condition] and twice on the right [once in the

Incongruent and once in the Congruent condition]) with respect to the whole experimental

stimulation.

The experiment consisted of seven blocks in total, each lasting ∼5 min and consisting of 160 trials

(128 standards, 16 deviants presented to the left, 16 deviants presented to the right). Thus, the

stimulation without any breaks lasted maximally ~35 min in total and resulted overall in 112 left and

112 right deviants while half of them occurred in the Congruent condition and the other half in the

Incongruent one. The auditory stimuli were presented at 50 dB above sensation level. The auditory

stimulation was delivered via MEG compatible pneumatic in-ear headphones (SOUNDPixx, VPixx

technologies, Canada). The visual stimuli were presented inside the magnetically shielded room using

a projector (PROPixx DLP LED Projector), via a back projection screen and a mirror system.

Before the start of the experiment the individual sensation level was determined using a 200-ms 1000

Hz tone presented via a staircase threshold measurement procedure separately for the right and left

ear. Thereafter, the participant practiced the task with a fewer number of trials to get familiar with the

experiment. Deviants were chosen randomly from the pool of environmental sounds (N=56). They

were played backwards so that participants did not become familiar with those sounds.

MEG recording and preprocessing

Before MEG recording five head position indicator (HPI) coils were placed at the left and right

mastoid, on the upper right and lower right forehead, and on the left middle forehead. Anatomical

landmarks (nasion and left/right preauricular points), the HPI locations, and at least 300 headshape

points were sampled using a Polhemus FASTTRAK digitizer. Six pre-gelled and self-adhesive

Ag/AgCl electrodes (Ambu Neuroline 720) were attached to record the vertical and horizontal

electrooculogram (EOG) and the electrocardiogram (ECG). One additional electrode - placed at the

right shoulder - served as ground.

During data recording participants were in a seated position. A chin rest was used to avoid head

movements during and across different blocks of recording. The eye, heart and magnetic signals of the

brain were recorded at 1000 Hz (hardware filters: 0.03 - 330 Hz) in a standard passive magnetically

shielded room (AK3b, Vacuumschmelze, Germany) using a whole head MEG (Elekta Neuromag

Triux, Elekta Oy, Finland). Signals were captured by 102 magnetometers and 204 orthogonally placed

planar gradiometers at 102 different positions.

MEG data were preprocessed and analyzed offline using Fieldtrip (Oostenveld et al., 2011), an open

source toolbox for Matlab (www.mathworks.com) and custom-written functions. Maxfilter (version

2.2.15, Elekta Neuromag, Finland) was applied to the continuous MEG raw data using a signal space

separation algorithm (Taulu and Kajola, 2005; Taulu and Simola, 2006) provided by the MEG

manufacturer. This allowed 1) to automatically remove and interpolate the data of bad channels, and

2) to remove artifacts from the MEG signal (50 Hz line-noise and its harmonics, 16.7 Hz train noise,

and muscle activity with origins beyond the head). 3) It also allowed to compensate offline for

different head positions across blocks by realigning the data to a common standard head position (-

trans default Maxfilter parameter) based on the head position measured at the beginning of each

block. Thereafter, continuous data were visually inspected and sample points of extensive artifacts

were stored separately for each block and participant.

Two different pipelines served for further preprocessing of the MEG data. 1) Preprocessing of the

data was optimized for applying independent component analysis (ICA). Therefore, MEG data were

offline filtered using a 100 Hz lowpass filter (sinc FIR, kaiser window, order 3624, cutoff (-6 dB) 100

Hz) and a 1-Hz highpass filter (sinc FIR, kaiser window, order 908, cutoff (-6 dB) 1 Hz). After

rejecting sample points covering extensive artifacts, runica ICA was applied in order to identify

components originating from eye movements and heartbeat. 2) The second preprocessing pipeline was

optimized for further data analysis. In more detail, cleaned MEG data were offline filtered using a 35

Hz lowpass filter (sinc FIR, kaiser window, order 3624, cutoff (-6 dB) 35 Hz). Data were

downsampled to 256 Hz. Thereafter, a 0.1 Hz highpass filter was applied (sinc FIR, kaiser window,

order 9056, cutoff (-6 dB) 0.1 Hz).

Filtered continuous data (originating from the second preprocessing pipeline) were cleaned from the

ICs (on average 2 per participant). Data were segmented into 3-s epochs (-1.5 s to 1.5 s time-locked to

the onset of the sound) and cleaned from artifacts that were identified via visual inspection (see

above). For further analysis, we used deviant trials only. Trials where deviants occurred on the left or

right are referred to as Deviant Left or Deviant Right, respectively, hereafter. Only those deviant trials

entered the analysis for which participants responded once, correctly and within the predefined

response window (100 to 800 ms following the target). This resulted on average in 106.53/112 (SD =

3.38) Deviant Left trials and 106.93/112 (SD = 2.89) Deviant Right trials.

Data analyses

Behavioral Analysis

The following trials were excluded from the analysis of behavioral data: trials without responses,

trials including responses faster than 100 ms or slower than 800 ms after target onset, trials that

included more than one button press. Individual reaction times (RTs) and hit rates were calculated.

Statistical analyses. For the statistical analyses of behavioral data we used jamovi (version 1.2.22.0,

https://www.jamovi.org/). To determine the behavioral distraction effect on RTs and hit rates we

applied two-way repeated-measures analyses of variance (ANOVAs) using the factors sound type

(deviant, standard) and deviant location (left, right). Note that we used standards before the

corresponding deviants for the analyses. To determine the behavioral effect of the spatial shift of

attention on RTs and hit rates we applied two-way repeated-measures analyses of variance

(ANOVAs) using the factors congruence type (congruent, incongruent) and deviant side (left, right).

Note that for statistical purposes hit rates were transformed to rationalized arcsine units (rau) to

account for non-normality of proportion data (Studebaker, 1985). As an effect size measure we report

the generalized eta-squared (η^2_G; Bakeman, 2005). Where appropriate, we reported mean values (M)

and the standard error of the mean (SEM) in the format M ± SEM.

MEG Data Analyses

For the analyses of MEG data in sensor and source space we solely used the signal recorded at the

gradiometers. Data were detrended to remove slow frequency shifts.

Sensor space analyses.

Time-frequency analysis. Time-frequency analysis was performed on the preprocessed single trial

data between 1 and 30 Hz (1 Hz steps) using Morlet wavelets with a width of 5 cycles separately for

each participant and condition. The analysis time window was shifted in steps of 50 ms between -1.5

to 1.5 s relative to deviant onset. Note that power was first computed for each individual gradiometer.

In a second step, the power for each gradiometer pair was summed in order to yield an orientation

independent estimate. Thereafter, single trial spectral power was log transformed with a base of 10.

For each participant, trials were averaged separately for Deviant Left and Deviant Right.

In order to statistically examine alpha power modulations in dependence of the deviant side (Deviant

Left, Deviant Right) in both hemispheres, we performed non-parametric cluster-based dependent-

samples t-tests with Monte-Carlo randomization (Maris and Oostenveld, 2007). The tests were

restricted to the alpha frequency range (8 – 14 Hz) and the time window ranging from 0.2 to 0.6 s

following the deviant sound and preceding the target (cf. Feng et al., 2017; Störmer et al., 2016;

Weise et al., 2016). Since we had a priori hypotheses on the direction of the lateralized alpha power

modulations expected in each hemisphere, we applied paired-sample one-tailed Student's t-tests

separately for the left-hemispheric channels and the right-hemispheric channels. The Student's t-tests

were applied for each sensor-time-frequency bin separately for each hemisphere by testing the alpha

power values of Deviant Left against the alpha power values of Deviant Right. Based thereon clusters

were formed of neighboring sensors of the respective hemisphere and corresponding time-frequency

bins that exceeded an a priori chosen threshold ($p = 0.05$). The sum of the t-values within the

respective cluster was used as cluster-level statistic. The cluster with the maximum sum was

subsequently used as a test statistic. By permuting the data between the two conditions and

recalculating the test statistic 10,000 times, we obtained a reference distribution of maximum cluster t

values to evaluate the significance of the to-be-tested contrast. The power distributions between the

two conditions are considered to be significantly different from each other if not more than 5% of the

sum-t-values of all of the 10,000-permuted data are larger than the sum-t-value of the to-be-tested

cluster.

In addition, we ran the time frequency analysis a second time using the same parameter as described

above but with the preceding step that phase-locked activity, i.e. the evoked signal averaged across

trials of the respective condition, was subtracted from every single trial in order to increase sensitivity

to non-phase locked signals and to reduce contamination by phase-locked signals. The results are

comparable to those obtained via the time frequency analysis of phase-locked activity and can be

found in the supplementary material.

We further tested what caused the alpha power modulation observed in the data (Figure 3). Therefore,

we analyzed the time courses of ipsi- and contralateral alpha power, for the right centro-temporal and

the left parieto-occipital region of interest (ROI), respectively, that had been identified via the

preceding cluster-based permutation analysis (see above). The ROIs consisted each of three channels

that showed prominent deviant-related effects in the deviant–target interval. Data were pooled within

each ROI and across frequencies (right centro-temporal ROI: 8 – 14 Hz, left parieto-occipital ROI: 9-

13 Hz; see results). Alpha power was submitted to non-parametric cluster-based dependent-samples t-

tests with Monte-Carlo randomization (n = 10,000; Maris and Oostenveld, 2007) contrasting the data

at each sample point in the deviant-target interval (0.2 to 0.6 s following the deviant) with the mean

alpha power of the baseline period (-0.4 to -0.2 s; Feng et al., 2017). Since we had no a priori

hypotheses on the direction of the alpha power modulations expected in the respective ROI, we

applied paired-sample two-tailed Student's t-tests.

Source space analyses.

Source projection of time series data. For each participant, realistically shaped, single-shell head

models (Nolte, 2003) were computed by co-registering the participants' head shapes either with their

structural MRI (18 participants) or—when no individual MRI was available (12 participants)—with a

standard brain from the Montreal Neurological Institute (MNI, Montreal, Canada), warped to the

individual head shape. A grid with 3 mm resolution based on an

MNI template brain was morphed to fit the brain volume of each participant. The leadfield was

calculated for each grid point and the forward model was computed for gradiometers only.

To project the preprocessed single-trial sensor data into source space, i.e. to the points of the grid, we

applied the linearly constrained minimum variance (LCMV) spatial filter (Van Veen et al., 1997). We

followed a procedure described for single virtual sensors

(http://www.fieldtriptoolbox.org/tutorial/virtual_sensors/) and extended it to 146400 points covering

the grey matter of the brain. The covariance matrix across all trials (including Deviant Left and

Deviant Right conditions) was calculated and used to obtain a LCMV beamformer spatial filter for

each of the grid points (for a similar approach, see Neuling et al., 2015; Ruhnau et al., 2016). The

covariance window for the calculation of the beamformer filter was based on a 2-s time window

centered at deviant onset. To optimize the analysis in source space (i.e. increase spatial resolution

using a high-definition grid and at the same time compensate for computation time) we divided the

individual brain into 333 parcels (Gordon et al., 2016;

http://www.nil.wustl.edu/labs/petersen/Resources.html) based on the 'Anatomical Automatic

Labeling' [AAL] atlas (Tzourio-Mazoyer et al., 2002; provided by the fieldtrip toolbox). That is, for

each anatomical area, the spatial filters of the corresponding grid points were averaged to obtain one

spatial filter per parcel. The resulting 'averaged filter' was then multiplied with the sensor level data

to obtain time series data for each of the 333 parcels. A regularization parameter (lambda = 5%) was

applied.

Time-frequency analysis. For each participant, time-frequency analysis was performed on the single

trial data in source space between 1 and 30 Hz (1 Hz steps) using Morlet wavelets with a width of 5

cycles. The analysis time window was shifted in steps of 50 ms between -1.5 to 1.5 s relative to

deviant onset. For each participant, trials were averaged separately for Deviant Left and Deviant

Right.

To localize the deviant induced sensor level effects in source space, data were restricted to the time-

frequency ranges covering the alpha power effect (right centro-temporal ROI: 8 – 14 Hz and 0.3 - 0.5

s, left posterior ROI: 9 - 13 Hz and 0.2 - 0.6 s). Alpha power was averaged over frequency and time. T

values were calculated via a paired-sample one-tailed Student's t-test to each of the parcels and the

corresponding time-frequency bin, contrasting the alpha power values between Deviant Left and

Deviant Right. The respective labels of localized brain regions were identified with an anatomical

brain atlas (AAL atlas; Tzourio-Mazoyer et al., 2002) and a network parcellation atlas (Gordon et al.,

2016).

Results

Behavioral data

Task performance in terms of accuracy was in general very good, as indicated by the high hit rate

(>0.97) for targets regardless of whether they were preceded by deviants or standards (Figure 2A).

The outcome of the two-way repeated measures ANOVA on hit rates shows a significant main effect

for factor stimulus type ($F(1, 29) = 18.51$, $p < .001$, $\eta^2_G = .087$), with higher hit rates for deviant trials

(0.99 ± 0.003) than standard trials (0.97 ± 0.004). There was neither a significant main effect for

factor deviant side nor a significant interaction. This result may be linked to an increase in nonspecific

arousal upon the occurrence of a deviant (Näätänen, 1992) that facilitates task performance. Hit rates

did not differ when deviant and target occurred at the congruent vs. incongruent side (Figure 2B) as

indicated by the two-way repeated measures ANOVA yielding neither significant main effects for the

factors congruence type and deviant side nor a significant interaction.

As depicted in Figure 2A, reaction times (RTs) are prolonged when targets follow a deviant vs.

standard. This is substantiated by the two-way repeated measures ANOVA on RTs yielding a

significant main effect for stimulus type ($F(1, 29) = 39.68$, $p < .001$ $\eta^2_G = .037$; deviants: 296 ± 9 ms;

standards: 278 ± 8 ms). There was neither a significant main effect for factor deviant side nor a

significant interaction. Overall, these results point to a behavioral distraction effect due to an

attentional shift towards the task-distracting deviant. As depicted in Figure 2B, RTs are shorter when

deviant and target occurred at the congruent vs. incongruent side. This is substantiated by the two-way

repeated measures ANOVA on RTs yielding a significant main effect for congruence type ($F(1, 29) =$

4.59, $p < .041$ $\eta^2_G = .002$; congruent: 293 ± 9 ms; incongruent: 298 ± 10 ms). There was neither a

significant main effect for factor deviant side nor a significant interaction. Overall, these results point

to a *spatial* shift of attention towards the task-distracting deviant sound.

Figure 2: Bar plots of reaction times and hit rates in response to visual targets to determine a

distraction effect (A) or a shift of spatial attention (B) caused by the deviant. (A) Prolonged RTs and

higher hit rates for targets following deviants presented at the left or right side vs. standard before

deviant. (B) Shortened RTs for targets following deviants at the same vs. different side in the

Incongruent and Congruent condition, respectively. Data were analysed with respect to the deviant

side (left, right). Hit rates did not differ between conditions. SEMs are indicated.

421   **Alpha power analysis**

422   To probe whether attentional reorienting towards the location of the deviant sound, as indexed via the

423   current RT data, is also reflected in neural oscillatory activity we analyzed alpha-band (8 - 14 Hz)

424   power. We followed the methodological approach used for studying voluntary attention (Banerjee et

425   al., 2011; Deng et al., 2020; Haegens et al., 2011; Wöstmann et al., 2016) and contrasted alpha power

426   for the cued / attended target positions (attend left vs. attend right). Accordingly, we here contrasted

427   alpha power for the left and right attention-capturing, task-distracting deviants (Deviant Left condition

428   vs. Deviant Right condition). Since we had a priori hypotheses on the direction of the lateralized alpha

429   power modulations expected in each hemisphere, we applied the corresponding non-parametric

430   cluster-based permutation statistics using paired-sample one-tailed Student's t-tests separately for the

431   left-hemispheric and the right-hemispheric channels. Overall, the results clearly show that alpha

432   power to lateralized deviants is modulated in a spatially selective manner (Figure 3).

433   In more detail, the results for the left hemispheric channels show that alpha power is significantly

434   higher ipsilateral to the deviant location in the time window from about 0.2 s to 0.6 s following the

435   auditory deviant and preceding the visual target (see Fig. 3 A). It mainly covers the frequencies from

436   9 to 13 Hz. This effect is most pronounced over left parieto-occipital sensors ($p = .023$) as highlighted

437   via the scalp distribution of the alpha power modulation for the frequency range 9 - 13 Hz and the

438   time range 0.2 to 0.6 s following the deviant (see Fig. 3 C). The main generators of this sensor level

439   effect are estimated in occipital regions including primary and secondary visual cortex (see Fig. 3 E).

440   The results for the right hemispheric channels show that alpha power, including the frequencies 8 to

441   14 Hz, is significantly lower contralateral to the deviant location in the time window from about 0.3 s

442   to 0.5 s following the auditory deviant and preceding the visual target (see Fig. 3 B). This effect is

443   most pronounced over right centro-temporal sensors ($p = .012$) as highlighted via the scalp

444   distribution of the alpha power modulation for the frequency range 8 - 14 Hz and the time range 0.3 to

445   0.5 s following the deviant (see Fig. 3 D). The main generators of the sensor level effect are estimated

in parieto-occipital, auditory, and somatosensory regions as well as in regions belonging to the ventral

attention system (see Fig. 3 F).

Figure 3: Log10-transformed alpha power modulation to lateralized deviants based on group-

averaged data Deviant Left minus Deviant Right with a focus on the left hemispheric parieto-occipital

effect (left column) and the right hemispheric centro-temporal effect (right column). (A and B) Time

frequency representations highlighting (A) high alpha power ipsilateral to the deviant location at left-

hemispheric posterior gradiometers (average across three channels showing a prominent effect,

marked in C with a star) and (B) low alpha power contralateral to the deviant location at right-

hemispheric centro-temporal gradiometer (average across three channels showing a prominent effect,

marked in D with a star). The black box marks the time-frequency range (time: 0.2 - 0.6 s, frequency:

8 - 14 Hz) used for the statistical cluster analysis. (C and D) Topoplot of the alpha power distribution

(C: 9 -13 Hz, D: 8-14 Hz) in the time window showing a significant alpha power modulation (C: 0.2 -

0.6 s, D: 0.3 - 0.5 s). Stars indicate channels where the statistical effect was prominent. (E and F)

Source estimation of the (E) left hemispheric parieto-occipital alpha power effect and (F) the right

hemispheric centro-temporal alpha power effect.

To test whether the current alpha lateralization (see Fig. 3) is driven by a decrease or increase in

alpha-band power, we analyzed the time courses of alpha power separately for the right hemispheric

centro-temporal and the left hemispheric parieto-occipital region of interest and the corresponding

ipsi- and contralateral deviant. We contrasted the alpha power following the deviant with the mean

alpha power of the baseline period via a non-parametric cluster-based permutation statistic. Since we

had no a priori hypotheses on the direction of the alpha power modulations, we applied paired-sample

two-tailed Student's t-tests. The results show that the right/ipsilateral deviant causes a significant

increase in alpha power in the right-hemispheric centro-temporal ROI in the time range 0.2 to 0.4 s

after sound onset ($p = .017$, see Fig. 4 A). Further, we find that the left/ipsilateral deviant results in a

significant increase in alpha power in the left-hemispheric parieto-occipital ROI in the time range 0.2

to 0.3 s after sound onset ($p = .044$, see Fig. 4 B). Also, the right/contralateral deviant results in a

significant decrease in alpha power in the left-hemispheric parieto-occipital ROI in the time range 0.5

to 0.6 s after sound onset ($p = .044$, see Fig. 4 B)

Figure 4: What drives the alpha power modulations? The ipsilateral deviant conditions lead to a significant increase of alpha power compared with the baseline. Deviant-locked time courses of contra- and ipsilateral log10-transformed and group-averaged alpha power for (A) the right centro-temporal ROI (8 - 14 Hz) and (B) the left parieto-occipital ROI (9 -13 Hz). Red traces show deviant

ipsilateral, blue traces show deviant contralateral. Channels belonging to the ROI are marked with

stars in the corresponding channel layout. Significant time samples are indicated by the horizontal red

and blue traces (*p < .05). Upward directed arrows indicate an alpha power increase, the downward

directed arrow indicates an alpha power decrease. Gray areas indicate the baseline interval (-0.4 to -

0.2 s) and the test interval (0.2 to 0.6 s). The orange and green bar indicate the presentation time of

deviant and target, respectively.

Discussion

Our MEG study aimed to answer two questions concerning the cognitive and neural underpinnings of

behavioral distraction. 1) Does a shift of *spatial* attention due to a lateralized deviant sound contribute

to behavioral distraction? And 2) Does neural oscillatory alpha activity reflect this involuntary *spatial*

attention shift? Our reaction time and neural oscillatory MEG data provide a positive answer to each

of these questions that we will discuss in more detail in the following.

1) The shift of involuntary *spatial* attention contributes to behavioral distraction

As expected, our behavioral distraction effect - i.e. the longer reaction times to targets that followed a

deviant vs. a standard - shows that the deviant captured attention involuntarily from the task. This

strengthens previous findings that have been obtained via a more or less comparable behavioral

distraction task numerous times before (Bendixen et al., 2007; Parmentier et al., 2008; Schröger and

Wolff, 1998; Wetzel et al., 2012).

Importantly, we observed an advantage in response speed when deviants and targets occurred at the

same vs. opposite side. This clearly speaks in favor for a *spatial* shift of attention that has been

proposed though not directly tested earlier (Parmentier et al., 2008; Weise et al., 2016). That is, when

deviant and target occur at the same side the time penalties due to the spatial shift is lower as when

both events occur at opposite sides. This finding is in line with behavioral data obtained via the

exogenous cueing paradigm (McDonald et al., 2000) and receives further support from our MEG data

as will be discussed below.

Note that we do not claim that the spatial component of the attention shift fully accounts for the

overall behavioral distraction effect. In fact, several cognitive determinants may contribute to it. A

second potential determinant that could be at play in a crossmodal situation is the involuntary shift of

attention across sensory modalities, i.e. from vision to audition. If this holds, the time penalty

accumulated by the attentional shift from the visual to the auditory channel (i.e. upon the onset of the

deviant) and its re-orientation towards the visual channel (i.e. upon the onset of the target) would

contribute to behavioral distraction (Parmentier, 2014). The claim receives support from behavioral

studies showing that reaction times are slower for a target in an unexpected modality as opposed to a

target in an expected modality (Boulter, 1977; Spence et al., 2001).

The reaction time data seem to be at odds with the hit rates at a first glance. While the former indicate

a decrease in performance for targets following a deviant vs. standard, the latter rather reflect an

increase. However, this apparent contradiction can be reconciled when linking the response times to

the costs of the attentional shift towards the deviant (Schröger and Wolff, 1998; Parmentier et al.,

2008; for a review, see Parmentier 2014) and the hit rates to an increase in nonspecific arousal that

improves cognitive functioning (Näätänen, 1992). The current findings are well in line with previous

research on deviant distraction showing not only costs but also benefits (SanMiguel et al., 2010;

Wetzel et al., 2012, Wetzel et al, 2019).

2) Oscillatory alpha power reflects an involuntary shift of spatial attention

Our study is the first to show that a lateralized deviant sound modulates alpha power in a spatially

selective way in a narrow time window that cannot be attributed to stimulus-driven evoked activity

(Figure 3). Similarly to recent EEG data collected during an exogenous spatial cueing paradigm (Feng

et al., 2017; Störmer et al., 2016), we observed a lateralized alpha power modulation over parieto-

occipital sensors that occurred relatively early and was rather short-lived, thereby closely matching

the characteristics of involuntary attention in timing and build-up phase (Corbetta and Shulman,

2002). Our whole-brain analyses approach estimated the main sources of the change in the alpha band

in the visual cortex, thus, confirming and extending earlier EEG data (Feng et al., 2017; Störmer et al.,

2016) via an MEG approach.

Of utmost importance, we found a left-hemispheric alpha power modulation over parieto-occipital

sensors that was higher ipsilateral vs. contralateral to the deviant location (Figure 3 A, C). This effect

was driven by an alpha power increase (relative to baseline) due to the left deviant (Figure 4 B). The

current finding is in line with that obtained in endogenous spatial cueing tasks where attention was

voluntarily guided to the location of either a visual (Kelly et al., 2006; Rihs et al., 2009; Worden et al.,

2000) or auditory target (Banerjee et al., 2011; ElShafei et al., 2018; Frey et al., 2014; Müller and

Weisz, 2012). In those tasks lateralized alpha power was larger over the ipsilateral than the

contralateral hemisphere relative to the attended target location. This alpha power modulation was

driven by 1) an alpha power decrease and/or an alpha power increase in the hemisphere contralateral

and/or ipsilateral to the attended side, respectively (e.g. Haegens et al., 2011), 2) a bilateral alpha

power decrease that was maximal over the hemisphere contralateral to the attended position (e.g. Thut

et al., 2006) or 3) a bilateral alpha power increase that was maximal over the hemisphere ipsilateral to

the attended side (e.g. Banerjee et al., 2011; Worden et al., 2000). What pattern is actually observed,

reflecting mechanisms of suppression or enhancement at play in attentional biasing, probably relies on

divergent task demands of the respective study (Kelly et al., 2006; Thut et al., 2006). Anyway, based

on those studies, it has been widely acknowledged that brain regions showing decreased alpha power

reflect the processing of task-relevant information in the voluntarily attended space. Brain regions

with increased alpha power indicate the inhibition of processes associated with task irrelevant

information in the unattended space (for a review, see Klimesch, 2012). In a similar vein our left-

hemispheric parieto-occipital alpha power increase due to the left deviant can be linked to the

suppression of irrelevant information in the unattended space i.e. where deviant-induced involuntary

spatial attention is *not* captured. This active inhibition of visual brain areas could free up resources for

the corresponding counterpart contralateral to the deviant site processing relevant information (Jensen

and Mazaheri, 2010; Meeuwissen et al., 2011) related to the attentional reorienting towards the

deviant location. This interpretation receives support from the current behavioral data, showing that

deviants capture spatial attention (Figure 2B). Further, it is in line with findings suggesting that a

sound can engage visual processing associated with its location (Feng et al., 2014; Feng et al., 2017;

McDonald et al., 2013; Störmer et al., 2016).

Importantly, the current pattern of results contradicts the prevailing view on alpha power in the

context of involuntary attention. Those earlier studies, which used exogenous spatial cueing

paradigms, showed a bilateral alpha power decrease that was more prominent in the hemisphere

contralateral to the task-irrelevant sound. Based thereon, the authors suggested that a shift in

involuntary attention exclusively facilitates the processing of targets on the cued side while the

processing of targets on the opposite side is not negatively impacted (Feng et al., 2017; Störmer et al.,

2016). This conclusion was substantiated by the characteristic pattern of alpha power modulation

preceding correct and incorrect discriminations of valid and invalid targets (Feng et al., 2017). Our

data - the increase of parieto-occipital alpha power - challenges that conclusion as well as the

prevalent notion that the "lack of an early inhibitory influence may represent a fundamental difference

between involuntary and voluntary attention" (Feng et al., 2017). The difference between the current

and previous findings might be due to the fact that the current study did not only apply an exogenous

spatial cueing task but in addition to that a distraction task, thus, resulting in different task demands

(Kelly et al., 2006; Thut et al., 2006).

3) Oscillatory alpha power indicates deviant-related processes beyond involuntary spatial

attention

In addition to the current evidence pointing to a spatial shift of attention, the current   right-

hemispheric alpha power modulation over centro-temporal sites (Figure 3 B, D) gives more insights

into the processing of deviant-related information in a crossmodal scenario. Likewise to the left

hemispheric visual alpha power modulation the right-hemispheric effect was driven by an increase in

alpha power relative to baseline due to the right/ipsilateral deviant (Figure 4 A). This reflects the

inhibition of the underlying brain regions, namely the parieto-occipital, auditory and somatosensory

areas as well as the ventral attention network (Figure 3 F). That active inhibition leads to increased

allocation of resources to their respective contralateral counterparts, responsible for the processing of

the relevant deviant-related information (Jensen and Mazaheri, 2010; Meeuwissen et al., 2011).

Accordingly, the left auditory and somatosensory cortex should be engaged in the processing of the

right deviant or in preparatory processes according to the anticipated behavioral relevance of the right

deviant, respectively. Based thereon, we may draw further conclusions about the nature of the

attentional reorienting in crossmodal scenarios. In that case, the current modulation in auditory alpha

power fits the notion that in crossmodal scenarios attention is not only shifted spatially but also across

modalities, i.e. from vision to audition (Parmentier, 2014; Weise et al., 2016). This line of

argumentation receives support from data pointing to a supramodal attention mechanism that operates

modality independent (Banerjee et al., 2011; Deng et al., 2020; Wöstmann et al., 2016). Taking this

for granted, when attention is shifted from vision to audition, auditory processing may be facilitated

while the processing of the visual target comes at a cost. The latter point fits to our behavioral

findings showing longer reaction times when targets follow a deviant vs. a standard (cf. Ruhnau et al.,

2013; Weise et al., 2016) that we do not exclusively attribute to the spatial shift of attention (see

above).

Moreover, the current modulation of auditory alpha power contributes to the slowly increasing

number of findings highlighting distinct alpha generators in the auditory cortex. So far, this has

mainly been shown when endogenous (spatial) cueing paradigms were employed (Billig et al., 2019;

Frey et al., 2014; ElShafei et al., 2018; Müller and Weisz, 2012; Weisz et al., 2014; Wöstmann et al.,

2016). Importantly, our data extend those results to the field of involuntary spatial attention.

The right-hemispheric alpha power modulation in regions related to somatosensory processing may

point to the allocation of resources in the contralateral counterpart to be ready for action if needed.

Though auditory deviants were not important for performing the task, the brain seems to classify

those sounds as potentially behavioral relevant. This receives support from the engagement of the

ventral attention network that is activated when behaviorally relevant events are detected (Corbetta et

al., 2000). Note that the current somatosensory alpha power modulation cannot be explained by

preparatory processes related to the task for two reasons. 1) Participants responded with both hands to

the left and right target while the button-response assignment was counterbalanced across participants.

2) The alpha power modulation occurs in the pre-target interval while the responding hand was only

determined as soon as the target stimulus appeared.

The fact that we observed a right hemispheric alpha power modulation in areas linked to the ventral

attention network fits to the right hemispheric bias of that network known from mainly fMRI studies

(e.g. Downar et al., 2000; Corbetta et al., 2000; Vossel et al., 2012). Given that this fronto-parietal

network has functionally been linked to stimulus-driven attentional reorienting (for reviews, see

Corbetta and Shulman, 2002; Corbetta et al., 2008), our MEG data establish a strong link between

neural oscillatory activity in the alpha range and involuntary attention. One reason why the current

MEG data show sources in the fronto-parietal network while previous EEG data didn't (Feng et al.,

2017; Störmer et al., 2016) might be due to the different stimulus material being used. Those earlier

studies utilized salient noise bursts, whereas we made use of rare and unpredictable environmental

sounds (e.g. baby crying, telephone ringing, etc.) with a rule-violating character being potentially

behaviorally relevant. In fact, the outcome of fMRI studies suggests that especially behaviorally

relevant events activate the ventral attention network while salient but noninformative events don't

(Downar et al., 2001). Further, the modulation of alpha power in the ventral attention network fits to

the view that this neural network is particularly engaged whenever environmental events call for a

change in the task and especially when these events appear rare as is the case in the current study (for

a review, see Corbetta and Shulman, 2008). Even though the visual targets only but not the deviants

were task-relevant, the deviants clearly put the task related to the targets temporarily on hold - as

shown via the behavioral distraction effect - what can be regarded as a temporary change in the task.

4) Conclusion

We here show that lateralized deviant sounds cause a shift of involuntary spatial attention and

contribute to behavioral distraction. This becomes evident 1) by reaction time data indicating that

responding is faster when the target follows the auditory deviant at the same side as compared to a

situation when both deviant and target occur at different sides. 2) Our MEG data clearly show that

oscillatory alpha power is shaped in a spatially selective manner. Most importantly,

the attention-capturing, task-distracting deviant sounds induced high visual alpha power in the left

hemisphere. These attention-related changes were driven by alpha power increases due to the left

deviant, suggesting that visual areas that process non-relevant information outside the locus of

involuntary attention are disengaged. Together with the behavioral data, our oscillatory data

strengthen the view that alpha power reflects a shift of involuntary spatial attention. Further, they

challenge the prevailing view that involuntary orienting to the location of the sound has - contrary to

voluntary attention - a merely facilitatory impact.

Acknowledgments

We thank Jonas Heilig for his help in collecting and preprocessing the data. We also thank Fred

Seifter for his technical support in the lab.

Funding: This work was supported by the Deutsche Forschungsgemeinschaft (DFG, WE 6085/1-1).

Data and code availability

The preprocessed MEG and behavioral data used to generate the figures and calculate the statistics are

available at the OSF public repository (https://osf.io/qt95u/). The corresponding data analysis code is

available at the corresponding author's GitLab repository (https://gitlab.com/aweise/crossfrog-final-

analysis). The original raw data are available upon reasonable request from the corresponding author.

References

Bakeman, R. (2005). Recommended effect size statistics for repeated measures designs.

Behavior Research Methods, *37*(3), 379–384. https://doi.org/10.3758/bf03192707

Banerjee, S., Snyder, A. C., Molholm, S., & Foxe, J. J. (2011). Oscillatory alpha-band

mechanisms and the deployment of spatial attention to anticipated auditory and visual target

locations: Supramodal or sensory-specific control mechanisms? *The Journal of*

Neuroscience, *31*(27), 9923–9932. https://doi.org/10.1523/JNEUROSCI.4660-10.2011

Bendixen, A., Roeber, U., & Schröger, E. (2007). Regularity extraction and application in

dynamic auditory stimulus sequences. *Journal of Cognitive Neuroscience*, *19*(10), 1664–

1677. https://doi.org/10.1162/jocn.2007.19.10.1664

Billig, A. J., Herrmann, B., Rhone, A. E., Gander, P. E., Nourski, K. V., Snoad, B. F., …

Johnsrude, I. S. (2019). A sound-sensitive source of alpha oscillations in human non-

primary auditory cortex. *Journal of Neuroscience*, *39*(44), 8679–8689.

https://doi.org/10.1523/JNEUROSCI.0696-19.2019

Boulter, L. R. (1977). Attention and reaction times to signals of uncertain modality. *Journal of*

Experimental Psychology. Human Perception and Performance, *3*(3), 379–388.

https://doi.org/10.1037//0096-1523.3.3.379

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Cheal, M. L., & Gregory, M. (1997). Evidence of limited capacity and noise reduction with

single-element displays in the location-cuing paradigm. *Journal of Experimental*

Psychology. Human Perception and Performance, *23*(1), 51–71.

https://doi.org/10.1037//0096-1523.23.1.51

Corbetta, M., Kincade, J. M., Ollinger, J. M., McAvoy, M. P., & Shulman, G. L. (2000).

Voluntary orienting is dissociated from target detection in human posterior parietal cortex.

Nature Neuroscience, *3*(3), 292–297. https://doi.org/10.1038/73009

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in

the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215.

Corbetta, Maurizio, Patel, G., & Shulman, G. L. (2008). The reorienting system of the human

brain: From environment to theory of mind. *Neuron*, *58*(3), 306–324.

https://doi.org/10.1016/j.neuron.2008.04.017

Corral, M.-J., & Escera, C. (2008). Effects of sound location on visual task performance and

electrophysiological measures of distraction. *Neuroreport*, *19*(15), 1535–1539.

https://doi.org/10.1097/WNR.0b013e3283110416

Deng, Y., Choi, I., & Shinn-Cunningham, B. (2020). Topographic specificity of alpha power

during auditory spatial attention. *NeuroImage*, *207*, 116360.

https://doi.org/10.1016/j.neuroimage.2019.116360

Deng, Y., Reinhart, R. M., Choi, I., & Shinn-Cunningham, B. G. (2019). Causal links between

parietal alpha activity and spatial auditory attention. *ELife*, *8*.

https://doi.org/10.7554/eLife.51184

Downar, J., Crawley, A. P., Mikulis, D. J., & Davis, K. D. (2000). A multimodal cortical

network for the detection of changes in the sensory environment. *Nature Neuroscience*,

3(3), 277–283. https://doi.org/10.1038/72991

Downar, J., Crawley, A. P., Mikulis, D. J., & Davis, K. D. (2001). The effect of task relevance

on the cortical response to changes in visual and auditory stimuli: An event-related fMRI

study. *NeuroImage*, *14*(6), 1256–1267. https://doi.org/10.1006/nimg.2001.0946

Dufour, A. (1999). Importance of attentional mechanisms in audiovisual links. *Experimental

Brain Research*, *126*(2), 215–222. https://doi.org/10.1007/s002210050731

ElShafei, H. A., Bouet, R., Bertrand, O., & Bidet-Caulet, A. (2018). Two sides of the same coin:

Distinct sub-bands in the alpha rhythm reflect facilitation and suppression mechanisms

during auditory anticipatory attention. *ENeuro*, ENEURO.0141-18.2018.

https://doi.org/10.1523/ENEURO.0141-18.2018

Escera, C., Alho, K., Schröger, E., & Winkler, I. (2000). Involuntary attention and distractibility

as evaluated with event-related brain potentials. *Audiology and Neuro-Otology*, *5*(3–4),

151–166. http://dx.doi.org/10.1159/000013877

Escera, C., Alho, K., Winkler, I., & Näätänen, R. (1998). Neural mechanisms of involuntary

attention to acoustic novelty and change. *Journal of Cognitive Neuroscience*, *10*(5), 590–

604. http://dx.doi.org/10.1162/089892998562997

Feng, W., Störmer, V. S., Martinez, A., McDonald, J. J., & Hillyard, S. A. (2014). Sounds

activate visual cortex and improve visual discrimination. *The Journal of Neuroscience*,

34(29), 9817–9824. https://doi.org/10.1523/JNEUROSCI.4869-13.2014

Feng, W., Störmer, V. S., Martinez, A., McDonald, J. J., & Hillyard, S. A. (2017). Involuntary

orienting of attention to a sound desynchronizes the occipital alpha rhythm and improves

visual perception. *NeuroImage*, *150*, 318–328.

https://doi.org/10.1016/j.neuroimage.2017.02.033

Frey, J. N., Mainy, N., Lachaux, J.-P., Müller, N., Bertrand, O., & Weisz, N. (2014). Selective

modulation of auditory cortical alpha activity in an audiovisual spatial attention task. *The*

Journal of Neuroscience, *34*(19), 6634–6639. https://doi.org/10.1523/JNEUROSCI.4813-

13.2014

Gordon, E. M., Laumann, T. O., Adeyemo, B., Huckins, J. F., Kelley, W. M., & Petersen, S. E.

(2016). Generation and evaluation of a cortical area parcellation from resting-state

correlations. *Cerebral Cortex*, *26*(1), 288–303. https://doi.org/10.1093/cercor/bhu239

Haegens, S., Händel, B. F., & Jensen, O. (2011). Top-down controlled alpha band activity in

somatosensory areas determines behavioral performance in a discrimination task. *Journal of*

Neuroscience, *31*(14), 5197–5204. https://doi.org/10.1523/JNEUROSCI.5199-10.2011

Hanslmayr, S., Gross, J., Klimesch, W., & Shapiro, K. L. (2011). The role of α oscillations in

temporal attention. *Brain Research Reviews*, *67*(1–2), 331–343.

https://doi.org/10.1016/j.brainresrev.2011.04.002

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity:

Gating by inhibition. *Frontiers in Human Neuroscience*, *4*.

https://doi.org/10.3389/fnhum.2010.00186

Kelly, S. P., Lalor, E. C., Reilly, R. B., & Foxe, J. J. (2006). Increases in alpha oscillatory power

reflect an active retinotopic mechanism for distracter suppression during sustained

visuospatial attention. *Journal of Neurophysiology*, *95*(6), 3844–3851.

https://doi.org/10.1152/jn.01234.2005

Kleiner, M., Brainard, D., Pelli, D. (2007). "What's new in Psychtoolbox-3?" *Perception*, 36,

ECVP Abstract Supplement.

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored

information. *Trends in Cognitive Sciences*, *16*(12), 606–617.

https://doi.org/10.1016/j.tics.2012.10.007

Luck, S. J., Hillyard, S. A., Mouloua, M., & Hawkins, H. L. (1996). Mechanisms of visual-

spatial attention: Resource allocation or uncertainty reduction? *Journal of Experimental*

Psychology. Human Perception and Performance, *22*(3), 725–737.

https://doi.org/10.1037//0096-1523.22.3.725

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data.

Journal of Neuroscience Methods, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Matusz, P. J., Retsa, C., & Murray, M. M. (2016). The context-contingent nature of cross-modal

activations of the visual cortex. *NeuroImage*, *125*, 996–1004.

https://doi.org/10.1016/j.neuroimage.2015.11.016

McDonald, J. J., Teder-Sälejärvi, W. A., & Hillyard, S. A. (2000). Involuntary orienting to sound

improves visual perception. *Nature*, *407*(6806), 906–908. https://doi.org/10.1038/35038085

McDonald, John J., Störmer, V. S., Martinez, A., Feng, W., & Hillyard, S. A. (2013). Salient

sounds activate human visual cortex automatically. *The Journal of Neuroscience*, *33*(21),

9194–9201. https://doi.org/10.1523/JNEUROSCI.5902-12.2013

Meeuwissen, E. B., Takashima, A., Fernández, G., & Jensen, O. (2011). Increase in posterior

alpha activity during rehearsal predicts successful long-term memory formation of word

sequences. *Human Brain Mapping*, *32*(12), 2045–2053. https://doi.org/10.1002/hbm.21167

Müller, N., & Weisz, N. (2012). Lateralized auditory cortical alpha band activity and

interregional connectivity pattern reflect anticipation of target sounds. *Cerebral Cortex*,

22(7), 1604–1613. https://doi.org/10.1093/cercor/bhr232

Näätänen, R. (1992). *Attention and brain function*. Hillsdale, NJ: Erlbaum.

Neuling, T., Ruhnau, P., Fuscà, M., Demarchi, G., Herrmann, C. S., & Weisz, N. (2015).

Friends, not foes: Magnetoencephalography as a tool to uncover brain dynamics during

transcranial alternating current stimulation. *NeuroImage*, *118*, 406–413.

https://doi.org/10.1016/j.neuroimage.2015.06.026

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. *Physics in*

Medicine and Biology, *48*(22), 3637. https://doi.org/10.1088/0031-9155/48/22/002

Palva, S., & Palva, J. M. (2007). New vistas for alpha-frequency band oscillations. *Trends in*

Neurosciences, *30*(4), 150–158. https://doi.org/10.1016/j.tins.2007.02.001

Parmentier, F. B. R. (2014). The cognitive determinants of behavioral distraction by deviant

auditory stimuli: A review. *Psychological Research*, *78*(3), 321–338.

https://doi.org/10.1007/s00426-013-0534-4

Parmentier, F. B. R., Elford, G., Escera, C., Andrés, P., & San Miguel, I. (2008). The cognitive

locus of distraction by acoustic novelty in the cross-modal oddball task. *Cognition*, *106*(1),

408–432. https://doi.org/10.1016/j.cognition.2007.03.008

Rihs, T. A., Michel, C. M., & Thut, G. (2009). A bias for posterior alpha-band power

suppression versus enhancement during shifting versus maintenance of spatial attention.

NeuroImage, *44*(1), 190–199. https://doi.org/10.1016/j.neuroimage.2008.08.022

Ruhnau, P., Herrmann, B., Maess, B., Brauer, J., Friederici, A. D., & Schröger, E. (2013).

Processing of complex distracting sounds in school-aged children and adults: Evidence from

EEG and MEG data. *Frontiers in Psychology*, *4*. https://doi.org/10.3389/fpsyg.2013.00717

Ruhnau, P., Neuling, T., Fuscá, M., Herrmann, C. S., Demarchi, G., & Weisz, N. (2016). Eyes

wide shut: Transcranial alternating current stimulation drives alpha rhythm in a state

dependent manner. *Scientific Reports*, *6*, 27138. https://doi.org/10.1038/srep27138

SanMiguel, I., Linden, D., & Escera, C. (2010). Attention capture by novel sounds: Distraction

versus facilitation. *European Journal of Cognitive Psychology*, *22*(4), 481–515.

https://doi.org/10.1080/09541440902930994

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., …

Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with

human EEG alpha activity. *European Journal of Neuroscience*, *22*(11), 2917–2926. https://

doi.org/10.1111/j.1460-9568.2005.04482.x

Schröger, E. (1996). A neural mechanism for involuntary attention shifts to changes in auditory

stimulation. *Journal of Cognitive Neuroscience*, *8*, 527–539.

http://dx.doi.org/10.1162/jocn.1996.8.6.527

Schröger, E., & Wolff, C. (1998). Behavioral and electrophysiological effects of task-irrelevant

sound change: A new distraction paradigm. *Brain Research. Cognitive Brain Research*,

7(1), 71–87.

Shiu, L., & Pashler, H. (1994). Negligible effect of spatial precuing on identification of single

digits. *Journal of Experimental Psychology: Human Perception and Performance*, *20*(5),

1037–1054. https://doi.org/10.1037/0096-1523.20.5.1037

Spence, C., & Driver, J. (1997). Audiovisual links in exogenous covert spatial orienting.

Perception & Psychophysics, *59*(1), 1–22. https://doi.org/10.3758/BF03206843

Spence, C., Nicholls, M. E. R., & Driver, J. (2001). The cost of expecting events in the wrong

sensory modality. *Perception & Psychophysics*, *63*(2), 330–336.

https://doi.org/10.3758/BF03194473

Störmer, V., Feng, W., Martinez, A., McDonald, J., & Hillyard, S. (2016). Salient, irrelevant

sounds reflexively induce alpha rhythm desynchronization in parallel with slow potential

shifts in visual cortex. *Journal of Cognitive Neuroscience*, *28*(3), 433–445.

https://doi.org/10.1162/jocn_a_00915

Störmer, V. S. (2019). Orienting spatial attention to sounds enhances visual processing. *Current*

Opinion in Psychology. https://doi.org/10.1016/j.copsyc.2019.03.010

Studebaker, G. A. (1985). A „rationalized" arcsine transform. *Journal of Speech and Hearing*

Research, *28*(3), 455–462. https://doi.org/10.1044/jshr.2803.455

Taulu, S., & Simola, J. (2006). Spatiotemporal signal space separation method for rejecting

nearby interference in MEG measurements. *Physics in Medicine and Biology*, *51*(7), 1759–

1768. https://doi.org/10.1088/0031-9155/51/7/008

Taulu, Samu, & Kajola, M. (2005). Presentation of electromagnetic multichannel data: The

signal space separation method. *Journal of Applied Physics*, *97*(12), 124905. https://doi.org/

10.1063/1.1935742

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). Alpha-band

electroencephalographic activity over occipital cortex indexes visuospatial attention bias

and predicts visual target detection. *Journal of Neuroscience*, *26*(37), 9494–9502.

https://doi.org/10.1523/JNEUROSCI.0875-06.2006

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., …

Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a

macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*,

15(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Van Veen, B. D., van Drongelen, W., Yuchtman, M., & Suzuki, A. (1997). Localization of brain

electrical activity via linearly constrained minimum variance spatial filtering. *IEEE

Transactions on Bio-Medical Engineering*, *44*(9), 867–880.

https://doi.org/10.1109/10.623056

Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and ventral attention systems: Distinct

neural circuits but collaborative roles. *The Neuroscientist*, *20*(2), 150–159.

https://doi.org/10.1177/1073858413494269

Vossel, S., Weidner, R., Driver, J., Friston, K. J., & Fink, G. R. (2012). Deconstructing the

architecture of dorsal and ventral attention systems with dynamic causal modeling. *Journal

of Neuroscience*, *32*(31), 10637–10648. https://doi.org/10.1523/JNEUROSCI.0414-12.2012

Weise, A., Hartmann, T., Schröger, E., Weisz, N., & Ruhnau, P. (2016). Cross-modal distractors

modulate oscillatory alpha power: The neural basis of impaired task performance.

Psychophysiology, *53*(11), 1651–1659. https://doi.org/10.1111/psyp.12733

Weisz, N., Müller, N., Jatzev, S., & Bertrand, O. (2014). Oscillatory alpha modulations in right

auditory regions reflect the validity of acoustic cues in an auditory spatial attention task.

Cerebral Cortex, *24*(10), 2579–2590. https://doi.org/10.1093/cercor/bht113

Wetzel, N., Scharf, F., & Widmann, A. (2019). Can't Ignore—Distraction by task-irrelevant

sounds in early and middle childhood. *Child Development*, *90*(6), e819–e830.

https://doi.org/10.1111/cdev.13109

Wetzel, N., Widmann, A., & Schröger, E. (2012). Distraction and facilitation—Two faces of the

same coin? *Journal of Experimental Psychology. Human Perception and Performance*,

38(3), 664–674. https://doi.org/10.1037/a0025856

Worden, M. S., Foxe, J. J., Wang, N., & Simpson, G. V. (2000). Anticipatory biasing of visuospatial attention indexed by retinotopically specific alpha-band electroencephalography increases over occipital cortex. *The Journal of Neuroscience*, *20*(6), RC63.

Wöstmann, M., Herrmann, B., Maess, B., & Obleser, J. (2016). Spatiotemporal dynamics of auditory attention synchronize with speech. *Proceedings of the National Academy of Sciences*, *113*(14), 3873–3878. https://doi.org/10.1073/pnas.1523357113