5

Basal epidermis collective migration and local Sonic hedgehog signaling promote skeletal branching morphogenesis in zebrafish fins

Joshua A. Braunstein[1, 2, #], Amy E. Robbins[1, 2, #], Scott Stewart[1], Kryn Stankunas[1, 2, *]

Institute of Molecular Biology
Department of Biology
20 University of Oregon
273 Onyx Bridge
1318 Franklin Blvd
Eugene, OR 97403-1229
Office: (541) 346-7416
25 Fax: (541) 346-4854

* Correspondence: kryn@uoregon.edu

# Equal contributions

ABSTRACT

Adult zebrafish fins develop and robustly regenerate an elaborately branched bony ray skeleton. During caudal fin regeneration, basal epidermal-expressed Sonic hedgehog (Shh) locally promotes ray branching by partitioning pools of adjacent progenitor osteoblasts (pObs). We investigated if and how Shh signaling similarly functions during developmental ray branching. As during regeneration, *shha* is uniquely expressed by basal epidermal cells (bEps) overlying pOb pools at the distal aspect of outgrowing juvenile fins. Lateral splitting of each *shha*-expressing epidermal domain followed by the pOb pools precedes overt ray branching. We use *ptch2:Kaede* fish and Kaede photoconversion to identify short stretches of *shha+* bEps and tightly juxtaposed pObs as the active zone of Hh/Smoothened (Smo) signaling. Basal epidermal distal collective cell migration continuously replenishes each *shha+* domain with individual cells transiently expressing and responding to Shh. In contrast, pObs maintain Hh/Smo activity until differentiating. Hh/Smo inhibition using the small molecule BMS-833923 (BMS) prevents branching in all fins, paired and unpaired, with minimal effects on fin outgrowth or skeletal differentiation. Staggered addition of BMS indicates Hh/Smo signaling acts throughout the branching process. We use live time-lapse imaging and cell tracking to find Hh/Smo signaling restrains the distal movement of bEps by apparent 'tethering' to pObs. We conclude short-range Shh/Smo signaling enables ray branching by re-positioning pObs during both fin development and regeneration. We propose instructive basal epidermal collective migration and Shh/Smo-promoted heterotypic cell adhesion between bEps and pObs underlies this unique branching morphogenesis.

INTRODUCTION

Teleost fish fins have elaborately patterned skeletons comprised of bony rays, or lepidotrichia, that shape and structurally support fin appendages. The diversity and beauty of fish fins has long captivated aquarists while serving as a compelling model to consider morphological evolution. *Danio rerio* zebrafish are widely studied ray-finned teleost fish with branched rays in all paired and unpaired fins. For example, zebrafish caudal fins typically have 18 rays of which the central 16 branch into "daughter" rays (Figure 1 Supplement 1A). Juvenile fish form primary branches around 30 days post fertilization (dpf) followed by secondary and tertiary branches. Individual rays comprise two opposed hemi-rays that form cylindrical skeletal units segmented by joints and enveloped by a multilayered epidermis (Figure 1 Supplement 1B). Teleost fins and tetrapod vertebrate limbs evolved from a common ancestral appendage (Dahn et al., 2006; Freitas et al., 2006) with rays possibly sharing deep homology with digits (Nakamura et al., 2016). The relative simplicity of the fin's skeletal structure makes zebrafish a valuable model for understanding mechanisms of appendicular skeletal patterning.

Zebrafish fully regenerate adult fins including restoring a branched ray skeleton within two weeks of injury. Empowered by versatile genetic and other tools, zebrafish have become a leading model for appendage regeneration research. Collective studies implicate many of the same signaling pathways, including Wnt, Bmp, Fgf, and Hh, involved in tetrapod limb development (Reviewed in Sehring et al., 2019). Therefore, zebrafish fin development and regeneration provide complementary contexts to understand how cell signaling patterns the appendicular skeleton and how the same pathways are reactivated for repair. Further, zebrafish fins and their rays enable studies of fundamental developmental questions, including how

75   branched networks form – a common property of many organs including vasculature, lungs, kidneys, mammary glands, and pancreas.

Sonic hedgehog (Shh) signaling through its Smoothened (Smo) effector is one pathway dually involved in fin regeneration and appendage development. Shh is associated closely with

80   tetrapod limb skeletal patterning (Zuniga, 2015) and Shh/Smo pathway perturbations cause syndactyly and polydactyly (Anderson et al., 2012; Malik, 2012). Shh is the Zone of Polarizing Activity (ZPA) secreted morphogen that pre-patterns the limb field into distinct skeletal units, including digits (Chiang et al., 2001; Riddle et al., 1993; Saunders & Gasseling, 1968). Zebrafish *shha* expression studies suggest a ZPA patterns pectoral and other paired fins but not unpaired

85   fins including the caudal fin (Hadzhiev et al., 2007; Laforest et al., 1998). Nevertheless, *shha* is expressed in distal epidermal domains overlying each forming ray during caudal fin development and regeneration (Armstrong et al., 2017; Hadzhiev et al., 2007; Laforest et al., 1998; Y. Lee et al., 2009; Zhang et al., 2012). At least during regeneration, each *shha*-expressing domain splits prior to ray branching. We leveraged the highly specific Smo inhibitor BMS-833923 (BMS) to

90   show Shh/Smo specifically promotes ray branching during zebrafish fin regeneration (Armstrong et al., 2017). However, a potential Shh/Smo role during developmental ray branching and underlying mechanisms are unresolved.

During caudal fin regeneration, distal-moving basal epidermal cells (bEps) adjacent to

95   bony rays upregulate *shha* at the distal "progenitor zone" (Armstrong et al., 2017). Shh-expressing bEps activate Hedgehog/Smoothened (Hh/Smo) signaling in themselves and immediately adjacent progenitor osteoblasts (pObs) as marked by *patched2* (*ptch2*), which

encodes a Hh receptor and universal negative feedback regulator (Alexandre et al., 1996; Goodrich et al., 1996; Lorberbaum et al., 2016; Marigo et al., 1996). This short-range Hh/Smo signaling is required to split pOb pools and therefore ray branching without impacting proliferation or differentiation (Armstrong et al., 2017). While Hh/Smo signaling is active continuously, *shha*-expressing basal epidermal domains split laterally prior to pOb partitioning. We proposed Shh/Smo signaling might enhance physical associations between moving epidermal cells and pObs to enable the progressive partition of pOb pools (Armstrong et al., 2017).

Here, we explore the mechanisms underlying developmental fin ray branching in juvenile zebrafish. We show basal epidermal dynamics as well as Shh/Smo activity and function are largely the same as during regeneration. We use transgenic reporter lines for *shha* and its target gene *ptch2* to refine developmental expression profiles. Kaede photoconversion of *TgBAC(ptch2:Kaede)* fish reveals continuous Shh/Smo signaling in distal ray Shh+ basal epidermal domains and neighboring pObs. We inhibit Shh/Smo signaling using the small molecule BMS-833923 to show the pathway is largely dedicated to ray branching in all fins, including the paired pectoral fins. Shh+ bEps and pObs are closely apposed at the site of Shh/Smo signaling where a basement membrane is incompletely assembled. bEps constantly move distally, trafficking through while contributing to *shha*-expressing domains that split laterally prior to ray branching. We use live time-lapse imaging to demonstrate Shh/Smo signaling restrains basal epidermal distal migration, possibly by promoting transient adhesion of *shha*+ bEps to distal ray pObs. We conclude the collective migration of bEps, constantly distal with progressive lateral domain splitting, and their atypical use of local Shh/Smo signaling re-

positions pObs for skeletal branching during both fin development and regeneration. This reflects a unique branching morphogenesis process whereby movements of a neighboring cell type – the bEps – guides the tissue-forming cells – the pObs – into split pools.

    RESULTS

Shha expression is progressively restricted to distal ray basal epidermal domains that split preceding ray branching

Developing fins express *sonic hedgehog a* (*shha*) in single basal epidermal domains adjacent to the tip of each hemi-ray up to 20 days post fertilization (dpf) (Hadzhiev et al., 2007;

130   Laforest et al., 1998). During caudal fin regeneration, similar *shha*-positive basal epidermal domains split into distinct domains around 4 days post amputation (dpa) immediately preceding ray branching (Armstrong et al., 2017; Laforest et al., 1998; Y. Lee et al., 2009; Quint et al., 2002; Zhang et al., 2012). We expected a comparable pattern during fin development if branching during fin regeneration recapitulates developmental processes. We used the reporter

135   line *Tg(-2.4shha:gfpABC)^{sb15}*, which mimics endogenous *shha* expression (*shha:GFP*; Ertzer et al., 2007; Zhang et al., 2012), to monitor *shha* expression in live zebrafish from its emergence through primary ray branching in juveniles around 30 dpf. *shha:GFP* expression in the caudal region was restricted to the developing floor plate and notochord through 5 dpf, as previously established (Krauss et al., 1993). From 5-9 dpf, *shha:GFP* expression expanded into the caudal

140   fin fold primordium through a ventral gap of melanophores (Figure 1A). By 9-10 dpf, *shha:GFP+* cells were enriched along emerging immature rays (Figure 1B). At 10-12 dpf, *shha:GFP* expression became specifically associated with the distal aspect of maturing central rays while remaining along the lengths of immature peripheral rays (Figure 1C). *shha:GFP* was restricted to ray tips by 14 dpa, when all rays contained maturing, segmented bone (Figure 1D).

The *shha:GFP* domains began splitting around 30 dpf, immediately prior to branching of the corresponding ray (Figure 1I, I', white brackets). We used whole mount immunostaining for

GFP and the basal epidermal marker Tp63 (H. Lee & Kimelman, 2002) and 3D confocal reconstructions to confirm *shha*-driven GFP was expressed exclusively in basal epidermal cells (bEps) (Figure 1K, L; expanded data in Figure 1 Supplement 2). Where Tp63-marked cells were multi-layered (magenta dashed line, Figure 1K), only the innermost cells expressed GFP. Similar *shha* expression patterns, including *shha+* bEp domain splitting, support a common Shh/Smo-dependent mechanism for both developmental and regenerative ray branching.

Zebrafish have 3 unpaired (dorsal, anal, and caudal) and 4 paired (pectoral and pelvic) fin appendages, each with a branched dermoskeleton. We explored if *shha:GFP* domains split ahead of developmental ray branching in all fins. We imaged dissected fins from 51 dpf juvenile fish carrying *shha:GFP* and *Tg(runx2:mCherry)* (*runx2:mCherry*; Shannon Fisher Lab, unpublished), which marks *runx2+* pre-osteoblasts (pObs) and then perdures in differentiated Obs along ray lengths. *shha:GFP* was expressed at and slightly distal to the *runx2:mCherry+* pOb-defined fin boundary in all seven fins, including paired pectoral and pelvic fins (Figure 1 Supplement 3). In all cases, branching rays were tipped with two distinct *shha:GFP+* domains, suggesting shared Shh/Smo-promoted ray branching regardless of fin evolutionary or morphological divergence.

Ptch2 expression indicates Hh/Smo activity in basal epidermis and progenitor osteoblasts

We next assessed expression of *ptch2*, a Hh/Smo negative feedback regulator and activity marker (Alexandre et al., 1996; Goodrich et al., 1996; Lorberbaum et al., 2016; Marigo et al., 1996). Hh/Smo signaling induces *ptch2* in pObs and neighboring bEps during caudal fin regeneration (Armstrong et al., 2017; Quint et al., 2002). *ptch2* is also expressed in distal regions of late larval caudal fins, although cell-level expression is unresolved (Laforest et al., 1998). We

imaged the *TgBAC(ptch2:kaede)^{a4596}* reporter line (*ptch2:Kaede*; (Huang et al., 2012) at the same developmental time points we examined *shha:GFP*. *ptch2:Kaede* expression was confined to the notochord and floor plate until ~9 dpf (Figure 1E). By 9-10 dpf, *ptch2:Kaede* was associated with nascent rays in the ventrally expanding fin primordial (Figure 1F). At 10-12 dpf, *ptch2:Kaede* was detected the entire length of each ray (Figure 1G). This pattern, which includes past *ptch2* expression due to prolonged Kaede stability, persisted through 14 dpf with notably higher *ptch2:Kaede* expression associated with joints and distal ray tips, matching its pattern in regenerating fins ((Figure 1H; Armstrong et al., 2017). As with *shha:GFP*, *ptch2:Kaede+* domains split as ray branching initiated in 30-33 dpf juvenile fish (Figure 1J, J').

To define the cell types expressing *ptch2* during caudal fin development, we combined *ptch2:Kaede* with *runx2:mCherry* and *shha:GFP* to mark *runx2+* pObs and *shha+* bEps, respectively (Figure 1M-P and expanded data with single channel images in Figure 1 Supplement 4). Confocal imaging of live 19 dpf double transgenic larval fish showed *ptch2:Kaede* co-localized with distal fin *runx2:mCherry*-expressing pObs (Figure 1M, N). Only adjacent and further distal presumptive bEps additionally expressed *ptch2:Kaede*. We used photoconversion to discern Kaede from GFP and reveal non-pOb *ptch2:Kaede* co-localized with *shha:GFP*-expressing and nearby bEps (Figure 1O, P). Therefore, *ptch2* defines autocrine (in bEps) and short-range (in pObs and non-*shha+* bEps) Shh/Smo signaling during caudal fin development.

Active Shha/Smo signaling is restricted to outgrowing distal ray regions

We photoconverted distal ray ends of 25 dpf *ptch2:Kaede* caudal fins and re-imaged 24 hours later to distinguish actively produced Kaede from perduring reporter fluorescence (Figure 2A-C). New, unconverted Kaede was produced exclusively in short, discrete domains at the ray tips. Therefore, active Shh/Smo signaling appears narrowly focused in close proximity with *shha*-expressing bEps. We also observed a stretch of photoconverted Kaede+ cells distal to the ray tips in tissue newly formed over the 24-hour post-conversion period. We observed the same pattern during fin regeneration, suggestive of distal migration of previously Shh/Smo-responsive bEps that ceased *ptch2* expression when moving beyond the Shh/Smo active zone (Armstrong et al., 2017).

We used the potent Smo inhibitor BMS-833923 (Armstrong et al., 2017; Lin & Matsui, 2012; henceforth abbreviated BMS) to confirm *ptch2:Kaede* reports Shh/Smo activity during fin development. As expected, caudal fins of 25 dpf fish treated with 1.25 µM BMS and photoconverted 3 hours later produced no new Kaede 24 hours post-conversion (hpc; Figure 2 Supplement 1G-L). Curiously, we no longer observed distally displaced photoconverted Kaede+ bEps and the photoconverted Kaede associated with ray tips was reduced (Figure 2 Supplement 1F', L'). We surmise the 24-hour Shh/Smo-inhibition led distal-moving Kaede+ bEps to shed prematurely, leaving only photoconverted Kaede+ pObs. As such, Shh/Smo signaling may retard distal bEp collective movements.

Sustained Shh/Smo signaling promotes ray branching during fin development

We next investigated if Shh/Smo is required for ray branching during fin development as during regeneration (Armstrong et al., 2017). Treating *Tg(sp7:EGFP)^{b1212}* osteoblast reporter

fish (*sp7:EGFP*; (DeLaurier et al., 2010) with 0.63 µM BMS from 25 to 42 dpf blocked caudal fin ray branching (Figure 3A-D, *n*=5 per BMS and DMSO control groups). In contrast, Shh/Smo-inhibition did not disrupt caudal fin outgrowth or skeletal maturation, indicated by well-defined cylindrical bony ray segments complete with joints, of the central 16 rays (Figure 3 Supplement 1). Curiously, the non-branching principal peripheral rays uniquely were shorter in BMS-treated fish (Figure 3A, C white arrows and Figure 3 Supplement 1). BMS-treatment of 29 dpf *shha:GFP* fish exposed to EdU for 12 hours did not change the fraction of EdU+ intra-ray cells, i.e. pObs and mesenchyme nestled between the epidermal Shh domains of each hemi-ray (Figure 3 Supplement 2, *n*=5 per group). We conclude Shh/Smo signaling is largely dedicated to ray branching with minimal proliferation or bone maturation effects during both fin development and, as shown previously, regeneration (Armstrong et al., 2017).

Shh/Smo signaling may act transiently to initiate ray branching or continuously during the branching process. To distinguish between these possibilities, we staggered the start of BMS treatment to "before", "during", or "after" branching (Figure 3E), identified by *a priori* screening 24-35 dpf *sp7:EGFP* clutchmate fish. Expectedly, BMS-exposure initiated prior to ray branching prevented said rays from branching ("before" group, Figure 3F, G, G', *n*=4/4) and rays that had already fully branched remained so after BMS treatment ("after" group, J, K, K', *n*=4/4). However, rays that recently initiated branching ("during" group, Figure 3H) re-fused upon BMS exposure, forming "gapped" ray segments (Figure 3I, I', *n*=4/5). Therefore, sustained Shh signaling acts throughout ray branching morphogenesis rather than as a switch that initiates branching.

Shh/Smo signaling does not substantially contribute to initial fin ray patterning

240   *shha* and *ptch2* expression during the initial stages of ray formation (Figure 1B, E)

suggest Shh/Smo influences early fin skeletal patterning in addition to promoting later, juvenile-

stage ray branching. To explore this possibility, we inhibited Shh/Smo signaling from as early as

2 dpf, when the larval fin fold entirely comprises soft tissue absent of any ray structures. As

expected, *ptch2:Kaede*-marked Shh/Smo signaling was restricted to the notochord and floor plate

245   (Figure 3 Supplement 3A-C). Photoconversion experiments confirmed BMS fully inhibited

production of new *ptch2:Kaede* in 14 dpf larval caudal fins (Figure 3 Supplement 3D-E', total

n=33-44 per group), as with embryos, juvenile fins, and regenerating adult fins (Figure 2

Supplement 1G-L; (Armstrong et al., 2017). We treated *sp7:EGFP;runx2:mCherry* fish with

1.25 µM BMS from 2 until 14 dpf, when all 18 rays were clearly established. Their caudal fins

250   developed the correct complement of 18 rays with apparently normal length distribution across

the dorsal-ventral axis to define typical "V-shaped" fins (Figure 3 Supplement 3F-G'). As such,

Shh/Smo may have only minor or no roles in initial caudal fin skeletal patterning.

Shh/Smo signaling promotes ray branching in all fins

255   We tested if Shh/Smo signaling promotes ray branching in all seven fins by treating

shha:GFP;runx2:mCherry fish with BMS starting at 21 dpf, prior to asynchronous ray branching

across fins. As expected, both DMSO control and BMS-treated fish showed *shha:GFP+*

domains at the distal end of every ray of all fins at 42 dpf (Figure 4). However, BMS-treated fish

showed no or, at best, severely delayed branching in all fins (*n*=6 per group). The rare, delayed

260   branching likely reflects incomplete Shh/Smo inhibition due to the challenge of sustaining

effective drug concentrations across a multi-week treatment. Nevertheless, the near absence of branchpoints demonstrates all fins employ a common Shh/Smo signaling-dependent mechanism.

Shha+ bEps and pObs are intimately associated in developing caudal fins

265   We next aimed to identify how sustained, local Shh/Smo signaling affects bEp and/or pOb cell behaviors to promote ray branching morphogenesis. The close proximity of these two Shh-responsive cell types suggested their movements might be physically coupled in a Shh/Smo-dependent manner to promote branching. To assess potential physical contacts between bEps and pObs, we first stained longitudinal sections of 32 dpf juvenile fin rays from *shha:GFP* fish with

270   GFP, the osteoblast marker Zns-5, and Laminin, a component of the epidermal-osteoblast separating basement membrane. As expected, Shha:GFP+ bEps were directly adjacent to pObs (Figure 5 Supplement 1). A thin Laminin-containing basement membrane separated pObs and the proximal-most Shha:GFP+ bEps that had recently arrived in the active zone and initiated *shha* expression. More distally, the double staining for Shha:GFP+ bEps and Zns-5+ pObs

275   produced even partially overlapping signal (Figure 5 Supplement 1D and D'), suggesting the two cell types are intimately associated. Here, the laminin+ basement membrane was less dense and sometimes fragmented, likely reflecting its nascent production (asterisks, D').

        We further explored the relative positioning of bEps and pObs at the onset of ray

280   branching by 3D confocal reconstructions of live imaged fins of 28 dpf *shha:GFP;runx2:mCherry* fish (Figure 5A-C). *shha:GFP+* bEps and *runx2:mCherry+* pObs were tightly juxtaposed in both hemi-rays of a single lepidotrichia (Figure 5 Movie 1). Focusing on one hemi-ray, we observed extensive apparent heterotypic surface contacts, including areas

where *shha:GFP+* bEps enshrouded a ridge of pObs (Figure 5 Movie 2). Single sagittal optical slices and reconstructed slice equivalents examined multi-dimensionally (Figure 5D-F) showed intertwined bEps and pObs unresolvable by conventional confocal microscopy.

We considered if Shh/Smo signaling promotes the juxtaposition of bEps and pObs. However, BMS treatment of *shha:GFP;runx2:mCherry* fish from 24-34 dpf did not alter the intimate association between Shh+ bEps and Runx2+ pObs in static images of live fins even though the same drug exposure prevented ray branching to 42 dpf (Figure 5 Supplement 2). As expected, Runx2+ pOb pools failed to split upon BMS exposure, even when *shha:GFP* domains still did. The *shha:GFP* domains of BMS-treated fish variably appeared to remain as one cluster per hemi-ray (Figure 5 Supplement 2E, J; 4/10 split, 6/10 un-split for dorsal ray 3). In contrast, our fin regeneration study indicated Shha-domain splitting is always Shh/Smo-independent (Armstrong et al., 2017). The difference could reflect difficulty visualizing the initial splitting of the smaller *shha:GFP* domains of juvenile fins. Further, constant *shha* production throughout fin growth, not just at branching, seemingly precludes a direct Shh/Smo role in periodically splitting bEp domains. Regardless, Shh/Smo signaling does not support ray branching by promoting close proximity between bEps and pObs per se.

Shh/Smo signaling restrains basal epidermal collective movements while adjacent to pObs

Shh/Smo inhibition appeared to increase the rate of bEp shedding due to accelerated distal collective movements (Figure 2 Supplement 1I, L). Therefore, we considered if Shh/Smo promotes transient adhesion (direct or indirect) between bEps and pObs that impedes bEp movements while they neighbor relatively stable pObs. Such regulated heterotypic adhesion,

which may not be evident by static imaging, coupled with force-generating bEp collective movements during *shha+* domain splitting, could re-position pOb pools over time. We time-lapse imaged caudal fins of *ptch2:Kaede* fish at late larval stages (22-24 dpf) to assess heterotypic cellular dynamics in outgrowing rays. *ptch2:Kaede+* bEps moved distally over *ptch2:Kaede+* pObs, which remained stationary over the 30 minute imaging period, and in more distal fin tissue (Figure 6 Movie 1, Figure 6 Supplement 1A-C). We used semi-automated tracking of individual *ptch2:Kaede+* bEps to determine *ptch2:Kaede+* bEps of BMS treated fish moved significantly faster (3-6 cells per fish and *n*=8 fish per group) (Figure 6 Supplement 1D). Therefore, Shh/Smo signaling restrains the distal movement of Shh/Smo-responsive bEps.

We imaged caudal fins of *shha:GFP*;*runx2:mCherry* larval fish with and without Shh/Smo inhibition to monitor movement dynamics of *shha:GFP*-expressing bEps relative to Runx2+ pObs (representative fish in Figure 6A-D' and Figure 6 Movie 2; all fish shown in Figure 6 Supplement 2). We resolved individual cells at higher detail in distal ray regions by capturing full confocal z-stacks every 2 minutes over 30 minutes. Assisted by semi-automated cell tracking, we noted bEps moved faster when beyond the field of pObs in DMSO-treated control fins. We observed slow moving bEps in contact with pObs before detaching and rapidly moving distally. In contrast, BMS exposure caused rapid distal bEp movement irrespective of proximal-to-distal position or proximity to pObs.

We quantified positional dynamics of individual bEps and plotted their average normalized speed compared to starting position relative to the end of ray-forming Runx2+ pObs (DMSO: *n*=5 fish, 26-38 cells per fish, total of 159 cells; BMS: *n*=4 fish, 26-41 cells per fish,

330   total of 135 cells). *shha:GFP*-expressing bEps located distal to pObs moved faster than pOb-associated bEps in control animals, producing a clear upward velocity shift at the pOb border (Figure 6E). In contrast, BMS treatment caused evenly distributed bEp velocities before and after the pOb-containing region. Taken together, we propose local Shh/Smo signaling enhances heterotypic cell adhesion that transiently restrains the continuous distal movement of bEps when

335   they pass directly over pObs. For ray branching morphogenesis, Shh/Smo-enhanced cell adhesion between pObs and successive waves of bEps could enable the pOb pool to gradually follow laterally splitting *shha*-expressing bEp domains. Eventually, the divided pOb pools would then form separate daughter rays connected at a branch point.

340   **DISCUSSION**

Basal epidermal movements and Shh/Smo signaling direct skeletal branching morphogenesis during zebrafish fin ray development and regeneration

Zebrafish fin ray branching provides an accessible context to define mechanisms of appendage patterning and skeletal morphogenesis. Our current and earlier study (Armstrong et

345   al., 2017) extends previous research to demonstrate the same Shh-dependent branching morphogenesis mechanism branches developing and regenerating rays. In both contexts, a gradually splitting domain of Shh-expressing basal epidermal cells (bEps) at the distal aspect of each outgrowing fin ray partitions the immediately adjacent pre-osteoblast (pOb) pool. Highly localized, continuous Shh/Smo activity allows a given ray's distal pOb population to gradually

350   follow the separating Shha+ basal epidermal domains. Eventually fully split, divided pOb pools continue to promote outgrowth of now two rays connected at a branch point. The shared mechanism of pOb positioning for ray branching underscores that fin regeneration re-activates developmental mechanisms. Strikingly, Smo-dependent Shh signaling appears largely dedicated to ray branching in all fins, unpaired and paired, with pathway inhibition producing minimal or

355   no effects on initial fin patterning, outgrowth, or skeletal differentiation during fin development or regeneration.

Our study of the readily observable developing caudal fin highlights how collective migration of Shha+ bEps positions pObs to generate branched rays. Our Kaede photoconversion

360   and time-lapse imaging show bEps continuously move distally in developing fins, activating *shha* expression upon reaching the distal zone that includes pObs. Individual bEps pass through the *shha*-expressing domain, down-regulate *shha* when moving beyond the pObs, and then are

seemingly shed from the end of the fin. In turn, proximal bEps enter the distal zone and activate *shha* to replenish the *shha*-expressing basal epidermal domain adjacent to pObs. Shha produces a

365   constant Smo-dependent response in neighboring pObs and an autocrine, transient response in *shha*-producing bEps as represented by upregulated *ptch2* in both cell types. This continuous, localized Hh/Smo signaling restrains bEp collective movement dynamics and promotes ray branching by enabling concomitant separation of pOb pools with lateral splitting *shha* basal epidermal domains.

The directly observed distal movement and likely shedding of fins' basal epidermis, as previously surmised during regeneration (Armstrong et al., 2017; Shibata et al., 2018) is intriguing both functionally and mechanistically. Functionally, continuous bEp replacement may enable rapid recovery from frequent environmental insults. Further, we propose bEp collective

375   movements, distal and lateral, contribute the force that maintains pOb alignment (Armstrong et al., 2017) and enables splitting of physically coupled pOb pools for ray branching. Distal epidermal movements are likely promoted by distributed proliferation across the fin, as seen during regeneration for the basal epidermis (Shibata et al., 2018) and overlying superficial epidermis (Chen et al., 2016; Shibata et al., 2018). In contrast, the cause of periodic lateral *shha*-

380   expressing bEp movements that split *shha*-defined bEp domains and thereby instruct ray branching is unknown.

Shh/Smo signaling is involved in branching morphogenesis of other organs, including the lung (Bellusci et al., 1997; Fernandes-Silva et al., 2017; Pepicelli et al., 1998) and submandibular

385   salivary gland (Jaskoll et al., 2004). In the lung, Shh/Smo also mediates interactions between

mesenchymal and epithelial populations although likely by promoting local proliferation and/or differentiation (Kim et al., 2015). Collective cell migration also is broadly implicated in branching morphogenesis, including for renal tubes, mammary glands and blood vessels (Ewald et al., 2008; Riccio et al., 2016; Spurlin & Nelson, 2017). Unlike those contexts, we propose a

390   neighboring cell type – basal epidermis – that does not directly contribute to the final tissue provides the instructive collective movements. This unusual arrangement may reflect regenerating fin pObs having a mesenchymal state during patterning before returning to their differentiated epithelial state (Stewart et al., 2014).

395   **Shh/Smo signaling may position pre-osteoblasts by promoting their adhesion to moving basal epidermal cells**

Continuous Shh/Smo signaling through fin development and retained splitting of Shh-expressing basal epidermal domains when the pathway is inhibited indicate Shh/Smo has a permissive role in ray branching. We favor a model whereby Shh/Smo's function is to promote

400   transient cell adhesion between bEps and pObs. The transient nature may result from the moving bEps rapidly terminating their *shha* expression and *ptch2*-defined Shh/Smo activity. A slight lateral component to bEp movements away from the midline of each forming ray would then successively tug contacting pObs to follow. Over the course of several days, pObs eventually are pulled into two pools. The pOb pools become sufficiently and irreversibly separated to now

405   generate branched daughter rays.

Our 3-D reconstructions showing Shh-expressing bEps and Runx2-expressing pObs likely share extensive and intimate physical contacts are consistent with this heterotypic cell

adhesion model. Importantly, bEps and pObs would remain adjacent but not necessarily adhered

410   when Shh/Smo signaling is blocked. As such, our novel time-lapse imaging of developing caudal

fins provides key functional support by showing Shh/Smo signaling impedes bEp distal

movements. Notably, Shh-expressing bEps accelerate when they pass beyond Shh-responding

pObs. Chemical inhibition of Shh/Smo signaling significantly increases overall Ptch2-positive

bEp migration rates and eliminates the characteristic velocity decrease when Shh-expressing

415   bEps pass adjacent to pObs. While inhibiting Shh/Smo signaling accelerates individual bEp cell

movements, a steady-state Shh-expressing basal epidermal domain persists and at least partially

splits. However, we propose the pObs cannot follow without Shh/Smo signaling to promote

adhesion with bEps and therefore remain as a single pOb pool that forms an unbranched ray.

Longer-term live imaging monitoring lateral movements of both bEps and pObs would help test

420   this model.

We favor Shh/Smo-promoted adhesion-based skeletal positioning over alternative

hypotheses for additional reasons. First, Shh/Smo-dependent cell proliferation is not observed

during development or regeneration (Armstrong et al., 2017), arguing against a model whereby

425   Shh promotes localized proliferation at the margins of a given pOb pool progressively dividing

it. Notably, previous conclusions that Shh/Smo signaling is a pro-proliferative factor in

regenerating fins (Y. Lee et al., 2009; Quint et al., 2002) may reflect off-target cyclopamine

effects (Armstrong et al., 2017). In contrast, fins develop and regenerate to normal size when

using BMS-833923 to block Shh/Smo signaling with the intriguing exception of the two

430   peripheral rays and their surrounding tissue. Second, all Shh/Smo-responsive cells remain

outwardly specified upon Shh/Smo inhibition, including osteoblasts that still differentiate to

produce ray skeletal units complete with joints. Any additional Shha or other Hedgehog ligand roles, if existent, would seem Smo-independent, as with Indian Hedgehog A (Ihha) and bone maturation during fin regeneration (Armstrong et al., 2017). Third, we use chemical genetics to

435   map the Shh/Smo time-of-function and show ray branching requires persistent Shh/Smo signaling from initial hints of Shh-expressing basal epidermal domain splitting until daughter rays are fully separated. Mechanistically, Shh/Smo signaling seemingly promotes a continuously emergent process and not a periodic switch-like event for either distal bEp movement or Shha-expressing bEp domain splitting.

Short range Shh promoting cell adhesion may be a common Shh/Smo signaling mode

Our proposed short range Shh/Smo signaling mode promoting heterotypic cell adhesion differs from Hh's more typical role as a gradient-forming morphogen. Providing precedence, Hh acts on neighboring cells in several well-established contexts. For example, Hh famously

445   mediates interactions between directly adjacent cells during *Drosophila* embryo segment boundary formation (Ingham, 1993). Short-range Shh/Smo signaling also occurs in vertebrates, including mammalian hair follicle development (Sato et al., 1999; Woo et al., 2012), avian limb patterning (Sanders et al., 2013), and zebrafish retina development (Shkumatava et al., 2004). Perhaps most germane, *shha+* epidermal cells organize directly underlying Hh-responsive

450   dermal cells during zebrafish scale morphogenesis (Aman et al., 2018).

Shh/Smo has also been tied to cell adhesion in other settings. For example, Hh's archetypal role in *Drosophila* wing disc compartment boundary establishment (Ayers et al., 2010) may be through increased "cell bonding" (Rudolf et al., 2015). Shh alters neural crest cell

455   adhesion and migration during avian neural tube morphogenesis (Fournier-Thibault et al., 2009; Jarov et al., 2003; Testaz et al., 2001). Further, misregulated Shh/Smo signaling is linked to invasive cell migration associated with liver, breast, ovarian, and skin cancers (J.-S. Chen et al., 2013; Q. Chen et al., 2014; Hanna & Shevde, 2016; Zeng et al., 2017).

460   Identification, characterization, and manipulation of Shh/Smo-upregulated adhesion molecules are key to strengthen our heterotypic cell adhesion model for ray branching. One intriguing possible mechanism is that Shh/Smo-upregulated Patched directly binds to non-secreted Shh retained on bEp surfaces to increase high-affinity contacts between pObs and bEps. This mechanism could apply elsewhere given Patched is an evolutionary-conserved Shh/Smo-

465   target gene (Alexandre et al., 1996; Goodrich et al., 1996; Lorberbaum et al., 2016; Marigo et al., 1996) while placing Patched in the curious position as both a Shh/Smo effector and negative feedback regulator. Alternatively, Shh/Smo activity could increase expression of more traditional cell adhesion factors or promote cell features (e.g. shape, polarity, or interconnectivity) that indirectly favor heterotypic adhesion.

Fin ray branching as an ancestral mechanism of Shh-mediated appendage patterning and skeletal morphogenesis

How vertebrates pattern skeletal appendages (fins, limbs) is a textbook question of evolutionary and developmental biology. Fin rays comprise dermal bone opposed to

475   endochondral bone found in tetrapod limbs with rays considered lost in tetrapod lineages. However, teleost fin dermal skeleton and tetrapod digits may share deep evolutionary homology (Nakamura et al., 2016). If so, our demonstration Shh/Smo signaling supports developmental ray

branching morphogenesis is intriguing given Shh's long-appreciated but mechanistically distinct role in vertebrate digit patterning. Shh is the secreted morphogen produced by the zone of polarizing activity (ZPA) at the posterior edge of developing limb buds that directs anterior-to-posterior patterning of skeletal elements including digits (Cohn et al., 2002; Riddle et al., 1993). Polarized *shha* expression in zebrafish pectoral fin buds indicates paired fins may follow ZPA-like skeletal patterning (Akimenko & Ekker, 1995; Krauss et al., 1993; Neumann et al., 1999). In contrast, caudal fin primordia lack polarized *shha* (Hadzhiev et al., 2007; Laforest et al., 1998, our results). Consistently, we found disrupting Shh/Smo signaling even prior to formation of the caudal fin field does not alter the initial complement of 18 rays. Moreover, we demonstrate Shh/Smo signaling is required for ray branching in all fins, whether paired or unpaired. The unpaired medial fins (dorsal, caudal, anal) evolved prior to paired fin appendages (Dahn et al., 2006; Desvignes et al., 2018; Freitas et al., 2006; Larouche et al., 2017). Therefore, Shh-dependent ray branching may reflect an ancestral skeletal morphogenesis mechanism that predates emergence of ZPA-based appendage patterning.

Shh/Smo signaling promotes skeletal morphogenesis in many contexts. In zebrafish, Shh/Smo patterns craniofacial dermal bones, as illustrated by the opercle (Huycke et al., 2012), and both developing and regenerating scales (Aman et al., 2018). Shh/Smo signaling also impacts mesenchymal cell movements to pattern bird feathers, another albeit non-ossified skin appendage (Li et al., 2018). Shh further supports patterning of the axial skeleton (Chiang et al., 1996; Choi et al., 2012; Dworkin et al., 2016; Hu et al., 2015; Hu & Helms, 1999; Jeong et al., 2004; Swartz et al., 2012) as well as teeth (Ahn et al., 2010; Dassule et al., 2000; Seppala et al., 2017). Our discovery Shh/Smo signaling enables neighboring cells to position pObs during fin

ray branching suggests similar mechanisms act in other skeletal patterning contexts. If so, manipulating Shh/Smo pathway to position therapeutically delivered or endogenous progenitor cells could enhance skeletal regenerative medicine.

MATERIALS AND METHODS

Zebrafish

Danio rerio zebrafish were maintained in 28-29°C circulating fish water within the University of Oregon Aquatic Animal Care Services (UO AqACS) fish facility. The following lines were used: wildtype AB, *Tg(sp7:EGFP)[b1212]* (DeLaurier et al., 2010)*, TgBAC(ptch2:Kaede)[a4596]* (Huang et al., 2012), *Tg(-2.4shha:gfp:ABC)[sb15]* [previously known as *Tg(-2.2shh:gfp:ABC)*] (Ertzer et al., 2007; Shkumatava et al., 2004), *Tg(runx2:mCherry)* (Shannon Fisher Lab, unpublished). The University of Oregon Institutional Animal Care and Use Committee (IACUC) approved zebrafish experiments.

Microscopy

Larval and juvenile fish were anesthetized with 74 μg/ml tricaine (MS-222, Syndel) in fish facility system water. Fish or dissected fins were transferred immediately to a 35 mm glass bottom FluoroDish plate (World Precision Instruments). Two or three drops of 1% low-melt agarose, stored at 38°C and cooled before application, were placed on the caudal fin. Fins were quickly flattened to the FluoroDish with a single-hair paintbrush before the agarose hardened. The following microscopes were used: Nikon Eclipse T*i*-E widefield and Nikon Eclipse T*i2*-E with Yokogawa CSU-W1 spinning disk attachments, and Zeiss LSM 880 laser scanning confocal microscope. Confocal image stacks were processed using Imaris software to generate single optical slice digital sections, surface renderings, and 3D reconstructions. Adobe Photoshop was used to adjust levels with identical image acquisition and processing settings for a given experiment. Live fish promptly were euthanized or returned to tanks after imaging.

Kaede photoconversion and imaging

ptch2:Kaede fish were anesthetized and placed on FluoroDish plates as described above. Fins
530   were viewed with a Nikon Eclipse T*i*-E widefield microscope or Nikon T*i*2-E/ Yokogawa CSU-
W1 spinning disk confocal microscope. Kaede-expressing regions of interest (ROIs) were
photoconverted using a metal halide light source and DAPI excitation filter or with 405 nm laser
illumination from 10 seconds to 2 minutes, depending on ROI size and fish age. Before and after
images were acquired to ensure complete photoconversion of Kaede from green (518 nm) to red
535   (580 nm) emission. Fish were returned to system water and then similarly re-imaged after
defined periods.

BMS-833923 treatments

BMS-833923 ("BMS", Cayman Chemicals) was dissolved in DMSO to a concentration of 6.3-
540   12.5 mM. This stock was diluted to a final concentration of 0.63-1.25 μM in system fish water
for both larval and juvenile zebrafish treatments. Equal volumes of DMSO were used for control
group treatments.

To test Hh/Smo requirements for caudal fin ray branching, 25 dpf *sp7:EGFP* fish (*n*=6 per
545   group) with unbranched rays were treated initially for 24 hours in BMS or DMSO-alone water
and then returned to standard housing. Fish were exposed to BMS for 4 hours every other day
until the experimental end point at 42 dpf. To assess Hh/Smo roles in all fin appendages,
shha:GFP;runx2:mCherry fish were treated with 1.25 μM BMS or DMSO (*n*= 6 per group) from
21 to 42 dpf. Fins were dissected and imaged as described above.

For staggered-start juvenile fish treatments, 25 dpf *sp7:EGFP* fish were anesthetized, fluorescently screened, and sorted into groups of those having caudal fins with "unbranched" rays or fins in which branching had initiated but was incomplete ("during"). Fish from the two groups were then treated with BMS as described above. Untreated clutchmate *sp7:EGFP* fish

555   were returned to standard housing and screened every other day until all fish had developed branched rays. Drug treatment of the "branched" group of fish was started at 35 dpf. All treatments ended at 42 dpf, when fins were mounted and imaged as described. For BMS-treated fish (unbranched, during branching, and branched), $n=4$ or 5 fish per group with $n=2$ or 3 for DMSO-treated control groups.

For early larval development studies, *sp7:EGFP;runx2:mCherry* and *ptch2:Kaede* fish were bathed in 1.25 µM BMS starting at 2 dpf alongside DMSO-treated controls ($n=33-44$ fish per group, per clutch). The same drug exposure regiment described for juvenile fish was used. From 2-4 dpf, larvae were treated in 40 mL embryo media in petri dishes. From 5-14 dpf, fish were

565   drug-exposed in beakers containing 125 mL embryo media. Drug efficacy on larval fish was assessed by photoconverting distal fin ROIs of *ptch2:Kaede* fins (photoconversion methods described above) at 13 dpf and re-imaging those regions at 14 dpf ($n=3-5$ per group). All fish were screened for skeletal patterning phenotypes by widefield microscopy. Across clutches, 35/44 (79.5%) BMS-treated larvae developed normally (9/44 or 20.5% were runted) compared to

570   26/33 (78.8%) DMSO-treated larvae (7/33 or 21.2% were runted). The ~20% incidence of developmentally delayed larvae was likely caused by extended periods in 250 mL beakers instead of larger nursery tanks. Regardless, nearly all larvae irrespective of size in both groups developed the normal complement of 18 caudal fin rays.

Ray morphometrics

575 Ray lengths were assessed for *sp7:EGFP* clutch mates treated from 25-42 dpf with BMS-833923 or DMSO (experiment described above, *n*=6 per group). Using Fiji-ImageJ software, the Principal Peripheral Ray (unbranching lateral ray) and Dorsal Ray 3 (longest branching ray) were measured from 42 dpf endpoint caudal fin images from the proximal base of the fin to the distal fin end. Fin widths were used to normal for body size but did not differ between DMSO

580 and BMS groups. Raw and normalized data were graphed with GraphPad Prism V8 and significance assessed with a Student's unpaired t-test.

Whole mount immunostaining

shha:GFP caudal fins were harvested at 22-23 dpf and immediately fixed in 4% PFA/PBS

585 overnight at 4°C or for 4 hours at room temperature. Fins were washed extensively in PBS + 0.1% Tween-20 and blocked in 1x PBS, 1% Triton X-100, 5% Normal Goat Serum, and 10% DMSO buffer overnight at 4°C. Fins were incubated with primary antibodies in blocking buffer overnight at 4°C. Primary antibodies were anti-GFP (1:1000; AVES, GFP-1020), anti-Tp63 (1:100; Thermo Fisher, PA5-36069) and anti-Runx2 (1:100; Santa Cruz Biotechnology, sc-

590 101145). Fins were washed in a high-salt 500 mM NaCl buffer for 30 min followed by extensive washes in PBS + 0.1% Tween-20. Secondary antibody incubations using Alexa Fluor conjugates (Thermo Fisher) were performed overnight protected from light at 4°C at a concentration of 1:1000 in blocking buffer. Fins were then washed extensively in PBS + 0.1% Tween-20, nuclei stained with Hoechst (Thermo Fisher), and mounted with SlowFade Diamond Antifade (Thermo

595 Fisher).

Paraffin section immunostaining

Dissected 32 dpf *shha:GFP* caudal fins were fixed in 4% PFA/PBS overnight at 4°C. After

extensive PBS washing, fins were decalcified for 4 days in 0.5M EDTA, pH 8.0 with daily

600   solution changes. Fins then were dehydrated in an ethanol series and tissue cleared with xylenes

prior to longitudinal embedding in paraffin wax. 7 μm sections were cut on a Leica RM255

microtome. Antigen retrieval was performed on rehydrated sections using 1 mM EDTA + 0.1%

Tween-20 for 5 minutes in a pressure cooker. Following PBS washes, sections were blocked in

1x PBS, 10% nonfat dry milk, 2% normal goat serum, and 4% fetal bovine serum for a minimum

605   of 1 hour. Sections were incubated overnight at 4°C with primary antibodies in blocking

solution. Primary antibodies were: anti-GFP (1:3000; AVES, GFP-1020), anti-Tp63 (1:100;

Thermo Fisher, PA5-36039), anti-Laminin (1:40; Sigma, L9393), and anti-Zns5 (1:5, ZIRC).

Sections were washed in PBS containing 500 mM NaCl + 0.1% Tween-20. Alexa Fluor

conjugated secondary antibodies (Thermo Fisher) were diluted 1:1000 in blocking buffer and

610   incubated for 1 hour at room temperature protected from light. Sections were washed, nuclei

stained with Hoechst, and mounted with SlowFade Gold Antifade (Thermo Fisher). Images were

acquired on a Zeiss LSM 880 laser scanning confocal microscope and images processed with

Fiji-ImageJ, Imaris, and Adobe Photoshop.

615   **In vivo EdU incorporation assays**

29 dpf *shha:GFP* juvenile fish were treated with DMSO or 1.25   M BMS for 4 hours in groups

of *n*=5. Anesthetized fish were injected intraperitoneally with 5   l of 1 mg/mL EdU (Thermo

Fisher) in sterile PBS, monitored for recovery for 10 minutes in fresh facility water, and then

returned to treatment tanks. 12 hours post-injection, caudal fins were amputated and fixed for 4

620   hours at room temperature in 4% PFA/PBS. Fins were washed thoroughly with PBS and blocked

overnight at 4 C in PBS/1% Triton X-100/5% Normal Donkey Serum/10% DMSO. EdU signal

was detected with Click-iT Plus Alexa Fluor 647 Picolyl Azide (ThermoFisher) at 2.5 l/ mL

according to the manufacturer's protocol. Following EdU detection, whole-mount GFP

immunostaining and Hoechst nuclear staining was performed as described below. Whole mount

625   confocal images were acquired using a Zeiss 880 LSM and 3D reconstructions prepared using

Imaris. EdU+ and total intra-ray nuclei, i.e. from cells located in between the epidermal Shh

domains of each hemi-ray, were identified and scored for Rays 2 and 3 using the Imaris "Spots"

function and the following parameters: ROI around length of Shha:GFP+ domain, Quality

Threshold 0.642, cell diameter 3 microns. Quantification of EdU+ cells is expressed as the

630   number of EdU+ intra-ray cells over total number of Hoechst-stained nuclei.

Cell migration imaging and analysis

Fish were anesthetized sequentially in freshly prepared 74 μg/ml tricaine solution for 3 minutes

and monitored for slowed opercular movements. Anesthetized fish were transferred to a 35 mm

635   FluroDish plate and mounted in 3% low melt agarose as described earlier. Set agarose was

carefully removed from the most distal region of the caudal fin to allow for free movement of the

epidermis while the trunk remained adhered to the FluoroDish. 74 μg/ml of tricaine solution was

added to maintain anesthesia and cover the fin. After imaging, fish were returned to system water

to confirm recovery and then promptly euthanized. Fish that did not recover were excluded from

640   downstream cell migration analyses. We occasionally observed extremely rapid epidermal

movements in which entire *shha:GFP+* domains would be shed from rays in <15 min. We

suspect this phenomenon results from elevated stress, anesthesia intolerance, and/or damage from plate surface contact or agarose application. We excluded these animals from analyses.

645 For bulk cell migration assays, 22-24 dpf *ptch2:Kaede* fish were treated with 0.63 μM BMS or DMSO (*n*=8 per group). 24 hours later, fish were mounted and imaged with a Nikon Eclipse T*i*-E widefield microscope every 1 minute for 30 minutes. Imaris was used to automatically track cells for 3-6 single *ptch2:Kaede+* basal epidermal cells on dorsal rays 2-5 for each fish. All tracks were quality checked to confirm individual cell tracking. Individual average cell speeds

650 (*n*=38 cells per group) and then averages for each animal were determined. Statistical significance tests comparing all cells tracked (*n*=38 cells per group) and mean cell speed per fish (*n*=8 fish per group) used Student's unpaired t-tests.

For position-dependent cell migration assays, 21-24 dpf *shha:GFP;runx2:mCherry* fish were

655 treated with DMSO or 1.25 μM BMS for 24 hours prior to imaging. Fish were imaged every 2 minutes for 30 minutes with full z-stacks using a Nikon Ti2-E with a Yokogawa CSU-W1 SoRa spinning disk confocal unit. A single hemi-ray of Ray 2 or Ray 3 was analyzed for each time-lapse video. If both rays were captured, the ray with more pObs in frame was analyzed to avoid oversampling individuals. GFP+ cells were automatically tracked using Imaris software "Spots"

660 algorithms with the following parameters: estimated cell diameter 5 microns, maximum distance between frames 6 microns, maximum gap between frames 3 time points. Each cell track was quality checked using 3D reconstructions and edited if Imaris assigned multiple cells to one track or fragmented the track of a given cell. 26-41 cells were tracked across 9 fish (*n*=5 for DMSO and *n*=4 for BMS groups, respectively) for a total of *n*=159 for DMSO-treated and *n*=135 for

665 BMS-treated fish. Data was normalized for each fish by dividing the track speed of a single cell by the average of all cells tracked for that fish. Positional data was determined by setting the X-position of the most distal Runx2+ pOb as "0" and assigning a relative initial X-position for each *shha:GFP+* cell. Cells with a negative starting position were therefore pOb-associated while those with a positive starting position had already migrated beyond the pOb pool when video

670 acquisition began. Imaris was used to determine each cell's total X-displacement and track speed. Graphs were generated using GraphPad Prism V8. Fourth-order best-fit polynomial curves were added to position/speed graphs to help visualize data trends.

ACKNOWLEDGEMENTS

675 We thank the University of Oregon AqACS Facility for zebrafish care; the University of Oregon zebrafish community for support; A. Delaurier and C. Kimmel for *Tg(sp7:EGFP)b1212* fish; S. Megason for providing the *TgBAC(ptch2:Kaede)a4596* fish developed in A. Schier's lab and the *Tg(-2.4shha:gfp:ABC)sb15* fish developed in U. Strahle's lab; G. Crump for providing the *runx2:mCherry* fish generated by S. Fisher's group; and C. Kimmel, A. Saera-Vila, and the

680 Stankunas lab for input.

FOOTNOTES

Competing interests

None.

685 **Author contributions**

J.A.B, A.E.R. and K.S. designed experiments with input from S.S.; J.A.B and A.E.R. performed experiments; J.A.B, A.E.R. and K.S. prepared and wrote the manuscript.

Funding

The National Institutes of Health (NIH) provided research funding (1R01GM127761 (K. S. and

690 S. S.). J.A.B. had an Experiencing Science Practices through Research to Inspire Teaching (ESPRIT) award supported by the National Science Foundation's Robert Noyce Teacher Scholarship Program, a Mary G. Alden Scholarship, and an Institute of Molecular Biology Summer Scholarship. A.E.R. received support from the University of Oregon Genetics Training Program (5T32GM007413).

695 **Data and material availability**

Requests for materials should be addressed to K. S.

REFERENCES

Ahn, Y., Sanderson, B. W., Klein, O. D. and Krumlauf, R. (2010). Inhibition of Wnt signaling by Wise (Sostdc1) and negative feedback from Shh controls tooth number and patterning. *Development* **137**, 3221–3231.

Akimenko, M.-A. and Ekker, M. (1995). Anterior Duplication of the Sonic hedgehog Expression Pattern in the Pectoral Fin Buds of Zebrafish Treated with Retinoic Acid. *Dev. Biol.* **170**, 243–247.

Alexandre, C., Jacinto, A. and Ingham, P. W. (1996). Transcriptional activation of hedgehog target genes in Drosophila is mediated directly by the cubitus interruptus protein, a member of the GLI family of zinc finger DNA-binding proteins. *Genes Dev.* **10**, 2003–2013.

Aman, A. J., Fulbright, A. N. and Parichy, D. M. (2018). Wnt/β-catenin regulates an ancient signaling network during zebrafish scale development. *Elife* **7**, e37001.

Anderson, E., Peluso, S., Lettice, L. A. and Hill, R. E. (2012). Human limb abnormalities caused by disruption of hedgehog signaling. *Trends Genet.* **28**, 364–373.

Armstrong, B. E., Henner, A., Stewart, S. and Stankunas, K. (2017). Shh promotes direct interactions between epidermal cells and osteoblast progenitors to shape regenerated zebrafish bone. *Development* **144**, 1165–1176.

Ayers, K. L., Gallet, A., Staccini-Lavenant, L. and Thérond, P. P. (2010). The Long-Range Activity of Hedgehog Is Regulated in the Apical Extracellular Space by the Glypican Dally and the Hydrolase Notum. *Dev. Cell* **18**, 605–620.

Bellusci, S., Furuta, Y., Rush, M. G., Henderson, R., Winnier, G. and Hogan, B. L. (1997). Involvement of Sonic hedgehog (Shh) in mouse embryonic lung growth and morphogenesis. *Development* **124**, 53–63.

Chen, C.-H., Puliafito, A., Cox, B. D., Primo, L., Fang, Y., Di Talia, S. and Poss, K. D. (2016). Multicolor Cell Barcoding Technology for Long-Term Surveillance of Epithelial Regeneration in Zebrafish. *Dev. Cell* **36**, 668–680.

Chen, J.-S., Huang, X., Wang, Q., Huang, J.-Q., Zhang, L., Chen, X.-L., Lei, J. and Cheng, Z.-X. (2013). Sonic hedgehog signaling pathway induces cell migration and invasion through focal adhesion kinase/AKT signaling-mediated activation of matrix metalloproteinase (MMP)-2 and MMP-9 in liver cancer. *Carcinogenesis* **34**, 10–19.

Chen, Q., Xu, R., Zeng, C., Lu, Q., Huang, D., Shi, C., Zhang, W., Deng, L., Yan, R., Rao, H., et al. (2014). Down-Regulation of Gli Transcription Factor Leads to the Inhibition of Migration and Invasion of Ovarian Cancer Cells via Integrin β4-Mediated FAK Signaling. *PLoS One* **9**, e88386.

Chiang, C., Litingtung, Y., Harris, M. P., Simandl, B. K., Li, Y., Beachy, P. A. and Fallon, J. F. (2001). Manifestation of the Limb Prepattern: Limb Development in the Absence of Sonic Hedgehog Function. *Dev. Biol.* **236**, 421–435.

Chiang, C., Litingtung, Y., Lee, E., Young, K. E., Corden, J. L., Westphal, H. and Beachy, P. A. (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**, 407–413.

Choi, K.-S., Lee, C. and Harfe, B. D. (2012). Sonic hedgehog in the notochord is sufficient for patterning of the intervertebral discs. *Mech. Dev.* **129**, 255–262.

Cohn, M. J., Lovejoy, C. O., Wolpert, L. and Coates, M. I. (2002). Branching, segmentation and the metapterygial axis: pattern versus process in the vertebrate limb. *BioEssays* **24**, 460–465.

Dahn, R. D., Davis, M. C., Pappano, W. N. and Shubin, N. H. (2006). Sonic hedgehog function in chondrichthyan fins and the evolution of appendage patterning. *Nature* **445**, 311–314.

Dassule, H. R., Lewis, P., Bei, M., Maas, R. and McMahon, A. P. (2000). Sonic hedgehog regulates growth and morphogenesis of the tooth. *Development* **127**, 4775–4785.

DeLaurier, A., Eames, B. F., Blanco-Sánchez, B., Peng, G., He, X., Swartz, M. E., Ullmann, B., Westerfield, M. and Kimmel, C. B. (2010). Zebrafish sp7:EGFP: A transgenic for studying otic vesicle formation, skeletogenesis, and bone regeneration. *genesis* **48**, 505–511.

Desvignes, T., Carey, A., Braasch, I., Enright, T. and Postlethwait, J. H. (2018). Skeletal development in the heterocercal caudal fin of spotted gar (lepisosteus oculatus) and other lepisosteiformes. *Dev. Dyn.* **247**, 724–740.

Dworkin, S., Boglev, Y., Owens, H. and Goldie, S. J. (2016). The Role of Sonic Hedgehog in Craniofacial Patterning, Morphogenesis and Cranial Neural Crest Survival. *J. Dev. Biol.* **4**, 24.

Ertzer, R., Müller, F., Hadzhiev, Y., Rathnam, S., Fischer, N., Rastegar, S. and Strähle, U. (2007). Cooperation of sonic hedgehog enhancers in midline expression. *Dev. Biol.* **301**, 578–589.

Ewald, A. J., Brenot, A., Duong, M., Chan, B. S. and Werb, Z. (2008). Collective epithelial migration and cell rearrangements drive mammary branching morphogenesis. *Dev. Cell* **14**, 570–581.

Fernandes-Silva, H., Correia-Pinto, J. and Moura, R. S. (2017). Canonical Sonic Hedgehog Signaling in Early Lung Development. *J. Dev. Biol.* **5**, 3.

Fournier-Thibault, C., Blavet, C., Jarov, A., Bajanca, F., Thorsteinsdóttir, S. and Duband, J.-L. (2009). Sonic Hedgehog Regulates Integrin Activity, Cadherin Contacts, and Cell Polarity to Orchestrate Neural Tube Morphogenesis. *J. Neurosci.* **29**, 12506–12520.

Freitas, R., Zhang, G. and Cohn, M. J. (2006). Evidence that mechanisms of fin development evolved in the midline of early vertebrates. *Nature* **442**, 1033–1037.

Goodrich, L. V, Johnson, R. L., Milenkovic, L., McMahon, J. A. and Scott, M. P. (1996). Conservation of the hedgehog/patched signaling pathway from flies to mice: induction of a mouse patched gene by Hedgehog. *Genes Dev.* **10**, 301–312.

Hadzhiev, Y., Lele, Z., Schindler, S., Wilson, S. W., Ahlberg, P., Strähle, U. and Müller, F. (2007). Hedgehog signaling patterns the outgrowth of unpaired skeletal appendages in zebrafish. *BMC Dev. Biol.* **7**, 75.

Hanna, A. and Shevde, L. A. (2016). Hedgehog signaling: modulation of cancer properies and tumor mircroenvironment. *Mol. Cancer* **15**, 24.

Hu, D. and Helms, J. A. (1999). The role of sonic hedgehog in normal and abnormal craniofacial morphogenesis. *Development* **126**, 4873–4884.

Hu, D., Young, N. M., Li, X., Xu, Y., Hallgrimsson, B. and Marcucio, R. S. (2015). A dynamic Shh expression pattern, regulated by SHH and BMP signaling, coordinates fusion of primordia in the amniote face. *Development* **142**, 567–574.

Huang, P., Xiong, F., Megason, S. G. and Schier, A. F. (2012). Attenuation of Notch and Hedgehog Signaling Is Required for Fate Specification in the Spinal Cord. *PLoS Genet.* **8**, e1002762.

Huycke, T. R., Eames, B. F. and Kimmel, C. B. (2012). Hedgehog-dependent proliferation drives modular growth during morphogenesis of a dermal bone. *Development* **139**, 2371–2380.

Ingham, P. W. (1993). Localized hedgehog activity controls spatial limits of wingless transcription in the Drosophila embryo. *Nature* **366**, 560–562.

Jaskoll, T., Leo, T., Witcher, D., Ormestad, M., Astorga, J., Bringas, P., Carlsson, P. and Melnick, M. (2004). Sonic hedgehog signaling plays an essential role during embryonic salivary gland epithelial branching morphogenesis. *Dev. Dyn.* **229**, 722–732.

Jeong, J., Mao, J., Tenzen, T., Kottmann, A. H. and McMahon, A. P. (2004). Hedgehog signaling in the neural crest cells regulates the patterning and growth of facial primordia. *Genes Dev.* **18**, 937–951.

Kim, H. Y., Pang, M.-F., Varner, V. D., Kojima, L., Miller, E., Radisky, D. C. and Nelson, C. M. (2015). Localized Smooth Muscle Differentiation Is Essential for Epithelial

Bifurcation during Branching Morphogenesis of the Mammalian Lung. *Dev. Cell* **34**, 719–726.

800 **Krauss, R. S., Concordet, J. and Ingham, P. W.** (1993). A functionally conserved homolog of the Drosophila segment polarity gene hh is expressed in tissues with polarizing activity in zebrafish embryos. *Cell* **75**, 1431–1444.

Laforest, L., Brown, C. W., Poleo, G., Géraudie, J., Tada, M., Ekker, M. and Akimenko, M. A. (1998). Involvement of the sonic hedgehog, patched 1 and bmp2 genes in patterning of 805 the zebrafish dermal fin rays. *Development* **125**, 4175–4184.

Larouche, O., Zelditch, M. L. and Cloutier, R. (2017). Fin modules: an evolutionary perspective on appendage disparity in basal vertebrates. *BMC Biol.* **15**, 32.

Lee, H. and Kimelman, D. (2002). A Dominant-Negative Form of p63 Is Required for Epidermal Proliferation in Zebrafish. *Dev. Cell* **2**, 607–616.

810 **Lee, Y., Hami, D., Val, S. De, Kagermeier-Schenk, B., Wills, A. A., Black, B. L., Weidinger, G. and Poss, K. D.** (2009). Maintenance of blastemal proliferation by functionally diverse epidermis in regenerating zebrafish fins. *Dev. Biol.* **331**, 270–280.

Li, A., Cho, J.-H., Reid, B., Tseng, C.-C., He, L., Tan, P., Yeh, C.-Y., Wu, P., Li, Y., Widelitz, R. B., et al. (2018). Calcium oscillations coordinate feather mesenchymal cell 815 movement by SHH dependent modulation of gap junction networks. *Nat. Commun.* **9**, 5377.

Lin, T. L. and Matsui, W. (2012). Hedgehog pathway as a drug target: Smoothened inhibitors in development. *Onco. Targets. Ther.* **5**, 47–58.

Lorberbaum, D. S., Ramos, A. I., Peterson, K. A., Carpenter, B. S., Parker, D. S., De, S., Hillers, L. E., Blake, V. M., Nishi, Y., McFarlane, M. R., et al. (2016). An ancient yet 820 flexible cis-regulatory architecture allows localized Hedgehog tuning by patched/Ptch1. *Elife* **5**, e13550.

Malik, S. (2012). Syndactyly: phenotypes, genetics and current classification. *Eur. J. Hum. Genet.* **20**, 817.

Marigo, V., Scott, M. P., Johnson, R. L., Goodrich, L. V and Tabin, C. J. (1996). 825 Conservation in hedgehog signaling: induction of a chicken patched homolog by Sonic hedgehog in the developing limb. *Development* **122**, 1225–1233.

Nakamura, T., Gehrke, A. R., Lemberg, J., Szymaszek, J. and Shubin, N. H. (2016). Digits and fin rays share common developmental histories. *Nature* **537**, 225.

Neumann, C. J., Grandel, H., Gaffield, W., Schulte-Merker, S. and Nüsslein-Volhard, C. 830 (1999). Transient establishment of anteroposterior polarity in the zebrafish pectoral fin bud in the absence of sonic hedgehog activity. *Development* **126**, 4817–4826.

Quint, E., Smith, A., Avaron, F., Laforest, L., Miles, J., Gaffield, W. and Akimenko, M.-A. (2002). Bone patterning is altered in the regenerating zebrafish caudal fin after ectopic expression of sonic hedgehog and bmp2b or exposure to cyclopamine. *Proc. Natl. Acad. Sci.* **99**, 8713–8718.

Riccio, P., Cebrian, C., Zong, H., Hippenmeyer, S. and Costantini, F. (2016). Ret and Etv4 Promote Directed Movements of Progenitor Cells during Renal Branching Morphogenesis. *PLOS Biol.* **14**, e1002382.

Riddle, R. D., Johnson, R. L., Laufer, E. and Tabin, C. J. (1993). Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell* **75**, 1401–1416.

Rudolf, K., Umetsu, D., Aliee, M., Sui, L., Jülicher, F. and Dahmann, C. (2015). A local difference in Hedgehog signal transduction increases mechanical cell bond tension and biases cell intercalations along the Drosophila anteroposterior compartment boundary. *Development* **142**, 3845–3858.

Sanders, T. A., Llagostera, E. and Barna, M. (2013). Specialized filopodia direct long-range transport of SHH during vertebrate tissue patterning. *Nature* **497**, 628.

Sato, N., Leopold, P. L. and Crystal, R. G. (1999). Induction of the hair growth phase in postnatal mice by localized transient expression of Sonic hedgehog. *J. Clin. Invest.* **104**, 855–864.

Saunders, J. W. and Gasseling, M. T. (1968). Ectoderm-mesenchymal interactions in the origin of wing symmetry. *Ep. Interact.* 78–97.

Sehring, I. M. and Weidinger, G. (2019). Recent advancements in understanding fin regeneration in zebrafish. *WIREs Dev. Biol.* **9**, e367.

Seppala, M., Fraser, G. J., Birjandi, A. A., Xavier, G. M. and Cobourne, M. T. (2017). Sonic Hedgehog Signaling and Development of the Dentition. *J. Dev. Biol.* **5**, 6.

Shibata, E., Ando, K., Murase, E. and Kawakami, A. (2018). Heterogeneous fates and dynamic rearrangement of regenerative epidermis-derived cells during zebrafish fin regeneration. *Development* **145**, dev.162016.

Shkumatava, A., Fischer, S., Müller, F., Strahle, U. and Neumann, C. J. (2004). Sonic hedgehog, secreted by amacrine cells, acts as a short-range signal to direct differentiation and lamination in the zebrafish retina. *Development* **131**, 3849–3858.

Shubin, N. H. and Alberch, P. (1986). A morphogenetic approach to the origin and basic organization of the tetrapod limb. *Evol. Biol.* 319–387.

Spurlin, J. W. and Nelson, C. M. (2017). Building branched tissue structures: from single cell guidance to coordinated construction. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, 20150527.

Stewart, S., Gomez, A. W., Armstrong, B. E., Henner, A. and Stankunas, K. (2014). Sequential and Opposing Activities of Wnt and BMP Coordinate Zebrafish Bone Regeneration. *Cell Rep.* **6**, 482–498.

Swartz, M. E., Nguyen, V., McCarthy, N. Q. and Eberhart, J. K. (2012). Hh signaling regulates patterning and morphogenesis of the pharyngeal arch-derived skeleton. *Dev. Biol.* **369**, 65–75.

Testaz, S., Jarov, A., Williams, K. P., Ling, L. E., Koteliansky, V. E., Fournier-Thibault, C. and Duband, J.-L. (2001). Sonic hedgehog restricts adhesion and migration of neural crest cells independently of the Patched- Smoothened-Gli signaling pathway. *Proc. Natl. Acad. Sci.* **98**, 12521–12526.

Woo, W.-M., Zhen, H. H. and Oro, A. E. (2012). Shh maintains dermal papilla identity and hair morphogenesis via a Noggin–Shh regulatory loop. *Genes Dev.* **26**, 1235–1246.

Zeng, C., Chen, T., Zhang, Y. and Chen, Q. (2017). Hedgehog signaling pathway regulates ovarian cancer invasion and migration via adhesion molecule CD24. *J. Cancer* **8**, 786–792.

Zhang, J., Jeradi, S., Strähle, U. and Akimenko, M.-A. (2012). Laser ablation of the sonic hedgehog-a-expressing cells during fin regeneration affects ray branching morphogenesis. *Dev. Biol.* **365**, 424–433.

Zuniga, A. (2015). Next generation limb development and evolution: old questions, new perspectives. *Development* **142**, 3810–3820.

Figure 1. Basal epidermis-expressed *shha* and *ptch2*-defined responses in basal epidermis and progenitor osteoblasts become progressively distally restricted during caudal fin development.

Figure 1. Basal epidermal *shha* and *ptch2*-defined responses in basal epidermis and progenitor osteoblasts become progressively distally restricted during caudal fin development.

10   (A-J') Differential interference contrast and fluorescent overlay images of developing caudal fins of *shha:GFP* (A-D, I, I') and *ptch2:Kaede* (E-H, J, J') fish of indicated ages. White dotted lines outline the fin fold (A-F). Yellow boxes in (I, J) indicate the 2x zoom fields in (I', J'). White brackets (I-J') mark branched reporter domains in dorsal rays 2 and 3 preceding overt ray branching. (K-P) Single optical slices of caudal fin dorsal ray 3 in longitudinal (K, M ,O) and

15   transverse (L, N, P) planes derived from 3D-reconstructed whole mount confocal images of fluorescent reporter fish of indicated ages. (K, L) 23 dpf *shha:GFP* fin whole mount antibody stained for GFP (green) and the basal epidermis (bEp) marker Tp63 (magenta). The magenta dashed line outlines a representative Tp63+, GFP- cell that occasionally overlay the innermost bEp layer. Nuclei are stained with Hoechst (grey). (M-P) Single optical slice reconstructed

20   equivalents from live whole mount-imaged 19 dpf *ptch2:Kaede;runx2:mCherry* and *shha:GFP;ptch2:Kaede* caudal fins. (M, N) *ptch2:Kaede* (green) is in distal *runx2*-marked pObs (magenta; white brackets) and a thin layer of tightly associated adjacent and further distally-extending bEps (green brackets). (O, P) *ptch2:Kaede* (photoconverted; magenta) in pObs (magenta brackets) and co-localized with *shha:GFP*-expressing bEps (green, white brackets).

25   Scale bars are 250 μm in (A-J') and 10 μm in (K-P).

Figure 1 Supplement 1. Schematic of caudal fin skeletal anatomy.

Figure 1 Supplement 1. Schematic of caudal fin skeletal anatomy.

(A) Schematic of an adult zebrafish caudal fin. The caudal fin skeleton comprises 18 bony rays (lepidotrichia), of which the inner 16 branch at least once. (B) Cartoon rendering of a skeletal ray branch point. Two opposed hemi-cylindrical calcified hemi-rays form each ray. Branching produces two equally sized daughter rays. (C, D) Colored tracings of longitudinal (C) and transverse (D) sections through a branching lepidotrichia. Distal domains of *shha*-expressing basal epidermal cells (bEps, green) directly neighbor Runx2+ progenitor osteoblasts (pObs, magenta). Those bEps distally beyond pOb pools lose *shha* expression. *shha+* bEp and pOb domains both split peripherally as branching initiates. As the fin extends, the distal Runx2+ pOb pool generates differentiating Runx2/Sp7+ osteoblasts (purple) that eventually mature into proximal sp7+ bone-forming cells (blue).

Figure 1 Supplement 2. *shha* is expressed in a single layer of distal basal epidermal cells in developing caudal fins.

Figure 1 Supplement 2. *shha* is expressed in a single layer of distal basal epidermal cells in developing caudal fins.

3D reconstructed sectional views of dorsal ray 3 from a whole mount immunostained 23 dpf *shha:GFP* caudal fin. Panels show shha:GFP (green, A, E, I), the basal epidermal marker Tp63 (magenta, B, F, J), Hoechst-stained nuclei (Hoechst, C) and overlays (D, H, L; H, L are reproduced in main Figure 1). (A-D) Maximum intensity projection (MIP) of the "native" frontal view. (E-L) Reconstructed single optical slice equivalents showing longitudinal (E-H) and transverse (I-L) planar views of the dashed yellow boxed regions in (A-D). Shha:GFP cells define the innermost basal epidermal cell layer and all co-express Tp63. An occasional single-positive Tp63+ bEps (representative cell in magenta dashed lines, H) is found in a second, outer layer of bEps. Scale bars and orientations are indicated.

Figure 1 Supplement 3. *ptch2* is expressed in tightly associated layers of distal basal

60   epidermal cells and progenitor osteoblasts in developing caudal fins.

Figure 1 Supplement 3. *ptch2* **is expressed in tightly associated layers of distal basal epidermal cells and progenitor osteoblasts in developing caudal fins.**

(A-I) Whole mount confocal imaging of dorsal ray 3 from a live 19 dpf *ptch2:Kaede; runx2:mCherry* caudal fin. (A-C) Maximum intensity projection (MIP) of a frontal view. The progenitor osteoblast (pObs) pool is outlined with grey dashed lines. *runx2:mCherry*-marked pObs are in magenta (A) and *ptch2:Kaede* is in green (B). The overlay is shown in (C). (D-I) Reconstructed optical slice views of the region marked by yellow dashed lines in A-C. Overlays (F, I; reproduced in main Figure 1) show relatively proximal regions of *ptch2:Kaede* and *runx2:mCherry+* co-expressing pObs (white brackets) and an adjacent thin layer of *ptch2:Kaede+* basal epidermal cells (bEps). These bEps extend further distally from the pOb pool (green brackets). (J-R) Confocal images of dorsal ray 3 from a 19 dpf *shha:GFP;ptch2:Kaede* caudal fin in which the Kaede protein has been photoconverted from green to red fluorescence emission. (J-L) Frontal view MIP with osteoblast-populated region outlined with grey dashed lines. *shha:GFP* bEps (green) co-express *ptch2:Kaede* (magenta). (M-R) Reconstructed optical slices of the yellow dashed boxes in J-L. Overlays (O, R; reproduced in main Figure 1) demonstrate proximal regions contain single-positive *ptch2:Kaede+* pObs (magenta brackets) whereas distal regions include co-expressing *shha:GFP* and *ptch2:Kaede* bEps (white brackets). Scale bars are 50 μm in A-C, J-L and 10 μm in D-I, M-R.

Figure 1 Supplement 4. Distal *shha* expression is conserved across developing fins.

Figure 1 Supplement 4. Distal *shha* expression is conserved across developing fins.

Dissected fins from a representative 51 dpf *shha:GFP;runx2:mCherry* fish. Zebrafish have 7 fin appendages: the paired pectoral (A, B) and pelvic (C, D) fins and unpaired dorsal (E), anal (F), and caudal (F) fins. All have branched rays marked by *runx2:mCherry* (magenta) with *shha:GFP+* domains (green) at the distal end of each developing ray. All images are brightfield/GFP/mCherry overlays and sized to the same scale (1 mm scale bars).

Figure 2. Active Shh/Smo signaling is restricted to a narrow distal stretch of each developing fin ray.

Figure 2. Active Shh/Smo signaling is restricted to a narrow distal stretch of each developing fin ray.

(A-F) Whole mount fluorescent images of the distal aspect of the caudal fin of a 25 dpf *ptch2:Kaede* larval fish at the time of Kaede photoconversion from green to red (false colored magenta) fluorescence (0 hpc, A-C) and 24 hours later (24 hpc, D-F). The grey dashed octagon marks the photoconverted region of interest (ROI). The 0 hpc overlay (C) demonstrates complete Kaede photoconversion. The same fish imaged at 24 hpc displays a small patch of newly produced green Kaede (white dotted outlines in D, F) within the photoconverted ROI. The grey bracket in (F) marks distally displaced bEps retaining photoconverted Kaede. Slight splitting of the new green Kaede domain (D, F) indicates the onset of ray branching. (G) Schematic of the photoconversion time course. The fish shown is representative of *n=8*. Scale bars are 250 μm.

Figure 2 Supplement 1. BMS-833923 inhibits Shh/Smo signaling in developing caudal fins.

Figure 2 Supplement 1. BMS-833923 inhibits Shh/Smo signaling in developing caudal fins.

Expanded data from Figure 2. (A-L) Whole mount fluorescence images of the distal caudal fin of

25 dpf *ptch2:Kaede* fish treated with DMSO (A-F) or BMS-833923 (BMS; G-L), imaged at the

time of Kaede photoconversion (3 hours post treatment (hpt)) and 24 hours later (27 hpt). Grey

110   dashed octagons mark the photoconverted regions of interest (ROIs). BMS exposure prevented

production of new green fluorescent Kaede within the ROI (J-L). (F', L') Zoom view of distal

ray regions. Brackets mark presence or absence of photoconverted basal epidermal cells (bEps)

that migrated distally over the 24 hour "chase". These cells are missing in BMS-treated fish (L'),

likely due to accelerated movement and therefore shedding. The few green Kaede+ basal

115   epidermal cells in (J, L) migrated into the photoconverted region without producing new Kaede.

(M) Schematic of the time course for drug treatments, photoconversion, and imaging. Imaged

fish represent groups of *n*=8. Scale bars are 250 μm.

Figure 3. Sustained Shh/Smo signaling is required for ray branching in developing caudal

120   **fins.**

Figure 3. Sustained Shh/Smo signaling is required for ray branching in developing caudal fins.

(A-D) Differential interference contrast (DIC) and fluorescence images of caudal fins from DMSO (A, B) and BMS-833923 (BMS)-treated (C, D) 42 dpf *sp7:EGFP* osteoblast reporter fish. Dashed white lines outline dorsal ray 3 and daughter rays, when present. White arrowheads designate a present (A) or absent (B) branch point. White arrows mark ends of the principal peripheral rays. (E-K) Experimental schematic of (E) and resulting caudal fin images of 42 dpf *sp7:EGFP* fish from (F-K') staggering the start of BMS treatment to before (F-G'), during (H-I'), and after branch initiation (J-K'). Yellow dashed boxes outline dorsal ray 2 regions shown in (G', I' and K'). The yellow arrowhead in (G') designates where branching would have occurred without Shh/Smo-inhibition. The red arrowhead in (I') marks where a ray re-fused when BMS was added after branching had initiated. White arrowheads in (I' and K') indicate ray branch points. Images represent treatment groups (before, during, and after) of $n=4$ or 5 BMS-treated and $n=2$ or 3 DMSO controls. Ray re-fusions occur in 4 of 5 fish treated with BMS "during" ray branching in rays 2, 3, and 16. Scale bars are 250 µm or 1 mm, as indicated.

140 **Figure 3 Supplement 1. Shh/Smo signaling contributes to principal peripheral ray outgrowth.**

Figure 3 Supplement 1. Shh/Smo signaling contributes to principal peripheral ray outgrowth.

145   Whole mount caudal fin images of (A) DMSO- and (B) BMS-treated juvenile fish (exposed from 25-42 days post fertilization (dpf)) and (C) graphs of ray morphometrics. Yellow lines indicate measured ray lengths and fin widths. Dorsal ray 3 does not significantly differ in length between treatment groups ($p = 0.09$). In contrast, the non-branching principal peripheral ray 1 is uniquely shorter in BMS-treated versus DMSO control fish ($p < 0.0001$). As such, the ray length ratio

150   between ray 3 and principal peripheral ray 1 is higher in BMS-treated fish ($p < 0.0001$). Images are representatives of experimental groups of $n=6$. Ray length measurements are from the base of the fin and are normalized to fin width, which did not differ between groups. Statistical significance was determined using Student's unpaired t-tests.

Figure 3 Supplement 2. Shh/Smo signaling does not impact proliferation in developing fins.

Figure 3 Supplement 2. Shh/Smo signaling does not impact proliferation in developing fins.

160   (A, B) Confocal maximum intensity projection images of whole mount immunostained 29 dpf

developing *shha:GFP* caudal fins. Fish were treated with DMSO (A) or BMS-833923 (BMS)

(B) for 4 hours, IP injected with EdU, returned to drug, and fins harvested 12 hours later. GFP is

in green, Hoechst nuclear stain in blue, and EdU as detected with Click-iT Plus in white. (C)

Scatter plot graph showing the percent of intra-ray EdU+ cells, i.e. those located in between the

165   epidermal Shh domains of each hemi-ray, does not significantly differ between DMSO controls

and BMS-treated fish. *n=5* for each group. (D-I) 3D reconstructions of ray 3 domains marked by

yellow dashed boxes in (A, B). (D, G) Overlay of GFP and EdU, sagittal view. (E, H) GFP with

magenta spheres marking EdU+ cells located within the intra-ray space, detected and quantified

with Imaris software. (F, I) Longitudinal view of (E, H). Intra-ray EdU+ cells are located in

170   between Shh+ domains with epidermal cells excluded from automated scoring. Scale bars are

150 μm or 30 μm, as indicated.

Figure 3 Supplement 3. Shh/Smo signaling does not influence initial caudal fin patterning.

Figure 3 Supplement 3. Shh/Smo signaling does not influence initial caudal fin patterning.

175 (A-C) Whole mount differential interference contrast (DIC) and/or fluorescence images of a 2 dpf *ptch2:Kaede* embryo. The dashed yellow box marks the 1.5x zoomed region shown in (B, C). The primordial fin fold lacks rays or structural ray precursors. *ptch2:Kaede* expression is restricted to the notochord and does not expand into the fin fold. (D-E'). *ptch2:Kaede* larvae treated with DMSO or BMS-833923 (BMS) from 2 dpf, photoconverted distal fin regions at 13

180 dpf (grey dashed lines), and imaged 24 hours later (*n*=3-5 per group). BMS treatment blocked production of new green Kaede (yellow arrowhead, E') compared to controls (white arrowhead, D'). (F-G') Whole mount fluorescence images of *sp7:EGFP;runx2:mCherry* larvae treated with DMSO (F) or BMS (G) from 2-14 dpf. Both groups have caudal fins with the standard complement of 18 segmented rays, with 9 rays each on the dorsal and ventral lobes (yellow

185 brackets). BMS-treated fins exhibit shortened principal peripheral rays (yellow arrowhead, G') compared to DMSO controls (white arrowhead, F'). *n*=33 DMSO- and *n*=44 BMS-treated larvae from 2-14 dpf. Scale bars are 1 mm, 250 μm, or 500 μm, as marked.

Figure 4. Shh/Smo signaling promotes ray branching in all paired and unpaired fins.

Figure 4. Shh/Smo signaling promotes ray branching in all paired and unpaired fins.

190   Brightfield and fluorescence overlay images of isolated fins from *shha:GFP;runx2:mCherry*

     juvenile fish treated with DMSO (A-G) or 1.25 µM BMS-822923 (BMS; H-N) from 21-42 dpf.

     runx2:mCherry-labeled rays branch (white arrowheads) in all fins whereas rays of BMS-treated

     fish mostly fail to branch or have severely delayed branching (yellow arrowheads). *shha:GFP+*

     basal epidermal cells are restricted to distal ray tips under both conditions. *n=6* for each group.

195   Scale bars are 1 mm.

Figure 5. Shha+ bEps and pObs are intertwined in developing fins.

Figure 5. Shha+ bEps and pObs are intertwined in developing fins.

(A-F) Fluorescence widefield (A-C) or confocal (D-F) images of the dorsal caudal fin lobe of a 28 dpf *shha:GFP;runx2:mCherry* fish. (A-C) *shha:GFP*-expressing basal epidermal cells (green) overlay and extend distally from *runx:mCherry*-high progenitor osteoblasts (magenta; pObs). The overlay in (C) includes a brightfield image for context. (D-F) A single optical slice (E; sagittal; orientation key in E') and reconstructed longitudinal (D) and transverse (F) views of a distal ray region undergoing ray branching. *shha:GFP+* bEps (green) and *runx2:mCherry+* pObs (magenta) have overlapping signal at interfaces, indicating their tight juxtaposition. bEps and pObs tandemly separate into split pools during branching (white arrows). The grey dotted oval highlights a ridge of pObs nestled into a *shha:GFP+* bEp groove (Movie 2). Scale bars are 250 μm (A-C) and 10 μm (D-F).

Figure 5 Movie 1. A dynamic 3-D visualization of confocal-imaged caudal fin dorsal ray 3 from a 28 dpf *shha:GFP;runx2:mCherry* fish. *shha:GFP+* bEps (green) of both hemi-rays are beginning to branch. Underlying pObs (magenta) are pressed against bEps. 3D reconstruction of live confocal microscopy. Surfaces added with Imaris to visualize domain boundaries.

Figure 5 Movie 2. 3-D space exploration of the distal region of a confocal-imaged caudal fin hemi-ray from a 28 dpf *shha:GFP;runx2:mCherry* fish. The *shha:GFP+* bEp (green) domain has started splitting ahead of ray branching. A ridge of pObs (magenta) is nestled in a hollow of *shha:GFP+* bEps. Imaris-generated surfaces help visualize domain boundaries. Single optical slices of the same region are shown in Figure 5.

Figure 5 Supplement 1. Shha+ bEps are tightly associated with pObs in distal regions of incomplete basement membrane assembly.

Figure 5 Supplement 1. Shha+ bEps are tightly associated with pObs in distal regions of incomplete basement membrane assembly.

(A-D) Confocal images of the distal ray regions from immunostained longitudinal caudal fin sections of 32 dpf *shha:GFP* fish. GFP-expressing basal epidermal cells (bEps; green) and Zns-5-marked osteoblasts (white) are separated by a Laminin-defined basement membrane (BM; magenta). (D') 2x magnification of the yellow dashed box region in the (D) overlay. Asterisks mark areas where Laminin signal is less dense, indicating an incompletely assembled BM, and GFP and Zns-5 partially overlap (D'). Scale bars are 20 μm.

Figure 5 Supplement 2. Shh/Smo signaling does not influence Shh+ bEps and pOb juxtaposition.

Figure 5 Supplement 2. Shh/Smo signaling does not influence Shh+ bEps and pOb juxtaposition.

(A-J) Confocal or widefield fluorescence caudal fin images from *shha:GFP;runx2:mCherry* fish treated with DMSO (control) or 1.25 μM BMS-833923 (BMS) starting at 24 dpf (*n*=8 per group). (A-C, E-G) Single optical slices or equivalent 3D-reconstructed views of 34 dpf mid-treatment fish, imaged concurrently with active ray branching morphogenesis. *shha:GFP+* basal epidermal cells (bEps; green; A) are closely associated with *runx2:mCherry*-expressing progenitor osteoblasts (pObs; magenta). BMS-treated fish (E-G) maintain close Shh+ bEp and Runx2+ pOb contacts while lacking clear *shha:GFP* domain-splitting (*n*=4 per group). (D, H, I, J) Caudal fin images of *shha:GFP;runx2:mCherry* fish continued on drug treatment until 42 dpf (*n*=4 per group). (D, H, I) Widefield fluorescence and brightfield overlay caudal fin images. Unlike DMSO controls (D), BMS-treated fish do not develop branched rays, confirming drug efficacy (H). (I, J) Fin of a 21-42 dpf BMS-treated fish. The yellow dashed box indicates the GFP-alone magnified panel in (J). *shha:GFP* domains are variably split (two grey br ets, rays 2, 3), partially split (dashed grey bracket, ray 4) or unsplit (one grey bracket, rays 5-7). Scale bars are 20 μm (A-C, E-G), 1 mm (G, H, I) and 500 μm ack (J).

Figure 6. Shh/Smo signaling slows collective migration of *shha:GFP+* basal epidermal cells associated with pObs.

Figure 6. Shh/Smo signaling slows collective migration of *shha*-expressing basal epidermal cells associated with pObs.

(A-D') Frames from a time lapse movie showing dorsal ray 3 of the caudal fin from live mounted 24 dpf *shha:GFP;runx2:mCherry* fish treated with DMSO or BMS-833923 (BMS) for 24 hours prior to imaging. Images are maximum intensity projections (MIP) and show the start (0 min; A, A', C, C') and end (30 min; B, B', D, D') points. The Imaris-generated colored tracks show the progressive displacement of individual *shha:GFP+* bEps (green). Grey spheres show the starting or final positions of all tracked bEps. Grey dashed vertical lines in (A, B, C, D) indicate the distal most Runx2+ pOb (magenta), defined as position "0". (E, F) Scatterplot graphs showing the average speed of individual bEps, considering their net X-displacement and normalized to all scored cells of the given fish, relative to starting position for DMSO- (159 individual cells from five fish) and BMS-exposed fish (135 cells from four fish). Dot colors correspond to cells from a given fish. Scale bars are 50 μm.

Figure 6 Supplement 1. Shh/Smo signaling restrains the distal migration of basal epidermal cells.

Figure 6 Supplement 1. Shh/Smo signaling restrains the distal migration of basal epidermal cells.

(A-C) Still frames from 30 minute time lapse movies capturing basal epidermal (bEp) distal migration in a caudal fin of a live mounted 23 dpf *ptch2:Kaede* fish. The fish was DMSO-treated as a control group member for quantitative studies. Maximum intensity projections (MIPs) of a fin's dorsal lobe are shown at 0 minutes (A, start position), 16 minutes (B, halfway through video), and 30 minutes (C, end position). White and magenta arrowheads indicate a *ptch2:Kaede+* bEp and pOb, respectively. Yellow dashed boxes mark the 3x magnified distal ray region in (A'-C'). The representative bEp (white) and pOb (magenta) are outlined with dashed lines. The white Xs in (B', C') indicates the bEps' starting position at 0 minutes. Scale bars are 100 μm. (D) Scatterplot graph showing average cell migration speeds (arbitrary relative units) of individual *ptch2:Kaede+* bEps from fish treated with DMSO or BMS-833923 (BMS) for 24 hours and then imaged over 30 minutes. bEps of BMS-treated fish migrate faster than DMSO controls ($p < 0.0001$, Student's t-test). (E) Graph showing average cell migration rate by individual fish (as indicated by different colors) in BMS-treated fish are significantly faster ($p < 0.0006$, Student's t-test). *n*=3 to 6 cells per fish with 8 fish per treatment for a total of 38 tracked cells per group. Colors represent single bEps from the same animal.

Figure 6 Movie 1.

Time lapse movie of the dorsal lobe of the caudal fin of a live-mounted 23 dpf *ptch2:Kaede* fish treated with DMSO (control group) for 24 hours prior to imaging. Widefield fluorescence images were collected every 2 minutes over a 30 minute period. Still images and data analysis are in Figure 6 Supplement 1.

Figure 6 Supplement 2. Position- and Shh/Smo-dependent *shha:GFP*+ bEp migration rates of individual fish.

Figure 6 Supplement 2. Position- and Shh/Smo-dependent *shha:GFP+* bEp migration rates of individual fish.

Expanded data from Figure 6. (A-E', K-N') Maximum intensity projections of the final 30 minute frame of time lapse-imaged *shha:GFP;runx2:mCherry* fish (21-24 dpf) treated with DMSO (*n*=5) or BMS-833923 (BMS; *n*=4) for 24 hours. Panels show *runx2:mCherry* progenitor osteoblasts (pObs) only (A-E, K-N; magenta) or both pObs and *shha:GFP+* basal epidermal cells (A'-E', K'-N'; bEps; green;). Grey spheres mark semi-automatically tracked bEps with their cell displacement over 30 minutes indicated by multicolor tracks. (F-J, O-T) Average speed vs. starting position scatterplot graphs for each fish. Data points represent individual cell speeds. Non-parametric best-fit curves provide a visual guide. (S, T) Summary graphs reproduced from Figure 6 show combined data with added overall trends. Scale bars are 50 μm.

Figure 6 Movie 2.

Time lapse movie of caudal fin dorsal lobes of representative live-mounted *shha:GFP;runx2:mCherry* fish treated with DMSO (upper panel) or BMS (lower panel) for 24 hours prior to imaging. Shh+ bEps in green and Runx2+ pObs in magenta were imaged by fluorescence confocal microscopy. Multicolor lines indicate cell migration tracks for individual Shh+ bEps (marked by grey spheres) over time. Images were acquired every 2 minutes for 30 minutes. Still images and data analysis are in Figure 6.