Predicting clinical drug response from model systems by non-linear subspace-based transfer learning

Soufiane Mourragui[1,2], Marco Loog[2,3], Daniel J. Vis[1], Kat Moore[1], Anna G. Manjon[4], Mark A. van de Wiel[5,6], Marcel J.T. Reinders[2,7,8], Lodewyk F.A. Wessels[1,2,8]

[1]*Division of Molecular Carcinogenesis, Oncode Institute, The Netherlands Cancer Institute, Amsterdam, The Netherlands.*

[2]*Department of EEMCS, Delft University of Technology, Delft, The Netherlands.*

[3]*Department of Computer Science, University of Copenhagen, Copenhagen, Denmark.*

[4]*Division of Cell Biology, Oncode Institute, The Netherlands Cancer Institute, Amsterdam, The Netherlands.*

[5]*Epidemiology and Biostatistics, Amsterdam University Medical Center, Amsterdam, The Netherlands.*

[6]*MRC Biostatistics Unit, Cambridge University, Cambridge, United Kingdom.*

[7]*Leiden Computational Biology Center, Leiden University Medical Center, Leiden, The Netherlands.*

[8]*To whom correspondence should be addressed: Lodewyk F.A. Wessels (l.wessels@nki.nl) and Marcel J.T. Reinders (m.j.t.reinders@tudelft.nl).*

Abstract

Pre-clinical models have been the workhorse of cancer research for decades. While powerful, these models do not fully recapitulate the complexity of human tumors. Consequently, translating biomarkers of drug response from pre-clinical models to human tumors has been particularly challenging. To explicitly take these differences into account and enable an efficient exploitation of the vast pre-clinical drug response resources, we developed TRANSACT, a novel computational framework for clinical drug response prediction. First, TRANSACT employs non-linear manifold learning to capture biological processes active in pre-clinical models and human tumors. Then, TRANSACT builds predictors on cell line response only and transfers these to Patient-Derived Xenografts (PDXs) and human tumors. TRANSACT outperforms four competing approaches, including Deep Learning approaches, for a set of 15 drugs on PDXs, TCGA cohorts and 226 metastatic tumors from the Hartwig Medical Foundation data. For only four drugs Deep Learning outperforms TRANSACT. We further derived an algorithmic approach to interpret TRANSACT and used it to validate the approach by identifying known biomarkers to targeted therapies and we propose novel putative biomarkers of resistance to Paclitaxel and Gemcitabine.

Introduction

The accumulation of somatic alterations on the genome and epigenome transforms healthy cells into malignant tumor cells. Although these alterations are required for tumor growth, they also confer vulnerabilities on tumor cells. Some well-known examples of such genetic vulnerabilities are the amplification of ERBB2 in breast cancer[1], the BRAF^{V600E} mutation in skin melanoma[2] or the BCR/ABL fusion in leukemia[3]. These vulnerabilities have been successfully exploited clinically by

directing drugs against them. However, for the vast majority of cancer patients, no

clear biomarkers exist. Hence, expanding our arsenal of accurate biomarkers would

pave the way for personalized medicine, by identifying, for each patient, the most

effective drug[4].

In order to discover such biomarkers, pre-clinical models have been used extensively

in the past decades, either in the form of cell lines, patient-derived xenografts (PDX)

or organoids. This was partially fueled by the relative ease with which these model

systems can be subjected to drug screening. This has led to break-through discoveries

with broad clinical impact[5]. However, *Paul Valery*'s statement, "what is simple is

always wrong; what is not, is unusable"[6], also applies to these model systems.

Specifically, their simplicity also confers weaknesses: the lack of a micro-environment

in cell lines, and the absence of an immune system in cell lines, PDXs and organoids.

These shortcomings are further amplified by culture artefacts[7,8] that lead to a reduced

clinical significance of these models[9,10] and a high attrition rate in drug development[11].

Computational approaches that correct for these differences are therefore needed to

improve the identification of truly predictive biomarkers[12]. In the particular case of

cancer, these approaches are divided into two distinct categories. In the first category,

mechanistic models are developed on pre-clinical models and subsequently

"humanized" to focus on the similarities between pre-clinical models and human

tumors[13]. The second category approaches the problem in a statistical fashion. Using

molecular profiles and drug screens from large-scale panels of pre-clinical models[14,15],

cell line drug response predictors are inferred[16–18]. The resulting models are then

applied to predict the sensitivity of patients to certain drugs[19–21]. Although already

promising, these approaches either do not take into account the fundamental

differences between pre-clinical models and human tumors[22], or only model these

differences as a technical batch effect[19–21]. Recently, transfer learning and multi-task

learning approaches have been developed to explicitly take these differences into

account, either partially using tumor responses during training[23], or solely based on

pre-clinical labels[24] while only modeling differences linearly.

We present **TRANSACT** (**T**umor **R**esponse **A**ssessment by **N**on-linear **S**ubspace

Alignment of **C**ell-lines and **T**umors), a general framework for subspace-based

transfer learning[25–29] which enables the transfer of drug response predictors trained

on a source domain (e.g. cell lines and PDXs) to a target domain (e.g. human tumors).

TRANSACT employs the powerful mathematical framework of Kernel methods[30–35] to

capture both linear and non-linear molecular processes expressed in source and

target domains. First, we demonstrate that, compared to existing methods[20,21,24],

modeling non-linearities using TRANSACT improves drug response prediction in

PDXs after training on cell line responses only. We fix the hyperparameter controlling

the degree of non-linearity on the PDX data and then employ TRANSACT to transfer

predictors of drug response trained on cell lines to two human tumor datasets: primary

tumors from TCGA and metastatic lesions from the Hartwig Medical Foundation

(HMF). Specifically, we show a significant improvement in response prediction for five

drugs in TCGA and three drugs in HMF, compared to state-of-the art competing

approaches. Importantly, this performance improvement is attained without any

training on data from the human tumors. We finally employ the interpretability of our

approach to identify genes and pathways associated with drug response. We provide

a full mathematical derivation of our algorithm, a complete reproducible pipeline and

a fully open-source software package.

Results

TRANSACT: Generating non-linear manifold representations to transfer

predictors of response from pre-clinical models to tumors

TRANSACT compares genomic signals contained in the source (e.g. pre-clinical

models) and target (e.g. human tumors) datasets, and outputs a representation of

processes that are present in both datasets. The nature of this representation depends

on the similarity function K that characterizes the relationships between samples

(Methods). Depending on the similarity function employed, various types of non-linear

relationships can be modeled. For instance, in the case of a Gaussian similarity

function, these non-linearities include constant, linear, second and higher-order

interaction terms (Methods).

In a first step, TRANSACT computes the similarities between all samples (**Figure 1**A),

yielding three matrices: K_s, K_t and K_{st}, containing the similarities between source

samples, target samples, as well as source and target samples, respectively. Using

an eigen-decomposition of K_s and K_t, directions of maximal variance in the similarity

space are computed using *Kernel PCA*[36] (**Figure 1**B). This decomposition is

performed independently on the source and the target spaces and yields two

importance score matrices: one for the source non-linear principal components

(NLPCs) and one for the target NLPCs. These importance scores are *not* the projected

sample values — they actually correspond to loadings, in the sense that they

represent the geometric directions of the NLPCs in the sample similarity space (**Supp**

Figure 1A). We then quantify the geometric differences between all pairs of NLPCs from the source and target sets by computing the cosine similarity matrix (**Figure 1**C). In a subsequent step (**Figure 1**D), we align these two sets of NLPCs by using the notion of *principal vector*s illustrated in **Supp Figure 1**B. These principal vectors (PV) are pairs of vectors – one from the source, one from the target – ranked by decreasing similarity: the top PVs correspond to geometrically similar factors, while bottom PVs are pairs of almost orthogonal vectors. We restrict further analysis to the most similar PVs based on a similarity of at least 0.5 (Supp. Material). We perform, for each selected pair of PVs, an interpolation between the source and the target vectors and select one intermediate representation that best balances the contribution of the source and the target signals (**Figure 1**E, **Supp Figure 1**C). These vectors, called *consensus features*, define the consensus space, into which we project the source (pre-clinical) and target (tumor) samples. This projection yields, for all source and target samples, consensus feature values that can be used as input to any machine learning model to build predictors of drug response (**Figure 1**G, Method). In the case of a linear similarity function, TRANSACT reduces to PRECISE[24] (Subsection Supp. 8) and is fundamentally different from approaches such as Canonical Correlation Analysis (CCA)[37] (Subsection Supp. 9).

Non-linearities improve response prediction of predictors transferred from cell lines to patient-derived xenografts (PDXs)

When it comes to predicting drug response in one model system, it is known that inducing non-linearities can lead to improved performance[35], although linear methods remain competitive[18,38]. We investigated whether the introduction of non-linearities in the computation of sample similarities resulted in improved response prediction of

predictors trained on cell lines (source domain) and transferred to PDXs (target

domain). Since gene expression is known to have predictive power comparable to

other omics datasets combined[16,18,39,40], we restricted our analysis to the expression

of 1780 genes known to be related to cancer[41]. Using TRANSACT, we computed

consensus features for cell lines (1049 cell lines from 26 different tissues) and all PDXs

(399 samples from 5+ different tissues) (Methods). We projected the gene expression

data of all cell lines and all PDXs onto these consensus features. We employed Elastic

Net to train models of drug response with the projected cell line expression data as

input and the drug response data as output, which is quantified as the AUC, i.e. the

area under the drug response curve (Methods). We applied this trained predictor on

the projected PDX expression data and compared the predicted response to the

measured best average response by Spearman Correlation (**Figure 2**A). We made

use of the standard Gaussian similarity function (Methods) to vary the level of non-

linearity introduced. This similarity function is characterized by a single scaling factor

γ, whose size is directly proportional to the proportion of non-linearity introduced

(**Figure 2**B). We studied the predictive performance in PDXs for seven different values

of γ, ranging from a set of consensus features with an almost purely linear ($\gamma =$

1×10^{-5}) to an almost purely non-linear composition ($\gamma = 1 \times 10^{-2}$). We compared

the performance of TRANSACT to two approaches that do not perform domain

adaptation: *Elastic Net*[22] and *Deep Learning* regression (DL). We further compared it

to two state-of-the-art domain adaptation approaches: ComBat batch correction

followed by Deep Learning regression (*ComBat + DL*)[21] and *PRECISE*[24], a linear

domain adaptation approach. All models were trained to predict response to seven

different drugs (Erlotinib, Cetuximab, Gemcitabine, Afatinib, Paclitaxel, Trametinib and

Ruxolitinib) for which we had response data available for both PDX models and cell

lines (**Figure 2**C-I). For these seven drugs we observe, in general, a clear

improvement of approaches that employ domain-adaptation over those that do not,

indicating a necessity to correct the input signal when moving from the source to the

target domain. When comparing *Elastic Net* to *Deep Learning*, we observe that

modeling non-linearities results in a better transfer for five drugs. We further observe

an improvement of TRANSACT over *ComBat+DL*, suggesting that the correction

required from cell lines to PDXs is more complicated than correcting for a technical

batch effect. We observe that the introduction of non-linearity with TRANSACT

generally increases the predictive performance on PDXs when compared to a linear

similarity function (*PRECISE*). Specifically, we observe for several drugs that the

predictive performance increases with the scaling factor until a maximal performance

is reached ($\gamma = 10^{-4}$ for Erlotinib, Cetuximab and Afatinib and $\gamma = 10^{-3}$ for

Gemcitabine, Paclitaxel and Trametinib), after which the predictive performance

drops. We therefore decided to fix the scaling factor to the average of these two values

($\gamma^* = 5 \times 10^{-4}$) and employ the associated consensus space to transfer the predictors

of response to the tumor samples. As a further check, we analyzed the properties of

the consensus space obtained using γ^*. We observe a concentration of the offset

contribution in the top consensus features and an increasing proportion of non-linear

term contribution to lower order features (**Supp Figure 6**C). The UMAP[42] projection

of the consensus features shows a clear co-clustering of cell lines and PDXs of the

same tissue (**Supp Figure 6**D).

189 *Consensus features between cell lines (GDSC) and human tumors*

190 *conserve primary tumor information.*

With the scaling factor (γ) calibrated on PDX models, we moved to the clinical setting

to investigate domain adaptation between cell lines and two different human tumor

datasets: primary tumors from TCGA and metastatic lesions from the HMF. We

selected 30 consensus features in the GDSC-TCGA analysis (**Supp Figure 8**) and 20

in the GDSC-HMF analysis (**Supp Figure 9**). We arrived at these numbers by first

selecting NLPCs based on the inflexion point of the cumulative eigenvalues, and

subsequently only retaining PVs with a similarity above 0.5. We observe that the

consensus features computed between GDSC and TCGA (**Figure 3**A) and between

GDSC and HMF (**Figure 3**B) show the same proportion of non-linearity with a

concentration of offset and linearities in the top consensus features.

In order to visualize the structure retained in the consensus space, we embedded our

consensus scores into a 2D space using UMAP[42]. We observed that primary tumors

cluster together based on their tissue type (**Figure 3**C). Cell lines, however, show

different behaviors – most do cluster with the tumors from a similar tissue of origin,

while a group of cell lines cluster together and away from the tumors, regardless of

their tissue of origin, as observed in previous studies[43]. To quantify the degree of co-

clustering of cell lines and tumors, we compared distances between tumors and cell

lines from similar and non-similar tissues, and observed, as expected, a higher

similarity between tumors and cell lines from the same tissue (**Supp Figure 10**C).

Metastatic lesions show a weaker clustering based on the primary tumor's tissue of

origin (**Supp Figure 10**D-E). This is not unexpected, as the expression profiles are

derived from biopsy sites distant from the primary tissue. Of particular interest, we

observe the existence of a hematopoietic cell-line cluster that co-clusters with

metastatic samples from various biopsy sites. Most of these tumor samples (7 out of

12 samples) are lymph node metastases and most likely display a hematopoietic

expression profile due to blood infiltration in the samples (**Supp Figure 10**B).

Consensus features increase transfer of response predictors from cell

lines to primary tumors and metastatic lesions

To further validate our approach, we transferred response predictors from cell lines to

the TCGA and HMF collections of human tumors. First, we projected the GDSC and

TCGA expression data onto the GDSC-TCGA consensus features. Then we trained,

for each drug, a regression model using solely the cell line response data (measured

as AUC). These drug-specific regression models were then used to predict response

on the projected TCGA data. Finally, we compared the predicted response to the

known categorical clinical responses using a one-sided Mann-Whitney test and

computed the corresponding effect size. We trained models for seventeen different

drugs (**Table 1**A). We compared the performance of TRANSACT to the performance

obtained by four competing approaches (*Elastic Net, DL, ComBat+DL* and *PRECISE*)

(**Table 1**A, Methods). For the Deep Learning approaches (*DL* and *ComBat+DL*), we

selected the architecture and hyper-parameters for each drug by 5-fold cross-

validation on GDSC (Methods). We subsequently trained 50 models with different and

independent initializations (Method) and reported the median performance obtained

on TCGA. *Elastic Net* and *PRECISE* obtain significant associations for three and six

drugs, respectively, but neither approach ever outperforms all other approaches.

ComBat+DL and *DL* achieve significant associations for five and eight drugs,

respectively – however, both approaches outperform all others for only three and one

drugs, respectively. By contrast, TRANSACT achieves significant associations for

seven drugs and outperforms all other approaches for five drugs. For both deep

learning approaches, we observe an important dependency on the network

initialization (**Supp Figure 11**A, **Supp Figure 12**A). More importantly, we observe no

correlation between the training error achieved on the source domain and the

prediction accuracy on the target domain, making it impossible to select a proper

initialization solely based on the source domain performance (**Supp Figure 11**B,

Supp Figure 12B). Results obtained with TRANSACT, on the contrary, do not depend

on a random initialization.

For the HMF data, we repeated the steps above, while employing the GDSC-HMF

consensus features as well as the HMF and GDSC expression and response data.

We trained models for Irinotecan, Carboplatin (using Cisplatin GDSC response),

Trastuzumab (using Afatinib GDSC response), Gemcitabine, Paclitaxel and 5-

Fluorouracil. Across all approaches, we observe a significant association between the

predicted AUC and clinical responses for four of the six drugs (Irinotecan, Carboplatin,

Trastuzumab and Gemcitabine) (**Table 1**B, **Figure 4**A). *PRECISE* reaches

significance for two drugs, whereas *Elastic Net, DL* and *Combat+DL* reach

significance for a single drug. *TRANSACT*, by contrast, outperforms *PRECISE* on

three of the five drugs, and *ComBat + DL*, *Deep Learning* and *Elastic Net* on four

drugs. *TRANSACT* does not obtain a significant association for Paclitaxel. We do,

however, observe a trend that indicates a positive association between predicted and

clinical response. In contrast, all competing approaches fail to achieve any

association. **Figure 4**A provides a summary of the performance of TRANSACT and

two state-of-the-art approaches, *PRECISE* and *Combat + DL,* on both TCGA and HMF

cohorts. Finally, for the HMF cohort, we also observe a large variability in predictions

of the deep learning approaches as a result of network initialization (**Supp Figure 11**C,

Supp Figure 12C). As on the TCGA we do not observe any relationship between the

training error on the source domain data and the predictions on the target domain data

(**Supp Figure 11**D, **Supp Figure 12**D).

Two drugs, already standing out from the GDSC-PDX analysis, are of particular

interest. We first observe that Gemcitabine is consistently better predicted by

TRANSACT than by any other approaches (**Figure 4**B, **Table 1**A). When it comes to

Paclitaxel (**Figure 4**C, **Table 1**A), *TRANSACT* shows a clear improvement over

competing approaches in TCGA. *TRANSACT* is also the only method to recover a

positive association on HMF, although the association is not significant. Finally, for

Carboplatin (**Figure 4**C, **Table 1**A), we observe that *TRANSACT* outperforms all

approaches on TCGA, and on HMF. When taking the TCGA and HMF results together

– and noting that there is overlap in the drugs employed in these two cohorts – it is

interesting to observe that the best competing approach is *Deep Learning*. *Deep*

Learning achieves significant associations in 9 of the 23 comparisons while

ComBat+DL reaches significance in 6 trials. Based on this metric, *TRANSACT*

performs best, reaching significance in 11 of 23 trials. Furthermore, *Deep Learning*

outperforms all other approaches in only 3 comparisons, while *TRANSACT* delivers

increased performance in 8 comparisons. When also taking the performance on the

PDX models into account, we can safely conclude that *TRANSACT* is more accurate

in transferring drug response predictors.

Interpretability of consensus features confirms known mechanisms for

targeted therapies and unveils potential biomarkers of sensitivity for

cytotoxic drugs

We finally made use of the interpretability of our approach to mechanistically validate our predictors (Methods). We first validated targeted therapies with documented modes of action. We started with the TRANSACT predictor of response for Afatinib, a small molecule inhibitor of the EGFR family, which includes HER2 (**Figure 5**A). We performed a gene set enrichment analysis of the linear terms that constitutes to 80% of the predictor (**Supp. Table 1**). Most enriched gene sets are related to breast cancer subtypes as defined by *Charafe and colleagues*[44] where, contrary to the definition based on the intrinsic breast cancer subtypes, the Luminal subtype contains both ER+ and HER2+ tumors. The top ranked gene set amongst the genes associated with sensitivity (genes with a negative coefficient in the predictor) are genes associated with the "Luminal" subtypes (FDR < 0.001). Conversely, genes associated with resistance (genes with a positive coefficient in the predictor) show enrichment for the "Mesenchymal" molecular signatures, shared by basal and mesenchymal subtypes. This corresponds with HER2 negative samples, which is in line with our expectation as absence of the drug target would indicate lack of response. Similarly, in the TRANSACT response predictor for Gefitinib (EGFR inhibitor) the genes constituting the linear portion and associated with sensitivity (negative predictor coefficients) show an enrichment for genes downregulated in Gefitinib resistant tumors (**Figure 5**B, **Supp. Table 2**).

Cytotoxic drugs such as Gemcitabine or Paclitaxel have complex modes of actions involving different pathways, crosstalk between which remains challenging to understand. Since the predictions of these two drugs showed a significant association

in both PDXs and patients, we set out to interpret the mechanisms of sensitivity or

resistance inferred by our predictor. In Gemcitabine (**Supp. Table 3**), we observe that

over-expression of the CDC42 pathway is a significant marker of resistance (FDR =

0.012, **Figure 5**C.1.) together with pathways linked to microtubule formation and cell

migration (**Supp Figure 13**), both known to be promoted by CDC42[45]. Together, these

enriched pathways highlight CDC42 over-expression as a potential mechanism of

Gemcitabine resistance, which suggests the use of CDC42 inhibitors[46,47] for

Gemcitabine-resistant tumors. Another interesting finding is the significant enrichment

of TNFα signaling in the sensitive portion (FDR=0.046) (**Figure 5**C.1.). A clinical trial

has shown that co-administration of TNF with gemcitabine improves patient survival

and further inhibits tumor growth[48], lending additional credence to this finding. Last,

we observe a concentration of sensitive interactions involving BLK, a pro-apoptotic

Src-proto oncogene involved in B-cell signaling and differentiation (**Figure 5**C.2.).

Since hematopoietic cell lines respond better to Gemcitabine, these interactions can

either act as a tissue-type marker or could potentially represent a sensitive pathway.

Finally, we looked for enriched pathways in Paclitaxel predictor (**Figure 5**D.1., **Supp.**

Table 4) and observed three potential mechanisms of resistance. We first observe

that the resistant linear coefficients are significantly enriched in genes linked to

silencing of YBX1[49] (FDR=0.106), a gene associated with proliferation in certain tumor

types[50]. In ovarian cancer, YBX1 has been shown to regulate ABCB1 expression

levels, a gene related to Paclitaxel resistance[52–56]. Our pan-cancer analysis therefore

further supports the role of drug transporters in Paclitaxel resistance. Second, we

observe a significant resistant enrichment for PI3K activation (FDR=0.18), which is

corroborated by the observed activation of PI3K/AKT/mTOR signaling pathway in

Paclitaxel-resistant cancer cells[56,57]. Moreover, a recent investigation suggests that

PI3K catalytic subunits can regulate ABCB1 expression[58]. Finally, when it comes to

the non-linear part, we observe a concentration of Fibroblast growth factors

interactions in the resistant parts of non-linearities, in particular FGF3, FGF20 and

FGF8 and FGF4 (**Figure 5**D.2., **Supp. Table 4)**. This behavior, although suggested

by previous studies[59,60], is all the more interesting as cell lines do not contain any

micro-environment that would elicit such resistance.

Discussion

We introduced an approach to integrate pre-clinical and clinical data in a fully

unsupervised way. Our approach geometrically aligns sample-to-sample similarity

matrices and extracts directions of important variations for both datasets, without

requiring any sample-level pairing. By performing a geometrical alignment instead of

a direct distribution comparison, our approach limits the effect of a potential sample

selection bias. This geometrical alignment is implicitly performed in a space induced

by our similarity function, which enables the integration of various non-linearities,

corresponding to hypothesis made on the system. Although we restricted ourselves to

a Gaussian similarity, designing similarity functions that incorporate a wide range of

prior knowledge is a potentially promising avenue. Learning the similarity matrix, e.g.

using multiple kernel learning[61] or deep learning methods such as variational auto-

encoders[62], may also help increase performance.

TRANSACT compares directions computed using Kernel PCA, but our approach can

be extended to other basis expansion methods by modulating the way the coefficients

α^s and α^t are computed. More generally, our method is versatile, generalizable, and

can be applied beyond the scope of our study, e.g. to integrate single cell sequencing

data.

We showed that the consensus features can be used to build translatable predictors

of drug response. Although we do not require a strong covariate shift assumption as

in a previous study[63], we do assume the functions modeling the response from these

consensus features follow the same monotonicity in pre-clinical models and human

tumors. This assumption, albeit reasonable, may be questioned.

In this study, we limited ourselves to gene expression. Making use of other genomics

levels – e.g. mutations, copy number, methylation, chromatin accessibility – may help

refine the prediction by providing additional signal. The integration of our approach

with multi-omics integration strategies[64,65] may offer a solution to the translation of

multi-omics signatures.

Finally, we were able to predict response in patients that received a particular

treatment, either as monotherapy or in combination with other therapies, even though

the cell line predictors were trained on monotherapies only. We envision that large-

scale, cell line combination screens that take potential synergistic effects between

drugs into account will provide training data for the next generation of cell line

predictors to be transferred to human tumors in order to further enhance prediction of

patient response. Moreover, in this study, we focus on cytotoxic and targeted

therapies, still widely used in the clinic. The recent advent of immuno-therapies calls

for methods with the ability to predict the clinical response from model systems. This

requires model systems capable of mimicking the action of the immune system and

screening technologies able to measure the response for large panels. We believe

that our approach can be extended to such problems once data is made available.

Methods

Public data download and pre-processing

GDSC dataset, download and processing.

We made use of the GDSC1000 cell line panel[14], which contains complete molecular profiles for 1,049 cell lines (**Supp Figure 2**). Gene expression is provided in the form of both read counts and FPKM. For both settings, we corrected the dataset for library-size using TMM[66] and log-transformed the corrected read counts[67,68]. Finally, we performed a gene-level mean-centering and standardization.

Novartis PDXE dataset, download and processing.

We made use of NIBR PDXE dataset for patient-derived xenografts[15], which contains the gene expression profiles of 399 PDXs (**Supp Figure 3**). Gene expression is provided in the form of FPKM. We corrected for library-size using TMM[66] and log-transformed the corrected read counts[67,68]. Finally, we performed a gene-level mean-centering and standardization.

TCGA dataset, download and processing.

We made use of the TCGA dataset for analyzing human biopsies[69], which comprises 10,347 human tumors (**Supp Figure 4**). Gene expression is provided in the forms of both read counts and FPKM and we used the same pre-processing pipeline as for GDSC. Response data have been obtained from *Ding et al*[70]*.* Following *Ding et al,* for each drug, we consider the response to patients who were administered a particular drug either as monotherapy or in combination with other drugs.

 Hartwig Medical Foundation dataset (HMF) download and processing.

We validated our approach on a cohort of 1,049 patients provided by the Hartwig

Medical Foundation – referred to as *HMF* (**Supp Figure 5**A). Gene expression was

measured for each metastatic sample prior to indicated drug regimen. We used

MultiQC for quality control[71], salmon v1.0.0 for alignment to reference transcriptome[72],

and finally edgeR for gene-level quantification[73]. Comparison with results obtained

using STAR[74] and featureCounts[75] shows high degree of concordance (**Supp Figure**

5D) and we used this comparison to refine our filtering. Read counts were then

processed using the same pipeline as in GDSC and TCGA.

Drug response was measured in 802 unique metastatic samples using the RECIST

criteria. Response was measured differently for each patient (**Supp Figure 5**B) with

most patients having one single measure of response around 10 to 15 weeks after

treatment started (**Supp Figure 5**C). Since we are interested in the response of the

drug given the molecular characterization measured, we considered for each patient

the first response after treatment. Since most drugs are administered in combination,

we considered, for each drug, all the patients that received it, whether in combination

with other drugs or as monotherapy. For instance, in the case of Gemcitabine, we

predicted drug response for all patients that received Gemcitabine as part of their

treatment strategies.

Measure of drug response

In our different analysis, we rely on drug response measurements, either to train a

predictor (GDSC), or to validate it (PDXE, TCGA and HMF). For cell lines (GDSC),

drug response is measured as *Area Under the drug response Curve*, referred to as

AUC. For PDX, drug response is measured as *Best Average Response*, which

corresponds to the relative variation of tumor volume induced by a certain treatment.

For both AUC and Best Average Response, large values are associated with

resistance. For TCGA and HMF, clinical responses have been measured using the

RECIST criteria[76]. Based on various metrics, patients get assigned to one of the

following four groups: *Complete Response* (CR), *Partial Response* (PR), *Stable*

Disease (SD) and *Progressive Disease* (PD). Following the division used in previous

works[21,23], we divide TCGA patients in two categories: *Responders* (CR and PR) and

Non Responders (SD and PD). For HMF, we discriminate between each possible

couple: PR vs PD, PR vs SD and SD vs PD. Since only a couple of patients showed

a complete response, we did not consider these patients.

Mathematical settings

We denote by p the number of genes. We consider one source dataset $\mathcal{X}_s =$

$\{x_1^s, \ldots, x_{n_s}^s\} \subset \mathbb{R}^p$ and one target dataset $\mathcal{X}_t = \{x_1^t, \ldots, x_{n_t}^t\} \subset \mathbb{R}^p$ with corresponding

source and target data matrices $X_s \in \mathbb{R}^{n_s \times p}$ and $X_t \in \mathbb{R}^{n_t \times p}$.

We consider a similarity function K -- also called *kernel function* -- that assigns to two

samples a scalar value that is large for similar samples and small for dissimilar

samples. In this work, we assume the kernel to be positive semi-definite (p.s.d.), which

implies[77] that there exists a Hilbert space $\mathcal{H}$ and a mapping $\varphi: \mathbb{R}^p \mapsto \mathcal{H}$ such that

$$\forall x, y \in \mathbb{R}^p, \qquad K(x, y) = \langle \varphi(x), \varphi(y) \rangle_{\mathcal{H}}. \qquad (1)$$

In particular, we use two kernels:

▪ Linear kernel: $K^{linear}(x, y) = x^T y$.

▪ Radial Basis Function, also referred to as "Gaussian": $K_\gamma^{rbf}(x,y) =$

$\exp(-\gamma\|x-y\|^2)$, with $\gamma > 0$.

We denote by K_s the matrix of similarity between source samples, K_t between target

samples and K_{st} the matrix of similarity between source and target, formally:

$$
\begin{aligned}
K_s &= \left(K\left(x_i^s, x_j^s\right)\right)_{1\le i,j\le n_s} \\
K_t &= \left(K\left(x_i^t, x_j^t\right)\right)_{1\le i,j\le n_t}. \\
K_{st} &= \left(K\left(x_i^s, x_j^t\right)\right)_{\substack{1\le i\le n_s \\ 1\le j\le n_t}}
\end{aligned}
\tag{2}
$$

Implicit mean centering

458   In a linear-setting, it is standard to perform a gene-level centering prior to any statistical

459   analysis. In the case of non-linear analysis, we perform a feature-level centering in the

460   embedding space $\mathcal{H}$ implicitly using Equation (1). We perform this implicit centering

461   independently in source and target datasets. As shown in (Subsection Supp. 2), the

462   corresponding kernel matrices are:

$$
\begin{aligned}
\widetilde{K}_s &= C_{n_s} K_s C_{n_s} \\
\widetilde{K}_t &= C_{n_t} K_t C_{n_t} \text{ , with } C_n = I_n - \frac{1}{n} 1_n^T 1_n \ (n \ge 1). \\
\widetilde{K}_{st} &= C_{n_s} K_{st} C_{n_t}
\end{aligned}
\tag{3}
$$

Kernel PCA by eigen-decomposition of centered kernel matrix for

capturing directions of principal variance

Using the embedding introduced in Equation (1), the similarity matrices from Equation

(3) can be seen as sample-covariance matrices and therefore decomposed to

compute principal components inside the embedded space $\mathcal{H}$, a procedure known as

Kernel PCA[36]. We perform Kernel PCA on source and target data independently to

compute d_s and d_t principal components respectively. Kernel PCA on the source

dataset consists of an eigen-decomposition of the matrix $\widetilde{K}_s$, yielding $\alpha^s \in \mathbb{R}^{d_s \times n_s}$,

while Kernel PCA on the target dataset decomposes $\widetilde{K}_t$, yielding $\alpha^t \in \mathbb{R}^{d_t \times n_t}$ (Def

Supp 3.1).

Comparing and aligning pre-clinical and tumor non-linear principal

components

Similarly to the "cosine similarity matrix" in other related works[24,28], we define the *non-*

linear cosine similarity matrix $\boldsymbol{M}^K$ as the matrix that geometrically compares the source

NLPCs to the target NLPCs (Def Supp 5.1). This matrix can be computed as follow

(Prop Supp 5.2):

$$\boldsymbol{M}^K = \alpha^s \widetilde{K}_{st} \alpha^{t^T} = \alpha^s C_{n_s} K_{st} C_{n_t} \alpha^{t^T}. \tag{4}$$

In a first step of our domain adaptation approach, we use the matrix $\boldsymbol{M}^K$ to align

NLPCs, yielding *non-linear principal vectors* $s_1, .., s_d$ for the source and $t_1, .., t_d$ for the

target domains, with $d = \min(d_s, d_t)$ (Def Supp. 4.1). These principal vectors are pairs

of vectors: one linear combination of source NLPCs and one linear combination of

target NLPCs, ordered by decreasing similarity with the first pair being the most

similar. The computation of these PVs relies on the Singular Value Decomposition[78]

of $\boldsymbol{M}^K$, $\boldsymbol{M}^K = \beta^s \Sigma \beta^{t^T}$ that helps us define the source and target *sample importance*

loadings ρ^s and ρ^t as follows (Def. Supp. 5.4)

495
$$\rho^s = \beta^{s^T}\alpha^s \text{ and } \rho^t = \beta^{t^T}\alpha^t. \tag{5}$$

497 We also define the principal angles $\theta_1,..,\theta_d$ as follows (Def. Supp. 5.6)

498
$$\forall k \in \{1,..,d\}, \qquad \cos\theta_k = \Sigma_{k,k}. \tag{6}$$

Owing to the duality elicited in Equation (1), these principal vectors can be seen both

as functions that map samples in cell-view to a one-dimensional vector, or a vector

onto which we project the sample embedding (Proposition Supp. 5.8).

Interpolation between kernel principal vectors for balancing effect of

source and target

Each pair of principal vectors contains two vectors that are geometrically similar.

Projection on them will create two highly correlated covariates that would not be

optimal for subsequent statistical treatment. In order to compute one vector out of each

pair, we interpolate between the source and the target PV within each pair (Def Supp

6.2). For the k^{th} PV, the interpolation is modulated by a coefficient τ_k that ranges

between 0, when the interpolation returns the source PV, and 1, when the interpolation

returns the target PV. This interpolation between vectors within each PV pair relies on

two functions $\Gamma(\tau) = [\Gamma_1(\tau_1),..,\Gamma_d(\tau_d)]^T$ and $\xi(\tau) = [\xi_1(\tau_1),..,\xi_d(\tau_d)]^T$ defined as

(Definition Supp. 6.1):

516
$$\forall k \in \{1,..,d\}, \quad \Gamma_k(\tau_k) = \frac{\sin(1-\tau_k)\theta_k}{\sin\theta_k} \quad \text{and} \quad \xi_k(\tau_k) = \frac{\sin\tau_k\theta_k}{\sin\theta_k}. \tag{7}$$

518   For a set of d interpolation coefficients $[\tau_1, \ldots, \tau_d]$, we compute the projection of source

519   and target datasets $F(\tau) \in R^{(n_s + n_t) \times d}$ as follows (Theorem Supp. 6.6)

521
$$F(\tau) = \begin{bmatrix} K^s & K^{st} \\ K^{ts} & K^t \end{bmatrix} \begin{bmatrix} C_{n_s} & 0 \\ 0 & C_{n_t} \end{bmatrix} \begin{bmatrix} \rho^{sT} & 0 \\ 0 & \rho^{tT} \end{bmatrix} \begin{bmatrix} \Gamma(\tau) \\ \xi(\tau) \end{bmatrix}. \tag{8}$$

Such an interpolation between PVs balances the effect of source and target datasets.

We prove that, in the case of a linear kernel, our interpolation scheme is equivalent to

the one from previous approaches[26,79] (Supp Subsection 8).

Within each pair of PVs, we select one intermediate representation where the source

and target projections match the most. For the k^{th} PV-pair, we compare the source

and target projected data using a Kolmogorov-Smirnov statistic and select the

interpolation coefficient τ_k^* where the statistic is minimal. We obtain a set of optimal

interpolation coefficients $\tau^* \in [0,1]^d$ when, for each PV, source and target influence

are balanced. We call the corresponding vector *consensus features.* These consensus

features show the minimal difference between source and target domain, a theoretical

necessary condition for domain adaptation[80].

Prediction using Elastic Net

In order to predict drug response, we use Elastic Net regression[81]. Elastic Net is a

linear model that imposes two penalties on the coefficients to predict: an ℓ_1 penalty

that leads to a sparse model and an ℓ_2 penalty that jointly shrinks correlated features.

We chose Elastic Net first because it has repeatedly been shown in the drug response

prediction literature to give equivalent, if not better, predictive performance compared

to complex models[16,18,38]. Using a linear classifier limits the complexity and therefore

makes the transfer more robust in practice.

Taylor expansion of the similarity function for interpretability of the model

In the case of RBF, we perform a PCA in an infinite-dimensional feature space.

Although this space cannot be analytically computed, it can be approximated using a

Taylor expansion[82] (Subsection Supp. 7). For the q^{th} consensus feature, we

differentiate three kinds of contributions (Def Supp. 7.4):

- *Offset $\mathcal{O}_q$:* a Gaussian term that models the squared depth. For each sample

$x \in R^p$, it corresponds to $\exp(-\gamma \|x\|^2)$.

- *Linear contributions $\left(\mathcal{L}_{q,j}\right)_{1 \le j \le p}$:* a linear term, resembling the expression of one

gene. For each sample $x \in R^p$ and gene $j \in \{1,..,p\}$, it corresponds to

$x_j \exp(-\gamma \|x\|^2)$.

- *Interaction terms $\left(\mathcal{I}_{q,j,k}\right)_{1 \le j,k \le p}$:* an interaction term that expresses the product

of two genes. For each sample $x \in R^p$ and gene $j, k \in \{1,..,p\}$, it corresponds

to $x_j x_k \exp(-\gamma \|x\|^2)$.

These contributions can be computed from sample importance loadings of consensus

features (Prop. Supp. 7.7). We consider the contributions' sum-of-squares as a

geometrical proportion since these sum up to one (Def. Supp. 7.8).

In order to look for enrichment in a particular consensus feature, we look for

enrichment of particular gene sets[83]. Specifically, for the linear contribution, we

compute the loading of all linear terms (Equation Supp. 48) corresponding each to one

gene and we performed a Pre-Ranked gene set enrichment analysis with FDR

correction at 20% and 1000 permutations. Since these loadings correspond to a

Euclidean geometric proportion, we used a squared statistic to compare them.

Comparison to competing approaches

We compare TRANSACT to four different approaches. The two first approaches

consist in applying a regression model trained on a source dataset to a target dataset

without any correction; we use one linear Elastic Net model (referred to as *Elastic Net*)

and a non-linear neural network model (referred to as *Deep Learning*). In both cases,

we perform a grid-search 5-fold cross-validation on cell lines to select the model with

the best performance: on *ElasticNet* we vary the ℓ_1 ratio and the total regularization ;

on *Deep Learning*, following the protocol from *Sakellaropoulos et al*[21], we use an

hyperbolic-tangent activation function while varying the global network structure, the

ℓ_2 penalty and the input and output dropout levels (**Supp Table** 5).

The other two approaches first correct the signal and then train a regression model.

The third approach (referred to as *ComBat + Deep Learning*) reproduces the approach

from *Sakellaropoulos et al*[21] by first performing a ComBat technical batch effect

correction between source and target, and then applying a neural network on the

corrected signal, similar to *Deep Learning* (**Supp Table** 6). The last competing

approach, referred to as *PRECISE,* consists in using a linear similarity function

followed by an Elastic Net model, which is equivalent to PRECISE (**Supp Material**).

For the two deep learning approaches, we first performed cross-validation on the

source dataset (with or without correction) to select the hyper-parameters and the

network structures with the largest predictive performance. We then re-initialize the

network and train it on the complete GDSC dataset.

Availability of data and materials

TRANSACT is available as a Python 3.6 module (https://github.com/NKI-
CCB/TRANSACT). All our experiments are reproducible and use state-of-the-art
libraries[84–89] (https://github.com/NKI-CCB/TRANSACT_manuscript). The dataset(s)
supporting the conclusions of this article are available in the "transact_manuscript"
repository of the aforementioned GitHub page, except for the HMF data, which can
only be obtained upon request to the Hartwig Medical Foundation.

Acknowledgment

This publication and the underlying study have been made possible partly on the basis
of the data that Hartwig Medical Foundation and the Center of Personalised Cancer
Treatment (CPCT) have made available to the study.

We thank Mirrelijn van Nee (VUmc), Sander Canisius (NKI), Osman Kayhan (TU
Delft), Tycho Bismeijer (NKI), Stavros Makrodimitris (TU Delft), Tesa Severson (NKI),
Joe Siefert (NKI), Joris van de Haar (NKI), Ahmed Mahfouz (LUMC), Tamim Abdelaal
(TU Delft), David Tax (TU Delft), Bram Thijssen (NKI) and Ziqi Wang (TU Delft) for
useful discussions. We thank Mohammed Charrout (TU Delft), Marie Corradi (NKI),
Lisa Koob (NKI) and Wouter Kouw (TU Eindhoven) for careful reading of the
manuscript.
This work was supported by ZonMw TOP grant COMPUTE CANCER [40-00812-98-
16012].

Authors contribution

S.M., L.F.A.W., M.J.T.R., M.L. and M.A.vd.W. designed the study. S.M. performed the experiments. L.F.A.W., M.J.T.R. and M.L. supervised the experiments. S.M., L.F.A.W., M.J.T.R., M.L. and M.A.vd.W. analyzed the results. S.M. and M.L. developed the mathematical framework. S.M. developed the software package. S.M. and D.J.V. aligned the HMF data. K.M. and A.G. interpreted the GSEA results. S.M., L.F.A.W., M.J.T.R. and M.L. wrote the manuscript. All authors read and approved the manuscript.

Competing interests

L.F.A.W. received project funding from Genmab BV.

1.   Slamon, D. J. *et al.* Numb Er 11 Use of Chemotherapy Plus a Monoclonal

Antibody Against Her2. *N. Engl. J. Med.* **344**, 783–792 (2001).

2.   Davies, H. *et al.* Mutations of the BRAF gene in human cancer. *Nature* **417**,

949–954 (2002).

3.   Hughes, T. P. *et al.* Frequency of major molecular responses to imatinib or

interferon alfa plus cytarabine in newly diagnosed chronic myeloid leukemia.

N. Engl. J. Med. **349**, 1423–1432 (2003).

4.   Kalamara, A., Tobalina, L. & Saez-Rodriguez, J. How to find the right drug for

each patient? Advances and challenges in pharmacogenomics. *Curr. Opin.*

Syst. Biol. **10**, 53–62 (2018).

5.   Prahallad, A. *et al.* Unresponsiveness of colon cancer to BRAF(V600E)

inhibition through feedback activation of EGFR. *Nature* **483**, 100–104 (2012).

6.   Valery, P. *Mauvaises pensées et autres*. (1941).

7.   Ben-David, U. *et al.* Patient-derived xenografts undergo mouse-specific tumor

evolution. *Nat. Genet.* **49**, 1567–1575 (2017).

8.   Ben-David, U. *et al.* Genetic and transcriptional evolution alters cancer cell line

drug response. *Nature* **560**, 325–330 (2018).

9.   Gillet, J. P. *et al.* Redefining the relevance of established cancer cell lines to

the study of mechanisms of clinical anti-cancer drug resistance. *Proc. Natl.*

Acad. Sci. U. S. A. **108**, 18708–18713 (2011).

10. Gillet, J. P., Varma, S. & Gottesman, M. M. The clinical relevance of cancer

cell lines. *J. Natl. Cancer Inst.* **105**, 452–458 (2013).

11. Mak, I. W. Y., Evaniew, N. & Ghert, M. Lost in translation: Animal models and

clinical trials in cancer treatment. *Am. J. Transl. Res.* **6**, 114–118 (2014).

12. Brubaker, D. K. & Lauffenburger, D. A. Translating preclinical models to

humans. *Science (80-.).* **367**, 742–743 (2020).

13.   Webber, J. T., Kaushik, S. & Bandyopadhyay, S. Integration of Tumor

Genomic Data with Cell Lines Using Multi-dimensional Network Modules

Improves Cancer Pharmacogenomics. *Cell Syst.* **7**, 526-536.e6 (2018).

14.   Iorio, F. *et al.* A Landscape of Pharmacogenomic Interactions in Cancer. *Cell*

(2016). doi:10.1016/j.cell.2016.06.017

15.   Gao, H. *et al.* High-throughput screening using patient-derived tumor

xenografts to predict clinical trial drug response. *Nat. Med.* **21**, 1318–1325

(2015).

16.   Costello, J. C. *et al.* A community effort to assess and improve drug sensitivity

prediction algorithms. *Nat. Biotechnol.* **32**, 1202–1212 (2014).

17.   Ali, M. & Aittokallio, T. Machine learning and feature selection for drug

response prediction in precision oncology applications. *Biophys. Rev.* **11**, 31–

39 (2019).

18.   Jang, I. S., Neto, E. C., Guinney, J., Friend, S. H. & Margolin, A. A. Systematic

assessment of analytical methods for drug sensitivity prediction from cancer

cell line data. *Pacific Symp. Biocomput.* **23**, 1–7 (2013).

19.   Geeleher, P., Cox, N. J. & Huang, R. S. Clinical drug response can be

predicted using baseline gene expression levels and in vitro drug sensitivity in

cell lines. *Genome Biol.* **15**, 1–12 (2014).

20.   Geeleher, P. *et al.* Discovering novel pharmacogenomic biomarkers by

imputing drug response in cancer patients from large genomics studies.

Genome Res. **27**, 1743–1751 (2017).

21.   Sakellaropoulos, T. *et al.* A Deep Learning Framework for Predicting

Response to Therapy in Cancer. *Cell Rep.* **29**, 3367-3373.e4 (2019).

22. Kurilov, R., Haibe-Kains, B. & Brors, B. Assessment of modelling strategies for

drug response prediction in cell lines and xenografts. *Sci. Rep.* **10**, 2849

(2020).

23. Noghabi, H. S., Peng, S., Zolotareva, O., Collins, C. C. & Ester, M. AITL:

Adversarial Inductive Transfer Learning with input and output space adaptation

for pharmacogenomics. *bioRxiv* 2020.01.24.918953 (2020).

doi:10.1101/2020.01.24.918953

24. Mourragui, S., Loog, M., van de Wiel, M. A., Reinders, M. J. T. & Wessels, L.

F. A. PRECISE: a domain adaptation approach to transfer predictors of drug

response from pre-clinical models to tumors. *Bioinformatics* **35**, i510–i519

(2019).

25. Pan, S. J., Kwok, J. T. & Yang, Q. Transfer Learning via Dimensionality

Reduction.

26. Gong, B., Shi, Y., Sha, F. & Grauman, K. Geodesic flow kernel for

unsupervised domain adaptation. *Proc. IEEE Comput. Soc. Conf. Comput. Vis.*

Pattern Recognit. 2066–2073 (2012). doi:10.1109/CVPR.2012.6247911

27. Gopalan, R., Li, R. & Chellappa, R. Domain adaptation for object recognition:

An unsupervised approach. *Proc. IEEE Int. Conf. Comput. Vis.* 999–1006

(2011). doi:10.1109/ICCV.2011.6126344

28. Fernando, B., Habrard, A., Sebban, M. & Tuytelaars, T. Unsupervised visual

domain adaptation using subspace alignment. *Proc. IEEE Int. Conf. Comput.*

Vis. 2960–2967 (2013). doi:10.1109/ICCV.2013.368

29. Kouw, W. & Loog, M. A review of domain adaptation without target labels.

IEEE Trans. Pattern Anal. Mach. Intell. **PP**, 1–1 (2019).

30. Hofmann, T., Schölkopf, B. & Smola, A. J. Kernel methods in machine

698  learning. *Ann. Stat.* **36**, 1171–1220 (2008).

699 31. Vert, J.-P., Koji Tsuda & Schölkopf, B. *Kernel methods in computational*

700  *biology*. (MIT Press, 2004).

701 32. He, X., Folkman, L. & Borgwardt, K. Kernelized rank learning for personalized

702  drug recommendation. *Bioinformatics* **34**, 2808–2816 (2018).

703 33. Ammad-Ud-Din, M. *et al.* Drug response prediction by inferring pathway-

704  response associations with kernelized Bayesian matrix factorization.

705  *Bioinformatics* **32**, i455–i463 (2016).

706 34. Li, Y., Wu, F. X. & Ngom, A. A review on machine learning principles for multi-

707  view biological data integration. *Brief. Bioinform.* **19**, 325–340 (2018).

708 35. Paltun, B. G., Mamitsuka, H. & Kaski, S. Improving drug response prediction

709  by integrating multiple data sources : matrix factorization , kernel and network-

710  based approaches. **00**, 1–14 (2019).

711 36. Schölkopf, B., Smola, A. J. & Müller, K.-R. Nonlinear Component Analysis as a

712  Kernel Eigenvalue Problem. *Neural Comput.* **10**, 1299–1319 (1998).

713 37. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-

714  1902.e21 (2019).

715 38. Smith, A. M. *et al.* Standard machine learning approaches outperform deep

716  representation learning on phenotype prediction from transcriptomics data.

717  *BMC Bioinformatics* **21**, 1–18 (2020).

718 39. Aben, N., Vis, D. J., Michaut, M. & Wessels, L. F. A. TANDEM: A two-stage

719  approach to maximize interpretability of drug response models based on

720  multiple molecular data types. *Bioinformatics* **32**, i413–i420 (2016).

721 40. Aben, N. *et al.* ITOP: Inferring the topology of omics data. *Bioinformatics* **34**,

722  i988–i996 (2018).

41.   Hoogstraat, M. *et al.* Genomic and transcriptomic plasticity in treatment-naïve

ovarian cancer. *Genome Res.* **24**, 200–211 (2014).

42.   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation

and Projection for Dimension Reduction. (2018).

43.   Warren, A. *et al.* Global computational alignment of tumor and cell line

transcriptional profiles. *bioRxiv* 2020.03.25.008342 (2020).

doi:10.1101/2020.03.25.008342

44.   Charafe-Jauffret, E. *et al.* Gene expression profiling of breast cell lines

identifies potential new basal markers. *Oncogene* 2273–2284 (2006).

doi:10.1038/sj.onc.1209254

45.   Cau, J. & Hall, A. Cdc42 controls the polarity of the actin and microtubule

cytoskeletons through two distinct signal transduction pathways. *J. Cell Sci.*

118, 2579–2587 (2005).

46.   Guo, Y. *et al.* R-ketorolac targets Cdc42 and Rac1 and alters ovarian cancer

cell behaviors critical for invasion and metastasis. *Mol. Cancer Ther.* **14**, 2215–

2227 (2015).

47.   Del Mar Maldonado, M. & Dharmawardhane, S. Targeting rac and Cdc42 GT

pases in cancer. *Cancer Res.* **78**, 3101–3111 (2018).

48.   Murugesan, S. R. *et al.* Combination of human tumor necrosis factor-alpha

(hTNF-α) gene delivery with gemcitabine is effective in models of pancreatic

cancer. *Cancer Gene Ther.* **16**, 841–847 (2009).

49.   Basaki, Y. *et al.* Akt-dependent nuclear localization of Y-box-binding protein 1

in acquisition of malignant characteristics by human ovarian cancer cells.

Oncogene **26**, 2736–2746 (2007).

50.   Frye, B. C. *et al.* Y-box protein-1 is actively secreted through a non-classical

pathway and acts as an extracellular mitogen. *EMBO Rep.* **10**, 783–789

(2009).

51. Housman, G. *et al.* Drug resistance in cancer: An overview. *Cancers (Basel).*

6, 1769–1792 (2014).

52. Goldstein, L. J. MDR1 gene expression in solid tumours. *Eur. J. Cancer* **32**,

1039–1050 (1996).

53. Vaidyanathan, A. *et al.* ABCB1 (MDR1) induction defines a common

resistance mechanism in paclitaxel- and olaparib-resistant ovarian cancer

cells. *Br. J. Cancer* **115**, 431–441 (2016).

54. Christie, E. L. *et al.* Multiple ABCB1 transcriptional fusions in drug resistant

high-grade serous ovarian and breast cancer. *Nat. Commun.* **10**, 5–14 (2019).

55. Patch, A. M. *et al.* Whole-genome characterization of chemoresistant ovarian

cancer. *Nature* **521**, 489–494 (2015).

56. Chen, D. *et al.* Dual PI3K/mTOR inhibitor BEZ235 as a promising therapeutic

strategy against paclitaxel-resistant gastric cancer via targeting

PI3K/Akt/mTOR pathway article. *Cell Death Dis.* **9**, (2018).

57. Hu, L., Hofmann, J., Lu, Y., Mills, G. B. & Jaffe, R. B. Inhibition of

phosphatidylinositol 3′-kinase increases efficacy of paclitaxel in in vitro and in

vivo ovarian cancer models. *Cancer Res.* **62**, 1087–1092 (2002).

58. Zhang, L. *et al.* The PI3K subunits, P110α and P110β are potential targets for

overcoming P-gp and BCRP-mediated MDR in cancer. *Mol. Cancer* **19**, 1–18

(2020).

59. Gan, Y., Wientjes, M. G. & Au, J. L. S. Expression of basic fibroblast growth

factor correlates with resistance to paclitaxel in human patient tumors. *Pharm.*

Res. **23**, 1324–1331 (2006).

773 60. Kim, S. H. *et al.* BGJ398, a pan-FGFR inhibitor, overcomes paclitaxel

774   resistance in urothelial carcinoma with FGFR1 overexpression. *Int. J. Mol. Sci.*

775   **19**, (2018).

776 61. Manica, M., Cadow, J., Mathis, R. & Rodríguez Martínez, M. PIMKL: Pathway-

777   Induced Multiple Kernel Learning. *npj Syst. Biol. Appl.* **5**, (2019).

778 62. Lin, X., Duan, Y., Dong, Q., Lu, J. & Zhou, J. Deep variational metric learning.

779   in *ECCV* 714–729 (2018). doi:10.1007/978-3-030-01267-0_42

780 63. Rampášek, L. Latent variable models for drug response prediction and genetic

781   testing. *PhD Thesis* (2019). doi:.1037//0033-2909.I26.1.78

782 64. Wang, B. *et al.* Similarity network fusion for aggregating data types on a

783   genomic scale. *Nat. Methods* **11**, 333–337 (2014).

784 65. Sharifi-Noghabi, H., Zolotareva, O., Collins, C. C. & Ester, M. MOLI: Multi-

785   omics late integration with deep neural networks for drug response prediction.

786   *Bioinformatics* **35**, i501–i509 (2019).

787 66. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential

788   expression analysis of RNA-seq data. *Genome Biol.* **11**, (2010).

789 67. Dillies, M. A. *et al.* A comprehensive evaluation of normalization methods for

790   Illumina high-throughput RNA sequencing data analysis. *Brief. Bioinform.* **14**,

791   671–683 (2013).

792 68. Zwiener, I., Frisch, B. & Binder, H. Transforming RNA-Seq data to improve the

793   performance of prognostic gene signatures. *PLoS One* **9**, 1–13 (2014).

794 69. Weinstein, J. N. *et al.* The cancer genome atlas pan-cancer analysis project.

795   *Nat. Genet.* **45**, 1113–1120 (2013).

796 70. Ding, Z., Zu, S. & Gu, J. Evaluating the molecule-based prediction of clinical

797   drug responses in cancer. *Bioinformatics* **32**, 2891–2895 (2016).

71.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize

analysis results for multiple tools and samples in a single report. *Bioinformatics*

32, 3047–3048 (2016).

72.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon

provides fast and bias-aware quantification of transcript expression. *Nat.*

Methods **14**, 417–419 (2017).

73.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics **26**, 139–140 (2009).

74.   Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,

15–21 (2013).

75.   Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**,

923–930 (2014).

76.   Therasse, P. tumors. European Organization for Research and Treatment of

Cancer, National Cancer Institute of the United States, National Cancer

Institute of Canada. *J natl Cancer inst* **92**, 205–16 (2000).

77.   Aronszajn, N. Theory of Reproducing Kernels. *Trans. Am. Math. Soc.* **68**, 337

(1950).

78.   Golub, G. H. & Van Loan, C. F. *Matrix Computations*. (2013).

79.   Gopalan, R., Li, R. & Chellappa, R. by Generating Intermediate Data

Representations. *IEEE Trans. Pattern Anal. Mach. Intell.* **36**, 2288–2302

(2014).

80.   Ben-David, S. *et al.* A theory of learning from different domains. *Mach. Learn.*

79, 151–175 (2010).

81.   Zou, H. & Hastie, T. Regularization and variable selection via the elastic net

Hui. *J. Stat. Soc. Ser. B* **67**, 301–320 (2005).

82.   Steinwart, I., Hush, D. & Scovel, C. An Explicit Description of the Reproducing

Kernel Hilbert Spaces of Gaussian RBF Kernels. *IEEE Trans. Inf. Theory* **52**,

4635–4643 (2006).

83.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based

approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad.*

Sci. U. S. A. **102**, 15545–15550 (2005).

84.   Van Der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: A

structure for efficient numerical computation. *Comput. Sci. Eng.* **13**, 22–30

(2011).

85.   Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in

Python. *Nat. Methods* **17**, 261–272 (2020).

86.   Hunter, J. D. Matplotlib: A 2D graphics environment. *Comput. Sci. Eng.* **9**, 99–

104 (2007).

87.   Varoquaux, G. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn.*

Res. **19**, 29–33 (2015).

88.   Tietz, M., Fan, T. J., Nouri, D., Bossan, B. & skorch Developers. skorch: A

scikit-learn compatible neural network library that wraps PyTorch. (2017).

89.   Behdenna, A., Haziza, J., Azencott, C.-A. & Nordor, A. pyComBat, a Python

tool for batch effects correction in high-throughput molecular data using

empirical Bayes methods. *bioRxiv* (2020). doi:10.1101/2020.03.17.995431

Figure 1 TRANSACT: Generating non-linear manifold representations to transfer

predictors of response from pre-clinical models to human tumors. (**A**) Samples are

compared using a **similarity function** yielding similarity matrices between pre-clinical

models (source, K_s), between tumors samples (target, K_t) and between pre-clinical

models and tumors (K_{st}). (**B**) Using non-linear PCA, the pre-clinical and tumor

similarity matrices are independently decomposed into **non-linear principal**

components (NLPCs) geometrically represented by "sample importance scores"

(**Supp Figure 1A**) that represent the importance of each sample in each NLPC (α^s

and α^t, for source and target space, respectively). (**C**) Geometrical comparison of pre-

clinical and tumor NLPCs results in a **non-linear cosine similarity matrix M^K**. (**D**)

Alignment of NLPCs using the notion of **principal vectors** (**Supp Figure 1B**). (**E**)

Interpolation within each pair of vectors to select one vector per PV-pair that balances

the effect of pre-clinical and tumor signals: the **consensus features** (**Supp Figure**

1C). (**F**) Projection of each tumor and pre-clinical sample on the consensus features

to obtain **consensus scores**: scores that correspond to the activity of processes

conserved between tumors and pre-clinical models. (**G**) Finally, these scores can be

used as input to any predictive model, for instance to predict drug response based on

these consensus scores.

Figure 2 Impact of modeling non-linearities for drug response prediction transfer from cell lines to PDXs. (**A**) Main workflow of the prediction on PDXs. Using cell lines and PDX gene expression, we compute consensus features and project each dataset onto these. We then trained a regression model (Elastic Net) on cell lines projected scores to predict response, measured as Area Under the drug response Curve (AUC). Finally, we used this regression model to predict drug response in PDXs and correlate the predicted AUC to the known best average response. (**B**) Proportion of non-linearities induced by the Gaussian similarity function as a function of the scaling factor γ. For different values of γ, we computed the average contribution over all consensus features of offset, linear, interaction and higher order features (Methods). Offset is here to be understood as the exponential of the squared depth and does not correspond to a constant term. We finally evaluated the response prediction on the PDX models for different values of γ, and for four competing approaches: two methods (*ElasticNet* and *Deep Learning*) that do not model the transfer, and two transfer

learning methods (ComBat followed by Deep Learning, and *PRECISE*). We report

results for Erlotinib (**C**), Cetuximab (**D**), Gemcitabine (**E**), Afatinib (**F**), Paclitaxel (**G**),

Trametinib (**H**) and Ruxolitinib (**I**). Concordance on PDX is measured as the Spearman

correlation between predicted AUC and Best Average Response.

Figure 3 <u>Pan-cancer consensus features between cell lines and tumors conserve</u>

<u>tissue type information</u>. We used cell lines (GDSC) as source data and computed

two sets of consensus features with two different target datasets: primary tumors

(TCGA, A and C) and metastatic lesions (HMF, B and D). (**A**) Proportion of linear

and non-linear contributions to each of the 30 GDSC-to-TCGA consensus features.

(**B**) Proportion of linear and non-linear contributions to each of the 20 GDSC-to-HMF

consensus features. (**C**) UMAP plot of primary tumors (TCGA, 21 tissues) and cell

lines (GDSC, 22 tissues) projected on the consensus features, using the same

parameters as selected in **Figure 2**. (**D**) UMAP plot of metastatic lesions (HMF) and

cell lines, colored by primary tissue for both HMF and GDSC. For both UMAP plots,

the full legend can be found in **Supp Figure 10**A.

Figure 4 <u>Consensus features improve response prediction in patients.</u> (**A**) We used

consensus features computed between GDSC and each tumor dataset to train a

predictor for 17 drugs on TCGA and 5 drugs on HMG, using GDSC response only. We

then predicted the Area Under the drug response Curve (AUC) for each drug and

compared this predicted value to the observed clinical response using a one-sided

Mann-Whitney test (**Table 1**). We performed the same prediction tasks using two

state-of-the-art approaches (*ComBat + Deep Learning* and *PRECISE*, **Method**) and

summarized here the associations for the 3×22 predictors in a Volcano plot, with the

corresponding p-values on the y-axis and the effect-sizes on the x-axis. (**B**) Results

for the two Gemcitabine predictors. (**B**.1) For the Gemcitabine predictor on TCGA, we

compared predicted AUC for each patient to the known clinical response. (**B**.2) Results

for Gemcitabine predictor on HMF. (**C**.1) Results for Paclitaxel predictor on TCGA.

(**C**.2.) Results for Paclitaxel predictor on HMF. (**D**.1) Results for Carboplatin predictor

on TCGA. (**D**.2.) Results for Carboplatin predictor on HMF.

A. Gene Set Enrichment Analysis on linear portion (80%) of Afatinib TRANSACT predictor

B. Gene Set Enrichment Analysis on linear portion (80%) of Gefitinib TRANSACT predictor

C. Analysis of Gemcitabine TRANSACT predictor

C.1. Gene Set Enrichment Analysis on linear portion (72%)

C.2. Interaction terms (15%)

D. Analysis of Paclitaxel TRANSACT predictor

D.1. Gene Set Enrichment Analysis on linear portion (76%)

D.2. Interaction terms (12%)

Figure 5 Interpretability of TRANSACT consensus features recapitulates modes of

action for Afatinib and Gefitinib, and highlights mechanisms of sensitivity and

resistance to Gemcitabine and Paclitaxel. (**A**) For the Afatinib (HER-2 protein kinase

inhibitor) response model trained on GDSC samples projected on GDSC-to-TCGA

consensus features, we show the contribution of each gene to the predictor (left):

positive (negative) weights in the predictor indicate that high (low) expression of the

genes leads to resistance (sensitivity) represented by larger (smaller) AUCs. We

performed a PreRanked gene set enrichment analysis (GSEA) on these weights and

highlight genes associated to two significantly enriched gene sets: "Charafe [...]

Luminal vs Mesenchymal UP" (center-left) and "Charafe [...] Luminal vs Mesenchymal

DOWN" (center-right). The two triangles on the right illustrate these two gene-set

distributions. (**B**) Interpretation of the linear part of the Gefitinib (EGFR inhibitor)

model. We display gene weights ordered by contribution (left) and the ranks of genes

known to be down regulated in Gefitinib resistant tumors (right). (**C**) Interpretation of

Gemcitabine predictor. (*C.1.*) We display the positions of genes associated to two

significantly enriched gene sets: "CDC42 pathway", associated to positive weights

(resistance), and "TNFα-signalling", associated to negative weights (sensitivity). (*C.2.*)

Interaction terms make 15% of Gemcitabine predictor. We show the distribution of

their weights annotated with the ten largest weights (resistance) and ten smallest

weights (sensitivity). (**D**) Interpretation of Paclitaxel predictor. (*D.1.*) We display the

positions of genes associated to two significantly enriched gene sets, both to

resistance: "Basaki YBX1 target UP" and "constitutive signaling by PI3K-aberrant

signaling". (*D.2.*) Interaction terms for Paclitaxel, that make 12% of the predictor,

alongside the top-10 resistant and top-10 sensitive interactions.

A. GDSC-to-TCGA								
GDSC		**TCGA**		**p-val [effect-size] on TCGA**				
Drug	**Samples**	**Drug**	**Samples**	**Elastic Net**	**Deep Learning**	**DL + ComBat**	**PRECISE**	**TRANS ACT**
Afatinib	800	Trastuzumab	16	**0.089 [0.82]**	**0.034 [0.93]**	**0.048 [0.89]**	**0.026 [0.96]**	**0.016 [1.00]**
Bleomycin	856	Bleomycin	53	0.15 [0.63]	0.19 [0.61]	0.41 [0.53]	0.11 [0.66]	0.091 [0.67]
Cetuximab	868	Cetuximab	19	0.12 [0.67]	0.52 [0.50]	0.39 [0.54]	0.45 [0.52]	0.077 [0.70]
Cisplatin	764	Cisplatin	308	**1.1E-6 [0.69]**	**1.8E-4 [0.64]**	**1.1E-2 [0.59]**	**2.2E-5 [0.66]**	**3.9E-7 [0.70]**
Cisplatin	764	Carboplatin	166	**0.042 [0.58]**	0.10 [0.56]	0.53 [0.50]	**0.024 [0.59]**	**0.0035 [0.63]**
Cyclophosphamide	747	Cyclophosphamide	102	0.57 [0.48]	0.67 [0.45]	0.89 [0.35]	0.11 [0.65]	0.64 [0.46]
Docetaxel	665	Docetaxel	102	0.62 [0.48]	0.27 [0.54]	0.54 [0.49]	0.78 [0.46]	0.11 [0.58]
Doxorubicin	871	Doxorubicin	101	0.81 [0.45]	**0.0055 [0.66]**	0.60 [0.48]	0.99 [0.32]	0.69 [0.47]
Etoposide	880	Etoposide	84	0.21 [0.58]	**0.0032 [0.76]**	**0.0076 [0.73]**	**0.0071 [0.73]**	**0.027 [0.68]**
5-Fluorouracil	801	Fluorouracil	186	0.22 [0.54]	0.72 [0.47]	0.79 [0.46]	0.80 [0.46]	0.35 [0.52]
Gemcitabine	752	Gemcitabine	156	0.25 [0.53]	**0.021 [0.59]**	**0.0057 [0.62]**	**0.029 [0.59]**	**0.0047 [0.62]**
Irinotecan	796	Irinotecan	25	0.81 [0.39]	0.42 [0.53]	0.68 [0.44]	0.54 [0.49]	0.49 [0.51]
Oxaliplatin	724	Oxaliplatin	66	0.38 [0.52]	**0.022 [0.65]**	**0.0099 [0.68]**	**0.028 [0.64]**	**0.035 [0.64]**
Paclitaxel	753	Paclitaxel	160	**0.032 [0.59]**	**0.025 [0.60]**	0.28 [0.53]	0.10 [0.56]	**0.0042 [0.63]**
Pemetrexed	898	Pemetrexed	38	0.16 [0.60]	0.39 [0.53]	0.55 [0.49]	0.16 [0.59]	0.24 [0.57]
Temozolomide	746	Temozolomide	96	0.56 [0.49]	0.29 [0.55]	0.47 [0.51]	0.59 [0.48]	0.19 [0.58]
Vinorelbine	746	Vinorelbine	30	0.31 [0.57]	**0.043 [0.72]**	0.053 [0.71]	0.19 [0.61]	0.42 [0.53]
B. GDSC-to-HMF								
GDSC		**HMF**		**p-val [effect-size] on HMF**				
Drug	**Samples**	**Drug**	**Samples**	**Elastic Net**	**Deep Learning**	**DL + ComBat**	**PRECISE**	**TRANS ACT**
Afatinib	800	Trastuzumab	25	0.18 [0.70]	0.069 [0.81]	0.23 [0.67]	**0.021 [0.93]**	**0.032 [0.89]**
Irinotecan	796	Irinotecan	67	0.060 [0.73]	0.060 [0.73]	0.060 [0.73]	0.10 [0.69]	**0.020 [0.80]**
Cisplatin	764	Carboplatin	64	0.23 [0.58]	0.051 [0.67]	0.094 [0.64]	0.59 [0.67]	**0.0045 [0.78]**
5-Fluorouracil	801	Fluorouracil	65	0.065 [0.68]	0.12 [0.64]	0.16 [0.62]	0.21 [0.59]	0.24 [0.58]
Paclitaxel	753	Paclitaxel	45	0.56 [0.48]	0.32 [0.56]	0.39 [0.54]	0.39 [0.53]	0.061 [0.70]
Gemcitabine	752	Gemcitabine	50	**0.039 [0.71]**	**0.014 [0.76]**	**0.039 [0.71]**	**0.0089 [0.78]**	**0.0042 [0.81]**

Table 1 Results of TRANSACT compared to four competing approaches – *Elastic Net*,

Deep Learning, *ComBat+DL*, and *PRECISE* – for 17 drugs on TCGA and 5 drugs on

HMF. For each drug, we train 5 predictors and compare in each scenario the predicted

AUC to the known clinical response using one-side Mann-Whitney test. The clinical

response corresponds to the following division: Responders and Non-Responders for

TCGA, PR and PD for HMF (Methods). For each predictor, we report the p-value and

the effect size (Area under the ROC, effect size associated with the Mann-Whitney

test) under bracket. Bold cells correspond to significant association (pval < 0.05). Red

cells correspond to significant associations with the largest effect size across the three

methods.

Supp Figure 1 <u>Visual explanation of geometric alignment</u>. **A**: Difference between importance scores (α^s, α^t) and projected scores. Since the space induced by the similarity function K is intractable, we use a dual representation of the NLPC in terms of samples: the importance scores. To project samples on NLPCs, one needs to compute the similarity between this sample and all of the samples used to gauge the NLPC. The projected score is obtained by taking the vector-product between this similarity vector and the importance scores. The same rational holds principal vectors that are represented by γ^s and γ^t. **B**: Visual example of principal vectors (PV). We here consider 3 genes (features) and 2 NLPCs. The pre-clinical (source) and tumor (target) NLPCs intersect in one direction, which form the pair of closest vectors: the first PV forms the pair of the two red vectors – although these are identical. The second pair of PVs is defined orthogonally to the red pair. This defines the green vectors (with a swap in direction for visual purposes). These pairs reconstruct the original NLPC spaces and are ordered by similarity. **C**: Interpolation between PVs. For one pair of PVs – e.g. the green one in B – source and target vectors are different. In order to generate one robust vector out of these two and avoid redundancy, we draw an arc between these two vectors. We then project source and target datasets onto these interpolated vectors and select one intermediate representation where source and

target projected signals are maximally matched. This optimal intermediate vector is

called the consensus feature.   50

Supp Figure 2 <u>Composition of the GDSC dataset (cell lines).</u> We make use of the

GDSC1000 cell line panel[14]. **A**: Number of cell lines per tissue type. **B**: Number of cell

lines screened for each drug that we used in our experiments.

Supp Figure 3 <u>Composition of the NIBR PDXE dataset (patient derived xenografts).</u>

We make use of the NIBR PDXE patient derived xenograft panel[15]. **A**: Number of

PDXs per tissue type. **B**: Number of unique PDXs screened for each drug that we

used in our experiments.

Supp Figure 4 <u>Structure of the TCGA dataset (primary tumors).</u> We make use of the

TCGA dataset for primary tumors. **A**: Number of samples per cancer type. **B**: For each

drug, number of samples with known response.

Supp Figure 5 <u>Structure of the HMF dataset (metastatic lesions).</u> We make use of the

Hartwig Medical Foundation (HMF) dataset for metastatic lesions. (**A**) Number of

samples per cancer type (primary tumor location). (**B**) For each patient, number of

response measurements made. For further analysis, we considered the first response

measure – i.e. first measure after treatment start. (**C**) Histogram of number of weeks

between treatment start and response measurement. (**D**) For each protein coding gene, we measure the Spearman correlation between read counts obtained using Salmon and STAR alignment tools using all samples in the HMF dataset. We then ranked genes based on the obtained Spearman correlation (x-axis) and plotted it against the mean-expression of these genes obtained using Salmon (y-axis). Since lowly concordant genes tend to have low expression, we put a threshold at $corr = 0.5$ and discarded genes below this threshold. (**E**) After the previous selection, we computed the sample-level Pearson and Spearman correlations between read counts obtained with STAR and Salmon. All samples but five show a correlation above 0.8 – these were discarded. We finally further restricted to genes from the mini-cancer genome.

1014

Supp Figure 6 <u>Analysis of consensus features between cell lines (GDSC) and PDXs</u>

1015

<u>with $\gamma = 0.0005$.</u> We use a Gaussian similarity matrix with hyper-parameter $\gamma = 0.0005$

1016

and run TRANSACT. (**A**) Cosine similarity between the 20 top source and target

1017

NLPCs. (**B**) Similarity between principal vectors (blue line) alongside the similarity

1018

obtained after gene-level permutation on GDSC (boxplots). (**C**) For each consensus

1019

feature, proportion of offset, linear and interaction term. (**D**) UMAP of data projected

1020

on the consensus features, colored by tissue of origin. (**E**) For each tissue type in

1021

PDXs, we compare the distances between corresponding PDXs with cell lines from

1022

the same tissue of origin (blue), or from another tissue (orange). (**F**) For the first

1023

consensus feature, sorted contribution of each linear features (i.e. gene, left) and

1024

interaction terms (right). (**G**) For the second consensus feature, sorted contribution of

1025

each linear features (i.e. gene, left) and interaction terms (right).

1026

1027

1028

1029

1030

Supp Figure 7 <u>Tissue clustering without domain adaptation and with PRECISE</u>

<u>alignment between GDSC and PDXE</u>. (**A**) UMAP plot of cell lines and PDXs colored

by tissue type without any domain-adaptation. Data was normalized prior to

performing UMAP: cell lines and PDXs were independently mean-centered and scaled

to unit variance. (**B**) UMAP plot of cell lines and PDXs colored by tissue type after

projection on consensus features obtained with linear PRECISE. (**C**) Comparison of

distances between PDXs and cell lines from the same tissue type (blue) or from a

different tissue type (orange) without domain adaptation. (**D**) Comparison of distances

when using linear PRECISE. We zoom in on lung (NSCLC) without domain adaptation

(**E**), with linear PRECISE (**F**) or with TRANSACT (**G**) using same setting as in **Supp**

Figure 6.

1042

1043

1044

Supp Figure 8 <u>Choice of the number of NLPCs and consensus features between</u>

<u>GDSC and TCGA. </u>(**A**) Cumulative sum of eigenvalues of $\widetilde{K_s}$ (GDSC) with $\gamma^* =$

5×10^{-4}. The cumulative sum increases steeply, reaches an inflexion points and then

follows an almost-linear behavior. We select all the NLPCs before this almost-linear

zone, corresponding to 75 NLPCs. (**B**) Cumulative sum of eigenvalues of $\widetilde{K_t}$ (TCGA)

with $\gamma^* = 5 \times 10^{-4}$. Following similar reasoning as in (**A**), we restrict the study to the

first 150 NLPCs. (**C**) Similarity between PVs when 75 NLPCs are considered for GDSC

and 150 for TCGA. We observe that the 33 first PVs have a similarity above 0.5 (our

cut-off) and round the selection to 30 PVs.

1054

1055

1056

Supp Figure 9 <u>Choice of the number of NLPCs and consensus features between</u>

<u>GDSC and HMF.</u> (**A**) Cumulative sum of eigenvalues of $\widetilde{K_s}$ (GDSC) with $\gamma^* = 5 \times 10^{-4}$.

The cumulative sum increases steeply, reaches an inflexion points and then follows

an almost-linear behavior. We select all the NLPCs before this almost-linear zone,

corresponding to 75 NLPCs. (**B**) Cumulative sum of eigenvalues of $\widetilde{K_t}$ (HMF) with $\gamma^* =$

5×10^{-4}. Following similar reasoning as in (**A**), we restrict the study to the first 75

NLPCs. (**C**) Similarity between PVs when 75 NLPCs are considered for both GDSC

and HMF. We observe that the 21 first PVs have a similarity above 0.5 (our cut-off)

and round the selection to 20 PVs.

1066

1067

1068

Supp Figure 10 <u>Pan-cancer consensus features between cell lines and tumors</u>

<u>conserve tissue type information (Supplement of Figure 3)</u> **A**: Legend of UMAP plots

for **Figure 3**C-D. **B**: UMAP plot of HMF metastatic lesions (same as Figure 3D)

colored by metastatic site. **C**: In TCGA, for each tumor type, distance between tumors

and cell lines from similar (blue) and non-similar (orange) tissue. **D**: In HMF, for each

primary tumor type, distance between metastatic sample and cell line from similar and

non-similar tissue of origin. **E**: In HMF, for each metastatic site, distance between

metastatic sample and cell line from tissue of origin similar (blue) or dissimilar from

the metastatic site.

1078

Supp Figure 11 <u>Impact of initialization on results for the *Deep Learning* (*DL*)</u>

<u>approach.</u> For each drug on TCGA and HMF, we considered the architecture and the

set of hyper-parameters with the lowest Mean Squared Error on GDSC given an

initialization. We then randomly generated 50 independent initializations of the

resulting networks and trained them using the GDSC data. Each of these trained

networks was then employed to predict the TCGA or HMF response. The resulting

prediction accuracies (area under the ROC) are plotted for the different drugs on the

TCGA and HMF data. **A**: Deep Learning prediction accuracies for each drug on TCGA

1087 (green boxplots). For comparison, the results from TRANSACT are displayed (blue

1088 curve). **B**: Pearson correlation of the Mean Square Error of the predictor on GDSC to

1089 the Area under the ROC of the same predictor on TCGA. **C**: Boxplot of performances

1090 on each HMF drug for 50 different initializations. **D**: Pearson correlation on HMF

1091 between MSE (GDSC) and Area under the ROC (HMF).

1092

1093

GDSC-to-TCGA

GDSC-to-HMF

1094

Supp Figure 12 <u>Impact of initialization on results for the *ComBat+DL* approach.</u> For each drug on TCGA and HMF, we considered the architecture and the set of hyper-parameters with the lowest Mean Squared Error on GDSC given an initialization. We then randomly generated 50 independent initializations of the resulting networks and trained them using the GDSC data. Each of these trained networks was then employed to predict the TCGA or HMF response. The resulting predictions accuracies (area under the ROC) are plotted for the different drugs on the TCGA and HMF data. **A**: *ComBat+DL* prediction accuracies for each drug on TCGA (green boxplots). For comparison, the results from TRANSACT are displayed (blue curve). **B**: Pearson

1104   correlation of the Mean Square Error of the predictor on GDSC to the Area under the

1105   ROC of the same predictor on TCGA. **C**: Boxplot of performances on each HMF drug

1106   for 50 different initializations. **D**: Pearson correlation on HMF between MSE (GDSC)

1107   and Area under the ROC (HMF).

1108

1109

1110

1111

1112 **Supp Figure 13** <u>Pathway enriched for resistant linear coefficients in GDSC-to-TCGA</u>

1113 <u>Gemcitabine drug response predictor</u>. Additional pathways significantly enriched in

1114 the linear part of the GDSC-to-TCGA predictor.