REV-ERB Agonism Improves Liver Pathology in a Mouse Model of NASH

Kristine Griffett[*], Gonzalo Bedia-Diaz, Bahaa Elgendy, and Thomas P. Burris

Center for Clinical Pharmacology, Washington University School of Medicine and St. Louis College of Pharmacy, St. Louis, MO

[*]Corresponding Author
E-mail: kgriffett@wustl.edu

Abstract

Non-alcoholic fatty liver disease (NAFLD) affects a significant number of people worldwide and currently there are no pharmacological treatments. NAFLD often presents with obesity, insulin resistance, and in some cases cardiovascular diseases. There is a clear need for treatment options to alleviate this disease since it often progresses to much more the much more severe non-alcoholic steatohepatitis (NASH). The REV-ERB nuclear receptor is a transcriptional repressor that regulates physiological processes involved in the development of NAFLD including lipogenesis and inflammation. We hypothesized that pharmacologically activating REV-ERB would suppress the progression of fatty liver in a mouse model of NASH. Using REV-ERB agonist SR9009 in a mouse NASH model, we demonstrate the beneficial effects of REV-ERB activation that led to an overall improvement of hepatic health by suppressing hepatic fibrosis and inflammatory response.

Introduction

Among the metabolic disorders, non-alcoholic fatty liver disease (NAFLD) is considered a hepatic manifestation of metabolic syndrome (MetS) and it is one of the prominent health challenges of the twenty-first century as NAFLD is the most prevalent liver disease worldwide affecting 25-30% of the general population and its prevalence could reach 70-90% in specific populations with comorbidities such as morbid obesity or type 2 diabetes mellitus [1-2]. NAFLD can often progress to non-alcoholic steatohepatitis (NASH), which is associated with progressive liver disease [1]. NASH has been mainly associated with higher morbidity and mortality than other diseases in the NAFLD spectrum and, although there are pharmacological therapies under clinical investigation for treatment of NASH [2], no drugs are approved by the Federal Drug Administration (FDA) or the European Medicines Agency (EMA) for the NASH treatment [3].

Nuclear receptors (NRs) are transcription factors generally activated by ligands and involved in diverse biological processes such as cell growth and differentiation, apoptosis, gene expression during

tumor formation and metabolism. They bind to specific sequences of DNA allowing them to regulate the expression of adjacent genes. Many diseases including NASH are directly or indirectly related to nuclear receptor signaling and many NRs have become favored targets for drug discovery [4]. NRs play an important role in liver diseases and they are key modulators in the onset and progression NAFLD, including the peroxisome proliferator-activated receptors (PPAR) α/β/γ; liver X receptors (LXR) α/β; farnesoid X receptors (FXR); constitutive androstane receptor (CAR); and pregnane X receptor (PXR). All of these NRs form obligate heterodimers with retinoid X receptor (RXR) α/β/γ in order to modulate corresponding target genes in the nucleus [5,6].

REV-ERB nuclear receptors (REV-ERBα and REV-ERBβ) are transcriptional repressors that regulate a variety of physiological processes including lipogenesis, inflammation, circadian regulation, and muscle regeneration and are expressed in all tissues but has significantly higher expression in liver, skeletal muscle, adipose tissue, and brain [7]. Although REV-ERBs play a regulatory role in hepatic metabolism, inflammation and lipogenesis, these NRs have yet to be validated as a potential therapeutic target for liver disease [8–11]. Here, we show that REV-ERB pan-agonist SR9009 treatment in *ob/ob* mice fed a high-fat, high-fructose (NASH) diet has beneficial effects and may provide some translational groundwork for further developing REV-ERB agonists for metabolic diseases, specifically NAFLD.

Materials and Methods

Animals

Animal studies were performed as previously described [12–14]. Briefly, six-week old B6 V-Lep[ob]/J (*ob/ob*) male mice were purchased from Jackson Labs (Bar Harbor, ME). Upon receipt, mice were housed individually in standard cages with huts and immediately placed on NASH diet (D09100301; Research Diets) [15]. Mice were maintained on this diet throughout the experiment. Mice were handled and weighed weekly while acclimating to the diet. At 12-weeks of age, mice were assigned into weight-matched groups (n = 7) and dosing began. Mice were weighed daily and food-intake was monitored

daily. At the termination of the study, mice were fasted and euthanized by CO_2 and blood was collected

by cardiac puncture for clinical chemistry analysis at Scripps Florida Metabolism Core (Roche COBAS

instrument) and ELISA analysis (EMD Millipore). Tissues were collected and flash-frozen in liquid

nitrogen for gene expression, or placed in 4% Paraformaldehyde (PFA) in PBS for paraffin-embedding

(Saint Louis University Histology Core) or 10% Neutral-Buffered Formalin (NBF) for cryo-sectioning.

For intraperitoneal glucose tolerance test (ipGTT), mice were transferred to clean cages and fasted for

12 hours overnight. Mice were weighed, baseline blood glucose was taken, then mice were given an

injection of glucose solution in PBS (2g/kg body weight). Blood glucose levels were subsequently

repeated at 30-, 60-, 90-, and 120-minutes following the injection of glucose. Mean blood glucose levels

(mg/dL) are reported as well as the area under the curve (AUC) which was analyzed by two-tailed

student's t-test in Graphpad prism. Following the ipGTT, mice were given access to food *ad libitum*.

All animal work was approved by the Institutional Animal Care and Use Committee (IACUC) at

Washington University in St. Louis (Protocol #20180062).

Compounds and Dosing

SR9009 was formulated as 100mg/kg at 10mg/ml in 5% DMSO, 15% Cremophore EL (Sigma), 80%

PBS as previously described [16]. Both vehicle (5% DMSO, 15% CremophoreEL (Sigma), 80% PBS)

and SR9009 were filter sterilized (Millipore Steriflip) prior to dosing. Mice were given once daily i.p.

injections within an hour of "lights on" (ZT0-ZT1). Dosing was performed for 30 days by the same

researcher.

Gene Expression Analysis

Total RNA was isolated from liver using the trizol (Invitrogen) method [12]. Samples were analyzed by

QPCR using Fatty Liver and Fibrosis QPCR array plates (Bio-Rad; 384-well format) and Bio-Rad

supplied SYBR reagents (per manufacturer's protocol). Each sample was run in duplicate and analyzed

on the PrimePCR software supplied by Bio-Rad. Multiple reference genes were utilized (including

103 Gapdh, ActinB, and Cyclophillin) for analysis [14]. Results were plotted in GraphPad prism software as

104 Gene Regulation using mean +/- SEM.

Histology and Pathology Analysis

Livers were placed in 4% PFA at 4°C overnight and then were paraffin-embedded and sectioned at 10

μm onto slides at the Saint Louis University Histology Core Facility. H&E and Masson's Trichrome

staining were also performed at the core as a fee-for-service [12,14,17].. Stained sections (both H&E

and Masson's Trichrome) were sent to Reveal Biosciences (San Diego, CA) for quantification of fibrosis,

inflammation, steatosis, hepatocellular ballooning, and presence of Mallory Bodies utilizing AI-based

digital pathology and image processing. Briefly, whole slide images were generated using 3D Histech

Pannoramic SCAN and uploaded into imageDx™ software. Each scanned image was assessed for

quality and accuracy, then machine learning algorithms were applied to perform automated image quality

control assessments and quantitative measurements of disease features across the entire tissue. For

steatosis analysis, lipid regions within the H&E slides were identified and quantitated as a percent of the

total image analysis area. For hepatocyte ballooning, these hepatocytes were identified and quantitated

as the number of ballooning cells within the total image analysis area of the H&E stained sections.

Positive ballooning was identified based on cell diameter and prescence of disrupted cytoskeletal

structure. The number of ballooning cells was counted and expressed as density across the total area.

Inflammatory cells were identified and quantitated on H&E slides as the number of immune cells within

the total image analysis area. The number of immune cells was counted and measured as the immune

cell density, total number, and area of immune cells. For fibrosis, collagen and extracellular matrix fibers

were identified in the Massson's Trichrome-stained slides and quantitated as a percent of the total image

analysis area. H&E sections were also used to identify the presence of Mallory bodies within the total

image analysis areas.

Statistics

All data are expressed as mean +/- SEM (n = 4 or greater). All expression statistical analysis was performed using ANOVA with Tukey's post-hoc analysis in GraphPad prism software. Weekly mouse weights and food intake data was analyzed by two-way ANOVA followed by Sidak's multiple comparisons at the 95% confidence level. For ipGTT, area under the curve (AUC) was generated and the data was analyzed by two-tailed student's t-test in GraphPad prism software. Sample data generated by Reveal Biosciences was entered into Graphpad prism software and analyzed for statistical significance using an unpaired student's t-test (2-tailed). P-values are reported as follows: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, and **** $p \leq 0.0001$.

Results

Given that REV-ERBs have been demonstrated to play a regulatory role in hepatic lipid metabolism [18] as well as inflammation [16,18,19], we sought to examine the effects of pharmacologically activating REV-ERB in a mouse model of NASH and determine whether REV-ERB may be a therapeutically relevant target. We opted to utilize a diet-induced NASH model using *ob/ob* mice as previously described [14,15] as the time period for development of NAFLD with fibrosis (NASH) is relatively short as compared to other models. During the acclimation and NASH development period, mice were fed a diet that contains Primex as a trans-fat source, fructose, and cholesterol *ad libitum* and monitored for weight gain and food intake. These parameters were also monitored daily throughout the dosing period to validate that any weight-loss was not due to loss of appetite. As shown in Figure 1A, both groups (Vehicle and SR9009) gained weight throughout the experimental period, however the SR9009-treated group gained weight at a consistently slower rate. The slower weight gain was not due to lower food intake in the SR9009-treated mice since this group consistently consumed the same amount of food as the vehicle group (Figure 1B). After 30-days of dosing was completed, mice were euthanized, and we performed a variety of tissue and plasma analyses to determine whether SR9009 treatment had a beneficial effect in this model. While we did not see a significant effect in liver weight as a percentage

of total body weight (S1 Fig), we did observe a decrease in the fat mass of the SR9009-treated animals,

while lean mass was unchanged (Fig 1C). We were interested in determining whether SR9009 had any

utility in improving the hepatic health of these mice. In order to make this determination, we performed

clinical chemistry analysis on blood plasma samples to examine liver enzyme levels. ALT was

significantly decreased in the SR9009-treated group (Fig 1D). While not statistically significant, AST

levels in the SR9009 group were also trending lower than the vehicle group consistent with a benefit due

to SR9009 treatment (S1 Fig). These data suggest that that the amount of liver damage due to the diet

may be suppressed by treatment with REV-ERB agonist SR9009. In addition to effects on hepatic

enzymes, we also observed a significant decrease in fasted blood-glucose levels in the SR9009-treated

mice (Fig 1E). The *ob/ob* mouse model is typically hyperglycemic and addition of the high fat/high

fructose diet hyperglycemia can be particularly prominent. During the third week of dosing, we

performed an ipGTT on the mice (Fig 1F) and observed that while both groups of mice were

hyperglycemic, the SR9009-treated group responded better to the bolus of glucose. In fact, the difference

in AUC was statistically significant between the two groups (Fig 1F right panel) as analyzed by a two-

tailed student's t-test ($p = 0.027$), although final plasma insulin levels were not affected by the treatment

(S1 Fig). Thus, our observation that of lowered hyperglycemia was promising and potentially relevant

to human NASH patients who often present with co-morbidities such as obesity and diabetes [20–24].

Figure 1: REV-ERB agonist treatment of ob/ob NASH diet-fed mice.
Mouse weight (A) and food intake (B) were recorded daily and averaged weekly for each group. At the
termination of the experiment, mice were euthanized and blood and tissues were collected for analysis.
Panel C shows body composition of lean and fat mass for each group. ALT (D), fasting blood glucose
(E), ipGTT and area under curve (AUC) (F), circulating triglycerides (G), and circulating cholesterol
levels (H) were also analyzed from blood plasma.

Hyperlipidemia is also a comorbidity in NASH patients, therefore we also analyzed circulating

triglyceride levels (Fig 1G) and total cholesterol levels (Fig 1H) in both mouse groups. We also observed

significantly reduced total protein in the plasma of SR9009 mice as compared to vehicle-treated mice

(S1 Fig). Our overall impression from the clinical chemistry data and mouse observations is that SR9009

treatment may have beneficial effects in a NASH model.

As the SR9009 group maintained a lower body weight throughout the dosing period and had

significantly decreased circulating lipid levels in blood plasma, we investigated whether the SR9009-

treated mice had reduced hepatosteatosis (both macrovesicular and microvesicular) [12,14,17]. Figure

2A and 2B show the total lipid accumulation and percentage of tissue covered in lipid accumulation

based on the H&E stained liver sections for both vehicle- and SR9009-treated animals. There is no

change in either macrovesicular or microvesicular steatosis (S1 Table), suggesting that the beneficial

effects observed in this mouse model may not be attributable to REV-ERB's role in reduced fat mass.

191
Figure 2: Hepatosteatosis in *ob/ob* mice maintained on a NASH diet does not appear affected in
SR9009-treated group.
Digital pathology analysis suggests that SR9009 treatment in NASH mice may not suppress total hepatic
lipid area (A) or total percentage of lipids within the entire section (B).   While steatosis appears
unaffected by SR9009 treatment, the number of ballooning hepatocytes is decreased in the treated mice
although the effect does not reach significance (C). (D) H&E sections of representative livers for vehicle
(top) and SR9009 (bottom). Scale bar for vehicle section indicates 1.56 mm. Scale bar for SR9009
section indicates 3.11 mm. For steatosis quantitation, macro droplet > 65 μm^2; micro droplet ≤ 65 μm^2.
200
To validate the clinical chemistry data suggesting that the SR9009 reduced liver damage and

improved liver health, quantitated hepatocyte ballooning density from H&E stained liver sections. As

shown by Fig 2C, SR9009-treated mice have a lowered density of hepatocyte ballooning as compared to

vehicle-treated mice. While this was not statistically significant (*p = 0.0903*), it suggests that treatment

with SR9009 may have slowed the progression of NASH in these animals. We were intrigued that REV-

ERB agonism in this model did not significantly affect steatosis in the liver with its known role in lipid

metabolism, but appeared to suppress circulating cholesterol and reduce overall body fat mass. Therefore

we assessed the expression levels of several known REV-ERB target genes involved in hyperlipidemia

(*Dhcr24, ApoE,* and *ApoC3*) by QPCR and observed a significant decrease in these genes (S2 Fig)

suggesting that SR9009 activation of REV-ERB is suppressing cholesterol and lipid metabolism but may

not have efficacy in this model in which treatment started well after the development of NASH began.

While hepatic steatosis was not significantly affected by SR9009 treatment in this model, we further investigated clinical chemistry findings that SR9009 improves hepatic health by quantitating fibrosis and immune cell infiltration in liver sections (Fig 3A). We first evaluated the Masson's trichrome stained sections and identified the collagen and extracellular matrix fibers stained with deep blue within the whole slide image. Positive collagen fibers were then visualized using an overlaid green mask and measured as a percentage of the total image analysis area (S3 Fig). The calculated percent area of positive collagen (fibrosis) was significantly reduced in the SR9009-treated mice (Fig 3B) suggesting that the beneficial hepatic effects from REV-ERB agonism in this model may be due to suppressed fibrosis. Further analysis of H&E staining also indicated a reduced number of inflammatory foci and immune cell count, although these did not reach significance in our analysis (Fig 3C-D). Additionally, H&E sections indicated that all liver samples in this study showed the presence of Mallory Bodies, however the overall morphology of the SR9009-treated livers appeared improved as compared to the vehicle group (S1 Table). Overall, these results suggest that SR9009 treatment halted the progression of NASH in this mouse model by reducing hepatic inflammation and suppressing fibrosis.

Figure 3: SR9009 Significantly Reduced Fibrosis and Suppressed Inflammatory Activity.
(A) Masson's Trichrome staining of liver sections (representative samples). Vehicle pictured at the top shows significant collagen formation. (B) Quantitated analysis of the Masson's Trichrome staining for fibrosis as measured by percent of the total area. Immune Cell Density (C) and Immune Cell Count (D) are unchanged, but appear to be less active as fibrosis is significantly reduced in SR9009 samples. Scale bars indicate 1.56 mm.

To continue our investigation into whether pharmacological activation of REV-ERB is beneficial in an NASH model, we analyzed gene expression of inflammatory markers by QPCR from frozen mouse liver tissues. For this analysis, we focused on expression changes related to inflammatory markers that indicate progression of NAFLD towards a NASH pathology, specifically *IL-1α, IL-1β, Ifnγ, and TNFα.* Previous work from our lab [16] and others have shown that REV-ERBs regulate a variety of genes involved in the pathogenesis of metabolic diseases including those associated with inflammation. Specifically, we were interested to see whether we were suppressing the progression of NAFLD towards

NASH with SR9009 by not only alleviating liver damage as assessed in Figure 1D-H, but by also

suppressing hepatic inflammation in these animals, which was suggested in the H&E staining (Figure 2).

Indeed, when compared to the vehicle-treated group, the SR9009 mice display significantly lower levels

of expression of inflammatory cytokines, all of which have been implicated as biomarkers in NASH

progression (Figure 4A). Since we observed a significant reduction in fibrosis in the SR9009-treated

animals, we also assessed pro-fibrotic gene expression by QPCR. Fig 4B confirms that pharmacological

activation of REV-ERB significantly reduces the expression of various pro-fibrotic genes including

Col3A1, Acta2, STAT1, Mmp13, Timp1, and TGFβ [25]. Interestingly, several other pro-inflammatory

genes (*Col1A2* and *Agt*) are not affected by SR9009 treatment suggesting that REV-ERB may play a role

in the regulation of a select array pro-fibrotic genes involved in NASH progression. While these results

indicated a significant effect on pro-inflammatory and pro-fibrotic genes in liver tissues, we sought to

assess peripheral inflammation in these animals as well. We assessed plasma TNFα using levels at the

time of euthanasia and as shown in Figure 4C, SR9009-treated mice display significantly reduced

circulating TNFα levels. This suggests that activation of REV-ERB in a NASH model may dampen the

progression of NAFLD towards NASH and improve hepatic health by suppressing fibrosis and

inflammatory activity.

Figure 4: Expression of inflammatory markers are downregulated by SR9009 treatment in a mouse
model of NASH.
(A) Gene expression of inflammatory cytokines (*IL-1a, IL-1b, IFNg, TNFa,* and *NFkB)* in mouse liver
tissues that are involved in the pathogenesis of NAFLD-NASH. (B) Gene expression was also performed
for pro-fibrotic genes. As shown in the figure, several genes are significantly downregulated by SR9009
pharmacological activation of REV-ERB. (C) An ELISA for mouse TNFα was performed using plasma
samples from each mouse in triplicate and shows that SR9009 mice had significantly reduced circulating
TNFα.

Discussion

Over the last several years, many studies by our lab and others have demonstrated that the REV-ERBs

regulate a variety of genes involved in lipogenesis, metabolism, and inflammation (Fig 5). Several studies

have investigated potential of REV-ERB agonists as potential therapeutics for cardio-metabolic diseases

including atherosclerosis which is a common comorbidity with NAFLD or NASH [16,19].

Understanding the critical role that the REV-ERBs play in these pathways, we hypothesized that

pharmacological activation of REV-ERB, with the tool compound SR9009, in a mouse model of NASH

would provide beneficial metabolic effects in the liver leading to reduced NASH pathology. While the

SR9009 group did not gain weight at the same rate as the vehicle group and we cannot validate whether

this was a direct effect of the compound, although feeding and other behaviors were observed to be the

same for both groups throughout the study. Our data clearly suggests that SR9009 improved the

metabolic profile in the mice (reduced glucose levels and improved glucose tolerance) while also

improving hepatic health by suppressing inflammation that essential for the progression of NAFLD to

NASH. Most importantly, we observed a significant reduction in expression of pro-fibrotic genes in the

liver, which was consistent with reduced hepatic fibrosis.

Figure 5: The Nuclear Receptor REV-ERB Regulates Inflammation, Lipid Metabolism, and
Glucose Metabolism by Recruiting NCoR to Suppress Transcriptional Activation of Target
Genes.
This schematic demonstrates how the REV-ERB nuclear receptors regulate transcription of target genes
that are involved in inflammation, and lipid and glucose metabolism. Upon agonist binding to REV-
ERB, a conformational change occurs and allows for the nuclear receptor co-repressor complex (NCoR)
to bind and inhibit the transcription of target genes. As REV-ERBs are regulators of inflammation and
hepatic metabolism, it was hypothesized that pharmacological activation of REV-ERB in a NASH model
would improve overall hepatic health by suppressing genes involved in lipid metabolism. As indicated
by the results, SR9009 actually had little to no effect on lipid metabolism but improved overall clinical
indications of NASH in this model. Further investigation demonstrated that REV-ERB agonism in a
NASH model suppresses hepatic inflammation and fibrosis and shows therapeutic benefit in dampening
the progression of this disease. Image created in Biorender.
295

Interestingly, we did not observe a significant effect on hepatic steatosis or total lipid

accumulation, suggesting that the beneficial effects of REV-ERB agonism in this model may not be due

to REV-ERB's role in reduction of fat mass. Based on the quantitative analysis performed on the liver

sections, it appears that hepatocyte ballooning was somewhat decreased in the SR9009 group, although

this did not reach significance. This study was performed in a fairly short period of time (30 days) and

it is possible that an extended period of dosing may have a more beneficial effect on hepatic steatosis.

We originally designed our hypothesis based on our previous study [18] that demonstrated a significant

metabolic impact in hepatic lipogenesis and obesity in DIO mice. In this study, SR9009 was dosed twice

per day (at 100 mg/kg) and resulted in significant repression of lipogenic and metabolic genes, as well

as significant reduction in overall body weight. It is possible that in the current study, the once per day

dosing (100 mg/kg) that was utilized to reduce animal stress did not reach the maximal potential efficacy

in the NASH model.   We utilized SR9009 as a tool to activate REV-ERB in vivo and currently there are

limited pharmacological tools available, but in terms of a pharmacological agent SR9009 has relatively

poor potency (1 μM range), solubility, and pharmacokinetic properties that lead to a quick clearance time

that may attribute to lowered efficacy.   Novel REV-ERB compounds with improved potency and

pharmacokinetic profiles may provide efficacy in NASH studies to suppress hepatic metabolic activity,

inflammation, and fibrosis in the future.

In summary, our data suggests that REV-ERB is a potential therapeutic target to slow the

progression of NAFLD towards NASH. The REV-ERB agonist SR9009 displayed efficacy in reduction

of hepatic pathology associated with NASH. SR9009 is effective in suppressing clinical markers of liver

damage, circulating lipids, hepatic fibrosis and markers of inflammation. Our data suggests that REV-

ERB agonists may offer novel therapies for NAFLD or NASH in the future.

Acknowledgments
319

320 We thank Barb Nagel at Saint Louis University School of Medicine (St. Louis, MO) Histology Core for

321 paraffin embedding and performing histological staining of liver sections, Melissa Kazantzis at The

322 Scripps Research Institute in Jupiter, FL for performing clinical chemistry analysis on samples and

323 Sherry Burris for sectioning of liver samples.

References

1. Fazel Y, B. Koenig A, Sayiner M, D. Goodman Z, M. Younossi Z. Epidemiology and natural history of non-alcoholic fatty liver disease [Internet]. Metabolism. 2016. pp. 1017 – 1025. doi:https://doi.org/10.1016/j.metabol.2016.01.012

2. Polyzos SA, Kang ES, Boutari C, Rhee E-J, Mantzoros CS. Current and emerging pharmacological options for the treatment of nonalcoholic steatohepatitis. Metabolism. Elsevier BV; 2020; 154203. doi:10.1016/j.metabol.2020.154203

3. Konerman MA, Jones JC, Harrison SA. Pharmacotherapy for NASH: Current and emerging. Journal of Hepatology. Elsevier BV; 2018;68: 362–375. doi:10.1016/j.jhep.2017.10.015

4. Yang X, Gonzalez FJ, Huang M, Bi H. Nuclear receptors and non-alcoholic fatty liver disease: An update. Liver Research. Elsevier BV; 2020; doi:10.1016/j.livres.2020.03.001

5. Tanaka N, Aoyama T, Kimura S, Gonzalez FJ. Targeting nuclear receptors for the treatment of fatty liver disease. Pharmacology & Therapeutics. Elsevier BV; 2017;179: 142–157. doi:10.1016/j.pharmthera.2017.05.011

6. López-Velázquez JA, Carrillo-Córdova LD, Chávez-Tapia NC, Uribe M, Méndez-Sánchez N. Nuclear Receptors in Nonalcoholic Fatty Liver Disease. Journal of Lipids. Hindawi Limited; 2012;2012: 1–10. doi:10.1155/2012/139875

7. Preitner N, Damiola F, Luis-Lopez-Molina, Zakany J, Duboule D, Albrecht U, et al. The Orphan Nuclear Receptor REV-ERBα Controls Circadian Transcription within the Positive Limb of the Mammalian Circadian Oscillator. Cell. Elsevier BV; 2002;110: 251–260. doi:10.1016/s0092-8674(02)00825-5

8. Li T, Eheim AL, Klein S, Uschner FE, Smith AC, Brandon-Warner E, et al. Novel role of nuclear receptor rev-erbα in hepatic stellate cell activation: Potential therapeutic target for liver injury. Hepatology. Wiley; 2014;59: 2383–2396. doi:10.1002/hep.27049

9. Stujanna EN, Murakoshi N, Tajiri K, Xu D, Kimura T, Qin R, et al. Rev-erb agonist improves adverse cardiac remodeling and survival in myocardial infarction through an anti-inflammatory mechanism. Fan G-C, editor. PLOS ONE. Public Library of Science (PLoS); 2017;12: e0189330. doi:10.1371/journal.pone.0189330

10. Kojetin D.J. and Burris T.P. A Role for Rev-erbα Ligands in the Regulation of Adipogenesis. Current Pharmaceutical Design. Bentham Science Publishers Ltd.; 2011;17: 320–324. doi:10.2174/138161211795164211

11. Pariollaud M, Gibbs JE, Hopwood TW, Brown S, Begley N, Vonslow R, et al. Circadian clock component REV-ERBα controls homeostatic regulation of pulmonary inflammation. Journal of Clinical Investigation. American Society for Clinical Investigation; 2018;128: 2281–2296. doi:10.1172/jci93910

12. Griffett K, Solt LA, El-Gendy BE-DM, Kamenecka TM, Burris TP. A Liver-Selective LXR Inverse Agonist That Suppresses Hepatic Steatosis. ACS Chemical Biology. American Chemical Society (ACS); 2012;8: 559–567. doi:10.1021/cb300541g

13. Griffett K, Burris TP. Promiscuous activity of the LXR antagonist GSK2033 in a mouse model of fatty liver disease. Biochemical and Biophysical Research Communications. Elsevier BV; 2016;479: 424–428. doi:10.1016/j.bbrc.2016.09.036

14. Griffett K, Welch RD, Flaveny CA, Kolar GR, Neuschwander-Tetri BA, Burris TP. The LXR inverse agonist SR9238 suppresses fibrosis in a model of non-alcoholic steatohepatitis. Molecular Metabolism. Elsevier BV; 2015;4: 353–357. doi:10.1016/j.molmet.2015.01.009

15. Trevaskis JL, Griffin PS, Wittmer C, Neuschwander-Tetri BA, Brunt EM, Dolman CS, et al. Glucagon-like peptide-1 receptor agonism improves metabolic, biochemical, and histopathological indices of nonalcoholic steatohepatitis in mice [Internet]. American Journal of Physiology-Gastrointestinal and Liver Physiology. 2012. pp. G762–G772. doi:10.1152/ajpgi.00476.2011

16. Sitaula S, Billon C, Kamenecka TM, Solt LA, Burris TP. Suppression of atherosclerosis by synthetic REV-ERB agonist. Biochemical and Biophysical Research Communications. Elsevier BV; 2015;460: 566–571. doi:10.1016/j.bbrc.2015.03.070

17. Sengupta M, Griffett K, Flaveny CA, Burris TP. Inhibition of Hepatotoxicity by a LXR Inverse Agonist in a Model of Alcoholic Liver Disease. ACS Pharmacology & Translational Science. American Chemical Society (ACS); 2018;1: 50–60. doi:10.1021/acsptsci.8b00003

18. Solt LA, Wang Y, Banerjee S, Hughes T, Kojetin DJ, Lundasen T, et al. Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. Nature. Springer Science and Business Media LLC; 2012;485: 62–68. doi:10.1038/nature11030

19. Pourcet B, Zecchin M, Ferri L, Beauchamp J, Sitaula S, Billon C, et al. Nuclear Receptor Subfamily 1 Group D Member 1 Regulates Circadian Activity of NLRP3 Inflammasome to Reduce the Severity of Fulminant Hepatitis in Mice. Gastroenterology. Elsevier BV; 2018;154: 1449–1464.e20. doi:10.1053/j.gastro.2017.12.019

20. Petit J-M, Vergès B. GLP-1 receptor agonists in NAFLD. Diabetes & Metabolism. Elsevier BV; 2017;43: 2S28–2S33. doi:10.1016/s1262-3636(17)30070-8

21. Scorletti E, Byrne CD. Extrahepatic Diseases and NAFLD: The Triangular Relationship between NAFLD, Type 2-Diabetes and Dysbiosis. Digestive Diseases. S. Karger AG; 2016;34: 11–18. doi:10.1159/000447276

22. Obesity, Type 2 diabetes and NAFLD. Diabetic Medicine. Wiley; 2014;31: 1–3. doi:10.1111/dme.12377_1

23. Portillo-Sanchez P, Cusi K. Treatment of Nonalcoholic Fatty Liver Disease (NAFLD) in patients with Type 2 Diabetes Mellitus. Clinical Diabetes and Endocrinology. Springer Science and Business Media LLC; 2016;2. doi:10.1186/s40842-016-0027-7

420 24. Bonora E. Novel predictors of diabetes - NAFLD, diabetes and CVD. Endocrine Abstracts.
421   Bioscientifica; 2017; doi:10.1530/endoabs.49.s29.3

423 25. Feng G, Li X-P, Niu C-Y, Liu M-L, Yan Q, Fan L-P, et al. Bioinformatics analysis reveals novel
424   core genes associated with nonalcoholic fatty liver disease and nonalcoholic steatohepatitis
425   [Internet]. Gene. 2020. p. 144549. doi:https://doi.org/10.1016/j.gene.2020.144549

Supporting Information

S1 Fig. Additional clinical chemistry analysis of plasma from mouse NASH model. Percentage of Liver weight to total body weight, AST levels, and plasma insulin levels were not significantly changed by SR9009 treatment although total protein was significantly reduced in these mice.

S2 Fig. REV-ERB target genes involved in cholesterol and lipid metabolism are significantly downregulated by SR9009 treatment in a mouse model of NASH. We analyzed several REV-ERB target genes (*Dhcr24, ApoeE,* and *ApoC3*) by QPCR to determine expression level differences in the groups. As indicated by the graphs, all three target genes were significantly downregulated, suggesting that SR9009 treatment was suppressing these metabolic pathways but had poor efficacy for reducing steatosis.

S3 Fig. Whole liver section images showing Masson's Trichrome staining for fibrosis. Scale bars indicate 1.56 mm.

S1 Table. Summary of Histo-Pathological Analysis on Liver Sections.

A Total Lipid Area

B Lipid Area

C Ballooning Hepatocyte Density

D

Vehicle

SR9009