The eyes reflect an internal cognitive state embedded in the population activity of cortical neurons

Richard Johnston[1, 2, 3], Adam C. Snyder[4, 5, 6], Sanjeev B. Khanna[3, 7], Deepa Issar[1] and Matthew A. Smith[1, 2, 3]

[1]Department of Biomedical Engineering, Carnegie Mellon University, Pittsburgh, USA; [2]Neuroscience Institute, Carnegie Mellon University, Pittsburgh, USA; [3]Center for the Neural Basis of Cognition, Carnegie Mellon University, Pittsburgh, USA; [4]Department of Brain and Cognitive Sciences, University of Rochester, Rochester, USA; [5]Department of Neuroscience, University of Rochester, Rochester, USA; [6]Center for Visual Science, University of Rochester, Rochester, USA; [7] Department of Bioengineering, University of Pittsburgh, Pittsburgh, USA.

Corresponding author:
Matthew A. Smith, PhD
Carnegie Mellon University
Department of Biomedical Engineering and Neuroscience Institute
Mellon Institute, Room 115
Pittsburgh, PA 15213
Email: mattsmith@cmu.edu

Summary

Decades of research have shown that global brain states such as arousal can be indexed by measuring the properties of the eyes. Neural signals from individual neurons, populations of neurons, and field potentials measured throughout much of the brain have been associated with the size of the pupil, small fixational eye movements, and vigor in saccadic eye movements. However, precisely because the eyes have been associated with modulation of neural activity across the brain, and many different kinds of measurements of the eyes have been made across studies, it has been difficult to clearly isolate how internal states affect the behavior of the eyes, and vice versa. Recent work in our laboratory identified a latent dimension of neural activity in macaque visual cortex on the timescale of minutes to tens of minutes. This 'slow drift' was associated with perceptual performance on an orientation-change detection task, as well as neural activity in visual and prefrontal cortex (PFC), suggesting it might reflect a shift in a global brain state. This motivated us to ask if the neural signature of this internal state is correlated with the action of the eyes in different behavioral tasks. We recorded from visual cortex (V4) while monkeys performed a change detection task, and the prefrontal cortex, while they performed a memory-guided saccade task. On both tasks, slow drift was associated with a pattern that is indicative of changes in arousal level over time. When pupil size was large, and the subjects were in a heighted state of arousal, microsaccade rate and reaction time decreased while saccade velocity increased. These results show that the action of the eyes is associated with a dominant mode of neural activity that is pervasive and task-independent, and can be accessed in the population activity of neurons across the cortex.

     Introduction

In the fields of psychology and neuroscience, the eyes are often viewed as a window to the

brain. Much has been learned about cognitive processes, and their development, from studying the

action of the eyes [1–7]. In addition, a large body of research has shown that properties related to

the eyes can be used to index global brain states such as arousal, motivation and cognitive effort

[8–13]. The action of the eyes can be considered broadly in two contexts – the action of the pupil

when the eyes are relatively stable, and the action of the eyes when they move, be it voluntarily,

in response to novel objects in the visual field, or involuntarily during periods of steady fixation.

In each context, technological advancements in infrared eye-tracking have allowed rich insight

about a subject's global brain state to be surmised in a rapid, accurate and non-invasive manner

[14,15].

When the eyes are relatively stable, the size of the pupil changes in response to the amount of

light hitting the retina [16]. However, the pupil does not merely reflect accommodation. Instead,

the size of the pupil is controlled by a balance between parasympathetic and sympathetic pathways

that reflect both light-driven accommodation and central modulation [12]. Several studies have

shown that pupil size is associated with arousal [9–13]. In the mammalian brain, arousal has been

largely associated with the activity of the locus coeruleus (LC) [17–22]. This small structure in the

pons contains a dense population of noradrenergic neurons and is the primary source of

norepinephrine (NE) to the central nervous system [23–25]. Recent neurophysiological work

carried out in rodents and non-human primates has shown that pupil size is significantly associated

with the spiking responses of LC neurons [23,26–28]. Under conditions of heightened arousal,

increases in LC activity and NE concentration are accompanied by increases in pupil size [17–22].

Voluntary saccades occur ~3 times per second to bring novel objects onto the high-resolution

fovea [29]. The characteristics of these saccades, such as the reaction time to initiate the saccade,

and the velocity reached during the saccade, have similarly been used to index global changes in

arousal [8]. For example, when arousal is increased by delivering a startling auditory stimulus prior

to the execution of a saccade, reaction time decreases and saccade velocity increases [31–34].

Another metric that has been linked to global brain states, although to a lesser extent than reaction

time and saccade velocity, is microsaccade rate. These small involuntary saccades are generated

at a rate of 1-2Hz through the activity of neurons in the superior colliculus (SC) [34–37]. Evidence

suggests that microsaccade rate decreases with increased cognitive effort on a range of behavioral

tasks [38–41]. Taken together, these results suggest that the action of the eyes, be it when they are

relatively stable and when they move, can be used to index global brain states.

A number of studies have related the activity of single neurons in many regions of the brain to

pupil size [26,42], microsaccade rate [43–49], reaction time [50–55] and saccade velocity [56,57].

However, if changes in these eye metrics are driven by a shift in an underlying internal state, then

one might expect them all to be related to a common underlying neural activity pattern. Recent

work in our laboratory used dimensionality reduction to identify a dominant mode of neural

activity called slow drift that was related to behavior in a change detection task and present in both

visual and prefrontal cortex in the macaque [58]. If such a prevalent change in brain-wide neural

activity was truly reflective of a changing internal state, then it might also be reflective of wide-

ranging changes in behavior. This motivated us to ask if our measure of the brain's internal state

("slow drift") is related to the activity of the eyes in different behavioral tasks. We recorded the

spiking responses of populations of neurons in V4 while monkeys performed a change detection

task and PFC while the same subjects performed a memory-guided saccade task. On both tasks,

slow drift was associated with a pattern that is indicative of changes in the subjects' arousal level

over time. When pupil size increased, microsaccade rate and reaction time decreased while saccade

velocity increased. These results show that the collective action of the eyes is associated with a

dimension of neural activity that is pervasive and task-independent. They support the view that

slow drift indexes a global brain state that manifests in the movement of the eyes and the size of

the pupil.

<u>Results</u>

To determine if observation of the eyes could provide insight into the internal state

associated with slow drift, we recorded the spiking responses of populations of neurons in two

macaque monkeys using 100-channel "Utah" arrays. We recorded from neurons in 1) V4 while

the subjects performed an orientation-change detection task (Figure 1A); and 2) PFC while they

performed a memory-guided saccade task (Figure 1B). Data from each task has been published

previously [58,60]. However, the main goal of this study was to determine whether the neural

population activity was related to eye metrics in a predictable manner across tasks. Four eye

metrics were recorded during each session: pupil size, microsaccade rate, reaction time and

saccade velocity. These metrics were chosen because they have been used extensively to index

global changes in brain state [8–11], and can be measured easily and accurately with an infrared

eye tracker. We made each of these four measurements on every trial of both behavioral tasks

(Figure 1).

Figure 1. Experimental methods. (A) Change detection task. After an initial fixation period, a sequence of stimuli (orientated Gabors separated by fixation periods) was presented. The subjects' task was to detect an orientation change in one of the stimuli and make a saccade to the changed stimulus. We recorded neural activity from V4 while the animals performed the change detection task using 100-channel "Utah" arrays (inset). (B) Memory-guided saccade task. After an initial fixation period, a target stimulus was presented at 1 of 40 locations followed by a delay period. The central point was then extinguished prompting the subjects to make a saccade to the remembered target location. Neural activity was recorded in PFC while the animals performed the memory-guided saccade task using Utah arrays (inset). (C) Measuring eye metrics on the change detection task. Mean pupil size was recorded during stimulus periods, whereas microsaccade rate was measured during periods of steady fixation (except for the initial fixation period, see *Methods*). Microsaccades (emboldened in two-dimensional eye position and eye velocity space) were defined as eye movements that exceeded a threshold (dashed circle) for 6ms [103]. Reaction time was the time taken to make a saccade to the changed stimulus. Saccade velocity was the peak velocity of the saccade. (D) Measuring eye metrics on the memory-guided saccade task. Mean pupil size was recorded during the presentation of the target stimulus, whereas microsaccade rate was measured during the delay period. Reaction time was the time taken to make a saccade to the remembered target location. Saccade velocity was the peak of velocity of the saccade. RT = reaction time.

<u>Correlations between the eye metrics over time</u>

First, we investigated if the different measures of the eyes were themselves correlated over

time during performance of the behavioral task. A large body of work has shown that arousal is

associated with changes in the action of the eyes, be it when they are relatively stable or when they

move. For example, increases in arousal are typically accompanied by increases in pupil size and

saccade velocity, and concomitant decreases in microsaccade rate and reaction time [8–13,30–

33,38–41]. Given that arousal is a domain-general phenomenon, one might expect a similar pattern

to emerge on different behavioral tasks. To explore whether or not this was the case, we binned

our eye data using a 30-minute sliding window stepped every 6 minutes (Figure 2A and Figure

2B). The width of the window, and the step size, were chosen to isolate slow changes over time

based on previous research. They were the same as those used by Cowley et al. [58], which meant

direct comparisons could be made across studies. An example session from the same subject on

the change detection task and the memory-guided saccade task is shown in Figure 2C and Figure

2D, respectively. In the example sessions shown across both tasks a characteristic pattern was

observed that was indicative of slow changes in the subject's arousal level over time. Specifically,

a large tonic pupil size (measured during stimulus periods on the change detection task and target

presentations on the memory-guided saccade task) was associated with high saccade velocity,

shorter reaction times, and a lower rate of microsaccades.

Figure 2. Isolating slow fluctuations in the eye metrics. (A) Change detection task. Each data point corresponds to a measurement from a single trial. To isolate slow changes, the data was binned using a 30-minute sliding window stepped every 6 minutes (solid line) [58]. (B) Same as (A) but for the memory-guided saccade task. Note in (B) and (C) that data points with a SD ~3 times greater than the mean are not shown for illustration purposes. (C) Example session from Monkey 1 on the change detection task. Each metric has been z-scored for illustration purposes. (D) Example session from the same subject on the memory-guided saccade task.

Next, we explored if a similar pattern was found across all sessions and both animals. We

computed correlations (Pearson product-moment correlation coefficient) between all combinations

of the four eye metrics for each session, and compared that distribution of Pearson's r values to

shuffled distributions using permutation tests (two-sided, difference of medians). Consistent with

the pattern of results observed in several individual sessions (Figure S5A and Figure S6A), we

found significant interactions among the four eye metrics. Pupil size was significantly and

negatively correlated with microsaccade rate (median r = -0.46; p < 0.001) and reaction time

(median r = -0.18; p = 0.027) on the change detection task (Figure 3A and Figure S5B) and

memory-guided saccade task (Figure 3B and Figure S6B, median r = -0.56; p < 0.001 for

microsaccade rate and median r = -0.15; p = 0.011 for reaction time). We did not observe

significant across-session trends in the correlation between pupil size and saccade velocity (change

detection task: r = 0.03, p = 0.651; memory-guided saccade task: r = 0.09, p = 0.523 in Figure 3A-

B), although saccade velocity was itself correlated with reaction time in both tasks (change

detection task: r = -0.52, p < 0.001; memory-guided saccade task: r = -0.61, p < 0.001). Taken

together, these results demonstrate that changes in the subjects' arousal level were accompanied

by changes in the action of the eyes. This was true of pupil size, microsaccade rate, reaction time,

and to a lesser extent, saccade velocity, and suggests that these eye metrics may be used to index

global changes in brain state. This motivated us to ask next whether there were neural correlates

of these behavioral signatures.

Figure 3. Correlations between the eye metrics over time. (A) Change detection task. Correlation matrix showing median r values across sessions between the four eye metrics. (B) Same as (A) but for the memory-guided saccade task. In (A) and (B) actual distributions of r values were compared to shuffled distributions using two-sided permutation tests (difference of medians). p < 0.05*, p < 0.01**, p < 0.001***.

Correlation between the eye metrics and slow drift over time

Previously we reported a slow fluctuation in neural activity in V4 and PFC that we termed 'slow drift' [58]. We found that this neural signature was related to the subject's tendency to make impulsive decisions in a change detection task, ignoring sensory evidence (false alarms). Here, we wanted to investigate if the constellation of eye metrics we observed were associated with the neural signature of internal state that we termed 'slow drift.' To calculate slow drift, we binned spike counts in V4 (change detection task) and PFC (memory-guided saccade task) using the same 30-minute sliding window that had been used to bin the eye metric data (Figure 4A-B, see *Methods*). We then applied principal component analysis (PCA) to the data and estimated slow drift by projecting the binned residual spike counts along the first principal component (i.e., the

loading vector that explained the most variance in the data). Because the sign of the loadings in

PCA is arbitrary [61], the correlation between slow drift and a given eye metric in any session was

equally likely to be positive or negative. This was problematic since we were interested in whether

slow drift was associated with a characteristic pattern that is indicative of changes in the subjects'

arousal level over time i.e., increased pupil size and saccade velocity, and decreased microsaccade

rate and reaction time. In order to establish consistency across sessions in the sign of the

correlations, we constrained the slow drift in each session to have the same relationship to the

spontaneous (change detection task = fixation periods; memory-guided saccade task = delay

periods) and evoked (change detection task = stimulus periods; memory-guided saccade task =

target presentations) activity of the neurons (Figure S4, and see *Methods*). This served to align

slow drift across sessions such that an increase in the slow drift value was associated with neural

activity closer to the evoked pattern of response (i.e., typically higher firing rates).

We computed the slow drift of the neuronal population in each session using this method,

and then compared it to the four eye measures. An example session is shown in Figure 4C-D for

the same subject on the change detection task and the memory-guided saccade task, respectively

(same sessions as in Figure 2). On both tasks, we found a characteristic pattern in which slow drift

was positively associated with pupil diameter and saccade velocity, and negatively associated with

microsaccade rate and reaction time.

Figure 4. Calculating slow drift. (A) Change detection task. Three example neurons from a single session (Monkey 1). Each point represents the mean residual spike count during a 400ms stimulus period. The data was then binned using a 30-minute sliding window stepped every six minutes (solid line) so that direct comparisons could be made with the eye metrics. PCA was used to reduce the dimensionality of the data and slow drift was calculated by projecting binned residual spike counts along the first principal component. (B) Same as (A) but for the memory-guided saccade task (same subject). (C) Example session from Monkey 1 on the change detection task. Each metric has been z-scored for illustration purposes. (D) Example session from the same subject on the memory-guided saccade task.

Next, we explored if a similar pattern was found across sessions. We computed correlations

(Pearson product-moment correlation coefficient) between slow drift and the eye metrics. Actual

distributions of r values were then compared to shuffled distributions using permutation tests (two-

sided, difference of medians). Because the slow drift was aligned across sessions based on the

neural activity alone and not the behavior, the shuffled distributions are centered on a correlation

value of zero. Consistent with the pattern of results observed in several individual sessions (Figure

S7), we found that slow drift in V4 on the change detection task (Figure 5A) was positively

correlated with pupil diameter (median $r = 0.46$, $p < 0.001$) and saccade velocity (median $r = 0.26$,

$p < 0.001$), and negatively correlated with microsaccade rate (median $r = -0.28$, $p = 0.007$) and

reaction time (median $r = -0.19$, $p = 0.016$). An identical pattern of results was found on the

memory-guided saccade task (Figure 5B), where slow drift in PFC was positively correlated with

pupil diameter (median $r = 0.20$; $p = 0.029$) and saccade velocity (median $r = 0.17$, $p = 0.042$), and

negatively correlated with microsaccade rate (median $r = -0.42$; $p < 0.001$) and reaction time

(median $r = -0.27$; $p = 0.008$). These results show that pupil size, microsaccade rate, reaction time

and saccade velocity are associated with slow drift across tasks. Thus, the slow drift was associated

with a pattern that is indicative of changes in the subject's arousal levels over time.

Figure 5. Correlations between the eye metrics and slow drift over time. (A) Change detection task. Histograms showing actual and shuffled distributions of r values. (B) Same as (A) but for the memory-guided saccade task. In (A) and (B) median r values across sessions are indicated by dashed lines (colored lines = actual data; gray lines = shuffled data). Actual distributions of r values were compared to shuffled distributions using two-sided permutation tests (difference of medians). p < 0.05*, p < 0.01**, p < 0.001***.

Discussion

In this study, we investigated if the size of the pupil and the movement of the eyes could be taken as an external signature of an internal brain state, a low-dimensional neural activity pattern called slow drift [58]. There is strong evidence that internal brain states such as slow drift can be measured in the population spiking activity of neurons, and that measurements of the eyes can provide important context into the behavior of subjects on perceptual and decision-making tasks. Hence, we were keen to determine whether we could directly link a neural measure of internal brain state acquired from the spiking activity of a population of neurons with external features of

behavior. On two types of perceptual tasks, we found that slow drift was significantly correlated

with a pattern of eye metrics that was indicative of changes in the subjects' arousal level over time.

Our results show that the action of the eyes, be it when they are relatively stable or when they

move, is associated with a latent dimension of neural activity that is pervasive and task-

independent.

As described above, decades of research have shown that eye metrics are related to task

performance in a variety of contexts [8–13]. Heightened levels of arousal have been associated

with a large pupil. In a variety of work, the size of the pupil has been associated positively with

saccade velocity and negatively with microsaccade rate and reaction time [8–13,30–33,38–41].

Given that arousal is a global phenomenon, one might expect a common pattern across multiple

behavioral tasks. In the present study, we addressed this issue by investigating the relationships

between eye metrics on tasks designed to probe the mechanisms underlying perceptual decision

making (change detection task) and working memory (memory-guided saccade task). Results

showed that pupil size was negatively correlated with microsaccade rate and reaction time on both

tasks. Although no significant correlation was found between pupil size and saccade velocity, the

overall pattern of results suggests that each subject's arousal level was changing over time in a

task-independent manner. These findings support the view that measuring properties related to the

eyes can provide a non-invasive index of global brain states [8–13]. This motivated us to ask if

they are also associated with slow drift.

Most studies that have explored the relationship between eye metrics and neural activity

have used single-neuron (spike count, Fano factor) and pairwise statistics (r_{sc}). However, numerous

recent studies have shown that rich insight about cognitive processes (e.g., learning, decision

making, working memory, time perception) can be gained from analysis of the simultaneous

activity of populations of neurons [51,62–68]. In addition, recent work has shown that a low-

dimensional representation of neural activity in the mouse can be used to index global changes in

brain state. Stringer et al. [69] applied PCA to data recorded from more than 10,000 neurons and

found that fluctuations in the first principal component were significantly associated with

whisking, pupil size, and running speed. In this study, we investigated if slow drift, a dominant

mode of neural activity in macaque cortex is associated with the action of the eyes. On both tasks,

we found that it was correlated with a pattern of eye metrics that is indicative of changes in the

subject's arousal level over time. Our results, coupled with those of Stringer et al. [69], suggest

that much can be learned about global brain states, as well as cognitive processes, when high-

dimensional population activity is reduced to a low-dimensional subspace [70–73]. A key question

for future research is whether latent dimensions of neural activity in the cortex are associated with

activity in subcortical brain regions? One might expect this to be the case given that slow drift was

significantly correlated with pupil size on both tasks.

Evidence suggests that pupil size is associated with activity in the LC, a subcortical

structure that regulates arousal by releasing NE in a diverse manner throughout much of the brain

[74–77]. That slow drift can be used to index global brain states suggests that it might be associated

with LC activity. Although the LC is limited in size (~3mm caudorostrally) and buried deep within

the brainstem [78,79], several studies have successfully recorded from single neurons in LC of the

macaque [26,27,80–84]. In addition, LC can be activated using optogenetics [85–88], electrical

microstimulation [14,16, 58], and pharmacological manipulations [17,89,90]. Thus, it should be

possible to alter the course of slow drift in the cortex using some, if not all, of these methods. As

well as having a significant effect on pupil size, our results predict that activating the LC, directly

or indirectly, may lead to task-independent changes in microsaccade rate, reaction time and

saccade velocity.

Evidence suggests that microsaccades, reaction time and saccade velocity are associated

with neural activity in the SC [37]. This layered structure, located at the roof of the brain stem,

plays a critical role in transforming sensory information into eye movement commands. Population

recordings have been successfully performed in the SC using linear probes [91], and it thus might

be possible to identify dominant patterns of neural activity in SC using dimensionality reduction

[71]. Our results predict that slow drift in the cortex should be significantly correlated with slow

drift in the SC. However, this might depend upon the mixture of SC neurons in the recorded

population. We previously suggested that slow drift must be removed at some stage before motor

commands are issued to prevent unwanted eye movements [58]. Thus, one might not expect a

correlation between slow drift in the cortex and slow drift in deep-layer SC neurons that fire

vigorously prior to the execution of a saccade, and relay motor commands to downstream nuclei

innervating the oculomotor muscles [92]. An alternative possibility is that slow drift is not

removed, but instead occupies an orthogonal subspace in the SC that is not read out by downstream

nuclei. This is not beyond the realm of possibility given that an identical scheme appears to exist

in the skeletomotor system to stop preparatory signals reaching the muscles [93–96]. Further

research is needed to disentangle these possibilities.

Studies in the fields of psychology and neuroscience have mainly used pupil size to index

arousal, but metrics such as heart rate (HR) and galvanic skin response (GSR) are also associated

with global brain states. For example, Wang et al. [97] measured pupil size, HR and GSR while

human subjects viewed emotional face stimuli specifically designed to evoke fluctuations in

arousal. Results showed that all three metrics were positively correlated prior to the presentation

of the stimuli. That is, when pupil size was large, and the subjects were in heightened state of

arousal, HR and GSR increased. In addition, it has been suggested that pre-stimulus oscillations in

the alpha band can be used to index global brain states as they are inversely related to performance

on visual detection tasks. Several studies using electroencephalography (EEG) have shown that

the probability of detecting near-threshold stimuli increases when pre-stimulus power in the alpha

band is low [98–101]. Simultaneous recordings of spiking activity and EEG in awake behaving

monkeys are rare. However, our results predict that slow drift should be associated with pre-

stimulus alpha oscillations.

Research has also uncovered a link between alpha oscillations and microsaccades [102].

This effect has been attributed to changes in spatial attention, which have a profound effect on

microsaccade direction [106–110]. For example, Lowet et al. [49] found that attention-related

modulation of spiking responses, Fano factor and r_{sc} only occur following a microsaccade in the

direction of an attended stimulus [49]. In the present study, we found that slow drift was

significantly correlated with microsaccade rate on both tasks. This is unlikely to be explained by

changes in spatial attention as both Cowley et al. [58] and Rabinowitz et al. [109] found that slow

drift on a change detection task was not associated with the blocks of trials used to cue spatial

attention inside and outside the RF. In addition, the correlation between slow drift and

microsaccade rate in the present study was lower on the change detection task (r = -0.27) than the

memory-guided saccade task (r = -0.37). One would not have expected this to be the case if slow

drift was mediated by changes in spatial attention. These results raise the possibility that spatial

attention and arousal have differential effects on microsaccades. Attention might specifically affect

microsaccade direction, whereas arousal might affect microsaccade rate irrespective of direction.

Further research is needed to test this hypothesis.

In summary, we investigated if properties related to the eyes are associated with slow drift:

a low-dimensional pattern of neural activity that was recently identified in the macaque cortex by

Cowley et al. [58]. On both tasks, we found that slow drift was significantly associated with a

pattern of eye metrics that is indicative of changes in the subjects' arousal level over time. These

results demonstrate that the collective action of the eyes is associated with a latent dimension of

neural activity that is pervasive and task-independent. They suggest that slow drift can be used to

index global changes in brain state over time. Further research is necessary to determine the origins

of this slow drift in population activity. A key question for future work will be to determine the

mechanisms by which slow drift influences behavior in some instances (e.g., when arousal level

drives an urgent response) and is circumvented in others (e.g., when an accurate perceptual

judgement must be made regardless of arousal level).

<u>Methods</u>

<u>Subjects</u>

Two adult rhesus macaque monkeys (*Macaca mulatta*) were used in this study. Surgical

procedures to chronically implant a titanium head post (to immobilize the subjects' head during

experiments) and microelectrode arrays were conducted in aseptic conditions under isoflurane

anesthesia, as described in detail by Smith and Sommer [108]. Opiate analgesics were used to

minimize pain and discomfort during the perioperative period. Neural activity was recorded using

100-channel "Utah" arrays (Blackrock Microsystems) in V4 (Monkey 1 = right hemisphere;

Monkey 2 = left hemisphere) and PFC (Monkey 1 = right hemisphere; Monkey 2 = left

hemisphere) while the subjects performed the change detection task (Figure 1A). On the memory-

guided saccade task (Figure 1B), neural activity was only recorded in PFC (Monkey 1 = left

hemisphere; Monkey 2 = right hemisphere). The arrays comprised a 10x10 grid of silicon

microelectrodes (1 mm in length) spaced 400 μm apart. Experimental procedures were approved

by the Institutional Animal Care and Use Committee of the University of Pittsburgh and were

performed in accordance with the United States National Research Council's Guide for the Care

and Use of Laboratory Animals.

<u>Microelectrode array recordings</u>

Signals from each microelectrode in the array were amplified and band-pass filtered (0.3–

7500 Hz) by a Grapevine system (Ripple). Waveform segments crossing a threshold (set as a

multiple of the root mean square noise on each channel) were digitized (30KHz) and stored for

offline analysis and sorting. First, waveforms were automatically sorted using a competitive

mixture decomposition method [110]. They were then manually refined using custom time

amplitude   window   discrimination   software   ([111],   code   available   at

https://github.com/smithlabvision/spikesort), which takes into account metrics including (but not

limited to) waveform shape and the distribution of interspike intervals. A mixture of single and

multiunit activity was recorded, but we refer here to all units as "neurons". On the change detection

task, the mean number of V4 neurons across sessions was 70 (SD = 11) for Monkey 1 and 31 (SD

= 16) for Monkey 2, whereas the mean number of PFC neurons across sessions was 84 (SD = 15)

for Monkey 1 and 90 (SD = 20) for Monkey 2. On the memory-guided saccade task, the mean

number of PFC neurons across sessions was 54 (SD = 11) for Monkey 1 and 38 (SD = 19) for

Monkey 2.

<u>Visual stimuli</u>

Visual stimuli were generated using a combination of custom software written in

MATLAB (The MathWorks) and Psychophysics Toolbox extensions (Brainard, 1997; Pelli, 1997;

Kleiner et al., 2007). They were displayed on a CRT monitor (resolution = 1024 X 768 pixels;

refresh rate = 100Hz), which was viewed at a distance of 36cm and gamma-corrected to linearize

the relationship between input voltage and output luminance using a photometer and look-up-

tables.

<u>Behavioral tasks</u>

<u>Orientation-change detection task</u>

Subjects fixated a central point (diameter = 0.6°) on the monitor to initiate a trial (Figure

1A). Each trial comprised a sequence of stimulus periods (400ms) separated by fixation periods

(duration drawn at random from a uniform distribution spanning 300-500ms). The 400ms stimulus

periods comprised pairs of drifting full-contrast Gabor stimuli. One stimulus was presented in the

aggregate receptive field (RF) of the recorded V4 neurons, whereas the other stimulus was

presented in the mirror-symmetric location in the opposite hemifield. Although the spatial

(Monkey 1 = 0.85cycles/°; Monkey 2 = 0.85cycles/°) and temporal frequencies (Monkey 1 =

8cycles/s; Monkey 2 = 7cycles/s) of the stimuli were not optimized for each individual V4 neuron

they did evoke a strong response from the population. The orientation of the stimulus in the

aggregate RF was chosen at random to be 45 or 135°, and the stimulus in the opposite hemifield

was assigned the other orientation. There was a fixed probability (Monkey 1 = 30%; Monkey 2 =

40%) that one of the Gabors would change orientation by ±1, ±3, ±6, or ±15° on each stimulus

presentation. The sequence continued until the subject) made a saccade to the changed stimulus

within 700ms ("hit"); 2) made a saccade to an unchanged stimulus ("false alarm"); or 3) remained

fixating for more than 700ms after a change occurred ("miss"). If the subject correctly detected an

orientation change, they received a liquid reward. In contrast, a time-out occurred if the subject

made a saccade to an unchanged stimulus delaying the beginning of the next trial by 1s. It is

important to note that the effects of spatial attention were also investigated (although not analyzed

in this study) by cueing blocks of trials such that the orientation change was 90% more likely to

occur within the aggregate V4 RF than the opposite hemifield. Data exploring how spatial attention

effects diversity in the neural population code [59] and slow drift [58] have been published

previously.

<u>Memory-guided saccade task</u>

Subjects fixated a central point (diameter = 0.6°) on the monitor to initiate a trial (Figure

1B). After fixating within a circular window (diameter = 2.4° and 1.8° for Monkey 1 and Monkey

2, respectively) for 200ms, a target stimulus (diameter = 0.8°) was presented for 50ms (except for

1 session in which it was presented for 400ms). The target stimulus appeared at 1 of 8 angles

separated by 45°, and 1 of 5 eccentricities, yielding 40 conditions in total. After the target stimulus

had been presented, subjects were required to maintain fixation for a delay period. For Monkey 1,

the duration of the delay period was either 1) drawn at random from a distribution spanning 1200-

3750ms; or 2) fixed at 1400 or 2000ms. For Monkey 2, the duration of the delay period was 500ms.

If steady fixation was maintained throughout the delay period, the central point was extinguished

prompting the subjects to make a saccade to the remembered target location. The subjects had

500ms to initiate a saccade. Once the saccade had been initiated, they had a further 200ms to reach

the remembered target location. To receive a liquid reward, the subjects' gaze had to be maintained

within a circular window centered on the target location (diameter = 4 and 2.7° for Monkey 1 and

Monkey 2, respectively) for 150 ms. In a subset of sessions, the target was briefly reilluminated,

after the fixation point was extinguished and the saccade had been initiated, to aid in saccade

completion.

<u>Eye tracking</u>

Eye position and pupil diameter were recorded monocularly at a rate of 1000Hz using an

infrared eye tracker (EyeLink 1000, SR Research).

Microsaccade detection

Microsaccades were defined as eye movements that exceeded a velocity threshold of 6

times the standard deviation of the median velocity for at least 6ms [103,114,115]. They were

required to be separated in time by at least 100ms. In addition, we removed microsaccades with

an amplitude greater than 1° and a velocity greater than 100°/s. To assess the validity of our

microsaccade detection method, the correlation (Pearson product-moment correlation coefficient)

between the amplitude and peak velocity of detected microsaccades (i.e., the main sequence) was

computed for each session (Figure S1A and Figure S1B). The mean correlation between

microsaccade amplitude and peak microsaccade velocity across sessions was 0.86 (SD = 0.07) for

the change detection task and 0.83 (SD = 0.04) for the memory-guide saccade task (Figure S1C

and Figure S1D). These findings indicate that our microsaccade detection algorithm was robust

[116].

Eye metrics

Change detection task

Mean pupil diameter was measured during stimulus periods, whereas microsaccade rate

was measured during fixation periods (Figure 1C). We did not include the first fixation period

when measuring microsaccade rate in the change-detection task. As can be seen in Figure 1C, there

was an increase in eye position variability during this period resulting from fixation having been

established a short time earlier (300-500ms). Such variability was not present in proceeding

fixation periods. Reaction time and saccade velocity were measured on trials in which the subjects

were rewarded for correctly detected an orientation change. Reaction time was defined as the time

from when the change occurred to the time at which the saccade exceeded a velocity threshold of

100°/s. Saccade velocity was the peak velocity of the saccade to the changed stimulus. To isolate

slow changes in the eye metrics over time the data for each session was binned using a 30-minute

sliding window stepped every 6 minutes (Figure 2A).

<u>Memory-guided saccade task.</u>

Pupil size, microsaccade rate, reaction time and saccade velocity were measured on trials

in which the subjects received a liquid reward for making a correct saccade to the remembered

target location. Mean pupil diameter was measured during the presentation of the target stimulus,

whereas microsaccade rate was measured during the delay period (Figure 1D). Reaction time was

defined as the time from when the fixation point was extinguished to the time at which the saccade

reached a threshold of 100 °/s. Saccade velocity was the peak of velocity of the saccade to the

remembered target location. As in the change-detection task, the data for each session was binned

using a 30-minute sliding window stepped every 6 minutes (Figure 2B).

<u>Calculating slow drift</u>

<u>Orientation-change detection task</u>

The spiking responses of populations of neurons in V4 were measured during a 400ms

period that began 50ms after stimulus presentation (Figure 4A). To control for the fact that some

neurons had a preference for one orientation (45 or 135°) over the other residual spike counts were

calculated. We subtracted the mean response for a given orientation across the entire session from

individual responses to that orientation. To isolate slow changes in neural activity over time,

residual spike counts for each V4 neuron were binned using a 30-minute sliding window stepped

every 6 minutes [58]. PCA was then performed to reduce the high-dimensional residual data to a

smaller number of variables [71]. Slow drift in V4 was estimated by projecting the binned residual

spike counts for each neuron along the first principal component [58]. As described above, the

spiking responses of neurons in PFC were simultaneously recorded on the change detection task.

When PFC slow drift was calculated using the method described above, we found it to be

significantly associated with V4 slow drift (median r = 0.95, p < 0.001), consistent with previous

results [58]. On the change detection task, we investigated the relationships between the eye

metrics and V4 slow drift. However, a very similar pattern of results was found when slow drift

was calculated using simultaneously recorded PFC data (Figure S2). The only difference was the

correlation between slow drift and reaction time, which was not statistically significant (r = -0.09,

p = 0.495).

Memory-guided saccade task

The spiking responses of populations of neurons in PFC were measured during the delay

period (Figure 4B). To control for the fact that some neurons had a preference for one target

location over another, residual spike counts were calculated. We subtracted the mean response to

a given target location across the entire session from individual responses to that location. To

isolate slow changes in neural activity over time residual spike counts for each PFC neuron were

binned using a 30-minute sliding window stepped every 6 minutes. PCA was then performed, and

slow drift was estimated by projecting the binned residual spike counts for each neuron along the

first principal component.

Controlling for neural recording instabilities

To rule out the possibility that slow drift arose due to recording instabilities (e.g., the

distance between the neuron and the microelectrodes changing slowly over time) we only included

neurons with stable waveform shapes throughout a session. This was quantified by calculating

percent waveform variance for each neuron (Figure S3). First, the session was divided into 10 non-

overlapping time bins. A residual waveform was then computed for each time bin by subtracting

the mean waveform across time bins. The variance of each residual waveform was divided by the

variance of the mean waveform across time bins yielding 10 values (one of reach time bin). Percent

waveform variance was defined as the maximum value across time bins. Neurons with a percent

waveform variance greater than 10% were deemed as having unstable waveform shapes

throughout a session. They were excluded from all analyses, consistent with previous research

[58].

<u>Aligning slow drift across sessions</u>

As described above, slow drift was calculated by projecting binned residual spike counts

along the first principal component [58]. The weights in a PCA can be positive or negative [61],

which meant the sign of the correlation between slow drift and a given eye metric was arbitrary.

Preserving the sign of the correlations was particularly important in this study because we were

interested in whether slow drift was associated with a pattern that is indicative of changes in the

subjects' arousal levels over time i.e., increased pupil size and saccade velocity, and decreased

microsaccade rate and reaction time. Thus, we had to devise a method to align slow drift across

sessions. As expected, the mean evoked population response calculated during stimulus periods

on the change detection task was significantly higher than the mean spontaneous population

response calculated during fixation periods (Figure S4A). Similarly, on the memory-guided

saccade task, the mean evoked population response calculated during target presentations was

significantly higher than the mean spontaneous population response calculated during delay

periods (Figure S4B). To align slow drift for each individual session, we projected evoked and

spontaneous responses onto the first principal component (Figure S4C-D). The sign of the slow

drift was flipped if the mean projection value for the evoked responses was less than that for the

spontaneous responses – i.e., if the relationship described above for unprojected data did not hold

true.

<u>Acknowledgements</u>

D.I. was supported by National Institutes of Health (NIH) Grant T32 GM-008208 and the ARCS

Foundation Thomas-Pittsburgh Chapter Award. M.A.S. was supported by NIH Grants R01 EY-

022928, R01 MH-118929, R01 EB-026953, and P30 EY-008098; NSF Grant NCS 1734901; a

career development grant and an unrestricted award from Research to Prevent Blindness; and the

Eye and Ear Foundation of Pittsburgh. A.C.S. was supported by NIH grant K99EY025768. S.B.K.

was supported by NIH Grant T32 EY-017271. The authors would like to thank Ms. Samantha

Schmitt for assistance with surgery and data collection.

Figure S1. Assessing the validity of our microsaccade detection algorithm. The correlation between the amplitude and
peak velocity of detected microsaccades was computed for each session. (A) Change detection task. Scatter plot
showing the relationship between microsaccade amplitude and peak velocity for an example session on the change
detection task. (B) Same as (A) but for the memory-guided saccade task (same subject). (C) Histogram showing the
distribution of r values for the change detection task. (D) Same as (C) but for the memory-guided saccade task. In (C)
and (D) dashed lines indicate the mean r value across sessions. On both tasks, the mean correlation between
microsaccade amplitude and peak velocity was strong, indicating that our method of detecting microsaccades was
robust [116]. $p < 0.05*$, $p < 0.01**$, $p < 0.001***$.

592 Figure S2. Correlations between the eye metrics and PFC slow drift on the change detection task. Histograms showing
593 actual and shuffled distributions of r values. Median r values across sessions are indicated by dashed lines (colored
594 lines = actual data; gray lines = shuffled data). Actual distributions of r values were compared to shuffled distributions
595 using two-sided permutation tests (difference of medians). P < 0.05*, p < 0.01**, p < 0.001***.

606
607 Figure S3. Controlling for neural recording instabilities. (A) Percent waveform variance of four example neurons
608 recorded from Monkey 1 during a single session on the change detection task. (B) Same as (A) but for the memory-
609 guided saccade task (same subject).
610

Figure S4. Aligning slow drift across sessions. (A) Bar charts showing the mean spontaneous and evoked population response across sessions on the change detection task. Spontaneous activity was calculated during fixation periods, whereas evoked activity was calculated during stimulus periods. (B) Same as (A) but for the memory-guided saccade task. Spontaneous activity was recorded during the delay period, whereas evoked activity was recorded during the presentation of the target stimulus. In (A) and (B) the mean evoked population response was significantly higher for evoked than spontaneous activity (two-sided permutation tests, difference of means). (C) For each session, slow drift on the change detection task was aligned by projecting evoked and spontaneous responses onto the first principal component. The sign of the slow drift was flipped if the mean projection value for the evoked responses was less than that for the spontaneous responses i.e., if the relationship shown in (A) for unprojected data did not hold true. (D) Same as (C) but for the memory-guided saccade task. $p < 0.05*$, $p < 0.01**$, $p < 0.001***$. Error bars represent ± 1 SEM.

670 Figure S5. Correlations between the eye metrics on the change detection task. (A) Three example sessions from
671 Monkey 1. (B) Histograms showing actual and shuffled distributions of r values across sessions. Median r values are
672 indicated by dashed lines (blue lines = actual data; gray lines = shuffled data). Actual distributions of r values were
673 compared to shuffled distributions using two-sided permutation tests (difference of medians). $p < 0.05$*, $p < 0.01$**,
674 $p < 0.001$***.
675

Figure S6. Correlations between the eye metrics on the memory-guided saccade task. (A) Three example sessions from Monkey 1. (B) Histograms showing actual and shuffled distributions of r values. Median r values are indicated by dashed lines (blue lines = actual data; gray lines = shuffled data). Actual distributions of r values were compared to shuffled distributions using two-sided permutation tests (difference of medians). $p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$.

A)

Change detection task

B)

Memory-guided saccade task

Time (hours)

703 Figure S7. Correlations between slow drift and the eye metrics. (A) Three example sessions from Monkey 1 on the
704 change detection task. (B) Same as (A) but for the memory-guided saccade task (same subject).
705

711   <u>References</u>

712   1.   Hannula, D.E., Althoff, R.R., Warren, D.E., Riggs, L., Cohen, N.J., and Ryan, J.D. (2010).

713   Worth a glance: Using eye movements to investigate the cognitive neuroscience of

714   memory. Front. Hum. Neurosci. *4*, 1–16.

715   2.   Ryan, J.D., and Shen, K. (2020). The eyes are a window into memory. Curr. Opin. Behav.

716   Sci. *32*, 1–6. Available at: https://doi.org/10.1016/j.cobeha.2019.12.014.

717   3.   König, P., Osnabrück, U., Ossandón, J.P., Ehinger, B. V, Osnabrück, U., Gameiro, R.R.,

718   Osnabrück, U., and Kaspar, K. (2016). Eye movements as a window to cognitive

719   processes. J. Eye Mov. Res. *9*, 1–16.

720   4.   Eckstein, M.K., Guerra-Carrillo, B., Miller Singley, A.T., and Bunge, S.A. (2017).

721   Beyond eye gaze: What else can eyetracking reveal about cognition and cognitive

722   development? Dev. Cogn. Neurosci. *25*, 69–91. Available at:

723   http://dx.doi.org/10.1016/j.dcn.2016.11.001.

724   5.   Aslin, R.N. (2012). Infant eyes: A window on cognitive development. Infancy *17*, 126–

725   140.

726   6.   Hessels, R.S., and Hooge, I.T.C. (2019). Eye tracking in developmental cognitive

727   neuroscience – The good, the bad and the ugly. Dev. Cogn. Neurosci. *40*, 100710.

728   Available at: https://doi.org/10.1016/j.dcn.2019.100710.

729   7.   Rayner, K. (2009). Eye movements and attention in reading, scene perception, and visual

730   search.

731   8.   Di Stasi, L.L., Catena, A., Cañas, J.J., Macknik, S.L., and Martinez-Conde, S. (2013).

732   Saccadic velocity as an arousal index in naturalistic tasks. Neurosci. Biobehav. Rev. *37*,

733   968–975.

9.   Mathôt, S., and Van der Stigchel, S. (2015). New Light on the Mind's Eye: The Pupillary

Light Response as Active Vision. Curr. Dir. Psychol. Sci. *24*, 374–378.

10.   Larsen, R.S., and Waters, J. (2018). Neuromodulatory correlates of pupil dilation. Front.

Neural Circuits *12*, 1–9.

11.   Wang, C.A., and Munoz, D.P. (2015). A circuit for pupil orienting responses: Implications

for cognitive modulation of pupil size. Curr. Opin. Neurobiol. *33*, 134–140. Available at:

http://dx.doi.org/10.1016/j.conb.2015.03.018.

12.   Joshi, S., and Gold, J.I. (2019). Pupil size as a window on neural substrates of cognition.

Trends Cogn. Sci., 1–24. Available at: https://doi.org/10.1016/j.tics.2020.03.005.

13.   Mathôt, S. (2018). Pupillometry: Psychology, Physiology, and Function. *1*, 1–23.

14.   McCamy, M.B., Otero-Millan, J., Leigh, R.J., King, S.A., Schneider, R.M., Macknik,

S.L., and Martinez-Conde, S. (2015). Simultaneous recordings of human microsaccades

and drifts with a contemporary video eye tracker and the search coil technique. PLoS One

10, 1–20.

15.   Kimmel, D.L., Mammo, D., and Newsome, W.T. (2012). Tracking the eye non-invasively:

Simultaneous comparison of the scleral search coil and optical tracking techniques in the

macaque monkey. Front. Behav. Neurosci. *6*, 1–17.

16.   Campbell, F.W., and Gregory, A.H. (1960). Effect of Size of Pupil on Visual Acuity.

Nature *187*, 1121–1123.

17.   Aston-Jones, G., and Cohen, J.D. (2005). AN INTEGRATIVE THEORY OF LOCUS

COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal

Performance. Annu. Rev. Neurosci. *28*, 403–450.

18.   van den Brink, R.L., Pfeffer, T., and Donner, T.H. (2019). Brainstem Modulation of

Large-Scale Intrinsic Cortical Activity Correlations. Front. Hum. Neurosci. *13*, 1–18.

19. Berridge, C.W., and Waterhouse, B.D. (2003). The locus coeruleus-noradrenergic system:

Modulation of behavioral state and state-dependent cognitive processes. Brain Res. Rev.

42, 33–84.

20. Sara, S.J. (2009). The locus coeruleus and noradrenergic modulation of cognition. Nat.

Rev. Neurosci. *10*, 211–223.

21. Samuels, E., and Szabadi, E. (2008). Functional Neuroanatomy of the Noradrenergic

Locus Coeruleus: Its Roles in the Regulation of Arousal and Autonomic Function Part II:

Physiological and Pharmacological Manipulations and Pathological Alterations of Locus

Coeruleus Activity in Humans. Curr. Neuropharmacol. *6*, 254–285.

22. Chandler, D.J. (2016). Evidence for a specialized role of the locus coeruleus noradrenergic

system in cortical circuitries and behavioral operations. Brain Res. *1641*, 197–206.

Available at: http://dx.doi.org/10.1016/j.brainres.2015.11.022.

23. Breton-Provencher, V., and Sur, M. (2019). Active control of arousal by a locus coeruleus

GABAergic circuit. Nat. Neurosci. *22*, 218–228. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/30643295.

24. Schwarz, L.A., and Luo, L. (2015). Organization of the locus coeruleus-norepinephrine

system. Curr. Biol. *25*, R1051–R1056. Available at:

http://dx.doi.org/10.1016/j.cub.2015.09.039.

25. Szabadi, E. (2013). Functional neuroanatomy of the central noradrenergic system. J.

Psychopharmacol. *27*, 659–693.

26. Joshi, S., Li, Y., Kalwani, R.M., and Gold, J.I. (2016). Relationships between Pupil

Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex.

Neuron *89*, 221–234. Available at: http://dx.doi.org/10.1016/j.neuron.2015.11.028.

27. Varazzani, C., San-Galli, A., Gilardeau, S., and Bouret, S. (2015). Noradrenaline and

dopamine neurons in the reward/effort trade-off: A direct electrophysiological comparison

in behaving monkeys. J. Neurosci. *35*, 7866–7877.

28. Reimer, J., McGinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D.A., and

Tolias, A.S. (2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic

activity in cortex. Nat. Commun. *7*, 1–7. Available at:

http://dx.doi.org/10.1038/ncomms13289.

29. Kowler, E. (2011). Eye movements: The past 25years. Vision Res. *51*, 1457–1483.

Available at: http://dx.doi.org/10.1016/j.visres.2010.12.014.

30. DiGirolamo, G.J., Patel, N., and Blaukopf, C.L. (2016). Arousal facilitates involuntary eye

movements. Exp. Brain Res. *234*, 1967–1976.

31. Kristjánsson, Á., Vandenbroucke, M.W.G., and Driver, J. (2004). When pros become cons

for anti- versus prosaccades: Factors with opposite or common effects on different saccade

types. Exp. Brain Res. *155*, 231–244.

32. Castellote, J.M., Kumru, H., Queralt, A., and Valls-Solé, J. (2007). A startle speeds up the

execution of externally guided saccades. Exp. Brain Res. *177*, 129–136.

33. Deuter, C.E., Schilling, T.M., Kuehl, L.K., Blumenthal, T.D., and Schachinger, H. (2013).

Startle effects on saccadic responses to emotional target stimuli. Psychophysiology *50*,

1056–1063.

34. Rolfs, M. (2009). Microsaccades: Small steps on a long way. Vision Res. *49*, 2415–2441.

Available at: http://dx.doi.org/10.1016/j.visres.2009.08.010.

35. Hafed, Z.M., Chen, C.-Y., and Tian, X. (2015). Vision, Perception, and Attention through

the Lens of Microsaccades: Mechanisms and Implications. Front. Syst. Neurosci. *9*, 1–16.

Available at: http://journal.frontiersin.org/Article/10.3389/fnsys.2015.00167/abstract.

36.   Martinez-Conde, S., Otero-Millan, J., and MacKnik, S.L. (2013). The impact of

microsaccades on vision: Towards a unified theory of saccadic function. Nat. Rev.

Neurosci. *14*, 83–96.

37.   Gandhi, N.J., and Katnani, H.A. (2011). Motor Functions of the Superior Colliculus.

Annu. Rev. Neurosci. *34*, 205–231. Available at:

http://www.annualreviews.org/doi/10.1146/annurev-neuro-061010-113728.

38.   Gao, X., Yan, H., and Sun, H.J. (2015). Modulation of microsaccade rate by task difficulty

revealed through between- and within-trial comparisons. J. Vis. *15*, 1–15.

39.   Siegenthaler, E., Costela, F.M., Mccamy, M.B., Di Stasi, L.L., Otero-Millan, J.,

Sonderegger, A., Groner, R., Macknik, S., and Martinez-Conde, S. (2014). Task difficulty

in mental arithmetic affects microsaccadic rates and magnitudes. Eur. J. Neurosci. *39*,

287–294.

40.   Valsecchi, M., Betta, E., and Turatto, M. (2007). Visual oddballs induce prolonged

microsaccadic inhibition. Exp. Brain Res. *177*, 196–208.

41.   Valsecchi, M., and Turatto, M. (2009). Microsaccadic responses in a bimodal oddball

task. Psychol. Res. *73*, 23–33.

42.   Reimer, J., Froudarakis, E., Cadwell, C.R., Yatsenko, D., Denfield, G.H., and Tolias, A.S.

(2014). Pupil Fluctuations Track Fast Switching of Cortical States during Quiet

Wakefulness. Neuron *84*, 355–362. Available at:

http://dx.doi.org/10.1016/j.neuron.2014.09.033.

43.   Leopold, D.A., and Logothetis, N.K. (1998). Microsaccades differentially modulate neural

activity in the striate and extrastriate visual cortex. Exp. Brain Res. *123*, 341–345.

44.   Martinez-Conde, S., Macknik, S.L., and Hubel, D.H. (2000). Microsaccadic eye

828   movements and firing of single cells in the striate cortex of macaque monkeys. Nat.

829   Neurosci. *3*, 251–258.

830   45.   Bair, W., and O'Keefe, L.R. (1998). The influence of fixational eye movements on the

831   response of neurons in area MT of the macaque. Vis. Neurosci. *15*, 779–786.

832   46.   Snodderly, D.M., Kagan, I., and Moshe, G. (2001). Selective activation of visual cortex

833   neurons by fixational eye movements: Implications for neural coding. Vis. Neurosci. *18*,

834   259–277.

835   47.   Herrington, T.M., Masse, N.Y., Hachmeh, K.J., Smith, J.E.T., Assad, J.A., and Cook, E.P.

836   (2009). The effect of microsaccades on the correlation between neural activity and

837   behavior in middle temporal, ventral intraparietal, and lateral intraparietal areas. J.

838   Neurosci. *29*, 5793–5805.

839   48.   Chen, C.Y., Ignashchenkova, A., Thier, P., and Hafed, Z.M. (2015). Neuronal response

840   gain enhancement prior to microsaccades. Curr. Biol. *25*, 2065–2074. Available at:

841   http://dx.doi.org/10.1016/j.cub.2015.06.022.

842   49.   Lowet, E., Gomes, B., Srinivasan, K., Zhou, H., Schafer, R.J., and Desimone, R. (2018).

843   Enhanced Neural Processing by Covert Attention only during Microsaccades Directed

844   toward the Attended Stimulus. Neuron *99*, 207-214.e3. Available at:

845   https://doi.org/10.1016/j.neuron.2018.05.041.

846   50.   Steinmetz, N.A., and Moore, T. (2019). Changes in the Response Rate and Response

847   Variability of Area V4 Neurons During the Preparation of Saccadic Eye Movements.

848   1171–1178.

849   51.   Khanna, S.B., Snyder, A.C., and Smith, M.A. (2019). Distinct sources of variability affect

850   eye movement preparation. J. Neurosci. *39*, 4511–4526.

851   52.   Supèr, H., and Lamme, V.A.F. (2007). Strength of figure-ground activity in monkey

852   primary visual cortex predicts saccadic reaction time in a delayed detection task. Cereb.

853   Cortex *17*, 1468–1475.

854   53.   Hanes, D.P., and Schall, J.D. (1996). Neural control of voluntary movement initiation.

855   Science (80-.). *274*, 427–430.

856   54.   Cook, E.P., and Maunsell, J.H.R. (2002). Dynamics of neuronal responses in macaque MT

857   and VIP during motion detection. Nat. Neurosci. *5*, 985–994.

858   55.   Roitman, J.D., and Shadlen, M.N. (2002). Response of neurons in the lateral intraparietal

859   area during a combined visual discrimination reaction time task. J. Neurosci. *22*, 9475–

860   9489.

861   56.   Huang, X., and Lisberger, S.G. (2009). Noise correlations in cortical area MT and their

862   potential impact on trial-by-trial variation in the direction and speed of smooth-pursuit eye

863   movements. J. Neurophysiol. *101*, 3012–3030.

864   57.   O'Leary, J.G., and Lisberger, S.G. (2012). Role of the lateral intraparietal area in

865   modulation of the strength of sensory-motor transmission for visually guided movements.

866   J. Neurosci. *32*, 9745–9754.

867   58.   Cowley, B.R., Snyder, A.C., Acar, K., Williamson, R.C., Yu, B.M., and Smith, M.A.

868   (2020). Slow drift of neural activity as a signature of impulsivity in macaque visual and

869   prefrontal cortex. bioRxiv, 2020.01.10.902403. Available at:

870   https://www.biorxiv.org/content/10.1101/2020.01.10.902403v1.

871   59.   Snyder, A.C., Yu, B.M., and Smith, M.A. (2018). Distinct population codes for attention

872  in the absence and presence of visual stimulation. Nat. Commun. *9*, 4382. Available at:

873  http://www.nature.com/articles/s41467-018-06754-5.

874 60. Cowley, B.R., Snyder, A.C., Acar, K., Williamson, R.C., Yu, B.M., and Smith, M.A.

875  (2018). The role that a slow drift in V4 population activity plays in perceptual decision-

876  making.

877 61. Jollife, I.T., and Cadima, J. (2016). Principal component analysis: A review and recent

878  developments. Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. *374*.

879 62. Murray, J.D., Bernacchia, A., Roy, N.A., Constantinidis, C., Romo, R., and Wang, X.J.

880  (2017). Stable population coding for working memory coexists with heterogeneous neural

881  dynamics in prefrontal cortex. Proc. Natl. Acad. Sci. U. S. A. *114*, 394–399.

882 63. Harvey, C.D., Coen, P., and Tank, D.W. (2012). Choice-specific sequences in parietal

883  cortex during a virtual-navigation decision task. Nature *484*, 62–68.

884 64. Mante, V., Sussillo, D., Shenoy, K. V., and Newsome, W.T. (2013). Context-dependent

885  computation by recurrent dynamics in prefrontal cortex. Nature *503*, 78–84.

886 65. Wang, J., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible timing by temporal

887  scaling of cortical responses. Nat. Neurosci. *21*, 102–112. Available at:

888  http://dx.doi.org/10.1038/s41593-017-0028-6.

889 66. Sohn, H., Narain, D., Meirhaeghe, N., and Jazayeri, M. (2018). Bayesian computation

890  through cortical latent dynamics. bioRxiv, 465419. Available at:

891  https://www.biorxiv.org/content/early/2018/11/08/465419.

892 67. Remington, E.D., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible

893  Sensorimotor Computations through Rapid Reconfiguration of Cortical Dynamics.

894  Neuron *98*, 1005-1019.e5. Available at: https://doi.org/10.1016/j.neuron.2018.05.020.

895   68.   Sadtler, P.T., Quick, K.M., Golub, M.D., Chase, S.M., Ryu, S.I., Tyler-Kabara, E.C., Yu,

896   B.M., and Batista, A.P. (2014). Neural constraints on learning. Nature *512*, 423–426.

897   Available at: http://dx.doi.org/10.1038/nature13665.

898   69.   Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D.

899   (2018). Spontaneous behaviors drive multidimensional, brain-wide population activity.

900   bioRxiv *7893*, 306019. Available at:

901   https://www.biorxiv.org/content/early/2018/12/28/306019?%3Fcollection=.

902   70.   Williamson, R.C., Doiron, B., Smith, M.A., and Yu, B.M. (2019). Bridging large-scale

903   neuronal recordings and large-scale network models using dimensionality reduction. Curr.

904   Opin. Neurobiol. *55*, 40–47. Available at: https://doi.org/10.1016/j.conb.2018.12.009.

905   71.   Cunningham, J.P., and Yu, B.M. (2014). Dimensionality reduction for large-scale neural

906   recordings. Nat. Neurosci. *17*, 1500–1509.

907   72.   Ruff, D.A., Ni, A.M., and Cohen, M.R. (2018). Cognition as a Window into Neuronal

908   Population Space. Annu. Rev. Neurosci. *41*, 77–97.

909   73.   Gallego, J.A., Perich, M.G., Miller, L.E., and Solla, S.A. (2017). Neural Manifolds for the

910   Control of Movement. Neuron *94*, 978–984. Available at:

911   http://dx.doi.org/10.1016/j.neuron.2017.05.025.

912   74.   Waterhouse, B.D., and Chandler, D. (2012). Evidence for broad versus segregated

913   projections from cholinergic and noradrenergic nuclei to functionally and anatomically

914   discrete subregions of prefrontal cortex. Front. Behav. Neurosci. *6*, 1–9.

915   75.   Chandler, D.J., Gao, W.J., and Waterhouse, B.D. (2014). Heterogeneous organization of

916   the locus coeruleus projections to prefrontal and motor cortices. Proc. Natl. Acad. Sci. U.

917   S. A. *111*, 6816–6821.

918   76.   Cerpa, J.C., Marchand, A.R., and Coutureau, E. (2019). Distinct regional patterns in

919   noradrenergic innervation of the rat prefrontal cortex. J. Chem. Neuroanat. *96*, 102–109.

920   Available at: https://doi.org/10.1016/j.jchemneu.2019.01.002.

921   77.   Schwarz, L.A., Miyamichi, K., Gao, X.J., Beier, K.T., Weissbourd, B., Deloach, K.E.,

922   Ren, J., Ibanes, S., Malenka, R.C., Kremer, E.J., *et al.* (2015). Viral-genetic tracing of the

923   input-output organization of a central noradrenaline circuit. Nature *524*, 88–92.

924   78.   Sharma, Y., Xu, T., Graf, W.M., Fobbs, A., Sherwood, C.C., Hof, P.R., Allman, J.M., and

925   Manaye, K.F. (2010). Comparative anatomy of the locus coeruleus in humans and

926   nonhuman primates. J. Comp. Neurol. *518*, 963–971.

927   79.   German, D.C., and Bowden, D.M. (1975). Locus ceruleus in rhesus monkey (Macaca

928   mulatta): A combined histochemical fluorescence, Nissl and silver study. J. Comp.

929   Neurol. *161*, 19–29.

930   80.   Kalwani, R.M., Joshi, S., and Gold, J.I. (2014). Phasic Activation of Individual Neurons in

931   the Locus Ceruleus/Subceruleus Complex of Monkeys Reflects Rewarded Decisions to

932   Go But Not Stop. J. Neurosci. *34*, 13656–13669.

933   81.   Lehmann, S.J., and Corneil, B.D. (2016). Transient Pupil Dilation after Subsaccadic

934   Microstimulation of Primate Frontal Eye Fields. J. Neurosci. *36*, 3765–3776.

935   82.   Clayton, E.C., Rajkowski, J., Cohen, J.D., and Aston-Jones, G. (2004). Phasic activation

936   of monkey locus ceruleus neurons by simple decisions in a forced-choice task. J.

937   Neurosci. *24*, 9914–9920.

938   83.   Aston-Jones, G., Rajkowski, J., and Kubiak, P. (1997). Conditioned responses of monkey

939   locus coeruleus neurons anticipate acquisition of discriminative behavior in a vigilance

940   task. Neuroscience *80*, 697–715.

941   84.   Aston-Jones, G., Rajkowski, J., Kubiak, P., and Alexinsky, T. (1994). Locus coeruleus

942   neurons in monkey are selectively activated by attended cues in a vigilance task. J.

943   Neurosci. *14*, 4467–4480.

944   85.   Hayat, H., Regev, N., Matosevich, N., Sales, A., Paredes-rodriguez, E., Krom, A.J.,

945   Bergman, L., Li, Y., Lavigne, M., Kremer, E.J., *et al.* (2020). Locus coeruleus

946   norepinephrine activity mediates sensory-evoked awakenings from sleep.

947   86.   Carter, M.E., Yizhar, O., Chikahisa, S., Nguyen, H., Adamantidis, A., Nishino, S.,

948   Deisseroth, K., and De Lecea, L. (2010). Tuning arousal with optogenetic modulation of

949   locus coeruleus neurons. Nat. Neurosci. *13*, 1526–1535.

950   87.   Li, Y., Hickey, L., Perrins, R., Werlen, E., Patel, A.A., Hirschberg, S., Jones, M.W.,

951   Salinas, S., Kremer, E.J., and Pickering, A.E. (2016). Retrograde optogenetic

952   characterization of the pontospinal module of the locus coeruleus with a canine adenoviral

953   vector. Brain Res. *1641*, 274–290. Available at:

954   http://dx.doi.org/10.1016/j.brainres.2016.02.023.

955   88.   Quinlan, M.A.L., Strong, V.M., Skinner, D.M., Martin, G.M., Harley, C.W., and Walling,

956   S.G. (2019). Locus coeruleus optogenetic light activation induces long-term potentiation

957   of perforant path population spike amplitude in rat dentate gyrus. Front. Syst. Neurosci.

958   *12*, 1–14.

959   89.   Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., and Wang, Q. (2017). Dynamic

960   Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from

961   Sympathetic, Not Parasympathetic, Contributions. Cell Rep. *20*, 3099–3112. Available at:

962   https://doi.org/10.1016/j.celrep.2017.08.094.

90.   Vazey, E.M., and Aston-Jones, G. (2014). Designer receptor manipulations reveal a role

of the locus coeruleus noradrenergic system in isoflurane general anesthesia. Proc. Natl.

Acad. Sci. U. S. A. *111*, 3859–3864.

91. Massot, C., Jagadisan, U.K., and Gandhi, N.J. (2019). Sensorimotor transformation elicits

systematic patterns of activity along the dorsoventral extent of the superior colliculus in

the macaque monkey. Commun. Biol. *2*, 1–14. Available at:

http://dx.doi.org/10.1038/s42003-019-0527-y.

92. Sparks, D.L., and Hartwich-Young, R. (1989). The deep layers of the superior colliculus.

Rev. Oculomot. Res. *3*, 213–255.

93. Kaufman, M.T., Churchland, M.M., Ryu, S.I., and Shenoy, K. V. (2014). Cortical activity

in the null space: Permitting preparation without movement. Nat. Neurosci. *17*, 440–448.

94. Elsayed, G.F., Lara, A.H., Kaufman, M.T., Churchland, M.M., and Cunningham, J.P.

(2016). Reorganization between preparatory and movement population responses in motor

cortex. Nat. Commun. *7*, 1–15. Available at: http://dx.doi.org/10.1038/ncomms13239.

95. Ames, K.C., and Churchland, M.M. (2019). Motor cortex signals corresponding to the two

arms are shared across hemispheres, mixed among neurons, yet partitioned within the

population response. bioRxiv, 1–45.

96. Stavisky, S.D., Kao, J.C., Ryu, S.I., and Shenoy, K. V. (2017). Motor Cortical Visuomotor

Feedback Activity Is Initially Isolated from Downstream Targets in Output-Null Neural

State Space Dimensions. Neuron *95*, 195-208.e9. Available at:

http://dx.doi.org/10.1016/j.neuron.2017.05.023.

97. Wang, C.-A., Baird, T., Huang, J., Coutinho, J.D., Brien, D.C., and Munoz, D.P. (2018).

Arousal Effects on Pupil Size, Heart Rate, and Skin Conductance in an Emotional Face

Task. Front. Neurol. *9*, 1–13.

98.   Samaha, J., Iemi, L., and Postle, B.R. (2017). Prestimulus alpha-band power biases visual

988   discrimination confidence, but not accuracy. Conscious. Cogn. *54*, 47–55. Available at:

989   http://dx.doi.org/10.1016/j.concog.2017.02.005.

990   99.   Benwell, C.S.Y., Tagliabue, C.F., Veniero, D., Cecere, R., Savazzi, S., and Thut, G.

991   (2017). Prestimulus EEG power predicts conscious awareness but not objective visual

992   performance. eNeuro *4*, 1–17.

993   100.   Romei, V., Gross, J., and Thut, G. (2010). On the role of prestimulus alpha rhythms over

994   occipito-parietal areas in visual input regulation: Correlation or causation? J. Neurosci. *30*,

995   8692–8697.

996   101.   Van Dijk, H., Schoffelen, J.M., Oostenveld, R., and Jensen, O. (2008). Prestimulus

997   oscillatory activity in the alpha band predicts visual discrimination ability. J. Neurosci. *28*,

998   1816–1823.

102.   Bellet, J., Chen, C.Y., and Hafed, Z.M. (2017). Sequential hemifield gating of α-and β-

behavioral performance oscillations after microsaccades. J. Neurophysiol. *118*, 2789–

2805.

103.   Engbert, R., and Kliegl, R. (2003). Microsaccades uncover the orientation of covert

attention. Vision Res. *43*, 1035–1045.

104.   Hafed, Z.M. (2013). Alteration of Visual Perception prior to Microsaccades. Neuron *77*,

775–786. Available at: http://dx.doi.org/10.1016/j.neuron.2012.12.014.

105.   Hafed, Z.M., and Clark, J.J. (2002). Microsaccades as an overt measure of covert attention

shifts. Vision Res. *42*, 2533–2545.

106.   Rolfs, M., Engbert, R., and Kliegl, R. (2005). Crossmodal coupling of oculomotor control

and spatial attention in vision and audition. Exp. Brain Res. *166*, 427–439.

107. Laubrock, J., Engbert, R., and Kliegl, R. (2005). Microsaccade dynamics during covert

attention. Vision Res. *45*, 721–730.

108. Smith, M.A., and Sommer, M.A. (2013). Spatial and Temporal Scales of Neuronal

Correlation in Visual Area V4. J. Neurosci. *33*, 5422–5432. Available at:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4782-12.2013.

109. Rabinowitz, N.C., Goris, R.L., Cohen, M., and Simoncelli, E.P. (2015). Attention

stabilizes the shared gain of V4 populations. Elife *4*, 1–24.

110. Shoham, S., Fellows, M.R., and Normann, R.A. (2003). Robust, automatic spike sorting

using mixtures of multivariate t-distributions. J. Neurosci. Methods *127*, 111–122.

111. Kelly, R.C., Smith, M.A., Samonds, J.M., Kohn, A., Bonds, A.B., Movshon, J.A., and

Sing Lee, T. (2007). Comparison of Recordings from Microelectrode Arrays and Single

Electrodes in the Visual Cortex. J. Neurosci. *27*, 261–264.

112. Brainard, D.H. (1997). The Psychophysics Toolbox. Spat. Vis. *10*, 433–436.

113. Pelli, D.G. (1997). The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. Spat. Vis. *10*, 437–442.

114. Scholes, C., McGraw, P. V., Nyström, M., and Roach, N.W. (2015). Fixational eye

movements predict visual sensitivity. Proc. R. Soc. B Biol. Sci. *282*.

115. Loughnane, G.M., Newman, D.P., Tamang, S., Kelly, S.P., and O'Connell, R.G. (2018).

Antagonistic interactions between microsaccades and evidence accumulation processes

during decision formation. J. Neurosci. *38*, 2163–2176.

116. Zuber, B.L., Stark, L., and Cook, G. (1965). Microsaccades and the velocity-amplitude

relationship for saccadic eye movements. Science (80-.). *150*, 1459–1460.