Running title: desert adaptation *Peromyscus*

Limited evidence for parallel evolution among desert adapted *Peromyscus* deer mice

Authors: Jocelyn P. Colella[1,2,†*], Anna Tigano[1,2], Olga Dudchenko[3,4], Arina D. Omer[4], Ruqayya Khan[4,5], Ivan D. Bochkov[4,5], Erez L. Aiden[3,4,6,7], Matthew D. MacManes[1,2]

[1]University of New Hampshire, Molecular, Cellular, and Biomedical Sciences Department, Durham, NH 03824, USA; jcolella.jc@gmail.com (JPC), anna.tigano@unh.edu (AT) matthew.macmanes@unh.edu (MDM)
[2]Hubbard Genome Center, University of New Hampshire, Durham, NH 03824, USA
[3]Department of Computer Science, Department of Computational and Applied Mathematics; Center for Theoretical and Biological Physics, Rice University, Houston, TX 77030, USA; Olga.Dudchenko@bcm.edu (OD), erez@erez.com (ELA)
[4]The Center for Genome Architecture, Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX 77030, USA; arinadanaomer@yahoo.com (ADO), Ruqayya.Khan@bcm.edu (RK), ibochkov33@gmail.com (IDB)
[5]Department of Computer Science, Department of Computational and Applied Mathematics, Rice University, Houston, TX 77030, USA;
[6]Shanghai Institute for Advanced Immunochemical Studies, ShanghaiTech University, Shanghai 201210, China
[7]School of Agriculture and Environment, University of Western Australia, Perth, WA 6009, Australia
[†]Biodiversity Institute, University of Kansas, Lawrence, KS 66045, USA
* Corresponding Author: Jocelyn.Colella@unh.edu

ABSTRACT

Phenotypic plasticity enables an immediate response to changing conditions, but for most

species, evolutionary change through adaptation will be more important for long-term survival.

Warming climate and increasing desertification urges the identification of genes involved in

heat- and dehydration-tolerance to better inform and target biodiversity conservation efforts.

Comparisons among extant desert adapted species can highlight parallel or convergent patterns

of genome evolution through the identification of shared signatures of selection. We generate

chromosome-level genome assembly for the canyon mouse (*Peromyscus crinitus*) and test for

signature of parallel evolution by comparing signatures of selective sweeps across population-

level genomic resequencing data from another desert specialist deer mouse (*P. eremicus*) and

a widely-distributed habitat generalist (*P. maniculatus*), that may locally adapted to arid

conditions. We identify few shared candidate loci involved in desert adaptation and do not find

support for a shared pattern of parallel evolution. Instead, we hypothesize divergent molecular

mechanisms of desert adaptation among deer mice, potentially tied to species-specific historical

demography, which may limit or enhance adaptation. We identify a number of candidate loci

experiencing selective sweeps in the *P. crinitus* genome that are implicated in osmoregulation

(*Trypsin, Prostasin*) and metabolic regulation (*Kallikrein, eIF2-alpha kinase GCN2, APPL1/2*),

which may be important to accommodating hot and dry environmental conditions.

Key words: dehydration, desert, parallel evolution, *Peromyscus,* thermoregulation

21   **INTRODUCTION**

Increasing global temperatures and altered patterns of precipitation threaten biodiversity

worldwide (Moritz et al. 2008; Cahill et al. 2013; Urban 2015). Phenotypic plasticity enables an

immediate response to changing conditions but adaptation through evolutionary change will be

critical for the long-term survival of most species (Hoffman and Sgrò 2011; Cahill et al. 2013).

Range shifts upward in elevation and latitude have been documented in a number of terrestrial

species and interpreted as a response to warming (Chen et al. 2011; Tingley and Beissinger

2013; Freeman et al. 2018); however, responses vary even among closely-related species or

populations (Hoffman and Willi 2008; Moritz et al. 2008). Geographic range shifts are often

governed by the physiological limits of species, which are in part controlled by genetics and

have been shaped by neutral and selective evolutionary forces across many generations.

Population genomics methods enable genome-wide scans for selection to identify genes and

molecular pathways that may be involved in local adaptation (Bassham et al. 2018; Garcia-

Elfring et al. 2019). For species adapted to similar environments, parallel or convergent

evolution can be inferred if a greater number of genes or phenotypes share signatures of

selection than would be expected under a purely stochastic model of evolution (e.g., drift). The

same gene or suite of genes consistently tied to a specific adaptive phenotype across distantly

related taxa is consistent with a signal of convergent evolution. In contrast, for taxa that share a

recent common ancestor, signatures of selection at the same loci may reflect parallel selection

on either new mutations or shared ancestral variation or similar demographic histories, resulting

in the same phenotypic effect. Evidence of convergent or parallel evolution can highlight

common loci involved in shared adaptive phenotypes (Rundle et al. 2000; McDonald et al.

2009), while a lack of concerted evolution may identify idiosyncratic evolutionary strategies to

achieve the same phenotypic result.

45   As a model taxon (Dewey and Dawson 2001; Bedford and Hoekstra 2015) inhabiting

varied environments throughout North America, deer mice (genus *Peromyscus*) are a frequent

and productive subject of adaptation studies (e.g., physiological, Storz 2007; behavioral, Hu and

Hoekstra 2017; genetic, Cheviron et al. 2012; Storz and Cheviron 2016; Tigano et al. 2020).

Physiological similarity of deer mice to lab mice (*Mus musculus*) further broadens the

implications of evolutionary and ecological investigations of *Peromyscus* by linking to biomedical

sciences. The genus *Peromyscus* (N = 67 species; mammaldiversity.org) is hypothesized to be

the product of a rapid ecological radiation across North America (origin ~8 Mya, radiation ~5.71

Mya; Platt et al. 2015), evident in their varied ecological niches and rich species diversity

(Glazier 1980; Riddle et al. 2000; Bradley et al. 2007; Platt et al. 2015; Lindsey 2020).

Peromyscus display tremendous thermoregulatory plasticity and can be found in extreme

thermal environments, ranging from cold, high elevations (Pierce and Vogt 1993; Cheviron et al.

2012, 2014; Kaseloo et al. 2014; Garcia-Elfring et al. 2019) to arid, hot deserts (Riddle et al.

2000; MacManes 2017; Tigano et al. 2020). Thermoregulation and dehydration tolerance are

complex physiological traits, suggesting that several potential evolutionary routes could lead to

the same phenotypic outcome. Within this framework, comparisons among divergent

Peromyscus species adapted to similar environments may highlight shared adaptive

polymorphisms or disparate evolutionary paths central to achieving the same phenotype

(Cheviron et al. 2012; Ivy and Scott 2017; Hu and Hoekstra 2017; Storz et al. 2019). In cold

environments, endotherms rely on aerobic thermogenesis to maintain constant internal body

temperatures. Changes in both gene expression and the functional properties of proteins in deer

mice (*P. maniculatus*) adapted to high-altitude suggest that changes in multiple hierarchical

molecular pathways may be common in the evolution of complex physiological traits, such as

thermoregulation (Wichman and Lynch 1991; Storz 2007; Cheviron et al. 2012; Storz and

Cheviron 2016; Garcia-Elfring et al. 2019). Nonetheless, research focused on thermoregulatory

adaptations in high-elevation species may be confounded by concurrent selection on other traits

conferring fitness benefits, such as high hemoglobin oxygen-binding affinity (Storz and Kelly

2008; Storz et al. 2010; Natarajan et al. 2015), which is critically important given the low partial

pressure of oxygen (PaO$_2$) associated with high elevation environments. In hot environments,

endotherms are challenged with balancing heat dissipation, energy expenditure, and water

retention (Anderson and Jetz 2005), resulting in a different suite of behavioral, physiological,

and molecular adaptations that enable survival (Schwimmer and Haim 2009; Degen 2012;

Kordonowy et al. 2016), but may be confounded by acute or chronic dehydration.

Understanding the biochemical mechanisms that enable survival under extreme environmental

stress can provide important insights into the nature of physiological adaptation.

Rapid environmental and ecological differentiation among *Peromyscus* species positions

these small rodents as models for generating hypotheses surrounding species responses to

accelerated warming (Cahill et al. 2013) and the potential for repeated adaptation to similar

environments among closely related species. Numerous *Peromyscus* species are adapted to

life in hot deserts, with each species and population subject to distinct histories of demographic

variation and gene flow. These idiosyncratic histories have a direct impact on evolution, as

effective population sizes (N_e) are inextricably linked to the efficacy of selection and

maintenance of genetic diversity in wild populations (Charlesworth 2009). Contemporary or

historical gene flow may further help or hinder adaptive evolution through homogenization or

adaptive introgression, respectively (Coyne and Orr 2004; Morjan and Reiseberg 2004; Jones et

al. 2018; Tigano and Friesen 2016). Native to the American West, the canyon mouse (*P.*

crinitus, Fig. 1) is a xerocole, highly specialized to life in hot deserts. In the lab, *P. crinitus* can

survive in the absence of exogenous water, with urine concentration levels similar to that of

desert-adapted kangaroo rats (*Dipodomys merriami*; Abbott 1971; MacMillen 1972; MacMillen

and Christopher 1975; MacMillien 1983), but without equivalently specialized renal anatomy

(Issaian et al. 2012). Canyon mice also exhibit a lower-than-expected body temperature relative

to their size and can enter environmentally-induced torpor in response to drought, food

limitation, or extreme external temperatures (McNab 1968; McNab and Morrison 1963; Morhardt

and Hudson 1966; Johnson and Armstrong 1987), which facilitates survival in highly-variable

and extreme desert environments. These phenotypes persist for multiple generations in the lab

indicating they have a genetic basis (McNab and Morrison 1968). Cactus mice (*P. eremicus*) are

frequently sympatric with *P. crinitus* and share the same adaptations described above for *P.*

crinitus (Veal and Caire 1979; Kordonowy et al. 2017). Thus, we expect these two species to

exhibit similar patterns of molecular adaptation. These two desert specialists belong to a

monophyletic clade of deer mice, which also includes *P. merriami, P. californicus*, *P. eva*, and *P.*

fraterculus, and is estimated to have diverged around 5-6 Mya (Platt et al. 2015). Other

members of this clade exhibit similar adaptations to desert environments, including urine

concentration, reduced water requirements, and environmentally-induced torpor (McNab and

Morrison 1963; Veal and Caire 1979) suggesting that desert adaptation may represent the

ancestral state of this clade. In contrast, the habitat generalist *P. maniculatus* (North American

deer mouse) is phylogenetically basal to the two desert specialists examined here and has a

geographically widespread distribution across North America. *Peromyscus maniculatus* inhabits

a wide range of thermal environments, including hot southwestern deserts and cool, high

elevations, but desert specialists are not its closest relatives and the species is not generally

considered a xerocole. Locally adapted desert populations of *P. maniculatus* (subspecies *P. m.*

sonoriensis), however, may exhibit patterns of selection similar to that of desert specialists,

either through the parallel selective retention of functional ancestral polymorphisms or

convergent selection on new mutations. Whole-genome assemblies are publicly available for

both *P. eremicus* (PRJEB33593, ERZ119825; Tigano et al. 2020) and *P. maniculatus*

(GCA_003704035.1), which positions these species as ideal comparatives against *P. crinitus* to

identify genes and regulatory regions associated with desert adaptation, including those unique

to desert specialists *P. eremicus* and *P. crinitus.*

Here, we investigate genomic signatures of selection among desert adapted

Peromyscus. We contrast signatures of selective sweeps across three related *Peromyscus*

species, two desert specialists (*P. crinitus* and *P. eremicus*) and one habitat generalist collected

in an arid environment (*P. maniculatus*). We hypothesize that similar genes or functional groups

will be under selection in related desert specialist species (*P. eremicus* and *P. crinitus*), due to

their shared recent common ancestor and mutual association with hot, arid environments. In

contrast, we hypothesize that *P. maniculatus* will show idiosyncratic evolutionary responses,

with arid adaptation in this clade having evolved independently in response to local conditions. If

similar genes and pathways are under selection in all three species, it would suggest local

adaptation of *P. maniculatus* to desert conditions, and potentially, parallel or convergent

evolution among divergent *Peromyscus* clades. Given the evolutionary distance of *P.*

maniculatus to the two desert adapted species, a shared signature of selection across all three

species may also indicate that adaptive responses to desert conditions are predictable and can

occur repeatedly and potentially on short evolutionary timescales. Finally, we place selective

sweep analyses into an evolutionary framework to interpret the varied evolutionary trajectories

available to small mammals to respond to changing environmental conditions and to account for

demographic and gene flow events.

MATERIALS AND METHODS

De novo genome sequencing and assembly

Wild mice were handled and sampled in accordance with the University of New Hampshire and

University of California Berkeley's Institutional Animal Care and Use Committee (130902 and

R224-0310, respectively) and California Department of Fish and Wildlife (SC-008135) and the

American Society of Mammalogists best practices (Sikes and Animal Care and Use Committee

of the American Society of Mammalogists 2016).

For the assembly of the *P. crinitus* genome, DNA was extracted from a liver subsample

from an individual collected in 2009 from the Philip L. Boyd Deep Canyon Desert Research

Center (DRDC) in Apple Valley, California. To generate a high-quality, chromosome-length

genome assembly for this individual we extracted high-molecular-weight genomic DNA using a

Qiagen Genomic-tip kit (Qiagen, Inc., Hilden, Germany). A 10X Genomics linked-reads library

was prepared according to the manufacturers protocol and sequenced to a depth of 70X on a

HiSeq 4000 (Novogene, Sacramento, California, USA). 10X Genomics reads were *de novo*

assembled into contigs using *Supernova* 2.1.1 (Weisenfeld et al. 2017). To arrange scaffolds

into chromosomes, a Hi-C library for *P. crinitus* was constructed and sequenced from primary

fibroblasts from the T.C. Hsu Cryo-Zoo at the University of Texas MD Anderson Cancer Center.

The Hi-C data were aligned to the supernova assembly using *Juicer* (Durand et al. 2016). Hi-C

genome assembly was performed using the *3D-DNA* pipeline (Dudchenko et al. 2017) and the

output was reviewed using *Juicebox Assembly Tools* (Dudchenko et al. 2018). The Hi-C data

are available on www.dnazoo.org/assemblies/Peromyscus_crinitus, where they can be

visualized using *Juicebox.js*, a cloud-based visualization system for Hi-C data (Robinson et al.

2018).

Benchmarking Universal Single-Copy Orthologs (*BUSCO* v3, using the Mammalia odb9

database; Simão et al. 2015) and *OrthoFinder2* (Emms and Kelly 2015) were used to assess

genome quality and completeness. Genome sizes were estimated for each species using

abyss-fac (Simpson et al. 2009) and the *assemblathon_stats.pl* script available at:

https://github.com/ucdavis-bioinformatics/assemblathon2-analysis/. *RepeatMasker* v.4.0 (Smit

et al. 2015) was used to identify repetitive elements. The genome was annotated using the

software package *MAKER* (3.01.02; Campbell et al. 2014). Control files, protein, and transcript

data used for this process are available at https://github.com/macmanes-

lab/pecr_genome/tree/master/annotation. We used *Mashmap (-f one-to-one --pi 90 -s 300000*;

Jain et al. 2017, 2018) to assess syntenic conservation between *P. crinitus* and *P. maniculatus*

genomes and alignments were plotted with the script *generateDotPlot.pl. Peromyscus crinitus*

chromosomes were renamed and sorted using *seqtk* (github.com/lh3/seqtk) following the *P.*

maniculatus chromosome naming scheme.

For comparative genomics analyses, we generated low-coverage whole-genome

resequencing data for nine *P. crinitus* and five *P. maniculatus* individuals collected from arid

sites in southern California (Fig. 1; Table S1). *Peromyscus crinitus* samples were also collected

from the University of California (UC) DCDRC and *P. maniculatus* were collected further East

from the UC Motte Rimrock (MOT) and Elliot Chaparral Reserves (ELL; Fig. 1). We also used

publicly available low-coverage whole-genome resequencing data from 26 *P. eremicus*

individuals, also collected from DCDRC and MOT, that were prepared and sequenced in parallel

(Tigano et al. 2020). All samples were collected in 2009, with the exception of eight *P. eremicus*

samples that were collected in 2018. Animals were collected live in Sherman traps and a 25 mg

ear-clip was taken from each individual and stored at -80ºC in 95% ethanol. Animals were

sampled from arid areas with average monthly temperatures between 9-40ºC and mean annual

rainfall of 15-18 cm. The Biotechnology Resource Center at Cornell University (Ithaca, NY,

USA) prepared genomic libraries using the Illumina Nextera Library Preparation kit (e.g., skim-

seq). Libraries were sequenced at Novogene (Sacramento, CA, USA) using 150 bp paired-end

reads from one lane on the Illumina NovaSeq S4 platform. *fastp* v. 1 (Chen et al. 2018) was

used to assess read quality and trim adapters. Sequences from all samples and all species

were mapped to the *P. crinitus* reference genome with *BWA* (Li and Durbin 2010) to enable

comparative analyses, duplicates were removed with *samblaster v. 0.1.24* (Faust and Hall

2014), and alignments were indexed and sorted using *samtools v. 1.10* (Li et al. 2009).

196   *Population Genomics*

We used the software package *ANGSD v. 0.93* (Korneliussen et al. 2014) to call variants from

low-coverage population genomic data from the three species (26 *P. eremicus,* 9 *P. crinitus,* 5

P. maniculatus) with high confidence. First, an initial list of high-quality SNPs was identified by

analyzing all samples from the three species together using the settings: *-SNP_pval 1e-6 -*

minMapQ 20 -minQ 20 -setMinDepth 20 -minInd 20 -minMaf 0.01. Then, allele frequencies for

each of those high-quality SNPs were calculated independently for each species, with the

following filtering steps: a minimum of half (-*minInd*) *P. crinitus* and *P. eremicus* samples and all

P. maniculatus samples had to meet independent quantity (-*minMapQ*) and quality (-*minQ*)

thresholds for each variable site.

Differentiation among species was examined using a multidimensional scaling (MDS)

analysis in *ANGSD*. MDS plots were generated in *R* v.3.6.1 (R Core Team 2017) based on the

covariance matrix. Cook's D was used to identify outliers (Cook and Weisberg 1984; Williams

1987). As an additional measure of differentiation, we estimated weighted and unweighted

global F_{ST} values for each species pair using *realSFS* in *ANGSD*. *NGSadmix v. 33* (Skotte et al.

2013) was used to fit genomic data into K populations to parse species-level differences and

provide a preliminary screen for genomic admixture under a maximum-likelihood model.

Individuals with < 90% assignment to a particular species were considered putatively admixed.

To examine the impact of coverage on the detection of admixture we also evaluated coverage

distributions among admixed and non-admixed individuals. Nonetheless, expanded sample

sizes with greater sequencing depth will be necessary to detail patterns of population structure

and introgression. We tested K = 1 through K = (N - 1), where N is the number of total

individuals examined. *NGSadmix* was run for all species combined and again for each species

independently.

We used Pairwise sequential Markovian Coalescent (*PSMC v. 0.6.5-r67;* Li and Durbin

2011) to examine historical demographic changes through time for each species. *PSMC*

analyses are not suitable for low-coverage genomes, therefore we used the higher-coverage

reads used to generate the high-quality, chromosome-length assemblies for each species (*P.*

crinitus, assembly methods detailed above; *P. eremicus*, SAMEA5799953, Tigano et al. 2020;

P. maniculatus: GCA_003704035.1, Harvard University). Quality reads (q > 20; *Skewer*, Jiang

et al. 2014) were mapped to their respective *de novo* assembled reference to identify

heterozygous sites. Reference assemblies were then indexed in *BWA*. *Samblaster* removed

PCR duplicates and *picard* (http://broadinstitute.github.io/picard/) added a read group to the

resulting bam file and generated a sequence dictionary (*CreateSequenceDictionary*) from the

reference assembly. For each species, *samtools* was used to sort and index alignments, and

variants were called using *mpileup* in *bcftools v1.10.2* (*call*, Li et al. 2009). Consensus

sequences were called in *VCFtools v 0.1.16* (*vcf2fq*, Danecek et al. 2011). *PSMC* distributions

of effective population size (N_e) were estimated with 100 bootstrap replicates and results were

visualized with *gnuplot v. 5.2* (Williams and Kelley 2010), using perl scripts available at

github.com/lh3/psmc. Output was scaled by a generation time of 6 months (0.5 yr, Millar 1989;

Pergams and Lacy 2008) and a general mammalian mutation rate of 2.2 x 10^{-9}

substitutions/site/year (Kumar and Subramanian 2002).

239   *Tests for selection & convergence*

We used *Sweepfinder2* (Nielsen et al. 2005; Huber et al. 2016; DeGiorgio et al. 2016) to detect

recent selective sweeps as it is compatible with low-coverage whole-genome data. This method

performs a composite likelihood ratio (CLR) test to detect deviations from the neutral site

frequency spectrum (SFS) that may indicate recent positive selection. *Sweepfinder2* was run on

both variant and invariant sites (Huber et al. 2016) for each species, excluding sex

chromosomes. Sex-chromosomes were excluded for three reasons: (1) sex chromosome

evolution is both rapid and complex relative to autosomes, (2) we had different sample sizes of

each sex across species, and (3) desert adaptations, the focus of this study, are unlikely to be

sex-specific. We repeated *Sweepfinder2* analyses on *P. eremicus*, initially analyzed by Tigano

et al. (2020), using an improved annotation scheme based on *Peromyscus*-specific data rather

than *Mus musculus*. Allele frequencies were estimated in *ANGSD* independently for each

species and converted to allele counts, and the site frequency spectrum (SFS) was estimated in

Sweepfinder2 from autosomes only. Identification of sweeps were based on the pre-computed

SFS and the CLR was calculated every 10,000 sites. Per Tigano et al. (2020), a 10 kbp window

size was selected as a trade-off between computational time and resolution. CLR values above

the 99.9[th] percentile of the empirical distribution for each species were considered to be

evolving under a model of natural selection, hereafter referred to as significant sweep sites.

Smaller sample sizes produce fewer bins in the SFS and a lower number of rare alleles may

impact both the overall SFS and local estimate surrounding sweep sites; therefore, we explored

the impact of sample sizes on *Sweepfinder2* results by downsampling the number of genomes

analyzed for each species to five individuals (the total number of low-coverage genomes

available for *P. maniculatus*) and compared sweep results between downsampled and all-

sample datasets for the three smallest chromosomes: 21, 22, and 23.

For each species, mean Tajima's D was calculated across the entire genome in non-

overlapping windows of 10 kbp and 1 kbp in *ANGSD*. Nucleotide diversity (π) was also

calculated in 10 kbp and 1 kbp windows and corrected based on the number of sites genotyped

(variant and invariant) per window. Tajima's D and π are expected to be significantly reduced in

regions surrounding selective sweeps (Smith and Haigh 1974; Kim and Stephan 2002),

therefore we also used a Mann-Whitney test ($p < 0.05$, after a Bonferroni correction for multiple

tests) to measure significant deviations from the global mean in 1 kbp and 10 kbp flanking

regions surrounding significant sweep sites. We also examined D and π for flanking regions

surrounding 27 candidate genes identified in a previous transcriptomic investigation of *P.*

eremicus and potentially involved in dehydration tolerance (MacManes 2017; Table S2).

Candidate loci include aquaporins (N = 12), sodium-calcium exchangers (*SLC8a1*), and *Cyp4*

genes belonging to the Cytochrome P450 gene family (N = 14). We used custom python scripts

to functionally annotate (I) the closest gene to each significant sweep site, (II) the nearest

upstream and downstream gene, regardless of strand (sense/antisense), and (III) the nearest

upstream and downstream gene on each strand. Dataset I follows the general assumption that

proximity between a significant sweep site and a protein-coding gene suggests interaction.

Dataset II represents an extension of that model by encompassing the most proximal gene in

each direction. Because *Sweepfinder2* is based on unphased data mapped to a consensus

sequence and our data is unphased, we do not have information indicating on which strand a

significant sweep site occurs. Therefore, dataset III encompasses strand-uncertainty by

including the two nearest genes to a significant sweep site on both strands. It should be noted

that the genes identified in smaller datasets (I, II) are nested within the larger datasets (II, III)

and by definition, the larger datasets include more noise, which may dilute a signature of

parallel evolution, but may better capture the true signal of selection. Hence, it is important to

critically examine numerous hierarchical gene subsets. Without a linkage map, these analyses

remain exploratory and can be better refined with estimates of linkage disequilibrium, linkage

block sizes, and gene density in future investigations. We tested genes from each dataset for

functional and pathway enrichment in Gene Ontology (GO) categories using *Panther v. 15.0* (Mi

et al. 2017) and extracted GO terms for each enriched functional group. We used *Mus musculus*

as a reference and a Bonferroni correction for multiple tests ($p < 0.05$) to correct for false

discoveries. Enriched GO terms were summarized and visualized in *REVIGO* (Reduce and

Visualize Gene Ontology, Supek et al. 2011) implemented at:

http://revigo.irb.hr/index.jsp?error=expired. As a test for similar evolutionary responses to desert

environments, overlap in the gene names and enriched GO terms associated with significant

selective sweeps was assessed for each dataset. Overlap was visualized in the R package

VennDiagram (Chen and Boutros 2011). To test for convergence, we used a Fisher's Exact

Test ($p < 0.05$) in the *GeneOverlap* package (Shen 2016) in R to assess whether gene or

enriched GO term overlap between species was greater than expected based on the total

number of genes/GO terms in the genome. To determine if signatures of selection were driven

by differences in sequencing depth, we calculated local coverage in 10 kbp windows

surrounding significant sweep sites and averaged local coverage estimates across all sweeps

on a single chromosome, using a modification of:

https://github.com/AdamStuckert/Ranitomeya_imitator_genome/blob/master/GenomeAssembly/

[DuplicateOrthologWorkbook.md](). Local coverage surrounding significant sweep sites for each

chromosome were compared to the chromosomal average calculated for each species

(*samtools coverage --min-MQ 20, --region chr*).

To compare patterns of gene family expansion and contraction potentially involved in

adaptation within the genus *Peromyscus*, we analyzed 14 additional genomes, including ten

Peromyscus species and four near outgroup rodent species: *Microtus ochrogaster, Neotoma*

lepida, Sigmodon hispidus, and *Mus musculus* (Table S3). To prevent bias driven by variable

assembly qualities, samples with < 70% complete mammalian BUSCOs were excluded from

downstream analyses, resulting in the final analysis of ten species (Table S3). Groups of

orthologous sequences (orthogroups) were identified in *Orthofinder2*. Invariant orthogroups and

groups that varied by more than 25 genes across taxa (custom python script: ortho2cafe.py)

were excluded. Our rooted species tree, estimated in *Orthofinder2*, was used to calculate a

single birth-death parameter (lambda) and estimate changes in gene family size using *CAFE*

v.4.2.1 (Han et al. 2013). Results were summarized using the python script

cafetutorial_report_analysis.py available from the Hahn Lab at:

hahnlab.github.io/CAFÉ/manual.html.

323   **RESULTS**

324   *Chromosome-length genome assembly for* P. crinitus

Linked reads combined with Hi-C scaffolding produced a high-quality, chromosome-length

genome assembly for *P. crinitus.* Our assembly has a contig N50 of 137,026 bp and scaffold

N50 of 97,468,232 bp, with 24 chromosome-length scaffolds. The anchored sequences in the

three *Peromyscus* genome assemblies were as follows: *P. crinitus* genome ~2.27 Gb, *P.*

eremicus ~2.5 Gb, and *P. maniculatus* ~2.39 Gb (Table 1). Our assembly has high contiguity

and completeness and low redundancy, as demonstrated by the presence of 89.3% complete

BUSCOs, 0.9% of duplicates, and 9.0% missing, excluding unplaced scaffolds. As anticipated

based on karyotypic analyses (Smalec et al. 2019), we found no significant variation in

chromosome number or major interchromosomal rearrangements between *P. crinitus* and *P.*

maniculatus (Fig. S4). We annotated 17,265 total protein coding genes in the *P. crinitus*

genome. Similar to other *Peromyscus* species, LINE1 (long interspersed nuclear elements) and

LTR (long terminal repeats) elements comprised 22.7% of the repeats in the *P. crinitus* genome,

with SINEs (short interspersed nuclear elements) representing an additional 9.6% (Table S5).

Although similar to other *Peromyscus* species, *P. crinitus* has the greatest total repeat content

(> 37%; see Tigano et al. 2020 Supplementary Table 2).

341   *Population Genomics*

MDS analysis parsed the three species into three well-separated clusters and identified no

outliers or evidence of admixture (Fig. S6). *NGSadmix* identified all three species as a single

group (K = 1) with the highest likelihood, but a three-population model neatly parsed the three

species as expected (Fig. S7, Table S8). *NGSadmix* analysis, which is more sensitive to low

sample sizes than MDS analyses, showed putative admixture in *P. crinitus* with at least three

individuals displaying 11-27% ancestry from *P. eremicus* and additional material from *P.*

maniculatus (4-16%). Variable samples sizes may impact assignment certainty and expanded

sequencing of additional *Peromyscus* species and populations will be required to identify

potential sources of introgressed material. Four *P. eremicus* individuals had < 90% assignment

probability to the *P. eremicus* species cluster, with a maximum of 15% assignment to a different

species cluster. Identification of admixture in both species was not biased by differences in

coverage, as low (2X), medium (8X), and high coverage (17X) samples were found to be

admixed at a < 90% assignment threshold (Fig. S9). No *P. maniculatus* individuals were

identified as admixed.

PSMC estimates of historical demography (> 10 kya, Nadachowska-Brzyska et al. 2016)

show greater variance and a higher overall N_e for *P. crinitus* relative to *P. eremicus* (Fig. 2).

Demographic estimates for *P. maniculatus* are included as an additional comparison but should

be interpreted with caution as they are based on sequence data from a captive-bred individual

and may not accurately reflect the demography of wild populations.

362   *Selection & convergence*

Sweepfinder2 results were generally consistent across downsampled (N = 5) and all-sample

datasets (9 *P. crinitus,* 26 *P. eremicus*). For example, there were no significant sweep sites

identified on chromosome 21 for *P. crinitus* using either dataset, and although more sweeps

were significant on chromosome 23 for the downsampled dataset (16 vs. 8), 15 of the 16

sweeps were proximal to a single protein coding gene (*S15A3*) that was also identified as

experiencing a significant sweep when using all available data. All genes proximal to significant

sweep sites in the downsampled *P. crinitus* dataset were also identified when all samples were

analyzed. Six additional genes were identified as experiencing selective sweeps when the

complete dataset was evaluated. Results for *P. eremicus* were slightly less consistent: identical

numbers of sweep sites were detected on chromosomes 22 and 23 with additional significant

sweep sites identified on chromosome 21 when all samples were examined. The majority of

sweep sites on chromosome 21 in *P. eremicus* were proximal to *G3P*, which represented a

significant sweep in both downsampled and all-sample datasets; however, there were eight

additional protein coding genes proximal to significant sweep sites that were detected using the

downsampled dataset, but not detected when all samples were included in the analysis (e.g.,

Peripherin-2, BICRAL, Mrps18b). We hypothesize that population structure may increase

inconsistency in these results, as the 26 *P. eremicus* samples represent two populations (Motte

and DCDRC) and three distinct collection events (Motte 2018, DCDRC 2009 and DCDRC 2018;

Table S1), whose representation in the reduced dataset may vary due to different sample sizes

and random selection of downsampled individuals. While the study design does not allow us to

distinguish between type 1 and type 2 statistical errors, we hypothesize that inconsistencies are

related to differences in statistical power, with greater power in the full datasets relative to the

subsampled datasets.

Within *P. crinitus* we identified a total of 209 significant sweep sites (Table S10), with

104 sites localized on chromosomes 9 and 16 experiencing major selective sweeps (Fig. 3). We

found 239 total significant sweep sites for *P. eremicus* (Table S11, Fig. S12). Despite the large

size of chromosome 1 and strong signature of selective sweeps in *P. eremicus*, we found no

significant sweep sites on this chromosome for *P. crinitus*. Finally, we identified a total of 213

significant sweep sites for *P. maniculatus* (Table S13), with 103 sites located on chromosome 4.

Despite general chromosomal synteny among *Peromyscus* species (Fig. S4, Tigano et al.

2020), the chromosomal distribution of sweep sites differed among species. For example, *P.

eremicus* had at least one significant sweep detected on every chromosome, while sweeps

were only detected on 8 or 13 chromosomes in *P. maniculatus* and *P. crinitus*, respectively. We

found a number of sweep sites were concentrated on chromosome 9 for both desert specialist

species, with additional significant sweep sites for *P. crinitus* localized on chromosome 16 (Fig.

3, S12; Table S10-11). Sweeps in *P. eremicus* were widespread across the genome, with a

large peak (56 significant sweep sites) on chromosome 1 (Table 11; Fig. S12). *Peromyscus*

maniculatus sweeps fell primarily on chromosomes 4 and 20 (Table S13). The chromosomal

distribution of significant sweep sites does not appear to be driven by differences in coverage

(Table S14). Average sequencing depth for 10 kbp windows surrounding each significant sweep

site did not differ significantly from the global average sequencing depth for *P. crinitus* (p = 0.25)

or *P. maniculatus* (p = 0.28). Local (10 kbp window) coverage surrounding significant sweeps

for *P. eremicus* were less consistent (Table S10, S11, S13; Supplementary Results). Fourteen

of 232 significant sweep sites (6%) in *P. eremicus* exhibited extreme local sequencing depths of

0 or >1,000, leading to a significant difference between overall mean sequencing depth and

sequencing depth surrounding sweep sites (p = 0.03), with sweep windows exhibiting higher

sequencing depths on average (73 vs. 25). If the 14 anomalous values are excluded,

sequencing coverage does not differ (p = 0.37) for *P. eremicus*.

The effect of a selective sweep extends beyond the specific site identified as the target

of positive selection; hence, putative outliers (CLR > 99.9%) are indicative of a sweep in the 10

kbp window but the specific nucleotides under selection cannot be identified. As coding genes

represent only a small proportion of the genome, if a sweep site does not fall in one of these

regions we assume that the target of selection may be a regulatory element affecting gene

expression of proximal coding genes. Under this assumption, we hierarchically examined

protein coding genes most proximal to each sweep site and chose not to set a distance

threshold, as regulatory elements are known to affect genes up to hundreds of kb away (e.g.,

Wallbank et al. 2016). On average the distance from a sweep site to the nearest coding gene

was 45 kbp in *P. crinitus* (range: 31 - 439,122 bp, median = 5 kbp) and much greater for both *P.*

maniculatus (average: 152 kbp; range: 190 - 513,562 bp, median = 111 kbp) and *P. eremicus*

(average: 117 kbp; range: 38 - 1,423,027 bp; median: 35 kbp), despite high assembly qualities

for all species and identical methods of gene annotation. For both *P. eremicus* and *P.*

maniculatus, only two significant sweep sites were localized within protein-coding genes (*P.*

eremicus: Meiosis-specific with OB domain-containing protein, Harmonin; *P. maniculatus:*

Dehydrogenase/reductase SDR family member 7B and Zinc finger protein 217; Table 2). In

contrast, for *P. crinitus* 12 significant sweep sites fell within 19 distinct candidate loci, many of

which code for multiple alternatively spliced transcripts (Table 2). Among the significant sweep

sites localized within *P. crinitus* coding sequences, we identified 19 enriched GO terms (3

Biological Process [BP], 9 Molecular Function [MF], 7 Cellular Component [CC]), with

functionality ranging from 'proteolysis' to 'hydrolase' activity (Fig. 4; Table S15). Functional

examination of candidate loci identified solute regulation as a key function, with genes

pertaining to calcium (*Trypsin-2* [*PRSS2*]) and zinc (*Kallikrein-4* [*KLK4*]) binding and sodium

regulation (*Prostasin* [*PRSS8*]) indicated as under selection.

Here we report the results for dataset II, as this dataset (i) ensures the inclusion of the

most proximal gene under selection, by including the most proximal gene on each strand, and

(ii) reduces noise associated with dataset III, which includes four genes proximal to each sweep

site. Results for datasets I and III are addressed in more detail in the Supplementary

Information. Examination of dataset II in *P. crinitus* identified 121 unique genes and 26 enriched

GO terms (8 Biological Processes [BP], 10 Molecular Functions [MF], 8 Cellular Component

[CC]), with functionality pertaining to metabolism (e.g., 'protein metabolic process',

'organonitrogen compound metabolic process', 'peptide metabolic process') and ribosomes (Fig.

4; Tables S10, S15). For *P. eremicus,* we identified 202 unique genes and 14 enriched GO

terms (0 BP, 1 MF, 13 CC) associated with selective sweeps, with functionality centered around

ribosomes (Table S11, S16). For *P. maniculatus,* we identified 215 unique genes and eight

enriched GO terms (0 BP, 1 MF, 7 CC) associated with selective sweeps (Table S13, S17). Two

genes and seven enriched GO terms that were proximal to sweep sites were shared between

the two desert specialists, but the number of shared genes was not significantly different from

what is expected by chance alone. Functional enrichment of *P. eremicus* and *P. maniculatus*

across all datasets was limited to ribosomes (e.g., 'structural constituent of ribosome', 'cytosolic

ribosome', 'ribosomal subunit'; Fig. 4; Table 3, S15-17). In contrast, functionality of enriched GO

terms for *P. crinitus* centered on metabolic processes, including protein breakdown, hydrolysis,

and cellular functionality (e.g., 'organelle', 'intracellular', 'cytoplasm'; Fig. 4; Table S15), in

addition to ribosomes.

Peromyscus eremicus and *P. maniculatus* shared significant overlap (p < 0.05) in

enriched GO terms across all hierarchical data subsets (I, II, III; Fig. 5). Significant overlap of

enriched GO terms was also detected between *P. crinitus* and both other *Peromyscus* species

for datasets II and III only, with no overlap detected for dataset I (Fig. 5). Significant overlap

between desert specialists *P. eremicus* and *P. crinitus* was only detected in dataset III. Overall,

GO terms and genes associated with ribosomal functionality were frequently shared among all

species examined, but a unique pattern of selection was not shared among desert specialists.

Species tree estimates (Fig. S18) were consistent with previous phylogenetic

investigations (Bradley et al. 2007). *Peromyscus crinitus* and *P. eremicus* are sister in our

species tree, but note that a number of intermediate taxa remain unsampled (e.g., *P. merriami*,

P. californicus). Among the species examined here, the two desert specialists are part of a

larger clade of desert adapted *Peromyscus* to the exclusion of *P. maniculatus*. The *maniculatus*

clade is comprised of *P. leucopus, P. polionotus*, and *P. maniculatus,* and the *nasutus-attwateri*

clade is most basal within *Peromyscus* (Fig. S18), consistent with Platt et al. (2015). For the

Peromyscus genus, we found 19,925 gene families that had experienced contractions, 502

expansions, and 12 families that were evolving rapidly. However, we found no gene families

experiencing significant expansions, contractions, or rapid evolution below the genus level.

Average Tajima's D (1 kbp windows) was negative for all species and ranged from -0.69

to -1.61. *Peromyscus crinitus* had the lowest Tajima's D value and *P. maniculatus* the highest

(Fig. S19-20). Global pairwise F_{ST} between species ranged from 0.20-0.27 (unweighted: 0.12-

0.17). Mean global π (1 kbp windows) was 0.005 (±0.005) for *P. crinitus,* 0.007 (± 0.007) for *P.*

eremicus, and 0.012 (± 0.010) for *P. maniculatus* (Fig. S21). Both Tajima's D and π for 1 and 10

kbp flanking regions surrounding significant selective sweep sites were significantly higher than

the global average for each species (Table S20). Only in *P. maniculatus* did we detect a

significant reduction in π surrounding significant sweep sites. Tajima's D for flanking regions

surrounding the *a priori* candidate loci identified by MacManes (2017) were also significantly

more positive in all three species (Table S20).

DISCUSSION

Continued and accelerating environmental change increases the exigency of accurately

predicting species responses to anthropogenic climate change. Adaptive evolutionary

responses vary among species and populations, even when subjected to similar environmental

selective pressures (Bi et al. 2015; Garcia-Elfring et al. 2019). Evidence of parallel *de novo*

molecular changes or selective retention of shared ancestral variation can highlight genes or

genomics regions, including but not limited to functional variants, haplotypes, or structural

features of the genome, that may be key to adaptation. Alternatively, the same adaptive

phenotype can evolve through alternative evolutionary strategies. Thus, it is possible, even

among related species, that adaptation to similar environmental conditions will not exhibit similar

patterns of molecular evolution despite similar adaptive phenotypes. We analyzed genome-wide

patterns of selective sweeps among three species of deer mice within the North American

genus *Peromyscus* to identify candidate loci involved in heat- and dehydration-tolerance. We

hypothesized that the desert specialists, *P. crinitus* and *P. eremicus,* would share genes or

pathways associated with selective sweeps that were not shared with phylogenetically ancestral

P. maniculatus. These patterns would be indicative of parallel selection (and therefore, parallel

evolution) on either *de novo* mutations or ancestral variation. Given the suite of desert

adaptations shared by these species, shared signatures of selection may relate to survival in

high-temperature, low-water environments. Additionally, we hypothesized that shared patterns

of selective sweeps and enriched functional groups across all three species, if present, would

highlight candidate loci underpinning local adaptation of *P. maniculatus* to arid conditions and

potentially identify common loci involved in the repeated evolution of desert adaptation.

Although the species examined here are monophyletic, the two desert specialists share a more

recent ancestor (Fig. S18) and there are number of unsampled taxa that phylogenetically

separate the desert specialists from *P. maniculatus* (e.g., *P. merriami*, *P. californicus*). For this

reason, we cannot distinguish between parallel and convergent evolution, and without evidence

of ancestral divergence followed by reconvergence, we will discuss shared signatures of

selection as parallel evolution hereafter.

Overall, we did not find support for parallel evolution among desert specialist species,

but we identified a number of candidate loci that may be important to desert adaptation in *P.*

crinitus. Instead of a shared mechanism of heat- and dehydration tolerance, we hypothesize

that the two desert specialists examined here may have adapted to similar environments

through divergent molecular mechanisms*,* with *P. crinitus* potentially responding through

genomic changes to protein coding genes and *P. eremicus* through transcriptional regulation of

gene expression. This hypothesis is based on the lack of overlap in selective sweeps between

desert specialists, the proximity of sweeps to protein coding genes in *P. crinitus* relative *to P.*

eremicus, and previous gene expression results for *P. eremicus* (Kordonowy and MacManes

2017; MacManes 2017). Molecular flexibility of thermoregulatory responses may have catalyzed

the radiation of *Peromyscus* in North America by enabling rapid exploitation of novel thermal

environments. Finally, the application of an evolutionary lens to the interpretation of genomic

patterns of selection, particularly one that integrates historical demography and gene flow, can

help parse varied evolutionary mechanisms (parallel vs. convergent, genomic vs.

transcriptomic) of molecular adaptation.

527   *Limited evidence of parallel evolution*

Identification of similar genes or functional groups under selection in different species adapted

to similar environments can provide evidence in support of parallel evolution. In contrast, we

found limited evidence of parallel evolution among desert-adapted *Peromyscus*. Few to no

enriched GO terms overlapped between desert specialists (Fig. 5). Only GO terms relating to

ribosomes (e.g., 'ribosome', 'ribosomal subunit', 'cystolic ribosome', etc.) overlapped between

all three *Peromyscus* species examined, with the most significant overlap in GO terms occurring

between *P. eremicus* and *P. maniculatus*. Although *P. maniculatus* are not generally xerocoles,

the individuals sequenced here were collected in arid regions of southern California (subspecies

P. m. sonoriensis)*.* Therefore, the shared signature of selection on ribosomes across all

examined *Peromyscus*, whether it reflects parallel evolution or the selective retention of shared

ancestral polymorphisms, may be associated with adaptation to hot and dry conditions or more

broadly relate to thermoregulatory plasticity among *Peromyscus* rodents. Few genes proximal to

selective sweeps were shared among all species, with only one instance of significant overlap:

ten genes were shared between the two desert specialists under dataset III (Fig. 5). Although

dataset III may be confounded by excess noise through the inclusion of additional protein

coding genes, this signature is potentially consistent with a parallel evolution. Again, many of

the genes shared between *P. crinitus* and *P. eremicus* are directly related to ribosomal

functionality (e.g., *RL36, RS26, RL15*) and also shared with *P. maniculatus*. Determining

whether these sweeps are the result of shared new mutations or ancestral variation and

whether selection on ribosomal functionality is unique to desert-adapted taxa or more broadly

relevant to the genus will require additional tests for selection and expanded taxonomic

sampling across the genus *Peromyscus*.

Cellular damage accumulates quickly in desert environments as a consequence of

increased thermal- and osmotic-stress (Lamitina et al. 2006; Burg et al. 2007). In response,

expression changes modulate osmoregulation by removing and replacing damaged proteins to

prevent cell death (Lamitina et al. 2006); hence, ribosomes, which play a critical role in protein

synthesis and degradation, are central to thermoregulatory responses (Porcelli et al. 2015).

Although we did not find significantly expanded or contracted gene families within the genus

Peromyscus, previous investigations of the entire Myodonta clade within Rodentia identified

multiple expanded or contracted gene families associated with ribosomes in *P. eremicus*

(Tigano et al. 2020). Here, ribosomes appear to be a potential target of parallel evolution in

desert-adapted *Peromyscus*, yet this genomic signature is not unique to this genus, nor to

desert-adapted species. First, the relative abundance of ribosome-associated genes throughout

the genome (>1000 GO annotations pertaining to ribosomes, Bult et al. 2019) may intrinsically

increase the representation of this functional group, especially at coarse resolution (10 kbp

windows). Second, selection on ribosomal functionality may be commonly experienced across

many species adapted to distinct thermal environments (metazoans; Porcelli et al. 2015).

Ribosomes are evolutionarily linked to the mitochondrial genomes of animals (Barreto and

Burton 2012; Bar-Yaacov et al. 2012) and accelerated mitochondrial evolution in animals has

led to compensatory, rapid evolution of ribosomal proteins (Osada and Akashi 2012; Barreto

and Burton 2013; Bar-Yaacov et al. 2012). Rapid mitochondrial diversification within

Peromyscus (Riddle et al. 2000; Bradley et al. 2007; Platt et al. 2015), coincident with the

ecological radiation of this genus (Lindsey 2020), suggests that equivalent, recent selection on

ribosomal proteins may be a key evolutionary innovation that enabled Peromyscine rodents to

successfully and quickly adapt to varied thermal environments. Alternatively, broad selection on

ribosomes across all species may also contribute to other, varied aspects of these species

biology. Comparisons among additional *Peromyscus* species will be necessary to test these

hypotheses in detail.

Evaporative cooling through sweating, panting, or salivating increases water loss and

challenges osmoregulatory homeostasis in a hot and dry climate (McKinley et al. 2018).

Thermal stress exacerbates dehydration by increasing evaporative water loss and if untreated,

can lead to cognitive dysfunction, motor impairment, and eventually death. In consequence,

osmoregulatory mechanisms are often under selection in extreme thermal environments

(MacManes and Eisen 2014; Marra et al. 2014). Consistent with the importance of

osmoregulation in desert species, four of the ten protein-coding genes that experienced a

significant selective sweep and were shared between desert specialist species (dataset III) are

involved in ion balance (Table 3). Proteins Trypsin-2 (*TRY2*) and Trimeric intracellular cation

channel type-B (*TM38B*) are associated with sweeps in both desert specialists and are involved

in calcium ion (Ca^{2+}) binding and release, respectively. DNA-directed RNA polymerase III

(*RPC1*) has also experienced a significant sweep in both desert specialists and influences

magnesium (Mg^{2+}) binding. Calcium and magnesium cations are among those essential for

osmoregulation (also, Na$^+$, K$^+$, Cl$^-$, HCO$_3^-$; Stockham and Scott 2008) and parallel selection on

these genes is consistent with the hypothesis that solute-carrier proteins are essential to

maintaining homeostasis in desert-specialized rodents (Marra et al. 2014; Kordonowy and

MacManes 2017). Additional genes involved in osmoregulation were identified as experiencing

selective sweeps only in *P. crinitus* (Table 2; Table S10). Prostatin (*PRSS8*), only found to be

under selection in *P. crinitus,* is critically responsible for increasing the activity of epithelial

sodium (Na$^+$) channels, which mediate sodium reabsorption through the kidneys (Narikiyo et al.

2002). Two more genes associated with Ca^{2+} regulation ([*PRSS2* and *TRYP*) and other genes

regulating zinc (*KLK4*) and iron (*NCOA4*) were also identified as targets of selective sweeps

exclusively in *P. crinitus.*

Genomic scans for selective sweeps based on the SFS are only one way to detect

signatures of parallel evolution and these methods can be sensitive to missing data, including

low-coverage and small sample sizes; thus, the putative roles of these candidate genes in

desert adaptation remains to be explored using additional methods with increased sequencing

depth (see Booker et al. 2017, Weigand and Leese 2018) and other experimental approaches

(e.g., MacManes 2017).

606   *Metabolic tuning: proteins-for-water or lipids-for-torpor?*

Hot deserts experience dramatic fluctuations in both food and water availability that challenge

species survival (Noy-Meir 1973; Silanikove 1994). Mammals accommodate high temperatures

by increasing body temperatures, to a point, and cold temperatures by aerobic thermogenesis

or metabolic suppression via the initiation of torpor or hibernation (Levesque et al. 2016). When

resources are scarce, metabolism relies exclusively on endogenous nutrients; carbohydrates

(e.g., sugars, glucose) are consumed immediately, then lipids, and eventually, proteins. Protein

oxidation has a low-energy return relative to lipid catabolism (Bar and Volkoff 2012), but yields

five times more metabolic water (Jenni and Jenni-Eiermann 1998; Gerson and Guglielmo

2011a, b; McCue et al. 2017). Therefore, in a low-water environment an early shift to protein

catabolism during periods of resource limitation may represent an important water source for

desert species (e.g., protein-for-water hypothesis; Mosin 1984; Jenni and Jenni-Eiermann,

1998; Gerson and Guglielmo, 2011a, b). Consistent with this hypothesis, we identified

numerous candidate genes that experienced selective sweeps in *P. crinitus* and that are

involved in the detection of metabolic-stress and shifts in metabolic fuel consumption. For

example, the gene eIF-2-alpha kinase GCN2 (*E2AK4*), which is responsible for sensing

metabolic stress in mammals and required for adaptation to amino acid starvation, experienced

the strongest selective sweep on chromosome 4 in *P. crinitus* (Fig. 3; Harding et al. 2003; Baker

et al. 2012; Taniuchi et al. 2016). Numerous candidate genes involved in oxidation

(Oxidoreductase NAD-binding domain-containing protein 1 [*Oxnad1*]), fat catabolism (Kallikrein-

6 [*KLK6*]), protein processing (Kallikrein-13 [*KLK13*]), and proteolysis (Kallikrein [*KLK4, KLK13*],

Trypsin [*PRSS2, TRYP, TRY2*], Chymotrypsin-like elastase family member 2A [*CELA2A*]) were

associated with significantly enriched GO terms in *P. crinitus*. Proteolysis was the most enriched

functional group in *P. crinitus* (Fig. 4; Table S15), potentially supporting the protein-for-water

hypothesis.

In contrast to the protein-for-water hypothesis, efficient lipid acquisition and storage may

be critical to enabling heat- and drought- induced torpor (Buck et al. 2002; Melvin and Andrews

2009), which allows long duration, low energy survival in desert adapted species, including

Peromyscus. Significant weight loss in experimentally-dehydrated *P. eremicus* and enhanced

thermogenic performance of high-altitude-adapted deer mice have been associated with

enhanced lipid metabolism (Cheviron et al. 2012; Kordonowy et al. 2016). At high altitudes,

increased lipid oxidation enables aerobic thermogenesis, but in hot deserts lipids may represent

a valuable energy source in a food-scarce environment (e.g., lipids-for-torpor hypothesis). Two

additional candidate genes, DCC-interacting protein 13-alpha and -beta (*APPL1*, *APPL2*),

experienced significant selective sweeps in *P. crinitus* and are important in glucose regulation,

insulin response, and fatty acid oxidation, potentially supporting the lipids-for-torpor hypothesis.

Laboratory manipulations of *APPL1* demonstrate protection against high-fat diet-induced

cardiomyopathy in rodents (Park et al. 2013) and *APPL2* is responsible for dietary regulation,

cold-induced thermogenesis, and cold acclimation (uniprot.org). Together, these genes play a

role in both obesity and dietary regulation. Both *APPL* genes are associated with obesity and

non-alcoholic fatty liver disease and their sweep signature in *P. crinitus* has relevant

connections to biomedical research that remain to be explored (Jiang et al. 2012; Barbieri et al.

2013). Physiological tests will be essential to determine whether desert-adapted deer mice

prioritize proteins or fats during periods of resource limitation (e.g., lipids-for-torpor) or extreme

dehydration (e.g., protein-for-water hypothesis).

Molecular rewiring of metabolic processes in response to environmental conditions has

been documented in a number of species (e.g., mammals, Velotta et al. 2020; birds, Xie et al.

2018; fruit flies, Mallard et al. 2018), but expression changes can also impact species

metabolism (Cheviron et al. 2012; Storz and Cheviron 2016). The capacity for rapid molecular

adaptation to distinct thermal environments through either transcriptomic regulation or changes

to protein coding genes, combined with thermoregulatory behavioral fine-tuning (e.g.,

nocturnality, aestivation, food caching, burrowing, dietary shifts), suggests there may be many

evolutionary strategies available for small mammals to accommodate increasing temperatures.

Anthropogenic change, however, is occurring at a rate that far outpaces the evolutionary

timescales on which these adaptations have naturally evolves; Thus, while metabolism and

metabolic plasticity represent fundamental phenotypes for anticipating species survival under

altered climate scenarios natural selection, alone they may be insufficient for species survival.

664   *Different evolutionary strategies, same result*

Diverse functional enrichment of the *P. crinitus* genome (Fig. 4), spanning metabolic and

osmoregulatory functions, in addition to the general functional enrichment of ribosomes,

identifies a number of candidate loci worthy of detailed examination. Additional, comparisons

across populations and environments will illustrate the influence of these loci and others in

thermoregulation, dehydration, and other adaptive traits. Significant selective sweeps that are

not shared among desert specialists, including most of the loci detected here, may still be

related to desert adaptation but could also be related to other aspects of this species biology.

There are multiple evolutionary routes to achieve environmental adaptation, most

notably through genomic changes in protein coding genes or transcriptional regulation of gene

expression. Lack of evidence for parallel evolution between desert specialists, the proximity of

significant selective sweeps to protein coding genes, diverse functional enrichment of *P. crinitus*

relative to *P. eremicus*, and previous gene expression results for *P. eremicus* (Kordonowy and

MacManes 2017; MacManes 2017) lead us to hypothesize alternative evolutionary strategies for

each desert specialist, each shaped by their independent demographic histories: *P. crinitus*

primarily through changes in protein coding genes and *P. eremicus* primarily through

transcriptional regulation. Evidence of many significant sweep sites in the *P. eremicus* genome,

located more distant from protein coding genes, and with functional enrichment restricted to

ribosomes, suggests that adaptation in this species may be driven more by selection on

regulatory or non-coding regions of the genome that impact gene expression, a hypothesis that

is consistent with transcriptomic investigations in this species (MacManes and Eisen 2014;

Kordonowy and MacManes 2017) and other *Peromyscus* and rodents (Cheviron et al. 2012;

Marra et al. 2014; Storz and Cheviron 2016). Without equivalent gene expression data for *P.

crinitus*, we cannot eliminate a similarly important role for transcriptional regulation and look

forward to testing this hypothesis in greater detail with RNAseq data. Transcriptional regulation

is a particularly useful mechanism for environmental acclimation, as these changes are more

transient relative to genomic changes and can enhance phenotypic flexibility (Garrett and

Rosenthal 2012; Rieder et al. 2015; Liscovitch-Brauer et al. 2017). Reduced variation is

expected near selective sweeps and can encompass tens to thousands of adjacent nucleotides

depending on recombination and the strength of selection (Fay and Wu 2000; Carlson et al.

2005), yet counter to expectations, Tajima's D and nucleotide diversity for regions flanking

putative selective sweeps were significantly higher than the global average for most

comparisons (Table S20). The same observation, elevated Tajima's D and nucleotide diversity

surrounding selective sweeps, was also made in *P. eremicus* (Tigano et al. 2020). This

counterintuitive pattern holds across different window sizes (1 kbp, 10 kbp) and warrants further

investigation.

Placing the results of selective sweep analyses within an evolutionary framework is also

critical to interpreting adaptive evolutionary responses. The deer mouse genus *Peromyscus*

originated approximately 8 Mya, followed by a massive radiation around 5-6 Mya that led to the

divergence of a monophyletic clade now comprised of desert adapted taxa, although the

ancestral state of this clade remains unknown. These desert adapted species may have

colonized arid environments through either a single or multiple invasions, with further

interspecific divergence thereafter (Platt et al. 2015). The expansion of North American deserts

following the conclusion of the last glacial maximum (~11 Kya; Pavlik et al. 2008) constrains the

adaptive timescales of contemporary desert species. The consistently stable and low historical

effective population size of *P. eremicus* suggests that his species has harbored less genetic

variation for selection to act on, despite similar levels of contemporary diversity relative to *P.*

crinitus (Fig. 2). In consequence, adaptive evolution of *P. eremicus* is likely to have been more

impacted by genetic drift (Allendorf 1986; Masel 2011) relative to *P. crinitus*, which historically

has a larger effective population size and therefore a broader pool of variation for selection to

act upon across evolutionary timescales, which could explain the higher diversity of genes and

enriched GO terms compared to *P. eremicus*. Within this context, changes in regulatory

elements that mediate gene expression may have been a more efficient means of

environmental adaptation available to *P. eremicus* (Allendorf 1986; Neme and Tautz 2016;

Mallard et al. 2018), whereas the larger historical effective population size of *P. crinitus* is more

conducive to the maintenance of higher levels of genetic diversity and may have enabled the

rapid evolution of protein coding sequences through mutational stochasticity, the reduced

impact of genetic drift, a larger pool of standing genetic variation, and potentially, gene flow.

Peromyscus crinitus experienced a historical demographic bottleneck prior to the formation of

North American deserts; Nevertheless, the recovered effective population size of *P. crinitus* is

much larger than *P. eremicus* and consistent with low levels of detected admixture in *P. crinitus*

(Fig. S7, Table S8). Negative Tajima's D values can indicate population expansion following a

bottleneck, consistent with both the demographic history of *P. crinitus* and putative admixture in

this species. Evidence of a historical bottleneck is also reinforced by moderate to high levels of

nucleotide diversity in *P. crinitus.* Repeated growth and contraction of rivers in the American

Southwest during Pleistocene glacial-interglacial cycles (0.7-0.01 Mya; Muhs et al. 2003; Van

Dam and Matzke 2016) would have provided iterative opportunities for connectivity and

introgression between incompletely-isolated *Peromyscus* species. Historical hybridization

between *P. crinitus* and one or more other Peromyscine species, likely unsampled here, may

have accelerated adaptation in *P. crinitus* through the rapid influx of novel mutation

combinations through adaptive introgression, a hypothesis that warrants further investigation

through expanded taxonomic sampling and explicit tests of adaptive introgression. Low-

coverage whole-genome resequencing is optimal for population genomics investigations

(O'Rawe et al. 2015; da Fonseca et al. 2016), but limits detailed analyses of historical

introgression. We look forward to testing this hypothesis with expanded population sampling

and increased sequencing depth. Finally, linkage disequilibrium decay is also weaker in larger

populations, where recombination is higher, therefore it's possible that the shorter distance

between significant sweep sites and the nearest coding gene in *P. crinitus* could be due to the

larger historical population sizes of this species relative to *P. eremicus.* However, the

evolutionary scales of *PSMC* and *Sweepfinder2* do not overlap, *PSMC* characterizes historical

demography beyond 10 kya, whereas selective sweeps have occurred recently. Overall,

incorporating an evolutionary perspective into the interpretation of selection patterns has

important implications for understanding species responses to changing climate, as historical

demography and gene flow, in addition to selection, shape genetic diversity over evolutionary

timescales.

750   *Conclusion*

Contrasting patterns of selective sweeps and evolutionary histories between different species

experiencing similar environmental pressures can provide powerful insights into the adaptive

potential of species. We used comparative and population genomic analyses of three

Peromyscus species to identify candidate loci that may underlie adaptations to desert

environments. Candidate loci identified in *P. crinitus* serve to inform future investigations

focused on predicting potential for adaptation and identifying the causes of warming-related

population declines (Cahill et al. 2013). The identification of numerous targets of selection within

P. crinitus highlights multiple molecular mechanisms (metabolic switching, osmoregulatory

tuning) associated with physiological responses to deserts that warrant further investigation..

Our approach demonstrates the importance of placing genomic selection analyses into an

evolutionary framework to anticipate evolutionary responses to change.

763   **ACKNOWLEDGEMENTS**

We thank A. S. Westbrook for computational support; the Premise computing cluster at the

University of New Hampshire, where all analyses were conducted; the Biotechnology Resource

Center at Cornell University for preparation of whole-genome resequencing libraries;

Christopher Tracy for access to the Boyd Deep Canyon Reserve; Jim Patton for desert field

expertise; Sen Pathak, Asha Multani, Richard Behringer for providing the fibroblast samples

from the T.C. Hsu Cryo-Zoo at the University of Texas M.D. Anderson Cancer Center; DNA Zoo

for generating Hi-C data; Pawsey Supercomputing Centre with funding from the Australian

Government and the Government of Western Australia for computational support of the DNA

Zoo effort and the Museum of Southwestern Biology at the University of New Mexico for loaned

tissue materials; two anonymous reviewers whose comments significantly improved the

manuscript. This work was funded by the National Institute of Health National Institute of

General Medical Sciences to MDM (1R35GM128843).

777   **DATA AVAILABILITY STATEMENT**

The draft assembly data are housed on the European Nucleotide Archive (ENA) under project

ID PRJEB33592. The Hi-C data is available on SRA (SRX7041777, SRX7041776,

SRX7041773) under the DNA Zoo project accession PRJNA512907. The *P. crinitus* genome

assembly is available at https://www.dnazoo.org/assemblies/Peromyscus_crinitus. Whole-

genome resequencing data for *P. crinitus* are available on ENA under project ID PRJEB35488.

Custom python scripts and other bash scripts used in analysis are available at:

https://github.com/jpcolella/Peromyscus_crinitus.

REFERENCES

Abbott, K. (1971). Water economy of the canyon mouse, *Peromyscus crinitus stephensi.*

Comparative Biochemistry and Physiology, 38A: 37–52.

Allendorf, F. (1986). Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology,* 5:

181–190.

Anderson, K., & Jetz, W. (2005). The broad-scale ecology of energy expenditure of

endotherms. *Ecology Letters,* 8: 310–318.

Baker, B., Nargund, A., Sun, T., & Haynes, C. (2012). Protective coupling of mitochondrial

function and protein synthesis via the eIF2α kinase GCN-2. *PLoS Genetics,* 8: e1002760.

Bar, N., & Volkoff, H. (2012). Adaptation of the physiological, endocrine and digestive system

functions to prolonged food deprivation in fish. In Comparative physiology of fasting,

starvation, and food limitation (pp. 69–90). New York: Springer.

Bar-Yaacov, D., Blumberg, A., & Mishmar, D. (2012). Mitochondrial-nuclear co-evolution and its

effects on OXPHOS activity and regulation. *Biochimica et Biophsyica Acta (BBA)-Gene*

Regulatory Mechanisms, 1819: 1107–1111.

Barbieri, M., Esposito, A., Angellotti, E., Rizzo, M., Marfella, R., & Paolisso, G. (2013).

Association of genetic variation in adaptor protein APPL1/APPL2 loci with non-alcoholic fatty

liver disease. *PLOS ONE,* 8: e71391.

Barreto, F., & Burton, R. (2013). Evidence for compensatory evolution of ribosomal proteins in

response to rapid divergence of mitochondrial rRNA. *Molecular Biology and Evolution,* 30:

310–314.

Bassham, S., Catchen, J., Lescak, E., von Hippel, F., & Cresko, W. (2018). Repeated selection

of alternatively adapted haplotypes creates sweeping genomic remodeling in stickleback.

Genetics, 209: 921–939.

Bedford, M., & Hoekstra, H. (2015). The natural history of model organisms: *Peromyscus* mice

as a model for studying natural variation. *Elife,* 4: e06813.

Bi, K., Linderoth, T., Singhal, S., Vanderpool, D., Patton, J., Nielsen, R., … Good, J. (2015).

Temporal genomic contrasts reveal rapid evolutionary responses in an alpine mammal during

recent climate change. *PLoS Genetics,* 15: e1008119.

Booker, T., Jackson, B., & Keightley, P. (2017). Detecting positive selection in the genome.

BMC Biology, 15. doi: https://doi.org/10.1186/s12915-017-0434-y

Bradley, R., Durish, N., Rogers, D., Miller, J., Engstrom, M., & Kilpatrick, C. (2007). Toward a

molecular phylogeny for *Peromyscus*: Evidence from mitochondrial cytochrome-b

sequences. *Journal of Mammalogy,* 88: 1146–1159.

Buck, M., Squire, T., & Andrews, M. (2002). Coordinate expression of PDK4 gene: a means of

regulating fuel selection in a hibernating mammal. *Physiological Genomics,* 8: 5–13.

Bult, C. J., Blake, J. A., Smith, C. L., Kadin, J. A., Richardson, J. E., and the Mouse Genome

Database Group. (2019). Mouse Genome Database (MGD). *Nucleic Acids Research*, 8:

D801-D806.

Burg, M., Ferraris, J., & Dmitrieva, N. (2007). Cellular response to hyperosmotic stresses.

Physiological Reviews, 87: 1441–1474.

Cahill, A., Aiello-Lammens, M., Fisher-Reid, M., Hua, X., Karanewsky, C., Yeong Ryu, H., …

Wiens, J. (2013). How does climate change cause extinction? *Proceedings of the Royal*

Society B: Biological Sciences, 280: 20121890.

Campbell, M., Holt, C., Moore, B., & Yandell, M. (2014). Genome annotation and curation using

MAKER and MAKER-P. *Current Protocols in Bioinformatics,* 48: 4–11.

Carlson, C., Thomas, D., Eberle, M., Swanson, J., Livingston, R., Rieder, M., & Nickerson, D.

(2005). Genomic regions exhibiting positive selection identified from dense genotype data.

Genome Research, 15: 1553–1565.

Charlesworth, B. (2009). Effective population size and patterns of molecular evolution and

variation. *Nature Reviews Genetics,* 10: 195–205.

Chen, H., & Boutros, P. (2011). VennDiagram: a package for the generation of highly-

customizable Venn and Euler diagrams in R. *BMC Bioinformatics,* 12: 35

Chen, I., Hill, J., Ohlemueller, R., Roy, D., & Thomas, C. (2011). Rapid range shifts of species

associated with high levels of climate warming. *Science*, 333: 1024–1026.

Chen, S., Zhou, Y., Chen, Y., & Gu, H. (2018). fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics,* 34: i884–i890.

Cheviron, Z., Bachman, G., Connaty, A., McClelland, G., & Storz, J. (2012). Regulatory

changes contribute to the adaptive enhancement of thermogenic capacity in high-altitude

deer mice. *Proceedings of the National Academy of Sciences,* 109: 8635–8640.

Cheviron, Z., Connaty, A., McClelland, G., & Storz, J. (2014). Functional genomics of

adaptation to hypoxic cold-stress in high-altitude deer mice: transcriptomic plasticity and

thermogenic performance. *Evolution,* 68: 48–62.

Cook, R., & Weisberg, S. (1984). Residuals and influence in Regression. Wiley.

Coyne, J. A., & Orr, H. A. (2004). Speciation. Sunderland, MA: Sinauer Associates.

da Fonseca, R. R., Albrechtsen, A., Themudo, G., Ramos-Madrigal, J., Sibbesen, J., Maretty,

L., … Pereira, R. (2016). Next-generation biology: sequencing and data analysis approaches

for non-model organisms. *Marine Genomics*, 30: 3–13. doi: 10.1016/j.margen.2016.04.012

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … 1000

Genome Project Data Process Subgroup. (2011). The variant call format and VCFtools.

Bioinformatics, 27: 2156–2158.

Degen, A. (2012). Ecophysiology of small desert mammals. Springer Science & Business

Media.

DeGiorgio, M., Huber, C., Hubisz, M., Hellmann, I., & Nielsen, R. (2016). SweepFinder2:

increased sensitivity, robustness and flexibility. *Bioinformatics,* 32: 1895–1897.

Dewey, M., & Dawson, W. (2001). Deer mice: "the *Drosophila* of North American mammalogy.

Genesis, 29: 105–109.

Dudchenko, O., Batra, S., Omer, A., Nyquist, S., Hoeger, M., Durand, N., … Aiden, E. (2017).

De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length

scaffolds. *Science,* 356: 92–95.

Dudchenko, O., Shamim, M., Batra, S., Durand, N., Musial, N., Mostofa, R., … Aiden, E. (2018).

The Juicebox Assembly Tools module facilitates de novo assembly of mammalian genomes

with chromosome-length scaffolds for under $1000. *BioRxiv.* doi:

https://doi.org/10.1101/254797

Durand, N., Shamim, M., Machol, I., Rao, S., Huntley, M., Lander, E., & Aiden, E. (2016). Juicer

provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell*

Systematics, 3: 95–98.

Emms, D. M., & Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. *Genome Biology,* 16:

157.

Faust, G., & Hall, I. (2014). SAMBLASTER: fast duplicate marking and structural variant read

extraction. *Bioinformatics,* 30: 2503–2505.

Fay, J., & Wu, C.-I. (2000). Hitchhiking under positive Darwinian selection. *Genetics*, 155:

1405–1413.

Freeman, B., Lee-Yaw, J., Sunday, J., & Hargreaves, A. (2018). Expanding, shifting and

shrinking: The impact of global warming on species' elevational distributions. *Global Ecology*

and Biogeography, 27: 1268–1276.

Fumagalli, M., Vieira, F., Linderoth, T., & Nielsen, R. (2014). ngsTools: methods for population

genetics analyses from next-generation sequencing data. *Bioinformatics,* 30: 1486–1487.

Garcia-Elfring, A., Barrett, R., & Millien, V. (2019). Genomic signatures of selection along a

climatic gradient in the northern range margin of the white-footed mouse (*Peromyscus*

leucopus). *Journal of Heredity,* 110: 684–695.

Garrett, S., & Rosenthal, J. (2012). RNA editing underlies temperature adaptation in K+

channels from polar octopuses. *Science,* 335: 848–851.

Gerson, A., & Guglielmo, C. (2011a). Flight at low ambient humidity increases protein

catabolism in migratory birds. *Science,* 333: 1434–1436.

Gerson, A., & Guglielmo, C. (2011b). House sparrows (*Passer domesticus*) increase protein

catabolism in response to water restriction. *American Journal of Physiology,* 300: R925–

R930.

Glazier, D. (1980). Ecological shifts and the evolution of geographically restricted species of

North American *Peromyscus* (mice). *Journal of Biogeography*, 7: 63–83.

Han, M., Thomas, G., Lugo-Martinez, J., & Hahn, M. (2013). Estimating gene gain and loss

rates in the presence of error in genome assembly and annotation using CAFE 3. *Molecular*

Biology and Evolution, 30: 1987-1997.

Harding, H., Zhang, Y., Zeng, H., Novoa, I., Lu, P., Calfon, M., … Ron, D. (2003). An integrated

stress response regulates amino acid metabolism and resistance to oxidative stress.

Molecular Cell, 11: 619–633.

Hoffmann, A., & Sgrò, C. (2011). Climate change and evolutionary adaptation. *Nature,* 470:

479–485.

Hoffmann, A., & Willi, Y. (2008). Detecting genetic responses to environmental change. Nature

Reviews: *Genetics,* 9: 421–432.

Hu, C., & Hoekstra, H. (2017). *Peromyscus* burrowing: A model system for behavioral evolution.

Seminars in Cell and Developmental Biology, 61: 107–114.

Huber, C., DeGiorgio, M., Hellmann, I., & Nielsen, R. (2016). Detecting recent selective sweeps

while controlling for mutation rate and background selection. *Molecular Ecology,* 25: 142–

156.

Issaian, T., Urity, V., Dantzler, W., & Pannabecker, T. (2012). Architecture of vasa recta in the

renal inner medulla of the desert rodent *Dipodomys merriami*: potential impact on the urine

concentrating mechanism. *American Journal of Physiology - Regulatory, Integrative and*

Comparative Physiology, 303: R748–R756.

Ivy, C., & Scott, G. (2017). Control of breathing and ventilatory acclimatization to hypoxia in

deer mice native to high altitudes. *Acta Physiologica,* 221: 266–282.

Jain, C., Koren, S., Dilthey, A., Phillippy, A. M., Aluru, S. (2018) A fast adaptive algorithm for

computing whole-genome homology maps. *Bioinformatics*, 34: i748-i756.

Jain, C., Dilthey A, Koren, S, Aluru, S., Phillippy, A. M. (2017) A fast approximate algorithm for

mapping long reads to large reference databases. In: Sahinalp S. (eds) *Research in*

Computational Molecular Biology. RECOMB 2017. Lecture Notes in Computer Science,

10229. Springer, Cham.

Jenni, L., & Jenni-Eiermann, S. (1998). Fuel supply and metabolic constraints in migrating birds.

Journal of Avian Biology, 29: 521–528.

Jiang, H., Lei, R., Ding., S.-W., Zhu, S. (2014). Skewer: a fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. *BMC Bioinformatics*, 15: 182.

Jiang, S., Fang, Q., Yu, W., Zhang, R., Hu, C., Dong, K., … Jia, W. (2012). Genetic variations in

APPL2 are associated with overweight and obesity in a Chinese population with normal

glucose tolerance. *BMC Medical Genetics,* 13: 22.

Johnson, D., & Armstrong, D. (1987). *Peromyscus crinitus. Mammalian Species,* 287: 1–8.

Jones, M., Mills, L., Alves, P., Callahan, C., Alves, J., Lafferty, D., … Good, J. (2018). Adaptive

introgression underlies polymorphic seasonal camouflage in snowshoe hares. *Science,* 360:

1355–1358.

Kaseloo, P., Crowell, M., & Heideman, P. (2014). Heritable variation in reaction norms of

metabolism and activity across temperatures in a wild-derived population of white-footed

mice (*Peromyscus leucopus*). *Journal of Comparative Physiology B, 184*(4): 525–534.

Kim, Y., & Stephan, W. (2002). Detecting a local signature of genetic hitchhiking along a

recombining chromosome. *Genetics,* 160: 765–777.

Kordonowy, L., & MacManes, M. (2017). Characterizing the reproductive transcriptomic

correlates of acute dehydration in males in the desert-adapted rodent, *Peromyscus eremicus*.

BMC Genomics, 18: 473.

Kordonowy, L., Lombardo, K., Green, H., Dawson, M., Bolton, E., LaCourse, S., & MacManes,

M. (2016). Physiological and biochemical changes associated with acute experimental

dehydration in the desert adapted mouse, *Peromyscus eremicus. Physiological Reports,* 5:

e13218.

Korneliussen, T., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *Bioinformatics*, 15: 356.

Kumar, S., & Subramanian, S. (2002). Mutation rates in mammalian genomes. Proceedings of

the National Academy of Sciences, 99: 803–808.

Lamitina, T., Haung, C., & Strange, K. (2006). Genome-wide RNAi screening identifies protein

damage as a regulator of osmoprotective gene expression. *Proceedings of the National*

Academy of Sciences of the United States of America, 103: 12173–12178.

Levesque, D., Nowack, J., & Stawski, C. (2016). Modelling mammalian energetics: the

heterothermy problem. *Climate Change Responses,* 3: 7.

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics,* 26: 589–595.

Li, Heng, & Durbin, R. (2011). Inference of human population history from whole genome

sequence of a single individual. *Nature,* 475: 493–496.

Li, H., Handsaker, B., Wysoker, A., Fennel, T., Ruan, J., Homer, N., … 1000 Genome Project

Data Process Subgroup. (2009). The sequence alignment/map format and SAMtools.

Bioinformatics, 25: 2078–2079.

Lindsey, L. (2020). Utilizing genomic applications to examine patterns of diversification in

deermice (Rodentia: Cricetidae: *Peromyscus*). Texas Tech Dissertation.

965 Liscovitch-Brauer, N., Alon, S., Porath, H., Elstein, B., Unger, R., Ziv, T., … Eisenberg, E.

966   (2017). Trade-off between transcriptome plasticity and genome evolution in Cephalopods.

967   *Cell,* 169: 191–202.

968 MacManes, M., & Eisen, M. (2014). Characterization of the transcriptome, nucleotide sequence

969   polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus.*

970   *PeerJ,* 2: e642.

971 MacManes, M. (2017). Severe acute dehydration in a desert rodent elicits a transcriptional

972   response that effectively prevents kidney injury. American Journal of Physiology. *Renal*

973   *Physiology,* 313: F262–F272.

974 MacMillen, R., & Christopher, E. (1975). The water relations of two populations of non-captive

975   desert rodents. In Environmental physiology of desert organisms ((N. F. Hadley, ed.): pp.

976   117–137). Stroudsburg, Pennsylvania: Dowden, Hutchinson, and Ross.

977 MacMillen, R. (1972). Water economy of nocturnal desert rodents. *Symposia of the Zoological*

978   *Society of London,* 31: 147–174.

979 MacMillen, R. (1983). Water regulation in *Peromyscus. Journal of Mammalogy,* 64: 38–47.

980 Mallard, F., Nolte, V., Tobler, R., Kapun, M., & Schlötterer, C. (2018). A simple genetic basis of

981   adaptation to a novel thermal environment results in complex metabolic rewiring in

982   *Drosophila. Genome Biology,* 19: 1-15. doi: 10.1186/s13059-018-1503-4

983 Marra, N., Romero, A., & DeWoody, J. (2014). Natural selection and the genetic basis of

984   osmoregulation in heteromyid rodents as revealed by RNA-seq. *Molecular Ecology,* 23:

985   2699–2711.

986 Masel, J. (2011). Genetic drift. *Current Biology,* 21: R837–R838.

987 McCue, M., Sandoval, J., Beltran, J., & Gerson, A. (2017). Dehydration causes increased

988   reliance on protein oxidation in mice: a test of the protein-for-water hypothesis in a mammal.

989   *Physiological and Biochemical Zoology,* 90: 359–369.

McDonald, M., Gehrig, S., Meintjes, P., Zhang, X.-X., & Rainey, P. (2009). Adaptive divergence

in experimental populations of *Pseudomonas fluorescens*. IV. Genetic constraints guide

evolutionary trajectories in a parallel adaptive radiation. *Genetics,* 183: 1041–1053.

McKinley, M., Martelli, D., Pennington, G., Trevaks, D., & McAllen, R. (2018). Integrating

competing demands of osmoregulatory and thermoregulatory homeostasis. *Physiology,* 33:

170–181.

McNab, B. (1963). The influence of fat deposits on the basal rate of metabolism in desert

homoiotherms. *Comparative Biochemistry and Physiology,* 26: 337–343.

McNab, B., & Morrison, P. (1963). Body temperature and metabolism in subspecies of

Peromyscus from arid and mesic environments. *Ecological Monographs,* 33: 63–82.

Melvin, R., & Andrews, M. (2009). Torpor induction in mammals: Recent discoveries fueling

new ideas. *Trends in Endocrinology and Metabolism,* 20: 490–498.

Mi, H., Huang, X., Muruganujan, A., Tang, H., Mills, C., Kang, D., & Thomas, P. (2017).

PANTHER version 11: expanded annotation data from Gene Ontology and Reactome

pathways, and data analysis tool enhancements. *Nucleic Acids Research,* 45: F183-D189.

Millar, J. (1989). Reproduction and development. In Advances in the Study of *Peromyscus*

(Rodentia) (eds Kirkland GI Jr, Layne JN, pp. 169–232). Lubbock, Texas: University of Texas

Press.

Morhardt, J., & Hudson, J. (1966). Daily torpor induced in white-footed mice (*Peromyscus* spp.)

by starvation. *Nature,* 212: 1046–1047.

Moritz, C., Patton, J., Conroy, C., Parra, J., White, G., & Beissinger, S. (2008). Impact of a

century of climate change on small-mammal communities in Yosemite National Park, USA.

Science, 322: 261–264.

Morjan, C., & Rieseberg, L. (2004). How species evolve collectively: implications of gene flow

and selection for the spread of advantageous alleles. *Molecular Ecology,* 13: 1341–1356.

Mosin, A. (1984). On the energy fuel in voles during their starvation. *Comparative Biochemistry*
and Physiology, 77: 563–565.

Muhs, D., Reynolds, R., Been, J., & Skipp, G. (2003). Eolian sand transport pathways in the
southwestern United States: importance of the Colorado river and local sources. *Quaternary*
International, 104: 3–18.

Murray, G. G. R., Soares, A. E. R., Novak, B. J., Schaefer, N. K., Cahill, J. A., Baker, A. J., …
Shapiro, B. (2017) Natural selection shaped the rise and fall of passenger pigeon genomic
diversity. *Science* 17: 951-954.

Nadachowska-Brzyska, K., Burri, R., Linnéa, S., Hans, E. (2016). PSMC analysis of effective
population sizes in molecular ecology and its application to black-and-white *Ficedula*
flycatchers. *Molecular Ecology* 25: 1058-1072.

Narikiyo, T., Kitamura, K., Adachi, M., Miyoshi, T., Iwashita, K., Shiraishi, N., … Tomita, K.
(2002). Regulation of prostasin by aldosterone in the kidney. *The Journal of Clinical*
Investigation, 109: 401–408.

Natarajan, C., Hoffmann, F., Lanier, H., Wolf, C., Cheviron, Z., Spangler, M., … Storz, J. (2015).
Intraspecific polymorphism, interspecific divergence, and the origins of function-altering
mutations in deer mouse hemoglobin. *Molecular Biology and Evolution,* 32: 978–997.

Neme, R., & Tautz, D. (2016). Fast turnover of genome transcription across evolutionary time
exposes entire non-coding DNA to de novo gene emergence. *ELife,* 5: e09977.

Nielsen, R., Williamson, S., Kim, Y., Hubisz, M., Clark, A., & Bustamante, C. (2005). Genomic
scans for selective sweeps using SNP data. *Genome Research,* 15: 1566–1575.

Noy-Meir, I. (1973). Desert ecosystems: Environment and producers. *Annual Review of Ecology*
and Systematics, 4: 25–51.

O'Rawe, J., Ferson, S., & Lyon, G. (2015). Accounting for uncertainty in DNA sequencing data.
Trends in Genetics, 31: 61–66.

Osada, N., & Akashi, H. (2012). Mitochondrial-nuclear interactions accelerated compensatory

evolution: evidence from the primate cytochrome C oxidase complex. *Molecular Biology and*

Evolution, 29: 337.

Park, M., Wu, D., Park, T., Choi, C., Li, R.-K., Cheng, K., … Sweeney, G. (2013). APPL1

transgenic mice are protected from high-fat diet-induced cardiac dysfunction. *American*

Journal of Physiology: Endocrinology and Metabolism, 305: E795–E804.

Pavlik, B. M. (2008). The California deserts: An ecological rediscovery. Berkeley: University of

California Press.

Pergams, O., & Lacy, R. (2008). Rapid morphological and genetic change in Chicago-area

Peromyscus. Molecular Ecology, 17: 450–463.

Pierce, S., & Vogt, F. (1993). Winter acclimatization in *Peromyscus maniculatus gracilis P*

leucopus noveboracensis, and *P. l. leucopus. Journal of Mammalogy,* 74: 665–677.

Platt, II, R., Amman, B., Keith, M., Thompson, C., & Bradley, R. (2015). What is *Peromyscus*?

Evidence from nuclear and mitochondrial DNA sequences suggests the need for a new

classification. *Journal of Mammalogy,* 96: 708–719.

Porcelli, D., Butlin, R., Gaston, K., Joly, D., & Snook, R. (2015). The environmental genomics of

metazoan thermal adaptation. *Heredity,* 114: 502–514.

R Core Team. (2017). R: A language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing.

Riddle, B., Hafner, D., & Alexander, L. (2000). Phylogeography and systematics of the

Peromyscus eremicus species group and the historical biogeography of North American

warm regional deserts. *Molecular Phylogenetics and Evolution,* 17: 145–160.

Rieder, L., Savva, Y., Reyna, M., Chang, Y.-J., Dorsky, J., Rezaei, A., & Reenan, R. (2015).

Dynamic response of RNA editing to temperature in *Drosophila. BMC Biology,* 13: 1.

Robinson, J., Turner, D., Durand, N., Thorvaldsdottir, H., Mesirov, J., & Aiden, E. (2018).

Juicebox.js provides a cloud-based visualization system for Hi-C data. *Cell Systems,* 6: 256–

258.

Rosenblum, E., Parent, C., & Brandt, E. (2014). The molecular basis of phenotypic

convergence. *Annual Review of Ecology and Systematics,* 45: 203–226.

Rundle, H., Nagel, L., Boughman, J., & Schluter, D. (2000). Natural selection and parallel

speciation in sympatric sticklebacks. *Science,* 287: 306–308.

Schwimmer, H., & Haim, A. (2009). Physiological adaptations of small mammals to desert

ecosystems. *Integrative Zoology,* 4: 357–366.

Shen, L. (2016). GeneOverlap: R package for testing and visualizing gene overlaps. New York

City, New York: Ichan School of Medicine at Mount Sinai.

Sikes, R. S., & The Animal Care and Use Committee of the American Society of Mammalogists.

(2016). 2016 Guidelines of the American Society of Mammalogists for the use of wild

mammals in research and education. *Journal of Mammalogy,* 97: 663–688.

Silanikove, N. (1994). The struggle to maintain hydration and osmoregulation in animals

experiencing severe dehydration and rapid rehydration: the story of ruminants. *Experimental*

Physiology, 79: 281–300.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics,* 31: 32-10–3212.

Simpson, J., Wong, K., Jackman, S., Schein, J., Jones, S., & Birol, I. (2009). ABySS: a parallel

assembler for short read sequence data. *Genome Research,* 19: 1117–1123.

Skotte, L., Korneliussen, T., & Albrechtsen, A. (2013). Estimating individual admixture

proportions from next generation sequencing data. *Genetics,* 195: 693–702.

Smalec, B., Heider, T., Flynn, B., & O'Neill, R. (2019). A centromere satellite concomitant with

extensive karyotypic diversity across the *Peromyscus* genus defies predictions of molecular

drive. Chromosome Research: An International Journal on the Molecular, Supramolecular

and Evolutionary Aspects of Chromosome Biology. doi: 10.1007/s10577-019-09605-1

Smit, A., Hubley, R., & Green, P. (2013). RepeatMasker Open-4.0. Available at:

http://www.repeatmasker.org/

Smith, J., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical Research,*

23: 23–35.

Steppan, S., Adkins, R., & Anderson, J. (2004). Phylogeny and divergence-date estimates of

rapid radiations in Muroid rodents based on multiple nuclear genes. *Systematic Biology,* 53:

533–553.

Stockham, S., & Scott, M. (2008). Fundamental of veterinary clinical pathology. Ames, IA:

Wiley-Blackwell.

Storz, J., & Cheviron, Z. (2016). Functional genomic insights into regulatory mechanisms of

high-altitude adaptation. *Advances in Experimental Medicine and Biology,* 903: 113–128. doi:

10.1007/978-1-4988-7678-9_8.

Storz, J., & Kelly, J. (2008). Effects of spatially varying selection on nucleotide diversity and

linkage disequilibrium: insights from deer mouse globin genes. Genetics, 180: 367–379.

Storz, J. (2007). Hemoglobin function and physiological adaptation to hypoxia in high-altitude

mammals. *Journal of Mammalogy*, 88: 24–31.

Storz, J., Runck, A., Moriyama, H., Weber, R., & Fago, A. (2010). Genetic differences in

hemoglobin function between highland and lowland deer mice. *Journal of Experimental*

Biology, 213: 2565–2574.

Storz, J., Cheviron, Z., McClelland, G., & Scott, G. (2019). Evolution of physiological

performance capacities and environmental adaptation: insights from high-elevation deer mice

(*Peromyscus maniculatus*). *Journal of Mammalogy*, 100: 910–922.

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO summarizes and visualizes

long lists of gene ontology terms. *PloS One*, 6: e21800.

Taniuchi, S., Miyake, M., Tsugawa, K., Oyadormari, M., & Oyadomari, S. (2016). Integrated

stress response of vertebrates is regulated by four eIF2α kinases. *Scientific Reports,* 6:

32886.

Tigano, A., Colella, J., & MacManes, M. (2020). Comparative and population genomics

approaches reveal the basis of adaptation to deserts in a small rodent. *Molecular Ecology,*

29: 1300–1314.

Tigano, A. & Friesen, V. L. (2016). Genomics of local adaptation with gene flow. *Molecular*

Ecology, 25: 2144-2164.

Tingley, M., & Beissinger, S. (2013). Cryptic loss of montane avian richness and high

community turnover over 100 years. *Ecology,* 94: 598–609.

Urban, M. (2015). Accelerating extinction risk from climate change. *Science,* 348: 571–573.

Van Dam, M., & Matzke, N. (2016). Evaluating the influence of connectivity and distance on

biogeographical patterns in the south-western deserts of North America. *Journal of*

Biogeography, 43: 1514–1532.

Veal, R., & Caire, W. (1979). *Peromyscus eremicus. Mammalian Species*, 118: 1–6.

Velotta, J., Robertson, C., Schweizer, R., McClelland, G., & Cheviron, Z. (2020). Adaptive shifts

in gene regulation underlie a developmental delay in thermogenesis in high-altitude deer

mice. *Molecular Biology and Evolution*, msaa086. doi:

https://doi.org/10.1093/molbev/msaa086

Wagner, G., & Lynch, V. (2008). The gene regulatory logic of transcription factor evolution.

Trends in Ecology & Evolution, 23: 277–385.

Wallbank, R. W., Baxter, S. W., Pardo-Diaz, C., Hanly, J. J., Martin, S. H., Mallet, J.,

Dasmahapatra, K. K., Salazar, C., Joron, M., Nadeau, N., McMillan, W. O. (2016).

Evolutionary novelty in a butterfly wing pattern through enhancer shuffling. *PLoS biology*,

14(1): p.e.1002353.

Weigand, H., & Leese, F. (2018). Detecting signatures of positive selection in non-model

species using genomic data. *Zoological Journal of the Linnean Society,* 184: 528–583.

Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct

determination of diploid genome sequences. *Genome Research,* 27: 757–767.

Wichman, H., & Lynch, C. (1991). Genetic variation for seasonal adaptation in *Peromyscus*

leucopus: nonreciprocal breakdown in a population cross. *Journal of Heredity,* 82: 197–204.

Williams, T., & Kelley, C. (2010). Gnuplot 4.4: an interactive plotting program (Version 4.4).

Retrieved from http://gnuplot.sourceforge.net/

Williams, D. (1987). Generalized linear model diagnostics using the deviance and single case

deletions. *Applied Statistics,* 36: 181–191.

Xie, S., Yang, X., Wang, D., Zhu, F., Yang, N., Hou, Z., & Ning, Z. (2018). Thyroid

transcriptome analysis reveals different adaptive responses to cold environmental conditions

between two chicken breeds. PLOS ONE, 13: 2018.

TABLE AND FIGURE CAPTIONS

Figure 1. Geographic ranges of the three *Peromyscus* species examined in this study with major southwestern North American deserts denoted by diagonal hashing. *P. crinitus* range is in red, *P. eremicus* in blue, and *P. maniculatus* in yellow. Areas of sympatry denoted by color overlap: dark purple = yellow + red + blue and green = yellow + blue. Collection localities are labeled with white dots and include the Motte Rimrock Reserve (MOT), Elliot Chaparral Reserve (ELL), and Philip L. Boyd Deep Canyon Desert Research Center (DRDC).

Figure 2. Distributions of effective population size (N_e) through time for *P. crinitus* (red), *P. eremicus* (blue), and *P. maniculatus* (yellow) based on a generation time of 6 months (0.5 years) and a general mammalian mutation rate of 2.2×10^{-9} substitutions/site/year. Note that the *P. maniculatus* genome was sequenced from a captive individual and therefore does not reflect natural populations trends of this species. Composite likelihood ratio (CLR) scores for *P. crinitus* based on *Sweepfinder2* results. Values above the horizontal red line surpass the 99.9th percentile. The top five or fewer unique genes are labeled for each chromosome.

Figure 3. *REVIGO* plots of enriched functional groups for *P. crinitus* (top row) and *P. eremicus* (bottom row) based on functional annotation of the two nearest protein-coding genes to each site (dataset II) identified as the subject of a selective sweep. Darker colors indicate greater significance. MP = metabolic process, MB = membrane-bound.

Figure 4. Composite likelihood ratio (CLR) scores for *P. crinitus* based on *Sweepfinder2* results. Values above the horizontal red line surpass the 99.9th percentile. The top five or fewer unique genes are labeled for each chromosome.

Figure 5. Overlap in proximal gene names (top row) and enriched GO terms (bottom row) for datasets I (left column), II (center), and III (right). *indicates significant overlap between species

Table 1. Assembly stats, genome size, and global Tajima's D and pi (1 kbp windows) for each *Peromyscus* species.

Table 2. Significant sweep sites localized within protein-coding genes for each *Peromyscus* species. Species (Spp.), chromosome (Chr.) sweep position (Pos.), gene name, protein, general function (based on UniProt database: uniprot.org), and direction (Dir.) of gene transcription. Abbreviation definitions: thermogen./thermogenesis, neg./negative, pos./positive, reg./regulation, IGF/insulin-like growth factor, ISR/Integrated Stress Response. *no gene name alternative available.

Table 3. Functional annotation of proximal gene names and enriched GO (gene ontology) terms associated with significant selective sweeps and shared between desert-adapted *P. crinitus* and *P. eremicus*. * indicates gene names or GO terms also shared with *P. maniculatus*

TABLES

Species	N	Scaffold N50	Contig N50	Size (Gb)[a]	Size (Gb)[b]	Taj. D	π
P. crinitus	9	94,816,992	204,461	2.27	2.28	-0.69	0.005
P eremicus	26	119,957,392	76,024	2.45	2.54	-1.27	0.007
P. maniculatus	5	115,033,041	42,400	2.33	2.44	-1.62	0.012

[a] abyss-fac estimate
[b] assemblathon estimate

Table 2. Significant sweep sites localized within protein-coding genes for each *Peromyscus* species. Species (Spp.), chromosome (Chr.) sweep position (Pos.), gene name, protein, general function (based on UniProt database: uniprot.org), and direction (Dir.) of gene transcription. Abbreviation definitions: thermogen./thermogenesis, neg./negative, pos./positive, reg./regulation, IGF/insulin-like growth factor, ISR/Integrated Stress Response. *no gene name alternative available.

Spp.	Chr.	Pos.	Gene	Protein	General function	Dir.
P. maniculatus	4	145409180	ZNF217	Zinc finger protein 217	DNA-binding transcription factor, transcription regulation, zinc binding	-
	20	36260251	DHRS7B	Dehydrogenase/reductase SDR family member 7B	Oxidoreductase activity	-
P. eremicus	1	42451454	Ush1c	Harmonin	Mechanotransduction in cochlear hair cells	-
	8	67956	MEIOB	Meiosis-specific with OB domain-containing protein	Meiosis	+
P. crinitus	3	52113514	PRSS2	Trypsin-2	Calcium ion binding	+
			KLK13	Kallikrein-13	Protein processing, proteolysis, reg. of IGF	+
			PRSS8	Prostasin	Sodium balance	+
			KLK4	Kallikrein-4	Zinc ion binding, proteolysis	+
			PRTN3	Myeloblastin	Degrades collagen (I, III, IV), elastin, fibronectin, laminin, vitronectin; blood coagulation, immune response	+
			KLK14	Kallikrein-14	Varied (epidermis morphogenesis)	+
			TRYP_PIG*	Trypsin	Calcium ion binding, proteolysis	+
			KLK6	Kallikrein-6	Varied (collagen catabolism, tissue regen)	+
			CELA2A	Chymotrypsin-like elastase family member 2A	Cleavage and elastin hydrolase, proteolysis	+
	4	57673659	EIF2AK4	eIF-2-alpha kinase GCN2	Metabolic stress sensing protein kinase, role in ISR required for adaptation to amino acid starvation, protein synthesis repression	-
	6	66203934	Nes	Nestin	Brain, eye development (neg. reg. catalytic activity)	-
	9	23303147	DENND64	DENND64	Endocytic recycling pathway component	+
		23323150	DENND64	DENND64	Endocytic recycling pathway component	+
		43305800	Nynrin	NYNRIN	Nucleic acid binding	+
		22243007	Parg	Poly(ADP-ribose) glycohydrolase	Prevent detrimental accumulation of poly(ADP-ribose) upon prolonged replicative stress	-
			Parg	Poly(ADP-ribose) glycohydrolase	Prevent detrimental accumulation of poly(ADP-ribose) upon prolonged replicative stress	-
		22283012	NCOA4	Nuclear receptor coactivator 4	Androgen receptor (iron ion homeostasis)	-
		22303015	Oxnad1	Oxidoreductase NAD-binding domain-containing protein 1	Oxidoreductase activity	-
	18		APPL2	DCC-interacting protein 13-beta	Varied (cold acclimation, diet induced thermogen., glucose homeostasis, neg. reg. of insulin response/fatty acid oxidation/glucose import, pos. reg. of cold-induced thermogen.)	-
		450151	APPL1	DCC-interacting protein 13-alpha	Varied. (insulin receptor signaling pathway, pos. reg. of glucose	-

				import)	
		APPL2	DCC-interacting protein 13-beta	Varied (cold acclimation, diet induced thermogen., glucose homeostasis, neg. reg. of insulin response/fatty acid oxidation/ glucose import, pos. reg. of cold-induced thermogen.)	-
	460153	APPL1	DCC-interacting protein 13-alpha	Varied (insulin receptor signaling pathway, pos. reg. of glucose import)	-
23	28065942	Tctn1	Tectonic-1	Neural development	+
		Tctn1	Tectonic-1	Neural development	+

Table 3. Functional annotation of proximal gene names and enriched GO (gene ontology) terms associated with significant selective sweeps and shared between desert-adapted *P. crinitus* and *P. eremicus*. * indicates gene names or GO terms also shared with *P. maniculatus*

Data set	Dataset	Gene Name / GO term	Function	Protein/Class
Gene Names	I	none	-	-
	II	*BTD*	Hydrolase, biotin transport/metabolism	Biotinidase
		RL36	Ribosomal protein, translation	60S ribosomal protein L36
	III	*BTD*	Hydrolase, biotin transport/metabolism	Biotinidase
		ENV	Zn binding, virion attachment	Envelope glycoprotein
		H3X	DNA binidng, protein heterodimerization	Putative histone H3.X
		RL15	RNA binding, ribosome constituent, translation	60S ribosomal protein L15
		RL36	Ribosomal protein, translation	60S ribosomal protein L36
		RPC1	Zn/Mg binding, immune response	DNA-directed RNA polymerase III subunit RPC1
		RS2*	Ribosomal protein, enzyme binding	40S ribosomal protein S2
		RS26	mRNA binding, ribosome, translation	40S ribosomal protein S26
		TM38B	Rapid Ca2+ release, K+ channel, ossification	Trimeric intracellular cation channel type B
		TRY2	Ca2+ binding, collagen catabolism, proteolysis, cell growth	Trypsin-2
Enriched GO terms	I	none	-	-
	II	GO:0005622	intracellular	cellular component
		GO:0005840*	ribosome	cellular component
		GO:0022626*	cystolic ribosome	cellular component
		GO:0043226	organelle	cellular component
		GO:0043229	intracellular organelle	cellular component
		GO:0044391*	ribosomal subunit	cellular component
		GO:1990904*	ribonucleoprotein complex	cellular component
	III	GO:0003735*	structural constituent of ribosome	molecular function
		GO:0022626*	cystolic ribosome	cellular component
		GO:0022625*	cystolic large ribosomal subunit	cellular component
		GO:0044391*	ribosomal subunit	cellular component
		GO:0005840*	ribosome	cellular component
		GO:0015934	large ribosomal subunit	cellular component
		GO:1990904*	ribonucleoprotein complex	cellular component

FIGURES

Figure 1. Geographic ranges of the three *Peromyscus* species examined in this study with major southwestern North American deserts denoted by diagonal hashing. *P. crinitus* range is in red, *P. eremicus* in blue, and *P. maniculatus* in yellow. Areas of sympatry denoted by color overlap: dark purple = yellow + red + blue and green = yellow + blue. MOT = Motte Rimrock Reserve, ELL = Elliot Chaparral reserve, DRDC = Philip Boyd Deep Canyon Desert Research Center.

Figure 2. Distributions of effective population size (N_e) through time for *P. crinitus* (red), *P. eremicus* (blue), and *P. maniculatus* (yellow) based on a generation time of 6 months (0.5 years) and a general mammalian mutation rate of 2.2×10^{-9} substitutions/site/year. Note that the *P. maniculatus* genome was sequenced from a captive individual and therefore does not reflect natural populations trends of this species.

Figure 3. *REVIGO* plots of enriched functional groups for *P. crinitus* (top row) and *P. eremicus* (bottom row) based on functional annotation of the two nearest protein-coding genes to each site (dataset II) identified as the subject of a selective sweep. Darker colors indicate greater significance. MP = metabolic process, MB = membrane-bound.

Figure 4. Composite likelihood ratio (CLR) scores for *P. crinitus* based on *Sweepfinder2* results. Values above the horizontal red line surpass the 99.9th percentile. The top five or fewer unique genes are labeled for each chromosome.

Figure 5. Overlap in proximal gene names (top row) and enriched GO terms (bottom row) for datasets I (left column), II (center), and III (right). *indicates significant overlap between species