Multiple evolutionary pathways to achieve thermal adaptation in small mammals

Jocelyn P. Colella[1,2,†*], Anna Tigano[1,2], Olga Dudchenko[3,4,5], Arina D. Omer[3], Ruqayya Khan[3,4],

Ivan D. Bochkov[3,4], Erez L. Aiden[3,4,5,6,7], Matthew D. MacManes[1,2]

[1]University of New Hampshire, Molecular, Cellular, and Biomedical Sciences Department,

Durham, NH 03824, USA (JPC: Jocelyn.Colella@unh.edu; AT: Anna.Tigano@unh.edu; MDM:

Matthew.MacManes@unh.edu)

[2]Hubbard Genome Center, University of New Hampshire, Durham, NH 03824, USA

[3]The Center for Genome Architecture, Department of Molecular and Human Genetics,

Baylor College of Medicine, Houston, TX 77030, USA

[4]Department of Computer Science, Department of Computational and Applied Mathematics,

Rice University, Houston, TX 77030, USA

[5]Center for Theoretical and Biological Physics, Rice University, Houston, TX 77030, USA

[6]Shanghai Institute for Advanced Immunochemical Studies, ShanghaiTech University, Shanghai

201210, China

[7]School of Agriculture and Environment, University of Western Australia, Perth, WA 6009,

Australia

[†]University of Kansas Biodiversity Institute and Ecology and Evolutionary Biology Department,

Lawrence, KS 66045, USA

* Corresponding Author: Jocelyn.Colella@unh.edu

ABSTRACT

Rapid ecological radiations provide useful models for identifying instances of parallel evolution, which can highlight critical genomic architecture involved in shared adaptations. Thermoregulatory innovations have allowed deer mice of the genus *Peromyscus* to radiate throughout North America, exploiting extreme thermal environments from mountain tops to desert valleys, and positioning this taxon as a model for understanding thermal adaptation. We compare signatures of selective sweeps across population-level genomic resequencing data from two desert-adapted *Peromyscus* species (*P. crinitus* and *P. eremicus*) and a third, widely-distributed habitat generalist (*P. maniculatus*) to test for signatures of parallel evolution and identify shared genomic architecture involved in adaptation to hot deserts. We found limited evidence of parallel evolution. Instead, we identified divergent molecular mechanisms of adaptation to similar environments potentially tied to species-specific historical demography that may limit or enhance adaptive variation. We also identified numerous genes under selection in *P. crinitus* that are implicated in osmoregulation (*Trypsin, Prostasin*) and metabolic responses to desert life (*Kallikrein, eIF2-alpha kinase GCN2, APPL1/2*). Evidence of varied evolutionary routes to achieve the same phenotype suggest there may be many molecular trajectories for small mammals to accommodate anthropogenic climate change.

Key words: convergence, dehydration, desert, parallel evolution, *Peromyscus,* thermoregulation

INTRODUCTION

Increasing global temperatures and altered patterns of precipitation threaten biodiversity

worldwide (Moritz et al. 2008; Cahill et al. 2013; Urban 2015). Phenotypic plasticity enables an

immediate response to changing conditions but evolutionary change through adaptation will be

critical for the long-term survival of most species (Hoffman and Sgro 2011; Cahill et al. 2013).

Range shifts upward in elevation and latitude have been documented in a number of terrestrial

species and interpreted as a response to warming (Chen et al. 2011; Tingley and Beissinger

2013; Freeman et al. 2018); however, responses vary even among closely-related species or

populations (Hoffman and Willi 2008; Moritz et al. 2008). The physiological limits responsible for

organismal range shifts are in part governed by genetics, which can facilitate adaptation to

specific environmental conditions. Population genomic methods enable the identification of

genes and molecular pathways involved in local adaptation by scanning the genome for

signatures of selection (Bassham et al. 2018; Garcia-Elfring et al. 2019). For species that have

independently adapted to similar environments, parallel or convergent evolution can be inferred

if a greater number of genes or phenotypes share signatures of selection than would be

expected under a purely stochastic model of evolution (e.g., drift). Convergence typically implies

the evolution of similar adaptive responses independently among distantly related taxa in

response to similar environmental or ecological conditions; in turn, parallel evolution is defined

as the occurrence of similar adaptive changes in groups with common ancestry (Simpson 1961;

Wood et al. 2005). Convergent evolution is often presumed to be driven by different underlying

molecular mechanisms, whereas parallel evolution may be driven by similar mechanisms;

however, this generalization may not reflect reality (Arendt and Reznick 2008). Evidence of

parallel or convergent evolution can suggest a deterministic effect of selection and highlight

conserved genomic architecture involved in shared adaptive phenotypes (Rundle et al. 2000;

McDonald et al. 2009), while a lack of concerted evolution may identify novel evolutionary

strategies to achieve the same phenotypic result.

As a model taxon (Dewey and Dawson 2001; Bedford and Hoekstra 2015) inhabiting

varied environments throughout North America, deer mice in the genus *Peromyscus* are a

frequent and productive subject of classical adaptation studies (e.g., physiological, Storz 2007;

behavioral, Hu and Hoekstra 2017; genetic, Cheviron et al. 2012; Storz and Cheviron 2016;

Tigano et al. 2020). Physiological similarity of deer mice to lab mice (*Mus musculus*) further

broadens the implications of evolutionary and ecological investigations of *Peromyscus* by linking

relevant results to biomedical sciences. The genus *Peromyscus* (N = 67 species;

mammaldiversity.org) is hypothesized to be the product of a rapid ecological radiation across

North America, evident in their varied ecological niches and rich species diversity (Glazier 1980;

Riddle et al. 2000; Bradley et al. 2007; Platt et al. 2015; Lindsey 2020). Adaptive radiations are

useful natural experiments for identifying patterns of parallel or convergent evolution, or the lack

thereof. Short generation times and accelerated thermoregulatory evolution relative to other

mammals, among other adaptive responses, appear to have enabled *Peroymscus* rodents to

exploit extreme thermal environments, ranging from cold, high elevations (Pierce and Vogt

1993; Cheviron et al. 2012, 2014; Kaseloo et al. 2014; Garcia-Elfring et al. 2019) to arid, hot

deserts (Riddle et al. 2000; MacManes 2017; Tigano et al. 2020). Thermoregulation and

dehydration tolerance are complex physiological traits and there are several potential

evolutionary routes to achieve the same phenotypic outcome. Within this framework,

comparisons among divergent *Peromyscus* species adapted to similar environments may

highlight shared adaptive polymorphisms or disparate evolutionary paths central to achieving

the same phenotype (Cheviron et al. 2012; Ivy and Scott 2017; Hu and Hoekstra 2017; Storz et

al. 2019). In cold environments, endotherms rely on aerobic thermogenesis to maintain constant

internal body temperatures. Changes in both gene expression and the functional properties of

proteins in high-altitude adapted deer mice suggest that changes in multiple hierarchical

molecular pathways may be common in the evolution of complex physiological traits, such as

thermoregulation (Wichman and Lynch 1991; Storz 2007; Cheviron et al. 2012; Storz and

Cheviron 2016; Garcia-Elfring et al. 2019). Nonetheless, investigations of thermoregulation

among high-elevation species may be confounded by concurrent selection on hemoglobin

oxygen-binding affinity as a consequence of a reduction in the partial pressure of oxygen as

elevation increases substantially (Storz and Kelly 2008; Storz et al. 2010; Natarajan et al. 2015).

In hot environments, endotherms are challenged with balancing heat dissipation, energy

expenditure, and water retention (Anderson and Jetz 2005), resulting in a different suite of

behavioral, physiological, and molecular adaptations that enable survival (Schwimmer and Haim

2009; Degen 2012; Kordonowy et al. 2016), but may be confounded by acute or chronic

dehydration. Understanding the biochemical mechanisms that enable survival under extreme

environmental stress can provide important insights into the nature of physiological adaptation.

Rapid thermoregulatory and ecological diversification among *Peromyscus* species

(origin ~8 Mya, radiation ~5.71 Mya; Platt et al. 2015) positions these small rodents as models

for anticipating species responses to accelerated warming (Cahill et al. 2013). Desert specialist

phenotypes have evolved repeatedly during the course of the *Peromyscus* radiation, with each

species and populations therein subject to distinct histories of demographic variation and gene

flow. These idiosyncratic histories can have a direct impact on evolution, as effective population

sizes are inextricably linked to the efficacy of selection and maintenance of genetic diversity in

wild populations (Charlesworth 2009). Further, contemporary or historical gene flow may help or

hinder adaptive evolution through homogenization or adaptive introgression, respectively

(Coyne and Orr 2004; Morjan and Reiseberg 2004; Jones et al. 2018; Tigano and Friesen

2016). Native to the American West, the canyon mouse (*P. crinitus*, Fig 1.) is well adapted to

desert life. In the lab, *P. crinitus* can survive in the absence of exogenous water, with urine

concentration levels similar to that of desert-adapted kangaroo rats (*Dipodomys merriami*;

Abbott 1971; MacMillen 1972; MacMillen and Christopher 1975; MacMillien 1983), but without

equivalently specialized renal anatomy (Issaian et al. 2012). Canyon mice also exhibit a lower-

than-expected body temperature relative to their size and can enter environmentally-mediated

torpor in response to drought, food limitation, or low external temperatures (McNab 1968;

McNab and Morrison 1963; Morhardt and Hudson 1966; Johnson and Armstrong 1987), which

facilitates survival in highly-variable and extreme desert environments. These phenotypes

persist for multiple generations in the lab indicating they have a genomic basis (McNab and

Morrison 1968). Cactus mice (*P. eremicus*) are related to *P. crinitus* and the two species are

frequently sympatric. Cactus mice exhibit similar adaptations to desert environments, including

urine concentration, reduced water requirements, and environmentally-induced torpor (Veal and

Caire 1979). In contrast, the habitat generalist *P. maniculatus* (North American deer mouse) is

geographically-widespread, native to both cool, high-elevations and hot southwestern deserts.

Whole-genome assemblies are publicly available for both *P. eremicus* (Tigano et al. 2020) and

P. maniculatus (Harvard University, Howard Hughes Medical Institute), which positions these

species as ideal comparatives against *P. crinitus* to identify genes and regulatory regions

associated with desert adaptation. Without fossil evidence of divergence and subsequent

convergence between desert-adapted *Peromyscus* species, similar patterns of selection are

interpreted as evidence of parallel evolution.

Here, we investigate genomic signatures of selection in desert-adapted *P. crinitus*. We

contrast signatures of selective sweeps across three related *Peromyscus* species, two desert

specialists (*P. crinitus* and *P. eremicus*) and one habitat generalist (*P. maniculatus*), to test for

signatures of parallel evolution. We hypothesize that similar genes or functional pathways will

be under selection in both desert-adapted species and not under selection in *P. maniculatus*,

providing a signature of parallel evolution. Finally, we place selective sweep analyses into an

evolutionary framework to interpret the varied evolutionary trajectories available to small

mammals to respond to changing environmental conditions and to account for demographic and

gene flow events.

104   **METHODS**

De novo genome sequencing and assembly

Wild mice were handled and sampled in accordance with the University of New Hampshire and

University of California Berkeley's Institutional Animal Care and Use Committee (130902 and

R224-0310, respectively) and California Department of Fish and Wildlife (SC-008135) and the

American Society of Mammalogists best practices (Sikes and Animal Care and Use Committee

of the American Society of Mammalogists 2016).

For the assembly of the *P. crinitus* genome, DNA was extracted from a liver subsample

from a *P. crinitus* individual collected in 2009 from the Philip L. Boyd Deep Canyon Desert

Research Center in Apple Valley, California. To generate a high-quality, chromosome-length

genome assembly for this individual we extracted high-molecular-weight genomic DNA using a

Qiagen genomic tip kit (Qiagen, Inc, Hilden, Germany). A 10X Genomics linked-reads library

was prepared according to manufacturer protocol at Mount Sinai and sequenced to a depth of

70X on a HiSeq 4000 (Novogene, Sacramento, California, USA). 10X Genomics reads were *de*

novo assembled into contigs using *Supernova* 2.1.1 (Weisenfeld et al. 2017). To arrange the

scaffolds thus obtained in chromosomes, a Hi-C library for *P. crinitus* was constructed and

sequenced from primary fibroblasts from the T.C. Hsu Cryo-Zoo at the University of Texas MD

Anderson Cancer Center. The Hi-C data were aligned to the supernova assembly using *Juicer*

(Durand et al. 2016). Hi-C genome assembly was performed using the *3D-DNA* pipeline

(Dudchenko et al. 2017) and the output was reviewed using *Juicebox Assembly Tools*

(Dudchenko et al. 2018). The Hi-C data are available on

www.dnazoo.org/assemblies/Peromyscus_crinitus visualized using *Juicebox.js*, a cloud-based

visualization system for Hi-C data (Robinson et al. 2018).

Benchmarking Universal Single-Copy Orthologs (*BUSCO* v3, using the Mammalia odb9

database; Simão et al. 2015) and *OrthoFinder2* (Emms and Kelly 2015) were used to assess

genome quality and completeness. Genome sizes were estimated for each species using

abyss-fac (Simpson et al. 2009) and the *assemblathon_stats.pl* script available at:

https://github.com/ucdavis-bioinformatics/assemblathon2-analysis/. *RepeatMasker* v.4.0 (Smit

et al. 2015) was used to identify repetitive elements. The genome was annotated using the

software package *MAKER* (3.01.02; Campbell et al. 2014). Control files, protein, and transcript

data used for this process are available at https://github.com/macmanes-

lab/pecr_genome/tree/master/annotation. We used *Mashmap* (-f one-to-one --pi 90 -s 300000;

Jain et al. 2017, 2018) to assess and plot (generateDotPlot.pl) syntenic conservation between

P. crinitus and *P. maniculatus* genomes. *Peromyscus crinitus* chromosomes were renamed and

sorted using *seqtk* (github.com/lh3/seqtk) following the *P. maniculatus* chromosome naming.

For comparative genomic analyses, we generated low-coverage whole-genome

resequencing data for nine *P. crinitus* and five *P. maniculatus* individuals collected from syntopic

areas of southern California (Table S1). *Peromyscus crinitus* samples were collected from the

University of California (UC) Philip L. Boyd Deep Canyon Desert Research Center (DCDRC)

near Palm Desert, California, and *P. maniculatus* were collected further East from the UC Motte

Rimrock Reserve and Elliot Chaparral Reserves. In addition to these, we used publicly available

whole-genome resequencing data from 26 *P. eremicus* individuals, also collected from DCDRC

and Motte Rimrock Reserve and prepared and sequenced in parallel (Tigano et al. 2020). All

samples were collected in 2009, with the exception of eight *P. eremicus* samples which were

collected in 2018. Animals were collected live in Sherman traps and a 25 mg ear-clip was taken

from each individual and stored at -80ºC in 95% ethanol. Animals were sampled from arid areas

with average monthly temperatures between 9-40ºC and mean annual rainfall of 15-18 cm. The

Biotechnology Resource Center at Cornell University (Ithaca, NY, USA) prepared genomic

libraries using the Illumina Nextera Library Preparation kit (e.g., skim-seq). Libraries were

sequenced at Novogene (Sacramento, CA, USA) using 150 bp paired-end reads from one lane

on the Illumina NovaSeq S4 platform. *fastp* v. 1 (Chen et al. 2018) was used to assess read

quality and trim adapters. Sequences from all samples and all species were mapped with *BWA*

(Li and Durbin 2010) to the *P. crinitus* reference genome to enable comparative analyses,

duplicates removed with *samblaster v. 0.1.24* (Faust and Hall 2014), and alignments indexed

and sorted using *samtools v. 1.10* (Li et al. 2009).

Population Genomics

We used the software package *ANGSD v. 0.93* (Korneliussen et al. 2014) to call variants from

low-coverage population genomic data with high confidence with the general options: -

SNP_pval 1e-6, -minMapQ 20, -minQ 20, -setMinDepth 20, -minInd 20, -minMaf 0.01. ANGSD

was run across all species and again within each species, where we required a minimum of half

(*-minInd*) *P. crinitus* and *P. eremicus* samples and all *P. maniculatus* samples to meet

independent quantity (*-minMapQ*) and quality (*-minQ*) thresholds and sample representation for

each variable site in each species.

Differentiation among species was examined using a Principle Component Analysis

(PCA, see Supporting Materials; *ngsTools,* Fumagalli et al. 2014) and multidimensional scaling

(MDS) of principal components in *NGSadmix v. 33* (Skotte et al. 2013). MDS plots were

generated in *R* v.3.6.1 (R Core Team 2017) based on the covariance matrix. Cook's D was

used to identify MDS outliers, using the broken stick method to identify single samples with

undue influence (Cook and Weisberg 1984; Williams 1987). *NGSadmix* was used to fit genomic

data into K populations to parse species-level differences and provide a preliminary screen for

genomic admixture under a maximum-likelihood model. Nonetheless, expanded sample sizes,

including representatives from additional populations of each species are necessary to

thoroughly investigate patterns of population structure and introgression. We tested K = 1

through K = (N - 1), where N is the number of total individuals examined. *NGSadmix* was run for

all species combined and for each species independently. As an additional measure of

differentiation, we estimated weighted and unweighted global F_{ST} values for each species pair

using *realSFS* in *ANGSD.*

We used Pairwise sequential Markovian Coalescent (*PSMC v. 0.6.5-r67;* Li and Durbin

2011) to examine patterns of historical demography through time for each species. The original

reads used to generate the high-quality, chromosome-length assemblies for each species (*P.*

crinitus generated here; *P. eremicus*, SAMEA5799953, Tigano et al. 2020; *P. maniculatus*:

GCA_003704035.1, Harvard University) were mapped to their assembly reference to identify

heterozygous sites and indexed in *BWA*. *Samblaster* removed PCR duplicates and *picard*

(http://broadinstitute.github.io/picard/) added a read group to the resulting bam file and

generated a sequence dictionary (*CreateSequenceDictionary*) from the reference assembly.

Samtools was used to sort, index, and variants called (*mpileup*) for each species, with *bcftools*

v1.10.2 (*call*, Li et al. 2009) and *VCFtools v 0.1.16* (*vcf2fq*, Danecek et al. 2011). *PSMC*

distributions of effective population size (N_e) were estimated with 100 bootstrap replicates.

PSMC results were visualized through *gnuplot v. 5.2* (Williams and Kelley 2010), using perl

scripts available at github.com/lh3/psmc, and scaled by a generation time of 6 months (0.5 yr,

Millar, 1989; Pergams and Lacy 2008) and a general mammalian mutation rate of 2.2 x 10^{-9}

substitutions/site/year (Kumar and Subramanian 2002).

198   *Tests for selection & convergence*

To detect recent selective sweeps in low-coverage whole-genome data, we used *Sweepfinder2*

(DeGiorgio et al. 2016; Nielsen et al. 2005). *Sweepfinder2* was run on both variant and invariant

sites (Huber et al. 2016) for each species, excluding sex chromosomes. Sex-chromosomes

were excluded for three reasons: (1) sex chromosome evolution is both rapid and complex

relative to autosomes, (2) we had different sample sizes of each sex across species, and (3)

desert adaptations, the focus of this study, are unlikely to be sex-specific. We repeated

Sweepfinder2 analyses on *P. eremicus*, initially analyzed by Tigano et al. (2020), using an

improved annotation scheme based on *Peromyscus*-specific data rather than *Mus musculus*

genes. Allele frequencies were estimated in *ANGSD*, converted to allele counts, and the site

frequency spectrum (SFS) was estimated from autosomes only in *Sweepfinder2*. Sweeps were

estimated from the pre-computed SFS and the composite likelihood ratio (CLR) and alpha

values, indicating the strength of selection, were calculated every 10,000 sites. Per Tigano et al.

(2020), a 10 kb window size was selected as a trade-off between computational time and

resolution. CLR values above the 99.9[th] percentile of the empirical distribution for each species

were considered to be evolving under a model of natural selection, hereafter referred to as

significant sweep sites. Smaller sample sizes produce fewer bins in the SFS and a diminishing

number of rare alleles which may impact both the overall SFS and local estimate surrounding

testing sites; therefore, we explored the impact of sample sizes on *Sweepfinder2* results in the

Supporting Information.

For each species, mean Tajima's D was calculated across the entire genome in non-

overlapping windows of 10 kb and 1 kb in *ANGSD*. Nucleotide diversity (π) was also calculated

in 10 kb and 1 kb windows and corrected based on the number of sites genotyped (variant and

invariant) per window. Tajima's D and π are expected to be significantly reduced in regions

surrounding selective sweeps (Smith and Haigh 1974; Kim and Stephan 2002), therefore we

used a Mann-Whitney test ($p < 0.05$, after a Bonferroni correction for multiple tests) to measure

significant deviations from the global mean in 1 kb and 10 kb flanking regions surrounding

significant sweep sites and 27 candidate genes identified in previous studies (MacManes 2017;

Table S2). Candidate loci include aquaporins (N = 12), sodium-calcium exchangers (*SLC8a1*),

and *Cyp4* genes belonging to the Cytochrome P450 gene family (N = 14). We used custom

python scripts to functionally annotated (I) the closest gene to each significant sweep site, (II)

the nearest upstream and downstream gene, regardless of strand (sense/antisense), and (III)

the nearest upstream and downstream gene on each strand. Dataset I follows the general

assumption that proximity between a significant sweep site and a protein-coding gene suggests

interaction. Dataset II represents an extension of that model by encompassing the most

proximal gene in each direction. Because *Sweepfinder2* is performed on the consensus

sequence and our data is unphased, we do not have information indicating on which strand a

significant sweep site occurs. Therefore, dataset III encompasses strand-uncertainty by

including the two nearest genes to a significant sweep site on both strands. It should be noted

that the genes identified in smaller datasets (I, II) are nested within the larger datasets (II, III)

and by definition, the larger datasets include more noise which may dilute a signature of parallel

evolution, but may better capture the true signal of selection. Hence, it is important to critically

examine numerous hierarchical gene subsets. We tested genes from each dataset for functional

and pathway enrichment in Gene Ontology (GO) categories using *Panther v. 15.0* (Mi et al.

2017) and extracted GO terms for each enriched functional group. We used *Mus musculus* as a

reference and a Bonferroni correction for multiple tests ($p < 0.05$) to correct for false

discoveries. Enriched GO terms were summarized and visualized in *REVIGO* (Reduce and

Visualize Gene Ontology, Supek et al. 2011) implemented at:

http://revigo.irb.hr/index.jsp?error=expired. As a test for convergence, the overlap in the gene

names and enriched GO terms associated with significant selective sweeps was assessed for

each dataset. Overlap was visualized in the R package *VennDiagram* (Chen and Boutros 2011).

To test for convergence, we used a Fisher's Exact Test ($p < 0.05$) in the *GeneOverlap* package

(Shen 2016) in R to assess whether gene or enriched GO term overlap between species was

greater than expected based on the total number of genes in the genome.

To compare patterns of gene family expansion and contraction potentially involved in

adaptation within the genus *Peromyscus*, we analyzed 14 additional genomes, including ten

Peromyscus species and four near outgroup rodent species: *Microtus ochrogaster, Neotoma*

lepida, Sigmodon hispidus, and *Mus musculus* (Table S3)*.* To prevent bias driven by variable

assembly qualities, samples with < 70% complete mammalian BUSCOs were excluded from

downstream analyses, resulting in the final analysis of ten species. Groups of orthologous

sequences (orthogroups) were identified in *Orthofinder2*. Invariant orthogroups and groups that

varied by more than 25 genes across taxa (custom python script: ortho2cafe.py) were excluded.

Our rooted species tree, estimated in *Orthofinder2*, was used to calculate a single birth-death

parameter (lambda) and estimate changes in gene family size using *CAFE v.4.2.1* (Han et al.

2013). Results were summarized using the python script *cafetutorial_report_analysis.py*

available from the Hahn Lab: hahnlab.github.io/CAFÉ/manual.html.

RESULTS

Chromosome-length genome assembly for P. crinitus

Linked reads combined with Hi-C scaffolding produced a high-quality, chromosome-length

genome assembly for *P. crinitus*. Our assembly has a contig N50 of 137,026 bp and scaffold

N50 of 97,468,232 bp, with 24 chromosome-length scaffolds. The anchored sequences in the

three *Peromyscus* genome assemblies were as follows: *P. crinitus* genome ~2.27 Gb, *P.*

eremicus ~2.5 Gb, and *P. maniculatus* ~2.39 Gb (Table 1). Our assembly has high contiguity

and completeness and low redundancy, as demonstrated by the presence of 89.3% complete

BUSCOs, 0.9% of duplicates and 9.0% missing, excluding unplaced scaffolds. As anticipated

(Smalec et al. 2019), we found no significant variation in chromosome number or major

interchromosomal rearrangements between *P. crinitus* and *P. maniculatus* (Figure S4). We

annotated 17,265 total protein coding genes in the *P. crinitus* genome. Similar to other

Peromyscus species, LINES1 (long interspersed nuclear elements) and LTR (long terminal

repeats) elements comprised 22.7% of the repeats in the *P. crinitus* genome, with SINEs (short

interspersed nuclear elements) representing an additional 9.6% (Table S5). Although similar to

other *Peromyscus* species, *P. crinitus* has the greatest total repeat content (>37%; see Tigano

et al. 2020 Supplementary Table 2).

Population Genomics

MDS analysis parsed the three species into well-separated clusters and identified no outliers or

evidence of admixture (Fig. S6). Analysis of genetic structure identified all three species as a

single group (K = 1) with the highest likelihood. A three-population model neatly parsed the

three species, as expected (Fig. S7, Table S8). We found evidence of potential admixture in *P.*

crinitus with at least three individuals containing 11-27% ancestry from *P. eremicus* and

additional material from *P. maniculatus* (4-16%), although variable samples sizes may impact

assignment certainty and expanded sequencing of additional species and populations will be

required to identify the specific sources of introgressed material. Four *P. eremicus* individuals

had < 90% assignment probability to the *P. eremicus* species cluster, with a maximum of 15%

assignment to a different species cluster. Identification of admixture in both species is not

biased by differences in coverage, as low (2X), medium (8X), and high coverage (17X) samples

were found to be admixed at a < 90% assignment threshold. No *P. maniculatus* individuals were

identified as admixed. *PSMC* estimates of historical demography show greater variance and a

higher overall N_e for *P. crinitus* relative to *P. eremicus* (Fig. 2). Demographic estimates for *P.*

maniculatus are included as a reference but should be interpreted with caution as they are

based on a captive-bred individual and may not accurately reflect the demography of wild

populations.

Average Tajima's D (1 kb windows) was negative for all species and ranged from -0.69

to -1.61. *Peromyscus crinitus* had the lowest Tajima's D value and *P. maniculatus* the highest

(Fig. S9). Global pairwise F_{ST} was moderate between all species ranging from 0.20-0.27

(unweighted: 0.12-0.17). Mean global π (1 kb windows) was 0.005 (±0.005) for *P. crinitus,* 0.007

(± 0.007) for *P. eremicus*, and 0.012 (± 0.010) for *P. maniculatus* (Fig. S10).

307   *Selection & convergence*

Within *P. crinitus* we identified a total of 209 significant sweep sites (Table S11), with 104 sites

localized on chromosome 9 and 16 regions experiencing major selective sweeps (Fig. 3).

Despite the large size of chromosome 1, we found no significant sweep sites on this

chromosome for *P. crinitus*. We found 239 total significant sweep sites for *P. eremicus* (Table

S12), with 56 sites concentrated on chromosome 1. Finally, we identified a total of 213

significant sweep sites for *P. maniculatus* (Table S13), with 103 sites located on chromosome 4.

A significant selective sweep (CLR > 99.9%) affects the area surrounding the site tested

- including both protein coding and non-coding regions - and cannot identify the specific

nucleotides under selection. Under the assumption that proximity suggests interaction between

a functional region of the genome and a sweep site, we hierarchically examined protein coding

genes most proximal to each sweep site. On average the distance from a sweep site to the

nearest coding gene was 45 kbp in *P. crinitus* (range: 31 - 439,122 bp), and much greater for

both *P. maniculatus* (average: 152 kbp; range: 190 - 513,562 bp) and *P. eremicus* (average:

117 kbp; range: 38 - 1,423,027 bp), which may be partially explained by differences in assembly

quality. For both *P. eremicus* and *P. maniculatus*, only two significant sweep sites were

identified within protein-coding genes (*P. eremicus:* Meiosis-specific with OB domain-containing

protein [gene name: *MEIOB*], Harmonin [*Ush1c*]; *P. maniculatus:* Dehydrogenase/reductase

SDR family member 7B [*DRS7B*] and Zinc finger protein 217 [*ZN217*]). In contrast, for *P.

crinitus* 12 significant sweep sites fell within 19 distinct candidate loci, many of which code for

numerous alternatively spliced transcripts (Table 2). Among *P. crinitus* significant sweep sites

localized within coding sequences, we identified 19 enriched GO terms (3 Biological Process

[BP], 9 Molecular Function [MF], 7 Cellular Component [CC]), with functionality ranging from

'proteolysis' to 'hydrolase' activity (Fig. 4). Functional examination of those genes identifies

solute regulation as a key function, with genes pertaining to calcium (*Trypsin-2* [*PRSS2*]) and

zinc (*Kallikrein-4* [*KLK4*]) binding and sodium regulation (*Prostasin* [*PRSS8*]) identified as under

selection. Examination of the two nearest genes to each sweep site (dataset II: one upstream,

one downstream gene, regardless of strand) in *P. crinitus* identified 121 unique genes and 26

enriched GO (gene ontology) terms (8 Biological Processes [BP], 10 Molecular Functions [MF],

8 Cellular Component [CC]), with functionality pertaining to metabolism (e.g., 'protein metabolic

process', 'organonitrogen compound metabolic process', 'peptide metabolic process') and

ribosomes (Fig. 4; Tables S11-13). For *P. eremicus,* we identified 202 unique genes and 14

enriched GO terms (0 BP, 1 MF, 13 CC) associated with selective sweeps, with functionality

again centered around ribosomes. For dataset II, two genes and seven enriched GO terms were

shared between the two desert-adapted species and we found no relationship between genes

identified under selection between the two species. Functional enrichment of *P. eremicus* and *P.*

maniculatus across all datasets was limited to ribosomes (e.g., 'structural constituent of

ribosome', 'cytosolic ribosome', 'ribosomal subunit'; Fig. 4; Tables S14-16). In contrast,

functionality of enriched GO terms for *P. crinitus* centered on metabolic processes, including

protein breakdown, hydrolysis, and cellular functionality (e.g., 'organelle', 'intracellular',

'cytoplasm'; Fig. 4; Table S16), in addition to ribosomes. See Supporting Information for detailed

results for datasets I and III. *Peromyscus eremicus* and *P. maniculatus* shared significant

overlap (p < 0.05) in enriched GO terms across all hierarchical data subsets (I, II, III; Fig. 5).

Significant overlap of enriched GO terms was also detected between *P. crinitus* and both other

Peromyscus species for datasets II and III only, with zero overlap for dataset I (Fig. 5).

Significant overlap in the genes located near significant sweep sites among desert-adapted

Peromyscus was only detected in dataset III. Overall, *P. crinitus* consistently had greater

diversity of functional enrichment relative to the other two species and the GO terms and genes

involved in ribosomal functionality were frequently shared among all species.

Species tree estimates (Fig. S17) were consistent with previous phylogenetic

investigations (Bradley et al. 2007). *Peromyscus crinitus* and *P. eremicus* are related in our

species tree, however a number of intermediate taxa remain unsampled (e.g., *P. merriami*, *P.*

californicus). Among the species examined here, the desert adapted *crinitus-eremicus* clade is

related to a clade comprised of *P. leucopus, P. polionotus*, and *P. maniculatus,* with the

nasutus-attwateri clade most basal within *Peromyscus.* For the *Peromyscus* genus, we found

19,925 gene families that had experienced contractions, 502 expansions, and 12 families that

were evolving rapidly. However, we found no gene families experiencing significant expansions,

contractions, or rapid evolution within *Peromyscus* or below the genus level.

Counter expectations, Tajima's D and π surrounding significant selective sweep sites

were significantly higher than the global mean estimates for each species (Table S18). Only in

P. maniculatus did we detect a significant reduction in π surrounding significant sweep sites.

Mean Tajima's D surrounding *a priori* candidate loci were also significantly more positive in all

three species.

371   **DISCUSSION**

Continued and accelerating environmental change increases the exigency of accurately

anticipating species responses to anthropogenic climate change. Adaptive evolutionary

responses vary among species and populations, even when subjected to seemingly

synonymous environmental selective pressures (Bi et al. 2015; Garcia-Elfring et al. 2019), but

evidence of parallel or convergent evolution can highlight critical genomic architecture involved

in key adaptations. We broadly define convergence as the evolution of similar adaptive

responses among distantly related taxa in response to similar environmental or ecological

conditions and parallel evolution as the occurrence of similar adaptive changes in groups that

share recent common ancestry (Simpson 1961; Wood et al. 2005). We analyzed genome-wide

patterns of selective sweeps among three species of deer mice within the North American

genus *Peromyscus* to identify critical genomic architecture or alternative evolutionary pathways

to achieve desert adaptation. We hypothesized that desert specialists *P. crinitus* and *P.*

eremicus would share similar patterns of selective sweeps related to surviving high-

temperature, low-water environments indicative of parallel or convergent evolution. The species

examined here share a common ancestor and the two desert adapted taxa share an even more

recent ancestor (Fig. S17); however, there are number of unsampled species separating *P.*

crinitus from *P. eremicus* (e.g., *P. merriami*, *P. californicus*) and these two taxa are relatively

divergent (average Fst of 0.25; divergence > 1 Mya, Platt et al. 2015). Nonetheless, without

evidence of ancestral divergence followed by subsequent convergence (molecular or

phenotypic) we cannot disentangle convergence from parallel evolution within the context of this

investigation. Instead, we infer parallel evolution as the most parsimonious explanation for

coordinated selection among desert-adapted *Peromyscus* species. As such, both selection on

novel beneficial mutations (typically deemed convergence) and the coincident retention of

advantageous ancestral polymorphisms are interpreted as evidence in support of parallel

evolution, as we do not have ancestral state information to distinguish the former.

397   Overall, we rejected our hypothesis that desert adapted species share parallel

signatures of selective sweeps reflecting adaptation to similar environments. Instead, we

identified divergent molecular mechanisms of adaptation to desert environments, with *P. crinitus*

potentially responding primarily through genomic changes to protein coding genes and *P.*

eremicus through transcriptional regulation of gene expression. This result potentially

contradicts the generalization that convergent evolution is driven by different molecular

mechanisms and parallel evolution, by similar mechanisms (Arendt and Reznick 2008).

Molecular flexibility of thermoregulatory responses may have catalyzed the radiation of

Peromyscus in North America by enabling rapid exploitation of novel thermal environments.

Finally, the application of an evolutionary lens to the interpretation of genomic patterns of

selective sweeps, particularly one that integrates historical demography and gene flow, may

serve to inform which evolutionary mechanism (genomic vs. transcriptomic) will be most

efficacious to achieve similar adaptive phenotypes in other small mammals.

411   *Limited evidence of parallel evolution*

Identification of similar genes or functions under selection in different species adapted to similar

environments suggests a deterministic effect of natural selection (Rosenblum et al. 2014) and

provides evidence in support of parallel evolution. In contrast, we found limited evidence of

parallel evolution among desert-adapted *Peromyscus*. Few to no enriched GO terms overlapped

between desert-specialists (Fig. 5). Instead, GO terms relating to ribosomes (e.g., 'ribosome',

'ribosomal subunit', 'cystolic ribosome', etc.) broadly overlapped between all three *Peromyscus*

species examined, with the most significant overlap occurring between desert adapted *P.*

eremicus and generalist *P. maniculatus*. Although *P. maniculatus* are not desert specialists, the

individuals sequenced here were collected in arid regions of southern California; therefore, the

shared signature of selection on ribosomes across all three species may reflect broadly shared

adaptations to hot and dry conditions or relate to thermoregulatory plasticity among *Peromyscus*

rodents. We found few genes experiencing significant selective sweeps shared among

Peromyscus species, with only one significant relationship: 10 genes, among hundreds, were

shared between the two desert specialists (dataset III; Fig. 5) and were consistent with a

signature of parallel evolution. Nonetheless, many of the overlapping genes are directly related

to broad ribosomal functionality (e.g., *RL36, RS26, RL15, RS2*) also shared with *P. maniculatus*.

Further, selective sweeps are only one way to detect signatures of parallel evolution and this

hypothesis remains to be explored in greater detail using additional methods (Booker et al.

2017, Weigand and Leese 2018).

Although we found no significantly expanded or contracted gene families within the

genus *Peromyscus*, previous investigations of the entire Myodonta clade within Rodentia

identified multiple expanded or contracted gene families associated with ribosomes in *P.*

eremicus (Tigano et al. 2020). Ribosomes play a critical role in protein synthesis and

degradation. Cellular damage accumulates quickly in desert environments as a consequence of

increased thermal- and osmotic-stress (Lamitina et al. 2006; Burg et al. 2007). In response,

changes in gene expression modulate osmoregulation by removing and replacing damaged

proteins to prevent cell death (Lamitina et al. 2006). While ribosomes appear to be a target of

parallel evolution in desert-adapted *Peromyscus*, this genomic signature is not unique. Instead,

selection on ribosomal functionality may be convergent across many species adapted to distinct

thermal environments (metazoans; Porcelli et al. 2015). Ribosomes are evolutionarily linked to

the mitochondrial genomes of animals (Barreto and Burton 2012; Bar-Yaacov et al. 2012) and

accelerated mitochondrial evolution in animals has led to compensatory, rapid evolution of

ribosomal proteins (Osada and Akashi 2012; Barreto and Burton 2013; Bar-Yaacov et al. 2012).

Rapid mitochondrial diversification within *Peromyscus* (Riddle et al. 2000; Bradley et al. 2007;

Platt et al. 2015), coincident with the ecological radiation of this genus (Lindsey 2020), suggests

that equivalent, rapid selection on ribosomal proteins may be a key evolutionary innovation that

enabled Peromyscine rodents to successfully and quickly adapt to varied thermal environments.

Comparisons among additional desert- and non-desert-adapted *Peromyscus* species will be

necessary to test this hypothesis within an evolutionary framework. Although not unique to

desert-species, rapid ribosomal evolution may provide a common evolutionary mechanism to

respond to anthropogenic climate change.

Evaporative cooling through sweating, panting, or salivating increases water loss and

challenges osmoregulatory homeostasis while maintaining thermoregulation in a hot and dry

climate (McKinley et al. 2018). Thermal stress exacerbates dehydration by increasing

evaporative water loss and if untreated, can result in cognitive dysfunction, motor impairment,

and eventually death. In consequence, osmoregulatory mechanisms are often under selection in

extreme thermal environments (MacManes and Eisen 2014; Marra et al. 2014). Consistently,

40% of the 10 genes targeted by selective sweeps and shared between desert adapted

Peromyscus are involved in ion balance (Table 3). Proteins Trypsin-2 (*TRY2*) and Trimeric

intracellular cation channel type B (*TM38B*) are associated with selective sweeps in both desert-

adapted species and involved in calcium ion (Ca^{2+}) binding and release, respectively. DNA-

directed RNA polymerase III (*RPC1*) has also experienced a significant selective sweep in both

desert species and influences magnesium (Mg^{2+}) binding. Calcium and magnesium cations are

among those essential for osmoregulation (also, Na^+, K^+, Cl^-, HCO_3^-; Stockham and Scott 2008)

and parallel selection on these genes is consistent with the hypothesis that solute-carrier

proteins are essential to maintaining homeostasis in desert-specialized rodents (Marra et al.

2014; Kordonowy and MacManes 2017). Additional genes implicated in osmoregulation were

identified as experiencing selective sweeps only in *P. crinitus* (Table 2; Table S16). In addition

to those shared with *P. eremicus,* Prostatin (*PRSS8*), only under selection in *P. crinitus,* is

critically responsible for increasing the activity of epithelial sodium (Na^+) channels, which

mediate Na^+ reabsorption through the kidneys (Narikiyo et al. 2002). Two more genes

associated with Ca^{2+} regulation (Anionic Trypsin-2 [*PRSS2*] and Trypsin [*TRYP*]) and other

genes regulating zinc (*KLK4*) and iron (*NCOA4*) were also identified as targets of selective

sweeps exclusively in *P. crinitus.*

477   *Metabolic tuning: proteins-for-water or lipids-for-torpor?*

Hot deserts experience dramatic fluctuations in both food and water availability that challenge

species survival (Noy-Meir 1973; Silanikove 1994). Mammals accommodate high temperatures

by increasing their body temperature, up to a point, and cold temperatures by aerobic

thermogenesis or metabolic suppression and the initiation of torpor or hibernation (Levesque et

al. 2016). When resources are scarce, metabolism relies exclusively on endogenous nutrients;

carbohydrates (e.g., sugars, glucose) are consumed immediately, then lipids, and eventually,

proteins. Protein oxidation has a low energy return relative to lipid catabolism (Bar and Volkoff

2012), but yields five times more metabolic water (Jenni and Jenni-Eiermann 1998; Gerson and

Guglielmo 2011a, b; McCue et al. 2017). In a low-water, desert environment an early shift to

protein catabolism during periods of resource limitation may represent an important source of

water for desert species (e.g., protein-for-water hypothesis; Mosin 1984; Jenni and Jenni-

Eiermann, 1998; Gerson and Guglielmo, 2011a, b), as demonstrated for migrating birds.

Consistent with this hypothesis, we identified numerous candidate genes that experienced

selective sweeps in *P. crinitus* and that are involved in the detection of metabolic-stress and

shifts in metabolic fuel consumption. For example, the gene eIF-2-alpha kinase GCN2 (*E2AK4*),

which is responsible for sensing metabolic stress in mammals and required for adaptation to

amino acid starvation, experienced the strongest selective sweep on chromosome 4 in *P.*

crinitus and (Fig. 3; Harding et al. 2003; Baker et al. 2012; Taniuchi et al. 2016). Numerous

candidate genes involved in oxidation (Oxidoreductase NAD-binding domain-containing protein

1 [*Oxnad1*]), fat catabolism (Kallikrein-6 [*KLK6*]), protein processing (Kallikrein-13 [*KLK13*]), and

proteolysis (Kallikrein [*KLK4*, *KLK13*]*,* Trypsin [*PRSS2, TRYP, TRY2*]*,* Chymotrypsin-like

elastase family member 2A [*CELA2A*]) were significantly enriched in *P. crinitus*, with proteolysis

as the most enriched functional group (Fig. 4; Table S16), potentially supporting the protein-for-

water hypothesis.

For desert species, including both desert specialists examined here (Morhardt and

Hudson 1966; MacMillen 1983), heat- and drought-induced torpor enable long duration, low

energy survival. Lipid acquisition and storage are critical to the initiation and maintenance of

torpor (Buck et al. 2002; Melvin and Andrews 2009). Significant weight loss in experimentally-

dehydrated *P. eremicus* and enhanced thermogenic performance of high-altitude-adapted deer

mice have been associated with enhanced lipid metabolism (Cheviron et al. 2012; Kordonowy et

al. 2016). At high-latitudes, increased lipid oxidation enables aerobic thermogenesis, but in hot

deserts lipids may represent a valuable energy source in a food-scarce environment (e.g.,

lipids-for-torpor hypothesis).

Metabolic processes were enriched in *P. crinitus*, but not *P. eremicus.* Two additional

candidate genes DCC-interacting protein 13-alpha and -beta (*APPL1*, *APPL2*), experienced a

significant selective sweep in *P. crinitus* and are important in glucose regulation, insulin

response, and fatty acid oxidation, reinforcing the hypothesis that enhanced lipid oxidation may

be critical to thermoregulatory responses. Laboratory manipulations of *APPL1* demonstrate

protection against high-fat diet-induced cardiomyopathy in rodents (Park et al. 2013) and *APPL2*

is responsible for dietary regulation, cold-induced thermogenesis, and cold acclimation

(uniprot.org). Together, these genes play a role in both obesity and dietary regulation. Both

APPL genes are associated with obesity and non-alcoholic fatty liver disease and their sweep

signature in *P. crinitus* may have relevant connections to biomedical research that remain to be

explored (Jiang et al. 2012; Barbieri et al. 2013). Physiological tests will be essential to

determine whether desert-adapted deer mice prioritize proteins or fats during periods of

resource limitation (e.g., lipids-for-torpor) or extreme dehydration (e.g., protein-for-water

hypothesis).

Molecular rewiring of metabolic processes in response to environmental conditions has

been documented in a number of species (e.g., mammals, Cheviron et al. 2012, Velotta et al.

2020; birds, Xie et al. 2018; fruit flies, Mallard et al. 2018). Nonetheless, expression changes

can also impact metabolism (Cheviron et al. 2012; Storz and Cheviron 2016). The capacity for

rapid adaptation to distinct thermal environments through multiple evolutionary mechanisms,

combined with thermoregulatory behavioral fine-tuning (e.g., nocturnality, aestivation, food

caching, burrowing, dietary shifts), suggests there may be many more evolutionary responses

available for small mammals to accommodate changing climates than previously realized.

Therein, metabolism and metabolic plasticity represents a fundamental phenotype for

anticipating species survival under altered climate scenarios.

Different evolutionary trajectories, same result

There are multiple evolutionary pathways to achieve environmental adaptation, most notably

through genomic changes in protein coding genes or transcriptional regulation of gene

expression. Although patterns of gene expression remain to be explored for *P. crinitus*, our

comparative genomic investigation suggests alternative evolutionary strategies for each desert-

adapted *Peromyscus* species shaped by their demographic histories: *P. crinitus* primarily

through changes in protein-coding genes and *P. eremicus* primarily through transcriptional

regulation.

Diverse functional enrichment of the *P. crinitus* genome (Fig. 4), spanning metabolic and

osmoregulatory functions in addition to the general functional enrichment of ribosomes, points to

polygenic adaptation to desert life. Here, we have identified a number of candidate loci worthy of

detailed examination across populations and in a laboratory setting to understand the varied

influence of different loci in thermoregulation and dehydration tolerance. In contrast, evidence of

many significant sweep sites in the *P. eremicus* genome, generally located more distant from

protein coding genes, and with functional enrichment restricted exclusively to ribosomes,

suggests local adaptation in this species may have been driven more by selection on regulatory

or non-coding regions of the genome that disproportionately impact gene expression. Indeed,

transcriptomic investigations have identified significant expression changes implicated in

osmoregulation in *P. eremicus* (MacManes and Eisen 2014; Kordonowy and MacManes 2017)

and in thermoregulation in other *Peromyscus* species and rodents (Cheviron et al. 2012; Marra

et al. 2014; Storz and Cheviron 2016). Transcriptional regulation is a particularly useful

mechanism for environmental acclimation, as these changes are more transient relative to

genomic changes and can enhance phenotypic flexibility (Garrett and Rosenthal 2012; Rieder et

al. 2015; Liscovitch-Brauer et al. 2017). For example, transcription factors (TFs), which are a

common target of selection located outside of protein-coding genes, can coordinate the

expression of many genes, which allows multiple phenotypic changes to occur simultaneously

(Wagner and Lynch 2008). Reduced genomic variation is expected near selective sweeps and

can encompass tens to thousands of adjacent nucleotides depending on recombination and the

strength of selection (Fay and Wu 2000; Carlson et al. 2005), yet counter expectation, Tajima's

D and nucleotide diversity for regions flanking putative selective sweeps were significantly

higher than the global average for most comparisons (Table S18).

Placing the results of selective sweep analyses within an evolutionary framework is

critical to interpreting varied evolutionary responses to similar environments. The expansion of

North American deserts following the conclusion of the last glacial maximum (LGM, ~11 Kya;

Pavlik et al. 2008) constrains the evolutionary and adaptive timescales of contemporary desert

species. Assuming simultaneous colonization of southwestern deserts, the stable but low

effective population size of *P. eremicus* suggests two hypotheses: (I) selection has removed

variation in this species over time (Murray et al. 2017) or (II) *P. eremicus* historically harbored

less genetic variation for selection to act on, despite equivalent contemporary diversity relative

to *P. crinitus* (Table S18). As a consequence of demography, the evolution of *P. eremicus* is

likely to have been more impacted by genetic drift (Fig. 2; Allendorf 1986; Masel 2011), so we

expect genomic evolution to be slower in *P. eremicus* relative to *P. crinitus*, which historically

has a larger effective population size and broader pool of variation for selection to act upon.

Within this context, environmental adaptation could be achieved more efficiently for *P. eremicus*

through transcriptomic plasticity or changes in regulatory elements (Allendorf 1986; Neme and

Tautz 2016; Mallard et al. 2018). In contrast, the larger historical effective population size of *P.*

crinitus is more conducive to the maintenance of genetic diversity and rapid evolution of protein

coding sequences through mutational stochasticity, reduced impact of genetic drift, and

potentially, gene flow. *Peromyscus crinitus* experienced a historical demographic bottleneck

prior to the formation of North American deserts; Nevertheless, the recovered effective

population size of *P. crinitus* is much larger than *P. eremicus* and consistent with low levels of

detected admixture (Fig. S7, Table S8). Also consistent with a history of admixture, *P. crinitus*

has the least negative Tajima's D, while low nucleotide diversity may indicate that admixture

may not be recent. Repeated growth and contraction of rivers in the American Southwest during

Pleistocene glacial-interglacial cycles (0.7-0.01 Mya; Muhs et al. 2003; Van Dam and Matzke

2016) would have provided iterative opportunities for connectivity and introgression between

incompletely-isolated *Peromyscus* species. Historical hybridization between *P. crinitus* and one

or more other Peromyscine species, likely unsampled here, may have accelerated adaptation in

P. crinitus through the rapid influx of novel mutation combinations through adaptive

introgression, a hypothesis that warrants further investigation through expanded sampling and

tests of adaptive introgression. Low-coverage whole-genome resequencing is optimal for

comparative and population genomic investigations (O'Rawe et al. 2015; da Fonseca et al.

2016), but detailed analyses of historical introgression are limited and we look forward to testing

this hypothesis with expanded population sampling and increased sequencing depth. Overall,

incorporating an evolutionary perspective into the interpretation of selection patterns has

important implications for anticipating species responses to anthropogenic climate change, as

historical demography and gene flow, in addition to selection, are responsible for shaping

contemporary diversity.

605   *Conclusion*

Contrasting patterns of selective sweeps and evolutionary histories between different species

experiencing similar environmental pressures can provide powerful insights into the adaptive

potential of species. We used comparative and population genomic analyses of three

Peromyscus species to identify candidate loci that may underlie adaptations to desert

environments in North America that serve to inform future investigations focused on predicting

potential for adaptation and identifying the causes of warming-related population declines (Cahill

et al. 2013). The identification of numerous targets of selection within *P. crinitus* highlights

multiple molecular mechanisms (metabolic switching, osmoregulatory tuning) associated with

physiological responses to deserts that warrant further investigation. Even in species recently

adapted to similar environments we identified divergent evolutionary trajectories, with one

species accommodating desert conditions primarily through genomic changes within protein

coding genes and the other, through transcriptional regulation mediated by historical

demographic processes. Our approach demonstrates the importance of placing genomic

selection analyses into an evolutionary framework to anticipate evolutionary responses to

change.

AUTHOR CONTRIBUTIONS

JPC performed analyses and wrote the first version of the paper. AT collected whole genome resequencing data, designed the bioinformatic pipeline. MDM conceptualized the project, performed *de novo* genome assembly and annotation, and funded data generation. OD, ADO, RK, IB, and ELA generated and assembled Hi-C data for *P. crinitus* as part of the DNA Zoo consortium effort. All authors reviewed and edited the manuscript.

ACKNOWLEDGEMENTS

We thank A. S. Westbrook for computational support; the Premise computing cluster at the University of New Hampshire, where all analyses were conducted; the Biotechnology Resource Center at Cornell University for preparation of whole-genome resequencing libraries; Christopher Tracy for access to the Boyd Deep Canyon Reserve; Jim Patton for desert field expertise; Sen Pathak, Asha Multani, Richard Behringer for providing the fibroblast samples from the T.C. Hsu Cryo-Zoo at the University of Texas M.D. Anderson Cancer Center; DNA Zoo for generating Hi-C data; Pawsey Supercomputing Centre with funding from the Australian Government and the Government of Western Australia for computational support of the DNA Zoo effort and the Museum of Southwestern Biology at the University of New Mexico for loaned tissue materials. This work was funded by the National Institute of Health National Institute of General Medical Sciences to MDM (1R35GM128843).

DATA ACCESSIBILITY STATEMENT

The draft assembly data are housed on the European Nucleotide Archive (ENA) under project ID PRJEB33592. The Hi-C data is available on SRA (SRX7041777, SRX7041776, SRX7041773) under the DNA Zoo project accession PRJNA512907. The *P. crinitus* genome assembly is available at https://www.dnazoo.org/assemblies/Peromyscus_crinitus. Whole-genome resequencing data for *P. crinitus* are available on ENA under project ID PRJEB35488.

Custom python scripts and other bash scripts used in analysis are available at:

https://github.com/jpcolella/Peromyscus_crinitus.

LITERATURE CITED

Abbott, K. (1971). Water economy of the canyon mouse, *Peromyscus crinitus stephensi*.
Comparative Biochemistry and Physiology, 38A, 37–52.
Allendorf, F. (1986). Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology,*
5(2), 181–190.
Anderson, K., & Jetz, W. (2005). The broad-scale ecology of energy expenditure of
endotherms. *Ecology Letters, 8*(3), 310–318.
Arendt, J., & Reznick, D. (2008). Convergence and parallelism reconsidered: what have we
learned about the genetics of adaptation? *Trends in Ecology & Evolution, 23*(1), 25–32.
Baker, B., Nargund, A., Sun, T., & Haynes, C. (2012). Protective coupling of mitochondrial
function and protein synthesis via the eIF2α kinase GCN-2. *PLoS Genetics, 8*(6), e1002760.
Bar, N., & Volkoff, H. (2012). Adaptation of the physiological, endocrine and digestive system
functions to prolonged food deprivation in fish. In Comparative physiology of fasting,
starvation, and food limitation (pp. 69–90). New York: Springer.
Bar-Yaacov, D., Blumberg, A., & Mishmar, D. (2012). Mitochondrial-nuclear co-evolution and its
effects on OXPHOS activity and regulation. *Biochimica et Biophsyica Acta (BBA)-Gene*
Regulatory Mechanisms, 1819(9–10), 1107–1111.
Barbieri, M., Esposito, A., Angellotti, E., Rizzo, M., Marfella, R., & Paolisso, G. (2013).
Association of genetic variation in adaptor protein APPL1/APPL2 loci with non-alcoholic fatty
liver disease. *PLOS ONE, 8*(8), e71391.
Barreto, F., & Burton, R. (2013). Evidence for compensatory evolution of ribosomal proteins in
response to rapid divergence of mitochondrial rRNA. *Molecular Biology and Evolution, 30*(2),
310–314.
Bassham, S., Catchen, J., Lescak, E., von Hippel, F., & Cresko, W. (2018). Repeated selection
of alternatively adapted haplotypes creates sweeping genomic remodeling in stickleback.
Genetics, 209(3), 921–939.
Bedford, M., & Hoekstra, H. (2015). The natural history of model organisms: *Peromyscus* mice
as a model for studying natural variation. *Elife, 4*, e06813.
Bi, K., Linderoth, T., Singhal, S., Vanderpool, D., Patton, J., Nielsen, R., … Good, J. (2015).
Temporal genomic contrasts reveal rapid evolutionary responses in an alpine mammal during
recent climate change. *PLoS Genetics, 15*(5), e1008119.
Booker, T., Jackson, B., & Keightley, P. (2017). Detecting positive selection in the genome.
BMC Biology, 15(98). doi: https://doi.org/10.1186/s12915-017-0434-y
Bradley, R., Durish, N., Rogers, D., Miller, J., Engstrom, M., & Kilpatrick, C. (2007). Toward a
molecular phylogeny for *Peromyscus*: Evidence from mitochondrial cytochrome-b
sequences. *Journal of Mammalogy, 88*(5), 1146–1159.
Buck, M., Squire, T., & Andrews, M. (2002). Coordinate expression of PDK4 gene: a means of
regulating fuel selection in a hibernating mammal. *Physiological Genomics, 8*(1), 5–13.
Burg, M., Ferraris, J., & Dmitrieva, N. (2007). Cellular response to hyperosmotic stresses.
Physiological Reviews, 87(4), 1441–1474.
Cahill, A., Aiello-Lammens, M., Fisher-Reid, M., Hua, X., Karanewsky, C., Yeong Ryu, H., …
Wiens, J. (2013). How does climate change cause extinction? *Proceedings of the Royal*
Society B: Biological Sciences, 280(1750), 20121890.
Campbell, M., Holt, C., Moore, B., & Yandell, M. (2014). Genome annotation and curation using
MAKER and MAKER-P. *Current Protocols in Bioinformatics, 48*(1), 4–11.

Carlson, C., Thomas, D., Eberle, M., Swanson, J., Livingston, R., Rieder, M., & Nickerson, D.
(2005). Genomic regions exhibiting positive selection identified from dense genotype data.
Genome Research, 15(11), 1553–1565.
Charlesworth, B. (2009). Effective population size and patterns of molecular evolution and
variation. *Nature Reviews Genetics, 10*(3), 195–205.
Chen, H., & Boutros, P. (2011). VennDiagram: a package for the generation of highly-
customizable Venn and Euler diagrams in R. *BMC Bioinformatics, 12*(1), 35
Chen, I., Hill, J., Ohlemueller, R., Roy, D., & Thomas, C. (2011). Rapid range shifts of species
associated with high levels of climate warming. *Science, 333*(6045), 1024–1026.
Chen, S., Zhou, Y., Chen, Y., & Gu, H. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics, 34*(17), i884–i890.
Cheviron, Z., Bachman, G., Connaty, A., McClelland, G., & Storz, J. (2012). Regulatory
changes contribute to the adaptive enhancement of thermogenic capacity in high-altitude
deer mice. *Proceedings of the National Academy of Sciences, 109*(22), 8635–8640.
Cheviron, Z., Connaty, A., McClelland, G., & Storz, J. (2014). Functional genomics of
adaptation to hypoxic cold-stress in high-altitude deer mice: transcriptomic plasticity and
thermogenic performance. *Evolution, 68*(1), 48–62.
Cook, R., & Weisberg, S. (1984). Residuals and influence in Regression. Wiley.
Coyne, J. A., & Orr, H. A. (2004). Speciation. Sunderland, MA: Sinauer Associates.
da Fonseca, R. R., Albrechtsen, A., Themudo, G., Ramos-Madrigal, J., Sibbesen, J., Maretty,
L., … Pereira, R. (2016). Next-generation biology: sequencing and data analysis approaches
for non-model organisms. *Marine Genomics, 30*, 3–13. doi: 10.1016/j.margen.2016.04.012
Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … 1000
Genome Project Data Process Subgroup. (2011). The variant call format and VCFtools.
Bioinformatics, 27(15), 2156–2158.
Degen, A. (2012). Ecophysiology of small desert mammals. Springer Science & Business
Media.
DeGiorgio, M., Huber, C., Hubisz, M., Hellmann, I., & Nielsen, R. (2016). SweepFinder2:
increased sensitivity, robustness and flexibility. *Bioinformatics, 32*(12), 1895–1897.
Dewey, M., & Dawson, W. (2001). Deer mice: "the *Drosophila* of North American mammalogy.
Genesis, 29(3), 105–109.
Dudchenko, O., Batra, S., Omer, A., Nyquist, S., Hoeger, M., Durand, N., … Aiden, E. (2017).
De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length
scaffolds. *Science, 356*(6333), 92–95.
Dudchenko, O., Shamim, M., Batra, S., Durand, N., Musial, N., Mostofa, R., … Aiden, E. (2018).
The Juicebox Assembly Tools module facilitates de novo assembly of mammalian genomes
with chromosome-length scaffolds for under $1000. *BioRxiv.* doi:
https://doi.org/10.1101/254797
Durand, N., Shamim, M., Machol, I., Rao, S., Huntley, M., Lander, E., & Aiden, E. (2016). Juicer
provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell
Systematics, 3*(1), 95–98.
Emms, D. M., & Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biology, 16*(1),
157.
Faust, G., & Hall, I. (2014). SAMBLASTER: fast duplicate marking and structural variant read
extraction. *Bioinformatics, 30*(17), 2503–2505.
Fay, J., & Wu, C.-I. (2000). Hitchhiking under positive Darwinian selection. *Genetics, 155*(3),
1405–1413.
Freeman, B., Lee-Yaw, J., Sunday, J., & Hargreaves, A. (2018). Expanding, shifting and
shrinking: The impact of global warming on species' elevational distributions. *Global Ecology
and Biogeography, 27*(11), 1268–1276.

Fumagalli, M., Vieira, F., Linderoth, T., & Nielsen, R. (2014). ngsTools: methods for population genetics analyses from next-generation sequencing data. *Bioinformatics, 30*(10), 1486–1487.

Garcia-Elfring, A., Barrett, R., & Millien, V. (2019). Genomic signatures of selection along a climatic gradient in the northern range margin of the white-footed mouse (*Peromyscus leucopus*). *Journal of Heredity, 110*(6), 684–695.

Garrett, S., & Rosenthal, J. (2012). RNA editing underlies temperature adaptation in K+ channels from polar octopuses. *Science, 335*(6070), 848–851.

Gerson, A., & Guglielmo, C. (2011a). Flight at low ambient humidity increases protein catabolism in migratory birds. *Science, 333*(6048), 1434–1436.

Gerson, A., & Guglielmo, C. (2011b). House sparrows (*Passer domesticus*) increase protein catabolism in response to water restriction. *American Journal of Physiology, 300*(4), R925–R930.

Glazier, D. (1980). Ecological shifts and the evolution of geographically restricted species of North American *Peromyscus* (mice). *Journal of Biogeography, 7*(1), 63–83.

Han, M., Thomas, G., Lugo-Martinez, J., & Hahn, M. (2013). Estimating gene gain and loss rates in the presence of error in genome assembly and annotation using CAFE 3. *Molecular Biology and Evolution, 30*(8), 1987-1997.

Harding, H., Zhang, Y., Zeng, H., Novoa, I., Lu, P., Calfon, M., … Ron, D. (2003). An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Molecular Cell, 11*(3), 619–633.

Hoffmann, A., & Sgro, C. (2011). Climate change and evolutionary adaptation. *Nature, 470*(7335), 479–485.

Hoffmann, A., & Willi, Y. (2008). Detecting genetic responses to environmental change. Nature Reviews: *Genetics, 9*(6)*,* 421–432.

Hu, C., & Hoekstra, H. (2017). *Peromyscus* burrowing: A model system for behavioral evolution. *Seminars in Cell and Developmental Biology, 61*, 107–114.

Huber, C., DeGiorgio, M., Hellmann, I., & Nielsen, R. (2016). Detecting recent selective sweeps while controlling for mutation rate and background selection. *Molecular Ecology, 25*(1), 142–156.

Issaian, T., Urity, V., Dantzler, W., & Pannabecker, T. (2012). Architecture of vasa recta in the renal inner medulla of the desert rodent *Dipodomys merriami*: potential impact on the urine concentrating mechanism. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology, 303*(7), R748–R756.

Ivy, C., & Scott, G. (2017). Control of breathing and ventilatory acclimatization to hypoxia in deer mice native to high altitudes. *Acta Physiologica, 221*(4), 266–282.

Jain, C., Koren, S., Dilthey, A., Phillippy, A. M., Aluru, S. (2018) A fast adaptive algorithm for computing whole-genome homology maps. *Bioinformatics, 34*(17), i748-i756.

Jain, C., Dilthey A, Koren, S, Aluru, S., Phillippy, A. M. (2017) A fast approximate algorithm for mapping long reads to large reference databases. In: Sahinalp S. (eds) *Research in Computational Molecular Biology*. RECOMB 2017. Lecture Notes in Computer Science, 10229. Springer, Cham.

Jenni, L., & Jenni-Eiermann, S. (1998). Fuel supply and metabolic constraints in migrating birds. *Journal of Avian Biology, 29*(4), 521–528.

Jiang, S., Fang, Q., Yu, W., Zhang, R., Hu, C., Dong, K., … Jia, W. (2012). Genetic variations in APPL2 are associated with overweight and obesity in a Chinese population with normal glucose tolerance. *BMC Medical Genetics, 13*(1), 22.

Johnson, D., & Armstrong, D. (1987). *Peromyscus crinitus. Mammalian Species, 287*, 1–8.

Jones, M., Mills, L., Alves, P., Callahan, C., Alves, J., Lafferty, D., … Good, J. (2018). Adaptive introgression underlies polymorphic seasonal camouflage in snowshoe hares. *Science, 360*(6395), 1355–1358.

Kaseloo, P., Crowell, M., & Heideman, P. (2014). Heritable variation in reaction norms of
metabolism and activity across temperatures in a wild-derived population of white-footed
mice (*Peromyscus leucopus*). *Journal of Comparative Physiology B, 184*(4), 525–534.
Kim, Y., & Stephan, W. (2002). Detecting a local signature of genetic hitchhiking along a
recombining chromosome. *Genetics, 160*(2), 765–777.
Kordonowy, L., & MacManes, M. (2017). Characterizing the reproductive transcriptomic
correlates of acute dehydration in males in the desert-adapted rodent, *Peromyscus eremicus.*
BMC Genomics, 18(1), 473.
Kordonowy, L., Lombardo, K., Green, H., Dawson, M., Bolton, E., LaCourse, S., & MacManes,
M. (2016). Physiological and biochemical changes associated with acute experimental
dehydration in the desert adapted mouse, *Peromyscus eremicus. Physiological Reports,*
5(6), e13218.
Korneliussen, T., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation
Sequencing Data. *Bioinformatics, 15*(1), 356.
Kumar, S., & Subramanian, S. (2002). Mutation rates in mammalian genomes. Proceedings of
the National Academy of Sciences, 99(2), 803–808.
Lamitina, T., Haung, C., & Strange, K. (2006). Genome-wide RNAi screening identifies protein
damage as a regulator of osmoprotective gene expression. *Proceedings of the National*
Academy of Sciences of the United States of America, 103(32), 12173–12178.
Levesque, D., Nowack, J., & Stawski, C. (2016). Modelling mammalian energetics: the
heterothermy problem. *Climate Change Responses, 3*(1), 7.
Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics, 26*(5), 589–595.
Li, Heng, & Durbin, R. (2011). Inference of human population history from whole genome
sequence of a single individual. *Nature, 475*(7357), 493–496.
Li, H., Handsaker, B., Wysoker, A., Fennel, T., Ruan, J., Homer, N., … 1000 Genome Project
Data Process Subgroup. (2009). The sequence alignment/map format and SAMtools.
Bioinformatics, 25(16), 2078–2079.
Lindsey, L. (2020). Utilizing genomic applications to examine patterns of diversification in
deermice (Rodentia: Cricetidae: *Peromyscus*). Texas Tech Dissertation.
Liscovitch-Brauer, N., Alon, S., Porath, H., Elstein, B., Unger, R., Ziv, T., … Eisenberg, E.
(2017). Trade-off between transcriptome plasticity and genome evolution in Cephalopods.
Cell, 169(2), 191–202.
MacManes, M., & Eisen, M. (2014). Characterization of the transcriptome, nucleotide sequence
polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus.*
PeerJ, 2, e642.
MacManes, M. (2017). Severe acute dehydration in a desert rodent elicits a transcriptional
response that effectively prevents kidney injury. American Journal of Physiology. *Renal*
Physiology, 313(2), F262–F272.
MacMillen, R., & Christopher, E. (1975). The water relations of two populations of non-captive
desert rodents. In Environmental physiology of desert organisms ((N. F. Hadley, ed.), pp.
117–137). Stroudsburg, Pennsylvania: Dowden, Hutchinson, and Ross.
MacMillen, R. (1972). Water economy of nocturnal desert rodents. *Symposia of the Zoological*
Society of London, 31, 147–174.
MacMillen, R. (1983). Water regulation in *Peromyscus. Journal of Mammalogy, 64*(1), 38–47.
Mallard, F., Nolte, V., Tobler, R., Kapun, M., & Schlötterer, C. (2018). A simple genetic basis of
adaptation to a novel thermal environment results in complex metabolic rewiring in
Drosophila. Genome Biology, 19(119), 1-15. doi: 10.1186/s13059-018-1503-4
Marra, N., Romero, A., & DeWoody, J. (2014). Natural selection and the genetic basis of
osmoregulation in heteromyid rodents as revealed by RNA-seq. *Molecular Ecology, 23*(11),
2699–2711.

Masel, J. (2011). Genetic drift. *Current Biology, 21*(20), R837–R838.

McCue, M., Sandoval, J., Beltran, J., & Gerson, A. (2017). Dehydration causes increased
reliance on protein oxidation in mice: a test of the protein-for-water hypothesis in a mammal.
Physiological and Biochemical Zoology, 90(3), 359–369.

McDonald, M., Gehrig, S., Meintjes, P., Zhang, X.-X., & Rainey, P. (2009). Adaptive divergence
in experimental populations of *Pseudomonas fluorescens*. IV. Genetic constraints guide
evolutionary trajectories in a parallel adaptive radiation. *Genetics, 183*(3), 1041–1053.

McKinley, M., Martelli, D., Pennington, G., Trevaks, D., & McAllen, R. (2018). Integrating
competing demands of osmoregulatory and thermoregulatory homeostasis. *Physiology,*
33(3), 170–181.

McNab, B. (1963). The influence of fat deposits on the basal rate of metabolism in desert
homoiotherms. *Comparative Biochemistry and Physiology, 26*, 337–343.

McNab, B., & Morrison, P. (1963). Body temperature and metabolism in subspecies of
Peromyscus from arid and mesic environments. *Ecological Monographs, 33*(1), 63–82.

Melvin, R., & Andrews, M. (2009). Torpor induction in mammals: Recent discoveries fueling
new ideas. *Trends in Endocrinology and Metabolism, 20*(10), 490–498.

Mi, H., Huang, X., Muruganujan, A., Tang, H., Mills, C., Kang, D., & Thomas, P. (2017).
PANTHER version 11: expanded annotation data from Gene Ontology and Reactome
pathways, and data analysis tool enhancements. *Nucleic Acids Research, 45*(D1), F183-
D189.

Millar, J. (1989). Reproduction and development. In Advances in the Study of *Peromyscus*
(Rodentia) (eds Kirkland Gl Jr, Layne JN, pp. 169–232). Lubbock, Texas: University of Texas
Press.

Morhardt, J., & Hudson, J. (1966). Daily torpor induced in white-footed mice (*Peromyscus* spp.)
by starvation. *Nature, 212*(5066), 1046–1047.

Moritz, C., Patton, J., Conroy, C., Parra, J., White, G., & Beissinger, S. (2008). Impact of a
century of climate change on small-mammal communities in Yosemite National Park, USA.
Science, 322(5899), 261–264.

Morjan, C., & Rieseberg, L. (2004). How species evolve collectively: implications of gene flow
and selection for the spread of advantageous alleles. *Molecular Ecology, 13*(6), 1341–1356.

Mosin, A. (1984). On the energy fuel in voles during their starvation. *Comparative Biochemistry*
and Physiology, 77(3), 563–565.

Muhs, D., Reynolds, R., Been, J., & Skipp, G. (2003). Eolian sand transport pathways in the
southwestern United States: importance of the Colorado river and local sources. *Quaternary*
International, 104(1), 3–18.

Murray, G. G. R., Soares, A. E. R., Novak, B. J., Schaefer, N. K., Cahill, J. A., Baker, A. J., …
Shapiro, B. (2017) Natural selection shaped the rise and fall of passenger pigeon genomic
diversity. *Science 17*(358), 951-954.

Narikiyo, T., Kitamura, K., Adachi, M., Miyoshi, T., Iwashita, K., Shiraishi, N., … Tomita, K.
(2002). Regulation of prostasin by aldosterone in the kidney. *The Journal of Clinical*
Investigation, 109(3), 401–408.

Natarajan, C., Hoffmann, F., Lanier, H., Wolf, C., Cheviron, Z., Spangler, M., … Storz, J. (2015).
Intraspecific polymorphism, interspecific divergence, and the origins of function-altering
mutations in deer mouse hemoglobin. *Molecular Biology and Evolution, 32*(4), 978–997.

Neme, R., & Tautz, D. (2016). Fast turnover of genome transcription across evolutionary time
exposes entire non-coding DNA to de novo gene emergence. *ELife, 5*, e09977.

Nielsen, R., Williamson, S., Kim, Y., Hubisz, M., Clark, A., & Bustamante, C. (2005). Genomic
scans for selective sweeps using SNP data. *Genome Research, 15*(11), 1566–1575.

Noy-Meir, I. (1973). Desert ecosystems: Environment and producers. *Annual Review of Ecology*
and Systematics, 4(1), 25–51.

O'Rawe, J., Ferson, S., & Lyon, G. (2015). Accounting for uncertainty in DNA sequencing data.
Trends in Genetics, 31(2), 61–66.
Osada, N., & Akashi, H. (2012). Mitochondrial-nuclear interactions accelerated compensatory
evolution: evidence from the primate cytochrome C oxidase complex. *Molecular Biology and*
Evolution, 29(1), 337.
Park, M., Wu, D., Park, T., Choi, C., Li, R.-K., Cheng, K., … Sweeney, G. (2013). APPL1
transgenic mice are protected from high-fat diet-induced cardiac dysfunction. *American*
Journal of Physiology: Endocrinology and Metabolism, 305(7), E795–E804.
Pavlik, B. M. (2008). The California deserts: An ecological rediscovery. Berkeley: University of
California Press.
Pergams, O., & Lacy, R. (2008). Rapid morphological and genetic change in Chicago-area
Peromyscus. Molecular Ecology, 17(1), 450–463.
Pierce, S., & Vogt, F. (1993). Winter acclimatization in *Peromyscus maniculatus gracilis P*
leucopus noveboracensis, and *P. l. leucopus. Journal of Mammalogy, 74*(3), 665–677.
Platt, II, R., Amman, B., Keith, M., Thompson, C., & Bradley, R. (2015). What is *Peromyscus*?
Evidence from nuclear and mitochondrial DNA sequences suggests the need for a new
classification. *Journal of Mammalogy, 96*(4), 708–719.
Porcelli, D., Butlin, R., Gaston, K., Joly, D., & Snook, R. (2015). The environmental genomics of
metazoan thermal adaptation. *Heredity, 114*(5), 502–514.
R Core Team. (2017). R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.
Riddle, B., Hafner, D., & Alexander, L. (2000). Phylogeography and systematics of the
Peromyscus eremicus species group and the historical biogeography of North American
warm regional deserts. *Molecular Phylogenetics and Evolution, 17*(2), 145–160.
Rieder, L., Savva, Y., Reyna, M., Chang, Y.-J., Dorsky, J., Rezaei, A., & Reenan, R. (2015).
Dynamic response of RNA editing to temperature in *Drosophila. BMC Biology, 13*(1), 1.
Robinson, J., Turner, D., Durand, N., Thorvaldsdottir, H., Mesirov, J., & Aiden, E. (2018).
Juicebox.js provides a cloud-based visualization system for Hi-C data. *Cell Systems, 6*(2),
256–258.
Rosenblum, E., Parent, C., & Brandt, E. (2014). The molecular basis of phenotypic
convergence. *Annual Review of Ecology and Systematics, 45*, 203–226.
Rundle, H., Nagel, L., Boughman, J., & Schluter, D. (2000). Natural selection and parallel
speciation in sympatric sticklebacks. *Science, 287*(5451), 306–308.
Schwimmer, H., & Haim, A. (2009). Physiological adaptations of small mammals to desert
ecosystems. *Integrative Zoology, 4*(4), 357–366.
Shen, L. (2016). GeneOverlap: R package for testing and visualizing gene overlaps. New York
City, New York: Ichan School of Medicine at Mount Sinai.
Sikes, R. S., & The Animal Care and Use Committee of the American Society of Mammalogists.
(2016). 2016 Guidelines of the American Society of Mammalogists for the use of wild
mammals in research and education. *Journal of Mammalogy, 97*(3), 663–688.
Silanikove, N. (1994). The struggle to maintain hydration and osmoregulation in animals
experiencing severe dehydration and rapid rehydration: the story of ruminants. *Experimental*
Physiology, 79(3), 281–300.
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics, 31*(19), 32-10–3212.
Simpson, G. (1961). Principles of Animal Taxonomy. New York: Columbia University Press.
Simpson, J., Wong, K., Jackman, S., Schein, J., Jones, S., & Birol, I. (2009). ABySS: a parallel
assembler for short read sequence data. *Genome Research, 19*(6), 1117–1123.
Skotte, L., Korneliussen, T., & Albrechtsen, A. (2013). Estimating individual admixture
proportions from next generation sequencing data. *Genetics, 195*(3), 693–702.

Smalec, B., Heider, T., Flynn, B., & O'Neill, R. (2019). A centromere satellite concomitant with
extensive karyotypic diversity across the *Peromyscus* genus defies predictions of molecular
drive. Chromosome Research: An International Journal on the Molecular, Supramolecular
and Evolutionary Aspects of Chromosome Biology. doi: 10.1007/s10577-019-09605-1
Smit, A., Hubley, R., & Green, P. (2013). RepeatMasker Open-4.0. Available at:
http://www.repeatmasker.org/
Smith, J., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical Research,*
23(1), 23–35.
Steppan, S., Adkins, R., & Anderson, J. (2004). Phylogeny and divergence-date estimates of
rapid radiations in Muroid rodents based on multiple nuclear genes. *Systematic Biology,*
53(4), 533–553.
Stockham, S., & Scott, M. (2008). Fundamental of veterinary clinical pathology. Ames, IA:
Wiley-Blackwell.
Storz, J., & Cheviron, Z. (2016). Functional genomic insights into regulatory mechanisms of
high-altitude adaptation. *Advances in Experimental Medicine and Biology, 903*, 113–128. doi:
10.1007/978-1-4988-7678-9_8.
Storz, J., & Kelly, J. (2008). Effects of spatially varying selection on nucleotide diversity and
linkage disequilibrium: insights from deer mouse globin genes. Genetics, 180, 367–379.
Storz, J. (2007). Hemoglobin function and physiological adaptation to hypoxia in high-altitude
mammals. *Journal of Mammalogy, 88*(1), 24–31.
Storz, J., Runck, A., Moriyama, H., Weber, R., & Fago, A. (2010). Genetic differences in
hemoglobin function between highland and lowland deer mice. *Journal of Experimental*
Biology, 213(15), 2565–2574.
Storz, J., Cheviron, Z., McClelland, G., & Scott, G. (2019). Evolution of physiological
performance capacities and environmental adaptation: insights from high-elevation deer mice
(*Peromyscus maniculatus*). *Journal of Mammalogy, 100*(3), 910–922.
Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO summarizes and visualizes
long lists of gene ontology terms. *PloS One, 6*(7), e21800.
Taniuchi, S., Miyake, M., Tsugawa, K., Oyadormari, M., & Oyadomari, S. (2016). Integrated
stress response of vertebrates is regulated by four eIF2α kinases. *Scientific Reports, 6*,
32886.
Tigano, A., Colella, J., & MacManes, M. (2020). Comparative and population genomics
approaches reveal the basis of adaptation to deserts in a small rodent. *Molecular Ecology,*
29(7), 1300–1314.
Tigano, A. & Friesen, V. L. (2016). Genomics of local adaptation with gene flow. *Molecular*
Ecology, 25, 2144-2164.
Tingley, M., & Beissinger, S. (2013). Cryptic loss of montane avian richness and high
community turnover over 100 years. *Ecology, 94*(3), 598–609.
Urban, M. (2015). Accelerating extinction risk from climate change. *Science, 348*(6234), 571–
573.
Van Dam, M., & Matzke, N. (2016). Evaluating the influence of connectivity and distance on
biogeographical patterns in the south-western deserts of North America. *Journal of*
Biogeography, 43(8), 1514–1532.
Veal, R., & Caire, W. (1979). *Peromyscus eremicus. Mammalian Species, 118*(8), 1–6.
Velotta, J., Robertson, C., Schweizer, R., McClelland, G., & Cheviron, Z. (2020). Adaptive shifts
in gene regulation underlie a developmental delay in thermogenesis in high-altitude deer
mice. *Molecular Biology and Evolution*, msaa086. doi:
https://doi.org/10.1093/molbev/msaa086
Wagner, G., & Lynch, V. (2008). The gene regulatory logic of transcription factor evolution.
Trends in Ecology & Evolution, 23(7), 277–385.

Weigand, H., & Leese, F. (2018). Detecting signatures of positive selection in non-model
species using genomic data. *Zoological Journal of the Linnean Society, 184*(2), 528–583.

Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct
determination of diploid genome sequences. *Genome Research, 27*(5), 757–767.

Wichman, H., & Lynch, C. (1991). Genetic variation for seasonal adaptation in *Peromyscus*
leucopus: nonreciprocal breakdown in a population cross. *Journal of Heredity, 82*(3), 197–
204.

Williams, T., & Kelley, C. (2010). Gnuplot 4.4: an interactive plotting program (Version 4.4).
Retrieved from http://gnuplot.sourceforge.net/

Williams, D. (1987). Generalized linear model diagnostics using the deviance and single case
deletions. *Applied Statistics, 36*(2), 181–191.

Wood, T., Burke, J., & Rieseberg, L. (2005). Parallel genotypic adaptation: when evolution
repeats itself. *Genetica, 123*(1–2), 157–170.

Xie, S., Yang, X., Wang, D., Zhu, F., Yang, N., Hou, Z., & Ning, Z. (2018). Thyroid
transcriptome analysis reveals different adaptive responses to cold environmental conditions
between two chicken breeds. PLOS ONE, 13(1), 2018.

TABLES

Table 1. Assembly stats, genome size, and global Tajima's D and pi (1 kb windows) for each

Peromyscus species.

Species	N	Scaffold N50	Contig N50	Size (Gb)[a]	Size (Gb)[b]	Taj. D	π
P. crinitus	9	94,816,992	204,461	2.27	2.28	-0.69	0.005
P eremicus	26	119,957,392	76,024	2.45	2.54	-1.27	0.007
P. maniculatus	5	115,033,041	42,400	2.33	2.44	-1.62	0.012

[a] abyss-fac estimate
[b] assemblathon estimate

Table 2. Significant sweep sites localized within protein-coding genes for each *Peromyscus* species. Species (Spp.), chromosome (Chr.) sweep position (Pos.), gene name, protein, general function (based on UniProt database: uniprot.org), and direction (Dir.) of gene transcription. Abbreviation definitions: thermogen./thermogenesis, neg./negative, pos./positive, reg./regulation, IGF/insulin-like growth factor, ISR/Integrated Stress Response. *no gene name alternative available.

Spp.	Chr.	Pos.	Gene	Protein	General function	Dir.
P. maniculatus	4	145409180	ZNF217	Zinc finger protein 217	DNA-binding transcription factor, transcription regulation, zinc binding	-
	20	36260251	DHRS7B	Dehydrogenase/reductase SDR family member 7B	Oxidoreductase activity	-
P. eremicus	1	42451454	Ush1c	Harmonin	Mechanotransduction in cochlear hair cells	-
	8	67956	MEIOB	Meiosis-specific with OB domain-containing protein	Meiosis	+
P. crinitus	3	52113514	PRSS2	Trypsin-2	Calcium ion binding	+
			KLK13	Kallikrein-13	Protein processing, proteolysis, reg. of IGF	+
			PRSS8	Prostasin	Sodium balance	+
			KLK4	Kallikrein-4	Zinc ion binding, proteolysis	+
			PRTN3	Myeloblastin	Degrades collagen (I, III, IV), elastin, fibronectin, laminin, vitronectin; blood coagulation, immune response	+
			KLK14	Kallikrein-14	Varied (epidermis morphogenesis)	+
			TRYP_PIG*	Trypsin	Calcium ion binding, proteolysis	+
			KLK6	Kallikrein-6	Varied (collagen catabolism, tissue regen)	+
			CELA2A	Chymotrypsin-like elastase family member 2A	Cleavage and elastin hydrolase, proteolysis	+
	4	57673659	EIF2AK4	eIF-2-alpha kinase GCN2	Metabolic stress sensing protein kinase, role in ISR required for adaptation to amino acid starvation, protein synthesis repression	-
	6	66203934	Nes	Nestin	Brain, eye development (neg. reg. catalytic activity)	-
	9	23303147	DENND64	DENND64	Endocytic recycling pathway component	+
		23323150	DENND64	DENND64	Endocytic recycling pathway component	+
		43305800	Nynrin	NYNRIN	Nucleic acid binding	+
		22243007	Parg	Poly(ADP-ribose) glycohydrolase	Prevent detrimental accumulation of poly(ADP-ribose) upon prolonged replicative stress	-
			Parg	Poly(ADP-ribose) glycohydrolase	Prevent detrimental accumulation of poly(ADP-ribose) upon prolonged replicative stress	-
		22283012	NCOA4	Nuclear receptor coactivator 4	Androgen receptor (iron ion homeostasis)	-
		22303015	Oxnad1	Oxidoreductase NAD-binding domain-containing protein 1	Oxidoreductase activity	-
	18		APPL2	DCC-interacting protein 13-beta	Varied (cold acclimation, diet induced thermogen., glucose homeostasis, neg. reg. of insulin response/fatty acid oxidation/glucose import, pos. reg. of cold-induced thermogen.)	-
		450151	APPL1	DCC-interacting protein 13-alpha	Varied. (insulin receptor signaling pathway, pos. reg. of glucose	-

				import)	
		APPL2	DCC-interacting protein 13-beta	Varied (cold acclimation, diet induced thermogen., glucose homeostasis, neg. reg. of insulin response/fatty acid oxidation/ glucose import, pos. reg. of cold-induced thermogen.)	-
	460153	APPL1	DCC-interacting protein 13-alpha	Varied (insulin receptor signaling pathway, pos. reg. of glucose import)	-
23	28065942	Tctn1	Tectonic-1	Neural development	+
		Tctn1	Tectonic-1	Neural development	+

Table 3. Functional annotation of proximal gene names and enriched GO (gene ontology) terms associated with significant selective sweeps and shared between desert-adapted *P. crinitus* and *P. eremicus*. * indicates genes or GO terms also shared with *P. maniculatus*

Data set	Dataset	Gene Name / GO term	Function	Protein/Class
Gene Names	I	none	-	-
	II	*BTD*	Hydrolase, biotin transport/metabolism	Biotinidase
		RL36	Ribosomal protein, translation	60S ribosomal protein L36
	III	*BTD*	Hydrolase, biotin transport/metabolism	Biotinidase
		ENV	Zn binding, virion attachment	Envelope glycoprotein
		H3X	DNA binidng, protein heterodimerization	Putative histone H3.X
		RL15	RNA binding, ribosome constituent, translation	60S ribosomal protein L15
		RL36	Ribosomal protein, translation	60S ribosomal protein L36
		RPC1	Zn/Mg binding, immune response	DNA-directed RNA polymerase III subunit RPC1
		RS2*	Ribosomal protein, enzyme binding	40S ribosomal protein S2
		RS26	mRNA binding, ribosome, translation	40S ribosomal protein S26
		TM38B	Rapid Ca2+ release, K+ channel, ossification	Trimeric intracellular cation channel type B
		TRY2	Ca2+ binding, collagen catabolism, proteolysis, cell growth	Trypsin-2
Enriched GO terms	I	none	-	-
	II	GO:0005622	intracellular	cellular component
		GO:0005840*	ribosome	cellular component
		GO:0022626*	cystolic ribosome	cellular component
		GO:0043226	organelle	cellular component
		GO:0043229	intracellular organelle	cellular component
		GO:0044391*	ribosomal subunit	cellular component
		GO:1990904*	ribonucleoprotein complex	cellular component
	III	GO:0003735*	structural constituent of ribosome	molecular function
		GO:0022626*	cystolic ribosome	cellular component
		GO:0022625*	cystolic large ribosomal subunit	cellular component
		GO:0044391*	ribosomal subunit	cellular component
		GO:0005840*	ribosome	cellular component
		GO:0015934	large ribosomal subunit	cellular component
		GO:1990904*	ribonucleoprotein complex	cellular component

FIGURES & FIGURE LEGENDS

Figure 1. Geographic ranges of the three *Peromyscus* species examined in this study with major southwestern North American deserts denoted by diagonal hashing. *P. crinitus* range is in red, *P. eremicus* in blue, and *P. maniculatus* in yellow. Areas of sympatry denoted by color overlap: dark purple = yellow + red + blue and green = yellow + blue.

Figure 2. Distributions of effective population size (N_e) through time for *P. crinitus* (red), *P. eremicus* (blue), and *P. maniculatus* (yellow) based on a generation time of 6 months (0.5 years) and a general mammalian mutation rate of 2.2×10^{-9} substitutions/site/year. Note that the *P. maniculatus* genome was sequenced from a captive individual and therefore does not reflect natural populations trends of this species.

Figure 3. *REVIGO* plots of enriched functional groups for *P. crinitus* (top row) and *P. eremicus* (bottom row) based on functional annotation of the two nearest protein-coding genes to each site (dataset II) identified as the subject of a selective sweep. Darker colors indicate greater significance. MP = metabolic process, MB = membrane-bound.

Figure 4. Composite likelihood ratio (CLR) scores for *P. crinitus* based on *Sweepfinder2* results. Values above the horizontal red line surpass the 99.9[th] percentile. The top five or fewer unique genes are labeled for each chromosome.

Figure 5. Overlap in proximal gene names (top row) and enriched GO terms (bottom row) for datasets I (left column), II (center), and III (right). *indicates significant overlap between species