1

Developmental models reveal the role of phenotypic plasticity in explaining genetic evolvability

Miguel Brun-Usan[1,2*], Alfredo Rago[1,2], Christoph Thies[1], Tobias Uller[2], Richard A. Watson[1]

1- Institute for life sciences / Electronics and computer sciences. University of Southampton (UK)

2-Department of Biology, Lund University, 22362, Sweden.

*- Corresponding author:

E-mail: miguel.brun-usan@biol.lu.se

Running title: Adaptive plasticity and genetic evolvability

Keywords: Phenotypic plasticity, Evolvability, Genotype-phenotype-map, Phenotype-first hypothesis

Abstract

Biological evolution exhibits an extraordinary capability to adapt organisms to their environments. The explanation for this often takes for granted that random genetic variation produces at least some beneficial phenotypic variation for natural selection to act on. Such genetic evolvability could itself be a product of evolution, but it is widely acknowledged that the immediate selective gains of evolvability are small on short timescales. So how do biological systems come to exhibit such extraordinary capacity to evolve? One suggestion is that adaptive phenotypic plasticity makes genetic evolution find adaptations faster. However, the need to explain the origin of adaptive plasticity puts genetic evolution back in the driving seat, and genetic evolvability remains unexplained. To better understand the interaction between plasticity and genetic evolvability, we simulate the evolution of phenotypes produced by gene-regulation network-based models of development. First, we show that the phenotypic variation resulting from genetic and environmental change are highly concordant. This is because phenotypic variation, regardless of its cause, occurs within the relatively specific space of possibilities allowed by development. Second, we show that selection for genetic evolvability results in the evolution of adaptive plasticity and *vice versa*. This linkage is essentially symmetric but, unlike genetic evolvability, the selective gains of plasticity are often substantial on short, including within-lifetime, timescales. Accordingly, we show that selection for phenotypic plasticity is, in general, the most efficient and results in the rapid evolution of high genetic evolvability. Thus, without overlooking the fact that adaptive plasticity is itself a product of genetic evolution, this shows how plasticity can influence adaptive evolution and helps explain the genetic evolvability observed in biological systems.

Introduction

Understanding how evolution works is not complete by understanding natural selection; we also need to understand the generation of the phenotypic variation that natural selection will act on (Hallgrimsson and Hall 2005; Houle et al. 2010). This is challenging since developmental processes often make non-directed genetic variation give rise to phenotypic variation that is highly structured, with some variants appearing more frequently than others (Alberch 1982; Salazar-Ciudad et al. 2003; Kavanagh et al. 2007). If this developmental bias was aligned with the adaptive demands imposed by local environments (i.e., if the mutationally more accessible phenotypes are also the more adaptive ones), adaptive evolution would be greatly facilitated. Such genetic evolvability could itself be a product of past evolution (Uller et al. 2018), but the idea that natural selection would be able to improve genetic evolvability is problematic because the immediate selective gains of evolvability are small on short timescales (Wagner and Altenberg, 1996; Watson 2020).

One suggestion is that the high genetic evolvability is acquired through the capability of organisms to rapidly adapt to their environment during their lifetime (Crispo 2007). However, while adaptive plasticity often appears to 'take the lead' in adaptive evolution (e.g., Radersma et al. 2020), the idea that adaptive plasticity explains rapid genetic evolution overlooks the need to explain the origination of the adaptive plasticity that supposedly 'came first'. In this paper, we seek to better understand whether phenotypic plasticity can help to explain genetic evolvability without overlooking the fact that adaptive plasticity is itself a product of genetic evolution (Levis and Pfennig 2016).

Our approach follows the observation that the phenotypic consequences of genetic variation and environmental variation are not expected to be independent because the consequences of any such variations are channelled by the same underlying developmental mechanisms (Cheverud 1982; Salazar-Ciudad 2006). From this it follows that, if selection for phenotypic plasticity alters the structure of developmental interactions, this will alter how phenotypes respond to genetic mutations and hence affect evolvability. Population genetic models demonstrate that selecting for plasticity can increase genetic variation along the dimensions of the phenotype that are plastic (Ancel and Fontana 2000; Hansen 2006; Draghi and Whitlock 2012; Furusawa and Kaneko 2018). This seems to support a role for plasticity in shaping genetic evolvability. However, given that the interdependence of plasticity and evolvability on development is essentially symmetric, the reverse may be equally possible. That is, if selection for genetic evolvability alters the structure of developmental interactions, this should also alter how phenotypes respond to environmental variation. It thus remains an open question whether genetic evolvability is predominantly shaped by plasticity or *vice versa* (Schwander and Leimar 2011; Levis and Pfennig 2016).

To address this question, we study the relationship between plasticity and evolvability by
representing these phenomena in a common framework where the phenotypic effects and adaptive
consequences of genetic and environmental variation can be compared. The phenotype distribution
that is generated by genetic variation can be represented as a genotype-phenotype (GP) map: an
idealized representation of development that assigns a phenotype to each genotype (Alberch 1982;
Kirschner and Gerhart 1998; Jimenez et al. 2015). Genetic evolvability is high when the GP map
enables genetic change to produce phenotypes that are suitable for adaptation. Analogous to the GP
map, plasticity can be understood as an Environment-Phenotype (EP) map (aka reaction norm) that
associates each environment with its corresponding phenotype (West-Eberhard 2003; Salazar-
Ciudad 2007; Li et al. 2018). Plasticity is adaptive when the EP map enables individuals to produce
phenotypes that fit the requirements of the environment in which it finds itself. However, the GP
and EP maps do not exhaust the sources of phenotypic variation. Development is also sensitive to
non-genetic and non-environmental initial conditions such as biochemical templates, resources and
nutrients that are provided by the parents (Newman and Muller 2000; Jablonka and Lamb 2005).
The association of such elements with phenotypic variation is often referred to as parental effects
(Badyaev and Uller 2009). Here we emphasize the analogous role of such initial conditions in
structuring the phenotype distribution by referring to it as the PP (for 'Parental-Phenotype') map: a
mapping that associates a phenotype with the parentally inherited initial conditions required to
produce it.

To model the potential interdependence of these three maps (GP, EP and PP), we use several
different and widely used models of development based on gene regulatory networks (GRN). This
approach means that we neither assume that the three maps are independent nor that they are
related; rather, these are hypotheses we can test. To do this we apply the three different forms of
variation (i.e. genetic, environmental, parental) to the same developmental system (GRN), and
compare the resulting phenotypic effects. In our approach, genetic changes alter the regulatory
connections of the network, environmental changes alter the gene expression levels, and parental
effects alter the initial gene expression levels of the network (see Methods). Together, these
variables define the GRN dynamics which, when implemented and iterated in the models, result in
measurable phenotypes.

In principle, it could be the case that any concordance between the three maps in this model
could be intrinsic to the properties of GRNs and arise without selection, or observed only in GRNs
subjected to particular selective conditions, or not observed at all. Likewise, it is not known whether
selection can, for example, change the GP map produced by a GRN without altering the EP or PP
maps, or more generally, whether the effects of selecting for one map has consequences for the
properties of the others. Lastly, even if it is the case that selection for one map can determine the

evolution of any other map, it could be the case that one of the maps is easier to change with
selection than the others. Therefore, the aims of this paper are 1) to model the potential
interdependence of the GP, EP and PP maps; 2) to determine how the evolution of one affects the
evolution of the other, and whether this evolution is symmetric; and 3) to establish whether or not
the three maps are equally responsive to selection.

Methods

To model how the GP, EP and PP maps can arise and interact, it is necessary to represent
developmental systems in a way that allow for genetic, environmental and parental inputs. We
therefore expand on three different and widely used GRN-based models of development, each one
entailing a different complexity and biological realism (Fig. 1A-C). From the simpler to the more
complex, these models are:

-**Basic GRN model.** This model (based on Wagner 1994) represents a simple gene regulatory
network (GRN; Fig. 1A). It consists of N_g transcription factors that have continuous, positive
concentrations (vector $G=(g_1,…,g_{Ng})$; $gi≥0$ ∀ i), and regulate the expression of each other by binding
to cis-regulatory sequences on gene promoters. The regulatory interactions of this GRN are encoded
in the N_g x N_g matrix B, whose elements B_{ik} represent the effect of gene k on the transcription of
gene i. Positive elements ($B_{ik}>0$) represent activation and negative elements ($B_{ik}>0$) represent
inhibition. A binary (0 or 1) matrix M (N_g x N_g) encodes the GRN topology, so that the interaction
B_{ik} is only active if $M_{ik}=1$. The initial state of the vector G (represented by the vector G_0) accounts
for the initial state at the beginning of development, which is supposed to be parentally determined.
The vector E contains N_e environmental factors ($N_e≤N_g$) which affect the levels of gene expression
by activating or repressing them (with an intensity of $0≤E_i≤1$).

Developmental dynamics are attained by changes in gene concentration over a number of
developmental iterations (t_{dev}), and the phenotype is recorded as the steady-state expression levels of
two arbitrarily chosen genes in t_{dev} (Fig. 1A). Only viable (temporally stable) phenotypes are
considered: normalized G, G^*, must remain the same within a threshold of 10^{-2} over an interval of
$t_{dev}/10$ developmental time units ($|G^*_{0.9·tdev}-G^*_{tdev}|≤10^{-2}$). The gene-gene interactions within the GRN
follow a non-linear, saturating Michaelis-Menten dynamics (a special type of Hill function), so that
the concentration of the gene i changes over developmental time according to the following
differential equation:

$\dfrac{\partial g_i}{\partial t} = \dfrac{R(h_i)}{K_M + R(h_i)} - \mu g_i + \xi$ (1)

where $h_i = \sum\limits_{j=1}^{Ng} M_{ij} B_{ij} g_j + E_j$ (2)

and $R(h^i)$ is the Ramp function ($R(x)=x, \forall\ x\geq 0$ and 0 otherwise) which prevents negative
concentrations in gene products resulting from inhibiting interactions, and K_M is the Michaelis-
Menten coefficient. Without loss of generality, we set $K_M=1$ (other choices of K_M or specific Hill
functions are known not to affect the results, see Salazar-Ciudad et al. 2000). Eq. (1) was
numerically integrated using the Euler method $(\delta_t=10^{-2})$. All genes and gene products (but not
environmental factors) are degraded with a decay term $\mu=0.1$. Noise is introduced in the system
through the term $\xi \sim U(-10^{-2}, 10^{-2})$.

-GRN + Multilinear model. This model (based on Draghi and Whitlock 2012), can be viewed as a
multi-linear model of phenotypic determination (Hansen 2006) that is added to a basic GRN-model
(Wagner 1994) (Fig. 1B). The key difference with the previous model lies in how each phenotypic
trait is generated. Rather than being the expression level of one element of the GRN, each trait T_i,
$i=(1,2)$ receives a contribution from each transcription factor according to a linear coefficient:

$T_i = \sum\limits_{j=1}^{Ng} Z_{ij} g_j$ (3)

where the factor Z_{ij} represents the contribution of the j^{th} gene to the i^{th} trait ($-1<Z_{ij}<1$). Note that the
Z matrix encoding the linear coefficients is separated from the matrix B encoding the GRN itself. In
this paper, the evolutionary implications of the correlations between maps are reported on the basis
of this model.

-Lattice model. This reaction-diffusion model (based on Salazar-Ciudad et al. 2000 and Jimenez et
al. 2015) represents a simple developmental model that implements multicellular phenotypes in an
explicitly spatial context (Fig. 1C). The model describes on a one-dimensional row of N_c non-motile
cells ($N_c=16$ in our case), whose developmental dynamics is determined by a GRN (as described in
the basic model) that is identical for all cells. Interaction between the different cells is achieved
through cell-cell signalling involving extracellular diffusion of morphogens ($N_g/3$ of the GRN
elements are considered to be diffusible morphogens). Each of these morphogens has a specific

diffusion rate D_i $(0<D_i<1)$ and follows Fick's second law. Zero-flux boundary conditions are used.

Thus, the concentration of gene i over developmental time now is calculated as:

176
$$\frac{\partial g_{ij}}{\partial t} = \frac{R(h_{ij})}{K_M + R(h_{ij})} - \mu g_{ij} + \xi + D_i \nabla^2 g_{ij} \qquad (4)$$

In most works that use this model, the phenotype is conceptualised as the expression pattern of one

of the constituent genes along the row of cells (e.g. Salazar-Ciudad et al. 2000; Jimenez et al. 2015).

Here, for the sake of comparability with the other models used, we set T_1 and T_2 as the average

concentration of gene 1 in the first and last two cells of the organism ($T_1=(g_{1,1}+g_{1,2})/2$, and $T_2=(g_{1,Nc-}$

$_1+g_{1,Nc})/2)$.

Fig. 1. Experimental overview. Conceptual depiction of the three GRN-based models used in this work: **(A)** A pure GRN model
where the (two-trait) phenotype is the steady-state concentration of two arbitrary genes. **(B)** GRN + Multilinear model, where each
phenotypic trait is calculated as the weighted sum of all the elements within the steady-state GRN. **(C)** Lattice model, where the
phenotype is conceptualised as the steady-state expression pattern of one of the constituent genes (Gen 5 in this example) along a
one-dimensional row of cells that can communicate between them through cell-cell signalling. In all of these models, phenotypic
variation is created by perturbing one or several elements in the core GRN: Perturbations can be introduced in the strength of gene-
gene interaction (i.e. as genetic mutations, **D)**; in some environmental cue may regulate some environmentally-sensitive gene **(E)**; or
in the (maternally inherited) initial concentrations of each GRN element **(F)**. Perturbations on each of these three different sources of
phenotypic variation (one element of the GRN perturbed at a time) will produce a collection of two-trait phenotypes (i.e., hind- and
fore-limb lengths). If these phenotypes are plotted in a two-trait (T_1-T_2) morphospace, they can reveal the structure of the parameter-
to-phenotype maps (D-F, right panels). The linear slopes of these maps can be used as a coarse description of these maps, allowing
for map-to-map comparisons of random (Fig. 2) and evolved GRNs (Figs 3-5).

While these models differ in complexity, all three feature a GRN at their core, which
provides an intuitive way to link each of the constituent elements of the GRN to a different source
of phenotypic variation: (i) changes in gene-gene interaction strengths in the GRN can be
conceptualised as an effect of genetic variation; (ii) changes in the environmentally sensitive
elements in the GRN (sensor nodes and diffusion rates) as an effect of environmental variation; and
(iii) changes in the initial concentration of each transcription factor in the GRN as an inherited
initial state (Fig. 1D-F). This approach exhausts the ways in which a specific GRN can vary (these
variations do not alter the GRN topology (i.e. the M matrix), which here is assumed to evolve much
slower than the inputs, Salazar-Ciudad et al. 2003).

With the described settings, all the models used in this paper produce a single, 2-trait (2-
dimensional) phenotype for each combination of inputs. Thus, a set of phenotypes (i.e. a phenotype
distribution) can be generated by introducing variation in those inputs. These phenotype
distributions represented in a 2D morphospace are considered maps: those resulting from variation
in the genetic inputs are GP maps, whereas those resulting from environmental perturbations or
perturbations in the initial conditions are considered as EP and PP maps, respectively. The 2D
representation of the whole set of phenotypes that a given developmental mechanism can create
from the sum of all perturbations is referred to as a general phenotype distribution, or GPD (i.e., the
GP, EP and PP maps are all contained within this GPD, Fig. S1).

Results

GP, EP and PP mappings are correlated in randomly generated GRNs. We first explore the
inter-dependence between GP, EP, and PP maps in a large ($n>10^6$) ensemble of randomly generated
GRNs. We then separately introduced random variation (10 input values $0<x<1$) in the genetic,
environmental and parental (i.e., initial conditions) inputs of each of these GRNs, and compared the
resulting phenotypic distributions, that is, the resulting GP, EP and PP maps.

We estimated the similarity between these three maps by testing whether or not variation in
the genetic, environmental or parental inputs produce similar covariation between the two traits
(Fig. 1D-F), using the linear slopes in the phenotypic morphospace as basic descriptors of the
different maps. Pairwise comparisons between the slopes caused by variation in genetic,
environmental, or parental inputs were all significantly positive (Pearson $r \geq 0.3$; Fig. 2A). This
demonstrates that GP, EP and PP mappings are not independent in random GRNs. Note that this
positive correlation between maps does not imply that the map-specific slopes themselves are
positive; only that their slopes, which can be positive or negative, are similar (a comparison
between maps that does not consider the direction of the slope is necessary since genetic

evolvability is concerned with the sensitivity to random mutations, not their direction; Watson
2020). This interdependence across different mappings is stronger for large and densely connected
GRNs (Fig. S2), and is robust to more detailed measures of map-to-map similarity, such as
Euclidean distances (Fig. S3). GRNs showing zero or negative correlation between different
mappings also exist, but they are less frequent (Fig. 2B).

Fig. 2. Phenotypic distributions arising from genetic, epigenetic or environmental perturbations are not independent. In a
large random ensemble of GRNs ($n=10^6$), systematic parametric variations were introduced into each of their elements. Each
perturbation on an element generates a collection of phenotypes in a two-trait morphospace (a *XPM* map), characterised by a linear
slope S_{XPM} (see Fig. 1D-F). **(A)** For each GRN, we compare these slopes, two by two, searching for their correlations in the two-slope
spaces (note that these are not two-trait morphospaces). Each dot is a GRN, and the yellow shaded region contains 90% of the GRNs.
Correlations are significant (Pearson $r>0.3$) for every combination of maps considered. **(B)** Histograms showing the probability
distribution of maps with developmental insensitivity to the first ($Sx\leq0.01$; yellow) or second ($Sy\leq0.01$; red) type of inputs; and of
correlated (parallel slopes); anti-correlated (perpendicular slopes) and non-correlated slopes (otherwise). Each of these cases
correspond to the sector of a hypothetical circumference engulfing all points of (A), as exemplified in the coloured circumference,
and the relative frequency represents the probability of each point to be located within each sector. **(C)** The complexities of the
parameter-to-phenotype maps (i.e., how non-linear they are, see Methods); rather than between their linear slopes are also positive
(Pearson $r>0.56$). In (C), the colour represents slope similarity: similar slopes (black colour) are associated to simpler (i.e., more

To eliminate the possibility that these observed correlations were caused by similarities in

the input values, rather than in the structure of the GRNs, we gradually randomize the input

parametric values while recording the correlations between maps (Fig. S4). This procedure revealed

that the correlations do not depend on particular choices of the input parameters. In contrast,

correlations are extremely sensitive to parametric changes in the GRN topology, suggesting that the

observed similarity between maps is caused by the structure of the GRN connections (i.e.,

'developmental mechanism' *sensu* Salazar-Ciudad 2000; Jimenez et al. 2015) rather than the

structure of the input perturbations.

The complexity of GP, EP and PP maps are correlated in randomly generated GRNs. We also

investigated whether or not the GP, EP and PP maps exhibit similar complexity in random GRNs.

We defined map complexity as the degree of non-linearity in phenotypic response to inputs. This

captures the intuition that a linear slope is less complex than a U-shaped response, which is itself

simpler than a W-shaped response. Comparing the map complexities between the 10^6 random GRNs

reveals that map complexities are, on average, positively correlated (Pearson $r \geq 0.56$; Fig. 2C and

S2). In other words, if a map (e.g., GP) is simple, the other maps (EP and PP) will be simple too,

and they will exhibit very similar slopes. In contrast, if a map is complex, other maps too are likely

to be complex, and their slopes will be less similar (Figs. 2C and S2).

To ensure that these observed correlations between map complexities are not a general

property of pairs of input-output maps, we analysed a large ensemble of random mathematical

functions (polynomials of known degree ≤ 4) using the same tools that we used for calculating map

complexity. This analysis verified that the correlations do not arise between pairs of randomly

selected functions unless they belong to the same complexity class (polynomial degree) (Fig. S5).

How a network topology creates similarity between map slopes and complexities can be

better understood by looking at the whole set of developmentally attainable phenotypes (general

phenotypic distribution: GPD), which can be revealed by means of massive and unspecific

parametric perturbations (see Methods and Fig. S1). This procedure shows that each generative

network creates a distinctive GPD with a highly anisotropic and discontinuous structure. This

structure forces individual maps to be aligned in the same direction simply because many

phenotypic directions of change are either very unlikely or developmentally impossible (Figs. S1,

S2, S4 and S5).

Positive map-to-map correlations in both slopes and map complexities were found in all

three considered models of phenotypic determination (Fig. S2). However, the correlation

coefficients are higher and more variable for complex models involving more than pure-GRN
dynamics (Figs. 1B-C and Fig. S2).

Evolving only one of the GP, EP or PP maps changes the phenotypic biases across the other
maps. After exploring the map-to-map correlations in random GRNs, we next wanted to address
whether or not adaptive changes within one map (i.e., changes in the covariation between traits) are
able to induce similar changes to the other maps. To do so, we performed three sets of selection
simulations using the developmental model of intermediate complexity (GRN + multilinear). In
each simulation, we allowed only one of the three different maps (henceforth the "selected map") to
evolve in response to selection. We refer to the other maps as the "non-selected" maps (see SI for
details).

At each evolutionary time step, we introduce variation only in the input associated with the
selected map (i.e., genetic, environmental or parental inputs). In response to that variation, each
individual develops a set of phenotypes that is compared to an arbitrary (linear) target map to
determine the individual's fitness. Thus, the entire phenotype distribution produced by the selected
map is accessible to natural selection (i.e., fine-grained selection). On the contrary, the inputs of the
non-selected maps were kept fixed (no variation) during simulations, so that these maps remain
effectively "invisible" to natural selection (Fig. 3A). Once each evolutionary simulation reached a
steady state, we assessed if there were any changes in the non-selected maps. We did this by
collapsing the variation for the selected map to a single input value ($x\sim U(0,1)$) and introducing
variation to each of the non-selected maps. This experimental setup guarantees that any observed
changes in non-selected maps can be attributed to indirect effects of direct selection on the selected
map.

The results revealed that evolving any one map modifies the other maps as well, introducing
in them the same adaptive phenotypic biases as observed in the selected map (Pearson *r>0.3*, Fig.
3A). This holds true for every map combination (Fig. 3B) and across the entire range of parameters
we tested (Fig. S2). However, the phenotype biases in the non-selected maps are not as strong ($|r|$
<0.5 in some cases) as in the map under selection (*r=0.99*), and exhibit substantial temporal
variation (Fig. 4). The results in Figure 3 illustrate the outcome of selecting for a linear map with a
slope *S=1* in the two-trait morphospace, but simulations with *S=-1* or with changing selective
pressures yielded similar results (Figs. 4 and S6).

Fig. 3. Evolving a single map creates similar phenotypic distributions in the other maps. A population whose individuals
initially exhibit a random phenotypic distribution in $t=0$ **(A**, small panels) is evolved to fit a target phenotypic distribution ($S^T=1$)
using as an input just one kind of phenotypic determinant (i.e., genetic, environmental, or parental variation). Other targets ($S^T=-1$)
give similar results (see Fig. S6). In each generation, one individual is exposed to 10 different input values ($0<x<1$) of a single
phenotypic determinant (the colour of each dot in A-B represents value of this input). This parametric variation produces a set of ten
potential phenotypes whose slope is compared to the target to evaluate the individual's fitness (see Methods). After 10^5 generations in
a mutation-selection-drift scenario (where other sources of phenotypic variation are frozen), the population has a narrow phenotypic
distribution in the evolved map (A, large panels). In **(B)** we uncover variation in the other maps by introducing parametric variation
($0<x<1$) in the phenotypic determinants that were kept fixed during the evolutionary trial. Results reveal that selection on a single
map creates significant side-effect phenotypic distributions in the other maps that are not the target of selection. **(C)** Correlations in
the side-effect maps are significant across all parameter values at which the parameter of the evolved map is frozen. $p=64$
individuals; $n=30$ replicates, GRN + Multilinear model.

Maps evolve faster under fine-grained selection than under coarse-grained selection. In the
previous experiments, each evolving population was allowed to sample a wide range of genetic,
parental or environmental inputs in each generation, and selection therefore acted on a wide range
of phenotypic outputs. In other words, we assumed a very fine-grained selection. Several studies
show that adaptive plasticity readily evolves when selection is fine-grained (Levins 1966; Via and
Lande 1985; De Jong 1995), although it is not essential (Rago et al. 2019). Whether or not a similar
effect occurs for GP and PP mappings is unknown. To address this, we explored the ability of every
map to adapt to a target map under different levels of selective grain, ranging from very fine-
grained selection (where individuals can experience several inputs within their lifetime) to the very

coarse-grained cases in which there is just one input per generation and this input only shifts every
n generations.

Fig. 4. Side-effect phenotypic distributions are able to track shifting targets. These plots show how initially unstructured populations are able to adaptively evolve a specific target distribution (a single map with a defined slope of $S^T=1$, solid lines), which creates as a side-effect correlated phenotype distributions in the other maps (dashed lines). The middle point corresponds to the steady-state situation shown in Fig. 3. For these plots, the target slope has been shifted to $S^T=-1$ at generation $t\approx10^4$, showing how the maps that evolve as a side-effect are able to "follow" the one that is being selected. This pattern is similar for every map considered as long as the selective grain is the same. $p=64$ individuals; $n=30$ replicates, GRN + Multilinear model.

As Fig. 5 shows, all maps evolve more efficiently under fine-grained selection than under coarse-grained selection. Furthermore, the ability to adapt to the target map escalates sharply around a grain value of 1 (Figs. 5 and S8). Under the metrics adopted here (see SI), this is the value where single individuals experience on average more than one input per generation. This implies that it is much easier to evolve a map efficiently, and thus to affect the other maps, if there is within-lifetime variation in the inputs to that map.

In most organisms, individuals can experience different environmental inputs during its lifetime but are limited to a single genotype and a single set of parentally inherited initial conditions. As a result, the EP map would be the one most intensely sculpted by natural selection.

Because of this, the EP map can exercise a stronger influence on the other maps than *vice versa* when all the three maps evolve simultaneously (Fig. S7).

Besides selective grain, the ability of GRNs to evolve a target map also depends on the complexity of the target map (i.e., how non-linear is the phenotypic response to input variation). Simple (i.e., linear) maps can easily evolve with moderate fine-grained selection (≈2 inputs per lifetime) whilst evolving more complex (i.e., quadratic or cubic) map requires an increasing number of inputs per lifetime (Fig. S8).

Fig. 5. Map evolvability depends on selective grain, and it is maximal for Environment-Phenotype maps. In figures 3 and 4, simulations assumed that natural selection could act on the entire map. This corresponds to very fine-grained selection since we define selective grain as the average number of parameter-phenotype points that be experienced by a single individual in each generation (and hence "seen" by natural selection, see Methods). In this experiment, the assumption about high fine-grainedness has been relaxed. For each level of selective grain, the ability of natural selection to evolve a linear map with an arbitrary slope is recorded as the long-term fitness. Results for environment-phenotype (*EP*) map are depicted in red; for genotype-phenotype (*GP*) map in blue and for parental-phenotype (*PP*) map in yellow. These plots show that the ability to adapt to a target slope increases non-linearly with selective grain, and that maximal efficiency is achieved when selection is fine-grained (>1, shadowed areas), which corresponds to scenarios in which single individuals can experience more than one input per generation. Such high levels of selective grain are typically only attainable for Environment-Phenotype (*EP*) maps (see main text). Points correspond to individual replicates, and dashed lines to averages over the *n=30* replicates. For each replicate, the target map is a linear function of arbitrary non-zero slope. Euclidean-distance (*ED*)-based fitness. *p=64* individuals; *t=10⁴* generations, GRN + Multilinear model.

Discussion

Understanding how the processes that generate phenotypic variation interacts with natural selection is necessary to explain and predict the course of evolution (Alberch 1982; Kavanagh et al. 2007; Salazar-Ciudad and Marin-Riera 2013; Uller et al. 2018). While it is easy to understand that any developmental bias aligned with adaptive demands would facilitate adaptation, it is not obvious how these biases originate, nor how they might change or be maintained over evolutionary time.

Phenotypic adaptation can precede genetic adaptation, and it has been suggested that plasticity
therefore facilitates genetic evolution (reviewed in West-Eberhard 2003). However, trying to
explain genetic evolvability by presupposing the existence of adaptive plasticity overlooks the fact
that adaptive plasticity is itself a product of genetic evolution. If explaining adaptive plasticity
requires past genetic evolution to have already produced adaptive phenotypic responses to particular
environmental cues, this does not help to explain genetic evolvability itself. The idea that plasticity
and evolvability are intrinsically linked through development provides a way that selection for one
can result in the evolution of the other, as studied here. Our aims have been to explore this linkage
using mechanistic models of developmental dynamics and thus explore the evolutionary
consequences of the relationship between plasticity and evolvability.

Our results show that a concordance between the GP, EP and PP maps is intrinsic to
developmental dynamics based on GRNs (with or without selection). Because of this concordance,
selection for any map will affect the other maps in an essentially symmetric fashion. However, the
efficiency of natural selection in sculpting the different maps is not symmetric: while the immediate
selective gains of evolvability are small on short timescales (e.g., to individuals), the selective gains
of plasticity can be large. Accordingly, selection for plasticity is much more effective in changing
the GRN and genetic evolvability than direct selection for genetic evolvability. Thus, without
overlooking the fact that adaptive plasticity is itself a product of genetic evolution, we show how
the genotype-phenotype (GP) map can be adaptively shaped by selection for phenotypic plasticity,
suggesting that adaptation to environmental variation helps explain the remarkable genetic
evolvability of organisms in nature.

The results show that the phenotypic effects of genetic, parental and environmental sources
of variation are typically similar across a wide range of assumptions. This finding generalises
previous research that found similar correlations on other theoretical grounds or for more restricted
scenarios, such as selection for developmental robustness to environmental perturbation (Ancel and
Fontana, 2000; Salazar-Ciudad 2006; Salazar-Ciudad 2007; Draghi and Whitlock 2012; Furusawa
and Kaneko 2018). Moreover, since the concordance appears in randomly generated regulation
networks, it does not require the concourse of past selection to produce this concordance. Rather,
stability analysis revealed that correlations between GP, EP, and PP maps are caused by the structure
of the network itself and its associated dynamical properties. Although this result might be expected
for statistical models that assume linearity and additivity (e.g., the equality $P=GxE$ from
quantitative genetics, Lande 1982), it is non-trivial for mechanistic models like the ones presented
here since these involve non-linear interactions between the inputs. These interactions create a non-
uniform space of phenotypic possibilities (a generalised phenotype distribution; GPD) that includes
regions of the morphospace that most of the parameter combinations map onto (i.e., phenotypic

attractors; Furusawa and Kaneko 2018), and 'forbidden' regions that cannot be attained by any
parameter combination. These features of the GPD impose strong limitations on the phenotypic
variation that is possible, and the shape of the genotype-, environment-, and parental-phenotype
maps will be similar since they share the same attractors.

These shared attractors explain why a population that has evolved a specific (e.g., EP) map
will show similar biases in all its maps, even when those have not been selected for. However, this
dependence would not make plasticity exercise a disproportionate effect on genetic evolvability
unless there was some asymmetry that makes selection for properties of the EP map more efficient
than selection for properties of the GP or PP maps. We show that this crucial asymmetry follows
from differences in the temporal timescale ('grain') of environmental, parental and genetic variation
that is input to these maps. Specifically, the variational properties of a map evolve faster when
individuals experience multiple inputs, and hence can develop multiple selectable phenotypes,
during their lifetime (Levins 1966; Via and Lande 1985; De Jong 1995). While individuals can
experience multiple environments during their lifetime, they do not experience multiple genotypes
or initial conditions (e.g., the distribution of phenotypes produced under genetic variation is a
property of a family, population or lineage, not an individual). As a result, selection for GP and PP
maps should typically be more coarse-grained and less efficient than for EP maps. This general
property of natural selection causes adaptive EP maps to generally evolve more readily than
adaptive GP (and PP) maps, even though they depend on the same developmental dynamics.

These results generate predictions can be tested empirically, for example, by means of
experimental evolution. One particularly useful approach would be to select populations in
environments of different variability (i.e., selective grain), which should result in populations with
different EP maps. The prediction is that the finer the selective grain, the more the structure of the
GP map will resemble that of the EP map, which can be tested using mutation accumulation
experiments or genetic engineering. Whether or not such changes in the GP map changes the
capacity for future adaptation could be tested by exposing populations to new selective regimes that
are more or less structurally similar to what the population initially adapted to. Other experimental
and comparative approaches could also test one or several of the predictions of the relationship
between EP, PP, and GP maps (e.g., Lind et al. 2015; Noble et al. 2019; Radersma et al. 2020).

While our results demonstrate that natural selection on phenotypic plasticity would cause the
GP map to evolve, they also show that the ability to evolve a certain map is severely limited by the
map complexity itself, with complex (e.g., cubic) maps requiring highly fine-grained selection. This
would render very complex EP (and GP) maps unreachable by adaptive evolution even in the most
fine-grained scenarios. However, complex maps are known to exist, which suggests that other non-
selective processes, such as developmental system drift (True and Haag 2001; Jimenez et al. 2015),

may play an important role in developmental evolution (Alberch 1982; Salazar-Ciudad and Marin-Riera 2013). This possibility is compatible with our results, which simply state that whenever adaptive developmental biases do exist, they will be predominantly generated through selection for phenotypic plasticity. Furthermore, since each trait needs to maintain its function as other parts of the organism develop and grow, selection for plasticity can be even more fine-grained that expected (our models do not fully capture this developmental dependence because they record the individual fitness after a fixed developmental time).

These results shed light on whether or not plasticity can exercise a predominant role in adaptive evolution, a hypothesis with a long and contentious history in evolutionary biology (see Weber and Depew 2003; West-Eberhard 2003; Schwander and Leimar 2011; Laland et al. 2014; Uller et al. 2019; Parsons et al. 2020). While adaptive modification of environmentally induced phenotypes can make plasticity appear to 'take the lead' in evolution without any link between plasticity and genetic evolvability (Radersma et al. 2020), the evolutionary change in the GP map caused by adaptive plasticity suggests that evolution is particularly likely to proceed where plasticity leads. Over longer timescales, this process provides a biologically plausible mechanism for the internalisation of environmental information, resulting in developmental biases whose structure 'mirrors' the structure of the selective environment (Riedl 1978), and thereby facilitating further adaptations through genetic modification of environmentally induced phenotypes (Draghi and Whitlock 2012; Watson et al. 2014; Rago et al. 2019; Parsons et al. 2020). Without denying the importance of other (adaptive and non-adaptive) processes, this constitutes a strong argument for a role of plasticity in shaping the path of evolution.

Acknowledgements:

The authors thank J.F. Nash, D. Prosser, J. Caldwell, L.E. Mears, I. Hernando-Herraez and R. Zimm for helpful discussion and comments.

Author contributions:

M.B.U., T.U. and R.A.W. conceived the idea, which was later refined by all the authors. M.B.U., A.R., Ch.T. and R.A.W. established the experimental design. M.B.U. wrote the software and conducted the *in silico* experiments. All authors contributed to the interpretation of the results. M.B.U., A.R., T.U. and R.A.W. wrote the manuscript.

Competing interests:

The authors declare no competing interests.

References

Alberch, P. 1982. Developmental constraints in evolutionary processes. In Evolution and development (pp. 313-332). Springer, Berlin, Heidelberg.

Ancel, L. W. and Fontana, W. 2000. Plasticity, evolvability, and modularity in RNA. J. Exp. Zool. 288:242-283.

Badyaev, A. V., and T. Uller. 2009. Parental effects in ecology and evolution: mechanisms, processes and implications. Philos. Trans. Roy. Soc. B. 364:1169-1177.

Crispo, E. 2007. The Baldwin effect and genetic assimilation: revisiting two mechanisms of evolutionary change mediated by phenotypic plasticity. Evolution. 61:2469-2479.

Cheverud, J. M. 1982. Phenotypic, genetic, and environmental morphological integration in the cranium. Evolution 36:499-516.

De Jong, G. 1995. Phenotypic plasticity as a product of selection in a variable environment. Am. Nat. 145:493-512.

Draghi J. A., and M. C. Whitlock. 2012. Phenotypic plasticity facilitates mutational variance, genetic variance, and evolvability along the major axis of environmental variation. Evolution 66:2891-2902.

Furusawa, C., and K. Kaneko. 2018. Formation of dominant mode of evolution in biological systems. Phys. Rev. E. 97:042410.

Hallgrímsson, B., and B. K. Hall. 2005.Variation: A central concept in biology. Elsevier academic press.

Hansen, T. F. 2006. The evolution of genetic architecture. Annu. Rev. Ecol. Evol. Syst. 37:123-157.

Houle, D, Govindaraju D. R., and S. Omholt. 2010. Phenomics: the next challenge. Nature Rev. Genet. 11:855–866.

Jablonka, E. and M. J. Lamb. 2005. Evolution in Four Dimensions: Genetic, Epigenetic, Behavioral, and Symbolic Variation in the History of Life. MIT Press.

Jiménez, A., Cotterell, J., Munteanu, A. and J. Sharpe. 2015. Dynamics of gene circuits shapes evolvability. Proc. Natl. Acad. Sci. USA. 112:2103-2108.

Kavanagh, K. D., Evans, A. R. and J. Jernvall. 2007. Predicting evolutionary patterns of mammalian teeth from development. Nature. 449:427-432.

Kirschner, M., and J. Gerhart. 1998. Evolvability. Proc. Natl. Acad. Sci. USA. 95:8420-8427.

Laland, K., Uller, T., Feldman, M., Sterelny, K., Müller, G. B., Moczek, A. P., and E. Jablonka. 2014. Does evolutionary theory need a rethink?. Nature News. 514:162-164.

Lande, R. 1982. A quantitative genetic theory of life history evolution. Ecology. 63:607-615.

Levins, R. 1966. The strategy of model building in population biology. Am. Sci. 54:421-431.

Levis, N. A., and D. W. Pfennig. 2016. Evaluating 'plasticity-first'evolution in nature: key criteria and empirical approaches. Trends Ecol. Evol. 31, 563-574.

Li, X., Guo, T., Mu, Q., Li, X., and J. Yu. 2018. Genomic and environmental determinants and their interplay underlying phenotypic plasticity. Proc. Natl. Acad. Sci. USA. 115:6679-6684.

Lind, M. I., Yarlett, K., Reger, J., Carter, M. J., and Beckerman, A. P. 2015. The alignment between phenotypic plasticity, the major axis of genetic variation and the response to selection. Proc. R. Soc. B: Biol. Sci. 282(1816):20151651.

Newman, S. A., and G. B. Müller. 2000. Epigenetic mechanisms of character origination. J. Exp. Zool. 288:304-317.

Noble, D. W., Radersma, R., and T. Uller. 2019. Plastic responses to novel environments are biased towards phenotype dimensions with high additive genetic variation. Proc. Natl. Acad. Sci. USA. 116:13452-13461.

Parsons, K. J., McWhinnie, K., Pilakouta, N., and L. Walker. 2020. Does phenotypic plasticity initiate developmental bias?.Evol. Dev.22:56–70.

Radersma, R., Noble, D.W. and Uller, T. 2020. Plasticity leaves a phenotypic signature during local adaptation. Evolution Letters. doi:10.1002/evl3.185

Rago, A., Kouvaris, K., Uller, T., and R. A. Watson. 2019. How adaptive plasticity evolves when selected against. PLoS Comp. Biol. 15:e1006260.

Riedl, R. 1978. Order in Living Organisms. Wiley, New York.

Salazar-Ciudad, I., Garcia-Fernandez, J., and R. V. Solé. 2000. Gene networks capable of pattern formation: from induction to reaction–diffusion. J. Theor. Biol. 205:587-603.

Salazar-Ciudad, I., Jernvall, J., and S. A. Newman. 2003. Mechanisms of pattern formation in development and evolution. Development 130:2027-2037.

Salazar-Ciudad, I. 2006. Developmental constraints vs. variational properties: how pattern formation can help to understand evolution and development. J. Exp. Zool. B Mol. Dev. Evol. 306:107-125.

Salazar-Ciudad, I. 2007. On the origins of morphological variation, canalization, robustness, and evolvability. Int. Comp. Biol. 47:390-400.

Salazar-Ciudad, I. and M. Marin-Riera. 2013. Adaptive dynamics under development-based genotype-phenotype maps. Nature. 497:361–364.

Schwander, T., and O. Leimar. 2011. Genes as leaders and followers in evolution. Trends Ecol. Evol. 26:143-151.

True, J. R., and E. S. Haag. 2001. Developmental system drift and flexibility in evolutionary trajectories. Evol. Dev. 3:109-119.

Uller, T., Moczek, A. P., Watson, R. A., Brakefield, P. M., and K. N. Laland. 2018. Developmental bias and evolution: A regulatory network perspective. Genetics 209:949-966.

Uller, T., Feiner, N., Radersma, R., Jackson, I. S., and Rago, A. 2019. Developmental plasticity and evolutionary explanations. Evolution & Development, 22:47-55.

Via, S., and R. Lande. 1985. Genotype-environment interaction and the evolution of phenotypic plasticity. Evolution 39:505-522.

Wagner, A. 1994. Evolution of gene networks by gene duplications: a mathematical model and its implications on genome organization. Proc. Natl. Acad. Sci. USA. 91:4387-4391.

Wagner, G. P., and Altenberg, L. 1996. Perspective: complex adaptations and the evolution of evolvability. Evolution 50:967-976.

Watson, R. A., Wagner, G. P., Pavlicev, M., Weinreich, D. M., and R. Mills. 2014. The evolution of phenotypic correlations and "developmental memory". Evolution 68:1124-1138.

Watson, R.A. 2020. Evolvabiity. In Evolutionary Developmental Biology: A Reference Guide. Nuño de la Rosa, L. and Müller, G.B. Eds. Springer

Weber, B. H., and Depew, D. J. 2003. Evolution and learning: The Baldwin effect reconsidered. Mit Press.

West-Eberhard, M. J. 2003. Developmental plasticity and evolution. Oxford University Press.