The transcription factor Pdr802 regulates *Cryptococcus neoformans* Titan cell production and pathogenicity

Julia C. V. Reuwsaat[1,2], Heryk Motta[1], Daniel P. Agustinho[2], Holly Brown[2], Andrew L. Chang[2], Livia Kmetzsch[1,3,4], and Tamara L. Doering[2,4].

[1]Molecular Biology of Pathogens Laboratory, Biotechnology Center, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

[2]Department of Molecular Microbiology, Washington University School of Medicine, Washington University, St. Louis, Missouri, USA.

[3]Department of Molecular Biology and Biotechnology, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

[4] Corresponding authors: liviak@cbiot.ufrgs.br, doering@wustl.edu

Running head: Role of Pdr802 in *Cryptococcus neoformans* virulence.

ABSTRACT

Cryptococcus neoformans is a ubiquitous, opportunistic fungal pathogen that kills almost

200,000 people worldwide each year. It is acquired when mammalian hosts inhale the

infectious propagules; these are deposited in the lung and, in the context of

immunocompromise, may disseminate to the brain to cause lethal meningoencephalitis.

Once inside the host, *C. neoformans* cells develop a variety of features that help them

avoid immune recognition, including display of a polysaccharide capsule that impairs

phagocytosis and the production of giant (Titan) cells. The transcription factor Pdr802 is

one regulator of these responses to the host environment. Expression of the

corresponding gene is highly induced under host-like conditions *in vitro* and is critical for

C. neoformans dissemination and virulence in a mouse model of infection. Direct targets

of Pdr802 include the calcineurin targets Had1 and Pmc1, which are important for *C.*

neoformans virulence; the transcription factor Bzp4, which regulates cryptococcal

melanization and capsule thickness; and 35 transmembrane transporters. Notably, a

strain engineered to lack Pdr802 showed a dramatic increase in Titan cells, which are not

phagocytosed and have diminished ability to directly cross biological barriers. This

explains the reduced dissemination of *pdr802* mutant cells to the central nervous system

and the consequently reduced virulence of this strain. The role of Pdr802 as a negative

regulator of titanisation is thus critical for cryptococcal pathogenicity.

IMPORTANCE

The pathogenic yeast *Cryptococcus neoformans* presents a worldwide threat to human health, especially in the context of immunocompromise, and current antifungal therapy is hindered by cost, limited availability, and inadequate efficacy. After the infectious particle is inhaled, *C. neoformans* initiates a complex transcriptional program that integrates cellular responses and enables adaptation to the host lung environment. Here we describe the role of the transcription factor Pdr802 in the response to host conditions and its impact on *C. neoformans* virulence. We identified direct targets of Pdr802 and also discovered that it regulates cellular features that influence movement of this pathogen from the lung to the brain, where it causes fatal disease. These findings significantly advance our understanding of a serious disease.

INTRODUCTION

Cryptococcosis is a fungal infection caused by *Cryptococcus neoformans* and *Cryptococcus gattii. C. neoformans* is a ubiquitous opportunistic pathogen that infects mainly immunocompromised patients, while *C. gattii* is capable of infecting immunocompetent individuals (1). Cryptococcosis causes 180,000 deaths worldwide each year, including roughly 15% of all AIDS-related deaths (2), and is initiated by the inhalation of spores or desiccated yeast cells. In immunocompetent individuals, this typically causes an asymptomatic pulmonary infection that is controlled by the host immune response, although a population of *C. neoformans* may remain latent for extended periods of time (3, 4).

Under conditions of immunocompromise, cryptococci disseminate from the lung to the brain. Mechanisms that have been suggested to mediate fungal crossing of the blood-brain barrier (BBB) include transcellular migration, in which the yeast cells enter and exit vascular endothelial cells (5–8); paracellular movement, in which they cross the BBB at junctions between endothelial cells (9–11); and 'Trojan horse' crossing, whereby macrophages harboring *C. neoformans* enter the brain (12). Cryptococcal meningoencephalitis is difficult to treat and frequently lethal, for reasons that include lack of drug availability and high cost (13, 14).

The ability of *C. neoformans* to survive and proliferate in the lung, and subsequently disseminate to the brain, depends on the expression of multiple virulence traits, such as secreted factors (15, 16), a polysaccharide capsule that surrounds the cell wall (17), and the production of giant (Titan) cells (18, 19). For example, the secreted pigment melanin associates with the cell wall, where its antioxidant properties protect fungal cells from reactive oxygen species produced as a host immune defense (20–23). Urease, a metalloenzyme that converts urea to ammonia and CO_2, is secreted extracellularly and may affect the course of infection by modulating environmental pH and damaging host tissue structure (10, 11, 24).

The capsule, composed primarily of large polysaccharides (25–27), is a key cryptococcal virulence factor that impairs phagocytosis by immune cells (28–33). This dynamic entity changes its size and structure during interactions with the host or external environment (34–38), contributing to fungal adaptation (39). Capsule polysaccharides that are shed from the cell enable diagnosis of cryptococcal infection and also impede host responses (33, 40).

Titan cells are a cryptococcal morphotype that has been variously characterized as

having cell wall diameter greater than 10 or 15 µm or capsule diameter that exceeds 30

µm (19, 41, 42). These cells can be polyploid and produce normal-size cells during

infection (18, 43, 44). Titan cell formation is triggered by exposure to the host

environment, including nutrient starvation, reduced pH, and hypoxia (45–47), although

the extent of titanisation depends on the host immune response and the duration of

infection (43, 48). Titan cell production appears to benefit the development of *C.*

neoformans infection in the lungs, since there is reduced internalization of these large

cells by host phagocytes and they are more resistant to oxidative stress than normal-size

cells (18, 44). Some of these effects may be explained by the highly cross-linked capsule

and thickened cell wall of Titan cells (49). In contrast, Titan cells show impaired

dissemination to the brain (18, 44).

C. neoformans experiences a dramatic change in conditions upon entering a host,

including altered nutrient levels and pH. To adapt to the new environment, cryptococci

activate a network of transcription factors (TFs) (37, 50). For example, imbalances in ion

homeostasis trigger responses from the TFs Zap1 (51), Cuf1 (52), Pho4 (53), Cir1 (54),

and Crz1 (55). Alkaline pH stimulates expression of the TF Rim101, which enables

growth under basic conditions and other stresses such as high salt and iron limitation; it

also promotes the association of capsule polysaccharide with the cell and the formation

of Titan cells (45, 56).

Overlapping TF circuits regulate cryptococcal virulence determinants, including

polysaccharide capsule production and melanin synthesis. For example, Usv101, an

important regulator of capsule thickness and polysaccharide shedding, also regulates

three other TFs (Gat201, Crz1, and Rim101) and multiple polysaccharide-related

enzymes (57). Gat201 further regulates additional virulence-related transcription factors

and the anti-phagocytic protein Blp1 (58), while Crz1 plays a central role in the

maintenance of plasma membrane and cell wall stability (55, 59, 60). Crz1 expression is

also modulated by the calcineurin signalling pathway, which is required for normal yeast

growth at 37°C, virulence, and sexual reproduction (61). A group of TFs, including

Usv101, Bzp4, Hob1, and Mbs1 (57, 62), act together to regulate melanin production;

deletion of Bzp4 also alters capsule production (50).

In this study, we investigated the TF Pdr802. The corresponding gene is highly

variable, based on its rate of non-synonymous mutations, which suggests it is evolving

rapidly (63). Pdr802 has been implicated in *C. neoformans* virulence (37, 50, 64), but its

specific role and targets are not known. We discovered that Pdr802 is induced in host-like

conditions, is a negative regulator of Titan cell formation, and influences capsule

thickness and phagocytosis by macrophages. These functions, as well as its regulation of

HAD1, *PMC1* and *BZP4*, whose products act in cell wall remodeling and virulence factor

production, make Pdr802 critical for both lung survival and dissemination to the brain.

RESULTS

The role of Pdr802 in *C. neoformans* virulence

The importance of Pdr802 in *C. neoformans* virulence has been demonstrated in

multiple experimental models. Liu and collaborators first reported in 2008 that partial

deletion of *PDR802* reduced *C. neoformans* infectivity in a competition assay of pooled

C. neoformans strains (64). In 2015, Maier *et al* showed that a *pdr802* deletion mutant

had reduced virulence when tested individually in a short-term mouse model of infection

(37). Later that year, Jung and colleagues reported that Pdr802 was necessary for full

virulence in both wax moth larvae and short-term mouse infection using pooled strains

(50). Most recently, Lee and collaborators showed that Pdr802 was required for brain

infection (65).

To further investigate the role of Pdr802 in pathogenesis, we complemented a

complete deletion strain in the KN99α background that we had previously generated

(*pdr802*) (37) with the intact gene at its native locus (*PDR802*). To examine targets of

Pdr802, we also constructed a strain that expresses the protein fused to mCherry at its

N-terminus (Figure S1A). All of these strains lacked or expressed RNA encoding *PDR802*

or its modified forms as expected (Figure S1B) and, importantly, *PDR802* was expressed

at wild-type levels in the complemented and modified strains (Figure S1C).

We next assessed the long-term survival of C57BL/6 mice infected with the

parental wild-type strain (KN99α), the deletion mutant (*pdr802*), or the complemented

mutant (*PDR802*). In this model, mice infected with the parent or complemented strains

survived for roughly three weeks, while those infected with the deletion mutant showed a

striking increase in survival: all animals survived for at least 65 days and over half

survived to the end of the study (100 days; Figure 1A). The lung burden measured at the

time of death for *pdr802*-infected mice in this study was approximately 100-fold lower

than that of wild type infections (Figure S2A), demonstrating the importance of this TF in

C. neoformans virulence. Mean brain burden at the time of death was more similar

between mutant and wild type infections (Figure S2A), although we did note some

heterogeneity in this measure for *pdr802*-infected mice; animals sacrificed at around two

months of infection showed brain burden similar to WT levels (red symbols), while brain

burden of mice sacrificed at day 100 ranged between zero fungal cells and WT level

(blue symbols).

We next examined the time course of fungal proliferation in the lungs. As

expected, the burdens of WT and the complemented mutant strains increased steadily

over an 18-day interval (Figure 1B), eventually reaching roughly five logs over the

inoculum. Towards the end of this time period, these cells were also detected in the blood

and brain (Figure S2B). In contrast, the lung burden of *pdr802* remained close to the

original inoculum throughout this period, with no mutant cells detected in the blood or

brain. At a late time point of *pdr802* infection (75 days), we again noted some

heterogeneity of fungal burden: one mouse had high lung burden with no dissemination,

another had high lung burden with moderate brain burden, and the third had extremely

low lung burden with no dissemination (Figure S2C). No colony-forming units (CFU) were

detected in the blood of *pdr802*-infected mice at any point during infection. Together,

these results suggest that even though the *pdr802* strain is generally hypovirulent and

remains at low levels in the lung, it can occasionally reach the brain and, given enough

time, accumulate there (see Discussion).

Given the dramatic effects of Pdr802 on fungal virulence, we wondered about the

specific biological processes in which this transcription factor is involved. We first

examined the behavior of the *pdr802* strain *in vitro*, focusing on stress conditions that

might be encountered in the host. We saw no differences in growth of the *pdr802* mutant

compared to WT cells under conditions that challenge cell or cell wall integrity, including

sorbitol, high salt, cell wall dyes, caffeine, sodium dodecyl sulfate (SDS), or ethanol

(Figure S3A-C). The mutant also showed no altered susceptibility to elements of the host

response, such as nitrosative or oxidative stresses, or in melanin production, which is

implicated in cryptococcal survival and pathogenic potential (15, 20, 21). All of these

results held whether growth was at 30°C, 37°C, or 37°C in the presence of 5% CO_2,

which was recently described as an independent stress for *C. neoformans* (66) (Figure

S3A-C). Finally, the mutant showed no difference from wild-type cells in secretion of

urease (Figure S3D), which has also been implicated in *C. neoformans* virulence and

brain dissemination (10, 16, 24).

Pdr802 is regulated by "host-like" conditions

We next tested the growth of the *pdr802* mutant under conditions more like those

encountered inside the mammalian host, using tissue culture medium (DMEM) at 37°C in

the presence of 5% CO_2. We found that although the *pdr802* mutant grew like WT in rich

medium (YPD), it grew poorly in DMEM (Figure S3E-F). To test whether the mutant cells

were dead or just static after growth in DMEM, we plated aliquots on solid medium to

measure CFU over time (Figure 2A). The *pdr802* culture showed a dramatic decrease in

viability compared to WT and the complemented strain, which was greatest in the first 24

h. This is the same time frame in which expression of the *PDR802* gene shows a striking

increase, as measured by RNA-Seq (Figure 2B).

Another important feature that is induced by growth in DMEM at 37°C and 5% CO_2

is the polysaccharide capsule, which we previously reported to be regulated by Pdr802,

based on negative staining with India ink (37). Fluorescence microscopy confirmed

increased capsule thickness of the mutant, which reverted to WT in the complemented

strain (Figure 3A). To quantify this change, we took advantage of a semi-automated

assay that we have developed (Figure S4), which measures capsules on a population

scale (Figure 3B) and is therefore very sensitive. This analysis showed that the capsule

thickness of *pdr802* cells resembles that of the well-studied hypercapsular mutant *pkr1*

(37, 67, 68) and is completely restored to WT by complementation at the native locus

(Figure 3C). Previous studies suggest that capsule thickness upon induction reflects the

size of the dominant capsule polymer (glucuronoxylomannan; GXM) (69–71), which can

be analyzed by agarose gel migration and blotting with anti-capsule antibodies (69).

Consistent with the difference we observed in capsule thickness by imaging, this method

showed decreased mobility of GXM from *pdr802* as capsule induction progressed (Figure

S3G).

To validate the observations that we had made in standard 'host-like' conditions

based on synthetic tissue culture medium, we conducted similar studies in mouse serum

at 37°C and 5% CO_2. These conditions induced an even more pronounced hypercapsular

phenotype of the *pdr802* mutant (Figures 4A and 4B), as well as reduced cell viability

(Figure 4C) and increased cell wall diameter (Figure 4D).

We were intrigued by the enlarged cell body and capsule of the *pdr802* mutant

cells in host-like conditions *in vitro* and decided to examine these phenotypes *in vivo*. For

these studies, we isolated fungal cells from the lungs of mice at various times of infection

and assessed their morphology by negative staining (Figure 5A). At each time point, the

mutant cell wall diameters were larger than the controls, but while cell wall diameter size

was stable for WT and complemented strains throughout the infection period, cell wall

diameter of the deletion mutant trended larger at the end of the infection period (Figure

5B). In contrast, mutant capsule thickness, although initially larger than that of control

cells, remained stable through the period (~15 μm), while capsule thickness of control

cells increased to that level or beyond (Figure 5C). Over time, therefore, the ratio of

capsule thickness to cell wall diameter for WT and *PDR802* cells steadily increased,

while it remained roughly constant for the mutant (Figure S5A). Furthermore, although

the capsule diameter of *pdr802* cells consistently exceeded that of WT and

complemented cells, their sizes became more comparable late in infection (Figure S5B).

Pdr802 negatively regulates Titan cells formation

We were particularly interested in the cell size phenotype of *pdr802* because Titan

cells have been strongly implicated in cryptococcal pathogenesis (18). By any definition

of this morphotype (cell walls greater than 10 or 15 μm in diameter or capsules greater

than 30 μm in diameter), our mutant cell populations were strikingly enriched in Titan

cells. For example, the fraction of mutant cells that met the capsule diameter criterion for

Titan cells (30% to 60%) greatly exceeded that of control cells (0% to 30%) (Figure S5C).

To specifically test Titan cell formation by the *pdr802* strain, we subjected mutant

cells to *in vitro* conditions that induce this process (47) and analyzed the resulting

population by flow cytometry. Consistent with our *in vivo* observations, Titan cells

constituted a much larger fraction of the population in the mutant culture (13.2%) than in

the WT culture (1.62%) (Figure 6).

Titan cells are poorly engulfed by host phagocytes (18, 43), which may reflect their

increased size as well as alterations in capsule and cell wall (49). We observed this

reduced uptake for all strains after growth in conditions that favor titanisation (Figure 7,

Titan vs YPD). Also, all strains showed a reduction in phagocytosis after capsule

induction in DMEM (Figure 7, DMEM vs YPD), which is not surprising because the

capsule is antiphagocytic (29). Notably, the reduction in uptake was greatest for the

pdr802 mutant in both of these conditions, even though it showed normal engulfment

when all strains were grown in the control condition (YPD). This is likely because the

mutant culture is both hypercapsular and enriched in Titan cells.

Pdr802 regulates transmembrane transporters and transcription factors

To identify potential direct targets of Pdr802, we next performed ChIP-Seq

analysis. We compared DNA sequences immunoprecipitated by anti-mCherry mAb from

cells expressing mCherry-Pdr802, which grow similarly to WT (Figure S6), and untagged

cells. Both strains were grown for 24 hours in DMEM at 37°C and 5% CO_2, as this

condition induces *PDR802* expression roughly 45-fold compared to control growth in YPD

(Figure 2).

Using 2-fold-enrichment over control as a cutoff value, we identified 656 peaks in

mCherry-Pdr802 genomic DNA. Of these, 540 occurred within 1,000 bp upstream of

transcription start sites (Data Set S1, Sheet 1), which we used as an approximation of

regulatory regions. Gene Ontology (GO) enrichment analysis of the corresponding 540

genes suggested a broad regulatory role for Pdr802, as the largest target group (35

transmembrane transporters) comprised just under 9% of the total targets (Data Set S1,

Sheets 7 and 8). We next applied Discriminative Regular Expression Motif Elicitation

(DREME) (72) to the complete set of upstream regions to identify potential Pdr802

binding motifs (Figure S7A), which were notable for repeated TC (GA) doublets. Finally,

we used qRT-PCR to measure the expression of three putative target genes, including

BZP4, in WT and *pdr802* cells after growth in DMEM. We observed expression changes

consistent with Pdr802 regulation (Figure S7B, see Discussion).

A transcriptional profiling study of *C. neoformans* cells exposed to Titan cell

inducing conditions *in vitro* was recently published (46). Because we had observed that

the absence of *PDR802* was associated with an increase in Titan cell formation, we

compared these results to our ChIP-Seq data. Of the 562 genes reported to decrease

expression during Titan cell formation *in vitro*, 44 (8%) were identified in our ChIP-Seq

study; 40 of these contained a Pdr802 binding motif within 1,000 bp of the start site (Data

Set S2, Sheet 1). The largest functional group of these 40 genes encodes proteins

related to regulation of gene expression (Data Set S2, Sheets 1-   2). Several of these

putative Pdr802 targets encode transcription factors, including *STB4*, *ECM2201*, *and*

FZC34. Stb4 was recently shown to be important for cryptococcal infection of the brain

(65), Fzc34 is important for proper cryptococcal capsule formation (50), and Ecm2201 is

essential for capsule formation and implicated in *C. neoformans* metal ion transport and

chitin synthesis (37). The next largest group consists of genes encoding transmembrane

transporters (Data Set S2, Sheets 1-2). Interestingly, three of these transporters

(encoded by *PTP1, CNAG_07869*, and *CNAG_00598*) are upregulated during

macrophage ingestion, as is *PDR802* (73).

A somewhat higher fraction of the genes reported to increase expression during

Titan cell formation was identified in our ChIP-Seq study (44 of 421; 10.5%), with 37 of

them exhibiting a Pdr802 binding within 1,000 bp of the start site (Data Set S2, Sheet 3).

The largest group of these 37 genes encodes proteins related to general cellular

metabolic processes (Data Set S2, Sheets 2-3). Among these are two genes that encode

enzymes implicated in glucan synthesis and degradation (*CNAG_00393* and

CNAG_05138), consistent with the importance of cell wall remodeling and capsule

synthesis in Titan cell production (49). Another potential target encodes the transcription

factor Zfc3, which is important for *C. neoformans* capsule formation (50) and brain

infectivity (65) (Data Set S2, Sheet 3).

We also examined transporters and TFs implicated as Pdr802 targets that are not

related to Titan cell formation. Interestingly, Prd802 co-immunoprecipitated with promoter

regions of genes encoding the glucose transporters Hxt1 and Hxs1, as well as the inositol

transporters Itr2 and Itr3 (Figure 8A). Both of these hexoses serve as carbon sources for

C. neoformans and the activity of these transporters promotes cryptococci virulence and

brain infectivity (74–77). During *C. neoformans* engulfment by macrophages, *PDR802* is

upregulated, together with *HXS1* and *ITR2*, while *HXT1* is downregulated (73). During *C.*

neoformans infection in mice, *PDR802* and *ITR3c* are upregulated, in both lung and brain

(65).

Among TFs, Pdr802 binds upstream sequences of the genes that encode Mln1

and Fzc14 (Figure 8B). Null mutants lacking either of these TFs are hypercapsular and

show increased sensitivity to the antifungal drug fluconazole (50). Pdr802 also interacts

with the promotor of the gene encoding Bzp4 (Figure 8B), which, as mentioned above,

regulates two virulence factors (capsule (50) and melanin (50, 62)) and is required for

cryptococcal pathogenicity (50). We found that *BZP4* expression was repressed by

Pdr802 under host-like conditions (Figure S7B).

The most highly-enriched peak in our Pdr802 studies was in the regulatory

sequence of the gene encoding Fatty acyl-CoA synthetase and RNA processing-

associated Kinase 1-like 02 (Frk102; Figure 8C), whose product is involved in cadmium

sulphate tolerance and coping with endoplasmic reticulum stress, but is not important for

C. neoformans virulence (78).

Pdr802 regulates two calcineurin target genes

The calcineurin signaling pathway is activated by calcium and governs stress

response and virulence in *C. neoformans* (79–81). A significant portion of calcineurin

signaling is mediated by the transcription factor Crz1 (55) mentioned above, which is

highly responsive to temperature (55, 60) and important for cryptococcal virulence (55).

Other calcineurin targets, however, are regulated independently of Crz1 (55, 61). This

group includes the phosphatase Had1 (55, 61), which was previously shown to be

important for *C. neoformans* virulence (82). The *HAD1* gene contains three Pdr802

binding motifs within 1,000 bp of its start site (Data Set S2, Sheet 4) and was also

identified in our ChIP study (Figure 9A). *HAD1* expression increases upon deletion of

PDR802 (Figure 9B), further supporting that it is a Pdr802 target.

Another target of calcineurin is the vacuolar calcium transporter Pmc1 (55, 61),

which is required for *C. neoformans* virulence and dissemination to the brain (83, 84).

PMC1 was identified in our ChIP-Seq analysis (Figure 9C) and has two Pdr802 binding

motifs in its promoter (Data Set S2, Sheet 4). We found that Pdr802 deletion results in

increased levels of *PMC1* expression after 3 hours of growth in host-like conditions

(Figure 9D), although this effect dissipates by 24 hours. Because Pmc1 is important for

maintenance of normal cryptococcal Ca^{2+} concentrations (83), we wondered whether this

initial increase would imbalance intracellular calcium levels. We found that cytosolic

calcium levels in the *pdr802* mutant are higher than those observed in WT and

complemented strains after 24 hours of growth in DMEM (although they are below those

of a *crz1* null mutant), which may reflect a rebound from the transient increase in *PMC1*

expression that represses accumulation of cytosolic calcium levels. In rich medium

(YPD), however, only the *crz1* deletion strain accumulated cytosolic calcium (Figure 9E),

which reinforces the idea that Pdr802 acts primarily in host-like conditions.

Since we had observed that Pdr802 regulates at least two targets of calcineurin, we

compared gene expression profiles of a calcineurin mutant (55) to our ChIP-Seq data set.

Of the 393 genes that were differentially expressed in the calcineurin mutant under

thermal stress, 45 were identified in our ChIP-Seq study and. Notably, 8 genes in this

group encode transmembrane transporters and 37 contain a Pdr802 binding motif within

1,000 bp of the start site (Data Set S2, Sheet 5).

DISCUSSION

The host lung environment encountered early in infection presents multiple

challenges to *C. neoformans* survival, including nutrient limitation, high pH, and oxidative

stress (61, 85, 86). Cryptococcal cells lacking Pdr802 are unable to proliferate in this

context, and only survive for a few days in tissue culture media. Several factors may

contribute to these defects. One may be reduced ability of this mutant to compensate for

nutrient limitation, suggested by the multiple membrane transporters implicated as

Pdr802 targets by our ChIP-Seq analysis. Because these studies do not indicate the

direction in which gene expression is altered, we examined the expression of *PDR802*

and its potential targets in published RNA-Seq data sets. We noted that *PDR802*

expression in *C. neoformans* cells rises 3-fold upon internalization by macrophages (73),

while its target *HXS1*, which encodes a high affinity glucose transporter, is upregulated 11-fold. *HXS1* expression is known to be induced by glucose limitation, so this is consistent with nutritional deprivation within phagocytes (75). In contrast, the expression of the hexose transporter *HXT1*, normally induced by glucose (74, 75), is downregulated (2.5-fold) in this context (73).

Our measurements of cytosolic calcium show that calcium availability is also modulated by Pdr802. This is likely achieved through its regulation of the calcium transporter Pmc1, demonstrated by our ChIP and qRT-PCR studies. Since calcium ion is a major second messenger in eukaryotic cells, the accumulation of Ca^{2+} in *pdr802* cells could affect multiple processes central to its host interactions, including stress responses and cell wall integrity (59, 60, 80, 83, 87). Pmc1 and at least 38 other genes that are targets of Pdr802 are also downstream targets of calcineurin (55, 61), suggesting that Pdr802 is a downstream effector of calcineurin in the context of host infection. Pmc1 is also regulated by Crz1 (55).

C. neoformans dissemination to the brain is the main driver of patient mortality (2). We found that dissemination of *pdr802* cells is significantly impaired, although they do occasionally reach the brain. These observations can be explained by a combination of factors. First, the limited accumulation of the *pdr802* mutant in the lungs, discussed above, may directly affect dissemination (88). Second, this strain survives poorly in mouse serum, as demonstrated directly by our culture experiments and indirectly by our inability to detect it in the blood of infected mice (even 75 days after infection). This might occur because the cells do not reach the blood or because they are rapidly eliminated, consistent with previous observations (89). Third, the thick capsules of the *pdr802* mutant

reduce its ability to reach the central nervous system. This is true whether fungal brain

entry occurs directly, by the movement of free fungi across the BBB, or indirectly, via a

Trojan horse mechanism that requires macrophage uptake (12); such uptake is impeded

by enlarged capsules, independent of cell size (29). Fourth, inositol transporters, two of

which we found to be regulated by Pdr802, are required for normal *C. neoformans*

dissemination to the brain (76). Notably, expression of one of these, *ITR2*, is upregulated

during *C. neoformans* engulfment by macrophages (2.4-fold), like *PDR802* (3-fold),

although the other, *ITR3c*, was not differentially expressed under these conditions (73).

Interestingly, despite all of these obstacles to dissemination, cells that do reach the brain

are able to proliferate to wild-type levels.

Titan cells are a robust and persistent morphotype of *C. neoformans* that

contribute to yeast virulence (43). We showed that cells lacking Pdr802 demonstrate

increased formation of Titan cells *in vivo* and *in vitro*, suggesting that this TF is a novel

repressor of this process. Although Titan cells enhance aspects of cryptococcal

pathogenesis (18, 90), their overproduction negatively impacts dissemination to the brain

due to their resistance to phagocytosis by macrophages (18, 43) and decreased

penetration of biological barriers (18).

Under conditions of nutrient limitation, such as those encountered during host

infection, Pdr802 regulates a variety of transmembrane transporters to moderate the

effects of this environment. These conditions also stimulate Titan cell formation (45–47).

In the absence of Pdr802, this compensation does not occur, and Titan cell formation

increases. Pdr802 may also indirectly regulate Titan cell formation by regulating other

TFs that impact this process, such as Zfc3. This protein (also known as Cqs2) is

upregulated during Titan cell formation *in vitro* (46) and is further regulated by Qsp1 (91,

92), a quorum sensing molecule that inhibits Titan cell formation (45, 46).

Our studies *in vitro*, *ex vivo*, and *in vivo* show that Pdr802 downregulates capsule,

a feature that is highly induced in the host or host-like conditions and depends on

environmental factors that include nutrient availability (93). Capsule thickness could be

modulated by Pdr802's repression of the TF Bzp4 (Figure S7B), which is downregulated

when *C. neoformans* is engulfed by macrophages (concomitant with *PDR802*

upregulation) (73). Our ChIP-Seq results also suggest that Pdr802 interacts with

sequences upstream of four other capsule-related transcription factors (Ecm2201, Mln1,

Zfc3 and Zfc14) (37, 50). Finally, dysregulation of intracellular calcium concentration also

affects capsule production (83).

Our results show that Pdr802 influences key cryptococcal phenotypes that

influence virulence, including nutrient acquisition, titanisation, and capsule production.

We have further identified multiple genes that are central in these processes and are

directly or indirectly regulated by Pdr802. Some of these targets are also regulated by

calcineurin (e.g. Had1 and Pmc1) or by another important TF, Gat201 (e.g. Ecm2201,

CNAG_05867, CNAG_01552 and CNAG_01040) (58). The expression of *PDR802* itself,

along with that of one target, *ITR3C*, is also regulated by another TF, Hob1 (65). The

crosstalk between all of these regulatory mechanisms remains to be dissected.

Nonetheless, it is evident that Pdr802 is critical for both survival in the lung and

dissemination to the brain, thus explaining its role in cryptococcal virulence.

MATERIALS AND METHODS

Strain construction and cell growth

We previously reported the *PDR802* deletion mutant (*pdr802*) in the KN99α strain background (94) that was used in this work (37). Complementation of this mutant with the wild-type gene at the native locus (*PDR802*) and construction of a strain that expresses Pdr802 with N-terminal mCherry (mCherry-Pdr802) are detailed in the Supplementary Methods. For all studies, *C. neoformans* strains were inoculated from single colonies into YPD medium (2% [wt/vol] dextrose, 2% [wt/vol] Bacto peptone and 1% [wt/vol] yeast extract in double-distilled water [ddH$_2$O]) and grown overnight at 30°C with shaking at 230 rpm before further handling as detailed below. To assess viability during growth in tissue culture medium, overnight cultures were washed with phosphate-buffered saline (PBS), diluted to 10^6 cells/ml in DMEM (Sigma, S6429), plated (1 ml/well) in triplicate in 24-well plates, and incubated at 37°C and 5% CO$_2$. At the indicated times cells were mixed thoroughly, diluted in PBS, and plated on YPD agar (YPD medium, 2% agar [wt/vol]) for assessment of colony-forming units (CFU). To assess viability during growth in mouse serum, mouse blood was collected as described under "Animal experiments" and YPD-grown cryptococcal cells (10^3) were incubated in 100 µl of serum in 96-well plates for 24 h at 37°C and 5% CO$_2$ and CFU counted as above.

Animal experiments

All animal protocols were approved by the Washington University Institutional Animal Care and Use Committee (reference 20170131) or Comissão de Ética no Uso de animais – CEUA (reference 30936), and care was taken to minimize handling and discomfort. For survival studies, groups of five 4- to 6-week-old female C57BL/6 mice (The Jackson Laboratory) were anesthetized by subcutaneous injection of 1.20 mg ketamine and 0.24

mg xylazine in 120 µl sterile water and intranasally infected with 5×10^4 cryptococcal

cells. The mice were monitored and humanely sacrificed when their weight decreased to

below 80% of initial weight, at which point organ burden was assessed. The lungs and

brains were harvested, homogenized, diluted and plated on YPD agar. The resulting CFU

were enumerated and survival differences were assessed by Kaplan-Meier analysis.

For timed organ burden studies, *C. neoformans* overnight cultures were centrifuged

(1,000 x *g* for 3 min), washed with sterile PBS, and resuspended in PBS to 1×10^6

cells/ml. Groups of three 4- to 6-week-old female C57BL/6 mice (Centro Multidisciplinar

para Investigação Biológica na Área da Ciência em Animais de Laboratório, CEMIB)

were anesthetized as above and intranasally infected with 5×10^4 cryptococcal cells, and

monitored as above. At set time points post-infection (see text), mice were sacrificed and

fungal burden was assessed from organs (as above) or blood (obtained by cardiac

puncture). Organ burden was analyzed by Kruskal-Wallis test with Dunn's multiple

comparison *post hoc* test for each day post-infection.

To assess cryptococcal viability in mouse serum, 6 BALB/c mice were anesthetized with

isoflurane and blood was collected from the retro-orbital space using a sterile capillary

tube. Collected blood was incubated at 37°C for 30 min and serum was isolated by

centrifugation at 1,000 x *g* for 15 min and then heat-inactivated at 56 °C for 30 min.

Capsule analysis

To qualitatively assess capsule thickness, strains were grown on YPD medium for 16 h

and washed with PBS, and 10^6 cells were incubated in DMEM or mouse serum for 24 h

at 37°C and 5% CO_2. After incubation, cells were fixed in 4% paraformaldehyde and

washed three times with PBS. *C. neoformans* cells were placed on glass slides and

mixed with similar volumes of India ink and the capsule was measured as previously

described (95).

For population-level capsule measurement, *C. neoformans* strains were grown overnight

in YPD, washed with PBS, and diluted to 10^6 cells/ml in DMEM. 150 µl aliquots were then

plated in quadruplicate in the middle 32 wells of a poly-L-lysine coated 96-well plate

(Fisher 655936) and incubated at 37°C and 5% CO_2. After 24 hours, the cells were

washed with PBS and incubated with 150 µl of a staining mixture (100 µg/ml Calcofluor

white to stain cell walls, 50 µg/ml of the anticapsular monoclonal antibody 302 conjugated

to Alexa Fluor 488 (Molecular probes), and 1.5% goat serum in PBS) for 30 minutes at

room temperature in the dark. The cells were washed again with PBS, fixed with 4%

formaldehyde for 10 minutes at room temperature, washed with PBS, and each well

refilled with 150 µl PBS. The cells were imaged using a BioTek Cytation 3 imager, which

automatically collected 100 images per well in a grid pattern at the well center. Image

files were prepared for analysis with the GE InCell Translator and assembled into .xdce

image stacks for analysis with the GE INCell Developer Toolbox 1.9. Cell wall and

capsule images were first filtered to remove background noise and border objects and

then cells were identified using shape-based object segmentation (3-pixel kernel, 50%

sensitivity) followed by watershed clump breaking to prevent apparent connectivity

caused by incomplete segmentation. Target linking was performed to assign each cell

wall object to one capsule object based on known 1:1 pairing and location, generating a

target set. Capsule and cell wall object diameters were calculated for each target set

(hundreds to thousands per well), and the difference between each pair of measurements

was defined as the capsule thickness. Data were normalized by the difference in capsule

thickness between uninduced and induced WT cells, which were included in each

experiment, and compared to hypercapsular (*pkr1*) (37) and hypocapsular (*ada2*) (96)

control strains in each experiment. Capsule sizes were compared by One-Way ANOVA

with Dunnett's multiple comparison *post hoc* test.

To measure capsule thickness of cryptococcal cells grown in the lungs of infected mice,

lung homogenates were filtered through a cell strainer with 40 µm pores using a syringe

plunger, fixed in 3.7% formaldehyde, and used for India ink staining and measurement as

above. For the visualization of KN99α and *PDR802* cells from mouse lungs after 18 days

of infection, the tissue was treated with 50 µg/ml DNAse I for 30 min at 37°C.

GXM immunoblotting was conducted as previously described (69). Briefly, 10^6 cells/ml

were grown in DMEM for 24 and 48 h and in CIM (0.17% [wt/vol] yeast nitrogen base

without amino acids and ammonium sulfate, 0.15% [wt/vol] asparagine, 2% [wt/vol]

glucose, 12 mM $NaHCO_3$, 35 mM MOPS [morpholinepropanesulfonic acid], pH 7.1)

medium or YPD for 24 h. Culture supernatant fractions were then resolved by gel

electrophoresis on 0.6% agarose, transferred onto nylon membranes, and probed with 1

µg/ml anti-GXM antibody 302 or 18B7.

Phenotypic assays

For stress plates, cryptococcal cells were grown overnight in YPD, washed with PBS, and

diluted to 10^7 cells/ml in PBS. Aliquots (3 µl) of 10-fold serial dilutions were spotted on

YPD or YNB agar supplemented with various stressors (sorbitol, NaCl, $CaCl_2$, LiCl,

Congo Red, Calcofluor white, caffeine, SDS, $NaNO_2$, H_2O_2 and ethanol) in the

concentrations indicated in the figures. Melanization was tested on plates made by

mixing 10 ml of 2X minimal medium (2 g/L L-asparagine, 1 g/L $MgSO_4 \cdot 7H_2O$, 6 g/L

KH_2PO_4, 2 g/L thiamine, 2 mM L-3,4-dihydroxyphenylalanine [L-DOPA] and 0.1%

dextrose was added for melanization induction or 0.5% for melanization inhibition) with

10 ml of 2% agar-water per plate. A control strain lacking the ability to melanize was used

as a control (*lac1*) (97). For the solid urease assay, 10 µl of a 10^7 cells/ml suspension in

water was plated on Christensen's urea solid media (1 g/L peptone, 1 g/L dextrose, 5 g/L

NaCl, 0.8 g/L KH_2PO_4, 1.2 g/L Na_2HPO_4, 0.012 g/L phenol red and 15 g/L agar, pH 6.8).

Plates were incubated at 30°C or 37°C.

Titan cells

508   Titan cell induction was performed in 1x PBS supplemented with 10% heat inactivated

509   Fetal Calf Serum (FCS) for 72 hours at 37°C and 5% CO_2 as recently described (47) and

510   quantified by flow cytometry as previously reported (45, 46).

Phagocytosis

J774.16 cells were prepared for uptake experiments by seeding (10^5 cells/well) in a 96-

well plate and incubating in DMEM supplemented with 10% Fetal Bovine Serum (FBS) at

37°C and 5% CO_2 for 24 h. *C. neoformans* cells were prepared for uptake experiments

by inoculating an overnight culture in YPD into either DMEM or Titan cell induction

medium (47) and growing at 37°C and 5% CO_2 for 24 or 72 h, respectively. To initiate the

study, cryptococcal cells were washed with PBS and opsonized with anti-capsular

antibody 18B7 (1 µg/ml) for 1 h at 37°C while macrophages were activated with 50 nM

phorbol myristate acetate (PMA) for 1 h at 37°C and 5% CO_2; 10^6 cryptococcal cells were

then incubated with the macrophages for 2 h at 37°C and 5% CO_2. The wells were then

washed three times with warm PBS and the macrophages lysed with 0.1% Triton in PBS

and plated for CFU as above. Fold-change in CFU was assessed by comparison to the

CFU of opsonized cells. One-Way ANOVA with Dunnett's multiple comparison *post hoc*

test was used to compare phagocytosis of *pdr802* and *PDR802* strains with that of

KN99α.

Chromatin Immunoprecipitation (ChIP)

ChIP studies were performed as previously described (37, 96). Briefly, wild type and N-

terminal-mCherry-Pdr802 strains were cultivated in DMEM for 24 hours at 37°C and 5%

CO_2. The cells were then fixed with formaldehyde, lysed by mechanical bead-beating,

and the cell debris removed by centrifugation. The supernatant fraction was sheared by

sonication, centrifuged, and an aliquot was reserved as 'Input'. The remaining material

was incubated with rabbit IgG anti-mCherry antibody (Abcam, ab213511) tethered to

protein A sepharose ('IP') or sepharose alone ('Mock') overnight at 4°C. The beads were

then washed, incubated at 65°C to reverse DNA-DNA and DNA-protein crosslinks and

the DNA recovered by phenol/chloroform/isoamyl alcohol (25:24:1) extraction, ethanol

precipitation, and resuspension in nuclease-free water.

Samples were submitted to the Washington University Genome Technology Access

Center for library preparation and DNA samples were sequenced using the Illumina

Nextseq platform. The first replicate was sequenced using paired-end 2x75-bp reads and

replicates 2 and 3 were sequenced using single-end 75-bp reads; the minimum coverage

obtained was ~16x. The quality of the reads was evaluated by FastQC (98). Fastq files

were aligned to the KN99 genome (99) using NextGenMap 0.5.3 (100). SAM files were

converted to bam, reads were sorted and indexed, and read duplicates were removed

from the final bam files using samtools (101). Samtools was also used to filter out reads

with a mapping quality lesser than 20 phreds to guarantee single alignment of the reads.

Peaks were called using MACS2 (2.1.1.20160309) (102), filtered by size (maximum

threshold 5 kb and no minimum), and annotated using Homer 4.8 (103). The significant

peaks were chosen using the cutoff of fold enrichment above 2 and adjusted $p < 0.05$.

The Pdr802 binding motif was defined using DREME (72) and Gene Ontology was

performed using Revigo (104). The read coverage of each peak was obtained using

Samtools (101).

qRT-PCR

RNA from wild-type and *pdr802* cells grown in DMEM at 37°C with 5% CO_2 for 3 or 24

hours was extracted with TriZol® reagent (Invitrogen, MA, USA) according to the

manufacturer's instructions. RNA integrity was assessed by electrophoresis on 1%

agarose and quantification was performed by absorbance analysis using a NanoDrop™

2000 spectrophotometer (Thermo Fisher Scientific). cDNAs were prepared from DNase

(Promega, WI, USA)-treated total RNA samples (290 ng) using ImProm-II reverse

transcriptase (Promega) and oligo-dT. Quantitative real-time PCR (qRT-PCR) was

performed on a Fast 7500 real-time PCR system (Applied Biosystems, MA, USA) with the

following thermal cycling conditions: 95°C for 10 min followed by 50 cycles of 95°C for 15

s, 55°C for 15 s, and 60°C for 60 s. Platinum® SYBR® green qPCR Supermix

(Invitrogen) was used as the reaction mix and supplemented with 5 pmol of each primer

(see Data Set S2, Sheet 6, for sequences) and 8 ng of cDNA template for a final volume

of 20 µl. All experiments were performed in biological triplicate and each sample was

analyzed in triplicate for each primer pair. Melting curve analysis was performed at the

end of the reaction to confirm the presence of a single PCR product. Data were

normalized to levels of *ACT1*, which was included in each set of PCR experiments.

Relative expression was determined using the $2^{-\Delta Ct}$ method.

Intracellular calcium measurement

To measure intracellular free Ca^{2+}, yeast cells were cultured overnight in YPD at 30°C with shaking, washed three times with deionized water, diluted to 10^6 cells/ml in DMEM (Sigma, S6429), plated (1 ml/well) in triplicate in 24-well plates, and incubated at 37°C and 5% CO_2 for 24 hours. At the indicated times, cells were mixed thoroughly, diluted in PBS containing 2 µM Fluo4-AM (ThermoFisher), incubated at 30°C for 30 min, and analysed using flow cytometry. The overnight culture was used as a control and treated as above.

Data availability

Data is available at the NCBI Gene Expression Omnibus (accession number GSE153134).

REFERENCES

1. Kwon-Chung KJ, Fraser JA, Doering TL, Wang ZA, Janbon G, Idnurm A, Bahn Y-S. 2014. *Cryptococcus neoformans* and *Cryptococcus gattii*, the Etiologic Agents of Cryptococcosis. Cold Spring Harb Perspect Med 4:a019760.

2. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect Dis 17:873–881.

3. Ballou ER, Johnston SA. 2017. The cause and effect of *Cryptococcus* interactions with the host. Curr Opin Microbiol 40:88–94.

4.   Sabiiti W, May RC. 2012. Mechanisms of infection by the human fungal pathogen

Cryptococcus neoformans. Future Microbiol 7:1297–1313.

5.   Chang YC, Stins MF, McCaffery MJ, Miller GF, Pare DR, Dam T, Paul-Satyasee M,

Kim KS, Kwon-Chung KJ. 2004. Cryptococcal yeast cells invade the central

nervous system via transcellular penetration of the blood-brain barrier. Infect

Immun 72:4985–4995.

6.   Huang S-H, Long M, Wu C-H, Kwon-Chung KJ, Chang YC, Chi F, Lee S, Jong A.

2011. Invasion of *Cryptococcus neoformans* into Human Brain Microvascular

Endothelial Cells Is Mediated through the Lipid Rafts-Endocytic Pathway via the

Dual Specificity Tyrosine Phosphorylation-regulated Kinase 3 (DYRK3). J Biol

Chem 286:34761–34769.

7.   Jong A, Wu C-H, Shackleford GM, Kwon-Chung KJ, Chang YC, Chen H-M,

Ouyang Y, Huang S-H. 2008. Involvement of human CD44 during *Cryptococcus*

neoformans infection of brain microvascular endothelial cells. Cell Microbiol

10:1313–1326.

8.   Maruvada R, Zhu L, Pearce D, Zheng Y, Perfect J, Kwon-Chung KJ, Kim KS. 2012.

Cryptococcus neoformans phospholipase B1 activates host cell Rac1 for traversal

across the blood-brain barrier. Cell Microbiol 14:1544–1553.

9.   Chen SHM, Stins MF, Huang S-H, Chen YH, Kwon-Chung KJ, Chang Y, Kim KS,

Suzuki K, Jong AY. 2003. *Cryptococcus neoformans* induces alterations in the

cytoskeleton of human brain microvascular endothelial cells. J Med Microbiol

52:961–970.

10. Olszewski MA, Noverr MC, Chen G-H, Toews GB, Cox GM, Perfect JR, Huffnagle

GB. 2004. Urease Expression by *Cryptococcus neoformans* Promotes

Microvascular Sequestration, Thereby Enhancing Central Nervous System

Invasion. Am J Pathol 164:1761–1771.

11. Shi M, Li SS, Zheng C, Jones GJ, Kim KS, Zhou H, Kubes P, Mody CH. 2010.

Real-time imaging of trapping and urease-dependent transmigration of

Cryptococcus neoformans in mouse brain. J Clin Invest 120:1683–1693.

12. Santiago-Tirado FH, Onken MD, Cooper JA, Klein RS, Doering TL. 2017. Trojan

Horse Transit Contributes to Blood-Brain Barrier Crossing of a Eukaryotic

Pathogen. MBio 8:e02183-16.

13. Jarvis JN, Bicanic T, Loyse A, Namarika D, Jackson A, Nussbaum JC, Longley N,

Muzoora C, Phulusa J, Taseera K, Kanyembe C, Wilson D, Hosseinipour MC,

Brouwer AE, Limmathurotsakul D, White N, van der Horst C, Wood R, Meintjes G,

Bradley J, Jaffar S, Harrison T. 2014. Determinants of Mortality in a Combined

Cohort of 501 Patients With HIV-Associated Cryptococcal Meningitis: Implications

for Improving Outcomes. Clin Infect Dis 58:736–745.

14. John R. Perfect, William E. Dismukes, Francoise Dromer, David L. Goldman, John

R. Graybill, Richard J. Hamill, Thomas S. Harrison, Robert A. Larsen, Olivier

Lortholary, Minh-Hong Nguyen, Peter G. Pappas, William G. Powderly, Nina Singh,

Jack D. Sobel TCS. 2010. Clinical Practice Guidelines for the Management of

Cryptococcal Disease: 2010 Update by the Infectious Diseases Society of America.

Chinese J Infect Chemother 50:291–322.

15. Nosanchuk JD, Casadevall A. 2003. The contribution of melanin to microbial

pathogenesis. Cell Microbiol 5:203–223.

16.   Singh A, Panting RJ, Varma A, Saijo T, Waldron KJ, Jong A, Ngamskulrungroj P,

Chang YC, Rutherford JC, Kwon-Chung KJ. 2013. Factors Required for Activation

of Urease as a Virulence Determinant in *Cryptococcus neoformans*. MBio

4:e00220-13.

17.   Chang YC, Kwon-Chung KJ. 1994. Complementation of a capsule-deficient

mutation of *Cryptococcus neoformans* restores its virulence. Mol Cell Biol 14:4912–

4919.

18.   Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chrétien F, Heitman J,

Dromer F, Nielsen K. 2010. Cryptococcal Cell Morphology Affects Host Cell

Interactions and Pathogenicity. PLoS Pathog 6:e1000953.

19.   Zaragoza O, Nielsen K. 2013. Titan cells in *Cryptococcus neoformans*: cells with a

giant impact. Curr Opin Microbiol 16:409–413.

20.   Wang Y, Aisen P, Casadevall A. 1995. *Cryptococcus neoformans* melanin and

virulence: Mechanism of action. Infect Immun 63:3131–3136.

21.   Mednick AJ, Nosanchuk JD, Casadevall A. 2005. Melanization of *Cryptococcus*

neoformans Affects Lung Inflammatory Responses during Cryptococcal Infection.

Infect Immun 73:2012–2019.

22.   Wang Y, Casadevall A. 1994. Susceptibility of melanized and nonmelanized

Cryptococcus neoformans to nitrogen- and oxygen-derived oxidants. Infect Immun

62:3004–7.

23.   Doering TL, Nosanchuk JD, Roberts WK, Casadevall A. 1999. Melanin as a

potential cryptococcal defence against microbicidal proteins. Med Mycol 37:175–

81.

24.   Cox GM, Mukherjee J, Cole GT, Casadevall A, Perfect JR. 2000. Urease as a

virulence factor in experimental cryptococcosis. Infect Immun 68:443–8.

25.   Bose I, Reese AJ, Ory JJ, Janbon G, Doering TL. 2003. A Yeast under Cover: the

Capsule of *Cryptococcus neoformans*. Eukaryot Cell 2:655–663.

26.   Wang ZA, Li LX, Doering TL. 2018. Unraveling synthesis of the cryptococcal cell

wall and capsule. Glycobiology 28:719–730.

27.   Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: the outer

layers of *Cryptococcus neoformans*. Mem Inst Oswaldo Cruz 113:e180040.

28.   Bulmer GS, Sans MD. 1967. *Cryptococcus neoformans*. II. Phagocytosis by human

leukocytes. J Bacteriol 94:1480–3.

29.   Bulmer GS, Sans MD. 1968. *Cryptococcus neoformans*. 3. Inhibition of

phagocytosis. 95:5–8.

30.   Tacker JR, Farhi F, Bulmer GS. 1972. Intracellular fate of *Cryptococcus

neoformans*. Infect Immun 6:162–7.

31.   Monari C, Bistoni F, Vecchiarelli A. 2006. Glucuronoxylomannan exhibits potent

immunosuppressive properties. FEMS Yeast Res 6:537–542.

32.   Hayes JB, Sircy LM, Heusinkveld LE, Ding W, Leander RN, McClelland EE, Nelson

DE. 2016. Modulation of Macrophage Inflammatory Nuclear Factor κB (NF-κB)

Signaling by Intracellular *Cryptococcus neoformans*. J Biol Chem 291:15614–

15627.

33. Fonseca FL, Nohara LL, Cordero RJB, Frases S, Casadevall A, Almeida IC,

Nimrichter L, Rodrigues ML. 2010. Immunomodulatory Effects of Serotype B

Glucuronoxylomannan from *Cryptococcus gattii* Correlate with Polysaccharide

Diameter. Infect Immun 78:3861–3870.

34. Garcia-Hermoso D, Dromer F, Janbon G. 2004. *Cryptococcus neoformans* Capsule

Structure Evolution *In Vitro* and during Murine Infection. Infect Immun 72:3359–

3365.

35. Guimarães AJ, Frases S, Cordero RJB, Nimrichter L, Casadevall A, Nosanchuk

JD. 2010. *Cryptococcus neoformans* responds to mannitol by increasing capsule

size *in vitro* and *in vivo*. Cell Microbiol 12:740–753.

36. Cordero RJB, Pontes B, Guimarães AJ, Martinez LR, Rivera J, Fries BC,

Nimrichter L, Rodrigues ML, Viana NB, Casadevall A. 2011. Chronological Aging Is

Associated with Biophysical and Chemical Changes in the Capsule of

Cryptococcus neoformans. Infect Immun 79:4990–5000.

37. Maier EJ, Haynes BC, Gish SR, Wang ZA, Skowyra ML, Marulli AL, Doering TL,

Brent MR. 2015. Model-driven mapping of transcriptional networks reveals the

circuitry and dynamics of virulence regulation. Genome Res 25:690–700.

38. Silveira CP, Piffer AC, Kmetzsch L, Fonseca FL, Soares DA, Staats CC, Rodrigues

ML, Schrank A, Vainstein MH. 2013. The heat shock protein (Hsp) 70 of

Cryptococcus neoformans is associated with the fungal cell surface and influences

the interaction between yeast and host cells. Fungal Genet Biol 60:53–63.

39.   Santos JRA, Holanda RA, Frases S, Bravim M, Araujo GDS, Santos PC, Costa

MC, Ribeiro MJA, Ferreira GF, Baltazar LM, Miranda AS, Oliveira DB, Santos

CMA, Fontes ACL, Gouveia LF, Resende-Stoianoff MA, Abrahão JS, Teixeira AL,

Paixão TA, Souza DG, Santos DA. 2014. Fluconazole Alters the Polysaccharide

Capsule of *Cryptococcus gattii* and Leads to Distinct Behaviors in Murine

Cryptococcosis. PLoS One 9:e112669.

40.   Alspaugh JA. 2015. Virulence mechanisms and *Cryptococcus neoformans*

pathogenesis. Fungal Genet Biol 78:55–58.

41.   García-Rodas R, de Oliveira H, Trevijano-Contador N, Zaragoza O. 2018.

Cryptococcal Titan Cells: When Yeast Cells Are All Grown up, p. 101–120. *In*

Current Topics in Microbiology and Immunology. Springer Verlag.

42.   Zhou X, Ballou ER. 2018. The *Cryptococcus neoformans* Titan Cell: From *In Vivo*

Phenomenon to *In Vitro* Model. Curr Clin Microbiol Reports 5:252–260.

43.   Zaragoza O, García-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodríguez-

Tudela JL, Casadevall A. 2010. Fungal Cell Gigantism during Mammalian Infection.

PLoS Pathog 6:e1000945.

44.   Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, Berman J,

Nielsen K. 2015. Polyploid Titan Cells Produce Haploid and Aneuploid Progeny To

Promote Stress Adaptation. MBio 6:e01340-15.

45.   Hommel B, Mukaremera L, Cordero RJBB, Coelho C, Desjardins CA, Sturny-

Leclère A, Janbon G, Perfect JR, Fraser JA, Casadevall A, Cuomo CA, Dromer F,

Nielsen K, Alanio A. 2018. Titan cells formation in *Cryptococcus neoformans* is

finely tuned by environmental conditions and modulated by positive and negative

genetic regulators. PLOS Pathog 14:e1006982.

46. Trevijano-Contador N, de Oliveira HC, García-Rodas R, Rossi SA, Llorente I,

Zaballos Á, Janbon G, Ariño J, Zaragoza Ó. 2018. *Cryptococcus neoformans* can

form titan-like cells *in vitro* in response to multiple signals. PLOS Pathog

14:e1007007.

47. Dambuza IM, Drake T, Chapuis A, Zhou X, Correia J, Taylor-Smith L, LeGrave N,

Rasmussen T, Fisher MC, Bicanic T, Harrison TS, Jaspars M, May RC, Brown GD,

Yuecel R, MacCallum DM, Ballou ER. 2018. The *Cryptococcus neoformans* Titan

cell is an inducible and regulated morphotype underlying pathogenesis. PLOS

Pathog 14:e1006978.

48. García-Barbazán I, Trevijano-Contador N, Rueda C, de Andrés B, Pérez-Tavárez

R, Herrero-Fernández I, Gaspar ML, Zaragoza O. 2016. The formation of titan cells

in *Cryptococcus neoformans* depends on the mouse strain and correlates with

induction of Th2-type responses. Cell Microbiol 18:111–124.

49. Mukaremera L, Lee KK, Wagener J, Wiesner DL, Gow NAR, Nielsen K. 2018. Titan

cell production in *Cryptococcus neoformans* reshapes the cell wall and capsule

composition during infection. Cell Surf 1:15–24.

50. Jung K-W, Yang D-H, Maeng S, Lee K-T, So Y-S, Hong J, Choi J, Byun H-J, Kim

H, Bang S, Song M-H, Lee J-W, Kim MS, Kim S-Y, Ji J-H, Park G, Kwon H, Cha S,

Meyers GL, Wang LL, Jang J, Janbon G, Adedoyin G, Kim T, Averette AK,

Heitman J, Cheong E, Lee Y-H, Lee Y-W, Bahn Y-S. 2015. Systematic functional

profiling of transcription factor networks in *Cryptococcus neoformans*. Nat Commun

6:6757.

51.   Schneider R de O, Fogaça N de SS, Kmetzsch L, Schrank A, Vainstein MH, Staats

CC. 2012. Zap1 Regulates Zinc Homeostasis and Modulates Virulence in

Cryptococcus gattii. PLoS One 7:e43773.

52.   Garcia-Santamarina S, Festa RA, Smith AD, Yu C-H, Probst C, Ding C, Homer

CM, Yin J, Noonan JP, Madhani H, Perfect JR, Thiele DJ. 2018. Genome-wide

analysis of the regulation of Cu metabolism in *Cryptococcus neoformans*. Mol

Microbiol 108:473–494.

53.   Lev S, Kaufman-Francis K, Desmarini D, Juillard PG, Li C, Stifter SA, Feng CG,

Sorrell TC, Grau GER, Bahn Y-S, Djordjevic JT. 2017. Pho4 Is Essential for

Dissemination of *Cryptococcus neoformans* to the Host Brain by Promoting

Phosphate Uptake and Growth at Alkaline pH. mSphere 2:e00381-16.

54.   Jung WH, Sham A, White R, Kronstad JW. 2006. Iron Regulation of the Major

Virulence Factors in the AIDS-Associated Pathogen *Cryptococcus neoformans*.

PLoS Biol 4:e410.

55.   Chow EWL, Clancey SA, Billmyre RB, Averette AF, Granek JA, Mieczkowski P,

Cardenas ME, Heitman J. 2017. Elucidation of the calcineurin-Crz1 stress

response transcriptional network in the human fungal pathogen *Cryptococcus*

neoformans. PLOS Genet 13:e1006667.

56.   O'Meara TR, Norton D, Price MS, Hay C, Clements MF, Nichols CB, Alspaugh JA.

2010. Interaction of *Cryptococcus neoformans* Rim101 and Protein Kinase A

Regulates Capsule. PLoS Pathog 6:e1000776.

57.   Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, Li LX,

769   Williams M, Crouch EC, Khader SA, Brent MR, Doering TL. 2016. Computational

770   Analysis Reveals a Key Regulator of Cryptococcal Virulence and Determinant of

771   Host Response. MBio 7:e00313-16.

772   58.   Chun CD, Brown JCS, Madhani HD. 2011. A Major Role for Capsule-Independent

773   Phagocytosis-Inhibitory Mechanisms in Mammalian Infection by *Cryptococcus*

774   *neoformans*. Cell Host Microbe 9:243–251.

775   59.   Moranova Z, Virtudazo E, Hricova K, Ohkusu M, Kawamoto S, Husickova V,

776   Raclavsky V. 2014. The CRZ1/SP1-like gene links survival under limited aeration,

777   cell integrity and biofilm formation in the pathogenic yeast *Cryptococcus*

778   *neoformans*. Biomed Pap 158:212–220.

779   60.   Lev S, Desmarini D, Chayakulkeeree M, Sorrell TC, Djordjevic JT. 2012. The

780   Crz1/Sp1 Transcription Factor of *Cryptococcus neoformans* Is Activated by

781   Calcineurin and Regulates Cell Wall Integrity. PLoS One 7:e51403.

782   61.   Park H-S, Chow EWL, Fu C, Soderblom EJ, Moseley MA, Heitman J, Cardenas

783   ME. 2016. Calcineurin Targets Involved in Stress Survival and Fungal Virulence.

784   PLOS Pathog 12:e1005873.

785   62.   Lee D, Jang E-H, Lee M, Kim S-W, Lee Y, Lee K-T, Bahn Y-S. 2019. Unraveling

786   Melanin Biosynthesis and Signaling Networks in *Cryptococcus neoformans*. MBio

787   10:e02267-19.

788   63.   Rhodes J, Desjardins CA, Sykes SM, Beale MA, Vanhove M, Sakthikumar S, Chen

789   Y, Gujja S, Saif S, Chowdhary A, Lawson DJ, Ponzio V, Colombo AL, Meyer W,

Engelthaler DM, Hagen F, Illnait-Zaragozi MT, Alanio A, Vreulink J-M, Heitman J,

Perfect JR, Litvintseva AP, Bicanic T, Harrison TS, Fisher MC, Cuomo CA. 2017.

Tracing Genetic Exchange and Biogeography of *Cryptococcus neoformans* var.

grubii at the Global Population Level. Genetics 207:327–346.

64. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic

Genetic Analysis of Virulence in the Human Fungal Pathogen *Cryptococcus*

neoformans. Cell 135:174–188.

65. Lee K-T, Hong J, Lee D-G, Lee M, Cha S, Lim Y-G, Jung K-W, Hwangbo A, Lee Y,

Yu S-J, Chen Y-L, Lee J-S, Cheong E, Bahn Y-S. 2020. Fungal kinases and

transcription factors regulating brain infection in *Cryptococcus neoformans*. Nat

Commun 11:1521.

66. Krysan DJ, Zhai B, Beattie SR, Misel KM, Wellington M, Lin X. 2019. Host Carbon

Dioxide Concentration Is an Independent Stress for *Cryptococcus neoformans* That

Affects Virulence and Antifungal Susceptibility. MBio 10:e01410-19.

67. D'Souza CA, Alspaugh JA, Yue C, Harashima T, Cox GM, Perfect JR, Heitman J.

2001. Cyclic AMP-Dependent Protein Kinase Controls Virulence of the Fungal

Pathogen *Cryptococcus neoformans*. Mol Cell Biol 21:3179–3191.

68. Hu G, Steen BR, Lian T, Sham AP, Tam N, Tangen KL, Kronstad JW. 2007.

Transcriptional Regulation by Protein Kinase A in *Cryptococcus neoformans*. PLoS

Pathog 3:e42.

69. Yoneda A, Doering TL. 2008. Regulation of *Cryptococcus neoformans* Capsule

Size Is Mediated at the Polymer Level. Eukaryot Cell 7:546–549.

70.   McFadden DC, Fries BC, Wang F, Casadevall A. 2007. Capsule Structural

813   Heterogeneity and Antigenic Variation in *Cryptococcus neoformans*. Eukaryot Cell

814   6:1464–1473.

815   71.   McFadden DC, De Jesus M, Casadevall A. 2006. The Physical Properties of the

816   Capsular Polysaccharides from *Cryptococcus neoformans* Suggest Features for

817   Capsule Construction. J Biol Chem 281:1868–1875.

818   72.   Bailey TL. 2011. DREME: Supp. Bioinformatics 27:1653–1659.

819   73.   Derengowski L da S, Paes HC, Albuquerque P, Tavares AHFP, Fernandes L,

820   Silva-Pereira I, Casadevall A. 2013. The Transcriptional Response of *Cryptococcus*

821   *neoformans* to Ingestion by *Acanthamoeba castellanii* and Macrophages Provides

822   Insights into the Evolutionary Adaptation to the Mammalian Host. Eukaryot Cell

823   12:761–774.

824   74.   Chikamori M, Fukushima K. 2005. A new hexose transporter from *Cryptococcus*

825   *neoformans*: Molecular cloning and structural and functional characterization.

826   Fungal Genet Biol 42:646–655.

827   75.   Liu T-B, Wang Y, Baker GM, Fahmy H, Jiang L, Xue C. 2013. The Glucose Sensor-

828   Like Protein Hxs1 Is a High-Affinity Glucose Transporter and Required for

829   Virulence in *Cryptococcus neoformans*. PLoS One 8:e64239.

830   76.   Liu T-B, Kim J-C, Wang Y, Toffaletti DL, Eugenin E, Perfect JR, Kim KJ, Xue C.

831   2013. Brain Inositol Is a Novel Stimulator for Promoting *Cryptococcus* Penetration

832   of the Blood-Brain Barrier. PLoS Pathog 9:e1003247.

77.   Xue C, Liu T, Chen L, Li W, Liu I, Kronstad JW, Seyfang A, Heitman J. 2010. Role

of an Expanded Inositol Transporter Repertoire in *Cryptococcus neoformans*

Sexual Reproduction and Virulence. MBio 1:e.00084-10.

78. Lee K-T, So Y-S, Yang D-H, Jung K-W, Choi J, Lee D-G, Kwon H, Jang J, Wang

LL, Cha S, Meyers GL, Jeong E, Jin J-H, Lee Y, Hong J, Bang S, Ji J-H, Park G,

Byun H-J, Park SW, Park Y-M, Adedoyin G, Kim T, Averette AF, Choi J-S, Heitman

J, Cheong E, Lee Y-H, Bahn Y-S. 2016. Systematic functional analysis of kinases

in the fungal pathogen *Cryptococcus neoformans*. Nat Commun 7:12766.

79. Kozubowski L, Lee SC, Heitman J. 2009. Signalling pathways in the pathogenesis

of *Cryptococcus*. Cell Microbiol 11:370–380.

80. Fox DS, Cruz MC, Sia RAL, Ke H, Cox GM, Cardenas ME, Heitman J. 2001.

Calcineurin regulatory subunit is essential for virulence and mediates interactions

with FKBP12-FK506 in *Cryptococcus neoformans*. Mol Microbiol 39:835–849.

81. Odom A. 1997. Calcineurin is required for virulence of *Cryptococcus neoformans*.

EMBO J 16:2576–2589.

82. Jung W-H, Son Y-E, Oh S-H, Fu C, Kim HS, Kwak J-H, Cardenas ME, Heitman J,

Park H-S. 2018. Had1 Is Required for Cell Wall Integrity and Fungal Virulence in

Cryptococcus neoformans. Genes|Genomes|Genetics 8:643–652.

83. Kmetzsch L, Staats CC, Cupertino JB, Fonseca FL, Rodrigues ML, Schrank A,

Vainstein MH. 2013. The calcium transporter Pmc1 provides Ca 2+ tolerance and

influences the progression of murine cryptococcal infection. FEBS J 280:4853–

4864.

84. Squizani ED, Oliveira NK, Reuwsaat JCV, Marques BM, Lopes W, Gerber AL, de

Vasconcelos ATR, Lev S, Djordjevic JT, Schrank A, Vainstein MH, Staats CC,

Kmetzsch L. 2018. Cryptococcal dissemination to the central nervous system

requires the vacuolar calcium transporter Pmc1. Cell Microbiol 20:e12803.

85. Chang AL, Kang Y, Doering TL. 2019. Cdk8 and Ssn801 Regulate Oxidative Stress

Resistance and Virulence in *Cryptococcus neoformans*. MBio 10:e02818-18.

86. Watkins R, King J, Johnston S. 2017. Nutritional Requirements and Their

Importance for Virulence of Pathogenic *Cryptococcus* Species. Microorganisms

5:65.

87. Cruz MC. 2001. Calcineurin is required for hyphal elongation during mating and

haploid fruiting in *Cryptococcus neoformans*. EMBO J 20:1020–1032.

88. Denham S, Brown J. 2018. Mechanisms of Pulmonary Escape and Dissemination

by *Cryptococcus neoformans*. J Fungi 4:25.

89. Sun D, Sun P, Li H, Zhang M, Liu G, Strickland AB, Chen Y, Fu Y, Xu J, Yosri M,

Nan Y, Zhou H, Zhang X, Shi M. 2019. Fungal dissemination is limited by liver

macrophage filtration of the blood. Nat Commun 10:4566.

90. Okagaki LH, Nielsen K. 2012. Titan Cells Confer Protection from Phagocytosis in

Cryptococcus neoformans Infections. Eukaryot Cell 11:820–826.

91. Summers DK, Perry DS, Rao B, Madhani HD. 2020. Coordinate genomic

association of transcription factors controlled by an imported quorum sensing

peptide in *Cryptococcus neoformans*. bioRxiv 2020.03.30.016329.

92. Tian X, He G-J, Hu P, Chen L, Tao C, Cui Y-L, Shen L, Ke W, Xu H, Zhao Y, Xu Q,

Bai F, Wu B, Yang E, Lin X, Wang L. 2018. *Cryptococcus neoformans* sexual

reproduction is controlled by a quorum sensing peptide. Nat Microbiol 3:698–707.

93. Srikanta D, Santiago-Tirado FH, Doering TL. 2014. *Cryptococcus neoformans* :

historical curiosity to modern pathogen. Yeast 31:47–60.

94. Nielsen K, Cox GM, Wang P, Toffaletti DL, Perfect JR, Heitman J. 2003. Sexual

Cycle of *Cryptococcus neoformans* var. *grubii* and Virulence of Congenic a and α

Isolates. Infect Immun 71:4831–4841.

95. Reuwsaat JCV, Motta H, Garcia AWA, Vasconcelos CB, Marques BM, Oliveira NK,

Rodrigues J, Ferrareze PAG, Frases S, Lopes W, Barcellos VA, Squizani ED,

Horta JA, Schrank A, Rodrigues ML, Staats CC, Vainstein MH, Kmetzsch L. 2018.

A Predicted Mannoprotein Participates in *Cryptococcus gattii* Capsular Structure.

mSphere 3: e00023-18.

96. Haynes BC, Skowyra ML, Spencer SJ, Gish SR, Williams M, Held EP, Brent MR,

Doering TL. 2011. Toward an Integrated Model of Capsule Regulation in

Cryptococcus neoformans. PLoS Pathog 7:e1002411.

97. Walton FJ, Idnurm A, Heitman J. 2005. Novel gene functions required for

melanization of the human pathogen *Cryptococcus neoformans*. Mol Microbiol

57:1381–1396.

98.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput

Sequence Data.

99. Friedman RZ, Gish SR, Brown H, Brier L, Howard N, Doering TL, Brent MR. 2018.

Unintended Side Effects of Transformation Are Very Rare in *Cryptococcus*

neoformans. Genes|Genomes|Genetics 8:815–822.

100. Sedlazeck FJ, Rescheneder P, von Haeseler A. 2013. NextGenMap: fast and

901   accurate read mapping in highly polymorphic genomes. Bioinformatics 29:2790–1.

902   101. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

903   Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence

904   Alignment/Map format and SAMtools. Bioinformatics 25:2078–2079.

905   102. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C,

906   Myers RM, Brown M, Li W, Liu XS. 2008. Model-based Analysis of ChIP-Seq

907   (MACS). Genome Biol 9:R137.

908   103. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C,

909   Singh H, Glass CK. 2010. Simple combinations of lineage-determining transcription

910   factors prime cis-regulatory elements required for macrophage and B cell identities.

911   Mol Cell 38:576–89.

912   104. Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO Summarizes and

913   Visualizes Long Lists of Gene Ontology Terms. PLoS One 6:e21800.

914   105. Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. Xylose donor transport

915   is critical for fungal virulence. PLOS Pathog 14:e1006765.

ACKNOWLEDGMENTS

We appreciate helpful discussions with Michael Brent, Charley Christian Staats, and

members of the Doering, Kmetzsch and Brent labs. We are grateful to Thomas Hurtaux,

Eamim Squizani, and Julia Sperotto for assistance with mouse experiments, Guohua

Chen for helping with spotting assays, and Jessica Plaggenberg for assistance with

library preparation and sequencing. We thank Marilene Henning Vainstein and her lab for

providing support with *in vivo* experiments. We also thank Liza Miller for comments on

the manuscript and Arturo Casadevall for providing the antibody anti-GXM (18B7).

These studies were supported by National Institutes of Health grants AI087794, AI36688,

and AI140979 to TLD and grants from Coordenação de Aperfeiçoamento de Pessoal de

Nível Superior (CAPES, Brazil), Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq, Brazil – Grant number 310510/2018-0), and Fundação de Amparo à

Pesquisa do Estado do Rio Grande do Sul (FAPERGS, Brazil) to LK. CAPES fully

supported JCVR during her studies in Brazil; her studies in the United States were

partially supported by this source (Advanced Network of Computational Biology -

RABICÓ - Biocomputational Grant 23038.010041/2013-13).

AUTHORS CONTRIBUTION

Conceived and designed the experiments: J.C.V.R., A.L.C., L.K. and T.L.D.;

Performed the experiments: J.C.V.R., H.M., H.B., and A.L.C;

Analyzed the data: J.C.V.R., A.L.C., D.P.A., L.K. and T.L.D.;

Contributed reagents and materials: L.K. and T.L.D.;

Wrote the paper: J.C.V.R, L.K. and T.L.D.

COMPETING INTERESTS

The authors declare no competing financial interests.

FIGURE LEGENDS

Figure 1. The transcription factor Pdr802 influences *C. neoformans* virulence. A.

Survival of C57BL/6 mice over time after intranasal inoculation with 5 x 10^4 cryptococci,

with sacrifice triggered by weight below 80% of peak. B. Mean +/- SD of total colony-

forming units (CFU) in lung tissue at various times post-infection. CFU inoculated for

each strain were 41,400 (KN99α), 42,800 (*pdr802*) and 26,600 (*PDR802*). p<0.05 for

pdr802 compared to the other strains at all time points.

Figure 2. *PDR802* expression is required for cell viability and induced during

growth in host-like conditions. A. Cells grown in DMEM at 37°C and 5% CO_2 were

sampled at the times indicated and plated on YPD to assess viability (measured by CFU

and plotted as fold-change from time 0). B. *PDR802* expression in KN99α cells grown in

DMEM at 37°C and 5% CO_2 was assessed by RNA-seq as in Li *et al.*, 2018 (105).

Figure 3. The *pdr802* mutant is hypercapsular. A. Representative

immunofluorescence micrographs of the indicated strains after growth in DMEM (37°C,

5% CO_2) for 24 hours. The capsule was stained with monoclonal antibody anti-GXM 302

conjugated with Alexa 488 (green) and the cell wall with Calcofluor White (blue). All

images are to the same scale; scale bar, 5 μm. B. Capsule thickness distribution for the

indicated strains. C. Mean +/- SD of capsule size, quantified as detailed in the Methods

and Figure S4, with *pkr1* (37) and *ada2* (96) shown as hypercapsular and hypocapsular

controls, respectively. ****, p<0.0001 compared to KN99α by one-way ANOVA with

posthoc Dunnett test.

Figure 4. Growth in mouse serum elicits increased capsule and cell   wall

diameter in the *pdr802* mutant. A. Light micrographs of the indicated strains after

growth in mouse serum (at 37°C, 5% CO_2) for 24 h and negative staining with India ink to

visualize the capsule. All images are to the same scale; scale bar, 5 μm. B. Mean +/- SD

of capsule thickness, assessed by measuring at least 50 cells per strain with ImageJ. C.

Cells grown as in Panel A were plated on YPD to assess CFU. Mean +/- SD of the fold-

change compared to 0 h is shown. D. Mean +/- SD of cell wall diameter, measured as in

B. ***, p<0.001 and ****, p<0.0001 for comparison of *pdr802* results to KN99α by one-

way ANOVA with posthoc Dunnett test.

Figure 5. Absence of *PDR802* yields enlarged cells and loss of capsule induction in

the context of animal infection. A. India ink staining of fungi isolated from the lungs of

mice infected with the indicated strains for various times (3, 7, 14 and 18 days). All

images are to the same scale; scale bar, 10 μm. B and C. Mean +/- SD of cell wall

diameter (B) and capsule thickness (C), assessed by measuring at least 50 cells per

strain with ImageJ. ****, p<0.0001 and ***, p<0.001 for comparison of *pdr802* results to

KN99α or *PDR802* by one-way ANOVA with posthoc Dunnett test for each day post-

infection.

Figure 6. Pdr802 is a negative regulator of titanisation. Left, cultures were subjected

to *in vitro* conditions that induce Titan cell formation and imaged with India Ink. All images

are to the same scale; scale bar, 10 µm. Images were selected so that each shows

multiple examples of Titan cells, not to reflect abundance of this morphotype. Right, the

percent of Titan cells (TC) in each culture was quantified using flow cytometry, gated as

indicated by the blue square. FSC, forward scatter; SSC, side scatter.

Figure 7. Deletion of *PDR802* affects phagocytosis after growth under conditions

that induce capsule and titan cell formation. *C. neoformans* strains were grown in

YPD (18 h), DMEM (24 h), or Titan-cell induction medium (72 h) and then incubated for 2

h with J774.16 mouse macrophages; host cells were then washed and lysed to assess

fungal burden by CFU. Data shown are normalized to the CFU of the initial inoculum. *,

$p < 0.01$ and **, $p < 0.001$ compared to KN99α by one-way ANOVA with posthoc Dunnett

test.

1001

Figure 8. Interactions of Pdr802 with specific gene promoters. Read coverage for

samples immunoprecipitated from tagged (light gray) and untagged (dark gray) strains.

The X-axis shows the sequence position (in bp) plotted relative to the first position of the

coding sequence (+1) for genes discussed in the text that encode transporters (A),

transcription factors (B), and Frk102 (C). Peaks were plotted using R software (ggplot2

package).

1008

Figure 9. Pdr802 participates in calcineurin signaling. A. ChIP-Seq results for the

HAD1 promoter, plotted as in Figure 8. B, qRT-PCR analysis of *HAD1* expression in the

KN99α and *pdr802* strains after growth for 3 or 24 hours in DMEM (37°C, 5% CO_2).

Results are from biological triplicates, normalized to *ACT1* expression. **, $p<0.01$ by one-

way ANOVA with posthoc Dunnett test. C. ChIP-Seq results for the *PMC1* promoter, as in

A. D. Analysis of *PMC1* expression as in B. *, $p<0.05$ by one-way ANOVA with posthoc

Dunnett test. E. Intracellular calcium measurement by flow cytometry using Fluo-4AM.

Each column shows the mean and standard deviation of three biological replicates. ***,

$p<0.001$ and ****, $p<0.0001$ by one-way ANOVA with posthoc Dunnett test.

1018

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Mutant strain construction and confirmation. A. Scheme for generating *C.*

neoformans strains in the KN99α background (middle) that either lack *PDR802* (*pdr802,*

top) or encode a tagged copy of the protein (*mCherry-PDR802*). B. Qualitative analysis of

gene expression in Panel A strains and the complemented *pdr802* mutant (*PDR802*)*.*

Cryptococcal mRNA isolated from cells grown in DMEM (37°C, 5% CO_2, 24 hours) was

used to generate cDNA; from this, segments of the genes indicated at the left were

amplified using the primers listed in Data Set S2, Sheet 6, and the products were

analyzed by agarose gel electrophoresis. Fragment sizes (in bp) are indicated and the

ladder bands shown are 400, 500, 650, 850 and 1000 bp for the top panel; 200, 300, 400,

500, and 650 bp for the middle panel; and 100, 200, and 300 bp for the bottom panel. C.

Quantitative analysis of *PDR802* expression. Samples of RNA isolated as in B were

analyzed for *PDR802* expression by qRT-PCR. All results were normalized to *ACT1*

expression. Each symbol represents a biological replicate, with the mean and standard

deviation also shown. ***, p<0.001 compared to KN99α by one-way ANOVA with posthoc

Dunnett test.

1035

Figure S2. Organ burdens. Mean +/- SD values of total colony-forming units (CFU) in

the indicated tissue of mice from the Figure 1 survival curve are shown. Each point is the

average value for a single animal at the time of death. For *pdr802* infections, red circles

represent mice sacrificed at days 65 and 69, while blue circles represent mice sacrificed

at the termination of the study (day 100). B. Mean +/- SD of total colony-forming units

(CFU) in the blood and brain at the indicated times post-infection. C. Mean +/- SD of total

colony-forming units (CFU) in the lung, blood and brain 75 days after infection with

pdr802. Each color represents one mouse.

1044

Figure S3. Characterization of *pdr802* cells. A-C.10-fold serial dilutions of WT, *pdr802*,

and *PDR802* cells were plated on the media shown and incubated at 30°C (A), 37°C (B),

or 37°C in the presence of 5% CO_2 (C). Nitrosative ($NaNO_2$) and oxidative (H_2O_2) stress

plates were prepared with YNB medium and melanization plates containing L-DOPA

were prepared as in the Methods; all other plates were prepared with YPD medium. *lac1*,

a control strain lacking the ability to melanize (97). D. Urease activity of the indicated

strains was evaluated using Christensen's urea solid medium (see Methods) at the

indicated temperatures. *ure1*, a control strain that does not produce urease (16). E-F.

Growth of the strains indicated in YPD at 30°C (E) or DMEM at 37°C and 5% CO_2 (F) was

assessed by OD_{600nm} at the times indicated. G. Conditioned medium from the indicated

strains was probed for the presence of GXM after growth in DMEM for 24 or 48 hours.

Immunoblotting was performed using the anti-GXM monoclonal antibody 302.

1057

Figure S4. Semi-automated assay for cryptococcal capsule imaging. A. Schematic

of applying this method to cryptococcal cells induced to form capsule by growth in DMEM

(37°C, 5% CO_2) for 24 h, followed by cell wall and capsule staining. Thousands of cells

may be imaged per well and analyzed automatically with software that annotates and

measures the capsule (annotated in dark blue) and cell wall (annotated in bright green).

See Methods for details. B. Capsule size distribution of WT cells after induction. Capsule

thickness for each cell is the difference between the paired diameters of the cell wall and

capsule, which is plotted with reference to the mean value. C and D. Mean and SD (C)

and cumulative percentage (D) analysis of WT compared to hyper and hypocapsular

control strains (here *pkr1* and *ada2*, respectively). Capsule thickness is in arbitrary units,

related to the pixels measured. E. The time required to analyze the capsule thickness of

1,000 cells by this method compared to manual assessment of India ink images.

1070

Figure S5. *PDR802* **deletion induces the formation of titan cells.** Mean +/- SD of (A)

capsule thickness relative to cell wall diameter and (B) capsule diameter, assessed by

measuring at least 50 cells per strain with ImageJ. ***, p<0.001 and ****, p<0.0001 for

comparison of *pdr802* results to KN99α or *PDR802* by one-way ANOVA with posthoc

Dunnett test for each day post-infection. C. Percent of Titan cells in the indicated strain,

evaluated using various published parameters: cell wall diameter bigger than 10 or 15 µm

(19) or capsule diameter bigger than 30 µm (41).

1078

1079 **Figure S6. Viability of tagged-Pdr802 strains in DMEM.** The indicated strains were

1080 grown in DMEM at 37°C and 5% CO_2 for the times shown and samples were tested for

1081 their ability to form colonies on YPD medium. Plotted is the fold-change in CFU relative to

1082 the initial culture.

1083

Figure S7. Pdr802 DNA-binding motif and target validation. A. Pdr802-binding motifs

determined using DREME (72). Primary and secondary hits are shown for analysis of

1,000 bp upstream of the initiating ATG (top two rows) or that region plus the next 200 bp

(lower two rows). RC, reverse complement. B. Expression of the indicated genes in

KN99α (filled bars) and *pdr802* (open bars) cells, grown in host-like conditions for the

times shown. qRT-PCR results are normalized to *ACT1* expression; mean and standard

deviation of three biological replicates for each time point are shown. **, p<0.01

compared to KN99α by one-way ANOVA with posthoc Dunnett test. *CNAG_03304*

encodes a hypothetical protein and *CNAG_07402* a pantothenate transporter.

1093

Data Set S1. Pdr802 ChIP-Seq data. Data in sheets 1-3 is limited to genes that showed

>2-fold enrichment when Chip-Seq was performed on strains expressing tagged versus

untagged Pdr802. Sheet 1, annotated peaks that occur in gene promoter regions. Sheet

2, ChIP-Seq data for Sheet 1. Sheet 3, all peaks, annotated (not restricted to promoter

regions). Sheet 4, ChIP-Seq data of Pdr802-specific peaks. Sheet 5, ChIP-Seq data of all

peaks in mCherry-Pdr802 samples. Sheet 6, ChIP-Seq data of all peaks in untagged

(WT) control samples. Sheet 7, GO analysis of genes enriched >2-fold in ChIP-Seq

studies. Sheet 8, transmembrane transporters identified as Pdr802 targets by ChIP-Seq.

Data Set S2. Pdr802 binding motifs, target analysis, and primers used in this study.

Sheet 1, Pdr802 binding motifs in genes that are down-regulated during Titan cell

formation *in vitro*. Sheet 2, gene ontology analysis for genes that are differentially

expressed during Titan cell formation *in vitro*. Sheet 3, Pdr802 binding motifs in genes

that are up-regulated during Titan cell formation *in vitro*. Sheet 4, Pdr802 binding motifs in

the promoters of *HAD1* and *PMC1*. Sheet 5, the intersection of Pdr802 targets and genes

that are Crz1-independent calcineurin targets under conditions thermal stress (55). Sheet

6, primers used in this study. Fold enrichment and adjusted p (q) values throughout are

for the ChIP-Seq results.

Supplementary Methods.

A

FIGURE 8