The transcription factor Pdr802 regulates Titan cell formation, quorum sensing, and pathogenicity of *Cryptococcus neoformans*

Julia C. V. Reuwsaat[1,2], Daniel P. Agustinho[2], Heryk Motta[1], Holly Brown[2], Andrew L. Chang[2], Michael R. Brent[3], Livia Kmetzsch[1,4,5], and Tamara L. Doering[2,5].

[1]Molecular Biology of Pathogens Laboratory, Biotechnology Center, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

[2]Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA.

[3]Center for Genome Sciences and Systems Biology, Washington University School of Medicine, and Departments of Computer Science and Genetics, Washington University, St. Louis, Missouri, USA.

[4]Department of Molecular Biology and Biotechnology, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.

[5]Corresponding authors: livia.kmetzsch@ufrgs.br, doering@wustl.edu

Running head: The role of Pdr802 in *Cryptococcus neoformans* virulence.

ABSTRACT

Cryptococcus neoformans is a ubiquitous, opportunistic fungal pathogen that kills almost

200,000 people worldwide each year. It is acquired when mammalian hosts inhale the

infectious propagules; these are deposited in the lung and, in the context of

immunocompromise, may disseminate to the brain and cause lethal meningoencephalitis.

Once inside the host, *C. neoformans* undergoes a variety of adaptive processes,

including secretion of virulence factors, expansion of a polysaccharide capsule that

impedes phagocytosis, and the production of giant (Titan) cells. The transcription factor

Pdr802 is one regulator of these responses to the host environment. Expression of the

corresponding gene is highly induced under host-like conditions *in vitro* and is critical for

C. neoformans dissemination and virulence in a mouse model of infection. Direct targets

of Pdr802 include the quorum sensing proteins Pqp1, Opt1 and Liv3; the transcription

factors Stb4, Zfc3 and Bzp4, which regulate cryptococcal brain infectivity and capsule

thickness; the calcineurin targets Had1 and Crz1, important for cell wall remodeling and

C. neoformans virulence; and additional genes related to resistance to host temperature

and oxidative stress, and to urease activity. Notably, cryptococci engineered to lack

Pdr802 showed a dramatic increase in Titan cells, which are not phagocytosed and have

diminished ability to directly cross biological barriers. This explains the limited

dissemination of *pdr802* mutant cells to the central nervous system and the consequently

reduced virulence of this strain. The role of Pdr802 as a negative regulator of Titan cell

formation is thus critical for cryptococcal pathogenicity.

IMPORTANCE

The pathogenic yeast *Cryptococcus neoformans* presents a worldwide threat to human health, especially in the context of immunocompromise, and current antifungal therapy is hindered by cost, limited availability, and inadequate efficacy. After the infectious particle is inhaled, *C. neoformans* initiates a complex transcriptional program that integrates cellular responses and enables adaptation to the host lung environment. Here we describe the role of the transcription factor Pdr802 in the response to host conditions and its impact on *C. neoformans* virulence. We identified direct targets of Pdr802 and also discovered that it regulates cellular features that influence movement of this pathogen from the lung to the brain, where it causes fatal disease. These findings advance our understanding of a serious disease.

INTRODUCTION

Cryptococcosis is a fungal infection caused by *Cryptococcus neoformans* and *Cryptococcus gattii. C. neoformans* is a ubiquitous opportunistic pathogen that infects mainly immunocompromised patients, while *C. gattii* is capable of infecting immunocompetent individuals (1). Cryptococcosis causes 180,000 deaths worldwide each year, including roughly 15% of all AIDS-related deaths (2), and is initiated by the inhalation of spores or desiccated yeast cells. In immunocompetent individuals, this typically leads to an asymptomatic pulmonary infection that is controlled by the host immune response, although a population of *C. neoformans* may remain latent for extended periods of time (3–5).

Under conditions of immunocompromise, cryptococci disseminate from the lung to

the brain. Mechanisms that have been suggested to mediate fungal crossing of the

blood-brain barrier (BBB) include transcellular migration, in which the yeast cells enter

and exit vascular endothelial cells (6–9); paracellular movement, in which they cross the

BBB at junctions between endothelial cells (10–12); and 'Trojan horse' crossing, whereby

macrophages harboring *C. neoformans* enter the brain (13). Cryptococcal

meningoencephalitis is difficult to treat and frequently lethal, for reasons that include the

availability and cost of therapy (14, 15).

The ability of *C. neoformans* to survive and proliferate in the lung, and

subsequently disseminate to the brain, depends on viability at mammalian body

temperature and the expression of multiple virulence traits; these include secreted factors

(16, 17), a polysaccharide capsule that surrounds the cell wall (18), and the production of

giant (Titan) cells (19, 20). One secreted molecule, the pigment melanin, associates with

the cell wall, where its antioxidant properties protect fungal cells from reactive oxygen

species produced as a host immune defense (21–25). Urease, a secreted metalloenzyme

that converts urea to ammonia and CO_2, may affect the course of infection by modulating

environmental pH and damaging host tissue structure (11, 12, 26).

The capsule, composed primarily of large polysaccharides (27–29), is a key

cryptococcal virulence factor that impairs phagocytosis by immune cells (30–35). This

dynamic entity changes its size and structure during interactions with the host or external

environment (36–39), contributing to fungal adaptation (40, 41). Capsule polysaccharides

that are shed from the cell enable diagnosis of cryptococcal infection and also impede

host responses (35, 42).

Titan cells display a cryptococcal morphotype that has been variously

characterized as having cell body diameter (excluding the capsule) greater than 10 or 15

µm or total cell diameter (including the capsule) that exceeds 30 µm (20, 43, 44). These

cells are polyploid and produce normal-size cells during infection (19, 45, 46). Titan cell

formation is triggered by exposure to the host environment, including nutrient starvation,

reduced pH, and hypoxia (47–49), although the extent of induction depends on the host

immune response and the duration of infection (45, 50). Titan cell production appears to

benefit the development of pulmonary *C. neoformans* infection, since these large cells

are less susceptible to internalization by host phagocytes and more resistant to oxidative

stress than normal-size cells (19, 46). Some of these effects may be explained by the

highly cross-linked capsule and thickened cell wall of Titan cells (51). In contrast to their

success in the lungs, Titan cells show impaired dissemination to the brain (19, 46).

C. neoformans experiences a dramatic change in conditions upon entering a host,

including altered nutrient levels and pH. To adapt to the new environment, cryptococci

activate a network of transcription factors (TFs) (39, 52). For example, imbalances in ion

homeostasis trigger transcriptional changes mediated by the TFs Zap1 (53), Cuf1 (54),

Pho4 (55), Cir1 (56), and Crz1 (57). Alkaline pH stimulates expression of the TF Rim101,

which enables growth under basic conditions and other stresses such as high salt and

iron limitation; it also promotes the association of capsule polysaccharide with the cell

and the formation of Titan cells (47, 58).

Overlapping TF circuits regulate cryptococcal virulence determinants, including

polysaccharide capsule production and melanin synthesis. For example, Usv101, an

important regulator of capsule thickness and polysaccharide shedding, also regulates

three other TFs (Gat201, Crz1, and Rim101) and multiple polysaccharide-related

enzymes (59). Gat201 further regulates additional virulence-related transcription factors

and the anti-phagocytic protein Blp1 (60), while Crz1 plays a central role in the

maintenance of plasma membrane and cell wall stability (57, 61, 62). Crz1 expression is

also modulated by the calcineurin signaling pathway, which is required for normal yeast

growth at 37°C, virulence, and sexual reproduction (63). A group of TFs, including

Usv101, Bzp4, Hob1, and Mbs1 (59, 64), act together to regulate melanin production;

deletion of Bzp4 also alters capsule (52).

In this study, we investigated the TF Pdr802. The corresponding gene has a high

rate of non-synonymous mutations, which suggests it is evolving rapidly (65). Pdr802 has

previously been implicated in *C. neoformans* virulence (39, 52, 66), but its specific role

and targets are not known. We discovered that Pdr802 is induced in host-like conditions,

is a negative regulator of Titan cell formation, and influences capsule thickness and

phagocytosis by macrophages. It also regulates genes whose products act in cell wall

remodeling, virulence factor production, resistance to host temperature and oxidative

stress, and quorum sensing. These functions make Pdr802 critical for cryptococcal

survival in the lung and dissemination to the brain.

RESULTS

The role of Pdr802 in *C. neoformans* virulence

The importance of Pdr802 in *C. neoformans* virulence has been demonstrated in

multiple experimental models. Liu and collaborators first reported in 2008 that partial

deletion of *PDR802* reduced *C. neoformans* infectivity in a competition assay of pooled

C. neoformans strains (66). In 2015, Maier *et al* showed that a *pdr802* deletion mutant

had reduced virulence when tested individually in a short-term mouse model of infection

(39). Later that year, Jung and colleagues reported that Pdr802 was required for full

virulence in both wax moth larvae and short-term mouse infection using pooled strains

(52). Most recently, Lee and collaborators showed that Pdr802 was required for brain

infection (67).

To further investigate the role of Pdr802 in pathogenesis, we complemented a

complete deletion strain in the KN99α background that we had previously generated

(*pdr802*) (39) with the intact gene at its native locus (*PDR802*). To examine targets of

Pdr802, we also constructed a strain that expresses the protein fused to mCherry at its

N-terminus (Figure S1A). All of these strains lacked or expressed RNA encoding *PDR802*

or its modified forms as expected (Figure S1B) and *PDR802* was expressed at wild-type

levels in the complemented and modified strains (Figure S1C).

We next assessed the long-term survival of C57BL/6 mice infected with the

parental wild-type strain (KN99α), the deletion mutant (*pdr802*), or the complemented

mutant (*PDR802*). In this model, mice infected with the parent or complemented strains

survived for roughly three weeks, while those infected with the deletion mutant showed a

striking increase in survival: all animals survived for at least 65 days and over half

survived to the end of the study (100 days; Figure 1A). The lung burden measured at the

time of death for *pdr802*-infected mice in this study was approximately 100-fold lower

than that of wild type infections (Figure S2A), demonstrating the importance of this TF in

C. neoformans virulence. Mean brain burden at the time of death was more similar

between mutant and wild type infections (Figure S2A), although we did note some

heterogeneity in this measure for *pdr802*-infected mice; animals sacrificed at around two

months of infection (red symbols) showed brain burden similar to WT levels, while brain

burden of mice sacrificed at day 100 (blue symbols) ranged between zero fungal cells

and WT level.

We next examined the time course of fungal proliferation in the lungs. As

expected, the burdens of WT and the complemented mutant strains increased steadily

over an 18-day interval (Figure 1B), eventually reaching roughly 10^5 times the original

inoculum. Towards the end of this period, these cells were also detected in the blood and

brain (Figure S2B). In contrast, the lung burden of *pdr802* remained close to the inoculum

throughout this period, with no mutant cells detected in the blood or brain. At a late time

point of *pdr802* infection (75 days), we again noted some heterogeneity of fungal burden:

one mouse had high lung burden with no dissemination, another had high lung burden

with moderate brain burden, and the third had extremely low lung burden with no

dissemination (Figure S2C). No colony-forming units (CFU) were detected in the blood of

pdr802-infected mice at any point during infection. These results suggest that even

though the *pdr802* mutant is generally hypovirulent and remains at low levels in the lung,

it can occasionally reach the brain and, given enough time, accumulate there (see

Discussion).

Given the dramatic effects of Pdr802 on fungal virulence, we wondered about the

specific biological processes in which this transcription factor is involved. We first

examined the behavior of the *pdr802* strain *in vitro*, including stress conditions that might

be encountered in the host. We saw no differences in growth of the mutant compared to

WT cells under conditions that challenge cell or cell wall integrity, including the presence

of sorbitol, high salt, cell wall dyes, caffeine, sodium dodecyl sulfate (SDS), or ethanol

(Figure S3A-C). The mutant also showed no altered susceptibility to elements of the host

response, such as nitrosative or oxidative stresses, or in melanin production. All of these

results held whether growth was at 30°C, 37°C, or 37°C in the presence of 5% CO_2,

which was recently described as an independent stress for *C. neoformans* (68) (Figure

S3A-C). Finally, the mutant showed no difference from wild-type cells in secretion of

urease at 30°C or 37°C (Figure S3D).

Pdr802 is regulated by "host-like" conditions

We next tested the growth of the *pdr802* mutant under conditions more like those

encountered inside the mammalian host, using tissue culture medium (DMEM) at 37°C in

the presence of 5% CO_2. We found that although the *pdr802* mutant grew like WT in rich

medium (YPD), it grew poorly in DMEM (Figure S4A-B). To test whether the mutant cells

were dead or just static after growth in DMEM, we plated aliquots on solid medium to

measure CFU over time (Figure 2A). The *pdr802* culture showed a dramatic decrease in

viability compared to WT and the complemented strain, which was greatest in the first 24

h. This is the same time frame in which expression of the *PDR802* gene shows a striking

increase in wild type cells, as measured by RNA-Seq (Figure 2B).

Another important feature that is induced by growth in DMEM at 37°C and 5% CO_2

is the polysaccharide capsule, which we previously reported to be regulated by Pdr802,

based on negative staining with India ink (39). Fluorescence microscopy confirmed

increased capsule thickness of the mutant, which reverted to WT in the complemented

strain (Figure 3A). To quantify this change, we took advantage of a semi-automated

assay that we have developed (Figure S5), which measures capsules on a population

scale (Figure 3B) and is therefore very sensitive. This analysis showed that the capsule

thickness of *pdr802* cells resembles that of the well-studied hypercapsular mutant *pkr1*

(39, 69, 70) and is completely restored to WT by complementation at the native locus

(Figure 3C). Previous studies suggest that capsule thickness upon induction reflects the

size of the dominant capsule polymer (glucuronoxylomannan; GXM) (71, 72), which can

be analyzed by agarose gel migration and blotting with anti-capsule antibodies (71).

Consistent with the difference we observed in capsule thickness by imaging, this method

showed decreased mobility of GXM from *pdr802* as capsule induction progressed (Figure

S4C).

To validate the observations that we had made in standard 'host-like' conditions

based on synthetic tissue culture medium, we conducted similar studies in mouse serum

at 37°C and 5% CO_2. These conditions induced an even more pronounced hypercapsular

phenotype of the *pdr802* mutant (Figures 4A and 4B), as well as reduced cell viability

(Figure 4C) and increased cell body diameter (Figure 4D).

We were intrigued by the enlarged cell body and capsule of the *pdr802* mutant

cells in host-like conditions *in vitro* and decided to examine these phenotypes *in vivo*. For

these studies, we isolated fungal cells from the lungs of mice at various times after

infection and assessed their morphology by negative staining (Figure 5A). At each time

point, the mean mutant cell body diameter was larger than that of the controls.

Additionally, while this parameter was stable for WT and complemented strains

throughout the infection period, it trended larger at the end of the infection period for the

deletion mutant (Figure 5B). In contrast, mutant capsule thickness, although initially

larger than that of control cells, changed little throughout the period, while capsule

thickness of control cells increased to that level or beyond (Figure 5C). Furthermore,

although the total cell diameter of *pdr802* cells consistently exceeded that of WT and

complemented cells, their sizes became more comparable late in infection (Figure S6A).

Over time, therefore, the ratio of total cell diameter to cell body diameter for WT and

PDR802 cells steadily increased, while it remained roughly constant for the mutant

(Figure S6B).

Pdr802 negatively regulates Titan cell formation

We were particularly interested in the cell size phenotype of *pdr802* because Titan

cells have been strongly implicated in cryptococcal pathogenesis (19). By any definition

of this morphotype (cell body diameter greater than 10 or 15 µm or total cell diameter

greater than 30 µm), our mutant cell populations were dramatically enriched in Titan cells

at every time of infection that we assessed (Figure S6C).

To specifically test Titan cell formation by the *pdr802* strain, we subjected mutant

cells to *in vitro* conditions that induce this process (49) and analyzed the resulting

population by flow cytometry. Consistent with our *in vivo* observations, Titan cells

constituted a much larger fraction of the population in the mutant culture (13.2%) than in

the WT and complemented cultures (1.62% and 1.40%, respectively) (Figure 6).

Titan cells are poorly engulfed by host phagocytes (19, 45, 73), which may reflect

their increased size as well as alterations in capsule and cell wall (51). We observed this

reduced uptake for all strains after growth in conditions that favor Titan cell formation

(Figure 7, Titan vs YPD). Also, all strains showed a reduction in phagocytosis after

capsule induction in DMEM (Figure 7, DMEM vs YPD), which is not surprising because

the capsule is antiphagocytic (31, 73). Notably, the reduction in uptake was greatest for

the *pdr802* mutant in both of these conditions, even though it showed normal engulfment

when all strains were grown in the control condition (YPD). This is likely because the

mutant culture is both hypercapsular and enriched in Titan cells.

Identification of direct, functional targets of Pdr802

To identify direct targets of Pdr802, we performed chromatin immunoprecipitation

followed by sequencing (ChIP-Seq). We then compared the DNA sequences

immunoprecipitated by anti-mCherry mAb from cells expressing mCherry-Pdr802, which

grow similarly to WT (Figure S7A), and untagged cells. Both strains were grown for 24

hours in DMEM at 37°C and 5% CO_2, as this condition induces *PDR802* expression

dramatically compared to standard YPD growth conditions (Figure 2B).

Using 2-fold-enrichment over control as a cutoff value for peaks with adjusted p

value <0.05, we identified 656 binding sites for mCherry-Pdr802 in genomic DNA. Of

these, 540 occurred within 1,000 bp upstream of transcription start sites (Data Set S1,

Sheets 1 and 2), which we used as an approximation of regulatory regions. Application of

Discriminative Regular Expression Motif Elicitation (DREME) (74) to this set of upstream

regions identified several putative Pdr802 binding motifs, which were highly enriched in

GA (TC) (Figure S7B). Notably, the ChIP-seq data also suggested self-regulation of

PDR802, as has been reported for other cryptococcal TFs (75, 76) (Figure S7C).

To complement our ChIP studies, we determined the set of genes regulated by

Pdr802 under host-like conditions by performing RNA-Seq of WT and *pdr802* cells after

growth for 24 h in DMEM at 37°C and 5% CO_2 (Data set 1, Sheet 3). We then used dual-

threshold optimization (DTO) to analyze the RNA-seq and ChIP-seq data sets together.

This statistical method allowed us to combine the evidence from binding and expression

studies to converge on a set of direct and functional TF targets (77). The Pdr802 target

genes yielded by this analysis include key players in multiple processes implicated in

cryptococcal virulence, including quorum sensing, Titan cell formation, and stress

resistance (Data set 1, Sheets 4 and 5).

Pdr802 represses Titan cell production through regulation of quorum sensing

proteins

The most striking phenotype we observed in cells lacking *PDR802* is the marked

increase in Titan cell formation. We therefore examined our DTO target list for genes

known to influence this phenotype, such as those involved in quorum sensing. Recent

studies have shown that the quorum sensing peptide Qsp1 is a negative regulator of

Titan cell formation (47, 48); Titan cell formation increases upon deletion of the gene

encoding this peptide (*QSP1*) or proteins that mediate its maturation and import (*PQP1*

and *OPT1*, respectively). We found that Pdr802 positively regulates *PQP1* and *OPT1*

gene expression (Table 1), consistent with its repression of Titan cell formation.

A study of *C. neoformans* cells exposed to Titan cell inducing conditions *in vitro*

reported that 562 genes were upregulated in this condition, while 421 genes were

downregulated (48). The overlap of these genes with our DTO set of Pdr802-regulated

genes included three TF genes *LIV3*, *STB4*, and *ZFC3* (48) (Data Set S2, Sheets 1 and

2). The first two are repressed during Titan cell induction while *ZFC3* (also known as

CQS2) is induced. Our analysis showed that Pdr802 positively regulates expression of

LIV3 and *STB4*, while it negatively regulates *ZFC3* (Table 1), in concordance with our

phenotypic observations of Titan cell formation. Notably, Liv3 and Zfc3 are responsive to

the peptide Qsp1 (75, 76) and are important for *C. neoformans* virulence, while Stb4

influences cryptococcal brain infection (67).

275 Pdr802 coordinates cryptococcal response to the host environment

C. neoformans deploys a variety of proteins to resist the many challenges it

experiences upon host entry, which include oxidative and temperature stress. Multiple

genes that are central to these responses were identified as direct, functional targets of

Pdr802 by our DTO analysis (Table 2). For example, Pdr802 induces the expression of

genes whose products detoxify reactive oxygen species (ROS), such as *CAT1*, *CAT2*,

and *SOD1* (78, 79), or participate in resistance to these compounds, such as *FZC34*,

MIG1, and *CCK1* (52, 80, 81) (Table 2). Both the kinase Cck1 (also known as Yck2) and

the TF Fzc34 have been implicated in cryptococcal virulence (80, 82).

As noted above, melanin has important anti-oxidant properties that promote

cryptococcal survival inside the host (16). Under host-like conditions, Pdr802 regulates

genes required for melanization, even though it melanizes normally *in vitro*. These genes

include *CAC1*, *PKC1*, *CUF1*, and *SNF5* (Table 2). Cac1 is an adenylyl cyclase

responsible for cyclic AMP (cAMP) production in *C. neoformans*, which plays a central

role in melanin synthesis as well as proper capsule production, mating and virulence (83).

The kinase Pkc1 induces production of the laccase (Lac1) that forms melanin and plays a

key role in resistance to oxidative and nitrosative stress (84, 85); the TF Cuf1 regulates

LAC1 expression and is important for cryptococcal virulence (86, 87); and *SNF5* is

required for full melanization (88). Melanin occurs in the fungal cell wall, which is another

key component in fungal stress resistance. Pdr802 is also a direct, functional regulator of

several genes whose products influence cell wall glycan content: two chitin deacetylases

(Cda3 and Mp98) and the mannoprotein MP88 (Table 2). Changes in mannose and chitin

occur in Titan cell walls (51).

Pdr802 positively regulates the expression of several proteins required for yeast

growth at 37ºC, including the kinases Kic1 and Ire1 (Table 2). Ire1 is a regulator of the

cryptococcal Unfolded Protein Response (UPR) pathway and lack of Ire1 or Kic1 impacts

C. neoformans virulence (80, 89). Pdr802 also modulates cryptococcal urease activity,

which is required for dissemination to the central nervous system (CNS) (11, 12), by

regulating the urea transporter Dur3 and other proteins that influence urease activity (e.g.

the kinases Fab1, Kin1, and Gut1 and the TF Hlh1) (Table 2). Deletion of *FAB1*, *KIN1*, or

HLH1 impair urease activity in *C. neoformans*, while *GUT1* disruption induces it (52, 80).

Above we documented the role of Pdr802 in capsule synthesis, which is

dramatically upregulated in the host environment in general and is further increased in

cells lacking this TF. We found that Pdr802 is a positive regulator of multiple genes that

have been implicated in reducing cryptococcal capsule thickness. These include the

kinases Arg2, Iks1, and Ksp1; the TF Fzc51; and the phosphodiesterase Pde2 (Table 2).

Notably, null mutants for those genes are hypercapsular, similar to *pdr802* cells (52, 80,

90). Pdr802 is a negative regulator of the TF Bzp4, which, as mentioned above, positively

regulates capsule (Table 2) (52).

Pdr802 regulates calcineurin target genes

The calcineurin signaling pathway is activated by calcium and governs stress

response and virulence in *C. neoformans* (91–93). One major mediator of calcineurin

signaling is the transcription factor Crz1 mentioned above, which is highly responsive to

temperature and influences cryptococcal virulence (57, 62). Upon intracellular calcium

influx calcineurin dephosphorylates Crz1, which then translocates to the nucleus and

regulates gene expression (57, 63). We found that Pdr802 binds the *CRZ1* gene

promoter and positively regulates its expression (Figure 8A, Table 2, and Data Set S2,

Sheet 3). Pdr802 also binds and regulates five other genes whose products are

dephosphorylated by calcineurin; these include the phosphatase Had1, which is

important for cryptococcal cell wall remodeling and virulence (Figure 8B, Table 2, and

Data Set S2, Sheet 3) (63, 94).

Because Crz1 helps maintain normal cryptococcal Ca^{2+} concentrations through the

regulation of calcium transporters (57), we wondered about the intracellular calcium

levels in *pdr802* cells. We found that after 24 hours of growth in DMEM, the level of

cytosolic calcium in the mutant significantly exceeded that of WT or complemented

strains (Figure 8C). It was still, however, below that of a *crz1* null mutant, supporting that

Pdr802 is not the sole regulator of *CRZ1* expression. Notably, *PDR802* deletion had no

effect in rich medium (YPD), which reinforces our hypothesis that Pdr802 acts primarily in

host-like conditions. To further explore the relationship of Pdr802 and calcineurin, we

compared published gene expression profiles of a calcineurin mutant (57) to our DTO

data set. Of the 393 genes that are differently expressed in the calcineurin mutant under

thermal stress, 26 are regulated by Pdr802 (Data Set S2, Sheet 4).

DISCUSSION

We have shown that Pdr802 is a potent regulator of cryptococcal responses to the

host environment. In this context, it influences the formation of capsule and Titan cells as

well as cellular responses to temperature and oxidative stress, acts as a downstream

effector of calcineurin, and modulates calcium availability. The last function is likely

achieved through its positive regulation of the transcription factor Crz1, which in turn

modulates the calcium transporters Pmc1 and Vcx1 (57). Since calcium ion is a major

second messenger in eukaryotic cells, its accumulation in *pdr802* cells affects multiple

processes central to host interactions, including stress responses, cell wall integrity, and

capsule size (61, 62, 92, 95, 96).

C. neoformans dissemination to the brain is the main driver of patient mortality (2).

We found that dissemination of *pdr802* cells is significantly impaired, although they do

occasionally reach the brain. These observations can be explained by a combination of

factors. First, the limited accumulation of the *pdr802* mutant in the lungs, due to factors

summarized above, may directly affect dissemination (97). Second, this strain survives

poorly in mouse serum, as demonstrated directly by our culture experiments and

indirectly by our inability to detect it in the blood of infected mice, even 75 days after

infection. The latter might be because the cells do not reach the blood or because they

are rapidly eliminated, consistent with previous observations (98). Third, the thick

capsules of the *pdr802* mutant reduce its ability to reach the brain. This is true whether

fungal entry occurs directly, by the movement of free fungi across the BBB, or indirectly,

via a Trojan horse mechanism that requires macrophage uptake (99); such uptake is

impeded by enlarged capsules, independent of cell size (31). Fourth, calcium imbalance

directly affects cryptococcal transmigration (100). Finally, *pdr802* cells show reduced

expression of genes required for urease activity, which promotes *C. neoformans*

dissemination to the CNS (11, 12, 100). Interestingly, despite all of these obstacles to

dissemination, mutant cells that do reach the brain are able to proliferate to wild-type

levels.

Titan cells are a robust and persistent morphotype of *C. neoformans* that

contributes to yeast virulence (45). We showed that cells lacking Pdr802 demonstrate

increased formation of Titan cells *in vivo* and *in vitro*, suggesting that this TF is a novel

repressor of this process. Although Titan cells enhance aspects of cryptococcal

pathogenesis (19, 101), their overproduction negatively impacts dissemination to the

brain due to their resistance to phagocytosis by macrophages (19, 45) and decreased

penetration of biological barriers (19).

Our combined analysis of DNA binding and gene expression data allows us to

understand the increase in Titan cell formation that occurs upon deletion of *PDR802*.

Under host-like conditions, Pdr802 positively regulates Pqp1, Opt1 and Liv3, all key

proteins in the cryptococcal quorum sensing pathway, which represses Titan cell

formation (47, 48). In the absence of this TF, quorum sensing is impaired, increasing

Titan cell formation. Pdr802 may also indirectly modulate Titan cell formation by

regulating other TFs that impact this process, such as Zfc3 (Cqs2) and Stb4.

We know that capsule, a key virulence factor, is typically highly induced in the host

or host-like conditions (102). Our studies *in vitro*, *ex vivo*, and *in vivo* show that Pdr802

normally reins in this process. This likely occurs via a combination of Pdr802's repression

of the TF Bzp4, which positively regulates capsule size, and the induction of other factors

(e.g. the TF Fzc1, the phosphodiesterase Pde2, and the kinases Ksp1, Arg2, and Iks1)

that negatively regulate capsule size (52, 80, 90).

Overall, we found that Pdr802 influences key cryptococcal phenotypes that

influence virulence, including quorum sensing, stress responses, Titan cell formation, and

capsule production (Figure 9). We have further identified multiple genes that are central

in these processes and are directly regulated by Pdr802. Some of these targets are also

regulated by calcineurin (e.g. Had1 and Crz1) or by another important TF, Gat201 (e.g.

Opt1, Liv3, Zfc3) (60, 75, 76). Finally, the expression of *PDR802* itself is regulated by the

TFs Gat201 and Hob1 (67, 76). The crosstalk between all of these regulatory

mechanisms remains to be dissected. Nonetheless, it is evident that Pdr802 is critical for

both survival in the lung and dissemination to the brain, thus explaining its role in

cryptococcal virulence.

MATERIALS AND METHODS

Strain construction and cell growth

We previously reported the *PDR802* deletion mutant (*pdr802*) in the KN99α strain

background (103) that was used in this work (39). Complementation of this mutant with

the wild-type gene at the native locus (*PDR802*) and construction of a strain that

expresses Pdr802 with N-terminal mCherry (mCherry-Pdr802) are detailed in the

Supplementary Methods. For all studies, *C. neoformans* strains were inoculated from

single colonies into YPD medium (2% [wt/vol] dextrose, 2% [wt/vol] Bacto peptone and

1% [wt/vol] yeast extract in double-distilled water [ddH$_2$O]) and grown overnight at 30°C

with shaking at 230 rpm before further handling as detailed below. To assess viability

during growth in tissue culture medium, overnight cultures were washed with phosphate-

buffered saline (PBS), diluted to 10^6 cells/ml in DMEM (Sigma, D6429), plated (1 ml/well)

in triplicate in 24-well plates, and incubated at 37°C and 5% CO_2. At the indicated times

cells were mixed thoroughly, diluted in PBS, and plated on YPD agar (YPD medium, 2%

agar [wt/vol]) for assessment of colony-forming units (CFU). To assess viability during

growth in mouse serum (prepared as below), YPD-grown cryptococcal cells (10^3) were

incubated in 100 µl of serum in 96-well plates for 24 h at 37°C and 5% CO_2 and CFU

assessed as above.

Animal experiments

All animal protocols were approved by the Washington University Institutional Animal

Care and Use Committee (reference 20170131) or Comissão de Ética no Uso de

Animais – CEUA (reference 30936), and care was taken to minimize handling and

discomfort. For survival studies, groups of five 4- to 6-week-old female C57BL/6 mice

(The Jackson Laboratory) were anesthetized by subcutaneous injection of 1.20 mg

ketamine and 0.24 mg xylazine in 120 µl sterile water and intranasally infected with 5 x

10^4 cryptococcal cells. The mice were monitored and humanely sacrificed when their

weight decreased to below 80% of initial weight or if they showed signs of disease, at

which point organ burden was assessed. The lungs and brains were harvested,

homogenized, diluted and plated on YPD agar. The resulting CFU were enumerated and

survival differences were assessed by Kaplan-Meier analysis.

For timed organ burden studies, *C. neoformans* overnight cultures were centrifuged

(1,000 x *g* for 3 min), washed with sterile PBS, and resuspended in PBS to 1 x 10^6

cells/ml. Groups of three 4- to 6-week-old female C57BL/6 mice (Centro Multidisciplinar

para Investigação Biológica na Área da Ciência em Animais de Laboratório, CEMIB)

were anesthetized as above and intranasally infected with 5 x 10^4 cryptococcal cells, and

monitored as above. At set time points post-infection (see text), mice were sacrificed and

fungal burden was assessed from organs (as above) or blood (obtained by cardiac

puncture). Organ burden was analyzed by Kruskal-Wallis test with Dunn's multiple

comparison *post hoc* test for each day post-infection.

To assess cryptococcal viability in mouse serum, 6 BALB/c mice were anesthetized with

isoflurane and blood was collected from the retro-orbital space using a sterile capillary

tube. Collected blood was incubated at 37°C for 30 min and serum was isolated by

centrifugation at 1,000 x *g* for 15 min and then heat-inactivated at 56 °C for 30 min.

Capsule analysis

To qualitatively assess capsule thickness, strains were grown in YPD medium for 16 h,

washed with PBS, and 10^6 cells were incubated in DMEM or mouse serum for 24 h at

37°C and 5% CO_2. After incubation, cells were fixed in 4% paraformaldehyde, washed

three times with PBS, and mixed with similar volumes of India ink for capsule

visualization and measurement as previously described (104).

For population-level capsule measurement, *C. neoformans* strains were grown overnight

in YPD, washed with PBS, and diluted to 10^6 cells/ml in DMEM. 150 µl aliquots were then

plated in quadruplicate in a poly-L-lysine coated 96-well plate (Fisher 655936) and

incubated at 37°C and 5% CO_2. After 24 hours, the cells were washed with PBS and

incubated with 150 µl of a staining mixture (100 µg/ml Calcofluor white to stain cell walls,

50 µg/ml of anticapsular monoclonal antibody 302 conjugated to Alexa Fluor 488

(Molecular probes), and 1.5% goat serum in PBS) for 30 minutes at room temperature in

the dark. The cells were washed again with PBS, fixed with 4% formaldehyde for 10

minutes at room temperature, washed with PBS, and each well refilled with 150 µl PBS.

The cells were imaged using a BioTek Cytation 3 imager, which automatically collected

100 images per well in a grid pattern at the well center. Image files were prepared for

analysis with the GE InCell Translator and assembled into .xdce image stacks for

analysis with the GE INCell Developer Toolbox 1.9. Cell wall and capsule images were

first filtered to remove background noise and border objects and then cells were identified

using shape-based object segmentation (3-pixel kernel, 50% sensitivity) followed by

watershed clump breaking to prevent apparent connectivity caused by incomplete

segmentation. Target linking was performed to assign each cell wall object to one

capsule object based on known 1:1 pairing and location, generating a target set. Capsule

and cell wall object diameters were calculated for each target set (hundreds to thousands

per well), and the difference between each pair of measurements was defined as the

capsule thickness. Data were normalized by the difference in capsule thickness between

uninduced and induced WT cells, which were included in each experiment, and

compared to hypercapsular (*pkr1*) (39) and hypocapsular (*ada2*) (105) control strains in

each experiment. Capsule sizes were compared by One-Way ANOVA with Dunnett's

multiple comparison *post hoc* test.

To measure capsule thickness of cryptococcal cells grown in the lungs of infected mice,

lung homogenates were filtered through a cell strainer with 40 μm pores using a syringe

plunger, fixed in 3.7% formaldehyde, and used for India ink staining and measurement as

above. For the visualization of KN99α and *PDR802* cells from mouse lungs after 18 days

of infection, the tissue was treated with 50 μg/ml DNAse I for 30 min at 37°C.

GXM immunoblotting was conducted as previously described (71). Briefly, 10^6 cells/ml

were grown in DMEM for 24 and 48h. Culture supernatant fractions were then resolved

by gel electrophoresis on 0.6% agarose, transferred onto nylon membranes, and probed

with 1 µg/ml anti-GXM antibody 302.

Phenotypic assays

For stress plates, cryptococcal cells were grown overnight in YPD, washed with PBS, and

diluted to 10^7 cells/ml in PBS. Aliquots (3 µl) of 10-fold serial dilutions were spotted on

YPD or YNB agar supplemented with various stressors (sorbitol, NaCl, $CaCl_2$, LiCl,

Congo Red, Calcofluor white, caffeine, SDS, $NaNO_2$, H_2O_2 and ethanol) in the

concentrations indicated in the figures. Melanization was tested on plates made by

mixing 10 ml of 2X minimal medium (2 g/L L-asparagine, 1 g/L $MgSO_4 \cdot 7H_2O$, 6 g/L

KH_2PO_4, 2 g/L thiamine, 2 mM L-3,4-dihydroxyphenylalanine [L-DOPA] and 0.1%

dextrose was added for melanization induction or 0.5% for melanization inhibition) with

10 ml of 2% agar-water per plate. A control strain lacking the ability to melanize was used

as a control (*lac1*) (88). For the solid urease assay, 10 µl of a 10^7 cells/ml suspension in

water was plated on Christensen's urea solid media (1 g/L peptone, 1 g/L dextrose, 5 g/L

NaCl, 0.8 g/L KH_2PO_4, 1.2 g/L Na_2HPO_4, 0.012 g/L phenol red and 15 g/L agar, pH 6.8).

Plates were incubated at 30°C or 37°C.

Titan cells

Titan cell induction was performed in 1x PBS supplemented with 10% heat inactivated

Fetal Calf Serum (FCS) for 72 hours at 37°C and 5% CO_2 as recently described (49) and

quantified by flow cytometry as previously reported (47, 48).

Phagocytosis

J774.16 cells were prepared for uptake experiments by seeding (10^5 cells/well) in a 96-

well plate and incubating in DMEM supplemented with 10% Fetal Bovine Serum (FBS) at

37°C and 5% CO_2 for 24 h. *C. neoformans* cells were prepared for uptake experiments

by inoculating an overnight culture in YPD into either DMEM or Titan cell induction

medium (49) and growing at 37°C and 5% CO_2 for 24 or 72 h, respectively. To initiate the

study, cryptococcal cells were washed with PBS and opsonized with anti-capsular

antibody 18B7 (1 µg/ml) for 1 h at 37°C while macrophages were activated with 50 nM

phorbol myristate acetate (PMA) for 1 h at 37°C and 5% CO_2; 10^6 cryptococcal cells were

then incubated with the macrophages for 2 h at 37°C and 5% CO_2. The wells were then

washed three times with warm PBS and the macrophages lysed with 0.1% Triton in PBS

and plated for CFU as above. Fold-change in CFU was assessed by comparison to the

CFU of opsonized cells. One-Way ANOVA with Dunnett's multiple comparison *post hoc*

test was used to compare phagocytosis of *pdr802* and *PDR802* strains with that of

KN99α.

Chromatin Immunoprecipitation (ChIP)

ChIP studies were performed as previously described (39, 105). Briefly, wild type and N-

terminal-mCherry-Pdr802 strains were cultivated in DMEM for 24 hours at 37°C and 5%

CO_2. The cells were then fixed with formaldehyde, lysed by mechanical bead-beating,

and the cell debris removed by centrifugation. The supernatant fraction was sheared by

sonication, centrifuged, and an aliquot was reserved as 'Input'. The remaining material

was incubated with rabbit IgG anti-mCherry antibody (Abcam, ab213511) tethered to

protein A sepharose ('IP') or sepharose alone ('Mock') overnight at 4°C. The beads were

then washed, incubated at 65°C to reverse DNA-DNA and DNA-protein crosslinks and

the DNA recovered by phenol/chloroform/isoamyl alcohol (25:24:1) extraction, ethanol

precipitation, and resuspension in nuclease-free water.

Samples were submitted to the Washington University Genome Technology Access

Center for library preparation and DNA samples were sequenced using the Illumina

Nextseq platform. The first replicate was sequenced using paired-end 2x75-bp reads and

replicates 2 and 3 were sequenced using single-end 75-bp reads; the minimum coverage

obtained was ~16x. The quality of the reads was evaluated by FastQC (106). Fastq files

were aligned to the KN99 genome (107) using NextGenMap 0.5.3 (108). SAM files were

converted to bam, reads were sorted and indexed, and read duplicates were removed

from the final bam files using samtools (109). Samtools was also used to filter out reads

with a mapping quality lesser than 20 phreds to guarantee single alignment of the reads.

Peaks were called using MACS2 (2.1.1.20160309) (110), filtered by size (maximum

threshold 5 kb and no minimum), and annotated using Homer 4.8 (111). The significant

peaks were chosen using the cutoff of fold enrichment above 2 and adjusted p value <

0.05 and read coverage of each peak was obtained using Samtools (109). Pdr802

binding motifs were identified using DREME (74); partial motifs were defined as at least 5

consecutive bp of the motif.

RNA-Seq and Dual-Threshold Optimization (DTO)

RNA from wild-type and *pdr802* cells grown for 24 hours in DMEM (37°C, 5% CO_2) was

isolated and sequenced as previously described (39). Briefly, cDNA samples were

sequenced using the Illumina Nextseq platform for single-end 1 x 75 bp reads and read

quality was evaluated by FastQC (106). Fastq files were aligned to the KN99 genome

(107) using Novoalign (112), SAM files were converted to bam, reads were sorted and

indexed, and read duplicates were removed from the final bam files using Samtools

(109). The number of reads mapped per gene was calculated using HTSeq (113) and

differential gene expression was analyzed with DESeq2 (114), using the Independent

Hypothesis Weighting (IHW) package to calculate the adjusted p-values (115). Dual-

Threshold Optimization (DTO) analysis was performed as recently described (77). This is

a method for simultaneously finding the best thresholds for significance in a TF binding

location dataset (e.g. ChIP) and a TF perturbation-response dataset (e.g. RNA-Seq of a

TF mutant). It works by trying out all pairs of thresholds for the two datasets, picking the

pair that minimizes the probability of the overlap between the bound and responsive gene

sets occurring by chance under a null model, and testing the significance of the overlap

by comparison to randomly permuted data. Our application of DTO to our ChIP and RNA-

Seq data yielded 1455 bound genes, 5186 responsive genes, and 1167 genes that were

both bound and responsive. Based on DTO, Pdr802 has an acceptable convergence

from binding and perturbation, with p value < 0.01 from the random permutation test and

minimum expected FDR less than or equal to 20% at 80% sensitivity. In addition to

requiring a statistically significant overlap between the ChIP-Seq and RNA-Seq gene

sets, we filtered out any genes for which traditional differential expression analysis

yielded an adjusted p-value ≤ 0.15 or absolute $\log_2$ of fold change ≥ 0.3, leaving 380

bound targets.

Intracellular calcium measurement

To measure intracellular free Ca^{2+}, yeast cells were cultured overnight in YPD at 30°C

with shaking, washed three times with deionized water, diluted to 10^6 cells/ml in DMEM

(Sigma, D6429), plated (1 ml/well) in triplicate in 24-well plates, and incubated at 37°C

and 5% CO_2 for 24 hours. At the indicated times, cells were mixed thoroughly, diluted in

PBS containing 2 µM Fluo4-AM (Thermo Fisher), incubated at 30°C for 30 min, and

analyzed using flow cytometry. The overnight culture was used as a control and treated

as above.

Data availability

ChIP-seq and RNA-seq data files are available at the NCBI Gene Expression Omnibus

under accession numbers GSE153134 and GSE162851, respectively.

REFERENCES

1. Kwon-Chung KJ, Fraser JA, Doering TL, Wang ZA, Janbon G, Idnurm A, Bahn Y-

S. 2014. *Cryptococcus neoformans* and *Cryptococcus gattii*, the Etiologic Agents of

Cryptococcosis. Cold Spring Harb Perspect Med 4:a019760.

2. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning

DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated

cryptococcal meningitis: an updated analysis. Lancet Infect Dis 17:873–881.

3. Ballou ER, Johnston SA. 2017. The cause and effect of *Cryptococcus* interactions

with the host. Curr Opin Microbiol 40:88–94.

4. Sabiiti W, May RC. 2012. Mechanisms of infection by the human fungal pathogen

Cryptococcus neoformans. Future Microbiol 7:1297–1313.

5. Garcia-Hermoso D, Janbon G, Dromer F. 1999. Epidemiological evidence for

dormant *Cryptococcus neoformans* infection. J Clin Microbiol 37:3204–3209.

6.   Chang YC, Stins MF, McCaffery MJ, Miller GF, Pare DR, Dam T, Paul-Satyasee M,

590   Kim KS, Kwon-Chung KJ. 2004. Cryptococcal yeast cells invade the central

591   nervous system via transcellular penetration of the blood-brain barrier. Infect

592   Immun 72:4985–4995.

593   7.   Huang S-H, Long M, Wu C-H, Kwon-Chung KJ, Chang YC, Chi F, Lee S, Jong A.

594   2011. Invasion of *Cryptococcus neoformans* into Human Brain Microvascular

595   Endothelial Cells Is Mediated through the Lipid Rafts-Endocytic Pathway via the

596   Dual Specificity Tyrosine Phosphorylation-regulated Kinase 3 (DYRK3). J Biol

597   Chem 286:34761–34769.

598   8.   Jong A, Wu C-H, Shackleford GM, Kwon-Chung KJ, Chang YC, Chen H-M,

599   Ouyang Y, Huang S-H. 2008. Involvement of human CD44 during *Cryptococcus*

600   *neoformans* infection of brain microvascular endothelial cells. Cell Microbiol

601   10:1313–1326.

602   9.   Maruvada R, Zhu L, Pearce D, Zheng Y, Perfect J, Kwon-Chung KJ, Kim KS. 2012.

603   *Cryptococcus neoformans* phospholipase B1 activates host cell Rac1 for traversal

604   across the blood-brain barrier. Cell Microbiol 14:1544–1553.

605   10.   Chen SHM, Stins MF, Huang S-H, Chen YH, Kwon-Chung KJ, Chang Y, Kim KS,

606   Suzuki K, Jong AY. 2003. *Cryptococcus neoformans* induces alterations in the

607   cytoskeleton of human brain microvascular endothelial cells. J Med Microbiol

608   52:961–970.

609   11.   Olszewski MA, Noverr MC, Chen G-H, Toews GB, Cox GM, Perfect JR, Huffnagle

610   GB. 2004. Urease Expression by *Cryptococcus neoformans* Promotes

611   Microvascular Sequestration, Thereby Enhancing Central Nervous System

Invasion. Am J Pathol 164:1761–1771.

12. Shi M, Li SS, Zheng C, Jones GJ, Kim KS, Zhou H, Kubes P, Mody CH. 2010.

Real-time imaging of trapping and urease-dependent transmigration of

Cryptococcus neoformans in mouse brain. J Clin Invest 120:1683–1693.

13. Santiago-Tirado FH, Onken MD, Cooper JA, Klein RS, Doering TL. 2017. Trojan

Horse Transit Contributes to Blood-Brain Barrier Crossing of a Eukaryotic

Pathogen. MBio 8:e02183-16.

14. Jarvis JN, Bicanic T, Loyse A, Namarika D, Jackson A, Nussbaum JC, Longley N,

Muzoora C, Phulusa J, Taseera K, Kanyembe C, Wilson D, Hosseinipour MC,

Brouwer AE, Limmathurotsakul D, White N, van der Horst C, Wood R, Meintjes G,

Bradley J, Jaffar S, Harrison T. 2014. Determinants of Mortality in a Combined

Cohort of 501 Patients With HIV-Associated Cryptococcal Meningitis: Implications

for Improving Outcomes. Clin Infect Dis 58:736–745.

15. John R. Perfect, William E. Dismukes, Francoise Dromer, David L. Goldman, John

R. Graybill, Richard J. Hamill, Thomas S. Harrison, Robert A. Larsen, Olivier

Lortholary, Minh-Hong Nguyen, Peter G. Pappas, William G. Powderly, Nina Singh,

Jack D. Sobel TCS. 2010. Clinical Practice Guidelines for the Management of

Cryptococcal Disease: 2010 Update by the Infectious Diseases Society of America.

Chinese J Infect Chemother 50:291–322.

16. Nosanchuk JD, Casadevall A. 2003. The contribution of melanin to microbial

pathogenesis. Cell Microbiol 5:203–223.

17. Singh A, Panting RJ, Varma A, Saijo T, Waldron KJ, Jong A, Ngamskulrungroj P,

Chang YC, Rutherford JC, Kwon-Chung KJ. 2013. Factors Required for Activation

of Urease as a Virulence Determinant in *Cryptococcus neoformans*. MBio

4:e00220-13.

18. Chang YC, Kwon-Chung KJ. 1994. Complementation of a capsule-deficient

mutation of *Cryptococcus neoformans* restores its virulence. Mol Cell Biol 14:4912–

4919.

19. Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chrétien F, Heitman J,

Dromer F, Nielsen K. 2010. Cryptococcal Cell Morphology Affects Host Cell

Interactions and Pathogenicity. PLoS Pathog 6:e1000953.

20. Zaragoza O, Nielsen K. 2013. Titan cells in *Cryptococcus neoformans*: cells with a

giant impact. Curr Opin Microbiol 16:409–413.

21. Wang Y, Aisen P, Casadevall A. 1995. *Cryptococcus neoformans* melanin and

virulence: Mechanism of action. Infect Immun 63:3131–3136.

22. Mednick AJ, Nosanchuk JD, Casadevall A. 2005. Melanization of *Cryptococcus*

neoformans Affects Lung Inflammatory Responses during Cryptococcal Infection.

Infect Immun 73:2012–2019.

23. Wang Y, Casadevall A. 1994. Susceptibility of melanized and nonmelanized

Cryptococcus neoformans to nitrogen- and oxygen-derived oxidants. Infect Immun

62:3004–7.

24. Doering TL, Nosanchuk JD, Roberts WK, Casadevall A. 1999. Melanin as a

potential cryptococcal defence against microbicidal proteins. Med Mycol 37:175–

81.

25.   Agustinho DP, Nosanchuk JD. 2017. Functions of Fungal MelaninsReference

Module in Life Sciences. Elsevier.

26.   Cox GM, Mukherjee J, Cole GT, Casadevall A, Perfect JR. 2000. Urease as a

virulence factor in experimental cryptococcosis. Infect Immun 68:443–8.

27.   Bose I, Reese AJ, Ory JJ, Janbon G, Doering TL. 2003. A Yeast under Cover: the

Capsule of *Cryptococcus neoformans*. Eukaryot Cell 2:655–663.

28.   Wang ZA, Li LX, Doering TL. 2018. Unraveling synthesis of the cryptococcal cell

wall and capsule. Glycobiology 28:719–730.

29.   Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: the outer

layers of *Cryptococcus neoformans*. Mem Inst Oswaldo Cruz 113:e180040.

30.   Bulmer GS, Sans MD. 1967. *Cryptococcus neoformans*. II. Phagocytosis by human

leukocytes. J Bacteriol 94:1480–3.

31.   Bulmer GS, Sans MD. 1968. *Cryptococcus neoformans*. 3. Inhibition of

phagocytosis. 95:5–8.

32.   Tacker JR, Farhi F, Bulmer GS. 1972. Intracellular fate of *Cryptococcus*

neoformans. Infect Immun 6:162–7.

33.   Monari C, Bistoni F, Vecchiarelli A. 2006. Glucuronoxylomannan exhibits potent

immunosuppressive properties. FEMS Yeast Res 6:537–542.

34.   Hayes JB, Sircy LM, Heusinkveld LE, Ding W, Leander RN, McClelland EE, Nelson

DE. 2016. Modulation of Macrophage Inflammatory Nuclear Factor κB (NF-κB)

Signaling by Intracellular *Cryptococcus neoformans*. J Biol Chem 291:15614–

15627.

35.   Fonseca FL, Nohara LL, Cordero RJB, Frases S, Casadevall A, Almeida IC,

Nimrichter L, Rodrigues ML. 2010. Immunomodulatory Effects of Serotype B

Glucuronoxylomannan from *Cryptococcus gattii* Correlate with Polysaccharide

Diameter. Infect Immun 78:3861–3870.

36.   Garcia-Hermoso D, Dromer F, Janbon G. 2004. *Cryptococcus neoformans* Capsule

Structure Evolution *In Vitro* and during Murine Infection. Infect Immun 72:3359–

3365.

37.   Guimarães AJ, Frases S, Cordero RJB, Nimrichter L, Casadevall A, Nosanchuk

JD. 2010. *Cryptococcus neoformans* responds to mannitol by increasing capsule

size *in vitro* and *in vivo*. Cell Microbiol 12:740–753.

38.   Cordero RJB, Pontes B, Guimarães AJ, Martinez LR, Rivera J, Fries BC,

Nimrichter L, Rodrigues ML, Viana NB, Casadevall A. 2011. Chronological Aging Is

Associated with Biophysical and Chemical Changes in the Capsule of

Cryptococcus neoformans. Infect Immun 79:4990–5000.

39.   Maier EJ, Haynes BC, Gish SR, Wang ZA, Skowyra ML, Marulli AL, Doering TL,

Brent MR. 2015. Model-driven mapping of transcriptional networks reveals the

circuitry and dynamics of virulence regulation. Genome Res 25:690–700.

40.   Santos JRA, Holanda RA, Frases S, Bravim M, Araujo GDS, Santos PC, Costa

MC, Ribeiro MJA, Ferreira GF, Baltazar LM, Miranda AS, Oliveira DB, Santos

CMA, Fontes ACL, Gouveia LF, Resende-Stoianoff MA, Abrahão JS, Teixeira AL,

Paixão TA, Souza DG, Santos DA. 2014. Fluconazole Alters the Polysaccharide

Capsule of *Cryptococcus gattii* and Leads to Distinct Behaviors in Murine

Cryptococcosis. PLoS One 9:e112669.

41.   Silveira CP, Piffer AC, Kmetzsch L, Fonseca FL, Soares DA, Staats CC, Rodrigues

ML, Schrank A, Vainstein MH. 2013. The heat shock protein (Hsp) 70 of

Cryptococcus neoformans is associated with the fungal cell surface and influences

the interaction between yeast and host cells. Fungal Genet Biol 60:53–63.

42.   Alspaugh JA. 2015. Virulence mechanisms and *Cryptococcus neoformans*

pathogenesis. Fungal Genet Biol 78:55–58.

43.   García-Rodas R, de Oliveira H, Trevijano-Contador N, Zaragoza O. 2018.

Cryptococcal Titan Cells: When Yeast Cells Are All Grown up, p. 101–120. *In*

Current Topics in Microbiology and Immunology. Springer Verlag.

44.   Zhou X, Ballou ER. 2018. The *Cryptococcus neoformans* Titan Cell: From *In Vivo*

Phenomenon to *In Vitro* Model. Curr Clin Microbiol Reports 5:252–260.

45.   Zaragoza O, García-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodríguez-

Tudela JL, Casadevall A. 2010. Fungal Cell Gigantism during Mammalian Infection.

PLoS Pathog 6:e1000945.

46.   Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, Berman J,

Nielsen K. 2015. Polyploid Titan Cells Produce Haploid and Aneuploid Progeny To

Promote Stress Adaptation. MBio 6:e01340-15.

47.   Hommel B, Mukaremera L, Cordero RJB, Coelho C, Desjardins CA, Sturny-Leclère

A, Janbon G, Perfect JR, Fraser JA, Casadevall A, Cuomo CA, Dromer F, Nielsen

K, Alanio A. 2018. Titan cells formation in *Cryptococcus neoformans* is finely tuned

by environmental conditions and modulated by positive and negative genetic

regulators. PLOS Pathog 14:e1006982.

48. Trevijano-Contador N, de Oliveira HC, García-Rodas R, Rossi SA, Llorente I,

Zaballos Á, Janbon G, Ariño J, Zaragoza Ó. 2018. *Cryptococcus neoformans* can

form titan-like cells *in vitro* in response to multiple signals. PLOS Pathog

14:e1007007.

49. Dambuza IM, Drake T, Chapuis A, Zhou X, Correia J, Taylor-Smith L, LeGrave N,

Rasmussen T, Fisher MC, Bicanic T, Harrison TS, Jaspars M, May RC, Brown GD,

Yuecel R, MacCallum DM, Ballou ER. 2018. The *Cryptococcus neoformans* Titan

cell is an inducible and regulated morphotype underlying pathogenesis. PLOS

Pathog 14:e1006978.

50. García-Barbazán I, Trevijano-Contador N, Rueda C, de Andrés B, Pérez-Tavárez

R, Herrero-Fernández I, Gaspar ML, Zaragoza O. 2016. The formation of titan cells

in *Cryptococcus neoformans* depends on the mouse strain and correlates with

induction of Th2-type responses. Cell Microbiol 18:111–124.

51. Mukaremera L, Lee KK, Wagener J, Wiesner DL, Gow NAR, Nielsen K. 2018. Titan

cell production in *Cryptococcus neoformans* reshapes the cell wall and capsule

composition during infection. Cell Surf 1:15–24.

52. Jung K-W, Yang D-H, Maeng S, Lee K-T, So Y-S, Hong J, Choi J, Byun H-J, Kim

H, Bang S, Song M-H, Lee J-W, Kim MS, Kim S-Y, Ji J-H, Park G, Kwon H, Cha S,

Meyers GL, Wang LL, Jang J, Janbon G, Adedoyin G, Kim T, Averette AK,

Heitman J, Cheong E, Lee Y-H, Lee Y-W, Bahn Y-S. 2015. Systematic functional

profiling of transcription factor networks in *Cryptococcus neoformans*. Nat Commun

6:6757.

53.   Schneider R de O, Fogaça N de SS, Kmetzsch L, Schrank A, Vainstein MH, Staats

CC. 2012. Zap1 Regulates Zinc Homeostasis and Modulates Virulence in

Cryptococcus gattii. PLoS One 7:e43773.

54.   Garcia-Santamarina S, Festa RA, Smith AD, Yu C-H, Probst C, Ding C, Homer

CM, Yin J, Noonan JP, Madhani H, Perfect JR, Thiele DJ. 2018. Genome-wide

analysis of the regulation of Cu metabolism in *Cryptococcus neoformans*. Mol

Microbiol 108:473–494.

55.   Lev S, Kaufman-Francis K, Desmarini D, Juillard PG, Li C, Stifter SA, Feng CG,

Sorrell TC, Grau GER, Bahn Y-S, Djordjevic JT. 2017. Pho4 Is Essential for

Dissemination of *Cryptococcus neoformans* to the Host Brain by Promoting

Phosphate Uptake and Growth at Alkaline pH. mSphere 2:e00381-16.

56.   Jung WH, Sham A, White R, Kronstad JW. 2006. Iron Regulation of the Major

Virulence Factors in the AIDS-Associated Pathogen *Cryptococcus neoformans*.

PLoS Biol 4:e410.

57.   Chow EWL, Clancey SA, Billmyre RB, Averette AF, Granek JA, Mieczkowski P,

Cardenas ME, Heitman J. 2017. Elucidation of the calcineurin-Crz1 stress

response transcriptional network in the human fungal pathogen *Cryptococcus*

neoformans. PLOS Genet 13:e1006667.

58.   O'Meara TR, Norton D, Price MS, Hay C, Clements MF, Nichols CB, Alspaugh JA.

2010. Interaction of *Cryptococcus neoformans* Rim101 and Protein Kinase A

Regulates Capsule. PLoS Pathog 6:e1000776.

59.   Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, Li LX,

Williams M, Crouch EC, Khader SA, Brent MR, Doering TL. 2016. Computational

Analysis Reveals a Key Regulator of Cryptococcal Virulence and Determinant of

Host Response. MBio 7:e00313-16.

60.   Chun CD, Brown JCS, Madhani HD. 2011. A Major Role for Capsule-Independent

Phagocytosis-Inhibitory Mechanisms in Mammalian Infection by *Cryptococcus*

neoformans. Cell Host Microbe 9:243–251.

61.   Moranova Z, Virtudazo E, Hricova K, Ohkusu M, Kawamoto S, Husickova V,

Raclavsky V. 2014. The CRZ1/SP1-like gene links survival under limited aeration,

cell integrity and biofilm formation in the pathogenic yeast *Cryptococcus*

neoformans. Biomed Pap 158:212–220.

62.   Lev S, Desmarini D, Chayakulkeeree M, Sorrell TC, Djordjevic JT. 2012. The

Crz1/Sp1 Transcription Factor of *Cryptococcus neoformans* Is Activated by

Calcineurin and Regulates Cell Wall Integrity. PLoS One 7:e51403.

63.   Park H-S, Chow EWL, Fu C, Soderblom EJ, Moseley MA, Heitman J, Cardenas

ME. 2016. Calcineurin Targets Involved in Stress Survival and Fungal Virulence.

PLOS Pathog 12:e1005873.

64.   Lee D, Jang E-H, Lee M, Kim S-W, Lee Y, Lee K-T, Bahn Y-S. 2019. Unraveling

Melanin Biosynthesis and Signaling Networks in *Cryptococcus neoformans*. MBio

10:e02267-19.

65.   Rhodes J, Desjardins CA, Sykes SM, Beale MA, Vanhove M, Sakthikumar S, Chen

Y, Gujja S, Saif S, Chowdhary A, Lawson DJ, Ponzio V, Colombo AL, Meyer W,

Engelthaler DM, Hagen F, Illnait-Zaragozi MT, Alanio A, Vreulink J-M, Heitman J,

Perfect JR, Litvintseva AP, Bicanic T, Harrison TS, Fisher MC, Cuomo CA. 2017.

Tracing Genetic Exchange and Biogeography of *Cryptococcus neoformans* var.

grubii at the Global Population Level. Genetics 207:327–346.

66.   Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic

Genetic Analysis of Virulence in the Human Fungal Pathogen *Cryptococcus*

neoformans. Cell 135:174–188.

67.   Lee K-T, Hong J, Lee D-G, Lee M, Cha S, Lim Y-G, Jung K-W, Hwangbo A, Lee Y,

Yu S-J, Chen Y-L, Lee J-S, Cheong E, Bahn Y-S. 2020. Fungal kinases and

transcription factors regulating brain infection in *Cryptococcus neoformans*. Nat

Commun 11:1521.

68.   Krysan DJ, Zhai B, Beattie SR, Misel KM, Wellington M, Lin X. 2019. Host Carbon

Dioxide Concentration Is an Independent Stress for *Cryptococcus neoformans* That

Affects Virulence and Antifungal Susceptibility. MBio 10:e01410-19.

69.   D'Souza CA, Alspaugh JA, Yue C, Harashima T, Cox GM, Perfect JR, Heitman J.

2001. Cyclic AMP-Dependent Protein Kinase Controls Virulence of the Fungal

Pathogen *Cryptococcus neoformans*. Mol Cell Biol 21:3179–3191.

70.   Hu G, Steen BR, Lian T, Sham AP, Tam N, Tangen KL, Kronstad JW. 2007.

Transcriptional Regulation by Protein Kinase A in *Cryptococcus neoformans*. PLoS

Pathog 3:e42.

71.   Yoneda A, Doering TL. 2008. Regulation of *Cryptococcus neoformans* Capsule

Size Is Mediated at the Polymer Level. Eukaryot Cell 7:546–549.

72.   Frases S, Pontes B, Nimrichter L, Viana NB, Rodrigues ML, Casadevall A. 2009.

811   Capsule of *Cryptococcus neoformans* grows by enlargement of polysaccharide

812   molecules. Proc Natl Acad Sci U S A 106:1228–1233.

813   73.   Gaylord EA, Choy HL, Doering TL. 2020. Dangerous Liaisons: Interactions of

814   *Cryptococcus neoformans* with Host Phagocytes. Pathogens 9:891.

815   74.   Bailey TL. 2011. DREME: Supp. Bioinformatics 27:1653–1659.

816   75.   Summers DK, Perry DS, Rao B, Madhani HD. 2020. Coordinate genomic

817   association of transcription factors controlled by an imported quorum sensing

818   peptide in *Cryptococcus neoformans*. PLOS Genet 16:e1008744.

819   76.   Homer CM, Summers DK, Goranov AI, Clarke SC, Wiesner DL, Diedrich JK,

820   Moresco JJ, Toffaletti D, Upadhya R, Caradonna I, Petnic S, Pessino V, Cuomo

821   CA, Lodge JK, Perfect J, Yates 3rd JR, Nielsen K, Craik CS, Madhani HD. 2016.

822   Intracellular Action of a Secreted Peptide Required for Fungal Virulence. Cell Host

823   Microbe 19:849–864.

824   77.   Kang Y, Patel NR, Shively C, Recio PS, Chen X, Wranik BJ, Kim G, Scott McIsaac

825   R, Mitra R, Brent MR. 2020. Dual threshold optimization and network inference

826   reveal convergent evidence from TF binding locations and TF perturbation

827   responses. Genome Res 30:459–471.

828   78.   Giles SS, Stajich JE, Nichols C, Gerrald QD, Alspaugh JA, Dietrich F, Perfect JR.

829   2006. The *Cryptococcus neoformans* catalase gene family and its role in

830   antioxidant defense. Eukaryot Cell 5:1447–1459.

79.   Cox GM, Harrison TS, McDade HC, Taborda CP, Heinrich G, Casadevall A,

Perfect JR. 2003. Superoxide dismutase influences the virulence of *Cryptococcus*

neoformans by affecting growth within macrophages. Infect Immun 71:173–80.

80. Lee K-T, So Y-S, Yang D-H, Jung K-W, Choi J, Lee D-G, Kwon H, Jang J, Wang

LL, Cha S, Meyers GL, Jeong E, Jin J-H, Lee Y, Hong J, Bang S, Ji J-H, Park G,

Byun H-J, Park SW, Park Y-M, Adedoyin G, Kim T, Averette AF, Choi J-S, Heitman

J, Cheong E, Lee Y-H, Bahn Y-S. 2016. Systematic functional analysis of kinases

in the fungal pathogen *Cryptococcus neoformans*. Nat Commun 7:12766.

81. Caza M, Hu G, Price M, Perfect JR, Kronstad JW. 2016. The Zinc Finger Protein

Mig1 Regulates Mitochondrial Function and Azole Drug Susceptibility in the

Pathogenic Fungus *Cryptococcus neoformans*. mSphere 1.

82. Chen Y, Toffaletti DL, Tenor JL, Litvintseva AP, Fang C, Mitchell TG, McDonald

TR, Nielsen K, Boulware DR, Bicanic T, Perfect JR. 2014. The *Cryptococcus*

neoformans transcriptome at the site of human meningitis. MBio 5:e01087-13.

83. Alspaugh JA, Pukkila-Worley R, Harashima T, Cavallo LM, Funnell D, Cox GM,

Perfect JR, Kronstad JW, Heitman J. 2002. Adenylyl cyclase functions downstream

of the Gα protein Gpa1 and controls mating and pathogenicity of *Cryptococcus*

neoformans. Eukaryot Cell 1:75–84.

84. Heung LJ, Kaiser AE, Luberto C, Del Poeta M. 2005. The role and mechanism of

diacylglycerol-protein kinase C1 signaling in melanogenesis by *Cryptococcus*

neoformans. J Biol Chem 280:28547–28555.

85. Gerik KJ, Bhimireddy SR, Ryerse JS, Specht CA, Lodge JK. 2008. PKC1 is

essential for protection against both oxidative and nitrosative stresses, cell integrity,

and normal manifestation of virulence factors in the pathogenic fungus

Cryptococcus neoformans. Eukaryot Cell 7:1685–1698.

86. Waterman SR, Hacham M, Hu G, Zhu X, Park Y-D, Shin S, Panepinto J, Valyi-

Nagy T, Beam C, Husain S, Singh N, Williamson PR. 2007. Role of a CUF1/CTR4

copper regulatory axis in the virulence of *Cryptococcus neoformans*. J Clin Invest

117:794–802.

87. Jiang N, Sun N, Xiao D, Pan J, Wang Y, Zhu X. 2009. A copper-responsive factor

gene *CUF1* required for copper induction of laccase in *Cryptococcus neoformans*.

FEMS Microbiol Lett 296:84–90.

88. Walton FJ, Idnurm A, Heitman J. 2005. Novel gene functions required for

melanization of the human pathogen *Cryptococcus neoformans*. Mol Microbiol

57:1381–1396.

89. Cheon SA, Jung K-W, Chen Y-L, Heitman J, Bahn Y-S, Kang HA. 2011. Unique

Evolution of the UPR Pathway with a Novel bZIP Transcription Factor, Hxl1, for

Controlling Pathogenicity of *Cryptococcus neoformans*. PLoS Pathog 7:e1002177.

90. Hicks JK, Bahn YS, Heitman J. 2005. Pde1 phosphodiesterase modulates cyclic

AMP levels through a protein kinase A-mediated negative feedback loop in

Cryptococcus neoformans. Eukaryot Cell 4:1971–1981.

91. Kozubowski L, Lee SC, Heitman J. 2009. Signalling pathways in the pathogenesis

of *Cryptococcus*. Cell Microbiol 11:370–380.

92. Fox DS, Cruz MC, Sia RAL, Ke H, Cox GM, Cardenas ME, Heitman J. 2001.

Calcineurin regulatory subunit is essential for virulence and mediates interactions

with FKBP12-FK506 in *Cryptococcus neoformans*. Mol Microbiol 39:835–849.

93.   Odom A. 1997. Calcineurin is required for virulence of *Cryptococcus neoformans*.

EMBO J 16:2576–2589.

94.   Jung W-H, Son Y-E, Oh S-H, Fu C, Kim HS, Kwak J-H, Cardenas ME, Heitman J,

Park H-S. 2018. Had1 Is Required for Cell Wall Integrity and Fungal Virulence in

Cryptococcus neoformans. Genes|Genomes|Genetics 8:643–652.

95.   Kmetzsch L, Staats CC, Cupertino JB, Fonseca FL, Rodrigues ML, Schrank A,

Vainstein MH. 2013. The calcium transporter Pmc1 provides Ca2+ tolerance and

influences the progression of murine cryptococcal infection. FEBS J 280:4853–

4864.

96.   Kmetzsch L, Staats CC, Simon E, Fonseca FL, de Oliveira DL, Sobrino L,

Rodrigues J, Leal AL, Nimrichter L, Rodrigues ML, Schrank A, Vainstein MH. 2010.

The Vacuolar Ca2+ Exchanger Vcx1 Is Involved In Calcineurin-Dependent Ca2+

Tolerance and Virulence in *Cryptococcus neoformans*. Eukaryot Cell 9:1798–1805.

97.   Denham S, Brown J. 2018. Mechanisms of Pulmonary Escape and Dissemination

by *Cryptococcus neoformans*. J Fungi 4:25.

98.   Sun D, Sun P, Li H, Zhang M, Liu G, Strickland AB, Chen Y, Fu Y, Xu J, Yosri M,

Nan Y, Zhou H, Zhang X, Shi M. 2019. Fungal dissemination is limited by liver

macrophage filtration of the blood. Nat Commun 10:4566.

99.   Santiago-Tirado FH, Doering TL. 2017. False friends: Phagocytes as Trojan horses

in microbial brain infections. PLOS Pathog 13:e1006680.

100. Squizani ED, Oliveira NK, Reuwsaat JCV, Marques BM, Lopes W, Gerber AL, de

Vasconcelos ATR, Lev S, Djordjevic JT, Schrank A, Vainstein MH, Staats CC,

Kmetzsch L. 2018. Cryptococcal dissemination to the central nervous system

requires the vacuolar calcium transporter Pmc1. Cell Microbiol 20:e12803.

101. Okagaki LH, Nielsen K. 2012. Titan Cells Confer Protection from Phagocytosis in

Cryptococcus neoformans Infections. Eukaryot Cell 11:820–826.

102. Srikanta D, Santiago-Tirado FH, Doering TL. 2014. *Cryptococcus neoformans*:

historical curiosity to modern pathogen. Yeast 31:47–60.

103. Nielsen K, Cox GM, Wang P, Toffaletti DL, Perfect JR, Heitman J. 2003. Sexual

Cycle of *Cryptococcus neoformans* var. *grubii* and Virulence of Congenic a and α

Isolates. Infect Immun 71:4831–4841.

104. Reuwsaat JCV, Motta H, Garcia AWA, Vasconcelos CB, Marques BM, Oliveira NK,

Rodrigues J, Ferrareze PAG, Frases S, Lopes W, Barcellos VA, Squizani ED,

Horta JA, Schrank A, Rodrigues ML, Staats CC, Vainstein MH, Kmetzsch L. 2018.

A Predicted Mannoprotein Participates in *Cryptococcus gattii* Capsular Structure.

mSphere 3:e00023-18.

105. Haynes BC, Skowyra ML, Spencer SJ, Gish SR, Williams M, Held EP, Brent MR,

Doering TL. 2011. Toward an Integrated Model of Capsule Regulation in

Cryptococcus neoformans. PLoS Pathog 7:e1002411.

106.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput

Sequence Data.

107. Friedman RZ, Gish SR, Brown H, Brier L, Howard N, Doering TL, Brent MR. 2018.

Unintended Side Effects of Transformation Are Very Rare in *Cryptococcus*

neoformans. Genes|Genomes|Genetics 8:815–822.

108. Sedlazeck FJ, Rescheneder P, von Haeseler A. 2013. NextGenMap: fast and

accurate read mapping in highly polymorphic genomes. Bioinformatics 29:2790–1.

109. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence

Alignment/Map format and SAMtools. Bioinformatics 25:2078–2079.

110. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C,

Myers RM, Brown M, Li W, Liu XS. 2008. Model-based Analysis of ChIP-Seq

(MACS). Genome Biol 9:R137.

111. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C,

Singh H, Glass CK. 2010. Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell identities.

Mol Cell 38:576–89.

112. Novocraft. www.novocraft.com.

113. Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-

throughput sequencing data. Bioinformatics 31:166–169.

114. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

115. Ignatiadis N, Klaus B, Zaugg JB, Huber W. 2016. Data-driven hypothesis weighting

increases detection power in genome-scale multiple testing. Nat Methods 13:577–

580.

116. Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. Xylose donor transport is critical for fungal virulence. PLOS Pathog 14:e1006765.

ACKNOWLEDGMENTS

We appreciate helpful discussions with Charley Christian Staats, Augusto Schrank, and members of the Doering, Kmetzsch and Brent labs. We are grateful for assistance from Thomas Hurtaux, Eamim Squizani, and Julia Sperotto with mouse experiments; Guohua Chen with spotting assays; Jessica Plaggenberg with library preparation and sequencing; Chase Mateusiak with RNA-seq data analysis; and Sandeep Acharya for performing the DTO analysis. We thank Marilene Henning Vainstein and her lab for providing support with *in vivo* experiments. We also thank Liza Miller for comments on the manuscript and Arturo Casadevall for providing the antibody anti-GXM (18B7).

These studies were supported by National Institutes of Health grant AI087794 to TLD and MRB; National Institutes of Health grants AI36688 and AI140979 to TLD; and grants from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil – Grant number 310510/2018-0), and Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS, Brazil) to LK. CAPES fully supported JCVR during her studies in Brazil; her studies in the United States were partially supported by this source (Advanced Network of Computational Biology - RABICÓ - Biocomputational Grant 23038.010041/2013-13).

AUTHORS CONTRIBUTION

Conceived and designed experiments: J.C.V.R., D.P.A., A.L.C., M.R.B., L.K. and T.L.D.;

Performed experiments: J.C.V.R., D.P.A., H.M., H.B., and A.L.C;

Analyzed data: J.C.V.R., D.P.A., A.L.C., M.R.B., L.K. and T.L.D.;

Contributed reagents and materials: M.R.B., L.K. and T.L.D.;

Drafted the paper: J.C.V.R, L.K. and T.L.D.

Revised the paper: J.C.V.R., D.P.A., M.R.B., L.K. and T.L.D.

COMPETING INTERESTS

The authors declare no competing financial interests.

TABLES

Table 1. Pdr802 targets involved in quorum sensing and Titan cell formation.

Table 2. Genes regulated by Pdr802 involved in adaptation to the host

environment.

FIGURE LEGENDS

Figure 1. The transcription factor Pdr802 influences *C. neoformans* virulence. A.

Survival of C57BL/6 mice over time after intranasal inoculation with 5×10^4 cryptococci of

the strains indicated, with sacrifice triggered by weight below 80% of peak. B. Mean +/-

SD of total colony-forming units (CFU) in lung tissue at various times post-infection. CFU

inoculated for each strain were 41,400 (KN99α), 42,800 (*pdr802*) and 26,600 (*PDR802*).

p<0.05 for *pdr802* compared to the other strains at all time points.

Figure 2. *PDR802* expression is required for cell viability and induced during

growth in host-like conditions. A. Cells grown in DMEM at 37°C and 5% CO_2 were

sampled at the times indicated and plated on YPD to assess viability (measured by CFU

and plotted as fold-change from time 0). B. *PDR802* expression in KN99α cells grown in

DMEM at 37°C and 5% CO_2 was assessed by RNA-seq as in Li *et al*., 2018 (116).

Figure 3. The *pdr802* mutant is hypercapsular. A. Representative

immunofluorescence micrographs of the indicated strains after growth in DMEM (37°C,

5% CO_2) for 24 hours. The capsule was stained with monoclonal antibody anti-GXM 302

conjugated with Alexa 488 (green) and the cell wall with Calcofluor White (blue). All

images are to the same scale; scale bar, 5 µm. B. Capsule thickness distribution for the

indicated strains. C. Mean +/- SD of capsule size, quantified as detailed in the Methods

and Figure S5, with *pkr1* (39) and *ada2* (105) shown as hypercapsular and hypocapsular

controls, respectively. ****, p<0.0001 compared to KN99α by one-way ANOVA with

posthoc Dunnett test.

1003

Figure 4. Growth in mouse serum elicits increased capsule thickness and cell body

diameter in the *pdr802* mutant. A. Light micrographs of the indicated strains after

growth in mouse serum (at 37°C, 5% CO_2) for 24 h and negative staining with India ink to

visualize the capsule. All images are to the same scale; scale bar, 5 μm. B. Mean +/- SD

of capsule thickness, assessed by measuring at least 50 cells per strain with ImageJ. C.

Cells grown as in Panel A were plated on YPD to assess CFU. Mean +/- SD of the fold-

change compared to 0 h is shown. D. Mean +/- SD of cell body diameter, measured as in

B. ***, p<0.001 and ****, p<0.0001 for comparison of *pdr802* results to KN99α by one-

way ANOVA with posthoc Dunnett test.

1013

Figure 5. Absence of *PDR802* yields enlarged cells and loss of capsule induction in

the context of animal infection. A. India ink staining of fungi isolated from the lungs of

mice infected with the indicated strains. Numbers at left indicate the days post-infection.

All images are to the same scale; scale bar, 10 μm. B and C. Mean +/- SD of cell body

diameter (B) and capsule thickness (C), assessed by measuring at least 50 cells per

strain with ImageJ. ****, p<0.0001 and ***, p<0.001 for comparison of *pdr802* results to

KN99α or *PDR802* by one-way ANOVA with posthoc Dunnett test for each day post-

infection.

1022

Figure 6. Pdr802 is a negative regulator of Titan cell formation. Left, cultures were

subjected to *in vitro* conditions that induce Titan cell formation and imaged with India Ink.

All images are to the same scale; scale bar, 10 μm. Images were selected so that each

shows multiple examples of Titan cells, not to reflect abundance of this morphotype.

Right, the percent of Titan cells (TC) in each culture was quantified using flow cytometry,

gated as indicated by the blue square. FSC, forward scatter; SSC, side scatter.

1029

Figure 7. Deletion of *PDR802* affects phagocytosis after growth under conditions

that induce capsule and Titan cell formation. The indicated *C. neoformans* strains

were grown in YPD (18 h), DMEM (24 h), or Titan-cell induction medium (72 h) and then

incubated for 2 h with J774.16 mouse macrophages; host cells were then washed and

lysed to assess fungal burden by CFU. Data shown are normalized to the CFU of the

initial inoculum. *, $p < 0.05$ and **, $p < 0.01$ compared to KN99α by one-way ANOVA with

posthoc Dunnett test.

1037

Figure 8. Pdr802 participates in calcineurin signaling. *Panels A-B*. Interactions of

Pdr802 with upstream regions of the indicated genes. The ratios ($\log_2$) of reads from

immunoprecipitated (IP) DNA to input DNA were calculated for 1,000 bp upstream of the

first coding nucleotide (+1); shown is the difference in these values between tagged and

untagged strains. Black triangles, complete Pdr802 DNA-binding motifs (Figure S7B);

gray triangles, partial motifs. C. Intracellular calcium measurement by flow cytometry

using Fluo-4AM. Each column shows the mean and standard deviation of three biological

replicates. ***, $p < 0.001$ and ****, $p < 0.0001$ by one-way ANOVA with posthoc Dunnett

test.

1047

Figure 9. Pdr802 mode of action. *Left panel*. When wild-type *C. neoformans* enters a

host, *PDR802* expression is induced and Pdr802 positively regulates elements of the

quorum sensing pathway (described in the text) as well as expression of TFs implicated

in this pathway (*LIV3*), brain infectivity (*STB4*), and Titan cell production (*ZFC3*). At the

same time, Pdr802 regulates two calcineurin targets (*CRZ1* and *HAD1*) and a variety of

other genes (see text). Shown are examples of genes involved in the response to

oxidative stress (*SOD1*), growth at 37°C (*KIC1*), urease activity (*GUT1*), capsule

production (*BZP4*), and cell wall remodeling (*MP98*). *Right panel*. In the absence of these

regulatory changes, *pdr802* cells are poorly equipped to survive the stress of the host

environment and are subject to increased intracellular calcium levels, dysregulation of

capsule production, and impaired stress resistance. As a result, the cryptococcal

population in the lung is smaller and is enriched in Titan cells and hypercapsular cells of

normal size, both of which demonstrate reduced phagocytosis by host cells and impaired

ability to cross biological barriers; these defects reduce dissemination to the central

nervous system.

1063

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Mutant strain construction and confirmation. A. Scheme for generating *C.*

neoformans strains in the KN99α background (middle) that either lack *PDR802* (*pdr802,*

top) or encode a tagged copy of the protein (*mCherry-PDR802*). B. Qualitative analysis of

gene expression in Panel A strains and the complemented *pdr802* mutant (*PDR802*).

Cryptococcal mRNA isolated from cells grown in DMEM (37°C, 5% CO_2, 24 hours) was

used to generate cDNA; from this, segments of the genes indicated at the left were

amplified using the primers listed in Data Set S2, Sheet 5, and the products were

analyzed by agarose gel electrophoresis. Fragment sizes (in bp) are indicated at right

and the ladder bands shown are 400, 500, 650, 850 and 1000 bp for the top panel; 200,

300, 400, 500, and 650 bp for the middle panel; and 100, 200, and 300 bp for the bottom

panel. C. Quantitative analysis of *PDR802* expression. Samples of RNA isolated as in B

were analyzed for *PDR802* expression by qRT-PCR. All results were normalized to *ACT1*

expression. Each symbol represents a biological replicate, with the mean and standard

deviation also shown. ***, $p < 0.001$ compared to KN99α by one-way ANOVA with posthoc

Dunnett test.

1080

Figure S2. Organ burdens. A. Mean +/- SD values of total colony-forming units (CFU) in

the indicated tissue of mice from the Figure 1 survival curve are shown. Each point is the

average value for a single animal at the time of death. For *pdr802* infections, red circles

represent mice sacrificed at days 65 and 69, while blue circles represent mice sacrificed

at the end of the study (day 100). B. Mean +/- SD of total colony-forming units (CFU) in

the blood and brain at the indicated times post-infection. C. Mean +/- SD of total colony-

forming units (CFU) in the lung, blood and brain 75 days after infection with *pdr802*. Each

color represents one mouse.

1089

Figure S3. Characterization of *pdr802* cells. *Panels A-C.*10-fold serial dilutions of WT,

pdr802, and *PDR802* cells were plated on the media shown and incubated at 30°C (A),

37°C (B), or 37°C in the presence of 5% CO_2 (C). Nitrosative (NaNO$_2$) and oxidative

(H$_2$O$_2$) stress plates were prepared with YNB medium and melanization plates containing

L-DOPA were prepared as in the Methods; all other plates were prepared with YPD

medium. *lac1*, a control strain lacking the ability to melanize (88). D. Urease activity of the

indicated strains was evaluated using Christensen's urea solid medium (see Methods) at

the indicated temperatures. *ure1*, a control strain that does not produce urease (17).

1098

Figure S4. Growth curves and capsule shedding. *Panels A-B.* Growth of the strains

indicated in YPD at 30°C (A) or DMEM at 37°C and 5% CO_2 (B) was assessed by

OD_{600nm} at the times shown. C. Conditioned medium from the indicated strains was

probed for the presence of GXM after growth in DMEM for 24 or 48 hours. Equal volumes

of culture supernatant were analyzed without normalization to cell density.

Immunoblotting was performed using the anti-GXM monoclonal antibody 302.

1105

Figure S5. Semi-automated assay for cryptococcal capsule imaging. A. Schematic

of applying this method to cryptococcal cells induced to form capsule by growth in DMEM

(37°C, 5% CO_2) for 24 h, followed by cell wall and capsule staining. Thousands of cells

may be imaged per well and analyzed automatically with software that annotates and

measures the capsule (annotated on the micrograph in blue) and cell wall (annotated in

bright green). See Methods for details. B. Capsule size distribution of WT cells after

induction. Capsule thickness for each cell is the difference between the paired diameters

of the cell wall and capsule, which is plotted here with reference to the mean value. C

and D. Mean and SD (C) and cumulative percentage (D) analysis of WT compared to

hyper and hypocapsular control strains (here *pkr1* and *ada2*, respectively). Capsule

thickness is in arbitrary units, related to the pixels measured. E. The time required to

analyze the capsule thickness of 1,000 cells by this method compared to manual

assessment of India ink images.

1119

Figure S6. *PDR802* deletion induces Titan cell formation. Mean +/- SD of (A) total cell

diameter and (B) the ratio of total cell to cell body diameters (diameter ratio), assessed by

measuring at least 50 cells per strain with ImageJ. **, $p<0.01$ and ****, $p<0.0001$ for

comparison of *pdr802* to KN99α or *PDR802* by one-way ANOVA with posthoc Dunnett

test for each day post-infection. C. Percent of Titan cells in the indicated strain, evaluated

using various published parameters: cell body diameter above 10 or 15 µm (20) or total

cell diameter above 30 µm (43).

1127

Figure S7. Pdr802 strain viability, putative DNA-binding motifs, and self-regulation.

A. The indicated strains were grown in DMEM at 37°C and 5% CO_2 for the times shown

and samples were tested for their ability to form colonies on YPD medium. Plotted is the

fold-change in CFU relative to the initial culture. B. Putative Pdr802-binding motifs

determined using DREME (74). Primary and secondary hits are shown for analysis of

1,000 bp upstream of the initiating ATG. C. Pdr802 self-regulation. The ratios (log_2) of

reads from immunoprecipitated (IP) DNA to reads from input DNA were calculated for

1,000 bp upstream of the first coding nucleotide (+1) of *PDR802*; shown is the difference

in these values between tagged and untagged strains. Red triangles, complete Pdr802

DNA-binding motifs (Figure S7B); blue triangles, partial motifs.

1138

Data Set S1. Pdr802 ChIP-Seq, RNA-Seq, and Dual-Threshold Optimization (DTO)

data. Sheet 1, peaks that occur in gene promoter regions that showed ≥2-fold enrichment

when Chip-Seq was performed on strains expressing tagged versus untagged Pdr802,

with annotation. Sheet 2, ChIP-Seq primary data of Pdr802-specific peaks. Sheet 3,

RNA-Seq data. Sheet 4, DTO filtered data. Sheet 5, DTO primary data. Sheet 6, ChIP-

Seq data of all peaks in mCherry-Pdr802 samples. Sheet 7, ChIP-Seq data of all peaks in

untagged (WT) control samples.

Data Set S2. Pdr802 target analysis and primers used in this study. Sheet 1, Pdr802

regulated genes that are down-regulated during Titan cell formation *in vitro* (48). Sheet 2,

Pdr802 regulated genes that are up-regulated during Titan cell formation *in vitro* (48).

Sheet 3, targets regulated by Pdr802 that are also dephosphorylated by calcineurin (63).

Sheet 4, the intersection of Pdr802 targets and genes that are Crz1-independent

calcineurin targets under conditions thermal stress (55). Sheet 6, primers used in this

study. Fold enrichment and adjusted p (q) values throughout are for the DTO results.

Supplementary Methods.

Table 1. Pdr802 targets involved in quorum sensing and Titan cell formation.

Biological process	CNAG	Gene name	ChIP-seq[1]	RNA-seq[2]	Description
Quorum sensing	00150	*PQP1*	1.38	-0.78	Peptidase
	03013	*OPT1*	1.25	-0.55	OPT small oligopeptide transporter
Titan cell formation	05835	*LIV3*	1.52	-0.84	Transcription factor
	05785	*STB4*	3.33	-1.71	Transcription factor
	05940	*ZFC3/CQS2*	2.23	0.68	Transcription factor

[1]Fold-change for mCherry-Pdr802 compared to WT
[2]$\log_2$ fold-change for *pdr802* compared to WT

Table 2. Genes regulated by Pdr802 involved in adaptation to the host environment.

Biological process	CNAG	Gene name	ChIP-seq[1]	RNA-seq[2]	Description
Oxidative stress resistance	04981	CAT1	1.47	-1.60	Catalase 1
	05256	CAT2	1.41	-0.42	Catalase 2
	01019	SOD1	2.59	-0.53	Superoxide dismutase [Cu-Zn]
	00896	FZC34	2.22	-0.82	Transcription factor
	06327	MIG1	1.72	-0.78	DNA-binding protein creA
	00556	CCK1	1.78	-0.48	Casein kinase I
Melanin and cell wall formation	03202	CAC1	1.60	-0.61	Adenylate cyclase
	01845	PKC1	2.00	-0.48	AGC/PKC protein kinase
	07724	CUF1	1.27	-0.67	Metal-binding regulatory protein
	00740	SNF5	1.28	-0.70	Swi/snf chromatin-remodeling subunit
	01239	CDA3	2.63	1.08	Chitin deacetylase 3
	01230	MP98	2.74	0.96	Chitin deacetylase 2
	00776	MP88	1.86	1.23	Immunoreactive mannoprotein
Growth at 37°C	00405	KIC1	1.88	-0.97	Ste/ste20/ysk protein kinase
	03670	IRE1	1.66	-1.14	IRE protein kinase
Urease activity	07448	DUR3	2.13	-3.05	Urea transporter
	01209	FAB1	2.29	-0.49	1-phosphatidylinositol-3-P 5-kinase
	01938	KIN1	1.61	-0.34	CAMK/CAMKL/KIN1 protein kinase
	01155	GUT1	2.96	0.62	Glycerol kinase
	00791	HLH1	1.71	-1.09	Transcription factor
Capsule thickness	02802	ARG2	1.56	-0.59	Inositol/phosphatidylinositol kinase
	06809	IKS1	2.28	-0.44	IKS protein kinase
	01905	KSP1	2.02	-0.64	Serine/threonine protein kinase
	02877	FZC51	1.75	-0.69	Transcription factor
	07470	PDE2	2.42	-1.80	High-affinity phosphodiesterase
	03346	BZP4	3.52	1.30	Transcription factor
Calcineurin signaling	00156	CRZ1	1.44	-0.49	Transcription factor
	01744	HAD1	2.26	0.47	Phosphatase

[1]Fold-change for mCherry-Pdr802 compared to WT
[2]Log$_2$ fold-change for *pdr802* compared to WT

A

B

A

B

A

B

C

A

B

C

D