Elucidation of global and local epidemiology of *Salmonella* Enteritidis through multilevel genome typing

Lijuan Luo[1], Michael Payne[1], Sandeep Kaur[1], Dalong Hu[1], Liam Cheney[1], Sophie Octavia[1], Qinning Wang[2], Mark M. Tanaka[1], Vitali Sintchenko[2,3] and Ruiting Lan[1,*]

[1]School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, New South Wales, Australia

[2]Centre for Infectious Diseases and Microbiology–Public Health, Institute of Clinical Pathology and Medical Research – NSW Health Pathology, Westmead Hospital, New South Wales, Australia

[3]Marie Bashir Institute for Infectious Diseases and Biosecurity, Sydney Medical School, University of Sydney, New South Wales, Australia

*Corresponding Author

Email: r.lan@unsw.edu.au

Phone: 61-2-9385 2095

Fax: 61-2-9385 1483

Abstract

Salmonella Enteritidis is a major foodborne pathogen that causes both local and international outbreaks

with complex transmission pathways. Invasive infections which are associated with multidrug

resistance are an increasing concern around the globe. However, the global epidemiological picture of

S. Enteritidis remains unclear due to the lack of a fine scale typing scheme based on genomic sequencing

data. Here, using the novel multilevel genome typing (MGT) approach, we have characterised the

genomic epidemiology of *S.* Enteritidis in unpreceded detail. We examined 26,670 publicly available

S. Enteritidis whole genome sequences from 86 countries over 101 years to reveal their spatial and

temporal distributions. Using MGT4 and MGT5, we identified globally prevalent and regionally

restricted STs. Source associated STs were identified, such as poultry associated MGT4-STs, which

were common in human cases in the USA. Temporal trends were observed in the UK with MGT5-STs

from 2014 to 2018, revealing both long lived endemic STs and the rapid expansion of new STs. Using

MGT3 to MGT6, we identified MDR associated STs to facilitate tracking MDR spread. The majority

of the global *S.* Enteritidis population fell within two predominant lineages with significant differences

in geographic distribution, outbreak frequency, antimicrobial resistance and mutation rate. An online

open MGT database has been established for unified international surveillance of *S.* Enteritidis. We

demonstrated that MGT provides a flexible and high-resolution genome typing tool for *S.* Enteritidis

surveillance and outbreak detection.

Keywords: *Salmonella* Enteritidis, MGT, global epidemiology, multidrug resistance, population

structure, mutation rate

Introduction

The infection of nontyphoidal *Salmonella* ranked as the second most common foodborne disease in

Europe and North America [1, 2], with *Salmonella* Enteritidis a dominant serovar in many countries [3,

4]. *S.* Enteritidis mainly causes human gastrointestinal infections leading to diarrhoea. However,

invasive infections manifesting as sepsis, meningitis and pneumonia, have recently been reported [5,

6]. As many *S.* Enteritidis strains causing invasive infections carry a plasmid which confers multidrug

resistance (MDR), the spread of these strains internationally is a major threat to global health [7, 8].

There has been limited studies of global epidemiology of *S.* Enteritidis genome types [5, 9] . And *S.*

Enteritidis has caused large-scale national and international outbreaks with complex transmission

pathways [3, 10, 11]. The geographic distribution, outbreak propensity, MDR and evolutionary

characteristics of different lineages of *S.* Enteritidis have not been systematically evaluated despite the

rapid expansion of global publicly available genome sequences. A rapid, stable and standardized global

genomic typing strategy for *S.* Enteritidis is required for the high-resolution and scalable surveillance

of outbreaks, tracking of international spread and MDR profiles of *S.* Enteritidis.

Single Nucleotide Polymorphism (SNP) address and Hierarchical Clustering of cgMLST (HierCC),

based on core genome multi-locus sequence typing (cgMLST), have been useful hierarchical genomic

typing methods for *S.* Enteritidis [12, 13]. Both SNP address and HierCC assign a series of cluster types

for each isolate based on a range of genetic distance thresholds. These cluster types, which are generated

based on the single-linkage clustering, can merge when additional data is added [14]. Thus, cluster

typing systems may not be stable when new isolates are added, which is problematic for long term

surveillance, especially for large databases in which the number of isolates is expanding quickly. In

contrast, sequence types (STs) are based on exact matching of genes between isolates, and thus are

stable, so that once a ST is assigned to an isolate it will never change [14]. It would be useful to develop

a surveillance system where each isolate is assigned with both stable STs and cluster types at varied

subtyping resolution.

Multilevel genome typing (MGT) is a novel genome sequence based typing method developed by

Payne *et al.* [14]. Based on the well-established multi locus sequence typing (MLST) concept [15, 16],

MGT is defined by multiple MLST schemes containing different numbers of loci which result in different resolutions. An MGT scheme and a public database were established for *Salmonella* Typhimurium where 7 to 5268 loci were included in MGT1 to 9 [14]. MGT1 corresponds to the classic 7-gene MLST scheme of *S. enterica* [16]. For each isolate, sequence types (STs) and clonal complexes (CCs) were called at each MGT level [14]. Thus, the subtyping resolution was gradually increased, which allowed isolates to be grouped together at different scales [14]. The stable STs at multiple MGT levels were found to be able to identify DT104, the MDR lineage of *S.* Typhimurium, as well as African invasive lineages [14]. MGT of *S.* Typhimurium also performed well in outbreak identification and source tracing [14].

Here, we describe an MGT scheme and public database for *S.* Enteritidis. By using MGT, we systematically examine the global and local epidemiology of *S.* Enteritidis over time, and evaluate the application of this scheme to the detection of outbreak clusters and the monitoring of multidrug resistant STs.

Results

Establishing the MGT system for *S.* Enteritidis

Based on 99% of coverage and 96% of identity (**Supplementary Fig. S1**), we defined the *S.* Enteritidis core genome, which included 3932 genes with 977 not found in the *Salmonella* core. Additionally, 1054 *S.* Enteritidis core intergenic regions were also defined (**Fig. 1a**) (**Supplementary Table S1**).

The first 8 schemes of *S.* Enteritidis MGT used the same 2955 core genes of *Salmonella* as the MGT of *S.* Typhimurium defined by Payne *et al*. [14]. The rationale of loci selection has been fully described previously [14]. MGT1 corresponded to the classic 7 gene MLST of *Salmonella* [16]. A total of 4986 loci were incorporated in the MGT9 scheme, including the 3932 *S.* Enteritidis core genes and 1054 core intergenic regions defined (**Fig. 1a**).

A total of 26,670 *S.* Enteritidis genomes with publicly available raw reads, collected from 26 source types in 86 countries during 1917 to 2018 (**Supplementary Fig. S2**), were analysed using the MGT scheme. At each MGT level, each isolate is assigned an ST and a CC. A CC is a group of STs with a

genetic distance threshold of one allele difference, as defined in the classic 7 gene MLST [15, 16]. The number of STs and CCs at each MGT level is shown in **Fig. 1b.** As the resolution of typing increased from MGT1 to MGT9, an increasing number of STs and CCs were assigned from MGT2 to MGT9. At MGT2, the 26,670 isolates were subtyped into 252 STs and 23 CCs. In contrast, MGT9 divided the isolates into 20,153 different STs and 14,441 CCs. The MGT scheme of *S.* Enteritidis is available through a global public online database (https://mgtdb.unsw.edu.au/enteritidis/).

The global epidemiology of *S.* Enteritidis by MGT

Of the 26,670 isolates typed by MGT, 25,207 have country metadata. By the 7-gene MLST (or MGT1) scheme, ST11 was the dominant type representing 94.8% of the isolates, followed by ST183 of 1.6%. ST183 is an endemic ST prevalent in Europe and was divided into two main phage types, PT11 and PT66 [17]. Using MGT, ST183 (or MGT1-ST183) can be divided into 7 STs at MGT2 level. Interestingly, 97% (137/141) of the PT11 isolates belonged to MGT2-ST3, and 100% (24/24) of the PT66 belonged to MGT2-ST82, highlighting the potential for backward compatibility of MGT with traditional typing data.

For the predominant ST11 isolates (or MGT1-ST11), we found that the optimal MGT level to describe their global epidemiology was MGT4, based on the distribution of each ST in different continents (**Supplementary Fig. S3**). At the MGT4 level, the 26,670 isolates were subtyped into 2,236 STs and 423 CCs. Among the 2,236 MGT4-STs, 163 STs were predominantly found in only one continent (defined as ≥70% from one continent). These 163 STs contained 20,341 of the 26,670 genomes (76.3%).

The distribution of STs varied between continents. MGT4-ST99, ST136, ST135, ST171 and ST163 were the most prevalent STs in North America (**Fig. 2**), while in Europe, MGT4-ST25, ST13, ST100 and ST31 were common. However, the North America and Europe *S.* Enteritidis sequences analysed were mostly obtained from the USA and UK, which represented 94.6% and 92.5% of isolates from these two continents. In Africa, MGT4-ST11 was more prevalent in West Africa and MGT4-ST16 in Central and Eastern Africa (**Supplementary Fig. S4**). These STs reflect the previously reported African

invasive infection lineages (West African clade and Central/Eastern African clade) [5]. Finally, MGT4-ST15 was a global ST which was observed in all continents.

These dominant STs can be further grouped into CCs which offered a more inclusive picture. Among the 423 MGT4-CCs at MGT4, 10 MGT4-CCs represented 94.1% of the all isolates. The top two CCs, MGT4-CC1 and MGT4-CC13 accounted for 88.0% of the isolates. MGT4-CC1 was prevalent in all six continents (**Fig. 2**). MGT4-CC13 was more common in North America and Europe. MGT4-CC101 was observed in different continents, with higher proportion in Asia. MGT4-CC11 and CC16 describe the African prevalent lineages that were associated with MDR and bloodstream-invasive infection [5].

National and local epidemiology of *S*. Enteritidis by MGT

As the majority of the *S*. Enteritidis sequences analysed in this study were from the USA and UK, we compared distribution of the STs between these two countries. We used MGT4 and MGT5 levels to describe the data. A total of 39 MGT4 STs with more than 50 isolates represented 75.1% of the USA and UK isolates and each country had its own specific types (**Fig. 3a**). MGT4-ST99, ST135 and ST170 were prevalent in the USA, whereas MGT4-ST15, ST25 and ST13 were more prevalent in the UK. MGT4-ST15 and ST25 were further subtyped into 34 MGT5-STs (more than 20 isolates each), of which the majority were prevalent in the UK, except for MGT5-ST412 and ST387 that were more prevalent in the USA (**Fig. 3b**).

To examine the relationship between MGT-STs, isolation source and location, we examined MGT4-STs of 4383 genome sequences from the USA which contained source and state metadata (**Fig. 4**). Of the 4383 genomes, 46.7% were from avian source while 43.1% were from humans. Six MGT4 STs (with more than 50 isolates each) were isolated from both human and avian sources including MGT4-ST99, ST135, ST136, ST160, ST198 and MGT4-ST25, while 3 STs including MGT4-ST15, ST163 and MGT4-ST170 were mainly from human sources. All the human only STs belonged to MTG4-CC1 whereas the mixed source STs were mostly of MGT4-CC13 origin (**Supplementary Table S2**). STs belonging to MGT4-CC13 were significantly more prevalent in avian sources than STs of MGT4-CC1 (P < 0.001, OR = 45.9) (**Supplementary Table S2**). *S*. Enteritidis isolate data from 48 states of the USA were available and ST frequencies were similar across the country (**Fig. 4**). In almost all states,

MGT4-ST99 was the dominant type, followed by MGT4-ST135 and MGT4-ST136, all of which belonged to MGT4-CC13.

UK isolates contained collection year and month metadata. There were 8,818 human *S.* Enteritidis isolates collected from March 2014 to July 2018 in the UK. We chose MGT5-STs to describe the monthly variation of *S.* Enteritidis in the UK. By MGT5, variation in the prevalence of MGT5-STs in different years and months was observed (**Fig. 5**). There were 13 MGT5-STs with more than 100 isolates, representing 46.3% of the 8,818 isolates. The top 5 STs over the entire period were MGT5-ST1, ST29, ST79, ST15 and MGT5-ST33, representing 30% of the total isolates. These STs showed differing temporal patterns. MGT5-ST1, ST29, ST79 and ST15 were consistently observed in each month across these four years while MGT5-ST15 included 94% of isolates previously reported as part of an outbreak [18]. MGT5-ST156 first appeared in April of 2012 in the UK, increased in frequency in June and July of 2014, and became rare after 2014. MGT5-ST156 contained all isolates reported as involved in an international outbreak in Europe [19] which explains the relatively short duration of this large ST. MGT5-ST423 was a dominant type from May to September of 2015, then became rare and was dominant again during the September to December of 2016.

Detection of potential large outbreak clusters of *S.* Enteritidis using MGT

MGT9 offered highest resolution for spread tracking and outbreak detection. By MGT9, there were 124 MGT9-STs including isolates from two or more countries each, indicating international spreading (**Supplementary Table S3**).

To detect potential outbreak clusters and to assess whether some STs were more likely to cause outbreaks, we identified clusters of isolates using MGT9 with no more than 2 allele differences which we refer as ODC2. At ODC2, the 26,670 isolates were grouped into 11,174 different ODC2 clusters, with 331 clusters each having more than 10 isolates (10645 isolates in total) and 42 large clusters each having more than 50 isolates (5240 in total). Further analysis of the 42 large clusters showed that these isolates were distributed into 13 STs, including MGT4-ST136 (6/42, 14%), ST25 (8/42, 19%), ST163 (2/42, 5%), ST171 (3/42, 7%), ST135 (3/42, 7%) and ST13 (3/42, 7%). Among the 42 large clusters, 71% (30/42) belonged to MGT4-CC13 representing 72% of the 5240 isolates in large clusters while 24%

(10/42) belonged to MGT4-CC1 representing 24% of isolates. MGT4-CC13 was found to be significantly associated with larger clusters of ODC2 (>= 50 isolates) compared to MGT4-CC1 (P < 0.001, OR = 2.26) (**Supplementary Table S4**).

Since the global data may be biased towards outbreak isolates that were preferentially sequenced, we used UK data from 2014 which included all human isolates in that year in their collection [20]. At MGT4, there were 17 ODC2 clusters of more than 50 isolates representing 1855 isolates in total, 88% of which belonged to 12 different MGT4-STs (**Supplementary Table S5**). MGT4-ST15 was the dominant type in clinical infections in UK, however only one ODC2 cluster of 62 isolates was found in this ST making up 3% of its isolates. MGT4-ST15 was significantly less likely to have large ODC2 clusters than the other STs (OR = 0.14, P value < 0.001). In contrast, MGT4-ST25, which was the second dominant type in clinical isolates in UK, 39% of the isolates belonged to 5 different ODC2 clusters of more than 50 isolates. MGT4-ST25 contained larger ODC clusters than other STs (OR = 4.31, P value < 0.001). The previous result showing MGT4-CC13 as more correlated with larger ODC2 clusters relative to MGT4-CC1 was also confirmed in the UK dataset (P < 0.001, OR = 6.24).

Antibiotic resistance (AR) profiling of *S.* Enteritidis

As nearly all of the isolates (100.0%, 26,665/26,670) harboured the aac(6')-Iaa_1 gene for aminoglycoside resistance, this gene was excluded for the antibiotic resistance analysis. A total of 2505 isolates (9.4% of the total 26,670 isolates) were found to harbor antibiotic resistant genes excluding for aac(6')-Iaa_1. The most frequent predicted class was tetracyclines (5.1%, 1,350/26,670), followed by beta-lactams (4.5%, 1,213/26,670) and aminoglycosides (2.8%, 741/2270). Among the 2505 isolates, 40.4% (1011/2505) of the AR isolates were MDR, harbouring genes conferring resistance to 3 or more different antibiotic classes. A total of 59.6% (1494/2505) of the AR isolates harboured genes conferring resistance to 1 to 2 different antibiotic classes. Although the selection of isolates for genome sequencing in some continents may be biased, African isolates were found to have highest proportion of AR/MDR isolates, followed by Asia and Oceania (**Supplementary Fig. S5**).

STs containing more than 10 isolates at different MGT levels were screened to identify MDR or AR associated STs (defined as >= 80% MDR isolates). Eleven STs from varied levels were identified as

MDR STs, representing 49% (659/1011) of the MDR isolates. These STs were mutually exclusive at different MGT levels. The top two STs, MGT3-ST15 and ST10, were two types that were prevalent in Africa. For MGT3-ST15, 99.2% of the isolates harboured resistance genes corresponding to as many as six different drug classes (**Table 1**). For MGT3-ST10, 86.5% of the isolates were MDR, and the antibiotic resistant patterns were similar to those of MGT3-ST15. MGT3-ST10 and ST15 represented 96.3% and 90.2% of the two Africa endemic lineages (MGT3-CC10 and CC15, respectively) as mentioned below. Among the other 9 MDR STs, 8 belonged to MGT4-CC1, and 93.5% of the isolates (243/260) harboured genes conferring resistance to as many as 8 classes of antibiotic drugs. Only one MDR ST (MGT6-ST2698) belonged to MGT4-CC13 with 89.7% (35/39) of the isolates harboured genes conferring resistance to 4 drug classes. MGT4-CC1 had significantly more isolates with MDR genes (5.6%, 474/8539) than MGT4-CC13 (0.7%, 107/14972) (Fisher exact test, P value < 0.001, OR=12.4).

A total of 707 isolates belonging to 47 different STs from MGT3 to MGT7, harboured genes conferring resistance to one or two different antibiotic classes including aminoglycosides, beta-lactams, tetracyclines and quinolones (**Supplementary Table S6**). Among the 47 STs, 34 STs (72%, 34/47) belonged to MGT4-CC1, representing 80% (564/707) of isolates, and 12 (26%, 12/47) STs belonged to MGT4-CC13 representing 18% (125/707) of the isolates. Again MGT4-CC1 had significantly more isolates with AR genes than MGT4-CC13 (Fisher exact test, P value < 0.001, OR=8.4).

Plasmid specific genes were identified from the PlasmidFinder database [21] and IncQ1, IncN, IncI1, IncX1plasmids were common in S. Enteritidis isolates with AR genes (**Supplementary Fig. S6**). Particularly, plasmid type IncQ1 was present in MGT3-ST15, ST10 and ST30, which harboured MDR genes up to 6 drug classes (**Table 1**). Plasmid type IncI1 was common in MGT3-ST10 and ST161 and plasmid type IncX1 was more common in the STs of MGT4-CC1.

Table 1. Antibimicrobial drug class and plasmid class of the MDR associated STs from differnet MGT levels.

MGT-ST *	MGT4-CCs	Total No. of isolates	No. of isolates with MDR (%)	% of the MDR isolates							% of isolates with plasmid									Continent ****
				Aminoglycoside	Beta-lactam	Sulphonamide	Tetracycline	Phenicol	Trimethoprim	Macrolide	Col156_1	ColRNAI_1	ColpVC_1	IncI1_1_Alpha	IncI2_1_Delta	IncN_1	IncQ1_1	IncX1_1	IncX1_4	
MGT3-ST15	CC16 ***	261	259 (99%)	98%	74%	97%	89%	95%	96%			1%		14%	1%		95%			Africa
MGT3-ST10	CC11 ***	141	122 (87%)	74%	84%	86%	82%	77%	82%			1%		95%	1%	4%	71%			Africa
MGT3-ST30	CC1	10	10 (100%)	100%		100%	100%	100%	100%								80%			EU
MGT3-ST161**	CC1	79	29 (37%)	30%	84%	35%	84%	32%	3%	3%	1%		100%	86%				1%		EU
MGT3-ST9	CC1	70	66 (94%)	93%	80%	91%	80%	1%	1%	1%	1%			3%					99%	EU/NA
MGT3-ST301	CC1	11	11 (100%)	100%	100%				100%									100%		NA
MGT4-ST354	CC1	39	39 (100%)	100%	100%	100%	62%				3%			5%					100%	NA/EU
MGT4-ST54	CC1	29	28 (97%)	90%	41%	90%	86%												97%	EU/NA/OC
MGT5-ST86	CC1	12	12 (100%)	100%	100%	100%								8%					100%	NA/EU
MGT5-ST682	CC1	10	10 (100%)	100%		100%	100%	90%						30%		100%		100%		NA
MGT6-ST2698	CC13	39	35 (90%)	90%	90%	90%	90%										90%	100%		NA
Total		701	621 (89%)																	

* MGT-STs with more than 80% of the isolated harboring AR genes to 3 or more drug types were identified. Isolates belonging to the MDR associated MGT-STs didn't overlap. Note aac(6')-Iaa_1 gene for aminoglycoside resistance, which was present in almost all isolates, was excluded for the AR analysis.

** MGT3-CC161 include 29 isolates with MDR genes (to 5 drug types) and 38 isolates with AR genes to 2 different drug types.

*** MGT4-CC16 and CC11 belonged to MGT3-CC15 and CC10, which were representative of the two African lineages.

**** The main continents that the MDR related STs were observed: EU refers to Europe, NA to North America and OC to Oceania.

The population structure and evolution of major *S*. Enteritidis STs/CCs

The majority of the STs from different MGT levels analysed mainly belonged to two MGT4 CCs, MGT4-CC1 and CC13. To describe the global phylogenetic structure of *S*. Enteritidis and explore the phylogenetic relationship of the two lineages, 1506 representative isolates were selected using representatives of MGT6-STs to encompass the diversity of the serovar. A previous study had suggested that *S*. Enteritidis has three clades, A, B and C. Clade A and C appeared to diverge earlier and were phylogenetically more distant to the global clade B than *S*. Gallinarum [9] (**Fig. 6a**). *S*. Enteritidis clade B and *S*. Gallinarum were sister clades. The vast majority of the genomes analysed in this study belonged to clade B.

Within clade B, we identified 10 main lineages, which were concordant with MGT-CCs from MGT1 to MGT4 (**Fig. 6b**). These lineages can be described at different MGT levels as shown in **Fig. 6b**. The lineages were consistent with the progressive division from lower to higher MGT levels grouped by CCs. MGT2-CC3 represent 98.8% of the Europe endemic MGT1-ST183 lineage [17]. At MGT3, there were 4 lineages named MGT3-CC10, CC15, CC18 and CC107. MGT3-CC10 and CC15 represented 100% of the two previously reported Africa lineages associated with invasive infection [5]. At MGT4, there were 5 lineages with two, MGT4-CC1 and MGT4-CC13, representing 88.0% of all 26,670 *S*. Enteritidis isolates. MGT4-CC30 and CC129, both of which were phylogenetically closer to MGT3-CC13, were two endemic lineages in Europe and North America, respectively. The population structure of *S*. Enteritidis defined by MGT, were generally in accordance with cgMLST HierCC HC100 (**Supplementary Fig. S7**).

We performed BEAST analysis using the *S*. Enteritidis core genome variation to determine the evolutionary rate of the clade B *S*. Enteritidis which was estimated to be 1.9×10^{-7} substitution/site/year (95% CI of 1.6×10^{-7} to 2.3×10^{-7}), corresponding to 0.8 SNPs per genome per year for the core genome (95% CI of 0.6 to 0.9 SNPs). The mutation rate of MGT4-CC13 lineage core genome was estimated to be 2.5×10^{-7} substitution/site/year (95% CI of 2.3×10^{-7} to 2.7×10^{-7}) or 1.0 SNP per genome per year (95% CI of 0.9 to 1.1 SNPs). The mutation rate of MGT4-CC1 lineage core genome was 1.7×10^{-7} substitution/site/year (95% CI of 1.6×10^{-7} to 2.0×10^{-7}), or 0.7 SNP per genome per year (95% CI of 0.6

to 0.8 SNPs). The mutation rate of MGT4-CC13 was significantly faster (1.44 times) than that of MGT4-CC1.

The most recent common ancestor (MRCA) of the 9 lineages belonging to MGT2-CC1, was estimated to have existed in the 1460s (95% CI 1323 to 1580) (**Supplementary Fig. S8**). The two global epidemic lineages MGT4-CC1 and CC13 are estimated to have diverged in around 1687 (95% CI 1608 to 1760). In 1869 (95% CI=1829-1900), MGT4-CC13 diverged into two sub-lineages of various MGT4-STs, with one more prevalent in North America than in other continents (labelled with red arrow) and the other sub-lineage more prevalent in Europe (labelled with blue arrow). We further estimated the population expansion of the two global epidemic lineages MGT4-CC1 and CC13 (**Supplementary Fig. S9**). For MGT4-CC1, there were two big expansions around 1950 and 1970. For MGT4-CC13, the population gradually increased until a more rapid expansion around 1970.

Discussion

S. Enteritidis is one of the most common foodborne pathogens causing large scale national and international outbreaks and food recalls [3, 22]. Understanding the population structure and genomic epidemiology of *S.* Enteritidis is essential for its effective control and prevention. In this study, we developed a new genomics tool, MGT, for *S.* Enteritidis and developed a database for international applications to achieve these objectives. We used MGT to describe its local and global epidemiology, and its population structure using 26,670 publicly available genomes and associated metadata

MGT for *S.* Enteritidis uncovers geographic, source and temporal epidemiological trends of *S.* Enteritidis within and between countries

A total of 26,670 isolates were successfully typed using the MGT scheme which allowed the examination of the local and global epidemiology of *S.* Enteritidis. The salient feature of the flexibility of the different levels of MGT has also been illustrated through this analysis. The lower resolution levels of MGT were found to be able to effectively describe the geographic variation in different continents, countries or regions. Of these levels we showed that MGT4-STs best described the global epidemiology of *S.* Enteritidis at continent level with some STs widely distributed in different continents while some others restricted in distribution. MGT4-ST15 was a global epidemic type which was prevalent in almost

all continents. In contrast, MGT4-ST16 (also described by MGT3-ST15) was prevalent in Central/Eastern Africa while MGT4-ST11 (also described by MGT3-ST10) was prevalent in West Africa, which agreed with a previous study [5].

In the USA, MGT4-CC13 and MGT-CC1 showed remarkable difference in their epidemiology. MGT4-CC13 STs including MGT4-ST99, ST136 and ST135 were the main cause of clinical infections and were common in poultry. The prevalent STs were generally similar in different states. In particular, MGT4-ST99, ST136 and ST135 were the dominant types in almost all states of the USA. These STs may be associated with poultry [23], as there were multistate outbreaks caused by the *S.* Enteritidis contaminated eggs [1]. Since poultry related products (i.e. eggs, chicken and turkey, especially eggs) are known to be the main source of *S.* Enteritidis infections or outbreaks in the USA [4], this isn't surprising, but an impressive demonstration of MGT.

On the other hand, MGT4-CC1 including MGT4-ST171, ST15 and ST163, were less prevalent in the USA. Nevertheless these STs were isolated from human infections but very rare from poultry source. More studies are required to explain the association between MGT4-CC13 but not MGT4-CC1 and poultry products. Beef, sprouts, pork, nuts and seeds can also be contaminated by *S.* Enteritidis [4]. And these non-poultry related foods may be the source of the MGT4-CC1 caused infections. The lack of potential source isolates within MGT4-CC1 STs highlights the need for comprehensive sequencing and epidemiological efforts across the food chain.

Temporal variation of UK *S.* Enteritidis was depicted clearly by MGT, showcasing the power of MGT for longer term epidemiology. From 2014, all clinical *S.* Enteritidis were routinely sequenced in the UK [20]. Some MGT5-STs were found to occur for a few months and then disappeared. Such a pattern of birth and death of a ST may be related to outbreaks. For example, MGT5-ST156, which increased substantially during June and July in 2014 and became rare after 2015, was associated with a reported large scale outbreak [19]. Other STs persisted for long periods of time. For example, MGT5-ST1 and ST29, appeared to be endemic isolates associated with local reservoirs. Therefore, the variation of MGT-STs across different seasons offered additional epidemiological signals for suspected

outbreaks and endemic infections of *S*. Enteritidis. The stable feature of STs avoids the cluster merging

issues of single linkage based clustering methods and ensures continuity for long-term surveillance [14].

MGT for *S*. Enteritidis facilitates outbreak detection, source tracing and evaluation of outbreak

propensity

Whole genome sequencing offered the highest resolution for outbreak detection and source tracing.

However, a standardised approach for prospective genomic surveillance is still lacking. cgMLST

offered potential for such standardisation and harmonisation of results between different providers.

However, resolution of cgMLST heavily depends on the diversity of the species. Our previous study

showed that for *S*. Typhimurium, serovar core genome offered higher resolution than species level core

genome for outbreak investigation [14]. We designed MGT9 for *S*. Enteritidis using 4986 loci, which

is around 2000 more loci than *Salmonella* core genes, thus increasing the resolution of subtyping [24].

MGT9 is designed to give the highest resolution for spread tracking and outbreak detection. In this

study, *S*. Enteritidis isolates belonging to the same MGT9-STs were identified in two or more different

countries, offering strong evidence of international transmission. To facilitate outbreak detection MGT9

STs were further grouped to ODCs. We used ODC2, which has a two allele difference cut-off, to

identify potential outbreak clusters. Although the data does not allow us to confirm whether any of these

ODC2 clusters was actual outbreaks, a number of known outbreaks from other studies fell into ODC2

clusters [22, 25]. However, outbreak detection may need to use dynamic threshold for cut-offs [25],

which requires further work for *S*. Enteritidis. By ODC2, 40% of the 26,670 isolates belonged to ODC2

clusters of more than 10 isolates, indicating the commonly occurrence of outbreaks by *S*. Enteritidis.

Importantly, we found the European and North American prevalent lineage MGT4-CC13 were

significantly more correlated with larger ODC2 clusters (>= 50 isolates each) than the global epidemic

lineage of MGT4-CC1 (Chi-Square test, P < 0.05). Thus, MGT4-CC13 is more likely to cause large

scale outbreaks than MGT4-CC1. In the past few years, the reported large scale outbreaks in Europe,

the majority of which were due to the contaminated eggs [18, 19, 22, 26], belonged to MGT4-CC13.

In the USA, MGT4-CC13 was the dominant lineage in both human infections and poultry product

contaminations, while MGT4-CC1 was relatively rare in poultry. The industrialised poultry/eggs

production and marketing would have facilitated the spread of MGT4-CC13 causing large scale outbreaks. Further studies are required to understand the underlying mechanisms how MGT4-CC13 has developed the capacity to cause large outbreaks.

ODCs offer a means for accurate detection of outbreaks at the highest genome resolution MGT9 [14]. Other studies used SNP address or cgMLST [18, 19, 26] with SNP cut-offs or HierCC clustering for outbreak detection. All three methods use the same single linkage clustering method for outbreak detection. MGT offers further advantage that potential outbreak clusters can be precisely defined by a stable MGT-ST identifier [14].

MGT for *S.* Enteritidis improves precision of detection and global tracking of MDR clones

The rise of AMR in *Salmonella* is a serious public health concern [5]. The global spread of AMR can be mediated by lateral transfer of resistance genes as well as clonal spread of resistant strains. This study systematically evaluated the presence of antibiotic resistant genes in the 26,670 genomes of *S.* Enteritidis, 9.4% of which harboured antibiotic resistant genes. We assessed resistance rates across STs from different MGT levels to identify drug resistant STs. Among the two global epidemic lineages, MGT4-CC1 was found to contain significantly more antibiotic resistant isolates than MGT4-CC13. We further identified STs that were associated with MDR. Eleven STs (>= 10 isolates each, 659 isolates in total) of different MGT levels were MDR. MGT can be used to track the spread of these isolates more precisely. These STs from different MGT levels were mutually exclusive, emphasising the flexibility of MGT for precise identification of MDR clones. MGT3-ST15 and MGT3-ST10 were representative of the previously reported Africa invasive infection related lineages [5], harboured resistant genes of up to 6 different antibiotic classes. The other STs, which were mainly from Europe and North America, harbor AR genes to up to 8 different types of drugs. MGT3-ST30 and MGT4-ST718 (5 isolates in total), which belonged to MGT4-CC1 and were observed in Europe, had been reported to be MDR phenotypically [27]. Significantly, isolates from these two STs were mainly isolated from blood samples [27]. These MDR STs correlated with blood stream infections should be monitored closely as they can potentially be more invasive [5].

Moreover, we identified multiple plasmids that were MDR associated. Plasmids are known to play a key role in the acquisition of drug resistance genes [28, 29]. MDR of African isolates in MGT3-ST10 and ST15 has been reported to be mediated by plasmids [5]. The West Africa lineage MGT3-ST10 isolates were reported to have IncI1 plasmids [30]. In this study, both IncI1 and IncQ1 plasmids were present in the Africa MGT3-ST10 and ST15 isolates as well as STs belonged to global epidemic lineage MGT4-CC1. IncQ1 plasmids, which are highly mobile and widely transferred among different genus of bacteria [31], are correlated with MDR of *Salmonella* [32, 33]. IncI1 plasmids were found responsible for the cephalosporin resistance of *Salmonella* and *Escherichia* in different countries [34-36]. Moreover, IncN and IncX1 plasmids were common in the MDR associated STs of MGT4-CC1 and CC13. The use of MGT would enhance the surveillance of drug resistance plasmids.

Except for the Africa invasive infection lineage STs, the majority of the MDR STs isolates belonged to MGT4-CC1 and were mainly from Europe and North America. It is likely due to less WGS data being publicly available from other continents. Africa and Asia *S.* Enteritidis isolates were found to have higher proportion of MDR, which are concordant with the frequent reports of MDR in *S.* Enteritidis isolates from these two continents [5, 29, 37]. With the expansion of the MGT database and genomic surveillance, more MDR related STs may be identified, which will help track and control the spread of MDR globally.

Defining the population structure of *S.* Enteritidis with increasing resolution and precision at different MGT levels

In this study, we defined the global population structure of *S.* Enteritidis using MGT-CCs. Three clades of *S.* Enteritidis were previously defined with clade A and C as outgroups to the predominant clade B [9], a classification supported by this study. We further identified 10 lineages within clade B, which can be represented by different CCs from MGT2 to MGT4 corresponding with time of divergence. Lower resolution level MGT CCs described old lineages well while higher resolution level MGT CCs described the recent derived lineages (**Supplementary Fig. S8**). Thus, the epidemiological characteristics of a lineage can be identified using STs and/or CCs at appropriate MGT level. MGT2-CC3 represent the Europe endemic MGT1-ST183 lineage, within which MGT2-ST3 and MGT2-ST82

were able to identify previously defined phage types PT11 and PT66 respectively [17]. MGT3-CC10 included all of the reported West Africa lineage isolates, and MGT3-CC15 included all of the reported Central/Eastern Africa lineage isolates as well as Kenyan invasive *S*. Enteritidis isolates [5, 8]. MGT3-ST10 and MGT3-ST15 (represented 96.3% of MGT3-CC10 and 90.2% of MGT3-CC15 respectively), were MDR with similar resistance patterns. Most importantly, the two major global epidemic lineages are clearly described by MGT4-CC1 and MGT4-CC13. MGT4-CC1 corresponds the global epidemic clade in the study of Feasey *et al*., and MGT4-CC13 to the outlier clade of that study [5]. MGT4-CC1 and CC13 correspond to lineage III and lineage V in Deng *et al*'s study [11]. Thus, by using STs and/or CCs of different MGT levels the population structure of *S*. Enteritidis can be described elegantly.

Interestingly, the two dominant lineages, MGT4-CC1 and CC13, showed different population expansion time and evolutionary dynamics. BEAST analyses revealed a core genome mutation rate for *S*. Enteritidis of 1.9×10^{-7} substitution/site/year which is similar to the genome mutation rate of 2.2×10^{-7} substitution/site/year estimated by Deng *et al* [11]. In contrast, the mutation rate of MGT4-CC13 was 1.5 times faster than MGT4-CC1. Population expansion of MGT4-CC1 occurred with two peaks at around 1950s and 1970s, while MGT4-CC13 population was increasing steadily until 1970s. The expansion around 1970s may be driven by the development of poultry industry that *S*. Enteritidis colonized in the poultry processing plants. It is concordant with Deng *et al*'s speculation that the expansion of *S*. Enteritidis population was correlated with the poultry farm industry [11, 38].

***S*. Enteritidis MGT scheme enables scalable nomenclature at different typing resolution**

The design of the *S*. Enteritidis MGT is based on the *S*. Typhimurium MGT scheme published previously [14]. The first eight levels (MGT1 to MGT8) of *S*. Enteritidis MGT scheme used the same loci as for *S*. Typhimurium. We defined the core genome of *S*. Enteritidis which was used as MGT9 with 4986 loci that include 3932 core genes and 1054 core intergenic regions. The subtyping resolution of MGT for *S*. Enteritidis substantially increased compared to the *Salmonella* cgMLST, which included 3002 core genes of *S. enterica* [24]. Our study suggests that the MGT levels 1 to 8 could be applied to all *Salmonella* serovars as a common scheme for the species, and only an additional serovar-specific

MGT9 scheme needs to be designed for individual serovars that require the highest resolution for outbreak investigations.

MGT is partly based on cgMLST, which has gained popularity as a genotyping tool [24] and both share the same basic principle of exact comparison of alleles. However, there are features that make MGT superior to cgMLST. For cgMLST, to provide varied clustering resolution of isolates, single linkage clustering methods are applied to cgMLST at various allelic difference cutoffs (eg, 5, 10, 200); such a system has been implemented in EnteroBase and is referred to as HierCC [12, 13]. A major disadvantage of HierCC is the merging of clusters when an intermediate isolate can be assigned to multiple groups. This scenario will almost certainly occur in a large database with ever expanding numbers of sequences [14]. This can be resolved by MGT-STs, which are based on exact comparison and offer stable nomenclature at different subtyping resolutions using different MGT levels. Additionally, CCs at each MGT level were able to cluster the STs with one allele difference and are concordant with phylogenetic lineages as shown in this study (**Fig. 6**). The major advantage of MGT is that STs/CCs at different levels of MGT offer flexibility for temporal and spatial epidemiological analysis using an underlying stable nomenclature of STs. The online database of *S.* Enteritidis MGT scheme offers an open platform for global public health communication, which facilitates the international transmission control and outbreak detection of *S.* Enteritidis.

Conclusions

In this study, we defined the core genome of *S.* Enteritidis and developed an MGT scheme with 9 levels. MGT9 offers the highest resolution and is suitable for outbreak detection whereas the lower levels (MGT1 to MGT8) are suitable for longer term epidemiology. At the MGT4 level, globally prevalent and regionally restricted STs were identified, which would facilitate the identification of the potential geographic source of *S.* Enteritidis isolates. Specific source associated STs were identified, such as poultry associated MGT4-STs, which were common in human cases in the USA. At the MGT5 level, temporal variation of STs was observed in *S.* Enteritidis infections from the UK, which reveal both long lived endemic STs and the rapid expansion of new STs. Using MGT3 to MGT6, we identified MDR associated STs to facilitate tracking of MDR spread. Additionally, certain plasmid types were found to be highly associated with the same MDR STs, suggesting plasmid-based resistance acquisition.

Furthermore, we evaluated the phylogenetic relationship of the STs by defining the population structure of *S*. Enteritidis. A total of 10 main lineages were defined in the globally distributed clade B *S*. Enteritidis. Of these, MGT4-CC1 and CC13 were the two dominant lineages, with significant differences in large outbreak frequency, antimicrobial resistance profiles and mutation rates. The online open database for *S*. Enteritidis MGT offers a unified platform for the international surveillance of *S*. Enteritidis. In summary, MGT for *S*. Enteritidis provides a flexible, high resolution and stable genome typing tool for long term and short long surveillance of *S*. Enteritidis infections.

Methods

***S*. Enteritidis sequences used in this study**

Raw reads of 26,670 *S*. Enteritidis whole genome sequences (WGS) downloaded from the European Nucleotide Archive (ENA) passed the quality filtering criteria by Quast [39, 40] (**Supplementary Table S7)**. The epidemiological metadata of those isolates were collated from NCBI and EnteroBase [24]. Trimmomatic v0.39 was used with default settings to trim raw reads [41]. The trimmed raw reads were then assembled using either SPAdes v3.13.0 for core genome defination or SKESA v2.3 for MGT typing [42, 43]. Quality assessment for the assemblies were performed using Quast v5.0.2 [39]. Thresholds of the assembly quality were in accordance with Robertson *et al* 's criteria [40]. Kraken v0.10.5 was used to double check any possible contamination in the assembled genomic sequences or problematic taxonomic identification [44]. Seqsero v1.0 and SISTR v1.0.2 were used to verify the serotype [45, 46].

Core gene definition of *S*. Enteritidis

The *S*. Enteritidis isolates from Enterobase (during May, 2018) were sub-sampled to identify representative isolates in three different ways depending on the size of the ribosomal sequence type (rST) [24]. For the isolates of the rSTs with less than 10 isolates each, all of the isolates were sampled. For the isolates of the rST with more than 10 and less than 350 isolates assigned, 10 isolates for each type were randomly sampled. For the isolates of the rST with more than 350 isolates (rST1425 and rST3888), 2% of the isolates for each type were sampled randomly while including isolates from as

many countries and collection years as possible. A total of 1801 genomes, which were selected to

represent 283 rSTs collected over 101 years (1917-2018) in 56 different countries.

The trimmed raw reads were then assembled using SPAdes v3.13.0 [42]. Prokka v1.12 was used to

predict and annotate genes [47]. Roary v3.11.2 was used for the determination of core genes, which

were defined as the genes present in 99% of the representative genomes. Core genes were identified as

present in each genome following the nucleotide identity threshold >= 96% and alignment coverage

threshold >= 99%. Those thresholds were assessed as the optimal identity by comparing the number of

core genes generated with different identity from 90% to 100% (**Supplementary Fig. S1**) [48]. Core

genes with duplicates in the Roary output (paralogues) were removed. Core intergenic regions were

defined using Piggy v3.11.2 based on the Roary outputs [49], and the nucleotide identity and coverage

thresholds were identical to those of core genes.

Establishment of MGT scheme for *S.* Enteritidis and MGT allele calling

The initial allele sequence of each locus used for MGT were extracted from the complete genome

sequence of *S.* Enteritidis strain P125109 (Genbank accession NC_011294.1). The first 8 levels of MGT

were identical to the MGT scheme for *S.* Typhimurium [14], in which MGT1 refers to the classical 7

gene MLST for the *Salmonella* genus [16]. Loci used for levels MGT2 to MGT8 were orthologous to

those in the *S.* Typhimurium MGT with the exception of one gene which was removed at MGT8 because

of a paralogue in *S.* Enteritidis. MGT9 contains all loci in MGT8 as well as the core genes and core

intergenic regions of *S.* Enteritidis.

The raw read processing and genome assembly procedures were the same as above, except that

SKESA v2.3 was used to assemble the raw reads and SISTR v1.0.2 was used for the molecular serotype

identification [43, 46]. SKESA v2.3 offers better per base accuracy and speed than SPADES and was

therefore used in the allele calling pipeline [43]. For the assemblies that passed quality control by the

criteria of Robertson *et al.* [40], a custom script was used to assign alleles, STs and CCs to those isolates

at levels from MGT2 to MGT9 [14]. Allele, ST, CC and outbreak detection cluster (ODC) assignments

were performed as previously described [14]. Briefly, if one or more allele difference was observed, a

new ST was assigned to the isolate. The same strategy was also used for assigning a new CC, if at least

two allele differences were observed comparing to any of STs in existing CCs in the database [14]. Outbreak detection cluster (ODC) is an extension of the CC clustering method, but it was performed only at MGT9 level [14]. For example, any two MGT9 STs with no more than 2, 5, or 10 allele differences were assigned the same ODC2, ODC5 or ODC10 type [14].

Geographic distribution and temporal variation analysis

We first determined which MGT level would be the most useful in describing the global epidemiology of *S.* Enteritidis by summarizing STs that contained at least 10 isolates for their distribution across continents. STs where more than 70% of isolates were from a single continent were identified at each level of MGT (**Supplementary Fig. S3a**). MGT4 STs matching this criterion had the highest percentage of all the levels at 76.3% of the 26,670 isolates, thus MGT4 was chosen to describe the continental distribution of *S.* Enteritidis. The USA states variation map and the UK monthly variation column chart were produced using Tableau v2019.2 [50].

Antibiotic resistance gene and plasmid specific gene identification

Abricate (https://github.com/tseemann/abricate) was used for identification of antibiotic resistance genes with ResFinder v3.0 database [51], and plasmid specific genes with PlasmidFinder v2.1 database [21]. The cut-off of the presence of a gene was set as both identity and coverage >= 90% [51]. At each MGT level, STs (>=10 isolates each) with more than 80% of isolates harbouring antibiotic resistance genes, were collected from MGT2 to MGT9 and were defined as high-antibiotic-resistant STs. The STs at each level of MDR were non-redundant meaning that no isolate was counted more than once.

Phylogenetic analysis

To define the major clades of *S.* Enteritidis globally, 1508 isolates were sampled based on the STs of MGT6 (random sampling of one isolate per ST with 3 or more isolates). A phylogenetic tree was constructed using ParSNP v1.2, which called core-genome SNPs [52, 53]. The branches with more than 10 isolates on the phylogenetic tree were collapsed together with iTOL v4 [54].

Average mutation rates were estimated for the global epidemic clade B of *S.* Enteritidis and the two dominant lineages, MGT4-CC1 and CC13. For the global epidemic clade B, a total of 90 isolates were randomly sampled based on MGT4-CCs and STs (covering the main STs for each CC). For the mutation rate of MGT4-CC1 or CC13, representative isolates were randomly sampled based on the higher resolution level MGT5 and collections years. For each MGT5-CC type, one isolate from each collection year were randomly sampled. SNPs derived from MGT9 alleles were then concatenated as a multi-alignment file and potential recombinant SNPs were removed using both Gubbins v2.0.0 and Recdetect v6.0 [55, 56]. Beast v1.10.4 was then used to estimate the mutation rate based on mutational SNPs [57]. A total of 24 combinations of clock and population size models were evaluated with the MCMC chain of 100 million states. Tracer v1.7.1 was used to identify the optimal model and to estimate population expansion over time [58]. For the predominant clade of *S.* Enteritidis including isolates of multiple lineages, uncorrelated relaxed lognormal clock combined with the Bayesian skyline model was chosen by the effective sample size (ESS) of both mean rate and joint probability of over 300. For the two dominant lineages of *S.* Enteritidis, MGT4-CC1 or CC13, the strict clock and Bayesian skyline models were found to be the optimal combination model by the ESS of mean rate and joint probability of over 100. The large size of each CC (more than 300 isolates for the Beast analysis) made it hard to reach 200 of ESS even for the optimal models.

Abbreviations

WGS: Whole genome sequence; MGT: Multilevel genome typing; ST: Sequence type; CC: Clonal complex; ODC: Outbreak detection cluster; rST: ribosomal sequence type; SNP: Single nucleotide polymorphism; MLST: Multi-locus sequence typing; cgMLST: core genome multi-locus sequence typing; HierCC: Hierarchical clustering of cgMLST. AR: Antibiotic resistance; MDR: Multi-drug resistance.

Acknowledgements

The authors thank Duncan Smith and Robin Heron for technical assistance.

Funding

This work was funded by a National Health and Medical Research Council project grant (grant number 1129713).

Authors' contributions

L.L., M.P., and R.L. designed the study. L.L. and M.P. set up the database. S.K and M.P. set up the website, M.P., R.L., D.H. L.C. S.O., Q.W., M.T., V.S. and S.K. provided critical analysis and discussions, L.L. wrote the first draft and all authors contributed to the final manuscript.

Availability of data and materials

The MGT database for *S.* Enteritidis is available in https://mgtdb.unsw.edu.au/enteritidis/.

Competing interests

The authors declare that they have no competing interests.

References

1. Dewey-Mattia D, Manikonda K, Hall AJ, Wise ME, Crowe SJ: Surveillance for Foodborne Disease Outbreaks - United States, 2009-2015. MMWR Surveill Summ. 2018;67(10):1-11.

2. ECDC: Salmonellosis - Annual Epidemiological Report for 2017. In: European Centre for Disease Prevention and Control. 2020.

3. Pijnacker R, Dallman TJ, Tijsma ASL, Hawkins G, Larkin L, Kotila SM, Amore G, Amato E, Suzuki PM, Denayer S *et al*: An international outbreak of *Salmonella enterica* serotype Enteritidis linked to eggs from Poland: a microbiological and epidemiological study. Lancet Infect Dis. 2019;19(7):778-786.

4. Snyder TR, Boktor SW, M'Ikanatha N M: Salmonellosis Outbreaks by Food Vehicle, Serotype, Season, and Geographical Location, United States, 1998 to 2015. J Food Prot. 2019;82(7):1191-1199.

5. Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, Deng X, Wigley P, Barquist L, Langridge GC, Feltwell T *et al*: Distinct *Salmonella* Enteritidis lineages associated with

enterocolitis in high-income settings and invasive disease in low-income settings. Nat Genet. 2016;48(10):1211-1217.

6. Mohan A, Munusamy C, Tan YC, Muthuvelu S, Hashim R, Chien SL, Wong MK, Khairuddin NA, Podin Y, Lau PS *et al*: Invasive *Salmonella* infections among children in Bintulu, Sarawak, Malaysian Borneo: a 6-year retrospective review. BMC Infect Dis. 2019;19(1):330.

7. Thung TY, Mahyudin NA, Basri DF, Wan Mohamed Radzi CW, Nakaguchi Y, Nishibuchi M, Radu S: Prevalence and antibiotic resistance of *Salmonella* Enteritidis and *Salmonella* Typhimurium in raw chicken meat at retail markets in Malaysia. Poult Sci. 2016;95(8):1888-1893.

8. Akullian A, Montgomery JM, John-Stewart G, Miller SI, Hayden HS, Radey MC, Hager KR, Verani JR, Ochieng JB, Juma J *et al*: Multi-drug resistant non-typhoidal *Salmonella* associated with invasive disease in western Kenya. PLoS Negl Trop Dis. 2018;12(1):e0006156.

9. Graham RMA, Hiley L, Rathnayake IU, Jennison AV: Comparative genomics identifies distinct lineages of *S*. Enteritidis from Queensland, Australia. PLoS One. 2018;13(1):e0191042.

10. Parn T, Dahl V, Lienemann T, Perevoscikovs J, De Jong B: Multi-country outbreak of *Salmonella* enteritidis infection linked to the international ice hockey tournament. Epidemiol Infect. 2017;145(11):2221-2230.

11. Deng X, Desai PT, den Bakker HC, Mikoleit M, Tolar B, Trees E, Hendriksen RS, Frye JG, Porwollik S, Weimer BC *et al*: Genomic epidemiology of *Salmonella enterica* serotype Enteritidis based on population structure of prevalent lineages. Emerg Infect Dis. 2014;20(9):1481-1489.

12. Zhou Z, Alikhan N-F, Mohamed K, Fan Y, Agama Study G, Achtman M: The EnteroBase user's guide, with case studies on *Salmonella* transmissions, *Yersinia* pestis phylogeny, and *Escherichia* core genomic diversity. Genome Res. 2020;30(1):138-152.

13. Ashton P, Nair S, Peters T, Tewolde R, Day M, Doumith M, Green J, Jenkins C, Underwood A, Arnold C *et al*: Revolutionising Public Health Reference Microbiology using Whole Genome Sequencing: *Salmonella* as an exemplar. bioRxiv. 2015:033225.

14. Payne M, Kaur S, Wang Q, Hennessy D, Luo L, Octavia S, Tanaka MM, Sintchenko V, Lan R: Multilevel genome typing: genomics-guided scalable resolution typing of microbial pathogens. Euro Surveill. 2020;25(20):1900519.

15. Feil EJ: Small change: keeping pace with microevolution. Nat Rev Microbiol. 2004;2(6):483-495.

16. Achtman M, Wain J, Weill FX, Nair S, Zhou Z, Sangal V, Krauland MG, Hale JL, Harbottle H, Uesbeck A *et al*: Multilocus sequence typing as a replacement for serotyping in *Salmonella enterica*. PLoS Pathog. 2012;8(6):e1002776.

17. Lawson B, Franklinos LHV, Rodriguez-Ramos Fernandez J, Wend-Hansen C, Nair S, Macgregor SK, John SK, Pizzi R, Nunez A, Ashton PM *et al*: *Salmonella* Enteritidis ST183: emerging and endemic biotypes affecting western European hedgehogs (Erinaceus europaeus) and people in Great Britain. Sci Rep. 2018;8(1):2449.

18. Kanagarajah S, Waldram A, Dolan G, Jenkins C, Ashton PM, Carrion Martin AI, Davies R, Frost A, Dallman TJ, De Pinna EM *et al*: Whole genome sequencing reveals an outbreak of *Salmonella* Enteritidis associated with reptile feeder mice in the United Kingdom, 2012-2015. Food Microbiol. 2018;71:32-38.

19. Hormansdorfer S, Messelhausser U, Rampp A, Schonberger K, Dallman T, Allerberger F, Kornschober C, Sing A, Wallner P, Zapf A: Re-evaluation of a 2014 multi-country European outbreak of *Salmonella* Enteritidis phage type 14b using recent epidemiological and molecular data. Euro Surveill. 2017;22(50):17-00196.

20. Chattaway MA, Dallman TJ, Larkin L, Nair S, McCormick J, Mikhail A, Hartman H, Godbole G, Powell D, Day M *et al*: The Transformation of Reference Microbiology Methods and Surveillance for *Salmonella* With the Use of Whole Genome Sequencing in England and Wales. Frontiers in public health. 2019;7(317):317.

21. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H: In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895-3903.

22. Dallman T, Inns T, Jombart T, Ashton P, Loman N, Chatt C, Messelhaeusser U, Rabsch W, Simon S, Nikisins S *et al*: Phylogenetic structure of European *Salmonella* Enteritidis outbreak

correlates with national and international egg distribution network. Microb Genom. 2016;2(8):e000070.

23. Nichols M, Stevenson L, Whitlock L, Pabilonia K, Robyn M, Basler C, Gomez T: Preventing Human *Salmonella* Infections Resulting from Live Poultry Contact through Interventions at Retail Stores. J Agric Saf Health. 2018;24(3):155-166.

24. Alikhan NF, Zhou Z, Sergeant MJ, Achtman M: A genomic overview of the population structure of *Salmonella*. PLoS Genet. 2018;14(4):e1007261.

25. Payne M, Octavia S, Luu LDW, Sotomayor-Castillo C, Wang Q, Tay ACY, Sintchenko V, Tanaka MM, Lan R: Enhancing genomics-based outbreak detection of endemic *Salmonella enterica* serovar Typhimurium using dynamic thresholds. Microb Genom. 2019.

26. Inns T, Lane C, Peters T, Dallman T, Chatt C, McFarland N, Crook P, Bishop T, Edge J, Hawker J *et al*: A multi-country *Salmonella* Enteritidis phage type 14b outbreak associated with eggs from a German producer: 'near real-time' application of whole genome sequencing and food chain investigations, United Kingdom, May to September 2014. Euro Surveill. 2015;20(16):21098.

27. Vidovic S, An R, Rendahl A: Molecular and Physiological Characterization of Fluoroquinolone-Highly Resistant *Salmonella* Enteritidis Strains. Front Microbiol. 2019;10:729.

28. Kaldhone PR, Carlton A, Aljahdali N, Khajanchi BK, Sanad YM, Han J, Deck J, Ricke SC, Foley SL: Evaluation of Incompatibility Group I1 (IncI1) Plasmid-Containing *Salmonella enterica* and Assessment of the Plasmids in Bacteriocin Production and Biofilm Development. Front Vet Sci. 2019;6(298):298.

29. Zhou X, Li M, Xu L, Shi C, Shi X: Characterization of Antibiotic Resistance Genes, Plasmids, Biofilm Formation, and In Vitro Invasion Capacity of *Salmonella* Enteritidis Isolates from Children with Gastroenteritis. Microb Drug Resist. 2019;25(8):1191-1198.

30. Aldrich C, Hartman H, Feasey N, Chattaway MA, Dekker D, Al-Emran HM, Larkin L, McCormick J, Sarpong N, Le Hello S *et al*: Emergence of phylogenetically diverse and fluoroquinolone resistant *Salmonella* Enteritidis as a cause of invasive nontyphoidal *Salmonella* disease in Ghana. PLoS Negl Trop Dis. 2019;13(6):e0007485.

31. Smalla K, Heuer H, Gotz A, Niemeyer D, Krogerrecklenfort E, Tietze E: Exogenous isolation of antibiotic resistance plasmids from piggery manure slurries reveals a high prevalence and diversity of IncQ-like plasmids. Appl Environ Microbiol. 2000;66(11):4854-4862.

32. Castellanos LR, van der Graaf-van Bloois L, Donado-Godoy P, Leon M, Clavijo V, Arevalo A, Bernal JF, Mevius DJ, Wagenaar JA, Zomer A *et al*: Genomic Characterization of Extended-Spectrum Cephalosporin-Resistant *Salmonella enterica* in the Colombian Poultry Chain. Front Microbiol. 2018;9:2431.

33. Mastrorilli E, Pietrucci D, Barco L, Ammendola S, Petrin S, Longo A, Mantovani C, Battistoni A, Ricci A, Desideri A *et al*: A Comparative Genomic Analysis Provides Novel Insights Into the Ecological Success of the Monophasic *Salmonella* Serovar 4,[5],12:i. Front Microbiol. 2018;9:715.

34. Wong MH, Kan B, Chan EW, Yan M, Chen S: IncI1 Plasmids Carrying Various blaCTX-M Genes Contribute to Ceftriaxone Resistance in *Salmonella enterica* Serovar Enteritidis in China. Antimicrob Agents Chemother. 2016;60(2):982-989.

35. Garcia-Fernandez A, Chiaretto G, Bertini A, Villa L, Fortini D, Ricci A, Carattoli A: Multilocus sequence typing of IncI1 plasmids carrying extended-spectrum beta-lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. J Antimicrob Chemother. 2008;61(6):1229-1233.

36. Kameyama M, Chuma T, Yokoi T, Yabata J, Tominaga K, Miyasako D, Iwata H, Okamoto K: Emergence of *Salmonella enterica* serovar infantis harboring IncI1 plasmid with bla(CTX-M-14) in a broiler farm in Japan. J Vet Med Sci. 2012;74(9):1213-1216.

37. Ma Y, Li M, Xu X, Fu Y, Xiong Z, Zhang L, Qu X, Zhang H, Wei Y, Zhan Z *et al*: High-levels of resistance to quinolone and cephalosporin antibiotics in MDR-ACSSuT *Salmonella enterica* serovar Enteritidis mainly isolated from patients and foods in Shanghai, China. Int J Food Microbiol. 2018;286:190-196.

38. Deng X, Shariat N, Driebe EM, Roe CC, Tolar B, Trees E, Keim P, Zhang W, Dudley EG, Fields PI *et al*: Comparative analysis of subtyping methods against a whole-genome-sequencing standard for *Salmonella enterica* serotype Enteritidis. J Clin Microbiol. 2015;53(1):212-218.

39. Gurevich A, Saveliev V, Vyahhi N, Tesler G: QUAST: quality assessment tool for genome assemblies. Bioinformatics. 2013;29(8):1072-1075.

40. Robertson J, Yoshida C, Kruczkiewicz P, Nadon C, Nichani A, Taboada EN, Nash JHE: Comprehensive assessment of the quality of *Salmonella* whole genome sequence data available in public sequence databases using the *Salmonella* in silico Typing Resource (SISTR). Microb Genom. 2018;4(2).

41. Bolger AM, Lohse M, Usadel B: Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014;30(15):2114-2120.

42. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD *et al*: SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19(5):455-477.

43. Souvorov A, Agarwala R, Lipman DJ: SKESA: strategic k-mer extension for scrupulous assemblies. Genome Biol. 2018;19(1):153.

44. Wood DE, Salzberg SL: Kraken: ultrafast metagenomic sequence classification using exact alignments. Genome Biol. 2014;15(3):R46.

45. Zhang S, Yin Y, Jones MB, Zhang Z, Deatherage Kaiser BL, Dinsmore BA, Fitzgerald C, Fields PI, Deng X: *Salmonella* serotype determination utilizing high-throughput genome sequencing data. J Clin Microbiol. 2015;53(5):1685-1692.

46. Yoshida CE, Kruczkiewicz P, Laing CR, Lingohr EJ, Gannon VP, Nash JH, Taboada EN: The *Salmonella* In Silico Typing Resource (SISTR): An Open Web-Accessible Tool for Rapidly Typing and Subtyping Draft *Salmonella* Genome Assemblies. PLoS One. 2016;11(1):e0147101.

47. Seemann T: Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30(14):2068-2069.

48. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J: Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics. 2015;31(22):3691-3693.

49. Thorpe HA, Bayliss SC, Sheppard SK, Feil EJ: Piggy: a rapid, large-scale pan-genome analysis tool for intergenic regions in bacteria. GigaScience. 2018;7(4):1-11.

1   50.   Tableau (version. 9.1). J Med Libr Assoc. 2016;104(2):182-183.

2   51.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM,

3   Larsen MV: Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother.

4   2012;67(11):2640-2644.

5   52.   Treangen TJ, Ondov BD, Koren S, Phillippy AM: The Harvest suite for rapid core-genome

6   alignment and visualization of thousands of intraspecific microbial genomes. Genome Biol.

7   2014;15(11):524.

8   53.   Price MN, Dehal PS, Arkin AP: FastTree 2--approximately maximum-likelihood trees for large

9   alignments. PloS one. 2010;5(3):e9490-e9490.

10   54.   Letunic I, Bork P: Interactive Tree Of Life (iTOL) v4: recent updates and new developments.

11   Nucleic Acids Res. 2019;47(W1):W256-W259.

12   55.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR:

13   Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome

14   sequences using Gubbins. Nucleic Acids Res. 2015;43(3):e15.

15   56.   Hu D, Liu B, Wang L, Reeves PR: Living Trees: High-Quality Reproducible and Reusable

16   Construction of Bacterial Phylogenetic Trees. Mol Biol Evol. 2020;37(2):563-575.

17   57.   Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A: Bayesian

18   phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol.

19   2018;4(1):vey016.

20   58.   Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA: Posterior Summarization in

21   Bayesian Phylogenetics Using Tracer 1.7. Syst Biol. 2018;67(5):901-904.

Figure 1. Makeup and assignments of each *S*. enteritidis MGT level. a. Number of loci included in each MGT level. The first eight levels are composed of *Salmonella* species core genes, which are orthologous to those of the *S*. Typhimurium MGT scheme, except for one gene at MGT8 which was excluded due to duplication in *S*. enteritidis. The MGT9 scheme includes core genes of S. Enteritidis (those not belonging to the *Salmonella* core coloured in yellow) and core intergenic regions (coloured in green). The number behind or within each bar refers to the number of loci included. **b.** The number of sequence types (ST) and clonal complexes (CC) assigned at each MGT level among the 26,670 genomes analysed. CC includes STs of no more than 1 allele difference. The numbers refer to the number of STs or CCs types assigned at each level, which were also indicated by colour. As MGT9 included 4986 loci with highest subtyping resolution, the 26,670 isolates were subtyped into 20,153 different ST types and 14,441 CC clustering types at MGT9.

Figure 2. Continental restricted STs of *S*. Enteritidis were identified within each CCs at MGT4. The numbers in each pie refer to the number of whole genome sequences (WGS) analysed in each continent. The inner ring shows the main MGT4-CCs (with proportions of 3% or more) in each continent. MGT4-CC1 and CC13 were the two predominant MGT4-CCs with MGT4-CC1 being more widely distributed in all six continents. The outer ring shows the common MGT4-STs in each continent (with proportions of 2% or higher) within corresponding CCs. MGT4-ST15 was the dominant ST in MGT4-CC1, which were widely distributed all over the world. Continental restricted STs were identified, especially for the STs of MGT4-CC13, which was more prevalent in Europe and North America.

Figure 3. USA and UK *S.* Enteritidis isolates can be distinguished by STs at MGT4 and MGT5. a. At MGT4, there were 39 STs (50 or more isolates each) representing 75.1% of the UK and USA isolates (22191 in total). MGT4-ST99, ST136 and ST135 were prevalent in the USA, MGT4-ST15, ST25 and ST13 were prevalent in the UK, while MGT4-ST15 and ST25 were mixed. **b.** MGT4-ST15 and ST25 can be subtyped into multiple STs at MGT5, the majority of which were prevalent in the UK, except for MGT5-ST412 and ST387 in the USA.

Figure 4. MGT4 ST proportions across states in the USA is similar but only a subset of STs are associated with avian hosts. For the USA *S*. Enteritidis isolates with metadata (4,383 isolates), a total of 3935 genomes were from either avian or human source, 95.0% (3738/3935) of which belonged to MGT4-CC1 and CC13. The MGT4-STs belonged MGT4-CC1 and CC13 of either avian or human source were shown in different states of the USA. Each pie chart illustrated the MGT-ST types and the size of the main STs represented with different colours. The size of the pie indicated the total number of isolates in each state. While CC13 appears to originate in birds before moving to humans, the reservoir of CC1 is not in birds. Origins for CC1 STs were also not observed in any other source type.

Figure 5. MGT5 STs allow identification of temporal patterns in strain diversity in the UK.
There were 13 MGT5-STs of more than 100 isolates shown with different colours. MGT5-ST156 in red was observed predominantly from June to August during 2014, and disappeared after 2014. MGT5-ST423 was a dominant type from May to September 2015, and became dominant again during the September to December of 2016.

Figure 6. Global phylogenetic structure of _S._ Enteritidis. A total of 1508 genomes were randomly sampled from each ST in MGT6. A phylogenetic tree was constructed by ParSNP v1.2, which called core-genome SNPs extracted from alleles. The tree scales indicated the 0.01 substitutions per locus. The number on internal branches represented the percentage of bootstrap support. **a.** Three clades were identified which were concordant with previous findings: The predominant clade, highlighted in yellow, is the global epidemic clade which was phylogenetically closer to _S._ Gallinarum than the other two clades in purple and green background which are limited in Oceania [9]. **b.** The main, yellow highlighted clade was expanded into a more detailed phylogeny with _S._ Gallinarium as an outgroup. 10 main lineages were defined where each lineage had more than 10 isolates. STs of the 7 gene MLST (or MGT1) and CCs from MGT2 to MGT4 were shown in different colours for each of the 10 lineages. The pie charts for each lineage represent the proportion of the isolates belonged to the same MGT4-CC shown on the right side.