Spatial summation of pain increases logarithmically

Wacław M. Adamczyk[1,2]*, Linn Manthey[1], Christin Domeier[1], Tibor M. Szikszay[1], Kerstin Luedtke[1,2]

[1]Institute of Health Sciences, Academic Physiotherapy, Pain and Exercise Research Luebeck (P.E.R.L.),

University of Luebeck, Luebeck, Germany.

[2]Laboratory of Pain Research, Institute of Physiotherapy and Health Sciences, The Jerzy Kukuczka

Academy of Physical Education, Katowice, Poland

*Corresponding author: Department of Health Sciences, Academic Physiotherapy, Pain and Exercise

Research Luebeck (P.E.R.L.), University of Luebeck, Luebeck, Germany, Ratzeburger Allee 160, 23562

Lübeck, Tel +49 451 3101 8541,

email: waclawadamczyk@gmail.com

Number of pages: 22

Number of tables: 5

Numbers of figures: 6

Word count: 7277

ABSTRACT

Pain intensity is difficult to predict. Mostly, because of modulatory processes underlying its formation. For example, when nociceptive stimulation occupies a larger body area, pain increases disproportionally. This modulation is called spatial summation of pain (SSp) and is responsible for coding pain intensity. To predict pain based on spatial variables, a profound understanding of the SSp effect is crucial. The aim of this study was i) to describe the SSp effect as a function of the size (or distance) of a stimulated area(s), ii) to investigate the effect of pain intensity on SSp and iii) to evaluate the influence of the SS type on the magnitude of SSp. Thirty-one healthy participants took part in a within-subject experiment. Participants were exposed to area- and distanced based SSp. In the former, electrocutaneous noxious stimuli were applied by up to 5 electrodes (5 areas) forming a line-like pattern at the ulnar side of the hand, while in the latter the same position and lengths of stimuli were used but only two electrodes were stimulated (5 separations). Each paradigm was repeated using pain of low, moderate and high intensity in a random and counterbalanced order. Each stimulus was assessed on a 0-100 scale. It was found that the pattern of increase in pain followed a logarithmic rather than a linear function. The dynamics of the pain increase were statistically different across pain intensities, with more summation occurring, if stimuli were calibrated to eliciting "high" pain. SSp was resistant to saturation in the area-based but not in the distance-based SSp, where 0.8cm separation between two electrodes produced a similar pain intensity as 1.6cm and 2.4cm. Results indicate that area-based SSp is more painful than distance-based SSp when low and moderate but not when high pain intensity is induced. Presented findings have important implications for all studies, in which the spatial dimension of pain is measured. When the area or separation between nociceptive stimulation increases, pain does not increase linearly. Furthermore, the pattern of the pain increase depends on i) intensity and ii) the number of sites of nociception. In conclusion, a logarithmic function should be considered when predicting the size of a nociceptive source. This pattern is indicative for inhibitory processes underlying SSp.

Key words: lateral inhibition; prediction model; nociception; pain modulation; temporal summation

INTRODUCTION

Pain is an adaptive response to noxious stimuli that protects humans from tissue damage. As in other sensory systems, nociceptive stimuli, which often -yet not always- lead to pain perception, are influenced by ascending (bottom-up) [54] and descending (top-down) [6,27,28] pain modulation. The same stimulus might be perceived as less or more painful depending on psychological, social and biological factors (see [1,13] for review). The latter category is particularly interesting as it reflects competitions between noxious stimuli targeting the neuroaxis with different temporal and spatial relations. For instance, research has shown that one stimulus is perceived as less painful if it is preceded by a heterotopic counter-conditioning stimulus [42,51], a phenomenon known as diffuse-noxious inhibitory-control (DNIC). However, if a pair of (or more) noxious stimuli occur concomitantly, they evoke pain facilitation, known as spatial summation of pain (SSp) [10–12,22,36,38,41,47,52].

Spatial summation of pain[1] has been studied in animals [6,14,43] and humans [10–12,22,36,38,41,47,52] using two distinct paradigms: 'distance-based' and 'area-based' SSp [41]. In the former, an increase in pain is reported when the distance between stimuli increases, while in the latter, more pain is reported when noxious stimuli are applied to a larger area. Spatial summation of pain is essential for the detection of pain [2], coding of pain intensity [7] and identification of pain quality [8].

Transferring this knowledge to a patient population, the most obvious examples are patients with chronic widespread pain (CWP). Research in this population has shown that the intensity of pain perception can be predicted by the number of painful body parts [46,48], therefore, it is likely that the SSp mechanism is altered in this clinical population. Nevertheless, previous attempts failed to show differences in SSp response profiles between healthy controls and chronic pain patients [15,18,45,47], which raised the question about the mechanism of SSp. Interestingly, previous studies in which other paradigms, also testing pain modulation, such as temporal summation [33], offset analgesia [49] or DNIC, showed distinct differences between chronic patients compared to healthy controls.

---

[1] Semantically, a more correct way to describe SSp in animal is spatial summation of nociceptive processing. This is due to the IASP definition of pain and its recent updated version, according to which, pain is a subjective experience.

The reason for this discrepancy is a relatively poor understanding of SSp in humans. Previous

work has been mostly limited to psychophysical testing of SSp, with some physical or body-related

manipulations made to study SSp mechanisms, e.g. skin types [16], modality [23,31], age [26], pain [47],

stimulus intensity [40], body location [39], sequence of testing, [31] or the type of paradigm used [38]

were considered. Notwithstanding to that, it is unclear where in the humans' neuroaxis the summation

occurs and how intensity shapes the magnitude of the SSp. Behavioral studies are a first line of research

that may approximate the underlying mechanisms of complex noxious processing.

In the past, SSp has been studied using rather static 'paradigms'. In that sense, area-based SSp

was provoked by comparing responses to a small and a large probe. For distance-based SSp, various

separations have been used to investigate this effect, often with small inter-spaced resolution, e.g. 5cm or

10cm steps (separations). Whether summation also occurs using smaller scales, has never been confirmed.

Considering mentioned limitations, a mathematical modeling of the SSp response was not possible, due to

the usually limited number of probes that were used.

The aim of this study was therefore to introduce a paradigm which investigates SSp as a function

of a stimulated area to create adequate pain predictions. Secondly, the experiment aimed to investigate the

effect of the experimentally induced pain intensity on the magnitude of SSp, described as a mathematical

function. It was hypothesized that pain increases less dynamically as the pain intensity increases. Such a

prediction is based on the evidence showing that more pronounced pain inhibition occurs, when more

severe [44,53] and larger [23,31] stimuli are used. The associated higher activation of pain-inhibitory

brain regions is linked to stimulus aversiveness. Lastly, we developed a study design to investigate SSp at

a micro-scale level, with stimuli forming a continuum of < less than 4cm in length. Such a design allows

to explore the relationship between the type of SSp (area-based, distance-based) and the effect of the

stimulus intensity, as well as their interactions.

MATERIALS AND METHODS

Study overview and structure

This study was based on a within-subject design to test the effect of SSp type and stimulus intensity on the pattern of pain summation. The structure of single session was similar to those described previously [32,37] and was subdivided into familiarization, calibration and the main data collection phase (SSp assessment). During the last phase, participants were assessed using two paradigms (area- and distance-based SSp) with three different pain intensities (low, moderate and high pain). One session was repeated three times.

The Ethics Committee of the University of Lübeck approved the protocol of this study (decision no. 19-303), which was preregistered in December 2019 at the osf.io platform (https://osf.io/qry9d), using the AsPredicted.org template. The study follows the principles of the Declaration of Helsinki, developed by the World Medical Association. Each participant was adequately informed about the objectives, methods, the anticipated benefits and potential risks and the discomfort, as well as any other relevant aspects of the study. A written informed consent was obtained from each participant before participation in the study.

Study population

Fifty-three volunteers signed up to participate in the study. Each volunteer was screened for eligibility. To be included in the study, participants had to be healthy (self-report), right handed, over the age of 18 years and have sufficient German or English language skills. Exclusion criteria were any acute or chronic pain, skin pathologies or tattoos in the area of the left hand, diagnosed neurological, cardiovascular or psychiatric diseases or any other disease requiring regular medication intake. Other exclusion criteria were pregnancy, metal implants or electronic devices in or on the body. The exclusion criteria were established to avoid any changes in pain perception and possible health-related risks during the application of electronically induced pain. Twenty-two volunteers were not considered eligible (reasons provided in Appendix 1) and 31 (15 females) participated and completed the experiment (mean age 26.2 ± 6.8 years, height 174.0 ± 9.9 cm, weight 68.9 ± 13.5 kg).

As it has been shown that both, caffeine and alcohol consumption, affect the sensation of pain

[24,50], participants were also asked to refrain from consumption on the day of participation. They were

further asked to not take any drugs 24 hours before participating in the study. In order to avoid a DNIC

effect, participants were asked to not engage in excessive sports that could cause muscle ache or other

pain 48 hours before participation in the study.

Experimental setting and materials

The trial took place in a quiet and temperature-controlled (20.5 ± 0.5 °C) laboratory. All

assessments were performed by the same examiner. The participants were set on a chair in front of a

monitor with a distance of 50cm. The 23.8-inch monitor was set up 1m above the floor and the chair was

50cm high. The participant was separated from the examiner by a moveable curtain. Pain was assessed

using a Numerical Rating Scale (NRS) after each stimulus application. The scale was anchored from 0

("no pain") to 100 ("most imaginable pain").

The electrocutaneous stimulation was produced by a Constant Current Stimulator (Digitimer,

model DS7A, United Kingdom, Welwyn) and a remote selector device (Digitimer, model D188, United

Kingdom, Welwyn) with 200μs duration of each electrocutaneous pulse and a capacity of 0 to 100 mA

with a maximum voltage of 400V, which is based on previous SSp studies [22,41]. External control of

DS7A and D188 was ensured via the Labjack U3 control device (LabJack Corporation, Lakewood, CO,

USA). The procedure was fully automatic and operated by the PsychoPy 2.0, open-source software [35].

Five wasp-like gold chloride sintered electrodes (Brainbox Ltd., Cardiff, UK) with 0.8cm diameter were

attached first to the armrest of the armchair, where the participants were seated and later to their hand.

Before attaching the hand electrodes, the ulnar edge of the hand was cleaned with an abrasive gel and an

alcohol wipe using a cotton pad to remove all particles, which could have a negative impact on the contact

between electrodes and skin.

Electrodes were placed just below the skin fold under the metacarpophalangeal joint of the fifth

phalanx (ulnar side of hand). The hand was placed in a neutral position between pronation and supination

and the elbow was rested at 90° of flexion. Electrodes were placed next to each other (with no space

between adjacent electrodes). Such an orientation allowed to study area-based SSp with up to 5 electrodes

activated simultaneously. For distance-based SSp, the same configuration was used. However, distanced-

based SSp was assessed when two electrodes (with randomly chosen distances between them), were

activated. In principle, area-based SSp is established when one manipulates the size of the stimulated area

and distance-based SSp, when different separations are used. Having this conceptual difference in mind,

the following definitions were used for clarity: Different areas were determined by the number of

electrodes activated (up to 5) and different separations were determined by the number of electrodes

forming the gap between two activated electrodes. As the electrodes were placed in a line, activation of 5

electrodes induced a stimulus of 4cm length (5 × 0.80cm) that served as the maximal area in area-based

paradigm. The same distance (4cm), formed by two outermost electrodes, was used as the maximal

separation for the distance-based paradigm (Fig. 2). Thus, technically, the stimuli had the same length,

regardless of the SSp type being assessed.

Familiarization and calibration

To check if the conduction under all 5 electrodes is accurate, an initial testing procedure was

performed. The electrodes were activated sequentially with a fixed intensity of 6.0 mA and the detection

of the stimuli were verbally reported by the participants. This procedure was also performed to make

participants familiar with electrocutaneous stimulation.

Calibration

Participants underwent 4 series of calibration trials to consider the individual differences in pain

perception and to determine the individual intensities for the subsequent assessment of SSp. The objective

of the calibration was to obtain pain corresponding to an NRS of 30 (low), 50 (moderate) and 70 (high

intensity). Thus, the influence of different pain conditions on the magnitude of SSp could be evaluated.

To calibrate the individual intensity levels, first the tactile (t) and pain (P) thresholds were

determined [4,5]. This was done with electrode 1 and 2 activated with a separation of 0cm (shortest

possible distance) for distance-based SSp and also with only one electrode activated (smallest possible

area) for the area-based SSp. Furthermore, t and P were determined for the largest distance possible,

which is equivalent to an activation of electrode 1 and 5 (distance-based SSp) and also with the largest

area, which is equivalent to an activation of all 5 electrodes (area-based SSp). For each calibration part,

the electrical current started from 0.0 mA and was increased by 0.5mA, until the participant verbally

reported a very first sensation (t). The intensity was further increased by 0.5mA until the first pain

perception was reported (P).

Based on these determined thresholds, participants received a random sequence of painful and

non-painful electrocutaneous stimuli of different intensities. The intensities were calculated using the

formulas depicted in Fig. 1. The participant was instructed to rate every stimulation on the NRS by

selecting a level that best corresponded to the perceived intensity. Based on these pain ratings, a

regression line for the stimulus-response function was individually plotted. The x-axis represents the

electrical current in mA, the y-axis the individual pain ratings.

Figure 1. Calibration phase. Calibration started from tactile (t) and pain threshold (P) determination. Based on these results, 7
stimuli of different intensities were applied in a random order. Then different pain intensities corresponding to low (30 on
Numeric Rating Scale, NRS), moderate (50) and high (70) pain were readout from regression lines plotted as a function of pain
and intensity (in mA). In the end, the given intensity was checked for accuracy in the pain induction of assumed intensity
(calibration check).

Through the formula obtained from the given regression line, a predicted intensity (in mA) for all

pain conditions (NRS 30, 50 and 70) was determined, using small (short) and large (long) configurations

(Fig 1). An average of predicted electrical currents was then tested in the calibration check [30]. The

calculated stimulus intensities for the different pain levels were applied during the check, if predicted

values from the regression models induced pain at the targeted level (pain of 30, 50 and 70, respectively).

The same final intensities were used for both SSp types, to compare pain summation between paradigms.

Spatial summation assessment

To assess SSp, different stimuli configurations were used, as described earlier. The line-like

pattern of 5 electrodes, allowed to apply 5 different stimulus intensities within each SSp type (see Fig. 2).

For the area-based paradigm, the control (one) electrode was activated separately or simultaneously with a

sum of up to 5 electrodes. Therefore, stimuli forming a filled line ranging from 0.80cm in length up to

4cm were applied. A series of 5 stimuli with jittered 5-7s inter-stimulus intervals (ISI) was applied in a

fully random sequence and repeated three times (>10s break between the series) for each pain condition,

i.e. low, moderate and high pain.

For the distance-based paradigm the same structure was maintained. Here the stimuli had the same

line-like pattern (from 0.80 to 4cm), however, the control electrode was activated separately or

simultaneously with a second electrode 0.0, 0.8, 1.6, or 2.4cm apart.

The order of SSp type, and the order of pain conditions (low, moderate, high) was randomly

assigned, yet counterbalanced across subjects. For each paradigm (area- and distance-based SSp), there

were 45 stimuli induced, representing 5 configurations $\times$ 3 series $\times$ intensities, so that each participant

received 90 stimuli in total during the assessment phase. As there were 3 sessions employed, each

participant received 270 stimuli.

219
220
Figure 2. Structure of the spatial summation assessment (SSp). A) For every pain condition (pain at the level of 30, 50, or 70
on 0-100 NRS scale), 5 different stimuli configurations were used. In the area-based (dark blue) SSp either one electrode or up to
5 electrodes were activated simultaneously. In the distance-based paradigm (cyan) either one electrode or two electrodes with a
varied separation were activated simultaneously. In principle, for both SSp types, stimuli formed line-like pattern, with filled or

unfilled areas (with different separations). The order of the pain conditions and electrode configurations were randomized and
counterbalanced across participants. **B)** For every pain condition there were three series consisting of 5 stimuli each. In total, 15
stimuli were applied for every pain intensity, thus 45 stimuli were used per SSp type (90 stimuli in total). The order of the stimuli
was randomized and counterbalanced across the assessment. AB, area-based SSp, DB, distance-based SSp, E, electrode. Note,
that the same intensities determined individually for the given condition (low, moderate and high pain), were used for both SSp
types.
231

Data processing and statistical analysis

Raw pain ratings were first extracted from the files generated by the PsychoPy software using

MATLAB R2017b (MathWorks Inc, Natick, Massachusetts). Descriptive statistics of all collected data

were presented as means and SD or their non-parametric equivalents. The analysis was based on a

complete dataset from 31 participants.

Spatial summation of pain was analysed in according to the following protocol: To reduce data for

the main analysis, a General Linear Model (GLM) was applied to test for significant difference in pain

perception between measurement sessions. Two GLM analyses were performed for area-based and

distance-based SSp, separately. The main SSp analyses were performed on the pooled dataset from three

sessions. Namely, a $2 \times 3 \times 5$ GLM analysis was performed with within-subject factors 'paradigm' (area-

based, distance-based SSp), 'intensity' (low, moderate, high) and 'stimulus configuration' (line of 0.8, 1.6,

2.4, 3.2, 4cm length). In case of significant main or interaction effects, post-hoc Bonferroni corrected

paired *t*-tests were applied to investigate the meaning of the effect.

To express SSp in one unified coefficient, linear ($y = b*x + a$) and logarithmic ($y = b*\log_{(x)} + a$)

curves were fitted to individual subjects for area-based and distance-based SSp in respect to stimulus

intensities. Thus, six linear and logarithmic curves were fitted to the data per subject using the Curve

Fitting Toolbox in MATLAB R2017b (MathWorks Inc, Natick, Massachusetts). Goodness of fit was

assessed visually and by comparing R^2 statistics generated by two different fittings using paired t-tests and

by one-sample t-tests against perfect fit of 1.0 value. The magnitude of SSp was expressed as the slope (β-

coefficient) of the curve where -in case of logarithmic distribution of the data- higher values mean more

pronounced SSp as the summation increases more dynamically together with the linear increase in the

stimulated area (or separation). To further demonstrate the SSp effect across different intensities, a GLM

model was used with β-coefficients as dependent variables and 'paradigm' (area-based and distance-based

SSp) and intensity (low, moderate, high) as two within-subject factors. Post-hoc corrected paired t-tests were applied if necessary.

Analyses were conducted using the STATISTICA data analysis software, version 13 (StatSoft Inc., Tulsa, OK, USA). The level of significance was set at $p < 0.05$ and Bonferroni correction was applied to control for family-wise error. When p-values did not exceed α levels after correction, it was marked as not significant (ns).

RESULTS

Thirty-one healthy participants took part and finished this experiment. Participants' characteristics are presented in Table 1. General Linear Model on 'pain ratings' as dependent variable and 'session' and 'stimuli' as within-subject factors revealed no significant effect of 'session', neither for area-based ($F_{(2,60)} = 0.61$, $p = 0.54$, $\eta_p^2 = 0.02$) nor for distance-based SSp ($F_{(2,60)} = 1.16$, $p = 0.32$, $\eta_p^2 = 0.04$). Thus, datasets were combined across sessions and subsequent analysis was performed on pooled dataset.

Table 1. Characteristics of study participants (n = 31)

Variable	Statistic
Female N (%)	15 (48.39)
Age (SD)	26.19 (± 6.83)
Palmar-wrist crease in cm (SD)	16.5 (± 1.4)
Hypothenar length in cm (SD)	6.58 (± 0.93)
Weight in kg (SD)	68.87 (± 13.46)
Height in cm (SD)	173.97 (± 9.88)
Fear of pain (SD)	3.23 (± 1.82)
PVAQ (SD)	39.39 (± 9.14)
PCS (SD)	9.58 (± 8.16)

SD, Standard Deviation, PVAQ, Pain Vigilance and Awareness Questionnaire, PCS, Pain Catastrophising Scale.

The main GLM analysis performed on pain ratings showed a significant main effect of 'SSp' ($F_{(1,30)} = 21.47$, $p < 001$, $\eta_p^2 = 0.42$), 'intensity' ($F_{(2,60)} = 245.47$, $p < 0.001$, $\eta_p^2 = 0.89$) and 'stimuli' ($F_{(4,120)} = 122.50$, $p < 0.001$, $\eta_p = 0.80$). These three main effects indicated that i) in general, area-based SSp was more painful compared to distance-based SSp (Fig. 3 and 4), ii) calibration was successful as participants discriminated pain levels significantly ($p < 0.001$), and iii) SSp was statistically significant as -in general- pain increased with the area or distance between two stimuli ($p < 0.001$, Fig. 3). Furthermore,

the GLM showed significant 'SSp' × 'intensity' ($F_{(2,60)} = 3.26$, $p < 0.05$, $\eta^2_p = 0.10$), 'intensity' × 'stimuli' ($F_{(8,240)} = 9.19$, $p < 0.001$, $\eta^2_p = 0.23$) interactions as well as an interaction between the factors 'SSp' and 'stimuli' ($F_{(4,120)} = 76.82$, $p < 0.001$, $\eta^2_p = 0.72$).

Figure 3. Spatial summation of pain (SSp). The pain intensity increases with increasing the stimulated area (AB) or distance (DB) between two stimuli in all pain conditions. Behaviour of area-based (AB) and distance-based (DB) SSp under the pain of low (left), moderate (middle) or high intensity (right). Note that in general, area-based SSp was more painful than distance-based but not when the highest intensity was used. Each figure presents mean values (dots) with standard errors of the mean (SE). ** $p < 0.01$.

Results of post-hoc tests following these interaction effects revealed that i) area-based was more painful than distance-based SSp but only when low ($p < 0.001$) or moderate ($p < 0.001$) but not when high intensity ($p = 0.16$) was applied (Fig. 4), ii) further increases in area or distance did not lead to a systematic increase in pain perception; regardless of the SSp type, the SSp effect was saturated when comparing the two outermost stimuli, i.e. 3.2cm vs. 4cm length (Fig. 3). The stimulation of the largest area consisting of 5 electrodes produced similar pain level compared to the slightly smaller area formed by 4 electrodes when low ($p = 0.06$), moderate ($p = 0.10$) or high ($p = 0.18$) intensity condition was considered, iii) the behaviour of SSp is different across SSp types: Area-based SSp was not saturated over the course of the linearly increased size of the stimulated area, as the pain level still increased significantly when the stimulus was enlarged (all p values < 0.01, except for 0.8 vs. 1.6cm stimuli - n.s. after correction). However, in the distance-based paradigm, pain was saturated and did not further increased when enlarging the distance, 0.8cm separation produced similar pain as 1.6cm (Fig. 3, $p = 0.99$) and 2.4cm ($p = 0.77$).

Figure 4. Pain intensity across spatial summation paradigms. Comparing the types of SSp, the area-based (AB) paradigm was perceived as more painful compared to the distance-based (DB) paradigm except for the highest intensity: here relatively the same level of pain was induced even though 2.5 times less area was stimulated in distance-based paradigm. ** $p < 0.01$.

Table 2. Descriptive statistics for pain ratings

Intensity	Area (separation)	Area-based		Distance-based	
		Mean	**SD**	**Mean**	**SD**
Low pain (30/100)	0.8 cm	15.64	8.31	15.64	7.57
	1.6 cm	30.29	11.54	28.87	11.54
	2.4 cm	39.37	12.96	32.56	12.19
	3.2 cm	44.70	13.73	32.57	12.08
	4.0 cm	50.03	14.19	34.76	14.12
Moderate pain (50/100)	0.8 cm	26.84	10.37	33.84	15.34
	1.6 cm	49.48	13.23	50.52	16.89
	2.4 cm	59.67	12.44	55.86	15.92
	3.2 cm	67.20	12.17	55.20	15.90
	4.0 cm	71.70	12.30	55.95	14.27
High pain (70/100)	0.8 cm	42.48	16.74	49.24	14.88
	1.6 cm	64.35	15.63	67.08	16.05
	2.4 cm	75.02	11.50	71.22	14.07
	3.2 cm	80.76	10.40	71.48	13.63
	4.0 cm	83.94	9.53	73.18	13.67

SD, Standard Deviation.

Lastly, GLM analysis showed a significant three-way 'SSp' × 'intensity' × 'stimuli' interaction ($F_{(8,240)} = 3.06$, $p < 0.01$, $\eta^2_p = 0.09$), indicating that the pattern of pain increase was different across the SSp types in respect to the intensity. Similar pain levels across SSp types were observed when the stimulus of 1.6cm (or 0cm separation between 2 stimuli) was applied with low ($p = 0.99$) moderate ($p = 1.00$) and high ($p = 0.43$) intensity. Interestingly, starting from the third level of stimuli (2.4cm stimuli

309 length), area-based was more painful than distance-based SSp (Fig. 3), however, when a single stimulus

310 was contrasted across SSp paradigms, area-based SSp was paradoxically less painful for moderate ($p <$

311 0.05, Fig. 3) and high intensity ($p < 0.05$, Fig. 3). This, however, might reflect procedural aspects of the

312 experimental paradigm such as higher stimuli contrast in area-based paradigm.

313 **Table 3. Curve fitting: one-sample t-tests results (mean R^2 value against value of 1)**

SSp	Intensity	N	Logarhitmic				Linear			
			R^2	SD	t	p	R^2	SD	t	p
	Low	31	0.86	0.11	-7.40	< 0.001	0.81	0.11	-9.31	< 0.000
AB	Moderate	31	0.86	0.09	-8.64	< 0.000	0.80	0.12	-9.74	< 0.000
	High	31	0.88	0.07	-9.52	< 0.000	0.79	0.11	-10.24	< 0.000
	Low	31	0.51	0.22	-12.49	< 0.000	0.45	0.22	-13.83	< 0.000
DB	Moderate	31	0.55	0.21	-11.86	< 0.000	0.47	0.23	-12.87	< 0.000
	High	31	0.60	0.17	-12.86	< 0.000	0.50	0.19	-14.37	< 0.000

AB, area-based, DB, distance-based, SSp, spatial summation of pain, SD, standard deviation

Studying pain behaviour across the full range of areas and distances allowed for visual assessment

of the stimulus-response functions (Fig. 3), which suggested that the pattern of pain increase followed

logarithmic function. When comparing R^2 coefficients for logarithmic and linear models by using one-

sampled (R^2 against referenced value of 1.0) and two sampled paired t -tests (R^2 across models), it was

found that i) lower t statistics were observed for the logarithmic compared to the linear fitting (Table 3)

and ii) that logarithmic curves showed significantly higher R^2 statistics compared to linear fits (Table 4).

Table 4. Curve fitting: paired t-tests results for logarithmic and linear comparisons

SSp type	Intensity	Mean difference	95% CI	t	p
	Low	0.05	(0.02 - 0.08)	3.46	< 0.01
AB	Moderate	0.06	(0.04 - 0.09)	4.95	< 0.000
	High	0.09	(0.06 - 0.12)	6.58	< 0.000
	Low	0.06	(0.03 - 0.09)	3.97	< 0.000
DB	Moderate	0.08	(0.05 - 0.11)	5.11	< 0.000
	High	0.10	(0.07 - 0.13)	6.46	< 0.000

AB, area-based, DB, distance-based spatial summation of pain, SSp, spatial summation of pain, CI, confidence interval

General Linear Model performed on beta coefficients from logarithmic fittings showed a

significant main effect for the factor 'SSp' ($F_{(1,30)} = 164.66$, $p < 0.001$, $\eta_p = 0.85$) and 'intensity' ($F_{(2,60)} =$

17.02, $p < 0.001$, $\eta^2_p = 0.36$), indicating that i) in-general, area-based SSp was characterized by higher

beta values and ii) that moderately painful ($p < 001$) and highly painful ($p < 001$) stimuli lead to a more

dynamic pain increase, compared to the least painful stimuli (Fig. 5). No difference between moderately

and highly painful stimuli was detected ($p = 0.81$, Fig. 5). Moreover, the GLM showed a significant 'SSp'

× 'intensity' interaction ($F_{(2,60)} = 5.18$, $p < 0.01$, $\eta^2_p = 0.15$) wherein post-hoc comparisons confirmed

that: i) in the area-based paradigm, moderately ($p < 0.01$) and highly painful stimuli ($p < 0.01$) led to

higher beta values than SSp induced via stimuli of the least painful intensity (Fig. 5), ii) distance-based

SSp showed significantly higher beta-values for condition with high pain compared to the least painful

stimuli, only ($p < 0.03$, Fig. 5). Non-linear fits showing that spatial summation of increases

logarithmically are presented in Fig. 6 and Table 5.

Figure 5. The magnitude of spatial summation of pain (SSp). Spatial summation of pain was expressed as beta coefficients of
non-linear logarithmic functions. Higher beta values for area-based (AB) compared to distance-based (DB) were observed. The
magnitude of SSp was intensity-dependent: when stimuli were calibrated to induce pain at the level of approximately 70 out of
100, the pain then increased more rapidly. Figure presents mean values with standard errors of the mean (SE). *$p < 0.05$, **$p <$
0.01, ***$p < 0.001$.

Figure 6. Spatial summation of pain increases logarithmically. The pain increases with increasing the area stimulated or the distance between two stimuli in a logarithmic fashion.

Table 5. Descriptive statistics for logarhitmic and linear fits for SSp

SSp type	Intensity	Logarhitmic				Linear			
		β		R^2		β		R^2	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD
AB	Low	21.26	8.26	0.86	0.11	8.32	3.29	0.81	0.11
	Moderate	27.98	9.64	0.86	0.09	10.74	3.68	0.80	0.11
	High	26.15	9.61	0.88	0.07	9.93	3.65	0.79	0.11
DB	Low	11.52	6.96	0.51	0.22	4.19	2.91	0.45	0.22
	Moderate	13.79	8.67	0.55	0.21	4.89	3.61	0.47	0.23
	High	14.57	7.00	0.60	0.17	5.23	2.77	0.50	0.19

345   AB, area-based, DB, distance-based spatial summation of pain, SSp, spatial summation of pain, SD, standard deviation

DISCUSSION

The aim of this study was to investigate two types of SSp and to compare the summation trajectories when using individually calibrated noxious electrocutaneous stimuli. Three novel findings are reported, relating to the processing of the spatial information in the pain system: It was found that the pain increases logarithmically as a non-linear function and that the pattern of pain increase was statistically different across three different pain intensities, however in a not anticipated direction. SSp became saturated in the area-based but not in the distance-based SSp. Moreover, saturation was more likely to be observed when lower intensities were applied. Such an observation was possible to capture using curve fitting. The summation of pain was expressed as a function and decomposed to the value of one logarithmic coefficient. Coefficients were of higher values when the baseline pain level was higher than 50. And finally, data from our paradigm showed that area-based SSp is more painful than distance-based SSp, however this effect depends on how the painful stimuli are.

The long history of research on SSp encountered difficulties from the very beginning. For example, early reports failed to show SSp at all, possibly due to the insufficient stimuli intensity [17,20]. Later reports confirmed that stimulus intensity shapes the magnitude of SSp [36,40,52], suggesting that larger SSp is more likely to be observed in higher stimulation intensities. The putative mechanism of this relative increase in summation was attributed to a more pronounced excitation of spinal nociceptors [40] at higher intensities. These studies, however, have shown thermal SSp, expressed as the raw difference in pain ratings between two [40,52] or a maximum of three conditions (areas or separations) [36]. The limited number of probes determined by the equipment used in previous studies, prevented from testing a predictive model. In the current study, five wasp-like electrodes were used for each SSp type, which allowed to investigate a summation pattern. It was found that a logarithmic function described more variance of the data compared to a linear fit. Interestingly, this pattern was observed for low, moderate and high pain conditions, confirming the validity of the model. The beta values were, however, higher when the intensity was calibrated to induce a high level of pain (>70/100), confirming preliminary observations, where lower SSp was observed for low pain intensities. This seems like a paradox. On the one hand, the

pain increase is i) disproportional and ii) non-linear, but on the other hand, more intense stimuli elicit a

larger SSp effect. The former results might indicate inhibitory processes underlying SSp, while the later

points towards the contribution of larger excitation waves. In some circumstances, the non-linear nature of

SSp might be of advantage for any living organism: Inhibitory processes saturate pain perception but not

for very intense pain. Considering that the pain level may in acute situations reflect the magnitude of

potential tissue damage, the bigger the damage (nociception) the bigger the pain increase. Thus, it seems

that the greater magnitude of the SSp at higher intensities serves a protective mechanism preventing the

individual from massive tissue damage.

The paradox described here may be explained by the variety of neural processes involved in the

human [3,38] and non-human [6,21] neuroaxis, including descending pain inhibition [6], lateral inhibition

[38] or even expectancies. Spatial summation has been originally linked with facilitation at the single

neuron level. For example, Sherrington (1906) has used this term to describe the generation of action

potential when subliminal -separated in space- stimuli were applied. Interestingly, first experiments on

SSp in rats showed spatial summation to be disproportional [6]. Bouhassira et al. [6] found that

convergent neurons exhibit enhanced electrical activity only when the spinal cord was sectioned and the

effect of the descending loop was thereby reduced. This finding, although limited to rat studies, indicates

that descending pain inhibition might play a role in the SSp magnitude and shape. Furthermore, a vast

majority of human studies found similar observations: SSp has been reported as subadditive, as the

reported pain marginally or modestly increased even though the stimulated area was doubled [9], tripled

[39], quadrupled [52], or even six times larger [26]. The subadditivity supports the role of descending pain

system. Indeed, the periaqueductal grey matter (PAG) has been found as core midbrain region of pain

modulation [19,25,34]. Enhanced activity of this region has been found in hypoalgesic or hyperalgesic

modulation, which presumably is reflect in its anatomical structure [29]. Whether PAG activation is

intensity-dependent is unclear and warrants future investigations.

The fact that a significant interaction between the type of SSp and the relative painfulness was

detected indicates that, potentially, lateral inhibition interplayed with SSp. For example, in the study by

Quevedo et al. [38], those trials were much more painful, in which only two points were separated by e.g.

8cm compared to a line stimulus applied by laser. In the current study, when the high intensity condition

was applied, there was no difference between area- and distance-based SSp which is partially in line with

this previous report. Although this current data did not reveal a similar trend as in the study by Quevedo et

al. [38], an interesting observation was, that despite a smaller number of nociceptors activated in

distanced-based SSp, an equal pain responses were noticed. The lack of replication of the full findings of

Quevedo et al. are methodological differences, mostly related to stimuli types and timing of stimuli

application. In that sense, for an inhibitory response, the temporal relation between spatially distinct

stimuli might be of important for the prediction of an accurate spatial summation effect.

For the first time, area-based and distance-based SSp were compared. Area-based SSp was not

saturated over the course of the linear increase in size of the stimulated area. Pain successfully increased,

when the area increased but not when the distance between two points became larger. Furthermore, this is

the first report which proved that SSp is a local effect as we used a continuum of areas, or distances, in a

line-like manner of up to 4cm, therefore the current findings complement previous SSp research using

laser, cold, pressure or electrocutaneous stimuli.

Although the current findings are robust in terms of internal validity, some aspects could have

affected the SSp estimations: Circular electrodes were placed close to each other with no space in

between. This solution was sufficient to manipulate area size; however, they were connected via only one

tangential point. Such a solution has been used previously, yet for future experiment it is recommended to

use square-like stimuli. Trains of pulses produced a strong pain at a relatively low intensity (<12 mA).

This could be a result of temporal summation processed as 'noise' during the spatial summation.

Notwithstanding to that, temporal facilitation was limited to single-stimulus only, and seems unlikely to

affect the pattern of spatial summation described in current study. The paradigm used in this study was

based on a random sequence of different stimulus types, with three repetitions each. Such a small number

of trials however, was compensated but three separate examination sessions for each participant, which

likely affected the precision of SSp estimations. Furthermore, as no difference between the three sessions

was detected it seems that the paradigm used and SSP effect itself is stable over-time.

Conclusions

Taken together, results presented here are an important step forward in the understanding of the

pain prediction in relation to spatial configurations of noxious stimuli. Our data indicates that the relative

"painfulness" of noxious stimuli affects the non-linear summation trajectory, however, the hypothesis that

reduced SSp magnitude is observed when high intensity of stimuli is used was not supported. It seems

however, that SSp increases logarithmically, which may suggest the SSp effect is unique mixture of

facilitatory and inhibitory processes.

ACHNOWLEDGEMENTS

Authors declare no conflicts of interest. Data presented here constitute a part of the preregistered

project (https://osf.io/qry9d) dedicated to test if the pain intensity affects magnitude and reliability of SSp.

Due to the significant amount of data generated in this project, reliability results will be presented in other

occasion.

REFERENCESS

[1]   Adams LM, Turk DC. Central sensitization and the biopsychosocial approach to understanding pain.
Journal of Applied Biobehavioral Research 2018;23:e12125.
[2]   Adriaensen H, Gybels J, Handwerker HO, Van Hees J. Latencies of chemically evoked discharges in human
cutaneous nociceptors and of the concurrent subjective sensations. Neurosci Lett 1980;20:55–59.
[3]   Apkarian AV, Gelnar PA, Krauss BR, Szeverenyi NM. Cortical Responses to Thermal Pain Depend on
Stimulus Size: A Functional MRI Study. Journal of Neurophysiology 2000;83:3113–3122.
[4]   Babel P, Adamczyk W, Swider K, Bajcar EA, Kicman P, Lisinska N. How Classical Conditioning Shapes
Placebo Analgesia: Hidden versus Open Conditioning. Pain Med 2018;19:1156–1169.
[5]   Bajcar EA, Adamczyk WM, Wiercioch-Kuzianik K, Bąbel P. Nocebo hyperalgesia can be induced by
classical conditioning without involvement of expectancy. PLoS ONE 2020;15:e0232108.
[6]   Bouhassira D, Gall O, Chitour D, Le Bars D. Dorsal horn convergent neurones: negative feedback triggered
by spatial summation of nociceptive afferents. Pain 1995;62:195–200.
[7]   Coghill RC, Mayer DJ, Price DD. The roles of spatial recruitment and discharge frequency in spinal cord
coding of pain: a combined electrophysiological and imaging investigation. Pain 1993;53:295–309.
[8]   Defrin R, Ohry A, Blumen N, Urca G. Sensory determinants of thermal pain. Brain 2002;125:501–510.
[9]   Defrin R, Ronat A, Ravid A, Peretz C. Spatial summation of pressure pain: effect of body region. Pain
2003;106:471–480.
[10]   Defrin R, Sheraizin A, Malichi L, Shachen O. Spatial summation and spatial discrimination of cold pain:
effect of spatial configuration and skin type. Pain 2011;152:2739–2745.
[11]   Defrin R, Tsedek I, Lugasi I, Moriles I, Urca G. The interactions between spatial summation and DNIC:
effect of the distance between two painful stimuli and attentional factors on pain perception. Pain
2010;151:489–495.
[12]   Defrin R, Urca G. Spatial summation of heat pain: a reassessment. Pain 1996;66:23–29.

[13]   Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL. Sex, gender, and pain: a review
of recent clinical and experimental findings. J Pain 2009;10:447–485.
[14]   Gall O, Villanueva L, Bouhassira D, Le Bars D. Spatial encoding properties of subnucleus reticularis
dorsalis neurons in the rat medulla. Brain Res 2000;873:131–134.
[15]   Goubert D, Danneels L, Graven-Nielsen T, Descheemaeker F, Meeus M. Differences in Pain Processing
Between Patients with Chronic Low Back Pain, Recurrent Low Back Pain, and Fibromyalgia. Pain
Physician 2017;20:307–318.
[16]   Granovsky Y, Raz N, Defrin R. Electrophysiological and psychophysical correlates of spatial summation to
noxious heat: the possible role of A-delta fibers. Exp Brain Res 2017;235:639–646.
[17]   Greene LC, Hardy JD. Spatial summation of pain. J Appl Physiol 1958;13:457–464.
[18]   Gruener H, Zeilig G, Laufer Y, Blumen N, Defrin R. Differential pain modulation properties in central
neuropathic pain after spinal cord injury. Pain 2016;157:1415–1424.
[19]   Gwilym SE, Keltner JR, Warnaby CE, Carr AJ, Chizh B, Chessell I, Tracey I. Psychophysical and
functional imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis
patients. Arthritis Rheum 2009;61:1226–1234.
[20]   Hardy JD, Wolff HG, Goodell H. Studies on pain. A new method for measuring pain threshold:
Observations on spatial summation of pain. J Clin Invest 1940;19:649– 657.
[21]   Hillman P, Wall PD. Inhibitory and excitatory factors influencing the receptive fields of lamina 5 spinal
cord cells. Exp Brain Res 1969;9:284–306.
[22]   Holbert MD, Pedler A, Camfermann D, Harvie DS. Comparison of spatial summation properties at different
body sites. Scand J Pain 2017;17:126–131.
[23]   Julien N, Marchand S. Endogenous pain inhibitory systems activated by spatial summation are opioid-
mediated. Neurosci Lett 2006;401:256–260.
[24]   Keogh E, Witt G. Hypoalgesic effect of caffeine in normotensive men and women. Psychophysiology
2001;38:886–895.
[25]   Lau BK, Vaughan CW. Descending modulation of pain: the GABA disinhibition hypothesis of analgesia.
Curr Opin Neurobiol 2014;29:159–164.
[26]   Lautenbacher S, Kunz M, Strate P, Nielsen J, Arendt-Nielsen L. Age effects on pain thresholds, temporal
summation and spatial summation of heat and pressure pain. Pain 2005;115:410–418.
[27]   Le Bars D, Dickenson AH, Besson JM. Diffuse noxious inhibitory controls (DNIC). II. Lack of effect on
non-convergent neurones, supraspinal involvement and theoretical implications. Pain 1979;6:305–327.
[28]   Le Bars D, Villanueva L, Bouhassira D, Willer JC. Diffuse noxious inhibitory controls (DNIC) in animals
and in man. Patol Fiziol Eksp Ter 1992:55–65.
[29]   Lei J, Sun T, Lumb BM, You H-J. Roles of the periaqueductal gray in descending facilitatory and inhibitory
controls of intramuscular hypertonic saline induced muscle nociception. Exp Neurol 2014;257:88–94.
[30]   Levy D, Abdian L, Dekel-Steinkeller M, Defrin R. Experimental evidence for weaker endogenous
inhibition of trigeminal pain than extra-trigeminal pain in healthy individuals. Cephalalgia 2018;38:1307–
1315.
[31]   Marchand S, Arsenault P. Spatial summation for pain perception: interaction of inhibitory and excitatory
mechanisms. Pain 2002;95:201–206.
[32]   Martin-Pichora AL, Mankovsky-Arnold TD, Katz J. Implicit versus explicit associative learning and
experimentally induced placebo hypoalgesia. J Pain Res 2011;4:67–77.
[33]   McPhee ME, Vaegter HB, Graven-Nielsen T. Alterations in pronociceptive and antinociceptive
mechanisms in patients with low back pain: a systematic review with meta-analysis. Pain 2020;161:464–
475.
[34]   Morgan MM, Sohn JH, Liebeskind JC. Stimulation of the periaqueductal gray matter inhibits nociception at
the supraspinal as well as spinal level. Brain Res 1989;502:61–66.
[35]   Peirce JW. PsychoPy--Psychophysics software in Python. J Neurosci Methods 2007;162:8–13.
[36]   Price DD, McHaffie JG, Larson MA. Spatial summation of heat-induced pain: influence of stimulus area
and spatial separation of stimuli on perceived pain sensation intensity and unpleasantness. J Neurophysiol
1989;62:1270–1279.
[37]   Price DD, Milling LS, Kirsch I, Duff A, Montgomery GH, Nicholls SS. An analysis of factors that
contribute to the magnitude of placebo analgesia in an experimental paradigm. Pain 1999;83:147–156.
[38]   Quevedo AS, Mørch CD, Andersen OK, Coghill RC. Lateral inhibition during nociceptive processing. Pain
2017;158:1046–1052.
[39]   Rasmussen VM, Ellehuus-Hilmersson C, Rotbøll-Nielsen P, Werner MU. Spatial summation of thermal
stimuli assessed by a standardized, randomized, single-blinded technique. Scand J Pain 2015;9:81–86.

[40] Raz N, Granovsky Y, Defrin R. Investigating the neural processing of spatial summation of pain: the role of
A-delta nociceptors. Exp Brain Res 2015;233:405–413.
[41] Reid E, Harvie D, Miegel R, Spence C, Moseley GL. Spatial summation of pain in humans investigated
using transcutaneous electrical stimulation. J Pain 2015;16:11–18.
[42] Roby-Brami A, Bussel B, Willer JC, Le Bars D. An electrophysiological investigation into the pain-
relieving effects of heterotopic nociceptive stimuli. Probable involvement of a supraspinal loop. Brain
1987;110:1497–1508.
[43] Sikandar S, Ronga I, Iannetti GD, Dickenson AH. Neural coding of nociceptive stimuli-from rat spinal
neurones to human perception. Pain 2013;154:1263–1273.
[44] Smith A, Pedler A. Conditioned pain modulation is affected by occlusion cuff conditioning stimulus
intensity, but not duration. Eur J Pain 2018;22:94–102.
[45] Staud R, Koo E, Robinson ME, Price DD. Spatial summation of mechanically evoked muscle pain and
painful aftersensations in normal subjects and fibromyalgia patients. Pain 2007;130:177–187.
[46] Staud R, Price DD, Robinson ME, Vierck CJ. Body pain area and pain-related negative affect predict
clinical pain intensity in patients with fibromyalgia. J Pain 2004;5:338–343.
[47] Staud R, Vierck CJ, Robinson ME, Price DD. Spatial summation of heat pain within and across dermatomes
in fibromyalgia patients and pain-free subjects. Pain 2004;111:342–350.
[48] Staud R, Weyl EE, Price DD, Robinson ME. Mechanical and heat hyperalgesia highly predict clinical pain
intensity in patients with chronic musculoskeletal pain syndromes. J Pain 2012;13:725–735.
[49] Szikszay TM, Adamczyk WM, Luedtke K. The Magnitude of Offset Analgesia as a Measure of
Endogenous Pain Modulation in Healthy Subjects and Patients with Chronic Pain - A Systematic Review
and Meta-analysis. Clin J Pain 2019; 35:189-204.
[50] Thompson T, Oram C, Correll CU, Tsermentseli S, Stubbs B. Analgesic Effects of Alcohol: A Systematic
Review and Meta-Analysis of Controlled Experimental Studies in Healthy Participants. J Pain
2017;18:499–510.
[51] Torta DM, Churyukanov MV, Plaghki L, Mouraux A. The effect of heterotopic noxious conditioning
stimulation on Aδ-, C- and Aβ-fibre brain responses in humans. Eur J Neurosci 2015;42:2707–2715.
[52] Weissman-Fogel I, Dror A, Defrin R. Temporal and spatial aspects of experimental tonic pain:
Understanding pain adaptation and intensification. Eur J Pain 2015;19:408–418.
[53] Williams AE, Rhudy JL. The influence of conditioned fear on human pain thresholds: does preparedness
play a role? J Pain 2007;8:598–606.
[54] Willis WD, Westlund KN. Neuroanatomy of the pain system and of the pathways that modulate pain. J Clin
Neurophysiol 1997;14:2–31.
551