Research article

How do sex biases in demography and disease characteristics affect the spread of

sexually transmitted infections?

Naerhulan Halimubieke[1*], Alistair Pirrie[2*], Tamás Székely[1,3], Ben Ashby[2]

[1] Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath,

UK

[2] Department of Mathematical Sciences, University of Bath, Bath, UK

[3] Department of Evolutionary Zoology and Human Biology, University of Debrecen, Debrecen,

Hungary

Corresponding author: Naerhulan Halimubieke; hn364@bath.ac.uk

*These authors contributed equally to the manuscript.

Abstract

The epidemiology of sexually transmitted infections (STIs) is inherently linked to host

mating dynamics. Studies across many taxa show that adult sex ratio, a major determinant of

host mating dynamics, is often skewed – sometimes strongly – toward males or females.

However, few predictions exist for the effects of skewed sex ratio on STI epidemiology, and

none when coupled with sex biased disease characteristics. Here we use mathematical

modelling to examine how interactions between sex ratio and disease characteristics affect

STI prevalence in males and females. Notably, we find that while overall disease prevalence

peaks at equal sex ratios, prevalence per sex peaks at skewed sex ratios. Furthermore, disease

characteristics, sex-biased or not, drive predictable differences in male and female STI

prevalence as sex ratio varies, with higher transmission and lower virulence generally

increasing differences between the sexes for a given sex ratio. These findings may be due to a

balance between increased per-capita mating in the less common sex, and a reduction in

mating rate – hence disease prevalence – at the population level. Our work reveals new

insights into how STI prevalence in males and females depends on a complex interaction

between host population demography and disease characteristics.

Keywords: sex ratio, sexual transmission, disease prevalence, STI, mating dynamics

1. Introduction

Sexually transmitted infections (STIs) – defined as any pathogen that is transmitted during copulation – are ubiquitous in the animal kingdom, often causing chronic infections with low recovery rates and reduced reproductive success, even sterility [1,2]. STIs typically exhibit contrasting epidemiological dynamics to non-STIs, with transmission likely to be frequency- rather than density-dependent (i.e. the number of sexual contacts per capita is invariant to population size). Another major difference is that STIs are primarily transmitted between the sexes (especially in non-human populations), which creates a natural bipartite network where the population is split into two disjoint sets (males and females) and connections only exist between individuals from distinct sets. In contrast, non-STI transmission networks are generally not bipartite. Sex ratio, which can vary greatly in non-human populations [3-6], will therefore likely have a profound impact on STI epidemiology compared to non-STI epidemiology, as it disrupts the structure of the contact network [7,8]. There is growing evidence shedding light on the relationship between host mating system structure and STI dynamics [9-11], much of which focuses on human populations [12,13]. However, the implications of variation in sex ratio for STI epidemiology are not well understood.

Evidence has shown that sex ratio has a crucial role in mating system variation in wild populations [14,15]. Sex ratio can vary at different stages of the life cycle, with the ratios of males to females at conception, at birth and during adult life (termed primary, secondary, and tertiary or adult sex ratio, respectively). Recent studies show that adult sex ratio may deviate from 1:1 in a variety of organisms: butterflies, reptiles and birds often have male-biased sex ratios with up to 90% of the population being male, whereas female-biased adult sex ratios are common in insects and mammals, with up to 100 females for every male [3-6]. Variation in the adult sex ratio can alter the mating opportunities of breeding males and females and

may select for different mating systems [16,17], the latter has significant consequences for

STI dynamics [18-21]. Interestingly, *Wolbachia* bacteria (transmitted vertically rather than

sexually in insect hosts) are known to distort the birth sex ratio, as it is maternally inherited

and kills male embryos, consequently, disrupt the mating dynamics of their hosts [7,22,23].

Adult sex ratio is clearly crucial for mating dynamics and hence STI transmission, but it is

shaped by multiple factors, including primary and secondary sex ratio as well as differential

survival during juvenile and adult stages [24]. STIs may cause differential mortality between

the sexes, potentially increasing or decreasing the ratio of adult males to females, causing

feedback between STI transmission and sex ratio [5]. In general, theory predicts that the rarer

sex in the population should exhibit higher STI prevalence since they have a higher per capita

mating rate, all else being equal [8]. For example, two-spot ladybirds *Adalia bipunctata*

exhibit male-biased patterns of STI prevalence in female-skewed populations [8]. A follow-

up experimental study by Pastok et al. [25] established that the presence of male-killing

bacteria at high prevalence skews the sex ratio towards a female-biased population, and

results in male-biased STI prevalence. Together, these studies suggest that there are likely to

be complex interactions between STIs and adult sex ratios. However, the extent to which STI

prevalence in males and females depends on sex ratio, demographic and disease

characteristics, which may vary between the sexes, has yet to be explored.

Many pathogens – both STIs and non-STIs - show sex bias in disease characteristics (e.g.

transmissibility, virulence), which can influence disease dynamics [26,27]. Two main

hypotheses have been proposed to explain sex-biased disease characteristics. The

physiological hypothesis emphasises that the interactions between sex hormones and the

immune system render one sex more susceptible to infection and disease [28]. For example,

in mammals, males have consistently weaker immune competence than females, and this

correlates with male-biased disease prevalence, mortality, and female-biased adult sex ratio

[29,30]. On the other hand, the *behavioural hypothesis* posits that sexual differences in

behaviour may cause sex-specific exposure to pathogens [31]. For instance, infections by

arthropods, helminths and unicellular parasites are often male-biased in mammals but not in

birds [32]. Sex differences are expected, since males tend to be more active in reproductive

behaviours such as combat for females, territorial defence and foraging, due to more intense

sexual selection in males than females, therefore increasing the chance of exposure to

pathogens. In human populations, gender-related behavioural differences may render one sex

more exposed to certain contagions (STIs or non-STIs). For example, Ebola haemorrhagic

fever outbreak that occurred in the Congo and Gabon in 2001-2002 suggested more men than

women were infected during the early stages of the outbreak because men spent more time

working away from home, where they were more likely to come into contact with wild

animals [33]. Regardless of whether sex biases in disease characteristics are due to

physiological or behavioural differences between males and females, there are likely to be

important implications for variation in disease prevalence between the sexes.

In the context of STIs, current empirical research in sex-biased disease characteristics is

mainly focused on humans. Some studies have established that there is a higher risk of

transmission of certain STIs from men to women than vice versa due to STIs being present in

ejaculate [34,35]; another study shows that human T-lymphotropic virus exhibits male-biased

virulence, potentially because transmissions occur during pregnancy, birth or breast-feeding,

and so selection has favoured lower virulence in women [36]. Empirical studies of sex-biased

STI characteristics in animals are limited, but mathematical models have shown how sex-

biased disease characteristics influence transmission dynamics (reviewed in [11], also see

[12]). However, the interplay between sex ratio and disease-characteristics (especially when

these are sex-biased) and their effects on STI prevalence have yet to be investigated. Here we

use mathematical modelling to examine the effects of sex biases in population demography

and disease characteristics on STI epidemiology. Specifically, we investigate how (1)

variation in the sex ratio at maturation; (2) sex-biases in transmission and disease-associated

mortality (virulence) in adults; and (3) the interactions between sex ratio at maturation and

disease characteristics, impact on STI prevalence in males and females.

2. Methods

We model the dynamics of an STI in a randomly mixing population. The epidemiological and

mating dynamics are described by the following system of ordinary differential equations:

117
$$\frac{dS_i}{dt} = \underbrace{b_i(c,q,s)}_{Births} - \underbrace{\beta_i[S_iI_j]}_{Infections} - \underbrace{dS_i}_{Natural\ mortality} + \underbrace{\gamma I_i}_{Recovery} \tag{1}$$

118
$$\frac{dI_i}{dt} = \underbrace{\beta_i[S_iI_j]}_{Infections} - \underbrace{dI_i}_{Natural\ mortality} - \underbrace{\alpha_i I_i}_{Mortality\ virulence} - \underbrace{\gamma I_i}_{Recovery} \tag{2}$$

119   where $i, j \in \{M,F\}$, $i \neq j$, correspond to males and females. The birth rates for each sex are

120
$$b_M(c,q,s) = \frac{bcs(1-qN)}{N}(S_M + I_M)(S_F + I_F), \tag{3}$$

121
$$b_F(c,q,s) = \frac{bc(1-s)(1-qN)}{N}(S_M + I_M)(S_F + I_F), \tag{4}$$

and: S_i and I_i (equivalently, S_j or I_j), are the number of susceptible and infected individuals,

respectively, in each sex; $N = S_M + S_F + I_M + I_F$ is the total population size; s is the sex

ratio at maturation (proportion of maturing individuals that are male), with $0 \leq s \leq 1$; $c \geq 0$ is

the pairwise rate at which males and females mate given frequency-dependent contact (see

equations 5-8); q represents the strength of density-dependence on births; d is the natural

mortality rate; α_i is the disease-associated mortality rate (virulence) in sex i; β_i is the

transmission rate from sex j to sex i per sexual contact; γ is the recovery rate; and $[X_iY_j]$ is

the mating rate between individuals in classes $X,Y \in \{S,I\}$.

For simplicity, we assume that: (i) both sexes have the same recovery rate from the STI; (ii)

both sexes have the same natural mortality rate; and (iii) the per-capita mating rate is

frequency-dependent, which means that larger populations do not have a higher per-capita

mating rate than smaller populations. We define the mating rates for each combination of the

population classes by:

$$[S_MS_F] = \frac{c\,S_M\,S_F}{N} \tag{5}$$

$$[S_MI_F] = \frac{c\,S_M\,I_F}{N} \tag{6}$$

$$[S_FI_M] = \frac{c\,S_F\,I_M}{N} \tag{7}$$

$$[I_MI_F] = \frac{c\,I_M\,I_F}{N} \tag{8}$$

140   The disease-free equilibrium $(S_M, S_F, I_M, I_F) = (S_M^*, S_F^*, 0, 0)$, of this system occurs at:

$$S_M^* = \frac{bcs(1-s) - d}{bcq(1-s)} \tag{9}$$

$$S_F^* = \frac{bcs(1-s) - d}{bcqs} \tag{10}$$

143   A newly introduced STI will spread in a susceptible population when the basic reproductive

144   ratio, R_0 is greater than 1, where:

$$R_0 = \frac{c}{S_F^* + S_M^*} \cdot \sqrt{\frac{\beta_M \beta_F S_M^* S_F^*}{(d+\gamma+\alpha_M)(d+\gamma+\alpha_F)}} \tag{11}$$

In order to guarantee that $R_0 > 1$, we require that $\frac{1}{2} - \psi < s < \frac{1}{2} + \psi$, where $\psi =$

$\sqrt{\frac{1}{4} - \frac{(d+\gamma+\alpha_M+\alpha_F)(d+\gamma)}{\beta_M \beta_F c^2}}$. To test the interactions between sex ratio and STI prevalence in

males and females, we start with no sex bias in STI characteristics. This means that males

and females are equally susceptible to the pathogen and the mortality caused by the pathogen

is same in both sexes ($\alpha_M=\alpha_F=\alpha$; $\beta_M=\beta_F=\beta$). STI prevalence in each sex is given as the

number of infected individuals in that sex divided by the total number of individuals in that

sex (at equilibrium, as indicated by asterisks):

$$D_M = \frac{I_M^*}{S_M^* + I_M^*} \tag{12}$$

$$D_F = \frac{I_F^*}{S_F^* + I_F^*} \tag{13}$$

To examine the impact of sex-biased disease characteristics on the prevalence of STIs in

males and females, we fix female susceptibility (β_F) and disease-related mortality (α_F) and

vary the corresponding parameters in males. The model is symmetric up to labelling, so the

results are analogous for the converse scenario. We primarily use numerical analysis to

determine disease prevalence as the endemic equilibrium of equations 1-2 is mostly

intractable to algebraic analysis.

3. Results

3.1 Non-sex-biased disease characteristics

We begin by exploring how sex ratio impacts on STI prevalence in males and females when

there are no sex-biases in disease characteristics (i.e. $\alpha_M=\alpha_F=\alpha$; $\beta_M=\beta_F=\beta$). Consistent with

previous theoretical and empirical work [7,8], the less common sex exhibits higher STI

prevalence at equilibrium (Fig. 1). Intuitively, when there are no sex biases in disease

characteristics, the overall prevalence of the STI is always maximised at an equal sex ratio

($s = 0.5$). However, both the extent to which STI prevalence differs between the sexes (i.e.

the difference between the male and female curves in Fig. 1a-b, shown in Fig. 1c-d) and the

point at which STI prevalence peaks in each sex (i.e. the sex ratio that maximises the male

and female curves, indicated by circles in Fig.1a-b) depends on the transmission rate and

virulence. Specifically, as the transmission rate increases or as the virulence decreases, the

difference between male and female STI prevalence for a given sex ratio tends to increase,

especially for more extreme sex ratios (Fig. 1c-d). Furthermore, for higher transmission

rates/lower virulence, the sex ratio at maturation for which disease prevalence peaks becomes

more extreme. Thus, while overall STI prevalence is qualitatively unchanged by variation in

the transmission rate and virulence (i.e. it always peaks at $s = 0.5$), STI prevalence per sex is

highly sensitive to disease characteristics.

Figure 1. Effects of sex ratio at maturation (proportion male), s, on STI prevalence when there are no sex biases

in disease characteristics: (a, c) variation in disease-related mortality, $\alpha_M = \alpha_F = \alpha$; (b, d) variation in

transmission probability, $\beta_M = \beta_F = \beta$. As disease-related mortality increases or the transmission probability

decreases, overall STI prevalence decreases, and the difference between male and female STI prevalence for a

given sex ratio decreases. (a, b) STI prevalence in males (solid line), females (dashed line), and overall (dotted

line). Markers show where STI prevalence peaks in males (black) and females (white). (c, d) Absolute

difference in STI prevalence between males (D_M) and females (D_F), corresponding to the absolute difference

between the solid and dashed lines in (a) & (b), with markers showing where the absolute difference peaks.

Other parameters are held constant throughout to ensure R_0 is greater than 1: $\delta = 0.1$; $\gamma = 0.2$; $c = 5$; $b = 1$;

$q = 0.001$.

We can see this analytically for variation in the transmission rate if we make the following

simplifying assumptions: (1) there is no disease-associated mortality ($\alpha_M = \alpha_F = 0$); (2) the

disease transmission rates are equal ($\beta_M = \beta_F = \beta$); and (3) there is no recovery ($\gamma = 0$).

These assumptions ensure that the adult sex ratio is equal to the sex ratio at maturation, which

greatly simplifies the analysis. First, we must calculate the equilibrium population size, N^*.

Let $M = S_M + I_M$ be the number of males in the population and $F = S_F + I_F$, be the number

of females in the population. Then at equilibrium, $M = sN^*$, and $F = (1 - s)N^*$. Now, $\frac{dN}{dt} =$

$\frac{dS_M}{dt} + \frac{dS_F}{dt} + \frac{dI_M}{dt} + \frac{dI_F}{dt}$, and evaluating this gives

$$\frac{dN}{dt} = \frac{bc(1 - qN)MF}{N} - dN \tag{14}$$

200   Substituting in $M = sN^*$ and $F = (1 - s)N^*$, and solving $\frac{dN}{dt} = 0$, the resulting (non-trivial)

201   equilibrium population size is:

$$N^* = \frac{bcs(1 - s) - d}{bcqs(1 - s)} \tag{15}$$

203   We can now substitute the equilibrium population size into the full system of equations,

204   giving the endemic equilibrium:

$$S_F^* = \frac{d(\beta c(1 - s) + d)\zeta}{\beta bc^2 qs(1 - s)} \tag{16}$$

$$S_M^* = \frac{S_F^*}{\zeta^2} \tag{17}$$

$$I_F^* = \frac{(\beta^2 c^2 s(1 - s) - d^2)\zeta}{\beta bc^2 qs(1 - s)} \tag{18}$$

$$I_M^* = \zeta I_F^* \tag{19}$$

209   where $\zeta = \frac{\beta c(1-s)-d}{\beta cs+d}$. The disease prevalence in males and females is therefore

$$D_M = \frac{I_M^*}{S_M^* + I_M^*} = \frac{\beta^2 c^2 s(1 - s) - d^2}{\beta cs(\beta c(1 - s) + d)} \tag{20}$$

211
$$D_F = \frac{I_F^*}{S_F^* + I_F^*} = \frac{\beta^2 c^2 s(1-s) - d^2}{\beta c(1-s)(\beta cs + d)} \qquad (21)$$

212   To find the peak in disease prevalence with respect to s, one must differentiate equations 20-

213   21 with respect to s and then find where they are equal to zero, i.e., find s_M^*, s_F^* such that

214     $\frac{d}{ds}(D_F)|_{s=s^*} = 0$ and $\frac{d}{ds}(D_M)|_{s=s^*} = 0$. For females, STI prevalence peaks at

215
$$s_F^* = \frac{\beta c + d - \sqrt{\beta cd + 2d^2}}{\beta c} = 1 + \frac{d - \sqrt{\beta cd + 2d^2}}{\beta c} \qquad (22)$$

216   and for males STI prevalence peaks at

217
$$s_M^* = \frac{-d + \sqrt{\beta cd + 2d^2}}{\beta c} \qquad (23)$$

(Note that the other root lies outside $s \in [0,1]$). Taking derivatives with respect to β, we find

that $\frac{ds_F^*}{d\beta} > 0$ and $\frac{ds_M^*}{d\beta} < 0$. Thus, as the transmission rate, β, increases, the sex ratio at which

STI prevalence peaks in each sex always becomes more extreme.

3.2 Sex-biased disease characteristics

We now consider the effects of sex-biased disease characteristics on STI prevalence.

Specifically, we investigate how STI prevalence (per sex and overall) varies when there are

biases in (1) female-to-male (β_M) or male-to-female (β_F) transmission, and/or (2) disease-

associated mortality (virulence) for males (α_M) or females (α_F). Since the only differences in

the model between males and females are these parameters, without loss of generality we fix

β_F and α_F whereas vary β_M and α_M accordingly.

Sex differences in transmission rate and in virulence influence disease prevalence in both

sexes, although the extent of their influence on males and females depends on sex ratio. A

higher female-to-male transmission rate relative to the male-to-female transmission rate

raises disease prevalence in both sexes, although interestingly the influence of sex ratio

differs between the sexes. With greater female-to-male transmission, the disease prevalence

peak shifts toward male-biased sex ratios in females whereas for males the change is

marginal (e.g. Fig. 2a-c). Similarly, lower disease-associated mortality in males relative to

females also universally increases disease prevalence, but leads to differential effects on

males and females with respect to sex ratio: the disease prevalence peak shifts toward female-

biased sex ratios in males whereas for males the disease prevalence peaks at more equal sex

ratios (Fig. 2b, e, h). These patterns are broadly consistent as we vary the degree of sex

biased transmission or virulence, with the exception that very high mortality virulence in

males relative to females cause the disease prevalence curves to peak at more equal sex ratios

(Fig. 3c-f).

Figure 2. STI prevalence in males (solid line), females (dashed line), and overall (dotted line) as a function of

the sex ratio at maturation (proportion male), s, when there are sex biases in STI transmission rate and

virulence. The parameters $\beta_F = 0.5$ and $\alpha_F = 0.3$ are held constant throughout, while $\beta_M \in \{0.2, 0.5, 1\}$ and $\alpha_F \in$

$\{0, 0.3, 0.6\}$ vary between panels. Other parameters as described in Fig. 1.

Figure 3. STI prevalence in (a, d) males, (b, e) females, and (c, f) overall as a function of the sex ratio at

maturation and: (a-c) female-to-male transmission rate (β_M); (d-f) mortality virulence in males (α_M). The red

lines show when the sex ratio at maturation is equal (solid) and when there is no sex bias in disease

characteristics (dashed): (a-c) $\beta_M = \beta_F = 0.5$; (d-f) $\alpha_M = \alpha_F = 0.3$. The white region in each panel corresponds to

conditions where the STI is unable to persist ($R_0 < 1$). The parameters $\beta_F = 0.5$ and $\alpha_F = 0.3$ are held constant

throughout, Other parameters as described in Fig. 1.

We see consistent patterns when transmission and virulence are varied simultaneously

(Fig. 4). By measuring the sex ratio at maturation, s, for which disease prevalence peaks, we

analyse how the interaction between transmission, virulence, and sex ratio leads to changes in

disease prevalence. When disease-associated mortality in males is relatively low ($\alpha_M < \alpha_F$),

variation in the female-to-male transmission rate (β_M) has little impact on the sex ratio where

disease prevalence peaks in males (Fig. 4a) and overall (Fig. 4c), but the effect in females is

much more pronounced, with the skew generally increasing for greater β_M (Fig. 4b). When

disease-associated mortality in males is relatively high ($\alpha_M > \alpha_F$), the sex ratio at maturation

where disease prevalence peaks in males and overall also increases with β_M, but to a lesser

extent than in females. It is also clear from the contours in Fig. 4 that variation in either

transmission or virulence can cause non-monotonic changes in the disease prevalence curves.

For example, the horizontal and vertical red lines in Fig. 4 (indicating $\beta_M = \beta_F$ or $\alpha_M = \alpha_F$)

intersect some of the contours twice, which means that varying either α_M or β_M can cause a

non-monotonic change in the skew of the disease prevalence curves. Together, these results

show that interactions between sex-biased disease characteristics and sex ratio can lead to

major changes in STI prevalence within each sex and overall, and that the differences

between male and female STI prevalence can be exacerbated by greater transmission rates or

lower mortality rates, in one or both sexes.

Figure 4. The sex ratio at maturation for which disease prevalence peaks in: (a) males, (b) females, and (c)

overall, as a function of disease-associated mortality for males (α_M) and female-to-male transmission rate (β_M).

The red lines show when there is no sex bias in transmission ($\beta_M=\beta_F=0.5$; dashed) and when there is no sex bias

in virulence ($\alpha_M=\alpha_F=0.3$; solid). The white region in the bottom right corner of each panel corresponds to

conditions where the STI is unable to persist ($R_0 < 1$). Other parameters as described in Fig. 1.

4. Discussion

Studies of STI epidemiology typically emphasise the importance of host mating system and population demography influencing STI dynamics [8-11,20,25]. Here we used mathematical modelling to examine the interaction between sex-biased population demography and disease-characteristics, and their effects on STI prevalence in males and females. At a fundamental level, our results demonstrate that the demographic conditions which lead to high STI prevalence are likely to be different between the sexes. Moreover, disease-characteristics, whether they be sex-biased or not, can increase or decrease differences in male and female STI prevalence. Further to this, our model shows there are interactions between disease characteristics and sex ratio that might change STI prevalence within each sex and overall.

In a population with a skewed adult sex ratio, the less common sex has more prospective mates and a higher per-capita mating rate than the more common sex. This pattern can be seen in human and non-human societies; for instance, in human communities with a male-skewed adult sex ratio, men are more likely to purchase sex and women are more likely to have multiple sex partners [37,38]. Consistently, in bird populations mating opportunities are related to adult sex ratios: in male-biased adult sex ratio, females have higher mating opportunities than males, whereas in female-skewed populations males re-mate faster than the females [39-41]. Hence, with high variance in mating success, many individuals of the more common sex may remain unmated or have a low mating rate, whereas the less common sex has a higher mating rate and hence greater exposure to infectious partners. Thus, equilibrium disease prevalence in the more common sex is expected to be lower than in the rarer sex, consistent with previous theoretical and empirical studies [7,8,10,11]. Here we

advance this research field by showing that the interactions between sex ratios and disease

characteristics may lead to complex patterns in disease prevalence.

Our model shows how STI prevalence – overall and in each sex – depends on both sex ratio

and disease characteristics. We have shown that even when there are no sex biases in disease

characteristics, increasing the transmission rate or decreasing mortality virulence generally

leads to a greater difference in STI prevalence between the sexes for a given sex ratio (Fig.

1c-d). We have also shown that the sex ratio for which disease prevalence peaks in each sex

depends on disease characteristics, again even when the disease itself shows no sex biases in

transmission or virulence. The reason for these patterns can be understood in terms of a

balance between an increase in the per-capita mating rate for the less common sex and a

decrease in overall disease prevalence as the sex ratio becomes more skewed. Suppose the

adult sex ratio is initially equal and that there are no biases in disease characteristics, hence

equilibrium STI prevalence is the same in both sexes. If we gradually remove males from the

population (similarly for females), then the per-capita mating rate for the remaining males

increases, and hence so does their risk of infection, at least initially. However, skewing the

sex ratio in either direction reduces the overall mating rate, and therefore always lowers

disease prevalence at the population level. As one continues to remove males from the

population, overall disease prevalence begins to fall more sharply because uninfected males

are increasingly less likely to mate with infected females. Thus, disease prevalence

eventually begins to fall in males as well. Once the population crosses a critical sex ratio, R_0

falls below 1 and the disease is driven extinct. Higher transmission and lower virulence both

increase overall disease prevalence and therefore buffer against the effects of skewed sex

ratios, which shifts the sex ratio at which disease prevalence peaks in each sex to more

extreme values. This can be seen in Fig. 1a-b, where the gradient of the overall disease

prevalence curves is flatter for higher values of β and lower values of α.

We also explored the effects of sex-biased transmission and virulence. Sex-biased disease

characteristics are common in wild populations as well as in human populations [31,42-44],

and previous studies have looked into the behavioural and physiological causes of such

disparities and their implications for population dynamics [27,28,31]. However, the

interaction between these biases and STI prevalence within each sex have previously been

overlooked. Our model suggests that sex-biased disease characteristics may independently

influence STI prevalence in a similar way as the unbiased transmission rate or virulence (see

above). However, the prevalence curves for each sex are no longer symmetrical when there

are biased STI characteristics (Fig. 2). Our model also reveals that interactions between sex

ratio and sex-biased disease characteristics can lead to differential and non-intuitive patterns

for STI prevalence. For example, while increasing the bias in transmission from females to

males causes STI prevalence in females to peak at a more extreme sex ratio, there are only

marginal effects on males, yet the effects of male-biased virulence on disease prevalence are

much more pronounced in males than females (as shown in Fig. 3). The model also predicts

that increasing or decreasing sex-biased disease characteristics may lead to non-monotonic

variation in STI prevalence. In the absence of mathematical modelling, it is unlikely that one

would be able to intuit such effects, but our results suggest that there are likely to be complex

interactions between sex-biased disease characteristics and population demography that

shapes STI epidemiological dynamics.

Our model is simple and makes few biological assumptions. Therefore, follow-up analyses

are necessary to investigate biological factors mediating STI prevalence that are not captured

in our model. For example, our model assumes no variance in mating rate, and hosts do not

form pair-bonds [46-49]. When one sex experiences higher variance in mating rate, the

mating population is effectively sex- skewed and so we would expect this to generally have a

similar effect on STI prevalence as seen in our model. If infection status impacts on mating

rate [45, 48-50], then the effects may be more complex, especially if mate choice is driven by

one sex. Previous models have explored how STIs might affect the evolution of host mating

dynamics [46-50], but the focus of the present study has been on STI prevalence. However,

given that there are interactions between sex ratio, sex-biased disease characteristics and

mating dynamics, there are likely to be evolutionary implications for these factors on hosts

and STIs, which should be the target of future research.

Our results have several implications for STIs in natural populations. First, when sex ratios

are skewed, we should expect STIs to be more prevalent in the rarer sex, although the relative

prevalence of co-circulating STIs in males and females will differ due to variation in

transmission and virulence (Fig. 1c-d). For example, in a female- skewed population, an STI

with a high transmission rate is likely to have a higher prevalence in males relative to females

compared to an STI with a low transmission rate. Second, our model predicts that variance in

the sex ratio between populations to drive STI prevalence in males and females, with

divergence from equal sex ratios generally increasing sex differences in prevalence (except at

extreme sex ratios). Finally, we should expect the interaction between sex ratio and sex-

biased transmission or virulence to consistently have a stronger effect on STI prevalence in

one sex than in the other.

5. Conclusion

To conclude, the objective of our model was to examine the interplay between sex ratio and disease characteristics and explore their effects on STI prevalence. Our key message is that while the less common sex is predicted to exhibit higher STI prevalence at equilibrium, disease characteristics such as transmission rate and virulence – sex-biased or not – combined with sex ratio, drive differential patterns in male and female STI prevalence.

Funding

This work is supported by Chinese Scholarship Council and the National Research, Development and Innovation Office of Hungary (grant no. ÉLVONAL KKP-126949, K-116310 to NH). University Research Scholarship Award from the University of Bath (to AP). Royal Society Wolfson Merit Award WM170050, APEX APX\R1\191045 and Leverhulme Trust (RF/ 2/ RFG/ 2005/ 0279, ID200660763 to TS). The Natural Environment Research Council (grant no. NE/N014979/1 to BA).

Author contributions

BA, NH and AP designed and analysed the model. NH and AP co-wrote the first draft, and all authors contributed to revisions of the manuscript.

Data availability

Source code is available in the Supplemental Material and in the following Github repository: https://github.com/ecoevogroup/Halimubieke_et_al_2020.

Competing interests

The authors declare that they have no competing interests.

References

1. Ashby B, Jordan JE, Knell RJ, Hurst GDD. 2019 Sexually transmitted infections in natural populations: what have we learnt from beetles and beyond? in Wilson PK, Fenton A, and Tompkins A (Eds.), *Wildlife disease ecology: linking theory to data and application (*pp. 187-222*)*. Cambridge, UK: Cambridge University Press.

2. Lockhart AB, Thrall PH, Antonovics J. 1996 Sexually transmitted diseases in animals: Ecological and evolutionary implications. *Biol. Rev. Camb. Philos. Soc.* **71**, 415-471.

3. Dyson EA, Hurst GD. 2004 Persistence of an extreme sex-ratio bias in a natural population. *Proc. Natl. Acad. Sci. USA* **101**, 6520-6523. (doi:10.1073/pnas.0304068101)

4. Liker A, Freckleton RPF, Székely T. 2013 The evolution of sex roles in birds is related to adult sex ratio. *Nat. Comm.* **4**, 1587.

5. Székely T, Weissing FJ, Komdeur J. 2014 Adult sex ratio variation: implications for breeding system evolution. *J. Evol. Biol.* **27**, 1500-1512. (doi: 10.1111/jeb.12415)

6. Bonnet X *et al*. 2016 A prison effect in a wild population: a scarcity of females induces homosexual behaviors in males. *Behav. Ecol.* **27**, 1206-1215.

7. Hurst GDD, Hurst LD, Majerus MEN. 1997 Cytoplasmic sex ratio distorters. In O'Neill SL, Hoffmann AA, Werren JH (Eds.), *Influential passengers: microbes and invertebrate reproduction* (pp. 125e154). Oxford, UK: Oxford University Press.

8. Ryder JJ *et al*. 2014 Disease epidemiology in arthropods is altered by the presence of nonprotective symbionts. *Am. Nat.* **183**, E89eE104. (http://dx.doi.org/ 10.1086/674827)

9. Thrall PH, Antonovics J, Bever JD 1997 Sexual transmission of disease and host mating systems: within-season reproductive success. *Am. Nat.* **149**, 485e506.

10. Thrall PH, Antonovics J, Dobson AP. 2000 Sexually transmitted disease in polygynous mating systems: prevalence and impact on reproductive success. *Proc. R. Soc. Lond. B* **267**, 1555e1563.

11. Ashby B, Gupta S. 2013 Sexually transmitted infections in polygamous mating systems. *Philos. Trans. R. Soc. B* **368**, 20120048. (doi: 10.1098/rstb.2012.0048)

12. Anderson RM, May RM. 1992 *Infectious diseases of humans*. Oxford, UK: Oxford University
Press.

13. Eames KTD, Keeling MJ. 2002 Modelling dynamic and network heterogeneities in the spread
of sexually transmitted diseases. *Proc. Natl. Acad. Sci.* **99**, 13330-13335. (doi:
10.1073/pnas.202244299)

14. Székely T, Weissing FJ. Komdeur J. 2014 Adult sex ratio variation: implications for breeding
system evolution. *J. Evol. Biol.* **27**, 1500-1512. (doi: 10.1111/jeb.12415)

15. Eberhart-Phillips LJ *et al*. 2017 Sex-specific early survival drives adult sex ratio bias in
snowy plovers and impacts mating system and population growth. *Proc. Natl. Acad. Sci. USA*
114, E5474-E5481. (doi: 10.1073/pnas.1620043114)

16. Emlen ST, Oring LW. 1977 Ecology, sexual selection, and the evolution of animal mating
systems. *Science* **197**, 215-223.

17. Rankin DJ, Kokko H. 2007 Do males matter? The role of males in population dynamics.
Oikos **116**, 335-348. (doi:10.1111/j.0030-1299.2007.15451.x)

18. Anderson RM, Blythe SP, Gupta S, Konings E. 1989 The transmission dynamics of the
human immunodeficiency virus type 1 in the male homosexual community in the United
Kingdom: the influence of changes in sexual behaviour. *Phil. Trans. R. Soc. Lond. B* **325**, 45-
98.

19. Anderson RM. 1991 The transmission dynamics of sexually transmitted diseases: the
behavioral component. In Wasserheit JN, Aral SO, Holmes KK.(ed.) *Research issues in*
human behavior and sexually transmitted diseases in the AIDS era, 38-60. Washington DC:
American Society for Microbiology.

20. Thrall PH, Antonovics J. 1997 Polymorphism in sexual versus non-sexual disease
transmission. *Proc. R. Soc. Lond. B* **264**, 581-587.

21. Thrall PH, Antonovics J, Wilson WG. 1998 Allocation to sexual vs. nonsexual disease
transmission. *Am. Nat.* **151**, 29-45.

22. Jiggins FM, Hurst GDD, Majerus MEN. 2000 Sex ratio distorting Wolbachia cause sex role
reversal in their butterfly hosts. *Proc. R. Soc. Lond. B* **267**, 69-73.

23. Charlat S *et al*. 2007 Male-Killing Bacteria Trigger a Cycle of Increasing Male Fatigue and

458   Female Promiscuity. *Curr. Biol.* **17**, 273-277. (doi: 10.1016/j.cub.2006.11.068)

459   24. Ancona S, Dénes FV, Krüger O, Székely T, Beissinger SR. 2017 Estimating adult sex ratios

460   in nature. *Phil. Trans. R. Soc. B* **372**, 20160313. (doi: 10.1098/rstb.2016.0313)

461   25. Pastok D. 2015 Causes of Spatial Variation in Parasite and Pathogen Pressure In insects. PhD

462   thesis, University of Liverpool.

463   26. Sheridan LAD, Poulin R, Ward DF, Zuk M. 2000 Sex differences in parasitic infections

464   among arthropod hosts: is there a male bias? *Oikos* **88**, 327–334.

465   27. Miller MR, White A, Wilson K, Boots M. 2007 The Population Dynamical Implications of

466   Male-Biased Parasitism in Different Mating Systems. *PLoS One* **2**, e624.

467   (doi:10.1371/journal.pone.0000624)

468   28. Zuk M, McKean KA. 1996 Sex differences in parasite infections: patterns and processes. *Int.*

469   *J. Parasitol.* **26**, 1009-1024.

470   29. Klein SL, Roberts CW. 2010 *Sex Hormones and Immunity to Infection*. Berlin: Springer

471   Verlag.

472   30. Lemaître JF *et al.* 2020 Sex differences in longevity and aging rates across wild mammals.

473   Proc. *Natl. Acad. Sci. USA* **117**, 8546-8553.

474   31. Brei B, Fish D. 2003 Comment on "Parasites as a viability cost of sexual selection in natural

475   populations of mammals". *Science* **300**, 55a.

476   32. Moore SL, Wilson K. 2002 Parasites as a viability cost of sexual selection in natural

477   populations of mammals. *Science* **297**, 2015-2018.

478   33. World Health Organization. 2007 Addressing Sex and Gender in Epidemic-Prone Infectious

479   Diseases. Geneva, Switzerland.

480   34. Coombs RW, Reichelderfer PS, Landay AL. 2003 Recent observations on HIV type-1

481   infection in the genital tract of men and women. *AIDS* **17**, 455-80.

482   35. Panchanadeswaran S *et al*. 2006 Gender differences in the prevalence of sexually transmitted

483   infections and genital symptoms in an urban setting in southern India. *Sex. Transm. Infect.* **82**,

484   491-495. (doi:10.1136/sti.2006.020768)

36. Úbeda F, Jansen VA. 2016 The evolution of sex-specific virulence in infectious diseases. *Nat.*

Comm. **7**, 13849. (doi:10.1038/ncomms13849)

37. Tucker JD *et al*. 2005 Surplus men, sex work, and the spread of HIV in China. *AIDS*. **19**, 539-

547. (doi: 10.1097/01.aids.0000163929.84154.87)

38. Bien CH, Cai Y, Emch ME, Parish W, Tucker JD. 2013 High adult sex ratios and risky sexual

behaviors: a systematic review. *PLoS One* **8**, e71580. (doi: 10.1371/journal.pone.0071580)

39. Székely T, Cuthill IC, Kis J. 1999 Brood desertion in Kentish plover sex differences in

remating opportunities. *Behav. Ecol.* **10**, 185-190. (doi: 10.1093/beheco/10.2.185)

40. Parra JE, Beltrán M, Zefania S, dos Remedios N, Székely T. 2014 Experimental assessment

of mating opportunities in three shorebird species. *Anim. Behav.* **90**, 83-90. (doi:

10.1016/j.anbehav.2013.12.030).

41. Carmona-Isunza MC, Küpper C, Serrano-Meneses MA, Székely T. 2015 Courtship behavior

differs between monogamous and polygamous plovers. *Behav. Ecol. Sociobiol.* **69**, 2035-

2042. (doi: 10.1007/s00265-015-2014-x)

42. Poulin R. 1996 Sexual inequalities in helminth infections: a cost of being a male? *Am. Nat.*

147, 287–295.

43. Schalk G, Forbes MR. 1997 Male biases in parasitism of mammals: effects of study type, host

age, and parasite taxon. *Oikos* **78**, 67-74.

44. Wilson K, Moore SL, Owens IPF. 2003 Response to comment on: ''Parasites as a viability

cost of sexual selection in natural populations of mammals''. *Science* **300**, 55B-55B.

45. Knell RJ. 1999 Sexually transmitted disease and parasite-mediated sexual selection. *Evolution*

53, 957-961.

46. Kokko H, Ranta E, Ruxton G, Lundberg P. 2002 Sexually transmitted disease and the

evolution of mating systems. *Evolution* **56**, 1091-1100.

47. Boots M, Knell RJ. 2002 The evolution of risky behaviour in the presence of a sexually

transmitted disease. *Proc. R. Soc. Lond. B* **269**, 585-589. (doi: 10.1098/rspb.2001.1932)

48. Ashby B, Boots M. 2015 Coevolution of parasite virulence and host mating strategies. *Proc.*

Natl. Acad. Sci. USA **112**, 13290-13295.

49. Ashby B. 2020 Antagonistic coevolution between hosts and sexually transmitted

infections. *Evolution* *74*, 43-56. (doi: 10.1111/evo.13883)

50. Loehle C. 1997 The adaptive significance of trunk inclination on slopes: a commentary.

Philos. Trans. R. Soc. B **264**, 1371-1374.