Research article

How do biases in sex ratio and disease characteristics affect the spread of sexually

transmitted infections?

Naerhulan Halimubieke[1*], Alistair Pirrie[2*], Tamás Székely[1,3], Ben Ashby[2]

[1] Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath,

UK

[2] Department of Mathematical Sciences, University of Bath, Bath, UK

[3] Department of Evolutionary Zoology and Human Biology, University of Debrecen, Debrecen,

Hungary

Corresponding author: Naerhulan Halimubieke ; hn364@bath.ac.uk

*These authors contributed equally to the manuscript.

Abstract

The epidemiology of sexually transmitted infections (STIs) is inherently linked to host

mating dynamics. Studies across many taxa show that adult sex ratio, a major determinant of

host mating dynamics, is often skewed - sometimes strongly - toward males or females.

However, few predictions exist for the effects of skewed sex ratio on STI epidemiology, and

none when coupled with sex biased disease characteristics. Here we use mathematical

modelling to examine how interactions between sex ratio and disease characteristics affect

STI prevalence in males and females. Notably, we find that while overall disease prevalence

peaks at equal sex ratios, prevalence per sex peaks at skewed sex ratios. Furthermore, disease

characteristics, sex-biased or not, drive predictable differences in male and female STI

prevalence as sex ratio varies, with higher transmission and lower virulence generally

increasing differences between the sexes for a given sex ratio. These findings may be due to a

balance between increased per-capita mating in the less common sex, and a reduction in

mating rate - hence disease prevalence - at the population level. Our work reveals new

insights into how STI prevalence in males and females depends on a complex interaction

between host population sex ratio and disease characteristics.

Keywords: sex ratio, sexual transmission, disease prevalence, STI, mating dynamics

1. Introduction

Sexually transmitted infections (STIs) – defined as any pathogen that is transmitted during copulation – are ubiquitous in the animal kingdom, often causing chronic infections with low recovery rates and reduced reproductive success, even sterility [1, 2]. STIs typically exhibit contrasting epidemiological dynamics to non-STIs, with transmission likely to be frequency- rather than density-dependent (i.e. the number of sexual contacts per capita is invariant to population size). Another major difference is that STIs are primarily transmitted between the sexes (especially in non-human populations), which creates a natural bipartite network where the population is split into two disjoint sets (males and females) and connections only exist between individuals from distinct sets. In contrast, non-STI transmission networks are generally not bipartite. Sex ratio, which can vary greatly in non-human populations [3-6], will therefore likely have a profound impact on STI epidemiology compared to non-STI epidemiology, as it disrupts the structure of the contact network [7, 8]. There is growing evidence shedding light on the relationship between host mating system structure and STI dynamics [9-11], much of which focuses on human populations [12, 13]. However, the implications of variation in sex ratio for STI epidemiology are not well understood.

Evidence has shown that sex ratio has a crucial role in mating system variation in wild populations [5, 14]. Sex ratio can vary at different stages of the life cycle, with the ratios of males to females at conception, at birth and during adult life (termed primary, secondary, and tertiary or adult sex ratio, respectively). Recent studies show that adult sex ratio may deviate from 1:1 in a variety of organisms: butterflies, reptiles and birds often have male-biased sex ratios with up to 90% of the population being male, whereas female-biased adult sex ratios are common in insects and mammals, with up to 100 females for every male [3-6]. Variation in the adult sex ratio can alter the mating opportunities of breeding males and females and

may select for different mating systems [15, 16], the latter has significant consequences for

STI dynamics [17-20]. Interestingly, *Wolbachia* bacteria (transmitted vertically rather than

sexually in insect hosts) are known to distort the birth sex ratio, as it is maternally inherited

and kills male embryos, consequently, disrupt the mating dynamics of their hosts [7, 21, 22].

Adult sex ratio is clearly crucial for mating dynamics and hence STI transmission, but it is

shaped by multiple factors, including primary and secondary sex ratio as well as differential

survival during juvenile and adult stages [23]. STIs may cause differential mortality between

the sexes, potentially increasing or decreasing the ratio of adult males to females, causing

feedback between STI transmission and sex ratio [5]. In general, theory predicts that the rarer

sex in the population should exhibit higher STI prevalence since they have a higher per capita

mating rate, all else being equal [8]. For example, two-spot ladybirds *Adalia bipunctata*

exhibit male-biased patterns of STI prevalence in female-skewed populations [8]. A follow-

up experimental study by Pastok et al. [24] established that the presence of male-killing

bacteria at high prevalence skews the sex ratio towards a female-biased population, and

results in male-biased STI prevalence. Together, these studies suggest that there are likely to

be complex interactions between STIs and adult sex ratios. However, the extent to which STI

prevalence in males and females depends on sex ratio and disease characteristics, which may

vary between the sexes, has yet to be explored.

Many pathogens – both STIs and non-STIs - show sex bias in disease characteristics (e.g.

transmissibility, virulence), which can influence disease dynamics [25, 26]. Two main

hypotheses have been proposed to explain sex-biased disease characteristics. The

physiological hypothesis emphasises that the interactions between sex hormones and the

immune system render one sex more susceptible to infection and disease [27]. For example,

in mammals, males have consistently weaker immune competence than females, and this

correlates with male-biased disease prevalence, mortality, and female-biased adult sex ratio

[28-30] On the other hand, the *behavioural hypothesis* posits that sexual differences in

behaviour may cause sex-specific exposure to pathogens [31]. For instance, infections by

arthropods, helminths and unicellular parasites are often male-biased in mammals but not in

birds [32]. Sex differences are expected, since males tend to be more active in reproductive

behaviours such as combat for females, territorial defence and foraging, due to more intense

sexual selection in males than females, therefore increasing the chance of exposure to

pathogens. In human populations, gender-related behavioural differences may render one sex

more exposed to certain contagions (STIs or non-STIs). For example, Ebola haemorrhagic

fever outbreak that occurred in the Congo and Gabon in 2001-2002 suggested more men than

women were infected during the early stages of the outbreak because men spent more time

working away from home, where they were more likely to come into contact with wild

animals [33]. Regardless of whether sex biases in disease characteristics are due to

physiological or behavioural differences between males and females, there are likely to be

important implications for variation in disease prevalence between the sexes.

In the context of STIs, current empirical research in sex-biased disease characteristics is

mainly focused on humans. Some studies have established that there is a higher risk of

transmission of certain STIs from men to women than vice versa due to STIs being present in

ejaculate [34, 35]; another study shows that human T-lymphotropic virus exhibits male-

biased virulence, potentially because transmissions occur during pregnancy, birth or breast-

feeding, and so selection has favoured lower virulence in women [36]. Empirical studies of

sex-biased STI characteristics in animals are limited, but mathematical models have shown

how sex-biased disease characteristics influence transmission dynamics (reviewed in [11],

also see [12]). However, the interplay between sex ratio and disease-characteristics

(especially when these are sex-biased) and their effects on STI prevalence have yet to be

investigated. Here we use mathematical modelling to examine the effects of biases in

population sex ratio and disease characteristics on STI epidemiology. Specifically, we

investigate how (1) variation in population sex ratio; (2) sex-biases in transmission and

disease-associated mortality (virulence) in adults; and (3) the interactions between sex ratio

and disease characteristics, impact on STI prevalence in males and females.

2. Methods

We model the dynamics of an STI in a randomly mixing population. The epidemiological and

mating dynamics are described by the following system of ordinary differential equations:

117
$$\frac{dS_i}{dt} = \underbrace{b_i(S_M, S_F, I_M, I_F)}_{Births} - \underbrace{\frac{\beta_i c S_i I_j}{N}}_{Infections} - \underbrace{dS_i}_{Natural\ mortality} + \underbrace{\gamma I_i}_{Recovery} \tag{1}$$

118
$$\frac{dI_i}{dt} = \underbrace{\frac{\beta_i c S_i I_j}{N}}_{Infections} - \underbrace{dI_i}_{Natural\ mortality} - \underbrace{\alpha_i I_i}_{Mortality\ virulence} - \underbrace{\gamma I_i}_{Recovery} \tag{2}$$

119   where $i, j \in \{M, F\}$, $i \neq j$, correspond to males and females. The birth rates for each sex are

120
$$b_M(S_M, S_F, I_M, I_F) = \frac{bcr(1 - qN)}{N}(S_M + I_M)(S_F + I_F), \tag{3}$$

121
$$b_F(S_M, S_F, I_M, I_F) = \frac{bc(1 - r)(1 - qN)}{N}(S_M + I_M)(S_F + I_F), \tag{4}$$

and: S_i and I_i (equivalently, S_j or I_j), are the number of susceptible and infected individuals,

respectively, in each sex; $N = S_M + S_F + I_M + I_F$ is the total population size; r is the sex

ratio at birth (the proportion of offspring that are male), with $0 \leq r \leq 1$; $c \geq 0$ is the pairwise

rate at which males and females mate given frequency-dependent contact; q represents the

strength of density-dependence on births; d is the natural mortality rate; α_i is the disease-

associated mortality rate (virulence) in sex i; β_i is the transmission rate from sex j to sex i

per sexual contact; γ is the recovery rate. The probability that a susceptible female mates with

an infected male (similarly, for the reverse scenario) is equal to $\frac{I_M}{S_M+I_M}$, which is obtained by

dividing the mating rate with infected males $\frac{cS_F I_M}{N}$ by the total mating rate for susceptible

females $\frac{cS_F(S_M+I_M)}{N}$.

For simplicity, we assume that: (i) both sexes have the same recovery rate from the STI; (ii)

both sexes have the same natural mortality rate; (iii) the per-capita mating rate is frequency-

dependent, which means that larger populations do not have a higher per-capita mating rate

than smaller populations; (iv) there is no structuring or choice in the mating system (mating is

random); (v) there is no juvenile period; and (vi) there is no effect of parental care on the

population dynamics.

The disease-free equilibrium $(S_M, S_F, I_M, I_F) = (S_M^*, S_F^*, 0, 0)$, of this system occurs at:

$$S_M^* = \frac{bcr(1-r)-d}{bcq(1-r)} \tag{5}$$

$$S_F^* = \frac{bcr(1-r)-d}{bcqr} \tag{6}$$

143 A newly introduced STI will spread in a susceptible population when the basic reproductive

144 ratio, R_0 is greater than 1 (R_0 is calculated as the maximum eigenvalue of the Jacobian of the

145 system, evaluated at the disease-free equilibrium), where:

$$R_0 = \frac{c}{S_F^*+S_M^*} \sqrt{\frac{\beta_M \beta_F S_M^* S_F^*}{(d+\gamma+\alpha_M)(d+\gamma+\alpha_F)}} \tag{7}$$

In order to guarantee that $R_0 > 1$, we require that $\frac{1}{2} - \psi < r < \frac{1}{2} + \psi$, where $\psi =$

$\sqrt{\frac{1}{4} - \frac{(d+\gamma+\alpha_M+\alpha_F)(d+\gamma)}{\beta_M \beta_F c^2}}$. To test the interactions between sex ratio and STI prevalence in

males and females, we start with no sex bias in STI characteristics. This means that males

and females are equally susceptible to the pathogen and the mortality caused by the pathogen

is same in both sexes ($\alpha_M = \alpha_F = \alpha$; $\beta_M = \beta_F = \beta$). STI prevalence in each sex is given as the

number of infected individuals in that sex divided by the total number of individuals in that

sex (at equilibrium, as indicated by asterisks):

$$D_M = \frac{I_M^*}{S_M^* + I_M^*} \qquad (8)$$

$$D_F = \frac{I_F^*}{S_F^* + I_F^*} \qquad (9)$$

To examine the impact of sex-biased disease characteristics on the prevalence of STIs in

males and females, we fix female susceptibility (β_F) and disease-related mortality (α_F) and

vary the corresponding parameters in males. The model is symmetric up to labelling, so the

results are analogous for the converse scenario. We primarily use numerical analysis to

determine disease prevalence as the endemic equilibrium of equations 1-2 is mostly

intractable to algebraic analysis.

3. Results

3.1 Non-sex-biased disease characteristics

We begin by exploring how sex ratio impacts on STI prevalence in males and females when

there are no sex-biases in disease characteristics (i.e. $\alpha_M = \alpha_F = \alpha$; $\beta_M = \beta_F = \beta$). Consistent with

previous theoretical and empirical work [7, 8], the less common sex exhibits higher STI

prevalence at equilibrium (Fig. 1). Intuitively, when there are no sex biases in disease

characteristics, the overall prevalence of the STI is always maximised at an equal sex ratio

($r = 0.5$). However, both the extent to which STI prevalence differs between the sexes (i.e.

the difference between the male and female curves in Fig. 1a-b, shown in Fig. 1c-d) and the

point at which STI prevalence peaks in each sex (i.e. the sex ratio that maximises the male

and female curves, indicated by circles in Fig.1a-b) depends on the transmission rate and

virulence. Specifically, as the transmission rate increases or as the virulence decreases, the

difference between male and female STI prevalence for a given sex ratio tends to increase,

especially for more extreme sex ratios (Fig. 1c-d). Furthermore, for higher transmission

rates/lower virulence, the sex ratio at birth for which disease prevalence peaks becomes more

extreme. Thus, while overall STI prevalence is qualitatively unchanged by variation in the

transmission rate and virulence (i.e. it always peaks at $r = 0.5$), STI prevalence per sex is

highly sensitive to disease characteristics.

Figure 1. Effects of sex ratio at birth (proportion male), r, on STI prevalence when there are no sex biases in disease characteristics: (a, c) variation in disease-related mortality, $\alpha_M = \alpha_F = \alpha$; (b, d) variation in transmission probability, $\beta_M = \beta_F = \beta$. As disease-related mortality increases or the transmission probability decreases, overall STI prevalence decreases, and the difference between male and female STI prevalence for a given sex ratio decreases. (a, b) STI prevalence in males (solid line), females (dashed line), and overall (dotted line). Markers show where STI prevalence peaks in males (black) and females (white). (c, d) Absolute difference in STI prevalence between males (D_M) and females (D_F), corresponding to the absolute difference between the solid and dashed lines in (a) & (b), with markers showing where the absolute difference peaks. Other parameters are held constant throughout to ensure R_0 is greater than 1: $\delta = 0.1$; $\gamma = 0.2$; $c = 5$; $b = 1$; $q = 0.001$.

We can see this analytically for variation in the transmission rate if we make the following

simplifying assumptions: (1) there is no disease-associated mortality ($\alpha_M = \alpha_F = 0$); (2) the

disease transmission rates are equal ($\beta_M = \beta_F = \beta$); and (3) there is no recovery ($\gamma = 0$).

These assumptions ensure that the adult sex ratio is equal to the sex ratio at birth, which

greatly simplifies the analysis. First, we must calculate the equilibrium population size, N^*.

Let $M = S_M + I_M$ be the number of males in the population and $F = S_F + I_F$, be the number

of females in the population. Then at equilibrium, $M = sN^*$, and $F = (1-r)N^*$. Now, $\frac{dN}{dt} =$

$\frac{dS_M}{dt} + \frac{dS_F}{dt} + \frac{dI_M}{dt} + \frac{dI_F}{dt}$, and evaluating this gives

$$\frac{dN}{dt} = \frac{bc(1-qN)MF}{N} - dN \tag{10}$$

201   Substituting in $M = sN^*$ and $F = (1-r)N^*$, and solving $\frac{dN}{dt} = 0$, the resulting (non-trivial)

202   equilibrium population size is:

$$N^* = \frac{bcr(1-r) - d}{bcqr(1-r)} \tag{11}$$

204   We can now substitute the equilibrium population size into the full system of equations,

205   giving the endemic equilibrium:

$$S_F^* = \frac{d(\beta c(1-r) + d)\zeta}{\beta bc^2 qr(1-r)} \tag{12}$$

$$S_M^* = \frac{S_F^*}{\zeta^2} \tag{13}$$

$$I_F^* = \frac{(\beta^2 c^2 r(1-r) - d^2)\zeta}{\beta bc^2 qr(1-r)} \tag{14}$$

$$I_M^* = \zeta I_F^* \tag{15}$$

210   where $\zeta = \frac{\beta c(1-r)-d}{\beta cs+d}$. The disease prevalence in males and females is therefore

$$D_M = \frac{I_M^*}{S_M^* + I_M^*} = \frac{\beta^2 c^2 r(1-r) - d^2}{\beta cr(\beta c(1-r) + d)} \tag{16}$$

$$212 \qquad D_F = \frac{I_F^*}{S_F^* + I_F^*} = \frac{\beta^2 c^2 r (1-r) - d^2}{\beta c (1-r)(\beta c r + d)} \qquad (17)$$

213 To find the peak in disease prevalence with respect to r, one must differentiate equations 16-

214 17 with respect to r and then find where they are equal to zero, i.e., find r_M^*, r_F^* such that

215 $\frac{d}{dr}(D_F)|_{r=r^*} = 0$ and $\frac{d}{dr}(D_M)|_{r=r^*} = 0$. For females, STI prevalence peaks at

$$216 \qquad r_F^* = \frac{\beta c + d - \sqrt{\beta c d + 2d^2}}{\beta c} = 1 + \frac{d - \sqrt{\beta c d + 2d^2}}{\beta c} \qquad (18)$$

217 and for males STI prevalence peaks at

$$218 \qquad r_M^* = \frac{-d + \sqrt{\beta c d + 2d^2}}{\beta c} \qquad (19)$$

219 (Note that the other root lies outside $r \in [0,1]$). Taking derivatives with respect to β, we find

220 that $\frac{dr_F^*}{d\beta} > 0$ and $\frac{dr_M^*}{d\beta} < 0$. Thus, as the transmission rate, β, increases, the sex ratio at which

221 STI prevalence peaks in each sex always becomes more extreme.

3.2 Sex-biased disease characteristics

We now consider the effects of sex-biased disease characteristics on STI prevalence.

Specifically, we investigate how STI prevalence (per sex and overall) varies when there are

biases in (1) female-to-male (β_M) or male-to-female (β_F) transmission, and/or (2) disease-

associated mortality (virulence) for males (α_M) or females (α_F). Since the only differences in

the model between males and females are these parameters, without loss of generality we fix

β_F and α_F whereas vary β_M and α_M accordingly.

Sex differences in transmission rate and in virulence influence disease prevalence in both

sexes, although the extent of their influence on males and females depends on sex ratio. A

higher female-to-male transmission rate relative to the male-to-female transmission rate

raises disease prevalence in both sexes, although interestingly the influence of sex ratio

differs between the sexes. With greater female-to-male transmission, the disease prevalence

peak shifts toward male-biased sex ratios in females whereas for males there is relatively

little change (e.g. Fig. 2a-c; Fig. 3a-c). Similarly, lower disease-associated mortality in males

relative to females also universally increases disease prevalence and shifts the peaks of the

disease prevalence curves. However, lower values of α_M generally cause the disease

prevalence peaks in each sex to shift towards lower values of r, with greater effects on male

than female STI prevalence (e.g. Fig. 2b, e, h; Fig. 3d-f).

Figure 2. STI prevalence in males (solid line), females (dashed line), and overall (dotted line) as a function of the sex ratio at birth (proportion male), r, when there are sex biases in STI transmission rate and virulence. The parameters $\beta_F = 0.5$ and $\alpha_F = 0.3$ are held constant throughout, while $\beta_M \in \{0.2, 0.5, 1\}$ and $\alpha_M \in \{0, 0.3, 0.6\}$ vary between panels. Other parameters as described in Fig. 1.

Figure 3. STI prevalence in (a, d) males, (b, e) females, and (c, f) overall as a function of the sex ratio at birth and: (a-c) female-to-male transmission rate (β_M); (d-f) mortality virulence in males (α_M). The red lines show when the sex ratio at birth is equal (solid) and when there is no sex bias in disease characteristics (dashed): (a-c) $\beta_M = \beta_F = 0.5$; (d-f) $\alpha_M = \alpha_F = 0.3$. The white region in each panel corresponds to conditions where the STI is unable to persist ($R_0 < 1$). The parameters $\beta_F = 0.5$ and $\alpha_F = 0.3$ are held constant throughout, Other parameters as described in Fig. 1.

We see consistent patterns when transmission and virulence are varied simultaneously

(Fig. 4). By measuring the sex ratio at birth, r, for which disease prevalence peaks, we

analyse how the interaction between transmission, virulence, and sex ratio leads to changes in

disease prevalence. When disease-associated mortality in males is relatively low ($\alpha_M < \alpha_F$),

variation in the female-to-male transmission rate (β_M) has little impact on the sex ratio where

disease prevalence peaks in males (Fig. 4a) and overall (Fig. 4c), but the effect in females is

much more pronounced, with the skew generally increasing for greater β_M (Fig. 4b). When

disease-associated mortality in males is relatively high ($\alpha_M > \alpha_F$), the sex ratio at birth

where disease prevalence peaks in males and overall also increases with β_M, but to a lesser

extent than in females. It is also clear from the contours in Fig. 4 that variation in either

transmission or virulence can cause non-monotonic changes in the disease prevalence curves.

For example, the horizontal and vertical red lines in Fig. 4 (indicating $\beta_M = \beta_F$ or $\alpha_M = \alpha_F$)

intersect some of the contours twice, which means that varying either α_M or β_M can cause a

non-monotonic change in the skew of the disease prevalence curves. Together, these results

show that interactions between sex-biased disease characteristics and sex ratio can lead to

major changes in STI prevalence within each sex and overall, and that the differences

between male and female STI prevalence can be exacerbated by greater transmission rates or

lower mortality rates, in one or both sexes.

Figure 4. The sex ratio at birth for which disease prevalence peaks in: (a) males, (b) females, and (c) overall, as

a function of disease-associated mortality for males (α_M) and female-to-male transmission rate (β_M). The red

lines show when there is no sex bias in transmission ($\beta_M = \beta_F = 0.5$; dashed) and when there is no sex bias in

virulence ($\alpha_M = \alpha_F = 0.3$; solid). The white region in the bottom right corner of each panel corresponds to

conditions where the STI is unable to persist ($R_0 < 1$). Other parameters as described in Fig. 1.

4. Discussion

Studies of STI epidemiology emphasise the importance of the host mating system for STI dynamics [8-11, 19, 24]. Since sex ratio influences mating systems, here we used mathematical modelling to examine the interaction between sex ratio and disease-characteristics, and their effects on STI prevalence in males and females. At a fundamental level, our results reveal how sex differences in STI prevalence within and between populations may be explained by both variation in sex ratio and disease characteristics. Moreover, our model shows that disease-characteristics, whether they be sex-biased or not, can increase or decrease differences in male and female STI prevalence.

In a population with a skewed adult sex ratio, the less common sex has more prospective mates and a higher per-capita mating rate than the more common sex. This pattern can be seen in human and non-human societies; for instance, in human communities with a male-skewed adult sex ratio, men are more likely to purchase sex and women are more likely to have multiple sex partners [37, 38]. Consistently, in bird populations mating opportunities are related to adult sex ratios: in male-biased adult sex ratio, females have higher mating opportunities than males, whereas in female-skewed populations males re-mate faster than the females [39-41]. Hence, with high variance in mating success, many individuals of the more common sex may remain unmated or have a low mating rate, whereas the less common sex has a higher mating rate and hence greater exposure to infectious partners. Thus, equilibrium disease prevalence in the more common sex is expected to be lower than in the rarer sex, consistent with previous theoretical and empirical studies [7, 8, 10, 11].

Our model shows how STI prevalence – overall and in each sex – depends on both sex ratio and disease characteristics. We have shown that even when there are no sex biases in disease

characteristics, increasing the transmission rate or decreasing mortality virulence generally

leads to a greater difference in STI prevalence between the sexes for a given sex ratio (Fig.

1c-d). We have also shown that the sex ratio for which disease prevalence peaks in each sex

depends on disease characteristics, again even when the disease itself shows no sex biases in

transmission or virulence. The reason for these patterns can be understood in terms of a

balance between an increase in the per-capita mating rate for the less common sex and a

decrease in overall disease prevalence as the sex ratio becomes more skewed. Suppose the

adult sex ratio is initially equal and that there are no biases in disease characteristics, hence

equilibrium STI prevalence is the same in both sexes. If we gradually remove males from the

population (similarly for females), then the per-capita mating rate for the remaining males

increases, and hence so does their risk of infection, at least initially. However, skewing the

sex ratio in either direction reduces the overall mating rate, and therefore always lowers

disease prevalence at the population level. As one continues to remove males from the

population, overall disease prevalence begins to fall more sharply because uninfected males

are increasingly less likely to mate with infected females. Thus, disease prevalence

eventually begins to fall in males as well. Once the population crosses a critical sex ratio, R_0

falls below 1 and the disease is driven extinct. Higher transmission and lower virulence both

increase overall disease prevalence and therefore buffer against the effects of skewed sex

ratios, which shifts the sex ratio at which disease prevalence peaks in each sex to more

extreme values. This can be seen in Fig. 1a-b, where the gradient of the overall disease

prevalence curves is flatter for higher values of β and lower values of α.

We also explored the effects of sex-biased transmission and virulence. Sex-biased disease

characteristics are common in wild populations as well as in human populations [29, 42-44],

and previous studies have looked into the behavioural and physiological causes of such

disparities and their implications for population dynamics [26, 27, 31]. However, the

interaction between these biases and STI prevalence within each sex have previously been

overlooked. Our model suggests that sex-biased disease characteristics may independently

influence STI prevalence in a similar way as the unbiased transmission rate or virulence (see

above). However, the prevalence curves for each sex are no longer symmetrical when there

are biased STI characteristics (Fig. 2). Our model also reveals that interactions between sex

ratio and sex-biased disease characteristics can lead to differential and non-intuitive patterns

for STI prevalence. For example, while increasing the bias in transmission from females to

males causes STI prevalence in females to peak at a more extreme sex ratio, there are only

marginal effects on males, yet the effects of male-biased virulence on disease prevalence are

much more pronounced in males than females (as shown in Fig. 3). The model also predicts

that increasing or decreasing sex-biased disease characteristics may lead to non-monotonic

variation in STI prevalence. In the absence of mathematical modelling, it is unlikely that one

would be able to intuit such effects, but our results suggest that there are likely to be complex

interactions between sex-biased disease characteristics and population sex ratio that shapes

STI epidemiological dynamics.

Our model is simple and makes few biological assumptions. Therefore, follow-up analyses

are necessary to investigate biological factors mediating STI prevalence that are not captured

in our model. First, previous studies have emphasised the importance of pair formation in STI

prevalence [45, 46], however, our model assumes mating is random, there is no variance in

mating rate, and hosts do not form pair-bonds [47-50]. This was important for the present

study as we wanted to focus on effects of sex ratio and disease characteristics, all else being

equal. However, future work would benefit from taking pair-formation into account, as a

series of studies in human STIs have shown the variations in form and duration of

partnerships may have an effect on pathogen prevalence [51-54]. Second, previous studies

have shown how variance in mating rate [10, 11] and mate choice [49, 50, 55] affect disease

dynamics, which in turn may influence host and STI evolution in non-intuitive ways. When

one sex experiences higher variance in mating rate, the mating population is effectively sex-

skewed and so we would expect this to generally have a similar effect on STI prevalence as

seen in our model. If infection status impacts on mating rate [47, 49, 50, 56], then the effects

may be more complex, especially if mate choice is driven by one sex.

Previous models have explored how STIs might affect the evolution of host mating dynamics

[46-50], but the focus of the present study has been on STI prevalence. However, given that

there are interactions between sex ratio, sex-biased disease characteristics and mating

dynamics, there are likely to be evolutionary implications for these factors on hosts and STIs,

which should be the target of future research. In particular, it is well-established that disease

prevalence is important for the evolution of host defence [57], and so biases in sex ratio or

disease characteristics are likely to influence selection for traits such as resistance, tolerance

or mate choice through STI prevalence [49, 50]. Additionally, since females are more

important than males for population growth in many species, female-biased disease

prevalence might be more likely to increase population extinction risk.

Our results have several implications for STIs in natural populations. First, when sex ratios

are skewed, we should expect STIs to be more prevalent in the rarer sex, although the relative

prevalence of co-circulating STIs in males and females will differ due to variation in

transmission and virulence (Fig. 1c-d). For example, in a female- skewed population, an STI

with a high transmission rate is likely to have a higher prevalence in males relative to females

compared to an STI with a low transmission rate. Second, our model predicts that variance in

the sex ratio between populations to drive STI prevalence in males and females, with

divergence from equal sex ratios generally increasing sex differences in prevalence (except at

extreme sex ratios). Finally, we should expect the interaction between sex ratio and sex-

biased transmission or virulence to consistently have a stronger effect on STI prevalence in

one sex than in the other.

5. Conclusion

To conclude, the objective of our model was to examine the interplay between sex ratio and

disease characteristics and explore their effects on STI prevalence. Our key message is that

while the less common sex is predicted to exhibit higher STI prevalence at equilibrium,

disease characteristics such as transmission rate and virulence – sex-biased or not – combined

with sex ratio, drive differential patterns in male and female STI prevalence.

Funding

This work is supported by Chinese Scholarship Council and The National Research,

Development and Innovation Office of Hungary (grant no. ÉLVONAL KKP-126949, K-

116310 to NH). University Research Scholarship Award from the University of Bath (to AP).

Royal Society Wolfson Merit Award WM170050, APEX APX\R1\191045 and Leverhulme

Trust (RF/ 2/ RFG/ 2005/ 0279, ID200660763 to TS). The Natural Environment Research

Council (grant no. NE/N014979/1 to BA).

Author contributions

BA, NH and AP designed and analysed the model. NH and AP co-wrote the first draft, and

all authors contributed to revisions of the manuscript.

Data availability

Source code is available in the Supplemental Material and in the following Github repository:

https://github.com/ecoevogroup/Halimubieke_et_al_2020.

Competing interests

The authors declare that they have no competing interests.

References

[1] Ashby, B., Jordan, J.E., Knell, R.J. & Hurst, G.D.D. 2019 Sexually transmitted infections in natural populations: what have we learnt from beetles and beyond? In *Wildlife disease ecology: linking theory to data and application* (eds. P.K. Wilson, A. Fenton & A. Tompkins), pp. 187-222. Cambridge, Cambridge University Press.

[2] Lockhart, A.B., Thrall, P.H. & Antonovics, J. 1996 Sexually transmitted diseases in animals: ecological and evolutionary implications. *Biological Reviews of the Cambridge Philosophical Society* **71**, 415-471. (doi:10.1111/j.1469-185X.1996.tb01281.x).

[3] Dyson, E.A. & Hurst, G.D.D. 2004 Persistence of an extreme sex-ratio bias in a natural population. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 6520-6523. (doi:10.1073/pnas.0304068101).

[4] Liker, A., Freckleton, R.P. & Székely, T. 2013 The evolution of sex roles in birds is related to adult sex ratio. *Nature Communications* **4**. (doi:10.1038/ncomms2600).

[5] Székely, T., Weissing, F.J. & Komdeur, J. 2014 Adult sex ratio variation: implications for breeding system evolution. *Journal of Evolutionary Biology* **27**, 1500-1512. (doi:10.1111/jeb.12415).

[6] Bonnet, X., Golubovic, A., Arsovski, D., Dordevic, S., Ballouard, J.M., Sterijovski, B., Ajtic, R., Barbraud, C. & Tomovic, L. 2016 A prison effect in a wild population: a

scarcity of females induces homosexual behaviors in males. *Behavioral Ecology* **27**,

1206-1215. (doi:10.1093/beheco/arw023).

[7] Hurst, G.D.D., Hurst, L.D. & Majerus, M.E.N. 1997 Cytoplasmic sex ratio distorters. In

Influential passengers: microbes and invertebrate reproduction (eds. S.L. O'Neill, A.A.

Hoffmann & J.H. Werren), pp. 125-154. Oxford, Oxford University Press.

[8] Ryder, J.J., Hoare, M.J., Pastok, D., Bottery, M., Boots, M., Fenton, A., Atkinson, D.,

Knell, R.J. & Hurst, G.D.D. 2014 Disease epidemiology in arthropods is altered by the

presence of nonprotective symbionts. *American Naturalist* **183**, E89-E104.

(doi:10.1086/674827).

[9] Thrall, P.H., Antonovics, J. & Bever, J.D. 1997 Sexual transmission of disease and host

mating systems: Within-season reproductive success. *American Naturalist* **149**, 485-506.

(doi:10.1086/286001).

[10] Thrall, P.H., Antonovics, J. & Dobson, A.P. 2000 Sexually transmitted diseases in

polygynous mating systems: prevalence and impact on reproductive success.

Proceedings of the Royal Society B: Biological Sciences **267**, 1555-1563.

(doi:10.1098/rspb.2000.1178).

[11] Ashby, B. & Gupta, S. 2013 Sexually transmitted infections in polygamous mating

systems. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368**.

(doi:10.1098/rstb.2012.0048).

[12] Anderson, R.M. & May, R.M. 1991 *Infectious diseases of humans*. Oxford, Oxford

University Press.

[13] Eames, K.T.D. & Keeling, M.J. 2002 Modeling dynamic and network heterogeneities in

the spread of sexually transmitted diseases. *Proceedings of the National Academy of*

Sciences of the United States of America **99**, 13330-13335.

(doi:10.1073/pnas.202244299).

[14] Eberhart-Phillips, L.J., Kupper, C., Miller, T.E.X., Cruz-Lopez, M., Maher, K.H., dos

Remedios, N., Stoffel, M.A., Hoffman, J.I., Kruger, O. & Székely, T. 2017 Sex-specific

early survival drives adult sex ratio bias in snowy plovers and impacts mating system

and population growth. *Proceedings of the National Academy of Sciences of the United*

States of America **114**, E5474-E5481.

[15] Emlen, S.T. & Oring, L.W. 1977 Ecology, sexual selection and evolution of mating

systems. *Science* **197**, 215-223. (doi:10.1126/science.327542).

[16] Rankin, D.J. & Kokko, H. 2007 Do males matter? The role of males in population

dynamics. *Oikos* **116**, 335-348. (doi:10.1111/j.2006.0030-1299.15451.x).

[17] Anderson, R.M., Blythe, S.P., Gupta, S. & Konings, E. 1989 The transmission dynamics

of the human immunodeficiency virus type-1 in the male-homosexual community in the

United Kingdom: the influence of changes in sexual behavior. *Philosophical*

Transactions of the Royal Society B-Biological Sciences **325**, 45-98.

(doi:10.1098/rstb.1989.0074).

[18] Anderson, R.M. 1991 The transmission dynamics of sexually transmitted diseases: the

behavioral component. In *Research issues in human behavior and sexually transmitted*

diseases in the AIDS era (eds. J.N. Wasserheit, S.O. Aral & K.K. Holmes), pp. 38-60.

Washington DC, American Society for Microbiology.

[19] Thrall, P.H. & Antonovics, J. 1997 Polymorphism in sexual versus non-sexual disease

transmission. *Proceedings of the Royal Society B-Biological Sciences* **264**, 581-587.

(doi:10.1098/rspb.1997.0083).

[20] Thrall, P.H., Antonovics, J. & Wilson, W.G. 1998 Allocation to sexual versus nonsexual

disease transmission. *American Naturalist* **151**, 29-45. (doi:10.1086/286100).

[21] Jiggins, F.M., Hurst, G.D.D. & Majerus, M.E.N. 2000 Sex-ratio-distorting Wolbachia

causes sex-role reversal in its butterfly host. *Proceedings of the Royal Society B-*

Biological Sciences **267**, 69-73. (doi:10.1098/rspb.2000.0968).

[22] Charlat, S., Reuter, M., Dyson, E.A., Hornett, E.A., Duplouy, A., Davies, N., Roderick,

G.K., Wedell, N. & Hurst, G.D.D. 2007 Male-killing bacteria trigger a cycle of

increasing male fatigue and female promiscuity. *Current Biology* **17**, 273-277.

(doi:10.1016/j.cub.2006.11.068).

[23] Ancona, S., Dénes, F.V., Krüger, O., Székely, T. & Beissinger, S.R. 2017 Estimating

adult sex ratios in nature. *Philosophical Transactions of the Royal Society B-Biological*

Sciences **372**. (doi:10.1098/rstb.2016.0313).

[24] Pastok, D. 2015 Causes of spatial variation in parasite and pathogen pressure in insects,

University of Liverpool.

[25] Sheridan, L.A.D., Poulin, R., Ward, D.F. & Zuk, M. 2000 Sex differences in parasitic

infections among arthropod hosts: is there a male bias? *Oikos* **88**, 327-334.

[26] Miller, M.R., White, A., Wilson, K. & Boots, M. 2007 The population dynamical

implications of male-biased parasitism in different mating systems. *Plos One* **2**.

(doi:10.1371/journal.pone.0000624).

[27] Zuk, M. & McKean, K.A. 1996 Sex differences in parasite infections: patterns and

processes. *Int J Parasitol* **26**, 1009-1023. (doi:10.1016/S0020-7519(96)80001-4).

[28] Klein, S.L. & Roberts, C.W. 2010 *Sex hormones and immunity to infection*. Berlin,

Springer Verlag.

[29] Lemaître, J.F., Ronget, V., Tidière, M., Allainé, D., Berger, V., Cohas, A., Colchero, F.,

Conde, D.A., Garratt, M., Liker, A., et al. 2020 Sex differences in adult lifespan and

aging rates of mortality across wild mammals. *Proceedings of the National Academy of*

Sciences of the United States of America **117**, 8546-8553.

(doi:10.1073/pnas.1911999117).

[30] Metcalf, C.J.E. & Graham, A.L. 2018 Schedule and magnitude of reproductive

investment under immune trade-offs explains sex differences in immunity. *Nature*

Communications **9**. (doi:10.1038/s41467-018-06793-y).

[31] Brei, B. & Fish, D. 2003 Comment on "Parasites as a viability cost of sexual selection in

natural populations of mammals". *Science* **300**, 55a. (doi:10.1126/science.1079746).

[32] Moore, S.L. & Wilson, K. 2002 Parasites as a viability cost of sexual selection in natural

populations of mammals. *Science* **297**, 2015-2018. (doi:10.1126/science.1074196).

[33] World Health Organization. 2007 Addressing sex and gender in epidemic-prone

infectious diseases. (Geneva, World Health Organization).

[34] Coombs, R.W., Reichelderfer, P.S. & Landay, A.L. 2003 Recent observations on HIV

type-1 infection in the genital tract of men and women. *Aids* **17**, 455-480.

(doi:10.1097/00002030-200303070-00001).

[35] Panchanadeswaran, S., Johnson, S.C., Mayer, K.H., Srikrishnan, A.K., Sivaram, S.,

Zelaya, C.E., Go, V.F., Solomon, S., Bentley, M.E. & D Celentano, D. 2006 Gender

differences in the prevalence of sexually transmitted infections and genital symptoms in

an urban setting in southern India. *Sexually Transmitted Infections* **82**, 491-495.

(doi:10.1136/sti.2006.020768).

[36] Úbeda, F. & Jansen, V.A.A. 2016 The evolution of sex-specific virulence in infectious

diseases. *Nature Communications* **7**. (doi:10.1038/ncomms13849).

[37] Tucker, J.D., Henderson, G.E., Wang, T.F., Huang, Y.Y., Parish, W., Pan, S.M., Chen,

X.S. & Cohen, M.S. 2005 Surplus men, sex work, and the spread of HIV in China. *Aids*

19, 539-547. (doi:10.1097/01.aids.0000163929.84154.87).

[38] Bien, C.H., Cai, Y., Emch, M.E., Parish, W. & Tucker, J.D. 2013 High adult sex ratios

and risky sexual behaviors: a systematic review. *Plos One* **8**.

(doi:10.1371/journal.pone.0071580).

[39] Székely, T., Cuthill, I.C. & Kis, J. 1999 Brood desertion in Kentish plover: sex

differences in remating opportunities. *Behavioral Ecology* **10**, 185-190.

[40] Parra, J.E., Beltran, M., Zefania, S., Dos Remedios, N. & Székely, T. 2014 Experimental

assessment of mating opportunities in three shorebird species. *Animal Behaviour* **90**, 83-

90. (doi:10.1016/j.anbehav.2013.12.030).

[41] Carmona-Isunza, M.C., Kupper, C., Serrano-Meneses, M.A. & Székely, T. 2015

Courtship behavior differs between monogamous and polygamous plovers. *Behavioral*

Ecology and Sociobiology **69**, 2035-2042. (doi:10.1007/s00265-015-2014-x).

[42] Poulin, R. 1996 Sexual inequalities in helminth infections: A cost of being a male?

American Naturalist **147**, 287-295.

[43] Schalk, G. & Forbes, M.R. 1997 Male biases in parasitism of mammals: Effects of study

type, host age, and parasite taxon. *Oikos* **78**, 67-74.

[44] Wilson, K., Moore, S.L. & Owens, I.P.F. 2003 Response to comment on "Parasites as a

viability cost of sexual selection in natural populations of mammals". *Science* **300**, 55b-

55b. (doi:10.1126/science.1081430).

[45] Dietz, K. & Hadeler, K.P. 1988 Epidemiological models for sexually-transmitted

diseases. *J Math Biol* **26**, 1-25. (doi:10.1007/BF00280169).

[46] Castillochavez, C. & Huang, W.Z. 1995 The logistic equation revisited: the two-sex

case. *Math Biosci* **128**, 299-316. (doi:10.1016/0025-5564(94)00077-D).

[47] Knell, R.J. 1999 Sexually transmitted disease and parasite-mediated sexual selection.

Evolution **53**, 957-961. (doi:10.2307/2640735).

[48] Boots, M. & Knell, R.J. 2002 The evolution of risky behaviour in the presence of a

sexually transmitted disease. *Proceedings of the Royal Society B-Biological Sciences*

269, 585-589. (doi:10.1098/rspb.2001.1932).

[49] Ashby, B. & Boots, M. 2015 Coevolution of parasite virulence and host mating

strategies. *Proceedings of the National Academy of Sciences of the United States of*

America **112**, 13290-13295. (doi:10.1073/pnas.1508397112).

[50] Ashby, B. 2020 Antagonistic coevolution between hosts and sexually transmitted

infections. *Evolution* **74**, 43-56. (doi:10.1111/evo.13883).

[51] Eames, K.T.D. & Keeling, M.J. 2006 Coexistence and specialization of pathogen strains

on contact networks. *American Naturalist* **168**, 230-241. (doi:10.1086/505760).

[52] Ong, J.B.S., Fu, X.J., Lee, G.K.K. & Chen, M.I.C. 2012 Comparability of Results from

Pair and Classical Model Formulations for Different Sexually Transmitted Infections.

Plos One **7**. (doi:10.1371/journal.pone.0039575).

[53] Kretzschmar, M. & Heijne, J.C.M. 2017 Pair formation models for sexually transmitted

infections: a primer. *Infectious Disease Modelling* **2**, 368-378.

(doi:10.1016/j.idm.2017.07.002).

[54] Leung, K.Y., Powers, K.A. & Kretzschmar, M. 2017 Gender asymmetry in concurrent

partnerships and HIV prevalence. *Epidemics-Neth* **19**, 53-60.

(doi:10.1016/j.epidem.2017.01.003).

[55] Kokko, H., Ranta, E., Ruxton, G. & Lundberg, P. 2002 Sexually transmitted disease and

the evolution of mating systems. *Evolution* **56**, 1091-1100. (doi:10.1111/j.0014-

3820.2002.tb01423.x).

[56] Loehle, C. 1997 The adaptive significance of trunk inclination on slopes: a commentary.

Proceedings of the Royal Society B-Biological Sciences **264**, 1371-1374.

(doi:10.1098/rspb.1997.0190).

[57] Haldane, J.B.S. 1949 Disease and evolution. *Supplement to La Ricerca Scientifica* **19**,

68-76.