Local Field Potentials in a Pre-motor Region Predict Learned Vocal Sequences

Daril E. Brown II[1, 2], Jairo I. Chavez[1], Derek H. Nguyen[1], Adam Kadwory[1],

Bradley Voytek [3, 5, 6, 7], Ezequiel Arneodo [4], Timothy Q. Gentner[2, 5, 7, 8, ¶], Vikash Gilja[1, 7, ¶*].

[1] Department of Electrical and Computer Engineering, University of California San Diego, 9500
Gilman Dr, La Jolla, CA, USA

[2] Department of Psychology, University of California San Diego, 9500 Gilman Dr, La Jolla, CA,
USA.

[3] Department of Cognitive Science, University of California San Diego, 9500 Gilman Dr, La
Jolla, CA, USA.

[4] Biocircuits Institute, University of California San Diego, 9500 Gilman Dr, La Jolla, CA, USA.

[5] Kavli Institute for Brain and Mind, 9500 Gilman Dr, La Jolla, CA, USA.

[6] Halıcıoğlu Data Science Institute, 9500 Gilman Dr, La Jolla, CA, USA.

[7] Neurosciences Graduate Program, University of California San Diego, 9500 Gilman Dr, La
Jolla, CA, USA.

[8] Neurobiology Section, Division of Biological Sciences, University of California San Diego,
9500 Gilman Dr, La Jolla, CA, USA.

37   *Corresponding Author: Vikash Gilja

38   E-mail: vgilja@eng.ucsd.edu

Abstract

Neuronal activity within the premotor region HVC is tightly synchronized to, and crucial for, the articulate production of learned song in birds. Characterizations of this neural activity typically focuses on patterns of sequential bursting in small carefully identified subsets of single neurons in the HVC population. Much less is known about population dynamics beyond the scale of individual neurons. There is a rich history of using local field potentials (LFP), to extract information about behavior that extends beyond the contribution of individual cells. These signals have the advantage of being stable over longer periods of time and have been used to study and decode complex motor behaviors, such as human speech. Here we characterize LFP signals in the HVC of freely behaving male zebra finches during song production, to determine if population activity may yield similar insights into the mechanisms underlying complex motor-vocal behavior. Following an initial observation that structured changes in the LFP were distinct to all vocalizations during song, we show that it is possible to extract time varying features from multiple frequency bands to decode both the identity of specific vocalization elements (syllables) and predict their temporal onsets within the motif. This demonstrates that LFP is a useful signal for studying motor control in songbirds. Surprisingly, the time frequency structure of HVC LFP is similar to well established oscillations found in both human and mammalian motor areas, suggesting that similar network solutions may emerge given similar constraints. This similarity in network dynamics, despite distinct anatomical structures, may give insight to common computational principles for learning and/or generating complex motor-vocal behaviors.

Author Summary

Vocalizations, such as speech and song, are a motor process that requires the coordination of several muscle groups that receive instructions from specific brain regions. In songbirds, HVC is a premotor brain region required for singing and it is populated by a set of neurons that fire sparsely during song. However, how HVC enables song generation is not well understood. Here we describe network activity in HVC that precedes the initiation of each vocal element during singing. We show that this network activity is similar to activity that has been documented in human, non-human primate, and mammalian premotor regions preceding and during muscle movements. This similarity in network activity adds to a growing body of literature that finds parallels between songbirds and humans in respect to the motor control of vocal organs. We also showed that this network activity can be used to predict both the identity of each vocal element (syllable) and when it will occur during song. Given the similarities of the songbird and human motor-vocal systems these results suggest that the songbird model could be leveraged to accelerate the development of clinically translatable speech prosthesis.

Introduction

Learned vocalizations, such as speech and song, are generated by the complex coordination of multiple muscle groups that control the vocal organs[1–3]. Like other voluntary movements this coordinated action arises from premotor brain regions [4–8] and is prepared prior to the physical initiation of movement [4,9,10]. How these behaviors are encoded during both preparation and generation remains an important topic of ongoing research. Developing an understanding for how the brain encodes complex sequential motor movements carries implications for the development of neuroprosthetics that aim to return or supplement lost motor function. In addition to their therapeutic benefits, such systems will help create new tools for examining the brain's mechanisms for learning and executing motor sequences.

At present, studying the motor encoding of speech and other complex motor movements in humans is difficult due to the intrinsic hurdles of doing invasive human clinical studies [11–15] and the complexity of human language [2,14]. Clinical studies in humans are difficult. The clinical setting constrains experimental studies with limits in how long and from where researchers can record data. The study of other motor movements is often supplemented by first studying simpler animal models such as non-human primates [16,17] and rodents [18–21]. These animal models fall short when it comes to more complex freely generated motor sequences such as speech. This is due to none of these models having the ability to learn vocal behavior resembling the complexity of human speech [2,22]. For this reason speech production studies, unlike other motor behavioral fields, have been limited exclusively to invasive [11,23,24] and non-invasive [25] clinical studies in humans. Previous work has found that the same cortical motor neurons that control hand and arm movements are also involved in the production of speech in patients with tetraplegia [26], providing preliminary support that human speech research could also be supplemented by a viable translational animal model. A natural candidate model is the songbird, which is widely used to study complex learned vocal behavior [27,28]. The zebra finch (Taeniopygia guttata), in particular, is known for its precisely timed sequentially structured song which is stable over the course of its adult lifetime. Experiments with this system have yielded insights into the acquisition [29,30], maintenance [31,32] and generation [8,33,34] of complex motor-vocal sequences.

The premotor nucleus HVC, used as a proper name, contains neurons with sparse spiking patterns precisely aligned to vocal production [8,35]. It is considered analogous to premotor cortex in human and non-human primates both functionally [36–39] and genetically [40,41]. HVC provides input to two pathways that lead to neurons within nuclei that control the vocal muscles (Fig 1A). The first is directly through RA in the posterior descending pathway (PDP), which is necessary for both the acquisition and production of learned motor-vocal behavior. For reference to mammalian anatomy, the PDP is analogous to a motor pathway that starts in the cerebral cortex and descends through the brain stem. The second, named the Anterior Forebrain Pathway (AFP), plays a strong role in acquisition only and projects indirectly through several nuclei analogous to a cortical pathway through the basal ganglia and thalamus in mammals [28]. These analogous regions share tens of hundreds of convergent genes despite their last shared ancestor being millions of years ago [41]. At a circuit level the similarities between the neural activity in humans and songbirds is much harder to compare. Single unit work and techniques, like those done in songbirds, are difficult to do within a clinical setting. In contrast given the accessibility of the system and a focus on fundamental physiology, songbird studies have largely focused on carefully identified neural units and their spiking activity, with limited examination of other neural features that may correlate with vocalization behavior. Examining neural features that can be recorded from both species and their relationship to motor-vocal behavior will enable direct comparison between

the neural activity in birdsong and human speech production. Clarifying whatever similarities (and
differences) that may exist and potentially bridging the gap between the two.
Local Field Potentials (LFP) are thought to reflect both aggregate local postsynaptic
activity and presynaptic inputs to the recording site [42]. Because stable single units can be difficult
to acquire and maintain in humans and non-human primates there is a rich history of literature
looking at both spiking and LFP for understanding, decoding and predicting motor production [43–
47]. At present most work with HVC in songbirds has focused almost exclusively on single unit
spiking activity [6,28,34,48,49], and limited work has focused on the structure of LFP and how
they relate to song production. The most detailed characterization of this signal in songbirds being
LFP's relation to interneuron synchrony [33]. This leaves a gap in the literature regarding the
structure of LFP activity in HVC and whether its characteristics have any similarities to LFP in
human, non-human primates, or mammalian premotor and motor regions.
We addressed this gap by chronically implanting and recording from freely behaving male
zebra finch (Fig 1B) and analyzing the LFP in relation to each bird's performed song (Fig 1C and
D). Our results show narrow-band oscillations similar to those reported in human, non-human
primate, and rodent motor electrophysiology literature. Further we provide evidence that phase
and amplitude modulation within these frequency bands is predictive of vocalization behavior.

Results

Adult male zebra finches (n=3) were chronically implanted with a lightweight custom 3D
printed microdrive (Fig 1B). Ensemble neural activity and vocal behavior were simultaneously
recorded using a laminar silicone probe and an omnidirectional microphone, respectively (Fig 1C
and 1D). A male zebra finch's learned song consists of 1-7 repetitions of a motif, each of which is
composed of a precisely ordered sequence of 3-10 discrete units called syllables. Song motifs are
also grouped into larger structures called bouts, which consist of multiple repetitions of motifs.
Syllables of each bird's learned song, other non-learned vocalizations, and tokens of non-vocal
intervals were segmented and annotated within periods of vocal behavior, referred to as Vocally
Active Periods (VAP), using acoustic landmarks (Fig 2). Temporal boundaries of VAPs were used
to define behaviorally relevant epochs in the simultaneously recorded neural signal. Subjects were
recorded for 1-10 hrs per day, and statistics of recorded behavior are documented in S1 and S2
Table. To maintain statistical power, we selected the two days with the most vocal activity from
each bird for the analyses reported here (S3-S5 Table). Results from days with fewer recorded
motifs were qualitatively similar.
156
157

Fig 1. Continuous electrical and audio recording of chronically implanted free behaving male zebra finch (Methods). (A) Four-Shank multi-site Si-Probes were chronically implanted targeting nucleus HVC. HVC provides input to two motor pathways that innervate neurons in the tracheosyringeal half of the hypoglosssal nucleus (nXIIts) that projects to vocal muscles. The posterior descending pathway (PDP) comprises a direct projection to the robust nucleus of the achopallium (RA) and is necessary for both the acquisition and production of learned vocal behavior (song). The Anterior Forebrain Pathway (AFP), provides an indirect pathway to RA, through Area X, the dorsolateral anterior thalamic nucleus (DLM), and LMAN, and is necessary for song acquisition. The PDP is homologous to a motor pathway in mammals that starts in the cerebral cortex and descends through the brain stem, while the AFP is homologous to a cortical pathway through the basal ganglia and thalamus. (B) Neural and audio recording apparatus. We recorded LFP and vocal activity from male zebra finches stimulated to sing by the presence of a female conspecific. (C) Exemplar sound pressure waveform (1.3 seconds in duration, top) from bird z007 above the corresponding spectrogram. (D) Ten randomly selected channels of simultaneously recorded neural activity aligned to the song during audio recording, after a 400 Hz low pass filter.

Fig 2: Temporal structure of vocal behavior. (A) Schematic showing the intermittent occurrence
of vocally active periods (VAPs, black bar) at sporadic intervals throughout a multiple-hours-long
continuous recording session of one male zebra finch (z007). (B) Zoomed in view of one 25 second
long VAP comprising several bouts, denoted by light blue rectangles above the sound pressure
waveform and corresponding spectrogram. (C) Zoomed in view of the first bout in segment (B)
showing introductory notes, typically repeated a variable number of times prior to the start of
the bout and labeled as 'i', other vocalizations not part of the courtship song, labeled 'C', and
syllables comprising the courtship song, labeled '1', '2', '3', '4', '5', based on their sequential order
in the motif.

Song-related LFP spectral changes in HVC

Fig 3A illustrates the spectrotemporal characteristics of the LFP amplitude during the preparation and production of bird song, with an exemplar single trial time-varying power spectral density (PSD) during the course of 3 separate bouts. Each frequency is scaled to its percentage of the mean amplitude during the respective VAP. Prior-to and during vocal production there is a broadband increase in power in all frequencies above roughly 50 Hz. This is coupled with rhythmic changes in frequencies below roughly 50Hz. These changes start just prior to song onset and end immediately after vocal production has stopped. Similar patterns emerged in each of the birds and within birds across days. Changes in LFP amplitude appeared to correspond with each instance of vocal behavior, including non-learned portions such as introductory notes.

The spectral structure of the LFP activity immediately prior to and during the production of learned song is highly consistent across bouts as shown when they are aligned to their initiation (Fig 3B) and termination (Fig 3C). These results were calculated by aligning to the start of the first syllable of the first motif in the bout (initiation) and the start of the last syllable of the last motif of the bout (termination) for all bouts recorded over the course of a single day. The aligned neural activity was then averaged for each frequency. As with the single trial time-varying PSD, there is an increase in amplitude for all frequencies above roughly 50 Hz and what appeared to be a drop in amplitude for frequencies below roughly 50 Hz. This figure illustrates the stereotypy of the time correlated spectral structure in HVC as it relates to the preparation and production of bird song. No dynamic time warping was used to align the behavior. Similarly consistent structure was found across all channels and across all subjects (S1 Fig).

As in previous work [31,33], we found a strong stereotyped 25-35 Hz oscillation that was consistent across renditions of motifs (Fig 1B and 1C). As shown in Fig 1A, patterned changes in spectrum occur over times scales longer than a single cycle for a given frequency below approximately 50 Hz. These oscillations were found to have contributions from both phase and amplitude [33], and occur over several frequencies besides 25-35 Hz. Which led us to ask: what other potential frequencies in the LFP might carry information about the preparation and production of song through either phase or amplitude changes.

Decoupling LFP power-spectrum reveals song related spectral features

A spectral decomposition technique previously used in human motor cortex was used [50] to determine which frequencies' amplitude changes were correlated with the production of birdsong. As shown in Fig 4A, on average there was a consistent decrease in the lower frequencies in alpha/beta range and a consistent increase in higher frequencies that are in the range often referred to as "high gamma" [51,52]. Following the steps of the spectral decomposition approach described in the methods, the principal component decomposition of these PSDs found Principle Spectral Components (PSC) similar to those previously reported in [50] (Fig 4). The most consistent PSC, which is the first principle component, was a spectrum wide-amplitude change that was consistent across channels and subjects. The second PSC peaked in the "alpha/low beta range", and in humans has been proposed to reflect the resting rhythms that decrease when cortical areas are activated [53]. Although the PSCs were learned without explicit knowledge of distinct vocally active and inactive time periods, the first three PSCs show that the power spectrum during vocal production is separable from vocally inactive periods (Fig 4C). This suggests that these frequencies could be used to detect the onset of vocal production.

Fig 3: Variation in HVC LFP power correlate with vocal behavior. (A) Time series power spectrum
of LFP activity aligned to the free vocal performance of a male zebra finch (z007) during the
production of three separate bouts. Colored bars above the spectrum annotate the vocal
behavior, with color representing a distinct syllable with the same color coding as Fig 2. (B) Time
series power spectrum (bottom) averaged across multiple renditions of similar bouts (n=27, top)
aligned to the start of the first motif in the bout. (C) As in (B) but aligned to the end of the last
motif in the bout. Above both is a behavioral raster showing the time course of the behavior
being averaged (n=27 bouts). No dynamic-time warping was used. To ensure the start and end
of the bout are unique time periods, only bouts with more than one motif in duration were used.
Behaviorally inconsistent bouts were excluded for clarity of visualization, including them does
not alter the result.

Fig 4: Naïve decomposition of LFP power spectra reveals song correlated features. (A) Averaged Power spectra during trials centered in 1 second intervals where the bird either initiates a motif, called vocally active (light blue), or does not vocalize, called vocally inactive (purple). The (B) The power spectra in Fig. 4A are normalized and naively decomposed (methods). The elements of the first principle spectral component (1st PSC, blue) is non-zero across all frequencies, likely due to the power law in LFP PSDs. The 2nd PSC, golden- yellow, peaks between 0-10 Hz. The 3rd PSC, brown, peaks between 10-30 Hz, but has variations across birds that extend into 50Hz. As the PSCs are all eigenvalues their signs do not matter when interpreting them. This structure is largely consistent across channels and across days. Note the other PSCs are not shown. (C) Projection of both the vocally active and vocally inactive trials into the first three PSCs. The color coding is the same as A.

Syllable onset is phase aligned to underlying LFP rhythms

We next calculated the Inter-Trial Phase Coherence (ITPC) of the spectrum (methods) from 2-200 Hz to determine which frequencies had stable song aligned structured changes in phase. A defining characteristic of zebra finch song is its stereotyped motif. Leveraging this stereotypy, we calculated the ITPC aligned to the first motif of all bouts recorded in a single day (n>28). Any LFP phase coherence occurring prior to song production can be attributed to the preparation of the subsequent vocalization. In two of the three birds, the first motif was almost always preceded by an introductory note. As each frequency and time sample are separate results with an attributed p-value they are all visualized in terms of their Rayleigh Z statistic to allow for comparisons. Fig 5A shows long lasting phase structure that precedes the onset of the bout by up 100 ms in frequencies lower than 50 Hz. This structure continues during the course of the bout and terminates soon after the end of the bout Fig 5B. Similar structures in the ITPC are observed across channels and across

bird subjects [S2 Fig]. The phase coherence is highly sensitive to jitter in the aligned behaviors as
shown in the more variable birds which motif was less deterministic in structure (S2 Fig).
To determine the frequencies where phase coherence is most consistent across individual
subjects we computed the normalized sustained Z Statistic (methods) for the two high yield days
for all subjects (Fig 5C). This revealed statistically reliable oscillations in the previously-
established 25-35 Hz range, as well at several other frequencies: 4-8 Hz, 8-12 Hz, 35-50 Hz, and
50-70. The lower frequencies exhibited longer periods of phase coherence through time than the
higher frequencies. These periods occurred over several cycles and would fall out of alignment
faster than a full cycle once song terminated (Fig 5B and S2 Fig). Strong phase coherence was
observed throughout the production of the motif without the need for dynamic time warping to
force motifs into the exact same time scale (Fig 5 and S2 Fig).
To better understand the dynamics underlying HVC population activity, we examined the
song-aligned LFP phase coherence in greater detail. One hypothesis, given the highly stereotyped
spectro-temporal structure of the motif including the brief gaps between each syllable, is that both
the vocal output and the underlying HVC dynamics are largely deterministic. That is, the observed
coherence may reflect an initial, single alignment at the start of the motif, that endures only by
virtue of the very low song-to-song variability. Alternatively, it could be that HVC population
activity reflects a more tightly controlled timescale, in which the observed oscillations are aligned
to shorter timescale vocal events such as syllables. To determine whether alignment is tied to the
motifs (hypothesis one), or unique to each syllable (hypothesis two), a smaller scale time alignment
was used. We first verified that phase locking was found when localized to each vocalization.
Syllable specific polar phase analysis, shown in Fig 6A, shows a strong phase preference to the
onset of each syllable in all of the previously determined frequency bands. These phases were
unique to each frequency band for each syllable. Similar results were seen for all subjects across
all days (S3-S6 Fig). We also found the same phase preference to vocalization onset for the
introductory note despite its huge variability in duration and structure beyond a single utterance.
Fig 6B and 6C demonstrate that this stereotyped phase structure occurs prior to and during each
vocalization and is statistically unlikely to have occurred by chance. To directly test the two
hypotheses, we compared the ITPC centered on each syllable after the first to the ITPC centered
on their stereotyped time within the motif (S7 Fig). Centering on the labeled syllable start time
yield significantly stronger ITPC compared to centering on the stereotyped, or averaged, start time
for each syllable within the motif (S11 Fig). Thus hypothesis two is the more likely source of this
phase locking characteristic.
305

Fig 5: LFP Inter-trial phase coherence during production of learned sequences. (A) ITPC of LFP aligned to the start of the first motif of the bout for z007, left, and (B) the end of the last motif of the bout, right, above both is a behavioral raster showing the time course of the behavior being averaged (n=28 bouts). No dynamic-time warping was used. To ensure the start and end of the bout are unique time periods, only bouts with more than one motif in duration were used. Behaviorally inconsistent bouts were excluded for clarity of visualization; including these bouts does not alter the result. (C) Normalized sustained Z- statistic of the ITPC for samples preceding the labeled start of the bout. The steps for calculating this metric are detailed in the methods. Data reflect all bouts from both high yield days for three birds.

Fig 6: Syllable specific inter-trial phase coherence reveals phase preference to syllable onset.
(A) Polar histogram of the phase for each LFP frequency band at the labeled start of all instances
of a given syllable over the course of one day, for one bird (z007). The number of instances
(n=100) are equal for all syllables, and set by the syllable class with the fewest renditions. (B) ITPC
resultant vector length for each frequency over time relative to the labeled start of each syllable
(0 ms) over the same instances as in (A). (C) Rayleigh Z-statistic of the ITPC over the same time
and frequencies as (B).

LFP features encode intended syllable identity

With potential LFP bands found through the decomposition of both phase and power we
next asked whether their spectral features were correlated to vocalization identity. If so, then these
features could be used to classify vocalization identity from neural activity alone. As the
dominating characteristic of these features were there consistent structure, a promising approach
was to create LFP templates that could be correlated with representative time traces. It was unclear
however, what the ideal time range and latency relative to song onset for this information might
be. To examine this, we conducted a hyper-parameter search using a LDA SVD classifier trained
on varying combinations of template duration in samples (bin width) and the number of
milliseconds prior to event onset from which the template was taken (offset). For most frequencies
found, multiple combinations of offset and binwidth were able to correctly classify between song
syllables, silent (non-vocal) periods, and —for the subject that performed them— introductory
notes (z020 & z007) significantly above binomial chance (Fig 7 and S10 Fig). The parameter
search for each frequency was done separately and found increasing performance with decreasing
offset. S6 and S7 Tables summarize the parameters in the search yielding the highest classification
accuracy. The information for vocal classification from lower frequencies were stable over longer
time spans, i.e. larger bin widths, and may be more useful for predicting song (bout) onset. Nearly
all frequencies we examined were useful for classifying prior neural activity as it relates to the
song it will produce downstream; however it is not immediately clear what component of these
oscillations, i.e. phase or amplitude, carries information regarding song identity.

To determine which component (phase or amplitude) of each frequency band provided
information about the vocalization identity, we used Hilbert transforms to selectively remove
either phase or power, while retaining only the contributions of power or phase information,
respectively. Information provided by a given component was inferred from classifier accuracy.
As shown in Fig 7A-C both phase and amplitude carry information for all frequencies. In general,
however, more information was encoded by phase for lower frequencies (below 30 Hz) and more
information was encoded via frequency amplitude in higher frequencies (above 30 Hz). Whereas
25-35 Hz appears to receive about equal performance when using information from either phase
or amplitude only. We also asked if the number of channels recorded were necessary for decoding
song identity by running a channel adding analysis for each frequency band. As shown in Fig 7A
the LDA classifier performed well above binomial chance with only 5-10 channels of neural
activity when classifying between song syllables, silence and introductory notes.
357

Fig 7: LFP phase and power provide independent and additive information for vocalization classification. (A-C) Channel dropping curves showing classifier performances with either (A) all LFP phase information removed, (B) all LFP power information removed, or (C) with both phase and power used as independent features. Each row corresponds to data from the highest yield day for each bird. z007 n=98 for each 7 classes (1, 2, 3, 4, 5, i (introductory note), Silence), z020 n=91 for each of 7 classes (1, 2, 3, 4, i (introductory note), Silence), and for z017 n=52* for each 8 classes (1, 2, 3, 4, 5, 6, 7, Silence). The results up to 15 channels are shown to allow for far comparison across subjects. Dark lines show the mean for each vocalization class, shaded areas give the standard deviation over the bootstrapped analysis using n=5,000 repetitions across 5 cross-validation folds. The p-value for all of the Binomial chances calculated for each bird was 0.05.

*The number of instances for each class was limited by Syllable 7, which is a intra-motif note.

Causal LFP Features Predict Syllable Onsets

The ability to accurately classify specific syllables using the features of the LFP, may be possible because a unique LFP structure associated with specific syllables or because both tend to co-occur at specific times within the motif. To dissociate these two possibilities, we examined whether the features optimized for determining syllable identity (Table 6 and 7) could also predict specific onset times of each separate vocalization within the full motif. To do this, we implemented a naive pattern-matching approach (Methods). Because of the well-documented and well-studied stereotyped structure of zebra finch's song, the mean error between the stereotyped start time of each syllable—relative to its first syllable—and the actual time the syllable occurred was used as a benchmark to test the performance of a predictor that used the LFP features as described before. An example of one motif and its corresponding confidences is shown in Fig 8A. Across all syllables for all birds the predictor that used all of the frequencies performed better than using the deterministic onset of the syllable only, Fig 8B. Similar performance was found for the 2 additional syllables for z017 as well (S11 Fig). These results mostly hold true across all days analyzed and were consistently found to be statistically significant (p<0.05) using the Wilcoxon signed rank test. There were only three cases in which the LFP predictor's performance failed to meet significance when using the Benjamini-Hochberg False Discovery Rate; they were syllable 2 & 4 for z020's 2nd high yield day and syllable 2 for z017's 2nd high yield day (S11 Fig). We next asked which frequency, if any, is best at predicting the onset of the syllables. As shown in Fig 8C, no single frequency performs best for predicting onset times across all frequencies and all subjects. Although the higher frequencies tend to do better and have lower variance than the lower frequencies, using all of the frequencies yields better performance than any one frequency across all subjects and sessions. There were poor prediction performances for 4-8 Hz, 8-12 and the 50-70 Hz for some syllables for certain birds. Further, results from 4-8 Hz were more likely to be statistically insignificant and performed worse than the stereotyped reference for most syllables.

Fig 8: Onset detection using LFP features. (A) Example motif from subject z007. Annotated behavior (top) using the same color scheme as in Fig 2, sound pressure waveform and the corresponding time-aligned spectrogram (middle), and the time-varying naïve confidence of the onset prediction (bottom) for each syllable in this example motif. Confidence signal traces are the same color as the syllable they are meant to predict. (B) Histogram of onset prediction times for each syllable relative to its annotated start time. The annotated start times are relative to the start of the first syllable of the motif the syllable occurred in. The histogram compares two approaches. The first, in pink, uses only behavioral information (the stereotyped onset relative to the start of the first syllable). The second, in blue, uses all the neural features to predict the start of the syllable. (C) Boxplot of onset prediction times relative to the labeled onset time for each bird. The order of each feature used is the same, going left to right: first is the stereotyped onset time using only the deterministic behavior, next is the results using all of the neural features, then each frequency band only in order from least to greatest (4-8 Hz, 8-12 Hz, 25-35 Hz, 35-50 Hz, and 50-70 Hz). Statistical significance was calculated using the one sided Wilcoxon signed-rank test, and * denotes results that are **not** statistically significant when using the Benjamini-Hochberg False Discovery Rate. All other results p<0.05 and q<0.05.

Discussion

We demonstrate that the local field potentials in zebra finch HVC carry significant information tied to vocal production. More specifically, the time-varying structure of this meso-scale signal, captured by both the power and phase of various LFP frequency bands, correlates with behaviorally-relevant timepoints during this complex, learned motor behavior. Our results extend previous studies in zebra finch HVC [33], by showing that LFP features can be used to detect, using simple methods, both when and what syllable the bird will sing. Prior to this previous study and our work, little was known about the relationship between LFP features and vocal production in birds. This limited the generalizability of the otherwise powerful birdsong model to other systems, including non-human primates and especially humans, where LFP and surface potentials are broadly used to measure the dynamics of cortical motor control [12,47,50,54]. In addition, we note that the bandwidth and features of the LFP signals investigated in this paper share similarities with LFP features tied to motor control in humans, non-human primates, and mammals. For example, the power spectrum components most closely tied to song in finches (Fig 4), match those documented in the human motor cortex during finger flexion [50]. We suggest that LFP recordings can serve as useful targets to further our understanding of songbird neurophysiology, and to more closely connect the powerful birdsong model to motor control studies in mammals, non-human primates and humans.

A consistent feature observed in the LFP of HVC in songbirds during singing is tight phase locking to vocal behavior. In addressing the question of what potentially gives rise to the observed song-aligned phase locking; we proposed two hypotheses. Hypothesis one where the phase locking is related to the larger motif sequence and hypothesis two which attributes the phase locking to each unique vocal units (syllables). In simpler terms is the phase locking a continuous process aligned to the start of the motif (Hypothesis one) or discrete process that aligns to each syllable (Hypothesis two). When directly testing the two hypotheses the second proved more statistically likely. While there is noticeable non-deterministic jitter in the brief gaps of silence between syllables that make up the motif, this may not be enough to completely disprove hypothesis one. However there are other notes that occur non-deterministically between motifs, referred to as intra-motif notes or "connectors" [55] and introductory notes for whose timing before the motif is variable. Looking at the phase polar plots of individual syllables would either show high selectivity of phase preference for all unique vocalizations if hypothesis two is correct or low levels of phase locking for motif syllables and a random distribution about the polar axis (null) for the intra-motif notes and introductory notes if hypothesis one is correct. As shown in Fig 6 and S7 Fig, each syllable has phase preference that is stable across every specific vocalization type including intra-motif notes (z017 & z020) (S4-S6 Fig) and introductory notes (z020 & z007) (Fig 6 and S3 Fig). This result indicates that precise phase preference is behaviorally relevant to both learned and non-learned vocalizations. Thus, it can be thought that the long template phase traces in lower frequencies that precede and persist during song production are a byproduct of the zebra finches' stereotyped song structure. This structure is composed of encoding at a smaller time scale centered on each individual syllable during song production. Similar phase preference to motor onset has been found in both human and non-human motor cortex [56].

Naturalistic motor behavior is intrinsically difficult to study due to its high degree of variability and limits on brain coverage in chronic experiments. Zebra finch are a desirable model species because they mitigate the first of these difficulties. Their song allows for repeated segments of the same sequence to be produced with two almost opposing characteristics: near perfect precision of each utterance and the natural irregularities that exist in all of nature. This provides a

dataset that has both high repetitions of each syllable class and a non-deterministic structure
beyond the motif level that facilitates detailed analyses. This model is a unique testbed for
approaches to predict the onset of vocalizations in other animals, particularly humans. The simple
onset prediction results where each syllable after the first were predicted using causal neural
features, described in Fig 8, were done using little-to-no parameter optimization and take into
account no information regarding the structure of the song. Better performance could be achieved
with more elegant approaches with minimal increases in computational complexity. Suggesting
that these features could be leveraged to predict song onset with low latency. At present the state
of the art for Human speech BCI's uses non-causal neural signals to synthesize sounds similar to
the intended speech [12,24,57]. While promising, the non-causal nature of these algorithms
introduces significant latency [57] and ultimately limits the quality of interaction with such a
prosthesis. Even further, the limits on how long researchers have access to neural signals impedes
the speed at which these offline techniques can be translated in real-time in a clinical setting. This
is where the songbird model can contribute to the BCI field; as an animal model to proof-of-
concept systems in which closed-loop interaction with prosthesis designs can be rigorously
studied. Systems derived from these proof-of-concepts will be highly valued when translated to
the human clinical setting. These features and approach provide a starting point for further analysis
to look to zebra finch as more than a model for vocal learning and sequence generation, but as a
model of vocal prediction and BCI development.
482
483
484

Methods

We implanted male zebra finch with laminar electrodes and simultaneously recorded their
vocal behavior and neural activity (Fig 1). Through a series of steps, explained in detail in the
Annotation and Alignment of Behavioral Data section, we found segments of the recording that
contained songs and hand annotated them using Praat (Fig 2). These behavioral labels were used
to analyze the neural activity in relation to specific classes of behavior to determine what
frequencies, if any, correlated to the behavior (Fig 3-6). Finally, the frequencies discovered were
used to classify and predict the onsets of the behaviors to clarify their relationship to the
vocalizations (Fig 7 & 8).
494
Subjects
All procedures were approved by the Institutional Animal Care and Use Committee of the
University of California (protocol number S15027). Electrophysiology data were collected from
three adult male zebra finches. Birds were individually housed for the entire duration of the
experiment and kept on a 14-h light-dark cycle (Fig 1A). Each day had one session that lasted
multiple hours and were unstructured as they depended on the subject's willingness to sing (Fig
2A). All available days were analyzed, however the two highest motif yielding days, hereafter
referred to as *high yield days*, were reported and used for statistical analysis. The full duration of
chronic recordings ranged from 5 to 19 days. The birds were not used in any other experiments.
504

Electrophysiology and audio recording

Electrophysiological recordings were gathered from 3 subjects, in which two were implanted with 16-channel probes and one with a 32-channel probe (Fig 1A). We used 4-shank, 16/32 site Si-Probes (Neuronexus), which were PEDOT-coated in-house. The probes were mounted on a custom designed printable microdrive [58] and implanted targeting nucleus HVC (Fig 1A & B). Audio (vocalizations) was recorded through a microphone (Earthworks M30) connected to a preamplifier (ART Tube MP) and registered temporally to ongoing neural activity (Fig 1)]. Extracellular voltage waveforms and pre-amplified audio were amplified and digitized at 30 kHz using an intan RHD2000 acquisition system, Open Ephys and custom software (Fig 1D) [59].

Analysis of electrophysiology data

Extracellular voltage traces were multiplexed and digitized at 30kHz on the headstage, and stored for offline analysis. **--offline--** They were then low-passed filtered at 400 Hz using a Hamming FIR filter and downsampled to 1 kHz. The group delay introduced by the filter is compensated by introducing a temporal shift to the filter output [60].

Annotation and alignment of behavioral data

The song of an adult male zebra finch can be partitioned and labeled in multiple ways. However, the most fundamental characteristic of their song that is agreed upon is their stereotyped motif. The motif is a consistent sequence of sounds interleaved with silent periods that is learned from a tutor while they are young [10,29,30]. Song motifs are also grouped into larger structures called bouts, which consist of multiple repetitions of motifs [10,29,55]. Depending on the definition used, bouts can include a short repeated vocalization that typically precedes the first motif of the bout. For the purpose of the analysis done, the start of the bout is explicitly defined to be the start of the first motif in that bout. This is due to the focus of understanding the correlation between LFP and learned motor sequences. Beyond the syllables of the motif male zebra finch may have an additional syllable or sequence of syllables they will optionally insert between successive repetitions of motifs. This is called a "connector" [55] or intra-motif note. In our recordings both z017 and z020 had intra-motif notes. Because z007 did not have any intra-motif notes, and therefore a more stereotyped song, it will be used for all of the empirical figures shown.

A custom template matching algorithm written in Python was used to find potential instances of vocal activity (using an exemplar motif as the template), which were then curated manually to rule out false positives [59]. The curated motif start times were grouped into larger time segments that ensured that the gap between each motif was no greater than 20 seconds (Fig 2A and 2B). These chunks of time are subsequently referred to as vocally active periods (VAP). These VAPs were then manually labeled using the Praat speech analysis software [61]. Vocal events were segmented by hand (Fig 2C) and labeled based on identity. There is a unique label for each individual syllable and introductory note (if the bird demonstrated one). All other male vocalizations were grouped together and labeled as a call. Calls are short, simple vocalizations that are produced by both sexes which mostly do not have learned components [30]. There were also two labels for silence in these VAPs: one for the gaps between song syllables, and another for silence not during song in the VAP.

To leverage these behavioral annotations for contextual analysis of their synchronously recorded neural activity, custom Python software was developed [BirdSongToolbox]. This

551 software added additional labels to the hand labels based on their sequence and context within the
552 larger bout structure; e.g. first or last motif in bout. Labeled vocalization segmentation times were
553 adjusted to the proper sampling rate for the downsampled neural data using this same software.
554 Finally, the software used these additional contextual labels to select particular event times to align
555 both vocal and neural data that fit certain criteria matching the particular vocalization of interest.
556

Principle spectral decomposition

Following work done in Human ECoG work, a principle component method was applied
to find consistent changes in LFP spectrum related to motor-vocal behavior. This method required
the calculation of the power spectral density (PSD) for windows of time during either vocal activity
or inactivity. Although the behavioral changes in birdsong are on a smaller timescale (20-100 ms)
than human limb movements, to keep methods consistent between studies the same window size
of 1 second was used for the trial length, τ_q. PSDs of vocally active trials centered on song activity,
which included introductory notes and calls, were analyzed alongside trials of neural activity
centered within larger periods of vocal inactivity longer than 2 seconds in duration. All PSDs were
calculated using the multitaper method with the time_frequency.psd_array_multitaper function
from the MNE-Python software package [60]. The frequency range was set from 0 to 200 Hz with
14 orthogonal slepian tapers. Each PSD, $P(f, \tau_q)$, was individually normalized using two steps:
each spectral sample was elementwise divided by the average across the ensemble, at each
frequency, and then the log was taken.

571
$$\bar{P}(f, \tau_q) = \ln\left(P(f, \tau_q)\right) - \ln\left(\frac{1}{N_q}\sum_{p=1}^{N_q} P(f, \tau_p)\right)$$

The label q refers to the times centered within periods of high vocal activity and vocal inactivity
(silence). The number of instances per class were balanced to be equal and their combined total
number of PSDs is denoted by N_q. The covariance matrix $C(f, \tilde{f})$ between frequencies of these
normalized PSDs were calculated: $C(f, \tilde{f}) = \sum_{\tau_q} \tilde{P}(f, \tau_q)\tilde{P}(\tilde{f}, \tau_q)$. The eigenvalues, λ_k, and
eigenvectors, $\vec{e}_k$, of this matrix elucidate common features during song production. These
eigenvectors, $\vec{e}_k$, are referred to as "Principal Spectral Components" (PSCs) [50].
578

Feature extraction

Feature extraction from the preprocessed data consisted of a series of steps that involved
first Common Average Referencing and then narrow band-pass filtering. We used the Hilbert
transform to compute the analytic signal. To extract oscillatory power the absolute value of the
analytical signal was used, which effectively removed all phase related information. To extract
oscillatory phase the instantaneous angle of the complex argument of the analytical signal was
used, and was further smoothed using a sine function. This effectively removed all information
related to power from the signal.
These signals, whether Hilbert transformed or not, were then sub-selected at times relative
to the label onset for a syllable using two parameters: bin width and offset. The bin width is the
number of samples used prior to the offset, and the offset is the number of samples, or milliseconds,
prior to the labeled start time to a specific syllable. All combinations of these hyperparameters
used offset which were prior to the start of the vocal behavior.

Band templates and Pearson correlation features. Using the optimized bin width and
offset for a particular frequency band, a template for that band was calculated by taking the mean
of the LFP traces of the training set of a specific behavior class. This template represents the
average LFP activity prior to and during the production of a particular vocal behavior. This
template was then used to extract features from a narrow-band LFP trace by taking the Pearson
correlation of the template from a segment of neural activity of the same length in samples. This
correlation value is set between -1 and 1. For the behavioral classification results, a segment of
the corresponding LFP frequency band that is set by the optimized bin width and offset is used as
the feature. For the onset detection analysis, the Pearson correlation for each sliding segment
equal in length to the optimal bin width is used to detect behavior that corresponds to its optimal
offset.

Computation and statistical testing of phase locking across renditions

We computed the Inter-trial phase coherence (ITPC) to assess trial-to-trial synchronization
of LFP activity with respect to time-locked syllables within the motif. The ITPC is a measure of
the phase synchronization of neural activity in a given frequency, calculated relative to critical
event times that are repeated over an experimental paradigm. To calculate the ITPC across trials,
for each channel, data were first filtered in 100 partially-overlapping narrow frequency bands that
were equal width in log-space. These filtered time series were then Hilbert transformed to extract
the analytic phase at each time point. Then, at each time point, for each channel and frequency
band, we calculated the phase-consistency across trials to estimate the ITPC. Here, 'trials' were
defined as the start or end of a specific type of vocal event, which could be further refined by its
context within the bout, e.g., first or last in bout. Once all instances of the event of interest were
selected, a window of time centered on each was implemented. The ITPC for each frequency and
time sample pair must be calculated independently. The equation for calculating the ITPC is
described as below, for n trials, if, $F_k(f, t)$ is the spectral estimate of trial k at frequency f and time
sample t

$$ITPC(f, t) = \frac{1}{n} \sum_{k=1}^{n} \frac{F_k(f, t)}{|F_k(f, t)|}$$

where $|x|$ represents the complex norm of x [62]. The spectral estimate, $F_k(f, t)$, is the instantaneous
phase of frequency f at time sample t. This was repeated for all narrow-band frequencies and time
samples within the event window. The ITPC value, or resultant vector length, scales from 0 to 1,
with 0 being the null hypothesis where phases are uniformly distributed about the polar axis and 1
being perfect synchrony. To determine the significance of this vector length, and to determine if it
could have been randomly sampled from a uniform distribution by chance, p-values were
calculated for each frequency and time sample. The mean resultant vector of the instantaneous
phase across trials for a specific time sample, and its corresponding bootstrapped p-value, were
calculated using the pycircstats toolbox. To enable visualization of the results for comparing ITPC
values while accounting for all of the frequency and time sample combinations, the Rayleigh Z
Statistic was calculated.

Rayleigh statistic The ITPC over the time length of a syllable or motif is evaluated in terms of the Rayleigh Z Statistic and is defined as $Z = \frac{R^2}{n}$ where R is the Rayleigh's R, $R = nr$ where r is the resultant vector length and n is the number of trials. P-values were estimated using the following equation [63–66], $P = exp\left[\sqrt{1 + 4n + 4(n^2 - R^2)} - (1 + 2n)\right]$.

Normalized sustained Rayleigh Z statistic. To determine which, if any, frequency bands could potentially be used as a feature to decode vocal behavior, a measure of how consistently significant a band stayed phase locked was created. As we don't expect the Rayleigh Z statistic across both channels and frequencies to be the exact same, we wanted to understand the relative significance across birds and channels. The measure ranks how consistently phase locked across time a frequency was during song production overall for one recording session. The Mean Rayleigh Z-Statistic of samples over the course of one oscillation for a set frequency normalized by the maximum Z-Statistic value for the entire time period analyzed was used and will be referred to as the Normalized Sustained Z Statistic. This analysis informed the frequency bands used for the classification and onset detection analysis.

Linear classifier

To classify behavior using the LFP features, a Linear Discriminant Analysis (LDA) model using Singular Value Decomposition (SVD) was trained using the scikit-learn toolbox [67] in Python. All priors for each class were set equal and all classes were balanced in the datasets used for classification analyses. Detailed information regarding the number of instances in each class can be found in S3-S5 Table. No shrinkage or regularization were used, however, the SVD optimizer was used to avoid ill-conditioned covariance matrices. Results were validated using 5-fold cross-validation. Templates for feature extraction were created by taking the mean across the training set. These templates were used to extract features from both the training set and the testing set. All frequencies were trained and tested independently of one another.

Channel adding curves were calculated using a bootstrap approach to determine how many channels were needed until additive information saturation. The channel adding analysis was repeated 5,000 times, changing the order in which each channel was included, with 5 fold cross validation. The order in which channels were added over the 5,000 repetitions was maintained across the folds to enable fair calculation of their validated mean. These are subsequently used to calculate the mean and standard deviation across repetitions.

Syllable onset detection

In order to analyze the temporal relationship between LFP and song production, we used a template matching approach to determine whether LFP can predict syllable onset. Each syllable of the motif excluding the first, had every labeled instance recorded in the same day split into training and testing sets using a 5-fold stratified split (80% training set and 20% testing) (S3-S5 Table). Both the template and the stereotyped onset of the syllable were calculated from the training set. Templates are the mean of the LFP traces of the training set. The stereotyped onset is the average time the syllable occurs in the training set with respect to the first syllable of the motif it occurred in. These templates were then used to compute the Pearson correlation across time for each of the motifs that contain the syllables of the test set, maintaining the temporal relationship of the optimal offset and the bin width for its respective frequency.

The prediction confidence of a single frequency band was calculated by first thresholding
the Pearson correlation values at zero, and taking the average of the resultant time series across
channels. The maximum confidence within a 100 millisecond window centered on the stereotyped
onset time is then used as the frequency's prediction for that instance of the syllable. To get the
results of a predictor that uses all of the frequencies the same steps as previously stated are
followed; only the prediction confidence across all frequencies and channels is used. The
prediction of the syllable onset using the bird's stereotyped behavior was determined by adding
the stereotyped onset time, calculated from the training set, to the actual start of the first syllable
for the motif the syllable occurred in. It is important to note two things with this approach; the first
is that neither approach receives information on the actual start time of the syllable in its respective
motif. The second is that this 100 millisecond window is significantly larger than the natural
variability of syllable onsets and gives the neural based predictor a more difficult task than the
baseline stereotyped comparison. Both predictions were then normalized by taking the difference
between the predicted time and the actual labeled start time of the syllable within the same motif.

Statistical significance of the results was calculated with a one-sided Wilcoxon signed-rank
test using the difference between the relative predictions of the stereotyped behavioral model and
the neural features models. Two null hypotheses were tested, first that there was no difference
between the stereotyped behavior and the prediction using neural features, and the second being
that the predictor using the neural features was closer to the actual labeled onset time. The second
null hypothesis requires that the LFP based predictor must outperform the behavior based
predictor. Results must pass both tests with a p-value less than .05 to be considered significant.
The Benjamini-Hochberg procedure was used to control the False Discovery Rate to account for
the multiple comparisons done. The procedure was implemented for each day treating the results
of each individual frequency and the predictor which uses all frequencies as separate results (n=6,
q=.05).
705

Acknowledgements

We would like to thank the Gilja, Gentner and Voytek Labs for feedback to help refine the
manuscript. We would like to especially thank Dr. Will Libertti III, Anna Mai, Abdulwahab
Alasfour, Tim Sainburg, Richard Gao, and Dr. Natalie Schaworonkow for insightful comments
and suggestions for improving the manuscript.
This work was supported by the U.S. NIH National Institute on Deafness and Other
Communication (R01DC008358, R01DC018446, R01DC018055), NIH National Institute of
General Medical Sciences (R01GM134363), National Science Foundation under grant BCS-
1736028, the Kavli Institute for the Brain and Mind (IRG #2016-004), the Office of
NavalResearch, Pew Latin American Fellowship in the Biomedical Sciences (E.A.), Halıcıoğlu
Data Science Institute Fellowship (J.I.C.), University of California – Historically Black Colleges
and Universities Initiative (D.E.B.). This material is based upon work supported by the National
Science Foundation Graduate Research Fellowship under Grant No. DGE-1650112. Any opinion,
findings, and conclusions or recommendations expressed in this material are those of the authors(s)
and do not necessarily reflect the views of the National Science Foundation.
721
722

References

1. Goller F, Cooper BG. Peripheral Motor Dynamics of Song Production in the Zebra Finch. Ann N Y Acad Sci. 2004;1016: 130–152. doi:10.1196/annals.1298.009

2. Jarvis ED. Evolution of vocal learning and spoken language. Science. 2019;366: 50–54. doi:10.1126/science.aax0287

3. Nowicki S. Vocal tract resonances in oscine bird sound production: evidence from birdsongs in a helium atmosphere. Nature. 1987;325: 53–55. doi:10.1038/325053a0

4. Churchland MM, Cunningham JP, Kaufman MT, Ryu SI, Shenoy KV. Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical Machine? Neuron. 2010;68: 387–400. doi:10.1016/j.neuron.2010.09.015

5. David A. Rosenbaum. Human Movement Initiation: Specification of Arm, Direction, and Extent. J Exp Psychol Gen. 1980;109: 444–474. doi:https://doi.org/10.1037/0096-3445.109.4.444

6. Fee MS, Kozhevnikov AA, Hahnloser RHR. Neural Mechanisms of Vocal Sequence Generation in the Songbird. Ann N Y Acad Sci. 2004;1016: 153–170. doi:10.1196/annals.1298.022

7. Ghez C, Hening W, Gordon J. Organization of voluntary movement. Curr Opin Neurobiol. 1991;1: 664–671. doi:10.1016/S0959-4388(05)80046-7

8. Kozhevnikov AA, Fee MS. Singing-Related Activity of Identified HVC Neurons in the Zebra Finch. J Neurophysiol. 2007;97: 4271–4283. doi:10.1152/jn.00952.2006

9. Daliparthi VK, Tachibana RO, Cooper BG, Hahnloser RH, Kojima S, Sober SJ, et al. Transitioning between preparatory and precisely sequenced neuronal activity in production of a skilled behavior. eLife. 2019;8: e43732. doi:10.7554/eLife.43732

10. Rajan R. Pre-Bout Neural Activity Changes in Premotor Nucleus HVC Correlate with Successful Initiation of Learned Song Sequence. J Neurosci. 2018;38: 5925–5938. doi:10.1523/JNEUROSCI.3003-17.2018

11. Bouchard KE, Mesgarani N, Johnson K, Chang EF. Functional organization of human sensorimotor cortex for speech articulation. Nature. 2013;495: 327–332. doi:10.1038/nature11911

12. Bouchard KE, Chang EF. Neural decoding of spoken vowels from human sensory-motor cortex with high-density electrocorticography. 2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society. Chicago, IL: IEEE; 2014. pp. 6782–6785. doi:10.1109/EMBC.2014.6945185

13. Martin S, Iturrate I, Millán J del R, Knight RT, Pasley BN. Decoding Inner Speech Using Electrocorticography: Progress and Challenges Toward a Speech Prosthesis. Front Neurosci. 2018;12: 422. doi:10.3389/fnins.2018.00422

761   14.   Rabbani Q, Milsap G, Crone NE. The Potential for a Speech Brain–Computer Interface
762   Using Chronic Electrocorticography. Neurotherapeutics. 2019;16: 144–165.
763   doi:10.1007/s13311-018-00692-2

764   15.   Sahin NT, Pinker S, Cash SS, Schomer D, Halgren E. Sequential Processing of Lexical,
765   Grammatical, and Phonological Information Within Broca's Area. Science. 2009;326: 445–
766   449. doi:10.1126/science.1174481

767   16.   Churchland MM, Santhanam G, Shenoy KV. Preparatory Activity in Premotor and Motor
768   Cortex Reflects the Speed of the Upcoming Reach. J Neurophysiol. 2006;96: 3130–3146.
769   doi:10.1152/jn.00307.2006

770   17.   Tanji J, Evarts EV. Anticipatory activity of motor cortex neurons in relation to direction of an
771   intended movement. J Neurophysiol. 1976;39: 1062–1068. doi:10.1152/jn.1976.39.5.1062

772   18.   Svoboda K, Li N. Neural mechanisms of movement planning: motor cortex and beyond.
773   Curr Opin Neurobiol. 2018;49: 33–41. doi:10.1016/j.conb.2017.10.023

774   19.   Erlich JC, Bialek M, Brody CD. A Cortical Substrate for Memory-Guided Orienting in the
775   Rat. Neuron. 2011;72: 330–343. doi:10.1016/j.neuron.2011.07.010

776   20.   Guo ZV, Li N, Huber D, Ophir E, Gutnisky D, Ting JT, et al. Flow of Cortical Activity
777   Underlying a Tactile Decision in Mice. Neuron. 2014;81: 179–194.
778   doi:10.1016/j.neuron.2013.10.020

779   21.   Li N, Chen T-W, Guo ZV, Gerfen CR, Svoboda K. A motor cortex circuit for motor planning
780   and movement. Nature. 2015;519: 51–56. doi:10.1038/nature14178

781   22.   Sainburg T, Theilman B, Thielk M, Gentner TQ. Parallels in the sequential organization of
782   birdsong and human speech. Nat Commun. 2019;10: 3636. doi:10.1038/s41467-019-
783   11605-y

784   23.   Angrick M, Herff C, Mugler E, Tate MC, Slutzky MW, Krusienski DJ, et al. Speech
785   synthesis from ECoG using densely connected 3D convolutional neural networks. J Neural
786   Eng. 2019;16: 036019. doi:10.1088/1741-2552/ab0c59

787   24.   Anumanchipalli GK, Chartier J, Chang EF. Speech synthesis from neural decoding of
788   spoken sentences. Nature. 2019;568: 493–498. doi:10.1038/s41586-019-1119-1

789   25.   Herff C, Schultz T. Automatic Speech Recognition from Neural Signals: A Focused
790   Review. Front Neurosci. 2016;10. doi:10.3389/fnins.2016.00429

791   26.   Stavisky SD, Willett FR, Wilson GH, Murphy BA, Rezaii P, Avansino DT, et al. Neural
792   ensemble dynamics in dorsal motor cortex during speech in people with paralysis. eLife.
793   2019;8: e46015. doi:10.7554/eLife.46015

794   27.   Brainard MS, Doupe AJ. What songbirds teach us about learning. Nature. 2002;417: 351–
795   358. doi:10.1038/417351a

796   28.   Nottebohm F. The Neural Basis of Birdsong. PLoS Biol. 2005;3: e164.
797   doi:10.1371/journal.pbio.0030164

29. Okubo TS, Mackevicius EL, Payne HL, Lynch GF, Fee MS. Growth and splitting of neural
sequences in songbird vocal development. Nature. 2015;528: 352–357.
doi:10.1038/nature15741

30. Woolley SMN. Early experience shapes vocal neural coding and perception in songbirds.
Dev Psychobiol. 2012;54: 612–631. doi:10.1002/dev.21014

31. Liberti WA, Markowitz JE, Perkins LN, Liberti DC, Leman DP, Guitchounts G, et al.
Unstable neurons underlie a stable learned behavior. Nat Neurosci. 2016;19: 1665–1671.
doi:10.1038/nn.4405

32. Poole B, Markowitz JE, Gardner TJ. The Song Must Go On: Resilience of the Songbird
Vocal Motor Pathway. Coleman MJ, editor. PLoS ONE. 2012;7: e38173.
doi:10.1371/journal.pone.0038173

33. Markowitz JE, Liberti WA, Guitchounts G, Velho T, Lois C, Gardner TJ. Mesoscopic
Patterns of Neural Activity Support Songbird Cortical Sequences. Ashe J, editor. PLOS
Biol. 2015;13: e1002158. doi:10.1371/journal.pbio.1002158

34. Picardo MA, Merel J, Katlowitz KA, Vallentin D, Okobi DE, Benezra SE, et al. Population-
Level Representation of a Temporal Sequence Underlying Song Production in the Zebra
Finch. Neuron. 2016;90: 866–876. doi:10.1016/j.neuron.2016.02.016

35. Schmidt MF. Pattern of Interhemispheric Synchronization in HVc During Singing Correlates
With Key Transitions in the Song Pattern. J Neurophysiol. 2003;90: 3931–3949.
doi:10.1152/jn.00003.2003

36. Bottjer SW, Johnson F. Circuits, hormones, and learning: Vocal behavior in songbirds. :
17.

37. Luo M, Perkel DJ. A GABAergic, Strongly Inhibitory Projection to a Thalamic Nucleus in
the Zebra Finch Song System. J Neurosci. 1999;19: 6700–6711.
doi:10.1523/JNEUROSCI.19-15-06700.1999

38. Nottebohm F, Stokes TM, Leonard CM. Central control of song in the canary,Serinus
canarius. J Comp Neurol. 1976;165: 457–486. doi:10.1002/cne.901650405

39. Vates GE, Vicario DS, Nottebohm F. Reafferent thalamo-"cortical" loops in the song
system of oscine songbirds. : 16.

40. Doupe AJ, Kuhl PK. BIRDSONG AND HUMAN SPEECH: Common Themes and
Mechanisms. Annu Rev Neurosci. 1999;22: 567–631. doi:10.1146/annurev.neuro.22.1.567

41. Pfenning AR, Hara E, Whitney O, Rivas MV, Wang R, Roulhac PL, et al. Convergent
transcriptional specializations in the brains of humans and song-learning birds. Science.
2014;346: 1256846–1256846. doi:10.1126/science.1256846

42. Buzsáki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents — EEG,
ECoG, LFP and spikes. Nat Rev Neurosci. 2012;13: 407–420. doi:10.1038/nrn3241

43. Carmena JM, Lebedev MA, Crist RE, O'Doherty JE, Santucci DM, Dimitrov DF, et al.
Learning to Control a Brain–Machine Interface for Reaching and Grasping by Primates.
Idan Segev, editor. PLoS Biol. 2003;1: e42. doi:10.1371/journal.pbio.0000042

44. Fetz EE. Operant Conditioning of Cortical Unit Activity. Science. 1969;163: 955–958.
doi:10.1126/science.163.3870.955

45. Gilja V, Nuyujukian P, Chestek CA, Cunningham JP, Yu BM, Fan JM, et al. A high-
performance neural prosthesis enabled by control algorithm design. Nat Neurosci.
2012;15: 1752–1757. doi:10.1038/nn.3265

46. Kubánek J, Miller KJ, Ojemann JG, Wolpaw JR, Schalk G. Decoding flexion of individual
fingers using electrocorticographic signals in humans. J Neural Eng. 2009;6: 066001.
doi:10.1088/1741-2560/6/6/066001

47. Miller KJ, Leuthardt EC, Schalk G, Rao RPN, Anderson NR, Moran DW, et al. Spectral
Changes in Cortical Surface Potentials during Motor Movement. J Neurosci. 2007;27:
2424–2432. doi:10.1523/JNEUROSCI.3886-06.2007

48. Kosche G, Vallentin D, Long MA. Interplay of Inhibition and Excitation Shapes a Premotor
Neural Sequence. J Neurosci. 2015;35: 1217–1227. doi:10.1523/JNEUROSCI.4346-
14.2015

49. Leonardo A. Ensemble Coding of Vocal Control in Birdsong. J Neurosci. 2005;25: 652–
661. doi:10.1523/JNEUROSCI.3036-04.2005

50. Miller KJ, Zanos S, Fetz EE, den Nijs M, Ojemann JG. Decoupling the Cortical Power
Spectrum Reveals Real-Time Representation of Individual Finger Movements in Humans.
J Neurosci. 2009;29: 3132–3137. doi:10.1523/JNEUROSCI.5506-08.2009

51. Brovelli A, Lachaux J-P, Kahane P, Boussaoud D. High gamma frequency oscillatory
activity dissociates attention from intention in the human premotor cortex. NeuroImage.
2005;28: 154–164. doi:10.1016/j.neuroimage.2005.05.045

52. Canolty RT, Edwards E, Dalal SS, Soltani M, Nagarajan SS, Kirsch HE, et al. High Gamma
Power Is Phase-Locked to Theta Oscillations in Human Neocortex. Science. 2006;313:
1626–1628. doi:10.1126/science.1128115

53. Pfurtscheller G, Graimann B, Huggins JE, Levine SP, Schuh LA. Spatiotemporal patterns
of beta desynchronization and gamma synchronization in corticographic data during self-
paced movement. Clin Neurophysiol. 2003;114: 1226–1236. doi:10.1016/S1388-
2457(03)00067-1

54. Jiang W, Pailla T, Dichter B, Chang EF, Gilja V. Decoding speech using the timing of
neural signal modulation. 2016 38th Annual International Conference of the IEEE
Engineering in Medicine and Biology Society (EMBC). Orlando, FL, USA: IEEE; 2016. pp.
1532–1535. doi:10.1109/EMBC.2016.7591002

55. Hyland Bruno J, Tchernichovski O. Regularities in zebra finch song beyond the repeated
motif. Behav Processes. 2019;163: 53–59. doi:10.1016/j.beproc.2017.11.001

56.   Miller KJ, Hermes D, Honey CJ, Hebb AO, Ramsey NF, Knight RT, et al. Human Motor
Cortical Activity Is Selectively Phase-Entrained on Underlying Rhythms. Behrens T, editor.
PLoS Comput Biol. 2012;8: e1002655. doi:10.1371/journal.pcbi.1002655

57.   G. Makin J, Edward F. Chang, A. Moses D. Machine translation of cortical activity to text
with an encoder-decoder framework. : 22.

58.   Ezequiel Arneodo. Software and hardware designs for chronic, high channel count
electrophysiology. Biocircuits Institute; 2016. Available:
https://github.com/singingfinch/bernardo.git

59.   Arneodo EM, Chen S, Gilja V, Gentner TQ. A neural decoder for learned vocal behavior.
Neuroscience; 2017 Sep. doi:10.1101/193987

60.   Gramfort A. MEG and EEG data analysis with MNE-Python. Front Neurosci. 2013;7.
doi:10.3389/fnins.2013.00267

61.   Paul Boersma, David Weenink. Praat: doing phonetics by computer. 2018. Available:
http://www.praat.org/

62.   Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. J Neurosci Methods. 2004;134: 9–
21. doi:10.1016/j.jneumeth.2003.10.009

63.   Berens P. CircStat: A MATLAB Toolbox for Circular Statistics. J Stat Softw. 2009;31.
doi:10.18637/jss.v031.i10

64.   Fisher NI. Statistical Analysis of Circular Data. 1st ed. Cambridge University Press; 1993.
doi:10.1017/CBO9780511564345

65.   Jammalamadaka SR, Sengupta A. Topics in circular statistics. River Edge, N.J: World
Scientific; 2001.

66.   Zar JH. Biostatistical analysis. 5th ed. Upper Saddle River, N.J: Prentice-Hall/Pearson;
2010.

67.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. Mach Learn PYTHON. 2011; 6.

899
900

Supporting information:

S1 Table. Overall Description of Subject Data

S2 Table. Detailed description of available labeled data

S3 Table. Behavior of subject z007 during high yield days

S4 Table. Behavior of subject z020 during high yield days

S5 Table. Behavior of subject z017 during high yield days

S6 Table. Bin Width Hyperparameter Search Results

S7 Table. Offset Hyperparameter Search Results

S1 Fig. Song correlated changes in LFP power across subjects and days

S2 Fig. Inter-trial phase coherence of LFP phase during production of learned sequences across subjects and days.

910
911

S3 Fig. Unique LFP phase preferences at each syllable onset (z007, day 1).

S4 Fig. Unique LFP phase preferences at each syllable onset (z020).

S5 Fig. Unique LFP phase preferences at each syllable onset (z007).

S6 Fig. Unique LFP phase preferences for sparsely used intra-motif note onset (z017).

S7 Fig. Phase preference to syllable phase preference is reflective to constitutive syllables rather than the larger motif structure.

916
917

S8 Fig. Detailed view of the syllable phase preference to syllable onset for z007.

S9 Fig. Both Phase and Power has independent/additive Information to classify syllable identity.

919
920

S10 Fig. Channel Dropping Curves for the second highest yield days across subjects.

S11 Fig. Onset detection across all high yield days.

Overall Description				
bird_id	# of Days Total	Motif Length	Intra Motif	Introductory Note?
z020	5	4	1	Y
z007	9	5	0	Y
z017	19?	7	2	N

923 **S1 Table: Overall Description of Subject Data**

Bird ID	Date	# of VAPs	Duration (hr. min. sec.)	# of Autodetected Motifs	# of VAEs
z020	day-2016-06-02	6	5 hr. 13 min 9.092 sec	34	6
z020	day-2016-06-03	27	7 hr. 32 min 18.526 sec	113	27
z020	day-2016-06-04	14	7 hr. 43 min 43.603 sec	24	14
z020	day-2016-06-05	15	3 hr. 9 min 32.552 sec	85	15
z020	day-2016-06-06	9	2 hr. 40 min 39.42 sec	44	9
z007	day-2016-09-07	10	2 hr. 32 mins 46.24 sec	37	10
z007	day-2016-09-09	5	2 hr. 19 min 5.739 sec	19	5
z007	day-2016-09-10	14	2 hr. 13 min 26.45 sec	71	14
z007	day-2016-09-11	20	5 hr. 14 min 40.959 sec	92	20
z007	day-2016-09-12	11	5 hr. 27 min 7.001 sec	44	11
z007	day-2016-09-13	5	1 hr. 18 min 5.539 sec	60	5
z017	day-2016-06-19	18	8 hr. 38 min 29.42 sec	97	18
z017	day-2016-06-21	16	10 hr. 8 min 10.959 sec	76	16

926 **S2 Table: Detailed Description of available labeled data**

z007			
Day 1			
Syllable	# of Instances	Duration (ms.) +/- std	Start within Motif relative to First Syllable (ms.) +/- std
1	72	100.084 +/- 3.558	-
2	72	55.406 +/- 2.877	125.694 +/- 3.101
3	71	164.915 +/- 3.296	221.830 +/- 4.087
4	71	120.870 +/- 2.654	448.069 +/- 6.111
5	71	53.958+/- 2.620	604.796 +/- 7.754
I	157	53.520+/- 4.662	-
Day 2			
Syllable	# of Instances	Duration (ms.) +/- std	Relative Start within Motif (ms.)
1	100	102.532 +/- 6.561	-
2	100	57.612 +/- 7.598	127.509 +/- 6.525
3	100	167.799 +/- 8.020	224.254 +/- 8.082
4	100	124.418 +/- 5.822	452.354 +/- 9.369
5	98	55.548 +/- 4.489	611.310 +/- 9.780
I	178	53.621 +/- 6.870	-

928 **S3 Table: Behavior of Subject z007 during High Yield Days**
929

z020			
Day 1			
Syllable	# of Instances	Duration (ms) +/- std	Start within Motif relative to First Syllable (ms) +/- std
1	160	38.882 +/- 4.716	-
2	160	132.902 +/- 8.230	65.039 +/- 5.053
3	160	49.861 +/- 5.099	203.551 +/- 10.469
4	91	110.964 +/- 3.476	347.219 +/- 23.193
I	235	42.081 +/- 5.792	-
Day 2			
Syllable	# of Instances	Duration (ms) +/- std	Relative Start within Motif (ms) +/- std
1	109	39.887 +/- 6.139	-
2	109	132.925 +/- 4.865	67.670 +/- 5.01
3	109	50.257 +/- 5.011	206.310 +/- 4.985
4	75	111.741 +/- 6.112	348.720 +/- 12.329
I	191	46.922 +/- 9.378	-

930 **S4 Table: Behavior of Subject z020 during High Yield Days**
931

z017			
Day 1			
Syllable	# of Instances	Duration (ms) +/- std	Start within Motif relative to First Syllable (ms) +/- std
1	97	50.692 +/- 5.341	-
2	97	57.389 +/- 6.629	73.109 +/- 5.061
3	97	117.752 +/- 7.757	157.192 +/- 7.368
4	97	106.304 +/- 5.832	307.916 +/- 9.093
5	97	127.465 +/- 4.870	420.749 +/- 9.343
6	97	154.392 +/- 6.617	591.343 +/- 13.956
7	52	72.894 +/- 8.067	809.046 +/- 21.541
Day 2			
Syllable	# of Instances	Duration (ms) +/- std	Relative Start within Motif (ms) +/- std
1	84	49.162 +/- 6.525	-
2	84	51.217+/- 9.207	75.058 +/- 6.447
3	86	114.807 +/- 15.198	155.762 +/- 12.082
4	83	106.109 +/- 8.912	306.191 +/- 18.28
5	82	127.825 +/- 11.710	420.302 +/- 22.832
6	66	154.634+/- 11.803	584.710 +/- 14.729
7	41	68.696 +/- 8.431	803.838 +/- 26.629

932 **S5 Table: Behavior of Subject z017 during High Yield Days**
933

Bird ID	Date	Best Bin Width (ms)				
		4-8 Hz	8-12 Hz	25-35 Hz	35-50 Hz	50-70 Hz
z020	day-2016-06-03	90	185	90	85	90
z020	day-2016-06-05	190	125	35	160	70
z007	day-2016-09-10	170	75	90	25	30
z007	day-2016-09-11	195	170	70	20	70
z017	day-2016-06-19	175	170	85	50	35
z017	day-2016-06-21	150	170	70	60	40

934   **S6 Table: Bin Width Hyperparameter Search Results**
935

Bird ID	Date	Best Offset (ms)				
		4-8 Hz	8-12 Hz	25-35 Hz	35-50 Hz	50-70 Hz
z020	day-2016-06-03	-10	-5	0	0	0
z020	day-2016-06-05	-25	-5	0	-20	0
z007	day-2016-09-10	-35	-5	-10	-10	0
z007	day-2016-09-11	-5	-5	-15	0	-15
z017	day-2016-06-19	-50	0	-10	-5	-5
z017	day-2016-06-21	-10	-15	-30	0	0

937   **S7 Table: Offset Hyperparameter Search Results**

S1 Fig: Song correlated changes in LFP power across subjects and days. Averaged spectrotemporal power activity aligned to the start of the first motif in the bout, left, and the last motif in the bout, right, for each recording day not shown in Fig 3. Above all results is a behavioral raster showing the time course of the behavior being averaged. (A) The averaged results for the second highest yielding day, designated Day 1, for z007 (n=25 Bouts). The other subjects results are show as follows; (B) z020's first high yield day (n=29 Bouts), (C) z020's second high yield day (n=25 Bouts), (D) z017's first high yield day (n=27 Bouts), and (E) z017's second high yield day (n=21 Bouts). Because z017 would end its bout on either syllable '5' or '6', the end of the bout was aligned to syllable '5'. No dynamic-time warping was used. To ensure the start and end of the bout are unique time periods, only bouts with more than one motif in duration were used. Behaviorally inconsistent bouts were excluded for clarity of visualization, however results are consistent when including them in the analysis.

S2 Fig: Inter-trial phase coherence of LFP phase during production of learned sequences across subjects and days. ITPC of LFP aligned to the start of the first motif in the bout, left, and the last motif in the bout, right, for each recording day not shown in Fig 3. Above all results is a Behavioral raster showing the time course of the behavior being averaged. (A) The averaged results for the second highest yielding day, designated Day 1, for z007 (n=25 bouts). The other subjects results are show as follows; (B) z020's first high yield day (n=29 bouts), (C) z020's second high yield day (n=25 bouts), (D) z017's first high yield day (n=27 bouts), and (E) z017's second high yield day (n=21 bouts). Because z017 would end its bout on either syllable '5' or '6', the end of the bout was aligned to syllable '5'. No dynamic-time warping was used. To ensure the start and end of the bout are unique time periods, only bouts with more than one motif in duration were used. Behaviorally inconsistent bouts were excluded for clarity of visualization, however results are consistent when including them in calculating the ITPC.

965
S3 Fig: Unique LFP phase preferences at each syllable onset (z007, day 1). (A) Polar histogram
of the phase for each LFP frequency band at the labeled start of all instances of a given syllable
over the course of one day (Day 1), for one bird (z007). The number of instances (n=71) are equal
for all syllables, and set by the syllable class with the fewest renditions. (B) ITPC resultant vector
length for each frequency over time relative to the labeled start of each syllable (0 ms) over the
same instances as in (A). (C) Rayleigh Z-statistic of the ITPC over the same time and frequencies
as (B).
973
974

S4 Fig: Unique LFP phase preferences at each syllable onset (z020). (A) Polar histogram of the phase for each LFP frequency band at the labeled start of all instances of a given syllable over the course of one day (Day 1), for one bird (z020). The number of instances (n=91) are equal for all syllables, and set by the syllable class with the fewest renditions. (B) ITPC resultant vector length for each frequency over time relative to the labeled start of each syllable (0 ms) over the same instances as in (A). (C) Rayleigh Z-statistic of the ITPC over the same time and frequencies as (B). (D) Polar histogram of the phase for each LFP frequency band at the labeled start of all instances of a given syllable over the course of one day (Day 2), for one bird (z020). The number of instances

984   (n=75) are equal for all syllables, and set by the syllable class with the fewest renditions. (E) ITPC
985   resultant vector length for each frequency over time relative to the labeled start of each syllable
986   (0 ms) over the same instances as in (D). (F) Rayleigh Z-statistic of the ITPC over the same time
987   and frequencies as (E).
988

S5 Fig: Unique LFP phase preferences at each syllable onset (z007). (A) Polar histogram of the phase for each LFP frequency band at the labeled start of all instances of a given syllable over the course of one day (Day 1), for one bird (z020). The number of instances (n=97) are equal for all syllables, and set by the syllable class with the fewest renditions. (B) ITPC resultant vector length for each frequency over time relative to the labeled start of each syllable (0 ms) over the same instances as in (A). (C) Rayleigh Z-statistic of the ITPC over the same time and frequencies as (B). (D) Polar histogram of the phase for each LFP frequency band at the labeled start of all instances

998   of a given syllable over the course of one day (Day 2), for one bird (z020). The number of instances
999   (n=82) are equal for all syllables, and set by the syllable class with the fewest renditions. (E) ITPC
1000 resultant vector length for each frequency over time relative to the labeled start of each syllable
1001 (0 ms) over the same instances as in (D). (F) Rayleigh Z-statistic of the ITPC over the same time
1002 and frequencies as (E).
1003

1004
S6 Fig: Unique LFP phase preferences for sparsely used intra-motif note onset (z017). (A) Polar
histogram of the phase for each LFP frequency band at the labeled start of all instances (n=52) of
syllable 7 over the course of one day (Day 1), for bird z017. (B) ITPC resultant vector length for
each frequency over time relative to the labeled start of syllable 7 (0 ms) over the same instances
as in (A). (C) Rayleigh Z-statistic of the ITPC over the same time and frequencies as (B). (D) Polar
histogram of the phase for each LFP frequency band at the labeled start of all instances (n=41) of
syllable 7 over the course of one day (Day 2), for bird z017. (E) ITPC resultant vector length for
each frequency over time relative to the labeled start of syllable 7 (0 ms) over the same instances
as in (D). (F) Rayleigh Z-statistic of the ITPC over the same time and frequencies as (E).

1014

S7 Fig: Phase preference to syllable phase preference is reflective to constitutive syllables
rather than the larger motif structure. Box plots of the differences in Rayleigh Statistic
between the syllable aligned ITPC and the ITPC aligned to that same syllables' stereotyped
onset time within the motif. This difference was determined by first combining the Z Statistic
for each channel centered at the designated time point to get a vector of all partially
overlapping frequencies for all channels. The Vector of stereotyped alignment was then
subtracted from the labeled onset alignment to get the difference for each frequency on every
channel. This was repeated for all syllable, excluding the first, with each syllable represented
with a specific color as indicated. (A) Shows the results for the two high yield days for z007, (B)
is the results for z020, and (C) is the results for z017. All instances of each syllable that were
preceded by the first syllable were used, which means that classes are not balanced and are
reflective of all existing data. To determine statistical significance the one sided Wilcoxon
signed-rank test was used with each frequency and channel pair. * denotes results that were
not statistically significant when using the using the Benjamini-Hochberg False Discovery Rate.
All other results p<0.05 and q<0.05.

1030

1031

S8 Fig: Detailed view of the syllable phase preference to syllable onset for z007. Detailed
rendering of the phase preference to syllable onset for z007 shown in Fig 6A. Each Row is a
different vocalization type which includes the five syllables of the motif and the introductory
note. Each column is a different frequency band and is organized top to bottom from least to
greatest. As such they are the polar plots for the (A) 4-8 Hz band, (B) 8-12 Hz band, (C) 25-35 Hz
band, (D) 35-50 Hz band, and (E) the 50-70 Hz band. The number of instances have been
balanced to match the class with the least number of instances (n=100) for each class.

1039

S9 Fig: Both Phase and Power has independent/additive Information to classify syllable identity.

Channel Dropping Curves of classifier performances with either (A) all information about phase removed, (B) all information about power removed, or (C) with both phase and power used as independent features. Each row corresponds to data from the highest yield day for each bird. z007 n=98 for each class n=7 (1, 2, 3, 4, 5, i, Silence), z020 n=91 for each class n = 6 (1, 2, 3, 4, i, Silence), and for z017 n=52* for each class n=8 (1, 2, 3, 4, 5, 6, 7, Silence). Error bars represent the standard deviation over the bootstrapped analysis using n=5,000 repetitions across 5 cross-validation folds. The p-value for all of the Binomial chances calculated for each bird was 0.05.

*The number of instances for each class was limited by Syllable 7, which is a intra-motif note.

1051

1052
1053

S10 Fig: Channel Dropping Curves for the second highest yield days across subjects

Channel Dropping Curves of classifier performances with either (A) all phase related information removed, (B) all power related information removed, or (C) with both phase and power used as independent features. Each row corresponds to data from the second highest yielding day for each bird. z007 n=71 for each class n=7 (1, 2, 3, 4, 5, i, Silence), z020 n=75 for each class n = 6(1, 2, 3, 4, i, Silence), and for z017 n=41* for each class n=8 (1, 2, 3, 4, 5, 6, 7, Silence). Error bars represent the standard deviation over the bootstrapped analysis using n=5,000 repetitions across 5 cross-validation folds. The p-value for all of the Binomial chances calculated for each bird was 0.05.

*The number of instances for each class was limited by Syllable 7, which is a intra-motif note.

1066
S11 Fig: Onset detection across all high yield days.
Boxplots of onset prediction times relative to the labeled onset time for each bird for every
syllable for two highest yielding days. Each Column reflects the result for syllable number within
the motif. Each row is for a specific bird with (A) corresponding to z007, (B) corresponding to
z020, and (C) corresponding to z017. The order of each feature used is the same, going left to
right: first is the stereotyped onset time using only the deterministic behavior, next is the results
using all of the neural features, then each frequency band only in order from least to greatest (4-
8 Hz, 8-12 Hz, 25-35 Hz, 35-50 Hz, and finally 50-70 Hz). The recording day designation number
refers to the chronological order that the recordings took place. Statistical significance was
calculated using the one sided Wilcoxon signed-rank test, and * denotes results that were **not**
statistically significant when using the Benjamini-Hochberg False Discovery Rate. All other results
p<0.05 and q<0.05.

49